1

# Integrative experimental/computational approach establishes active cellular protrusion as the primary driving force of phagocytic spreading by immune cells

7    Short Title: Computational analysis of immune cell spreading

10   Emmet A. Francis[1]*, Volkmar Heinrich[1]*

11   [1] Department of Biomedical Engineering, University of California Davis, Davis, CA, USA

12   * Corresponding author

13   E-mail: eafrancis@ucdavis.edu (EAF), vheinrich@ucdavis.edu (VH)

# Abstract

The dynamic interplay between cell adhesion and protrusion is a critical determinant of many forms of cell motility. When modeling cell spreading on adhesive surfaces, traditional mathematical treatments often consider passive cell adhesion as the primary, if not exclusive, mechanistic driving force of this cellular motion. To better assess the contribution of active cytoskeletal protrusion to immune-cell spreading during phagocytosis, we here develop a computational framework that allows us to optionally investigate both purely adhesive spreading ("Brownian zipper hypothesis") as well as protrusion-dominated spreading ("protrusive zipper hypothesis"). We model the cell as an axisymmetric body of highly viscous fluid surrounded by a cortex with uniform surface tension and incorporate as potential driving forces of cell spreading an attractive stress due to receptor-ligand binding and an outward normal stress representing cytoskeletal protrusion, both acting on the cell boundary. We leverage various model predictions against the results of a directly related experimental companion study of human neutrophil phagocytic spreading on substrates coated with different densities of antibodies. We find that the concept of adhesion-driven spreading is incompatible with experimental results such as the independence of the cell-spreading speed on the density of immobilized antibodies. In contrast, the protrusive zipper model agrees well with experimental findings and, when adapted to simulate cell spreading on discrete adhesion sites, it also reproduces the observed positive correlation between antibody density and maximum cell-substrate contact area. Together, our integrative experimental/computational approach shows that phagocytic spreading is driven by cellular protrusion, and that the extent of spreading is limited by the density of adhesion sites.

# Author Summary

To accomplish many routine biological tasks, cells must rapidly spread over different types of surfaces. Here, we examine the biophysical underpinnings of immune cell spreading during phagocytosis, the process by which white blood cells such as neutrophils engulf pathogens or other foreign objects. Our computational framework models the case in which a human neutrophil spreads over a flat surface coated with antibodies, which we also test experimentally in a companion paper. Our primary purpose is to assess whether phagocytic spreading is actively driven by protrusive forces exerted by the cell, or passively by adhesive forces acting between receptors in the cell membrane and antibodies on the surface. By directly comparing our model predictions to experimental results, we demonstrate that phagocytic spreading is primarily driven by protrusion, but the extent of spreading is still limited by the availability of binding sites. Our findings improve the fundamental understanding of phagocytosis and may also pave the way for future investigations of the balance between adhesion and protrusion in other forms of cell spreading, such as wound healing or cancer cell migration.

# Introduction

The ability of cells to spread on surfaces is critical to many biological processes, including cell migration, wound healing, and tissue formation. Immune cells are especially adept at rapid spreading; for instance, neutrophils commonly travel up to an order of magnitude faster in motility assays than fibroblasts or endothelial cells [1-5]. Vital immune cell functions, such as firm arrest at the endothelium and phagocytosis of pathogens, rely on this ability. A deeper understanding of the physical mechanisms driving cell spreading is essential for informed pharmaceutical strategies and new therapies targeting immune cells, such as current efforts to enhance phagocytosis for more effective clearance of cancer cells [6, 7].

Many experimental and theoretical studies have sought to unravel the physical principles of cell spreading by analyzing the initial growth of cell contact regions on a flat, adhesive substrate [8-11]. However, it often remains unclear whether any given type of cell spreading is primarily driven by passive forces due to adhesion or by active protrusive and/or contractile forces generated by the cell. In the simplest model, cell spreading is treated as a purely passive process, similar to a viscous droplet spreading on an adhesive substrate [8, 12, 13]. Alternative models assume that cytoskeleton-driven protrusion determines the rate of cell spreading, neglecting the effective attractive force due to ligand-receptor binding [10, 14, 15].

Here, we aim to illuminate fundamental cell-spreading mechanisms by focusing on phagocytic spreading. In conventional phagocytosis, an immune cell first adheres to the surface of a target particle such as a bacterium, often by binding to opsonic ligands such as immunoglobulin G (IgG) antibodies. The cell then spreads over the target surface until its membrane closes around the particle. On the other hand, when presented with an excessively

large target, the cell is unable to fully engulf the particle and continues to spread until its target-

contact area reaches a maximum [16, 17]. Such cases, including phagocytic spreading on flat

substrates, are termed frustrated phagocytosis. The relatively simple geometry of a cell

undergoing frustrated phagocytic spreading on a flat surface is particularly amenable to

theoretical analysis, especially when the cell spreads in an axisymmetric manner.

Reflecting the above-mentioned contrasting mechanistic notions about generic cell

spreading, two hypotheses have been put forward to explain phagocytic spreading, i.e., the

"Brownian zipper" and "protrusive zipper" models [18]. Both models concur that fresh contact

between the cell membrane and target surface results in a zipper-like adhesive attachment that is

essentially irreversible. However, regarding the driving force of cellular advancement, the

Brownian zipper model postulates that strong adhesion alone is responsible for pulling the cell

membrane onto the target surface. In contrast, the protrusive zipper model assumes that the

cytoskeleton actively generates protrusive force to push the front of the cell outwards.

We seek to establish which of these hypotheses is more realistic by developing

mathematical models of the mechanics of isotropic cell spreading and comparing our theoretical

predictions with experimental observations of human neutrophils spreading on flat, IgG-coated

surfaces. As reported in detail in a companion paper, our experimental design allowed us to

examine the spreading behavior of neutrophils on surfaces coated with different densities of IgG

[19]. Because higher IgG densities correspond to stronger adhesive forces, we reasoned that the

Brownian zipper model should predict a much stronger correlation between IgG density and rate

of contact-area growth ("spreading speed") than the protrusive zipper model. Here, we use

computer simulations of phagocytic spreading to verify this common-sense expectation and

further investigate mechanisms governing immune-cell spreading. Together, our integrative

experimental/theoretical approach establishes that the Brownian zipper model is inconsistent

with experimental results, whereas the spreading dynamics predicted by the protrusive zipper

model agree well with our experiments.

## Results

## Model implementation of key biophysical mechanisms of cell

## spreading

We consider a simple model cell that captures the basic geometrical and mechanical properties of

human neutrophils, the most abundant type of white blood cell and one of the prototypes of

professional phagocytes. This model cell consists of an axisymmetric, highly viscous fluid body

of constant volume that is surrounded by an elastic cortex with uniform tension (Fig 1). Termed

the "cortical shell, liquid core model", this theoretical concept has successfully reproduced the

passive response of human neutrophils to imposed deformations [20-22]. In the undeformed

reference state, the persistent tension of the cortex maintains a spherical cell shape with a typical

diameter of 8.5 μm. The nucleus of neutrophils is segmented into three to five lobes and appears

to play only a minor role during deformations, lending support to the approximation of the cell

interior as a homogeneous fluid. The effective viscosity ($\mu$) of the cytoplasm has been found to

be about 200,000 times the viscosity of water at room temperature in passive neutrophils [20,

21], and another ~3-times higher in active neutrophils [23]. It is well known to increase during

phagocytosis [24-26].  We use a viscosity value of $\mu = 200$ Pa-s in our passive Brownian zipper

model and a value of $\mu = 1660$ Pa-s in our active protrusive zipper model. The original

assumption of a constant cortical tension [20, 22] was later found to be approximately true only

for small deformations. The revised relationship between tension ($\tau$) and cell surface area ($A_{cell}$)

was shown to be roughly biphasic [27]. Assuming an isotropic and uniform tension, and in

overall agreement with a number of quantitative experimental studies [24, 27, 28], our

simulations incorporate a smoothed version of the following biphasic constitutive relationship:

132
$$\tau(A_{cell}) = \begin{cases} 0.01 + 0.16\left(\dfrac{A_{cell}}{A_{cell,0}} - 1\right)\dfrac{\text{mN}}{\text{m}} & \text{if} \quad \dfrac{A_{cell}}{A_{cell,0}} \leq 1.26 \\[4mm] 0.0516 + 0.545\left(\dfrac{A_{cell}}{A_{cell,0}} - 1.26\right)\dfrac{\text{mN}}{\text{m}} & \text{if} \quad \dfrac{A_{cell}}{A_{cell,0}} > 1.26 \end{cases}$$
(1)

133   Where $A_{cell,0}$ is the surface area of the resting neutrophil before spreading.

**Fig 1. Illustration of our computational model of isotropic cell spreading.**

(A) 3D rendering shows the geometry and defines coordinates for an axisymmetric spreading

cell on a flat surface. The enlarged inset (right) illustrates the stress balance at the free cell

boundary. (B) Adhesion stress effectively pulls the membrane onto the flat surface. The enlarged

insets conceptually depict how the adhesion stress is computed for different ligand densities $\rho_l$.

(C) Cortical tension and membrane curvature give rise to an effective inward normal stress. (D)

The protrusion stress acts normal to the cell membrane and is concentrated at the region of the

membrane closest to the substrate. (E) The cross-sectional snapshot of a simulation illustrates the

mesh composed of quadrilateral elements used in the calculations, with tighter element packing

closer to the flat surface. The computational domain only includes half of this mesh due to axial

symmetry. Boundary regions are labeled $\Gamma$, and the vertical dashed line is the symmetry axis ($r$

$= 0$) (F) This snapshot of a simulation illustrates the fluid velocity (vector field) and relative

pressure (heat map) computed for a given cell shape with known boundary stresses.

Movements of the viscous cytoplasm generally do not exceed velocities of 1 μm/s during

phagocytic cell spreading and are characterized by low Reynolds numbers. Such movements are

well described by the Stokes equations for creeping flow:

$$\mu \nabla^2 \mathbf{v} = \nabla p \tag{2}$$

along with the continuity equation for incompressible fluids:

$$\nabla \cdot \mathbf{v} = 0 \tag{3}$$

Here, $\mathbf{v}$ denotes the fluid velocity vector, and $p$ is fluid pressure. Together, these equations

govern the flow behavior of our model cell.

Interactions between the cell surface and the substrate can facilitate cell spreading

directly via adhesion, and indirectly through receptor-induced signaling that leads to protrusion.

Including the respective stresses, the moving boundary condition governing the free (i.e.,

nonadherent) part of the cell surface ($\Gamma_{free}$) balances up to four types of stress vectors: fluid stress

($\sigma_{fluid}$), adhesion stress ($\sigma_{adh}$), protrusion stress ($\sigma_{fluid}$), and cortical stress ($\sigma_{cortex}$) (Fig 1A):

$$\sigma_{fluid} + \sigma_{adh} + \sigma_{cortex} + \sigma_{prot} = \mathbf{0} \quad \text{on } \Gamma_{free} \tag{4}$$

The fluid stress vector $\boldsymbol{\sigma}_{fluid}$ encompasses the pressure difference across the cell surface

as well as the stress exerted on the surface by the moving cytoplasm. It is determined by the

constitutive relation for a Newtonian fluid:

166
$$\boldsymbol{\sigma}_{fluid} = \boldsymbol{\sigma}_{fluid,out} - \boldsymbol{\sigma}_{fluid,in} = (p - p_{out})\mathbf{n} - \mu\left(\nabla\mathbf{v} + (\nabla\mathbf{v})^{T}\right)\cdot\mathbf{n} \tag{5}$$

where $\mathbf{n}$ is the unit surface normal, and the dot in the last term on the right-hand side denotes

the inner product between a tensor and a vector. The extracellular medium has a much lower

viscosity than the intracellular fluid; therefore, Eq (5) neglects the medium's viscous

contribution and only accounts for its hydrostatic pressure denoted $p_{out}$.

Our implementation of the adhesion stress is based on the assumption that adhesion can

effectively be represented by a short-range attractive potential between the cell surface and the

ligand-coated substrate. We compute the effective adhesion stress at a given point on the cell

surface by summing over all pairwise interactions with substrate molecules that lie outside the

cell-substrate contact region at the time (Fig 1B). For a continuum of ligand molecules, the

resulting expression takes the form (derived in S6 Appendices)

177
$$\boldsymbol{\sigma}_{adh} = \sigma_o\rho_l \iint\limits_{\text{substrate}} \mathbf{f}_{unit}\left[\left(\frac{D_0}{D}\right)^{7} - \left(\frac{D_0}{D}\right)^{4}\right]dA \tag{6}$$

where $\rho_l$ is the ligand surface density and $\sigma_0$ is a scaling factor representing the strength of

interaction per unit area of the cell surface. The unit vector $\mathbf{f}_{unit}$ specifies the direction of a given

interaction, $D$ denotes the distance between the interacting points, and $D_0$ is a constant setting

the overall range of the adhesion potential (set to 50 nm in our simulations).

The origin of the cortical stress is the cortical tension, which resists expansion of the cell

surface area and acts along the curved surface of the cell (Fig 1D). The resulting stress acts

normal to the surface and is given by the product of the tension $\tau$ and the mean curvature of the

cell surface, as dictated by Laplace's law:

$$\boldsymbol{\sigma}_{cortex} = -\tau\left(\frac{\partial\phi}{\partial s} + \frac{\sin\phi}{r}\right)\mathbf{n} \tag{7}$$

Here, the term in parentheses is the mean curvature of the axisymmetric cell surface expressed in

the coordinates defined in Fig 1A.

The protrusion stress comprises outward forces exerted on the cell surface by the actively

remodeling cytoskeleton. Purposeful reorganization of intracellular structures requires tight

coordination by biochemical signaling cues, which in turn are induced by receptor-ligand

binding. Rather than attempting to implement these highly complex mechano-chemical

mechanisms, we capture their localized character in a semiempirical manner. Our model

postulates that the protrusive stress is greatest at the perimeter of the cell-substrate contact region

and decays as a function of the distance from this leading edge measured along the arc length ($s$,

Fig 1A) of the contour of the unbound cell surface. We assume that this stress acts normal to the

cell boundary, and that its decay along the free cell surface is exponential (Fig 1C):

$$\boldsymbol{\sigma}_{prot} = \sigma_{prot}e^{-\frac{s_{contact}-s}{s_0}}\mathbf{n} \tag{8}$$

In this equation, $s_{contact}$ is the arc length value at which the cell-surface contour makes contact

with the substrate, and $s_0$ is the characteristic length that sets the spatial range of the protrusion

stress.

Bearing in mind that cytoskeletal remodeling also is responsible for the behavior of the

cortical tension, and that the tension varies during cell spreading (Eq (1)), it is reasonable to

assume that the overall strength of the protrusion stress ($\sigma_{prot}$) varies in a similar fashion. In

addition to the spatial distribution of the cortical stress governed by Eq (8), we therefore also

postulate a deformation-dependent strength of the protrusive stress. Our choice of this

dependence, and its motivation, will be explained in a later section.

Finally, for the part of the cell surface that is in contact with the substrate, we enforce a

simple no-slip boundary condition, requiring that

$$\mathbf{v} = \mathbf{0} \quad \text{on } \Gamma_{adherent} \tag{9}$$

Each simulation begins with a model cell that rests on the surface and has formed a very

small initial contact footprint (area $< 0.1 \ \mu m^2$). The shape of the upper, free part of the cell at this

time is a spherical cap. We determine the fluid velocities and pressure distribution at each time

step by solving a perturbed form of the Stokes equations using the finite element method (FEM;

Fig 1E,F) as described previously [29]. We then use the computed velocities to evolve the cell

shape over a small time step. Details of our model implementation and solution method are given

in Methods and S6 Appendices.

This generic modeling framework of cell spreading can easily be adopted to represent

different mechanistic notions by adjusting the relative contributions of the adhesion and

protrusion stresses. Disregarding the protrusion stress altogether results in a mathematical

representation of the Brownian zipper model described in the Introduction. On the other hand,

the protrusive zipper model generally includes both the protrusion stress as well as the adhesion

stress. A purely protrusion-driven mode of spreading also can be tested by setting the adhesion

stress to zero, although one needs to bear in mind that irreversible adhesion still maintains the

cell-substrate contact region in this case. In all model versions, the adhesion stress can be varied

over several orders of magnitude to predict how the spreading behavior depends on the density

of ligand displayed on the substrate.

## Quantitative experimental results for the validation of model versions

The computational framework presented in the previous section can be adopted to simulate interactions by various cell types with targets of different axisymmetric geometries. Our present focus on human neutrophils spreading on flat substrates is motivated by a wealth of experimental results obtained for this particular scenario and presented in a companion paper [19]. In brief, we exposed cells to glass coverslips coated with IgG antibodies at densities ($\rho_{IgG}$) spanning ~3 orders of magnitude. The analysis of these highly controlled experiments included measurements of the cell-substrate contact area of spreading cells over time (Fig 2). Regardless of IgG density, we found that the area-versus-time curves typically were sigmoidal in shape (Fig 2B,C). Defining a cell's "spreading speed" as the fastest rate of contact-area growth (i.e., the slope of a sigmoidal fit to the area-versus-time curve at the inflection point), we found that all spreading cells increased their substrate-contact area at speeds of around 3 $\mu m^2$/s irrespective of the density of deposited IgG (Fig. 2D). The measured values of the maximum contact area of spreading cells, on the other hand, exhibited a moderate but significant increase at higher IgG densities (Fig 2E). These experimental results provide useful quantitative benchmarks for comparison with computer simulations and will be used in the following sections to evaluate the validity of different mechanistic notions about cell spreading.

**Fig 2. Summary of experimental results.**

(A) The contact region of a spreading neutrophil was imaged using reflection interference

contrast microscopy (RICM). Contact regions in this example are traced in yellow. The scale bar

in the first image denotes 10 μm. (B) Contact area is plotted as a function of time for the

spreading cell shown in (A). The spreading speed is defined as the maximum slope extracted

from a sigmoidal fit, and the maximum contact area is given by averaging the contact area at the

plateau. (C) Mean contact area is computed at discrete time values for aligned curves of cells

spreading on different densities of IgG. Mean quantified IgG densities are reported in the figure

legend as IgG molecules per $μm^2$. Error bars represent standard error of the mean. (D) Average

spreading speed was quantified for the different IgG densities. There is no significant difference

between mean slope values. (E) Maximum contact area increases slightly as a function of IgG

density. Statistically significant differences in maximum contact area values were verified using

ANOVA followed by a Tukey post hoc test. Error bars in both (D) and (E) indicate standard

deviation. The IgG density was quantified for each condition as described in [19].

An additional measure of interest can be inferred from the actual surface densities of IgG

used in the experiments. The most densely coated coverslips presented about 25,000 IgG

molecules per $\mu m^2$ and induced ~85% of deposited cells to spread. Remarkably, even at a low

IgG surface density of ~44 molecules per $\mu m^2$, about 30% of the deposited cells still spread in an

IgG-dependent manner. Assuming that all IgG molecules in the cell-substrate contact region

have formed bonds with the cell's Fcγ receptors, and using an average equilibrium constant of

$10^{-6}$ M for these receptor-ligand bonds [30], conservative (upper-limit) estimates of the adhesion

energy density of cell-substrate interactions on substrates coated with the high and low numbers

of IgG yield 1,500 $\mu J/m^2$ and 2.6 $\mu J/m^2$, respectively.

**Purely adhesion-driven spreading: predictions by the Brownian**

**zipper model**

We first tested the simplest version of our framework, the Brownian zipper model, by

simulating purely adhesion-driven spreading in the absence of any protrusion stress (Fig 3, S1

Movie). In this scenario, the model cell is passive and is pulled onto the substrate by short-range

attraction. This case is conceptually equivalent to a viscous droplet spreading on an adhesive

substrate.

**Fig 3. Predictions of the Brownian zipper model.**

(A) A model cell spreading on the highest tested ligand density (100%) quickly approaches a

spherical cap morphology. Time stamps are included for each snapshot. (B) Curves of contact

area vs. time show that the spreading rate decreases monotonically. The contact area ultimately

approaches steady-state values predicted by the Young-Dupre equation (dashed lines). Predicted

equilibrium shapes are included on the right. (C) Log-log plot of contact area vs. time is nearly

linear over the initial spreading phase. The slope of this line corresponds to the exponent of a

power law describing contact area growth as a function of time. The dashed line shows a slope of

0.5 for reference.

Varying the ligand density $\rho_l$ over two orders of magnitude, we found that the model
cells generally spread fastest immediately after first contacting the substrate. Afterwards, the
growth rate of the cell-substrate contact area decreased monotonically until the contact area
reached a steady-state plateau marking equilibrium (Fig 3B). As expected, the equilibrium
shapes were spherical caps. Log-log plots of the contact-area-versus-time predictions during the
initial, fast spreading phase were approximately linear, with slopes close to 0.5 (Fig 3C). Hence,
the contact-area growth initially obeys a power law, in agreement with various laws proposed to
describe droplet wetting dynamics [31] or passive cell spreading [8, 13]. In fact, the exponent of
0.5 is an excellent match to results of a previous study that modeled cells as highly viscous
droplets [13]. In S6 Appendices, Appendix B, we show how this exponent naturally arises during
early spreading of a passive model cell driven by adhesion stresses concentrated at the contact
perimeter.

Although the predictions of the Brownian zipper model agree well with the typical
theoretical behavior expected for this kind of passive model cell, they clearly differ from the
results of our phagocytic spreading experiments (Fig 2) in several important ways. First, the
simulated contact-area-versus-time curves lack the characteristic sigmoidal shapes observed in
experiments. Second, the predicted spreading curves exhibit a very strong dependence on the
density of encountered ligands, in stark contrast to the observed spreading behavior. This
dependence is evident in both the predicted spreading speeds, which increase dramatically with
ligand density, as well as the predicted maximum contact areas, which increase several-fold as a
function of the tested adhesive strengths.

312       Finally, the equilibrium shapes obtained for the adhesion energy densities used in the

313    simulations are inconsistent with the cell deformations observed on the experimentally tested

314    IgG densities. At equilibrium, the adhesion energy density ($\gamma$) of the Brownian zipper model cell

315    is related to the cortical tension $\tau$ and the contact angle $\theta_c$ via the Young-Dupre equation (Fig

316    4):

317
$$\gamma = \tau(1 - \cos\theta_c) \tag{10}$$

For a given value of the maximum cell-substrate contact area $A_c$, the equilibrium shape of the

model cell—a spherical cap—is uniquely defined by $A_c$ and the known, constant cell volume.

Thus, basic geometry allows us to calculate the contact angle $\theta_c$ as well as the total cell surface

area $A_{cell}$. Inserting the value of the latter into Eq (1) provides the cortical tension at equilibrium,

which then can be used in Eq (10) to obtain the adhesion energy density as a function of $A_c$ (Fig

4). As expected, the relationship between maximum contact area and adhesion strength predicted

by our numerical simulations agrees well with the Young-Dupre equation (Fig 3B, Fig S2 in S6

Appendices). The quantitative comparison of this theoretical prediction to actual adhesion energy

densities—estimated from the densities of deposited IgG molecules as explained earlier—

exposes entirely different behaviors between the model and experiments, however (Fig 4).

Especially in experiments performed at lower IgG densities, the conservatively estimated

adhesion energy densities were much lower than what would be required to form the measured

maximum contact areas solely by adhesion. In summary, the results of this section rule out

adhesion as the primary driving force of frustrated phagocytic spreading.

**Fig 4. Relationship between ligand density and maximum contact area, as predicted by the**

**Young-Dupre equation.**

The adhesion energy $\gamma$, and therefore the ligand density $\rho_{IgG}$, can be expressed as a function of

the contact area of the final spherical cap formed by a viscous droplet. The tested adhesion

strengths are marked as red dots. Actual experimental values are shown for comparison. Error

bars indicate standard deviation.

# Protrusion-dominated spreading: predictions of the protrusive

# zipper model

The protrusive zipper model incorporates the protrusion stress into the basic framework

considered in the previous section, enabling us to account for active force generation by the

model cell. Our semiempirical implementation is motivated by biological plausibility, and by the

goal to reproduce experimental observations with a minimum set of rules for the spatial

distribution (Eq (8)) and the deformation-dependent evolution of the protrusion stress during cell

spreading. To capture the presumed correlation between receptor-induced signaling and

protrusive force generation by the cytoskeleton, we express the strength of the protrusion stress

as a function of the cell-substrate contact area $A_c$. We assume that in the initial low-tension

regime, the protrusion stress grows linearly with $A_c$. Once the cortical tension increases more

steeply, the protrusion stress is assumed to grow faster as well. Ultimately, the strength of the

protrusion stress levels out at a maximum value of $\sigma_{prot,max} = 3.5$ kPa, within the range of

experimentally measured stresses exerted by parallel actin filaments [32, 33]. This behavior is

implemented in our model through a smoothed version of the following rule:

$$\sigma_{prot} = \sigma_{prot,max} \times \begin{cases} \left[0.05 + 0.3\left(\dfrac{A_c}{A_{c,trans}}\right)\right] & \text{if} & \dfrac{A_{cell}}{A_{cell,0}} \leq 1.26 \\[3mm] \left[0.35 + 0.65\left(\dfrac{A_c - A_{c,trans}}{120\,\mu m^2}\right)^{1/2}\right] & \text{if} & \dfrac{A_{cell}}{A_{cell,0}} > 1.26 \text{ and } A_c \leq A_{c,trans} + 120\,\mu m^2 \quad (11) \\[3mm] 1 & \text{if} & A_c > A_{c,trans} + 120\,\mu m^2 \end{cases}$$

where $A_{c,trans}$ is the contact area when the total cell surface area $A_{cell}$ is equal to $1.26 A_{cell,0}$ , the

transitional moment when the cortical tension starts rising more rapidly (Eq (1)).

We started examining the behavior of this model by first disregarding the adhesion stress

altogether. In this scenario, cell spreading is driven solely by protrusion, and the only role of

adhesion is to render cell-substrate contacts irreversible. We found that in this case, the model

cell begins spreading slowly while a distinctive leading edge gradually develops (Fig 5A, S2

Movie), resulting in a morphology that clearly differs from the spherical caps predicted by the

Brownian zipper model (Fig 3). Spreading slows noticeably at about 90 s, around the time the

protrusion stress reaches its maximum value. The resulting contact-area-versus-time curve has a

sigmoidal shape (Figure 5B,C). Plots of the protrusion stress and the cortical tension as a

function of time illustrate the qualitatively similar evolution of these two quantities, with the rise

of the tension lagging due to the initial "slack" of the cell surface in the low-tension regime (Eq

(1)).

**Fig 5. Predictions of the protrusive zipper model.**

(A) For purely protrusion-driven spreading, lamellar pseudopods form as the protrusion stress

increases over time. Time stamps are shown for the sample images. (B) Contact area increases

sigmoidally over time for the protrusive zipper model. The slope of the sigmoidal fit here is 3.0

$\mu m^2/s$ at the inflection point, which matches the average spreading speed observed

experimentally. (C) Protrusion stress (governed by Eq (11)) and cortical tension (governed by Eq

(1)) increase during cell spreading. (D) Varying the parameter $s_0$ determines the thickness of the

pseudopods, with skinny protrusions for $s_0 = 0.4$ µm (top) and coordinated global cell

deformation at $s_0 = 2.0$ µm (bottom); all sample profiles are shown at t = 120 s. For the

protrusive zipper simulations shown in all other cases, we used $s_0 = 0.8$ µm.

We chose the value $s_0 = 0.8$ µm for the characteristic length that defines the spatial range

of the protrusion stress (Eq (8)) in the simulations shown in Fig 5A-C. Because $s_0$ does not

appear to be a directly measurable quantity, it is instructive to inspect how its value affects the

spreading model cell. Snapshots from simulations using different values of $s_0$ demonstrate how

the cell morphology (here: at 120 s) depends on the spatial range over which the protrusion stress

acts (Fig 5D, S3 Movie). Unsurprisingly, the leading edge of the cell assumes the shape of a thin

lamella for small values of $s_0$ and becomes increasingly thicker at larger $s_0$-values. We note

though that aside from this varying thickness of cellular protrusions, the overall spreading

behavior is similar across different values of $s_0$.

We next repeated the simulation with the same settings as used for Fig 5 but now also

included the adhesion stress. We again examined the same set of adhesion strengths as used in

our tests of the Brownian zipper model (Fig 3). The simulations expose how, in this scenario,

cell-substrate adhesion promotes spreading, affecting the overall spreading behavior in a

quantitative manner but without altering it qualitatively. The model cells spread somewhat faster

and farther on substrates coated with higher ligand densities (Fig 6A), with slightly larger contact

angles between their leading edge and the substrate (Fig 6B, S4 Movie). However, the sigmoidal

character of the contact-area-versus-time curves is conserved through most of the tested adhesion

strengths, and both the spreading speeds as well as the maximum contact areas vary much less

than in the predictions of the Brownian zipper model (Fig 3).

**Fig 6. Predictions of the protrusive zipper model with varying adhesion stress.**

(A) Contact area increases faster and reaches higher values for higher ligand densities. The

protrusion-only curve is shown for reference as a dashed line. (B) The leading edge of model

cells spreading on $\rho_l = 0\%$ and $\rho_l = 100\%$ differ visibly at the sampled time points. On higher

ligand density, both the extent of spreading and the dynamic contact angle increase.

Clearly, the predictions of the protrusive zipper model are in much better agreement with

experimental observations than those of the Brownian zipper. A spreading cell morphology that

is characterized by a forward-bulging leading lamella (Figs. 5-6) matches the cross-sectional

profile of phagocytes engulfing large particles well [24, 34], in contrast to model cell shapes

whose leading edge coincides with the perimeter of contact between the cell and substrate (Fig

3). The timing of spreading and sigmoidal shapes of the contact-area-versus-time curves

predicted by the protrusive zipper model (Fig 5-6) agree with experiments as well, and the time-

dependent increase in protrusion stress is consistent with a biphasic signaling response measured

during FcγR-mediated phagocytosis [35]. Furthermore, both the predicted spreading speeds as

well as the maximum contact areas are comparable to the respective values measured in

experiments, and the slight increase of the maximum contact area at higher adhesion strengths is

consistent with our measurements (Fig 6A). On the other hand, the predicted increase of the

spreading speeds on higher ligand densities (Fig 6A) differs from our experimental data (Fig 2).

Therefore, this model still somewhat overestimates the contribution of the adhesion stress to the

dynamics of phagocytic spreading. Nevertheless, of the two considered models only the

protrusive zipper model succeeds in capturing the essential dynamics of experimentally observed

phagocytic spreading, establishing that active cellular protrusion rather than passive cell-

substrate adhesion is the dominating driving force of this type of cellular motion.

## 427  Protrusive zipper model with discrete adhesion sites: best match

## 428  between experiments and theory

The previous sections presented continuum models of cell spreading, resulting in

predictions that overestimated the role of cell-substrate adhesion even in the case where short-

range adhesion plays only a secondary role. In reality, however, adhesive ligands encountered by

a spreading cell reside at discrete substrate locations rather than presenting a contiguous adhesive

film. A pioneering study that considered discrete adhesion sites not only predicted dramatic

qualitative differences compared to continuum models of adhesion but even concluded that,

depending on the lateral spacing of discrete cell-substrate cross-bridges, "there is little or no

tendency for the contact to spread unless the surfaces are forced together" [36]. These

considerations prompted us to design and examine a version of the protrusive zipper model in

which the model cell can only form new attachments to the substrate at prescribed locations (Fig

7, S5 Movie).

**Fig 7. Predictions of the protrusive zipper model with discrete adhesion sites.**

(A) Two time series (times shown at the left) of simulation snapshots illustrate how the

progression of the model cell's leading edge is affected by the spacing of discrete adhesion sites.

Unoccupied binding sites are depicted by empty circles, whereas filled circles indicate that the

cell membrane is locally attached. (B) Contact area vs. time curves share similar slopes over

three orders of magnitude of ligand density, but plateau at different maximum contact areas. (C)

The ligand-density-dependent increase in maximum contact area agrees well with experimental

results obtained in frustrated phagocytosis experiments. Error bars indicate standard deviation.

In this model version, the spacing of adhesion sites is determined by the ligand density $\rho_l$

(the number of ligand molecules per $\mu m^2$), with the in-plane spacing given by $(\rho_l)^{-1/2}$.

Furthermore, we introduce two additional rules to capture realistic aspects of cell spreading on

discrete adhesion sites. First, we replace the continuous adhesion stress (Eq (6)) with the

following concept of discrete bond formation. We consider a new bond as "formed" as soon as

the distance between the next free ligand site and the closest point of the advancing cell

membrane reaches a threshold distance $d_{thresh}$. As a consequence of such bonding, we place the

cell surface into irreversible contact with this substrate site when carrying out computations at

the next time step. The value of the threshold distance is determined by the magnitude of thermal

membrane fluctuations. Assuming that the time scale of membrane fluctuations is generally

smaller than our time step, we define $d_{thresh}$ to be the root mean square height $<h^2>^{1/2}$ of

membrane fluctuations at equilibrium through [37]:

462
$$d_{thresh} = \sqrt{\langle h^2 \rangle} = \sqrt{\frac{k_B T}{4\pi\tau} \ln\left(1 + \frac{\tau A_{cell}}{\pi^2 \kappa_b}\right)} \tag{12}$$

where $k_B$ is Boltzmann's constant, $T$ is temperature, $A_{cell}$ is the cell surface area, and $\kappa_b$ is

the membrane bending modulus. According to this rule, increased cortical tension $\tau$ suppresses

membrane fluctuations, making it more difficult for the cell to form new adhesive connections at

later time points.

Second, because it seems unlikely that persistent receptor-ligand contacts induce

signaling indefinitely, we also assume that the protrusion stress can diminish over time. This

assumption is supported by experimental data showing that actin polymerization does not

continue indefinitely during phagocytosis if the cell encounters an unopsonized region of the

target particle [18, 38, 39]. We implement this behavior by postulating a transient protrusion

stress that initially is triggered by fresh cell-substrate contact (occurring at time $t_{bind}$) and then

decays exponentially according to the relation:

474
$$\tilde{\sigma}_{prot} = \sigma_{prot} e^{-\frac{t - t_{bind}}{t_0}} \qquad\qquad (13)$$

In this version of the model, $\sigma_{prot}$ in Eq (8) is replaced by $\tilde{\sigma}_{prot}$ defined in Eq (13).

To avoid possible bias due to the exact placement of discrete binding sites, we performed

each simulation multiple times, shifting all binding sites laterally by small increments without

changing their spacing. We then averaged the results of these individual simulations to predict

the cell-spreading behavior as a function of the chosen spacing—or density—of adhesion sites.

Overall, the predictions of this discrete protrusive zipper model (Fig 7) agree well with

those of the continuum model version. For instance, maximum cell-substrate contact areas

exhibit a similar dependence on the ligand density in both models. However, in this discrete

model, the sigmoidal contact-area-versus-time curves are aligned much more closely, reflecting

spreading speeds that hardly depend on the ligand density anymore (Fig 7B).

The behavior of model cells spreading on discrete adhesion sites agrees amazingly well

with that of real human neutrophils during frustrated phagocytosis. As in experiments, the

spreading speed remains remarkably consistent over a range of ligand densities spanning 3

orders of magnitude. Moreover, the moderate increase of the maximum contact area at higher

ligand densities successfully reproduces experimental observations (Fig 7C). These results

reinforce the conclusion that active cytoskeletal protrusion drives phagocytic spreading. They

also highlight that it can be important for realistic models of cell spreading on substrates coated

with specific adhesion molecules to account for the discrete distribution of these adhesion sites.

## Discussion

Quantitative understanding of the behavior of motile mammalian cells—from immune

cells to metastasizing cancer cells—is a prerequisite for transformative biomedical advances in

diagnosis, treatment, and therapy of many current and future health threats. Motile cells integrate

a complex variety of physical and chemical cues to carry out specific functions. Genuine

understanding of the underlying mechano-chemical processes not only requires an

interdisciplinary strategy but also integrative experimental/theoretical approaches. Translation of

mechanistic notions into mathematical models, and comparison of theoretical predictions with

experimental observations, often provides the strongest arguments for or against the validity of

biological hypotheses. Confidence in the predictive power of computer simulations requires that

the model assumptions be realistic and biologically plausible. Guided by this perspective, we

here examined fundamental biophysical mechanisms that potentially can facilitate spreading of

biological cells on a substrate.

Unpolarized cells usually begin spreading in a symmetric fashion [8, 40]. This stage is

often viewed as a passive process that is entirely driven by adhesion energy due to receptor-

ligand binding [8, 41]. An alternative class of models considers such spreading as dependent on

cytoskeletal activity [10, 14, 15]. In this study, we have developed a mechanical modeling

framework that allows us to establish whether frustrated phagocytosis, a specific form of

isotropic cell spreading, is primarily driven by adhesion (Brownian zipper model) or by

protrusive forces generated by the cell (protrusive zipper model). Our findings likely carry over

to other types of cell spreading inasmuch as key characteristics of adhesion and protrusion are

shared [42].

The incompatibility of our Brownian zipper simulations with experimental data

invalidates models of phagocytosis that treat neutrophil spreading as an adhesion-driven wetting

phenomenon [43]. Moreover, our findings invite a careful review of mathematical models of

other types of cell spreading that consider adhesion as the main driving force [8, 13]. A first

important prerequisite for the validity of such models is the use of realistic values of basic

mechanical parameters, such as the effective viscosity of the cell interior, the cortical tension,

and the adhesion energy density. For example, if the equilibrium cell-substrate contact areas

measured in our spreading experiments on low-to-moderate ligand densities were due to

adhesion, then the involved adhesion energies would have to be several orders of magnitude

higher than those estimated from our measurements of ligand density. Even receptor

accumulation in the contact region could not make up for this difference, because the smaller

number of available ligands limits the number of possible adhesion bonds. Furthermore, the

ability to simulate not only equilibrium situations but also the dynamical cell behavior, and to

leverage time-dependent experimental observations against the predictions of such models,

considerably raises confidence in a model's evaluation. As demonstrated in this study, such

comparison greatly benefits from the availability of experimental data obtained on substrates

presenting a large range of known densities of adhesive ligands.

It appears that adhesion-driven spreading is largely limited to situations where adhesion

is unusually strong, and where the cells—like red blood cells—or surrogate objects—like lipid

vesicles—are characterized by a low-viscosity interior and very low tension of their cortex or

membrane. Indeed, a recent mechanical modeling study found that Brownian membrane

fluctuations were only sufficient to drive very small amounts of cell spreading on a surface [44].

If there are cases in which motile mammalian cells spread predominantly due to adhesion, then

such spreading will likely occur on much longer timescales than considered here. Finally, it may

be possible that adhesion plays a more relevant role in the engulfment of very small particles

whose uptake does not require a significant expansion of the cell surface area. Irrespective of the

relevance of Brownian-zipper-type models in a particular practical situation, we note that the

excellent agreement of our theoretical treatment of purely adhesion-driven spreading with other

studies focusing on this form of motion [8, 13] strengthens confidence in our modeling approach

and its numerical implementation.

The only alternative mechanism by which cells can expand their region of contact with a

suitable substrate appears to be active, protrusion-dominated spreading. Many imaging studies

indeed have documented an increased density of F-actin at the protruding front of motile immune

cells [16, 45, 46], supporting the biological plausibility of the protrusive zipper concept.

Moreover, inhibition of actin polymerization with cytochalasin B or latrunculin A eliminates the

rapid phase of fibroblast spreading [10, 14, 47]. Similarly, chemical disruption of the actin

cytoskeleton generally inhibits phagocytosis by immune cells [28, 38]. We have previously

shown that extremely small concentrations of actin inhibitors actually can increase the size of

transient, chemotactic-like protrusions [28, 48]. However, the cortical tension was reduced in

such cases, and phagocytic spreading was not enhanced, highlighting the multiplicity of—likely

dichotomous—mechanical roles of F-actin during cell deformations.

Our simulations based on the protrusive zipper model not only reproduce the main

features of neutrophil spreading on IgG-coated substrates but, in the case of discrete adhesion

sites, they also agree very well with experimental data. We note that this agreement neither is

surprising, nor does it constitute solid proof of the validity of this model. The scarcity of

quantitative insights into the underlying mechanisms simply does not place sufficient constraints

on the design of computational models. Instead, our goal here was to explore what set of

minimalistic rules would need to be postulated to achieve a good match between theory and

experiment. The successful implementation of this approach allows us to propose likely

mechanistic details that can then provide guidance for future studies. Still, we are confident that

the protrusive zipper model paints a reasonably accurate picture of phagocytic spreading, not

only because of the elimination of the only alternative hypothesis, but also because our

semiempirical model is plausible, i.e., it does not contradict pertinent biophysical and biological

insights. For example, our implementation of the relationship between receptor-induced

signaling and the generation of protrusive force is consistent with the hitherto most successful

model of phagocytosis [24], as well as with previous measurements of the biphasic signaling

response from a macrophage-like cell line during Fcγ receptor-mediated phagocytosis [35].

Importantly, this implementation is independent of the exact mechanism by which intracellular

processes are translated into protrusive force; instead, it encompasses different mechanistic

notions such as the Brownian ratchet [49] and end-tracking motors [50].

An important outcome of our simulations of the protrusive zipper model is the difference

in predictions depending on whether we considered continuous or discrete cell-substrate

adhesion, in agreement with a previous theoretical study [36]. We are unaware of physiological

situations in which cell adhesion is not predominantly supported by specific receptor-ligand

bonds. Thus, the model version that is based on discrete binding sites is the more realistic version

of the protrusive zipper. Remarkably, it is also the version that yielded the best match between

theory and experimental observations. To achieve this agreement, we postulated that the

formation of new receptor-ligand bonds is coupled to the protrusion stress in a time-dependent

manner (Eq (13)), based on the assumption that receptor-induced signaling and subsequential

actin polymerization are not sustained indefinitely after a ligand activates the receptor. There are

multiple factors that could lead to this effective decay, such as the depletion of precursor

molecules needed for the production of signaling messengers (e.g. release of $IP_3$ will cease once

$PIP_2$ has been locally depleted [51]).

Perhaps the most important insight from the comparison of model versions based on

continuous versus discrete adhesion is that models based on continuous adhesion are likely to

misrepresent key aspects of the cell behavior, especially on lower ligand densities. For example,

such models cannot capture the possibility that too large ligand spacing may prevent the next

unbound receptor-ligand pair from approaching one another to within the distance at which they

are likely to form a bond. In our protrusive zipper model with discrete adhesion, variation of

substrate adhesivity by changing the density of adhesion sites primarily affects the maximum

cell-substrate contact area. This prediction agrees well with our experiments, unlike the

predictions by model versions that treat the ligand layer as a continuum.

The mechanisms governing isotropic cell spreading studied here are likely to also play a

crucial role in the directional motion of migrating cells. Many immune cells, for example, are

capable of executing both types of motion, as required during phagocytic spreading versus

chemotactic and other forms of migration. As revealed by our experiments and reproduced by

our discrete adhesion model, the rate of isotropic cell spreading is largely conserved on

substrates presenting a broad range of densities of adhesion sites. This finding exposes an

interesting difference to the prediction by models of migrating cells, i.e., that the crawling speed

of such cells has a maximum on substrates with intermediate adhesion strengths [52-54]. This

difference is a reminder that directional migration generally is a more complex type of motion

that involves, in addition to protrusion and adhesion, important mechanisms regulating

contraction and detachment of the rear of the migrating cell from the substrate. The positive

correlation between low-to-moderate ligand densities and the speed of motion predicted by

models of migrating cells can be attributed to either a direct scaling of protrusion with the

amount of adhesion [53], or a postulated increased actin retrograde flow at lower adhesion

strengths [52]. The predicted slowing of migrating cells at higher ligand densities is caused by an

effective frictional force due to the need to break stronger adhesive attachments at the cell's rear

[55, 56]. A noteworthy recent model of cell migration that decoupled friction from adhesive

attraction actually predicted faster movement for higher adhesion energies [41]. The excellent

agreement between our experiments and the protrusive zipper model suggests that the above

additional mechanisms included in models of migrating cells play at most a minor role in

isotropic phagocytic spreading. On the other hand, we believe that the main insights exposed by

our study provide important suggestions on how to treat the interplay between protrusion and

adhesion in models of migrating cells.

# Methods

## The perturbed Stokes equations

Assuming the cell interior can be approximated by a single, highly viscous, slow-moving Newtonian fluid, its motion obeys the Stokes equations for creeping flow with no body force, as given by Eq (2). Furthermore, for an incompressible fluid, the equation of continuity simplifies to the incompressibility condition (Eq (3)). Due to axial symmetry, there are three unknowns: $v_r$, $v_z$, and $p$, where $r$ and $z$ are cylindrical polar coordinates.

To solve this system of partial differential equations, we use the finite element method (FEM). Because directly solving the Stokes equations using FEM leads to numerical instabilities [57], we follow the procedure outlined by Drury and Dembo [29] to solve a perturbed form of the Stokes equations, in which the incompressibility condition (Eq (3)) is approximated as

$$\nabla \cdot (\mathbf{v} - \epsilon \nabla p) = 0 \tag{14}$$

In this form, the fluid is nearly incompressible, which allows us to avoid the well-known problems with using FEM for incompressible problems [57]. To accurately solve the Stokes equations, the value for $\epsilon$ must be sufficiently small. Specifically, we use the condition given in [29]:

$$\epsilon \leq \frac{h_{mesh}^2}{\mu} \tag{15}$$

where $h_{mesh}$ is the characteristic radius of a single quadrilateral element. We tested different values of $\epsilon$ in our models and found this condition to provide sufficient accuracy and convergence (Fig S5 in S6 Appendices).

Over the unbound portion of the cell, we specify Neumann boundary conditions,

corresponding to a stress balance (Eq (4)). For the adherent cell surface, we specify Dirichlet

boundary conditions, simply a no-slip condition in this case (Eq (9)). The pressure boundary

condition is given by assuming the normal component of the pressure gradient goes to zero at the

cell surface:

$$\nabla p \cdot \mathbf{n} = 0 \qquad (16)$$

## Computational details

At each time step, we first compute values for the surface curvature of the cell. Local

values of the angle of the normal $\phi$ are evaluated using cubic polynomial fits. We then

calculate the boundary stresses using the relationships outlined in the main text. The integral for

the effective adhesion stress (Eq (6)) is computed numerically as described in S6 Appendices.

The values for tension and protrusion stress at each time step are evaluated using smoothed

versions of the relationships given in Eq (1) and Eq (11) (see S6 Appendices for full forms).

To find the fluid velocity and pressure fields, we must simultaneously solve Eq (2) and

Eq (14) for $v_r$, $v_z$, and $p$. We employ the strategy outlined in [29], which is an adaptation of the

Uzawa method. Starting with an estimate for pressure $p_{est}$ given either by the previous time step

or as a uniform distribution prior to spreading, we solve for fluid velocities by treating the

pressure as known in Eq (2):

$$\mu \nabla^2 \mathbf{v} = \nabla p_{est} \qquad (17)$$

We then treat the fluid velocities as known ($\mathbf{v}_{est}$) and solve an altered version of equation (14):

$$\frac{1}{\mu}(p - p_{est}) = \nabla \cdot \mathbf{v}_{est} - \nabla \cdot (\epsilon \nabla p) \qquad (18)$$

The updated pressure is then inserted back into Eq (17) and the procedure is repeated until the

following condition is met:

665
$$\frac{\max\left(\left|p - p_{est}\right|\right)}{\max\left(\left|p - p_{out}\right|\right)} \leq 10^{-6} \tag{19}$$

666    This generally occurs in less than 50 iterations.

667    The new cell boundary is then given by

668
$$\begin{aligned} r(t+\Delta t) &= r(t) + v_r \Delta t \\ z(t+\Delta t) &= z(t) + v_z \Delta t \end{aligned} \tag{20}$$

The value for $\Delta t$ was set relative to the Courant number for each iteration, such that fluid flow

occurs over at most 10% of any individual element (S6 Appendices). In the continuum adhesion

case (no discrete adhesion sites), the contact line is updated at each time step by computing the

intersection between the cell boundary and the surface, then adding all points advected to $z \leq 0$

to the Dirichlet boundary condition (contact, no-slip) for the next time step. In the discrete

adhesion model, contact is determined by the distance threshold computed from Eq (12). The

contour is smoothed after each advection, and then small corrections are made to the cell contour

to compensate for any deviations from constant volume due to smoothing and numerical error.

This model was implemented in MATLAB R2020b (MathWorks) and the full code is

available at https://github.com/emmetfrancis/phagocyticSpreading. Individual simulations

generally run within 1-3 hours on a personal computer.

## Acknowledgements

We thank Dr. Robert Guy from UC Davis Mathematics for many helpful conversations on various iterations of our model and for assistance with the derivation found in S6 Appendices, Appendix B. We also thank Dr. Marc Herant for fruitful discussions on computational modeling of motile cells.

## References

1.  Henry SJ, Crocker JC, Hammer DA. Ligand density elicits a phenotypic switch in human neutrophils. Integr Biol (Camb). 2014;6(3):348-56. Epub 2014/02/01. doi: 10.1039/c3ib40225h. PubMed PMID: 24480897; PubMed Central PMCID: PMCPMC5850933.

2.  Hoang AN, Jones CN, Dimisko L, Hamza B, Martel J, Kojic N, et al. Measuring neutrophil speed and directionality during chemotaxis, directly from a droplet of whole blood. Technology (Singap World Sci). 2013;1(1):49. Epub 2014/05/09. doi: 10.1142/S2339547813500040. PubMed PMID: 24809064; PubMed Central PMCID: PMCPMC4010229.

3.  Ascione F, Vasaturo A, Caserta S, D'Esposito V, Formisano P, Guido S. Comparison between fibroblast wound healing and cell random migration assays in vitro. Exp Cell Res. 2016;347(1):123-32. doi: 10.1016/j.yexcr.2016.07.015. PubMed PMID: WOS:000383217500013.

701     4.      Czirok A. Endothelial cell motility, coordination and pattern formation during

702             vasculogenesis. Wiley Interdiscip Rev Syst Biol Med. 2013;5(5):587-602. Epub

703             2013/07/17. doi: 10.1002/wsbm.1233. PubMed PMID: 23857825; PubMed Central

704             PMCID: PMCPMC3737767.

705     5.      Schneider L, Cammer M, Lehman J, Nielsen SK, Guerra CF, Veland IR, et al.

706             Directional cell migration and chemotaxis in wound healing response to PDGF-AA are

707             coordinated by the primary cilium in fibroblasts. Cell Physiol Biochem. 2010;25(2-

708             3):279-92. Epub 2010/01/30. doi: 10.1159/000276562. PubMed PMID: 20110689;

709             PubMed Central PMCID: PMCPMC2924811.

710     6.      Morrissey MA, Williamson AP, Steinbach AM, Roberts EW, Kern N, Headley MB, et al.

711             Chimeric antigen receptors that trigger phagocytosis. Elife. 2018;7. doi: ARTN e36688

712     10.7554/eLife.36688. PubMed PMID: WOS:000435593500001.

713     7.      Feng MY, Jiang W, Kim BYS, Zhang CC, Fu YX, Weissman IL. Phagocytosis

714             checkpoints as new targets for cancer immunotherapy. Nat Rev Cancer. 2019;19(10):568-

715             86. doi: 10.1038/s41568-019-0183-z. PubMed PMID: WOS:000492743400012.

716     8.      Cuvelier D, Thery M, Chu YS, Dufour S, Thiery JP, Bornens M, et al. The universal

717             dynamics of cell spreading. Curr Biol. 2007;17(8):694-9. Epub 2007/03/24. doi:

718             10.1016/j.cub.2007.02.058. PubMed PMID: 17379524.

719     9.      Dubin-Thaler BJ, Giannone G, Dobereiner HG, Sheetz MP. Nanometer analysis of cell

720             spreading on matrix-coated surfaces reveals two distinct cell states and STEPs. Biophys

721             J. 2004;86(3):1794-806. Epub 2004/03/03. doi: 10.1016/S0006-3495(04)74246-0.

722             PubMed PMID: 14990505; PubMed Central PMCID: PMCPMC1304013.

10.   Xiong Y, Rangamani P, Fardin MA, Lipshtat A, Dubin-Thaler B, Rossier O, et al.

Mechanisms controlling cell size and shape during isotropic cell spreading. Biophys J.

2010;98(10):2136-46. Epub 2010/05/21. doi: 10.1016/j.bpj.2010.01.059. PubMed PMID:

20483321; PubMed Central PMCID: PMCPMC2872297.

11.   Reinhart-King CA, Dembo M, Hammer DA. The dynamics and mechanics of endothelial

cell spreading. Biophys J. 2005;89(1):676-89. Epub 2005/04/26. doi:

10.1529/biophysj.104.054320. PubMed PMID: 15849250; PubMed Central PMCID:

PMCPMC1366566.

12.   Carter SB. Haptotaxis and the mechanism of cell motility. Nature. 1967;213(5073):256-

60. Epub 1967/01/21. doi: 10.1038/213256a0. PubMed PMID: 6030602.

13.   Etienne J, Duperray A. Initial dynamics of cell spreading are governed by dissipation in

the actin cortex. Biophys J. 2011;101(3):611-21. Epub 2011/08/03. doi:

10.1016/j.bpj.2011.06.030. PubMed PMID: 21806929; PubMed Central PMCID:

PMCPMC3145312.

14.   Fardin MA, Rossier OM, Rangamani P, Avigan PD, Gauthier NC, Vonnegut W, et al.

Cell spreading as a hydrodynamic process. Soft Matter. 2010;6(19):4788-99. doi:

10.1039/c0sm00252f. PubMed PMID: WOS:000282008300022.

15.   Herant M, Dembo M. An integrative toy model of cell flattening, spreading, and ruffling.

Biorheology. 2015;52(5-6):405-14. Epub 2015/11/26. doi: 10.3233/BIR-14042. PubMed

PMID: 26600264.

16.   Kovari DT, Wei W, Chang P, Toro JS, Beach RF, Chambers D, et al. Frustrated

Phagocytic Spreading of J774A-1 Macrophages Ends in Myosin II-Dependent

Contraction. Biophys J. 2016;111(12):2698-710. Epub 2016/12/22. doi:

10.1016/j.bpj.2016.11.009. PubMed PMID: 28002746; PubMed Central PMCID:

PMCPMC5194617.

17.    Cannon GJ, Swanson JA. The macrophage capacity for phagocytosis. J Cell Sci.

1992;101 ( Pt 4):907-13. Epub 1992/04/01. PubMed PMID: 1527185.

18.    Heinrich V. Controlled One-on-One Encounters between Immune Cells and Microbes

Reveal Mechanisms of Phagocytosis. Biophys J. 2015;109(3):469-76. Epub 2015/08/06.

doi: 10.1016/j.bpj.2015.06.042. PubMed PMID: 26244729; PubMed Central PMCID:

PMCPMC4572503.

19.    Francis EA, Xiao H, Teng LH, Heinrich V. Mechanisms of frustrated phagocytic

spreading of human neutrophils on antibody-coated surfaces. 2022;

BIORXIV/2022/481104.

20.    Evans E, Yeung A. Apparent viscosity and cortical tension of blood granulocytes

determined by micropipet aspiration. Biophys J. 1989;56(1):151-60. Epub 1989/07/01.

doi: 10.1016/S0006-3495(89)82660-8. PubMed PMID: 2752085; PubMed Central

PMCID: PMCPMC1280460.

21.    Tran-Son-Tay R, Needham D, Yeung A, Hochmuth RM. Time-dependent recovery of

passive neutrophils after large deformation. Biophys J. 1991;60(4):856-66. Epub

1991/10/01. doi: 10.1016/S0006-3495(91)82119-1. PubMed PMID: 1742456; PubMed

Central PMCID: PMCPMC1260136.

22.    Yeung A, Evans E. Cortical shell-liquid core model for passive flow of liquid-like

spherical cells into micropipets. Biophys J. 1989;56(1):139-49. Epub 1989/07/01. doi:

10.1016/S0006-3495(89)82659-1. PubMed PMID: 2752083; PubMed Central PMCID:

PMCPMC1280459.

23.    Herant M, Marganski WA, Dembo M. The mechanics of neutrophils: synthetic modeling

770           of three experiments. Biophys J. 2003;84(5):3389-413. Epub 2003/04/30. doi:

771           10.1016/s0006-3495(03)70062-9. PubMed PMID: 12719267; PubMed Central PMCID:

772           PMCPMC1302898.

773    24.    Herant M, Heinrich V, Dembo M. Mechanics of neutrophil phagocytosis: experiments

774           and quantitative models. J Cell Sci. 2006;119(Pt 9):1903-13. Epub 2006/04/26. doi:

775           10.1242/jcs.02876. PubMed PMID: 16636075.

776    25.    Herant M, Lee CY, Dembo M, Heinrich V. Protrusive push versus enveloping embrace:

777           computational model of phagocytosis predicts key regulatory role of cytoskeletal

778           membrane anchors. PLoS Comput Biol. 2011;7(1):e1001068. doi:

779           10.1371/journal.pcbi.1001068. PubMed PMID: 21298079; PubMed Central PMCID:

780           PMCPMC3029235.

781    26.    Zak A, Merino-Cortes SV, Sadoun A, Mustapha F, Babataheri A, Dogniaux S, et al.

782           Rapid viscoelastic changes are a hallmark of early leukocyte activation. Biophys J. 2021.

783           Epub 2021/03/18. doi: 10.1016/j.bpj.2021.02.042. PubMed PMID: 33730552.

784    27.    Herant M, Heinrich V, Dembo M. Mechanics of neutrophil phagocytosis: behavior of the

785           cortical tension. J Cell Sci. 2005;118(Pt 9):1789-97. Epub 2005/04/14. doi:

786           10.1242/jcs.02275. PubMed PMID: 15827090.

787    28.    Lee CY, Herant M, Heinrich V. Target-specific mechanics of phagocytosis: protrusive

788           neutrophil response to zymosan differs from the uptake of antibody-tagged pathogens. J

789           Cell Sci. 2011;124(Pt 7):1106-14. Epub 2011/03/10. doi: 10.1242/jcs.078592. PubMed

790           PMID: 21385838; PubMed Central PMCID: PMCPMC3056606.

791    29.    Drury JL, Dembo M. Hydrodynamics of micropipette aspiration. Biophys J. 1999;76(1 Pt

792           1):110-28. Epub 1999/01/06. doi: 10.1016/S0006-3495(99)77183-3. PubMed PMID:

793           9876128; PubMed Central PMCID: PMCPMC1302505.

794    30.    Maenaka K, van der Merwe PA, Stuart DI, Jones EY, Sondermann P. The human low

795           affinity Fcgamma receptors IIa, IIb, and III bind IgG with fast kinetics and distinct

796           thermodynamic properties. J Biol Chem. 2001;276(48):44898-904. Epub 2001/09/07.

797           doi: 10.1074/jbc.M106819200. PubMed PMID: 11544262.

798    31.    Bonn D, Eggers J, Indekeu J, Meunier J, Rolley E. Wetting and spreading. Rev Mod

799           Phys. 2009;81(2):739-805. doi: 10.1103/RevModPhys.81.739. PubMed PMID:

800           WOS:000267197500008.

801    32.    Marcy Y, Prost J, Carlier MF, Sykes C. Forces generated during actin-based propulsion:

802           a direct measurement by micromanipulation. Proc Natl Acad Sci U S A.

803           2004;101(16):5992-7. Epub 2004/04/14. doi: 10.1073/pnas.0307704101. PubMed PMID:

804           15079054; PubMed Central PMCID: PMCPMC395911.

805    33.    Parekh SH, Chaudhuri O, Theriot JA, Fletcher DA. Loading history determines the

806           velocity of actin-network growth. Nat Cell Biol. 2005;7(12):1219-23. Epub 2005/11/22.

807           doi: 10.1038/ncb1336. PubMed PMID: 16299496.

808    34.    Lee C-Y, Thompson III GR, Hastey CJ, Hodge GC, Lunetta JM, Pappagianis D, et al.

809           *Coccidioides* Endospores and Spherules Draw Strong Chemotactic, Adhesive, and

810           Phagocytic Responses by Individual Human Neutrophils. PLoS ONE.

811           2015;10(6):e0129522. doi: 10.1371/journal.pone.0129522.

812    35.    Zhang Y, Hoppe AD, Swanson JA. Coordination of Fc receptor signaling regulates

813           cellular commitment to phagocytosis. Proc Natl Acad Sci U S A. 2010;107(45):19332-7.

Epub 2010/10/27. doi: 10.1073/pnas.1008248107. PubMed PMID: 20974965; PubMed

Central PMCID: PMCPMC2984174.

36.    Evans EA. Detailed mechanics of membrane-membrane adhesion and separation. II.

Discrete kinetically trapped molecular cross-bridges. Biophys J. 1985;48(1):185-92.

Epub 1985/07/01. doi: 10.1016/S0006-3495(85)83771-1. PubMed PMID: 4016208;

PubMed Central PMCID: PMCPMC1329388.

37.    Helfrich W, Servuss RM. Undulations, Steric Interaction and Cohesion of Fluid

Membranes. Nuovo Cimento D. 1984;3(1):137-51. doi: Doi 10.1007/Bf02452208.

PubMed PMID: WOS:A1984SF44800012.

38.    van Zon JS, Tzircotis G, Caron E, Howard M. A mechanical bottleneck explains the

variation in cup growth during FcgammaR phagocytosis. Mol Syst Biol. 2009;5:298.

Epub 2009/08/20. doi: 10.1038/msb.2009.59. PubMed PMID: 19690567; PubMed

Central PMCID: PMCPMC2736656.

39.    Griffin FM, Jr., Griffin JA, Leider JE, Silverstein SC. Studies on the mechanism of

phagocytosis. I. Requirements for circumferential attachment of particle-bound ligands to

specific receptors on the macrophage plasma membrane. J Exp Med. 1975;142(5):1263-

82. Epub 1975/11/11. doi: 10.1084/jem.142.5.1263. PubMed PMID: 1194852; PubMed

Central PMCID: PMCPMC2189973.

40.    McGrath JL. Cell spreading: the power to simplify. Curr Biol. 2007;17(10):R357-8. Epub

2007/05/16. doi: 10.1016/j.cub.2007.03.057. PubMed PMID: 17502086.

41.    Cao Y, Karmakar R, Ghabache E, Gutierrez E, Zhao Y, Groisman A, et al. Cell motility

dependence on adhesive wetting. Soft Matter. 2019;15(9):2043-50. Epub 2019/02/07.

doi: 10.1039/c8sm01832d. PubMed PMID: 30724956.

42.    Cougoule C, Wiedemann A, Lim J, Caron E. Phagocytosis, an alternative model system

for the study of cell adhesion. Semin Cell Dev Biol. 2004;15(6):679-89. Epub

2004/11/25. doi: 10.1016/j.semcdb.2004.09.001. PubMed PMID: 15561587.

43.    Tollis S, Dart AE, Tzircotis G, Endres RG. The zipper mechanism in phagocytosis:

energetic requirements and variability in phagocytic cup shape. BMC Syst Biol.

2010;4:149. Epub 2010/11/10. doi: 10.1186/1752-0509-4-149. PubMed PMID:

21059234; PubMed Central PMCID: PMCPMC2991294.

44.    DiNapoli KT, Robinson DN, Iglesias PA. A mesoscale mechanical model of cellular

interactions. Biophys J. 2021;120(22):4905-17. Epub 2021/10/24. doi:

10.1016/j.bpj.2021.10.021. PubMed PMID: 34687718; PubMed Central PMCID:

PMCPMC8633826.

45.    Nelsen E, Hobson CM, Kern ME, Hsiao JP, O'Brien Iii ET, Watanabe T, et al. Combined

Atomic Force Microscope and Volumetric Light Sheet System for Correlative Force and

Fluorescence Mechanobiology Studies. Sci Rep. 2020;10(1):8133. Epub 2020/05/20. doi:

10.1038/s41598-020-65205-8. PubMed PMID: 32424215; PubMed Central PMCID:

PMCPMC7234992.

46.    Fritz-Laylin LK, Riel-Mehan M, Chen BC, Lord SJ, Goddard TD, Ferrin TE, et al. Actin-

based protrusions of migrating neutrophils are intrinsically lamellar and facilitate

direction changes. Elife. 2017;6. Epub 2017/09/28. doi: 10.7554/eLife.26990. PubMed

PMID: 28948912; PubMed Central PMCID: PMCPMC5614560.

47.    Dubin-Thaler BJ, Hofman JM, Cai Y, Xenias H, Spielman I, Shneidman AV, et al.

Quantification of cell edge velocities and traction forces reveals distinct motility modules

during cell spreading. PLoS One. 2008;3(11):e3735. Epub 2008/11/18. doi:

10.1371/journal.pone.0003735. PubMed PMID: 19011687; PubMed Central PMCID:

PMCPMC2581916.

48.     Heinrich V, Lee C-Y. Blurred line between chemotactic chase and phagocytic

consumption: an immunophysical single-cell perspective. Journal of Cell Science.

2011;124(18):3041-51. doi: 10.1242/jcs.086413.

49.     Mogilner A, Oster G. Cell motility driven by actin polymerization. Biophys J.

1996;71(6):3030-45. Epub 1996/12/01. doi: 10.1016/S0006-3495(96)79496-1. PubMed

PMID: 8968574; PubMed Central PMCID: PMCPMC1233792.

50.     Dickinson RB, Caro L, Purich DL. Force generation by cytoskeletal filament end-

tracking proteins. Biophys J. 2004;87(4):2838-54. Epub 2004/09/30. doi:

10.1529/biophysj.104.045211. PubMed PMID: 15454475; PubMed Central PMCID:

PMCPMC1304702.

51.     Flannagan RS, Jaumouille V, Grinstein S. The cell biology of phagocytosis. Annu Rev

Pathol. 2012;7:61-98. Epub 2011/09/14. doi: 10.1146/annurev-pathol-011811-132445.

PubMed PMID: 21910624.

52.     Barnhart EL, Lee KC, Keren K, Mogilner A, Theriot JA. An adhesion-dependent switch

between mechanisms that determine motile cell shape. PLoS Biol. 2011;9(5):e1001059.

Epub 2011/05/12. doi: 10.1371/journal.pbio.1001059. PubMed PMID: 21559321;

PubMed Central PMCID: PMCPMC3086868.

53.     Ziebert F, Aranson IS. Effects of adhesion dynamics and substrate compliance on the

shape and motility of crawling cells. PLoS One. 2013;8(5):e64511. Epub 2013/06/07.

doi: 10.1371/journal.pone.0064511. PubMed PMID: 23741334; PubMed Central

PMCID: PMCPMC3669322.

54.    Copos CA, Walcott S, Del Alamo JC, Bastounis E, Mogilner A, Guy RD.

884           Mechanosensitive Adhesion Explains Stepping Motility in Amoeboid Cells. Biophys J.

885           2017;112(12):2672-82. Epub 2017/06/22. doi: 10.1016/j.bpj.2017.04.033. PubMed

886           PMID: 28636923; PubMed Central PMCID: PMCPMC5478966.

887    55.    Shao D, Levine H, Rappel WJ. Coupling actin flow, adhesion, and morphology in a

888           computational cell motility model. Proc Natl Acad Sci U S A. 2012;109(18):6851-6.

889           Epub 2012/04/12. doi: 10.1073/pnas.1203252109. PubMed PMID: 22493219; PubMed

890           Central PMCID: PMCPMC3344950.

891    56.    Kuusela E, Alt W. Continuum model of cell adhesion and migration. J Math Biol.

892           2009;58(1-2):135-61. Epub 2008/05/20. doi: 10.1007/s00285-008-0179-x. PubMed

893           PMID: 18488227.

894    57.    Hughes TJR. The Finite Element Method: Linear Static and Dynamic Finite Element

895           Analysis: Dover; 2000.

# Supporting Information

**S1 Movie. Simulations of purely adhesive cell spreading ("Brownian zipper") on substrates**

**presenting different densities of adhesion sites.** This sequence of animations shows side views

of model cells spreading on adhesive substrates coated with relative ligand densities ($\rho_l$) of 1%,

3%, 10%, 30%, and 100% for 400 s in the absence of protrusion stress.

**S2 Movie. Simulation of purely protrusion-driven cell spreading.** This animation shows a

side view of a model cell spreading on a flat substrate for 200 s. The cell motion is assumed to be

driven exclusively by active cytoskeletal protrusion at a maximum protrusion stress of $\sigma_{prot,max} =$

3.5 kPa. Cell-substrate contact, once established, is assumed to be permanent. All other

contributions of adhesive cell-substrate interactions to spreading are neglected.

**S3 Movie. Effect of the range of action of the protrusion stress on purely protrusion-driven**

**cell spreading.** This sequence of simulations shows side views of model cells spreading in the

absence of adhesion stress for 200 s. Increasing the range $s_0$ over which cytoskeletal protrusion

actively pushes the cellular pseudopod outwards (given in terms of the arc length of the free cell

contour, measured from the point of substrate contact) results in thicker pseudopods. In this

sequence, $s_0 = 0.4$ μm, 1.2 μm, and 2.0 μm, respectively. In all examples, $\sigma_{prot,max} \times s_0 = 2.8$

mN/m.

**S4 Movie. Minor role of cell-substrate adhesion stress in protrusion-dominated cell**

**spreading ("protrusive zipper").** This sequence of simulations compares magnified views of

purely protrusion-driven spreading (adhesion stress $\rho_l = 0\%$) with protrusion-driven spreading in

the presence of moderate and strong adhesion stress ($\rho_l = 10\%$ and 100%, respectively). In all

examples, $\sigma_{prot,max} = 3.5$ kPa.

**S5 Movie. Protrusive zipper simulations with discrete adhesion sites.** This sequence of

simulations shows a magnified view of protrusion-driven spreading ($\sigma_{prot,max} = 3.5$ kPa) on

substrates presenting different densities of discrete adhesion sites ($\rho_l = 30$ μm$^{-2}$, 100 μm$^{-2}$, 1,000

μm$^{-2}$, and 10,000 μm$^{-2}$). Open circles denote free adhesive binding sites, and filled circles depict

sites with engaged ligand.

**S6 Appendices. Derivation of important relationships and summary of numerical testing.**

(A) Adhesion stress calculation. (B) Derivation of a power law for contact area growth in passive

spreading. (C) Development of $C^1$ versions of tension and protrusion stress relations. (D) Details

of finite element implementation. (E) Summary of numerical testing.

**S7 Table. Summary of modeling parameters, with supporting references.**

# S6 Appendices

## Appendix A. Adhesion stress calculation

To compute the adhesion stress, we consider a potential acting between an adhesion molecule and a receptor in the cell membrane. Here, we use a 6-3 potential:

$$u_{adh}(D) = u_0 \left[ \left( \frac{D_0}{D} \right)^6 - 2 \left( \frac{D_0}{D} \right)^3 \right] \tag{S1}$$

where $D_0$ is the distance associated with the energy minimum and $u_0$ sets the magnitude of the binding energy. The force due to this single interaction is

$$f_{adh}(D) = -\frac{du_{adh}(D)}{dD} = f_0 \left[ \left( \frac{D_0}{D} \right)^7 - \left( \frac{D_0}{D} \right)^4 \right] \tag{S2}$$

where $f_0 = 6u_0/D_0$. As $D_0$ is the location of the energy minimum, we also see that $f_{adh}(D_0) = 0$. We use $D_0 = 50$ nm in our simulations. Realistically, one might expect this distance to be much smaller (molecular scale), but the longer range here effectively includes membrane fluctuations and filopodia, and effectively serves as an upper limit. Adhesion potentials with a range of tens to hundreds of nanometers are common when modeling cells and vesicles at the mesoscopic scale [1, 2].

To solve for the overall adhesion stress, we must integrate over the ligand-coated surface (surface coordinates $[r, \theta]$ ) and the region of membrane being considered (surface coordinates $[s, \psi]$ ) (Fig S1A).

**Fig S1. Adhesion force calculation.**
(A) The cell geometry is depicted from the side, as well as from the top, including definitions of coordinates used in the text. (B) Adhesion stress components are plotted as a function of arc length $s$ for the cell shape shown in part A given a relative ligand density of 10%.

The individual interaction forces must be specified as vectors, that is

$$\mathbf{f}_{single} = f_0 \left[ \left( \frac{D_0}{D} \right)^7 - \left( \frac{D_0}{D} \right)^4 \right] \mathbf{f}_{unit} \tag{S3}$$

The distance between a point on the membrane and a point on the surface is given by

$$D(r, \theta, s, \psi) = \sqrt{(r_m \cos \psi - r \cos \theta)^2 + (r_m \sin \psi - r \sin \theta)^2 + z_m^2} \tag{S4}$$

Note that the distance $z_m$ is adjusted from the $z$ coordinate used elsewhere so that the surface $z = 0$ corresponds to the zero-force distance $D_0$, i.e. $z_m = z + D_0$.

The direction of the force is expressed by a unit vector (we only calculate the $r$ and $z$ components, as the angular component cancels out due to symmetry)

$$\mathbf{f}_{unit} = \frac{1}{D} \begin{bmatrix} r_m \cos \psi - r \cos \theta \\ z_m \end{bmatrix} \tag{S5}$$

Considering a region of the cell membrane extending from $s = s_{cur} - l$ to $s = s_{cur} + l$ along the arc length and from $\psi = -\Delta\psi/2$ to $\psi = \Delta\psi/2$ in the angular direction, we compute the pairwise force by the total integral

$$\mathbf{f}_{adh} = \int_{-\Delta\psi/2}^{\Delta\psi/2} \int_{s_{cur}-\frac{l}{2}}^{s_{cur}+\frac{l}{2}} \left( \int_0^{2\pi} \int_{r_c}^{\infty} \mathbf{f}_{single} \rho_l r \, dr \, d\theta \right) \rho_R r_m \, ds \, d\psi \tag{S6}$$

where $\rho_l$ is the ligand density on the substrate and $\rho_R$ is the receptor density on the cell membrane. In practice, we lump $f_0$ and $\rho_R$ into a single parameter $\sigma_0 = f_0 \rho_R$, which effectively serves as a force per unit area of cell membrane.

The stress per unit area of membrane is given by

$$\boldsymbol{\sigma}_{adh} = \frac{\mathbf{f}_{adh}}{\int_{-\Delta\psi/2}^{\Delta\psi/2} \int_{s_{cur}-\frac{l}{2}}^{s_{cur}+\frac{l}{2}} r_m \, ds \, d\psi} \tag{S7}$$

In the limit $l\Delta\psi \to 0$, we obtain the final expression, as given in the main text

$$\boldsymbol{\sigma}_{adh} = \sigma_0 \rho_l \int_0^{2\pi} \int_{r_{edge}}^{\infty} \mathbf{f}_{unit} \left[ \left( \frac{D_0}{D} \right)^7 - \left( \frac{D_0}{D} \right)^4 \right] r \, dr \, d\theta \tag{S8}$$

Due to the embedded trigonometric functions, this integral is not readily carried out analytically unless we use a small angle estimate for $\theta$, which will not hold in general. Therefore, we evaluate it numerically in MATLAB by performing trapezoidal integration twice, using log-spaced meshes for both $r$ and $\theta$. This achieves essentially the same results as integration with MATLAB built-in functions, but computation time is reduced well over 10-fold.

An example of the adhesion stress components computed as a function of arc length $s$ along the cell contour shown in Fig S1A for $\rho_l = 10\%$ is shown in Fig S1B.

In practice, we modify the adhesion strength in simulations by varying the multiplicative factor $\sigma_0\rho_l$, but we cannot directly extract the ligand density $\rho_l$. However, as described in the main text, we can infer the adhesion energy per unit area $\gamma$ from the equilibrium contact area $A_c$ using simple geometry, conservation of volume, and the Young-Dupre equation. Using these estimated values of adhesion energy $\gamma(A_c)$ and the typical binding energy of a low affinity Fcγ receptor to IgG ($E_{bind} \approx 6 \times 10^{-20}$ J [3]), we can estimate the corresponding IgG density $\rho_{IgG,est}$

$$\rho_{IgG,est} = \frac{\gamma(A_c)}{E_{bind}} \tag{S9}$$

We perform a linear fit between $\rho_{IgG,est}$ and the relative adhesion strength $\rho_l / \rho_{l,max}$ to compute the effective IgG densities tested in our Brownian Zipper model (Fig S2B). From this fit, we find that $\rho_l = 10\%$ roughly corresponds to $\rho_{IgG,est} = 1{,}000$ µm$^{-2}$.

**Fig S2. Relationship between estimated IgG density and model adhesion strength**.
A linear fit relates model adhesion strength to IgG density based on the equilibrium contact areas we obtain from simulations.

## Appendix B. Derivation of a power law for contact area growth in passive spreading

To develop an approximate analytical prediction of the functional dependence of the contact area on time, we start from the overall energy balance for our passive model cell spreading on a flat surface with adhesion energy density $\gamma$, directly analogous to the energy balance in the spreading of a viscous droplet [4]. After canceling a factor of $2\pi$, this yields

$$\frac{\gamma}{2}\frac{\partial}{\partial t}(r_c^2) = 2\mu \iint e_{ij}e_{ij}r\mathrm{d}r\mathrm{d}z + \frac{\partial}{\partial t}\left(\tau \int r\mathrm{d}s\right) + \frac{\partial}{\partial t}\left(\iint pr\mathrm{d}r\mathrm{d}z\right) \qquad (S10)$$

where $r_c$ is the cell contact radius, $\mu$ is the effective cell viscosity, $e_{ij}$ are the components of the rate-of-strain tensor, $\tau$ is the cortical tension, and $p$ is the pressure inside the cell. The left-hand side corresponds to energy gain due to cell-surface adhesion. The first term on the right-hand side is the viscous dissipation rate [5], the second term is the work required to deform the cortex, and the third is the change in energy due to changes in pressure and/or volume. In our simulations volume is kept constant, and any energy required for pressure changes is much lower than for changes of the surface area or tension; therefore, we neglect the last term of Eq S10 in the following analysis.

To estimate the remaining terms in the energy balance, we assume that the passively spreading cell adopts the shape of a spherical cap with contact radius $r_c$, radius of curvature $R_{cell}$, and a constant volume (Fig S3A). This geometry defines the contact angle as:

$$\sin \theta_c = \frac{r_c}{R_{cell}} \qquad (S11)$$

We focus this analysis on initial contact area growth, during which the contact angle is small and we can treat $R_{cell}$ as a constant. Based on this geometry and features of the calculated flow profile (Fig S3A inset), we estimate the viscous dissipation rate and work required to deform the cortex as follows.

We first expand the dissipation term to include all non-zero components of the rate-of-strain tensor (all derivatives with respect to the polar angle are zero given axial symmetry):

$$\Pi = \mu \iint \left[2\left(\frac{\partial v_r}{\partial r}\right)^2 + 2\left(\frac{v_r}{r}\right)^2 + 2\left(\frac{\partial v_z}{\partial z}\right)^2 + \left(\frac{\partial v_r}{\partial z} + \frac{\partial v_z}{\partial r}\right)^2\right] r\mathrm{d}r\mathrm{d}z \qquad (S12)$$

The main effect of the adhesion force is to "pull down" the membrane onto the flat surface; that is, the z-component of the adhesion stress generally dominates (Fig 1B). Accordingly, we expect $v_z$ to dominate over $v_r$ in the portion of the cell closest to the substrate, which we do in fact see in our calculations (Fig S3A inset). Furthermore, due to incompressibility, we know

$$\frac{v_r}{r} + \frac{\partial v_r}{\partial r} + \frac{\partial v_z}{\partial z} = 0 \qquad (S13)$$

Therefore, if the contributions of $v_r$ are small, $\partial v_z/\partial z$ must also be small, and we are left with:

$$\Pi \approx \mu \iint \left(\frac{\partial v_z}{\partial r}\right)^2 r\,dr\,dz \qquad (S14)$$

As a test geometry for points near the surface, we consider a wedge shape, as shown in Fig S3B. The exact functional form of $v_z$ within this wedge cannot be readily inferred *a priori*. Instead, we identify characteristic length scales to estimate the velocity gradient. We expect the $z$-velocity to be highest close to the surface, that is, within the length scale set by the adhesion potential ($D_0$). For this region, the associated distance in the $r$ direction $\Delta r$ (Fig S3B) is given by

$$\Delta r = \frac{D_0}{\tan \theta_c} \approx \frac{D_0 R_{cell}}{r_c} \qquad (S15)$$

**Fig S3.** Energy balance calculations.
(A) Geometry of a spreading droplet during early spreading. The inset illustrates fluid velocities (arrows) at the boundary near the contact line as predicted by a typical Brownian zipper simulation. (B) Simplified geometry of a wedge used to represent a spreading droplet. The top panel defines variable used in the text. The bottom panel illustrates the manner of contact area growth implemented in our model. (C) Local dissipation per unit volume at the contact line for two early time points as computed by a Brownian zipper simulation at a ligand density of 10%. The dissipation is tightly confined to the point of contact. (D) Total viscous dissipation rate was directly computed using linear shape functions and plotted against $r_c^2 \times dr_c/dt$ on a log-log axis. Simulated values were binned and averaged for each interval on the x-axis. Error bars indicate standard deviation of binned data.

Before estimating the velocity gradient, we relate $v_z$ to the rate of contact area growth, $dr_c/dt$. Because there is no slip at the contact line itself, the only way for the contact area to increase is for nonadherent membrane to approach and touch the surface past the contact line. We consider a sample point close to the surface at which the membrane moves at maximum velocity $v_{max}$ in the negative $z$ direction (Fig S3B). If this point moves down from a distance $\delta z$ from the surface, then the contact line moves outward by a distance of $\delta r$ in the $r$-direction (Fig S3B), given by

$$\delta r = \frac{\delta z}{\tan \theta_c} \tag{S16}$$

In terms of velocities, this yields

$$\frac{dr_c}{dt} = \frac{v_{max}}{\tan \theta_c} \approx \frac{v_{max} R_{cell}}{r_c} \tag{S17}$$

Combining Eqs S15 and S17 and assuming the velocity gradient is highest in the direction tangent to the membrane, we approximate the sought velocity gradient:

$$\frac{\partial v_z}{\partial r} \approx \frac{v_{max}}{\Delta r} = \frac{r_c^2}{R_{cell}^2 D_0} \frac{dr_c}{dt} \tag{S18}$$

We insert these estimates into the dissipation term (Eq S14) and integrate over the wedge:

$$\Pi \approx \mu \left( \frac{r_c^2}{R_{cell}^2 r_0} \frac{dr_c}{dt} \right)^2 \int_{r_c}^{r_c + \Delta r} \int_{D_0 \frac{(r-r_c)}{\Delta r}}^{D_0} r\, dz\, dr \tag{S19}$$

$$= \mu \left( \frac{r_c^2}{R_{cell}^2 D_0} \frac{dr_c}{dt} \right)^2 \frac{D_0 \Delta r}{6} (3r_c + \Delta r)$$

Inserting the definition of $\Delta r$ from Eq S15 and assuming $\Delta r \ll 3r_c$, we find

$$\Pi \approx \Pi_0 r_c^4 \left( \frac{dr_c}{dt} \right)^2 \tag{S20}$$

where $\Pi_0 = \mu/(2R_{cell}^3)$. We verify the validity of this relationship by comparing it to numerically computed dissipation rates obtained in Brownian zipper simulations (Fig S3C-D). As expected, the dissipation rate per unit volume is highest in a small region near the contact line (Fig S3C). We integrate to compute the total dissipation rate at each time step, and then plot this as a function of $r_c^2(dr_c/dt)$ to compare it to our approximate analytical prediction from Eq S20 (Fig S3D). When plotted on a log-log axis, this curve shows a linear region with a slope of 2, in agreement with the scaling in Eq S20.

We next estimate the work required to deform the cortex, as given by the second term on the right-hand side of Eq S10. For a spherical cap of contact radius $r_c$ and height $h$:

$$\frac{\partial}{\partial t}\left(\tau \int r \, ds\right) = \frac{\partial}{\partial t}\left[\tau\left(r_c^2 + \frac{h^2}{2}\right)\right] \tag{S21}$$

Early in spreading, $h$ is approximately constant and tension changes slowly, therefore

$$\frac{\partial}{\partial t}\left[\tau\left(r_c^2 + \frac{h^2}{2}\right)\right] \approx 2\tau r_c \frac{dr_c}{dt} \tag{S22}$$

Inserting the approximations from Eq S20 and Eq S22 into the energy balance (Eq S10) and simplifying, we arrive at the following differential equation:

$$r_c^3 \left(\frac{dr_c}{dt}\right) = \frac{2(\gamma - \tau)}{\Pi_0} \tag{S23}$$

In early spreading, tension changes more slowly than $r_c$; therefore, it can be treated as a constant. This leads to the spreading law observed in our model as well as in a previous computational model of passive cell spreading [2]

$$r_c \propto t^{\frac{1}{4}} \rightarrow A_c \propto t^{\frac{1}{2}} \tag{S24}$$

## Appendix C. Development of C$^1$ versions of tension and protrusion stress relations

Our numerical calculations require versions of the piecewise relationships for tension (Eq 1) and protrusion stress (Eq 11) that have continuous first derivatives. We have developed C$^1$ versions of these relations by using polynomials to transition between each piecewise domain. For tension, we use

$$\tau(A_{cell}) = \begin{cases} 0.01 + 0.16\left(\dfrac{A_{cell}}{A_{cell,0}} - 1\right)\dfrac{\text{mN}}{\text{m}} & \text{for } \dfrac{A_{cell}}{A_{cell,0}} \leq 1.13 \\[2ex] 0.7405\left(\dfrac{A_{cell}}{A_{cell,0}} - 1\right)^2 - 0.0325\left(\dfrac{A_{cell}}{A_{cell,0}} - 1\right) + 0.0225 & \text{for } 1.13 < \dfrac{A_{cell}}{A_{cell,0}} \leq 1.39 \\[2ex] 0.0516 + 0.545\left(\dfrac{A_{cell}}{A_{cell,0}} - 1.26\right)\dfrac{\text{mN}}{\text{m}} & \text{for } \dfrac{A_{cell}}{A_{cell,0}} > 1.39 \end{cases} \tag{S25}$$

The C$^1$ version of protrusion stress requires two separate transition regions, which we achieve by using cubic polynomials:

$$\sigma_{prot} = \sigma_{prot,max} \begin{cases} \left[0.05 + 0.3\left(\dfrac{A_c}{A_{c,trans}}\right)\right] & \text{for } \dfrac{A_{cell}}{A_{cell,0}} \leq 1.13 \\[2ex] c_3 A_c^3 + c_2 A_c^2 + c_1 A_c + c_0 & \text{for } 1.13 < \dfrac{A_{cell}}{A_{cell,0}} \leq 1.39 \\[2ex] \left[0.35 + 0.65\left(\dfrac{A_c - A_{c,trans}}{120}\right)^{\frac{1}{2}}\right] & \text{for } \dfrac{A_{cell}}{A_{cell,0}} > 1.39 \ \& \ A_c \leq A_{c,mid} + 100 \\[2ex] d_3 A_c^3 + d_2 A_c^2 + d_1 A_c + d_0 & \text{for } A_{c,trans} + 100 < A_c \leq A_{c,trans} + 140 \\[1ex] 1 & \text{when } A_c > A_{c,trans} + 140 \end{cases} \tag{S26}$$

The $c$ and $d$ coefficients are found by solving the appropriate linear systems. Their values generally depend on the contact area $A_c$ at different values of surface area deformation; namely, at $A_{cell} = 1.13 \, A_{cell,0}$, $A_{cell} = 1.26 \, A_{cell,0}$, and $A_{cell} = 1.39 \, A_{cell,0}$. These are derived from sample runs of the simulation.

## Appendix D. Details of finite element implementation

The governing equations are given in Methods, referred to as the "velocity equation" (Eq 17) and the "pressure equation" (Eq 18) below. Both equations are readily cast into a Galerkin form, which then is rewritten in matrix form as is conventional for finite element methods.

For reference, we give the elemental contributions to the stiffness matrices and load vectors for each finite element system, following the notation used by Hughes [6]. $N_a$ is used to denote the finite element shape function associated with node $a$. We use bilinear quadrilateral shape functions and compute all integrals using two-by-two Gauss quadrature.

The velocity equation (Eq 17) gives rise to the linear system

$$\mathbf{K}\,\mathbf{v}_{all} = \mathbf{F} \tag{S27}$$

The vector $\mathbf{v}_{all}$ contains the velocity values $[v_r, v_z]$ at each node, and the global stiffness matrix $\mathbf{K}$ and global load vector $\mathbf{F}$ are assembled from elemental contributions. The contributions from nodes $a$ and $b$ belonging to element $e$ are given as follows (note the factor of $2\pi r$ due to axial symmetry, as well as the term $(N_a N_b / r^2)$ which arises when computing $\nabla \mathbf{v}$ in cylindrical coordinates)

$$K_{arbr,e} = \int \left(2\frac{\partial N_a}{\partial r}\frac{\partial N_b}{\partial r} + 2\frac{N_a N_b}{r^2} + \frac{\partial N_a}{\partial z}\frac{\partial N_b}{\partial z}\right) 2\pi r d\Omega \tag{S28}$$

$$K_{arbz,e} = \int \frac{\partial N_a}{\partial r}\frac{\partial N_b}{\partial z} 2\pi r d\Omega$$

$$K_{azbr,e} = \int \frac{\partial N_a}{\partial z}\frac{\partial N_b}{\partial r} 2\pi r d\Omega$$

$$K_{azbz,e} = \int \left( 2 \frac{\partial N_a}{\partial z} \frac{\partial N_b}{\partial z} + \frac{\partial N_a}{\partial r} \frac{\partial N_b}{\partial r} \right) 2\pi r d\Omega$$

For the load vector

$$F_{ar,e} = \int p_{est} \left( \frac{\partial N_a}{\partial r} + \frac{N_a}{r} \right) 2\pi r d\Omega + \int N_a \sigma_r 2\pi r d\Gamma \tag{S29}$$

$$F_{az,e} = \int p \frac{\partial N_a}{\partial z} 2\pi r d\Omega + \int N_a \sigma_z 2\pi r d\Gamma$$

where the second term in the load vector arises from the Neumann boundary condition (stress balance, where $[\sigma_r, \sigma_z]$ are the boundary stresses given by adding the adhesion stress, protrusion stress, and cortical stress) and only appears for those elements which belong to the free boundary of the cell. For elements in contact with the substrate, there is no contribution from the Dirichlet boundary condition as the enforced velocity is equal to zero.

The pressure equation (Eq 18) gives rise to the linear system

$$\mathbf{Q} \, \mathbf{p}_{all} = \mathbf{G} \tag{S30}$$

The vector $\mathbf{p}_{all}$ contains the pressure value $p$ at each node. The global stiffness matrix $\mathbf{Q}$ and global load vector $\mathbf{G}$ are assembled from elemental contributions given by

$$Q_{ab,e} = \int \left[ N_a N_b - \epsilon \left( \frac{\partial N_a}{\partial r} \frac{\partial N_b}{\partial r} + \frac{\partial N_a}{\partial z} \frac{\partial N_b}{\partial z} \right) \right] 2\pi r d\Omega \tag{S31}$$

$$G_{a,e} = \int N_a \left[ p_{est} - \frac{\partial v_r}{\partial r} - \frac{v_r}{r} - \frac{\partial v_z}{\partial z} \right] 2\pi r d\Omega \tag{S32}$$

The expression for $G_a$ incorporates the assumption that $\nabla p \cdot \mathbf{n}$ is zero along the cell boundary, as dictated by the boundary condition, and $\nabla^2 p = 0$ everywhere, which is satisfied exactly for an incompressible fluid. Derivatives of $[v_r, v_z]$ are computed using bilinear quadrilateral shape functions.

This system of equations is solved iteratively, as described in Methods. Examples of the fluid velocities and pressures obtained by the Brownian zipper model (Fig S4A) and the protrusive zipper model (Fig S4B) are included for illustration.

**Fig S4. Example velocity and pressure fields obtained by finite element computations.**
(A) Fluid velocity vector fields and pressure scalar fields (heat maps) are shown for purely adhesion-driven spreading with a ligand density of 10%. The magnitude of fluid velocities decay relatively quickly over time and the highest pressures are localized to the contact line in early spreading. (B) Fluid velocity vector fields and pressure scalar fields (heat maps) are shown for purely protrusion-driven spreading. The velocities remain of similar magnitude over time due to the time-dependent increase in protrusion stress.

## Appendix E. Numerical testing and optimization

The accuracy of numerical computations such as used in this study can be affected by several factors. In this appendix, we document a number of tests that we have carried out to verify the reliability of our numerical predictions. All tests of the Brownian zipper model are conducted for a ligand density ($\rho_l$) of 10%, and all tests of the protrusive zipper model are carried out in the absence of adhesion stress.

### I. Test for incompressibility

Solving the governing equations for the perturbed Stokes equations given in Methods (Eqs 17 and 18) yields an approximate satisfaction of incompressibility. The degree to which incompressibility is enforced depends on the magnitude of the perturbation parameter $\epsilon$ in Eq 18. Specifically, as $\epsilon$ goes to zero, the fluid becomes more incompressible. However, if $\epsilon$ is too small, there are numerical difficulties. Thus, it is worthwhile to evaluate how closely incompressibility is satisfied for different values of $\epsilon$. In our model, $\epsilon$ scales with the characteristic radius of an individual element $h_{mesh}$ and the effective viscosity $\mu$ [7]. As described in Eq 15 of the main text, we generally require that $\epsilon \leq h_{mesh}^2/\mu$ , but we can write more generally

$$\epsilon = \bar{\epsilon}\left(\frac{h_{mesh}^2}{\mu}\right) \tag{S33}$$

where $\bar{\epsilon}$ is a dimensionless scaling factor which should have a magnitude less than or equal to one to satisfy the above inequality. Here, we assess how this parameter affects the error in our model.

We quantify the degree of compressibility at different locations in the cell body in terms of the local rate of dilatation, obtained by integrating $\nabla \cdot \mathbf{v}$ over individual elements and dividing by the element volume $V_{el}$. That is,

$$\text{Dilatation rate } = \frac{\iiint (\nabla \cdot \mathbf{v}) dV}{V_{el}} \tag{S34}$$

This dilatation rate has units of inverse seconds and describes the rate of fluid expansion per unit volume. In general, we expect the largest departures from perfect incompressibility to occur in the immediate vicinity of the perimeter of the contact region, due to a well-known singularity that has been examined thoroughly for different cases of droplet spreading [4]. Focusing on this region, we have characterized how our choice of the value of $\epsilon$ (Eq S33) affects the dilatation rate (Fig. S5A). We evaluate the overall error of the model in terms of the mean square error (MSE) for the incompressibility condition

$$\text{MSE} = \frac{\iiint (\nabla \cdot \mathbf{v})^2 dV}{V_{tot}} \tag{S35}$$

The MSE initially depends only weakly on the value of $\epsilon$, but increases considerably at higher values, as expected (Fig S5B-C). Generally, a value of $\bar{\epsilon} = 1$ results in sufficiently low error while also avoiding numerical difficulties associated with lower values of $\bar{\epsilon}$.

**Fig S5. Testing for incompressibility.**
(A) Heat maps of the dilatation rate within the leading edge of the cell for purely protrusion-driven spreading at $t =$ 15s for different values of $\bar{\epsilon}$. (B) Volumetric mean square error is plotted as a function of $\bar{\epsilon}$ for Brownian Zipper simulations at $t = 15$ s. (C) Volumetric mean square error is plotted as a function of $\bar{\epsilon}$ for protrusion-only spreading at 15 s.

## II. Time-step testing

The optimal time step $\Delta t$ for many fluid mechanics simulations depends on the fluid velocity and the mesh resolution. A common reference for this is the Courant number, generally given by

$$C = \frac{v\Delta t}{l} \tag{S36}$$

where $l$ is a characteristic length and $v$ is a characteristic velocity. For numerical stability, $C$ should be much less than one.

We use Eq S36 to determine the appropriate time step for the minimum ratio of $l/v$ across all elements, that is

$$\Delta t = C \min\left(\frac{l}{v}\right) = C \min_{\text{all el}} \left( \min\left[ \frac{r_{span}}{v_{r,c}}, \frac{z_{span}}{v_{z,c}} \right] \right) \tag{S37}$$

where $r_{span}$ is the span of the element in the $r$ direction, $z_{span}$ is the span of the element in the $z$ direction, $[v_{r,c}, v_{z,c}]$ is the velocity vector at the centroid of the element, and $C$ is the desired Courant number. For instance, if $C = 0.1$, fluid flow crosses at most 10% of any individual element during a single time step. We performed a series of simulations using decreasing values of $C$ to establish at which Courant number our computations converge satisfactorily. Figure S6 shows examples of testing different time steps for the Brownian Zipper model (Fig S6A) and the protrusive zipper model with no adhesion stress (Fig S6B). For smaller time steps, the contact area curves become smoother and converge to a single solution. Generally, we find that $C \leq 0.1$ performs well in these cases, which is the value we choose for our simulations.

**Fig S6. Time-step testing.**
Contact area versus time for different Courant numbers for (A) the Brownian zipper model and (B) purely protrusion-driven spreading.

## III. Mesh refinement

We ensured that the results of our simulations do not depend significantly on the mesh resolution as follows. When constructing the mesh, we start by specifying the boundary nodes, choosing tight spacing near the substrate and gradually increasing spacing to 0.17 μm over the rest of the cell contour (Fig S7A). After parameterizing the contour, we use transfinite interpolation to construct the interior mesh ([8], Fig S7A). Using a global edge spacing of 0.17 μm generally results in about 2,000 elements. Here, we focus on refining the mesh spacing close to the surface where the stresses are most concentrated.

For both the Brownian zipper model and the protrusive zipper model, we tested minimum boundary point spacing values $\Delta s_{min}$ of 25 nm, 10 nm, 5 nm, and 2.5 nm. In both cases, we find

convergence for $\Delta s_{min} \leq 5$ nm (Fig S7B-C). For our simulations in the main paper, we use $\Delta s_{min}$ = 2.5 nm.

**Fig S7. Mesh refinement.**
(A) Illustration of mesh construction including our choice of relevant boundary spacing intervals. Convergence of the simulations is evaluated by inspection of contact-area-versus-time curves obtained with different minimum mesh spacings close to the substrate ($\Delta s_{min}$) for (B) the Brownian zipper model and (C) purely protrusion-driven spreading.

# References

1. Sukumaran S, Seifert U. Influence of shear flow on vesicles near a wall: A numerical study. Phys Rev E. 2001;64(1). doi: ARTN 011916 DOI 10.1103/PhysRevE.64.011916. PubMed PMID: WOS:000169907100090.

2. Etienne J, Duperray A. Initial dynamics of cell spreading are governed by dissipation in the actin cortex. Biophys J. 2011;101(3):611-21. Epub 2011/08/03. doi: 10.1016/j.bpj.2011.06.030. PubMed PMID: 21806929; PubMed Central PMCID: PMCPMC3145312.

3. Maenaka K, van der Merwe PA, Stuart DI, Jones EY, Sondermann P. The human low affinity Fcgamma receptors IIa, IIb, and III bind IgG with fast kinetics and distinct thermodynamic properties. J Biol Chem. 2001;276(48):44898-904. Epub 2001/09/07. doi: 10.1074/jbc.M106819200. PubMed PMID: 11544262.

4. Bonn D, Eggers J, Indekeu J, Meunier J, Rolley E. Wetting and spreading. Reviews of Modern Physics. 2009;81(2):739-805. doi: 10.1103/RevModPhys.81.739. PubMed PMID: WOS:000267197500008.

5. Aris R. Vectors, Tensors, and the Basic Equations of Fluid Mechanics: Dover Publications, Inc.; 1962.

6. Hughes TJR. The Finite Element Method: Linear Static and Dynamic Finite Element Analysis: Dover; 2000.

7. Drury JL, Dembo M. Hydrodynamics of micropipette aspiration. Biophys J. 1999;76(1 Pt 1):110-28. Epub 1999/01/06. doi: 10.1016/S0006-3495(99)77183-3. PubMed PMID: 9876128; PubMed Central PMCID: PMCPMC1302505.

8. Knupp PS, S.L. The Fundamentals of Grid Generation. 1st ed: CRC Press; 1993.

# S7 Table. Modeling parameters

| Parameter | Description | Value | Notes |
|---|---|---|---|
| $\tau_0$ | Resting cortical tension | 0.01 mN/m | Matches experimental value used by Herant et al [1] |
| $\tau_{max}$ | Maximum cortical tension | ~1 mN/m | In a reasonable range for high tensions measured during neutrophil phagocytosis [2, 3] |
| $\mu$ | Effective cytoplasmic viscosity | 200 Pa-s (passive) 1,660 Pa-s (active) | Matches experimentally measured values [4, 5] |
| $R_0$ | Initial cell radius | 4.25 μm | Radius of a human neutrophil |
| $\sigma_0$ | Adhesion stress constant (scaling factor) | 370 Pa | Together with the ligand density, this determines the adhesion strength |
| $\rho_{IgG,max}$ | Ligand density corresponding to 100% for Brownian Zipper model | 10,000 IgG/μm$^2$ | This density corresponds to about 600 μJ/m$^2$, which exceeds values derived from other cases of cell spreading [6] |
| $D_0$ | Zero adhesion force distance | 50 nm | Relatively large distance required for mesoscopic model, as standard in other continuum models [7, 8] |
| $\sigma_{prot,max}$ | Max. protrusion stress | 3,500 Pa | Actin filaments growing in parallel can achieve forces above 1 nN per μm$^2$ (> 1 kPa) [9, 10] |
| $s_0$ | Protrusive force range | 0.8 μm | Controls how the protrusion stress decays along the membrane (Eq 8) |
| $t_0$ | Characteristic time for decay of protrusion stress | 66 s | Used for discrete adhesion model (Eq 13) |

# References

1. Herant M, Lee CY, Dembo M, Heinrich V. Protrusive push versus enveloping embrace: computational model of phagocytosis predicts key regulatory role of cytoskeletal membrane anchors. PLoS Comput Biol. 2011;7(1):e1001068. doi: 10.1371/journal.pcbi.1001068. PubMed PMID: 21298079; PubMed Central PMCID: PMCPMC3029235.
2. Lee CY, Thompson GR, 3rd, Hastey CJ, Hodge GC, Lunetta JM, Pappagianis D, et al. Coccidioides Endospores and Spherules Draw Strong Chemotactic, Adhesive, and Phagocytic Responses by Individual Human Neutrophils. PLoS One. 2015;10(6):e0129522. doi: 10.1371/journal.pone.0129522. PubMed PMID: 26070210; PubMed Central PMCID: PMCPMC4466529.
3. Herant M, Heinrich V, Dembo M. Mechanics of neutrophil phagocytosis: behavior of the cortical tension. J Cell Sci. 2005;118(Pt 9):1789-97. doi: 10.1242/jcs.02275. PubMed PMID: 15827090.
4. Evans E, Yeung A. Apparent viscosity and cortical tension of blood granulocytes determined by micropipet aspiration. Biophys J. 1989;56(1):151-60. Epub 1989/07/01. doi: 10.1016/S0006-3495(89)82660-8. PubMed PMID: 2752085; PubMed Central PMCID: PMCPMC1280460.
5. Tran-Son-Tay R, Needham D, Yeung A, Hochmuth RM. Time-dependent recovery of passive neutrophils after large deformation. Biophys J. 1991;60(4):856-66. Epub 1991/10/01. doi: 10.1016/S0006-3495(91)82119-1. PubMed PMID: 1742456; PubMed Central PMCID: PMCPMC1260136.
6. Cuvelier D, Thery M, Chu YS, Dufour S, Thiery JP, Bornens M, et al. The universal dynamics of cell spreading. Curr Biol. 2007;17(8):694-9. Epub 2007/03/24. doi: 10.1016/j.cub.2007.02.058. PubMed PMID: 17379524.
7. Sukumaran S, Seifert U. Influence of shear flow on vesicles near a wall: A numerical study. Phys Rev E. 2001;64(1). doi: ARTN 011916 DOI 10.1103/PhysRevE.64.011916. PubMed PMID: WOS:000169907100090.
8. Etienne J, Duperray A. Initial dynamics of cell spreading are governed by dissipation in the actin cortex. Biophys J. 2011;101(3):611-21. Epub 2011/08/03. doi: 10.1016/j.bpj.2011.06.030. PubMed PMID: 21806929; PubMed Central PMCID: PMCPMC3145312.
9. Parekh SH, Chaudhuri O, Theriot JA, Fletcher DA. Loading history determines the velocity of actin-network growth. Nat Cell Biol. 2005;7(12):1219-23. Epub 2005/11/22. doi: 10.1038/ncb1336. PubMed PMID: 16299496.
10. Marcy Y, Prost J, Carlier MF, Sykes C. Forces generated during actin-based propulsion: a direct measurement by micromanipulation. Proc Natl Acad Sci U S A. 2004;101(16):5992-7. Epub 2004/04/14. doi: 10.1073/pnas.0307704101. PubMed PMID: 15079054; PubMed Central PMCID: PMCPMC395911.