Modelling the synergistic effect of bacteriophage and antibiotics on bacteria: killers and drivers of resistance evolution

Quentin J. Leclerc [1,2,3]*, Jodi A. Lindsay [3], Gwenan M. Knight [1,2]

[1] Centre for Mathematical Modelling of Infectious Diseases, Department of Infectious Disease Epidemiology, Faculty of Epidemiology & Population Health, London School of Hygiene & Tropical Medicine, UK

[2] Antimicrobial Resistance Centre, London School of Hygiene and Tropical Medicine, UK

[3] Institute for Infection & Immunity, St George's University of London, UK

* corresponding author: quentin.leclerc@lshtm.ac.uk (QJL)

ABSTRACT

Bacteriophage (phage) are bacterial predators that can also spread antimicrobial resistance (AMR) genes between bacteria by generalised transduction. Phage are often present alongside antibiotics in the environment, yet evidence of their joint killing effect on bacteria is conflicted, and the dynamics of transduction in such systems are unknown. Here, we combine *in vitro* data and mathematical modelling to identify conditions where phage and antibiotics act in synergy to remove bacteria or drive AMR evolution.

We adapt a published model of phage-bacteria dynamics, including transduction, to add the pharmacodynamics of erythromycin and tetracycline, parameterised from new *in vitro* data. We simulate a system where two strains of *Staphylococcus aureus* are present at stationary phase, each carrying either an erythromycin or tetracycline resistance gene, and where multidrug-resistant bacteria can be generated by transduction only. We determine rates of bacterial clearance and multidrug-resistant bacteria appearance, when either or both antibiotics and phage are present at varying timings and concentrations.

Although phage and antibiotics act in synergy to kill bacteria, by reducing bacterial growth antibiotics reduce phage production. A low concentration of phage introduced shortly after antibiotics fails to replicate and exert a strong killing pressure on bacteria, instead generating multidrug-resistant bacteria by transduction which are then selected for by the antibiotics. Multidrug-resistant bacteria numbers were highest when antibiotics and phage were introduced simultaneously.

The interaction between phage and antibiotics leads to a trade-off between a slower clearing rate of bacteria (if antibiotics are added before phage), and a higher risk of multidrug-resistance evolution (if phage are added before antibiotics), exacerbated by low concentrations of phage or antibiotics. Our results form hypotheses to guide future experimental and clinical work on the impact of phage on

37   AMR evolution, notably for studies of phage therapy which should investigate varying timings and

38   concentrations of phage and antibiotics.

40   AUTHOR SUMMARY

Bacteriophage ("phage") are viruses that can infect and kill bacteria, but also natural drivers of

antimicrobial resistance (AMR) evolution by transduction, when they carry non-phage DNA between

bacteria, including AMR genes. Phage are often present alongside antibiotics in the environment and

in humans, yet the joint dynamics of phage, antibiotics, and bacteria are unclear. Using laboratory

work and mathematical modelling, we show for the first time that depending on timing and

concentration, phage and antibiotics can either work together to kill bacteria faster, or phage can

generate multidrug-resistant bacteria by transduction which are then selected for by antibiotics. This

may be particularly important in the context of phage therapy, where phage are used to treat bacterial

infections, often alongside antibiotics. Our conclusions highlight the urgent need for clinical and

laboratory work to quantify the currently unknown contribution of phage to AMR evolution in

humans. Otherwise, we may be missing opportunities to reduce the global public health burden of

AMR.

INTRODUCTION

Bacteriophage (phage) are major bacterial predators and the most common organisms on the planet [1]. Phage are often present alongside antibiotics, naturally or in the context of antibacterial treatment [2–4], yet reports on their combined effect on bacteria are conflicting. Some previous studies showed that phage and antibiotics work synergistically to clear bacteria [5–8], while others have demonstrated that antibiotics reduce phage production [9,10]. This interaction has been extensively studied in biofilms to show that adding phage to disrupt the biofilm before adding antibiotics leads to the greatest synergistic killing [11–13]. However, whilst there is some work suggesting a similar timing-dependent synergistic killing effect on planktonic bacteria [6–8], this evidence is limited as only a few timings and concentrations of phage and antibiotics have been explored experimentally. This is further complicated by the fact that phage are also major drivers of bacterial evolution via horizontal gene transfer by transduction [14,15], which can notably contribute to antimicrobial resistance (AMR) spread [16,17]. If multidrug-resistant bacteria are generated by transduction, antibiotics present in the same environment may act as a synergistic selective pressure to increase their prevalence, yet to our knowledge the dynamics of transduction have not yet been investigated in such systems [18].

There are two types of transduction: specialised and generalised [19,20]. Specialised transduction occurs when a prophage sometimes picks up adjacent bacterial DNA upon excision from the bacterial chromosome, at the end of the lysogenic cycle. We also note lateral transduction as an important variation of specialised transduction, where phage packaging occurs before the prophage is excised from the bacterial chromosome, leading to transfer of DNA located further downstream of the prophage [21]. On the other hand, generalised transduction occurs during phage replication, when random non-phage DNA is packaged instead of phage DNA in a new phage particle. The resulting transducing phage is released upon bacterial lysis, and injects this DNA in another bacterium. Generalised transduction is likely the most important type of transduction in the context of AMR, as

it can lead to the horizontal transfer of any genetic material contained in a bacterium (including

plasmids, major vectors of AMR genes) [14].

A major bacterial pathogen often exposed to phage is *Staphylococcus aureus*, which at any given time

is colonising approximately 20% of humans [22]. Previous work suggests that all *S. aureus* carry

integrated prophage [23], and at least 50% of individuals colonised by *S. aureus* also carry free phage

capable of generalised transduction [24], the main mechanism of horizontal gene transfer for *S. aureus*

[23]. This is particularly relevant for methicillin-resistant *S. aureus* (MRSA), a group of *S. aureus* present

in both the ESKAPE list and the World Health Organization priority list of antibiotic-resistant bacteria

due to its large clinical burden [25–27]. *In vitro*, the generation rate of generalised transducing MRSA

phage carrying an AMR gene has been estimated to be approximately one per 10^8 new phage

produced, sufficient to consistently generate bacteria resistant to multiple antibiotics in less than 24h

[28], and *in vivo* transduction rates are likely to be even higher [29].

Understanding the dynamics of bacteria, antibiotics, phage and transduction is especially important

in the context of phage therapy, which aims to use phage as antibacterial agents, generally in

combination with antibiotics [4]. Phage therapy is currently investigated as a solution to counter the

threat of AMR, with ongoing clinical trials against MRSA infection [6,30–34]. Phage therapy guidelines

recommend that only phage with a limited ability to perform transduction should be used [19,35,36],

yet to our knowledge there is currently no technique to prevent phage from accomplishing generalised

transduction, which is fundamentally a mispackaging and thus a biological error similar to a mutation.

Hence, as previous reviews have highlighted, it is essential to explore the importance of this

mechanism, and identify conditions under which it could affect the outcome of therapy and lead to

multidrug-resistance evolution [18,35–37].

In this study, we aim to investigate the potential combined effect of antibiotics and phage capable of

generalised transduction on bacteria. Making conclusions about this multidimensional chequerboard

space of potential combinations and timings is difficult when using data from time-consuming, single-

scenario, *in vitro* experiments. Instead, we generate hypotheses to guide future experimental work by

simulating these conditions. We extend a previously published mathematical model of only phage-

bacteria dynamics, including generalised transduction, to newly incorporate the effect of antibiotics

on bacteria [28]. This model is parameterised using *in vitro* data from the same environment and set

of conditions as when it was originally developed [28], making it a reliable tool to infer the dynamics

governing this system.

We hypothesise that, depending on the timing and concentration at which they are added, phage and

antibiotics can either act synergistically to eradicate bacteria, or to create and select for multidrug-

resistant bacteria. We explore this in our model and generate guidelines to minimise the risk of

generating double-antibiotic-resistant *S. aureus* when two single-resistant strains are exposed to

antibiotics and phage, whilst maximising bacterial eradication.

MATERIALS AND METHODS

Laboratory methods

Bacterial strains and phage

Two *Staphylococcus aureus* strains were obtained from the Nebraska transposon library in the MRSA USA300 background [38]. These were NE327, with the *ermB* gene conferring resistance to erythromycin, and NE201KT7, a modified NE201 strain with a kanamycin resistance cassette replacing the *ermB* gene and a plasmid carrying the *tetK* gene conferring resistance to tetracycline. Horizontal gene transfer can only happen between these two strains via generalised transduction. When co-cultured with 80α phage, these give rise to double-resistant progeny (DRP) bacteria resistant to both erythromycin and tetracycline [28]. In the experiments conducted here, we used the double-resistant strain DRPET1, a DRP strain with a NE327 background generated during previous NE327, NE201KT7 and 80α co-cultures, containing both *ermB* and *tetK* genes [28]. Note that although 80α is a temperate phage, we have previously shown that lysogeny does not occur at a detectable level in our experiments, with a lysogenic frequency detection limit of 3.3×10^{-8} per non-lysogenic bacteria after 24h of co-culture, and that specialised transduction is unlikely to be responsible for horizontal gene transfer, due to the location of *tetK* on a plasmid and a ~1Mbp distance between *ermB* and the 80α integration site on the bacterial chromosome [28]. Hence the only interactions we consider between phage and bacteria in our environment are lysis and generalised transduction.

Time-kill curves

The growth conditions are identical to the ones previously used to generate data on lysis and transduction with these bacteria and phage [28]. Pre-cultures of each *S. aureus* bacterial strain (NE327, NE201KT7 and DRPET1) were separately generated overnight in 50mL conical tubes containing 10mL

of brain-heart infusion broth (BHIB, Sigma, UK). Unless otherwise stated, liquid cultures were incubated in a warm shaking water bath (37°C, 90 rpm). Each pre-culture was then diluted in phosphate-buffered saline (PBS) and mixed with 10mL of fresh BHIB in a 50mL conical tube to reach a starting concentration of 10^4 colony-forming units (cfu)/mL. The new culture was incubated for 2h to allow the bacteria to reach log-growth phase, following standard protocol for time-kill experiments [39]. Erythromycin or tetracycline was then added to the culture, at a concentration of either 0 (control), 0.25, 0.5, 1, 2, 4, 8, 16, or 32 mg/L. At 0, 1, 2, 3, 4, 6 and 24h after antibiotic addition, 30μl were sampled from the incubated culture, diluted in PBS, and plated on plain brain-heart infusion agar. The plates were incubated overnight at 37°C. Colonies on the plates were then counted to derive the concentration of bacteria in cfu/mL at the corresponding time point.

The experiment was repeated 3 times for each strain (NE327, NE201KT7 and DRPET1) and each antibiotic (erythromycin and tetracycline).

Minimum inhibitory concentration

We measured the minimum inhibitory concentrations (MIC) of erythromycin and tetracycline for NE327, NE201KT7 and DRPET1 by microbroth dilution [40]. Briefly, pre-cultures of each strain were generated overnight in Mueller-Hinton broth (MHB-II, Sigma, UK). Antibiotic stocks were generated in 50% ethanol at concentrations doubling from 0.25 to 256 mg/L, and 10μL of each dilution were added to separate wells on a 96-well plate. The overnight pre-cultures were diluted to a concentration of 10^5 cfu/mL, and 90μL were mixed with each antibiotic dilution in the 96-well plate. The plate was incubated overnight at 37°C, and the MIC were then determined by eye, identifying the lowest concentration of antibiotic which did not allow bacterial growth (i.e. the contents of the well were not turbid).

Modelling methods

All analyses were conducted in the R statistical analysis software [41]. The underlying code and data are available in a GitHub repository: https://github.com/qleclerc/phage_antibiotic_dynamics.

Mathematical model description

We extended a previously published mathematical model of phage-bacteria dynamics, including generalised transduction [28]. This original model focused on identifying the best method to capture the interaction between bacteria and phage *in vitro*, in the absence of antibiotics. This model was previously parameterised using the same bacterial and phage strains as in this study. Our three *S. aureus* strains are represented: B_E (erythromycin-resistant, corresponding to *in vitro* strain NE327), B_T (tetracycline-resistant, corresponding to NE201KT7), and B_{ET} (resistant to both erythromycin and tetracycline, corresponding to double-resistant bacteria) (Figure 1). The model also contains 80α lytic phage (P_L), which can give rise to transducing phage carrying either the erythromycin (P_E) or tetracycline (P_T) resistance gene. The transducing phage only carry an antibiotic resistance gene, and hence cannot lyse bacteria, but can give rise to B_{ET} bacteria via generalised transduction (Figure 1).

Antibiotic pharmacodynamics

We added two compartments to the model to track the concentration of antibiotics: C_E for erythromycin, and C_T for tetracycline. The concentrations are expressed in mg/L. We can simulate the addition of a chosen concentration of antibiotic at any given time, and the antibiotics decay at a constant rate γ_E and γ_T for erythromycin and tetracycline, respectively (Figure 1). The change in concentration of erythromycin over time is therefore expressed as

$$\frac{dC_E}{dt} = -\gamma_E \cdot C_E \ (1)$$

and the change in concentration of tetracycline over time is expressed as

179
$$\frac{dC_T}{dt} = -\gamma_T \cdot C_T \quad (2)$$

180   The antibiotic concentrations are then used to calculate $\varepsilon_{i,j}$, the effect of each antibiotic i (i $\in$ {E, T}) on

181   each bacterial strain j ($j \in$ {E, T, ET}), according to a pharmacodynamic relationship parameterised

182   through Hill equations [39,42], expressed as

183
$$\varepsilon_{i,j} = \mu_j^{max} \cdot \varepsilon_{i,j}^{max} \cdot \frac{C_i^{(H_{i,j})}}{EC50_{i,j}^{(H_{i,j})} + C_i^{(H_{i,j})}} \quad (3)$$

This commonly used function calculates the effect of an antibiotic using four parameters: the

maximum effect ($\varepsilon^{max}_{i,j}$), current concentration (C_i), half maximal effective concentration ($EC50_{i,j}$), and

a Hill coefficient ($H_{i,j}$). Note that $\varepsilon^{max}_{i,j}$ is relative to the maximum growth rate of the corresponding

strain j (μ^{max}_j). This was necessary to ensure that these values would work with our previous estimates

for bacterial growth, as these were slightly lower than the baseline growth rates estimated here (see

below – "Parameter estimation").

Figure 1: Mathematical model diagram. This model is an extension of our original model presented

in [28], with the inclusion of antibiotic effects. Each bacteria strain (B_E resistant to erythromycin, B_T

resistant to tetracycline, or B_{ET} resistant to both) can replicate (purple). The lytic phage (P_L) multiply

by infecting a bacterium and bursting it to release new phage (gold). This process can create

transducing phage (P_E or P_T) carrying a resistance gene (*ermB* or *tetK* respectively) taken from the

infected bacterium (dashed, green). These transducing phage can then generate new double-resistant

progeny (B_{ET}) by infecting the bacteria strain carrying the other resistance gene (solid, green). The

antibiotics, erythromycin (C_E) and tetracycline (C_T), decrease the growth rate of each bacteria strain to

varying extents, depending on their concentration and the resistance level of the strain (dotted, red

and orange). Phage and antibiotics can decay at a fixed rate (dotted, grey and black).

Bacterial growth and phage predation

203 The bacterial growth rate μ_ϑ ($\theta \in \{E, T, ET\}$, $N = B_E + B_T + B_{ET}$) is modelled using a logistic function given

204 by

$$\mu_\theta = \mu_\theta^{max} \cdot (1 - \frac{N}{N^{max}}) \qquad (4)$$

206 with a maximum growth rate of μ^{max}_ϑ and carrying capacity N^{max}.

207 Phage predation is modelled as a frequency-dependent process. Previous work has suggested that this

208 interaction is more biologically realistic than the more commonly used density-dependent interaction

209 [28]. At any given time, the proportion λ of phage successfully infecting bacteria is calculated as

$$\lambda = (1 - exp\left(-\beta \cdot N\right)) \qquad (5)$$

211 with β representing the rate of successful phage adsorption leading to bacterial lysis, and the

212 proportion φ_ϑ of bacteria successfully infected by the phage of type P_ϑ ($\theta \in \{L, E, T\}$) is equal to

213
$$\varphi_\theta = \left(1 - exp\left(-\lambda \cdot \frac{P_\theta}{N}\right)\right) \quad (6)$$

214 This interaction ensures that, for any given time, the number of phage infecting bacteria and the

215 number of bacteria infected by phage cannot be greater than the total number of phage and bacteria,

216 respectively.

217 In the model, phage burst size δ_θ for each bacteria strain θ (θ ∈ {E, T, ET}) decreases from the maximum

218 phage burst size δ^{max} as bacterial growth decreases, according to

219
$$\delta_\theta = \delta^{max} \cdot max\left(0, 1 - \frac{N}{N^{max}} - \frac{\varepsilon_{E,\theta}}{\mu_\theta^{max}} - \frac{\varepsilon_{T,\theta}}{\mu_\theta^{max}}\right) \quad (7)$$

This link between phage burst size and bacterial growth was identified as the most biologically

plausible explanation for the dynamics we have previously observed between the bacteria and phage

in our system [28]. This decrease happens as bacteria enter stationary phase, when the population

approaches carrying capacity N^{max} (as demonstrated previously [28,43,44]), but here the additional

effect of antibiotics must be included. To capture this, we use the same effective scaling as the

bacterial growth rate (Equation 4) with inclusion of the relative effect of antibiotics $\varepsilon_{E,\theta}$ and $\varepsilon_{T,\theta}$ in the

phage burst size estimation. Since antibiotics can kill bacteria, leading to a net negative growth rate,

we limit the multiplier value to 0 as a minimum, to prevent a negative burst size.

Bacteria model equations

The changes in bacteria numbers over time are given by

231
$$\frac{dB_E}{dt} = \mu_E \cdot \{B_E - [\omega \cdot (\varphi_L + \varphi_T) \cdot B_E]\} - (\varphi_L + \varphi_T) \cdot B_E - (\varepsilon_{E,E} + \varepsilon_{T,E}) \cdot B_E \quad (8)$$

{Change in B_E = growth of B_E – infections by P_L – infections by P_T - ery killing - tet killing}

$$\frac{dB_T}{dt} = \mu_T \cdot \{B_T - [\omega \cdot (\varphi_L + \varphi_E) \cdot B_T]\} - (\varphi_L + \varphi_E) \cdot B_T - (\varepsilon_{E,T} + \varepsilon_{T,T}) \cdot B_T \quad (9)$$

{Change in B_T = growth of B_T – infections by P_L – infections by P_E - ery killing - tet killing}

$$\frac{dB_{ET}}{dt} = \mu_{ET} \cdot [B_{ET} - (\omega \cdot \varphi_L \cdot B_{ET})] - \varphi_L \cdot B_{ET} + \varphi_E \cdot B_T + \varphi_T \cdot B_E - (\varepsilon_{E,ET} + \varepsilon_{T,ET}) \cdot B_{ET}$$

$$(10)$$

{Change in B_{ET} = growth of B_{ET} – infections by P_L + infections of B_T by P_E + infections of B_E by P_T - ery

killing - tet killing}

In this model, we assume that over one time step, bacterial replication and infection by phage are

mutually exclusive events, hence the repetition of the φ_ϑ terms ($\theta \in \{L, E, T\}$) on the right-hand side of

Equations 8-10. As we have shown previously, this biologically plausible assumption is necessary due

to the upper limits of the growth rate parameters being higher than the limits of the phage infection

parameters, preventing the bacterial population from decreasing due to phage killing if this

assumption is not included in the model [28]. The parameter ω represents the mutually exclusive

nature of these events over one time-step in the equations, and is therefore equal to 1 (i.e. one time-

step, with units of hours).

Some combinations of phage infection are not explicitly represented in the equations as they

represent a flow from one compartment towards the same compartment. P_L are the lytic phage,

capable of lysing all the bacteria, and are therefore included in all equations. However, the generalised

transducing phage P_E and P_T only carry a bacterial antibiotic resistance gene and cannot lyse bacteria.

Whilst they can infect any bacteria in our system to transfer the gene, this process will not affect

bacteria already carrying the corresponding gene. For example, if a bacterium B_E is infected by a phage

P_E, this bacterium remains only resistant to erythromycin, and therefore still belongs to the B_E

compartment.

Phage model equations

The change in phage numbers over time are calculated as

$$\frac{dP_L}{dt} = \varphi_L(t-\tau) \cdot \delta_E \cdot (1-\alpha) \cdot B_E \ +$$

$$\varphi_L(t-\tau) \cdot \delta_T \cdot (1-\alpha) \cdot B_T \ +$$

$$\varphi_L(t-\tau) \cdot \delta_{ET} \cdot (1-2 \cdot \alpha) \cdot B_{ET} \ -$$

$$\lambda \cdot P_L - \gamma_P \cdot P_L \qquad\qquad (11)$$

{Change in P_L = new P_L phage from B_E + new P_L phage from B_T + new P_L phage from B_{ET} −

P_L phage infecting bacteria − P_L decay}

$$\frac{dP_E}{dt} = \varphi_L(t-\tau) \cdot \delta_E \cdot \alpha \cdot B_E \ +$$

$$\varphi_L(t-\tau) \cdot \delta_{ET} \cdot \alpha \cdot B_{ET} \ -$$

$$\lambda \cdot P_E - \gamma_P \cdot P_E \qquad\qquad (12)$$

{Change in P_E = new P_E phage from B_E + new P_E phage from B_{ET} −

P_E phage infecting bacteria − P_E decay}

$$\frac{dP_T}{dt} = \varphi_L(t-\tau) \cdot \delta_T \cdot \alpha \cdot B_T \ +$$

$$\varphi_L(t-\tau) \cdot \delta_{ET} \cdot \alpha \cdot B_{ET} \ -$$

$$\lambda \cdot P_T - \gamma_P \cdot P_T \qquad\qquad (13)$$

{Change in P_T = new P_T phage from B_T + new P_T phage from B_{ET} −

274   *P_T phage infecting bacteria – P_T decay}*

In Equations 11-13, the latent period τ corresponds to the delay between bacterial infection and

burst, and the transduction probability α corresponds to the proportion of new phage released upon

burst which are transducing phage carrying the erythromycin or tetracycline resistance gene. Note

that a constant rate γ_P is used for phage decay (Equations 11-13), similar to the rates used for

antibiotic decay (Equations 1-2).

Parameter estimation

Parameters for bacterial growth and phage predation were originally obtained by fitting our model to

in vitro data for the same bacteria and phage strains as in this study [28]. Briefly, the parameters were

estimated using the Markov chain Monte Carlo Metropolis–Hastings algorithm. We ran the algorithm

with two chains until convergence was achieved (determined using the Gelman-Rubin diagnostic, with

the multivariate potential scale reduction factor < 1.2 [45]), then for each parameter we generated

50,000 samples from the posterior distributions [28].

Parameters for antibiotic effect were obtained in two steps using the least squares methods for model

fitting, which aims to minimise the squared difference between data and model output. Firstly, the

reduction in bacterial growth caused by a specific concentration of antibiotic ($\acute{\varepsilon}$) was obtained by

fitting a deterministic growth model given by

291
$$\frac{d\mathrm{B}}{dt} = \mu \cdot \left(1 - \frac{B}{N^{max}}\right) \cdot \mathrm{B} - \acute{\varepsilon} \cdot B \quad (14)$$

to the bacterial concentration B over time, estimated as the mean of three *in vitro* replicates.

$\acute{\varepsilon}$ was then scaled to growth ($\varepsilon = \acute{\varepsilon}/\mu$), to represent the relative impact of the antibiotic on bacterial

growth rather than an absolute value. This was necessary to ensure that these antibiotic impact values

would work with our previous estimates for bacterial growth, as these were slightly lower than the

baseline growth rates estimated here. Since we obtained growth curve time series data for 8

concentrations for each strain, we generated 8 estimates of antibiotic effect for each strain.

Secondly, the three parameters of the Hill equation [39,42] (maximum effect ε^{max}, Hill coefficient H,

and half maximum effective concentration $EC50$), which calculates the antibiotic effect as a function

of concentration, were estimated by fitting Equation 3 to the 8 antibiotic effects for each strain.

Model scenarios considered

Antibiotics alone

303 We start with an environment containing both single-resistant strains at stationary phase (10^9 cfu/mL).

304 This mirrors the within-host diversity we could expect to see during bacterial infections [24], and the

305 fact that bacteria most often live at stationary phase in the environment [46]. We first investigate the

306 effect of the presence of either or both erythromycin and tetracycline at concentrations of 1 mg/L,

307 similar to antibiotic concentrations measured *in vivo* during treatment [47,48].

Antibiotic and phage

We then consider scenarios where phage are present. We first artificially inactivate transduction, by

setting the corresponding model parameter to 0. We add 10^9 pfu/mL of phage, similar to a

concentration that could be added during phage therapy, and equivalent to a multiplicity of infection

(ratio of phage to bacteria) of 1, similar to what could be found naturally in the environment [1]. We

consider scenarios where the phage are present alone, alongside only one antibiotic, or alongside both

antibiotics. Phage and antibiotics are assumed to be introduced concurrently at the start, and the

simulations run for 24h.

We then repeat these scenarios, but with transduction enabled to levels that were previously

observed *in vitro*, with approximately 1 transducing phage carrying an AMR gene generated for each

10^8 new lytic phage [28].

We repeat the analyses above with either single-resistant strain present at 10^9 cfu/mL, and the other

at 10^6 cfu/mL (0.1%). This could correspond to a scenario where bacterial diversity is underestimated,

with only one type of resistance detected [24].

Antibiotic and phage level and timing variation

To further investigate the scenario where transduction is enabled and both single-resistant bacterial

strains have a starting concentration of 10^9 cfu/mL, we vary the timing of introduction for antibiotic

and phage, with up to 24h delay between their respective additions, as well as varying the

concentration of antibiotics between 0.2 and 2.2 mg/L, and phage between 10^5 and 10^{10} pfu/mL,

chosen to reflect realistic ranges [1,47,48]. We run these simulations for 48h after the introduction of

antibiotics or phage (whichever is added first).

Phage and bacteria parameters variation

Finally, we vary the probability for phage to perform generalised transduction to identify threshold

values which dictate whether multidrug-resistant bacteria appear above detectable levels within 24h.

To explore the sensitivity of our results to the parameter values estimated from this single

environment, we also conduct a partial rank correlation to assess the effect of varying phage predation

parameters (adsorption rate, latent period, burst size), bacteria growth rates, and antibiotic and phage

decay on total bacteria remaining after 48h, and maximum double-resistant bacteria generated. This

allows us to generalise our conclusions as these parameters likely vary for different combinations of

phage, antibiotics and bacteria. The ranges evaluated are shown in Table 1, and are either derived

from our initial parameterisation of the model [28], or from other studies [49,50].

RESULTS

In vitro pharmacodynamics of erythromycin and tetracycline

Erythromycin at all concentrations caused a decrease in erythromycin-sensitive NE201KT7 numbers over 6h, but only slowed down growth for erythromycin-resistant NE327 and double-resistant DRPET1, even at 32 mg/L (Figure 2). Tetracycline caused a decrease in bacterial numbers over 6h at concentrations greater than 0.5 mg/L for tetracycline-sensitive NE327, 8 mg/L for tetracycline-resistant NE201KT7, and 4 mg/L for DRPET1 (Figure 2). As a comparison, the minimum inhibitory concentrations (MICs) of erythromycin measured by microbroth dilution were 0.25, >256 and >256 mg/L for NE201KT7, NE327 and DRPET1 respectively. For tetracycline, the MICs were 32, 0.25 and 32 mg/L for NE201KT7, NE327 and DRPET1 respectively.

Figure 2: Growth curves of NE201KT7 (tetracycline-resistant, left), NE327 (erythromycin-resistant,

middle) and DRPET1 (double-resistant, right), exposed to varying concentrations of erythromycin

(top) or tetracycline (bottom). Solid lines show *in vitro* data, with error bars indicating mean +/-

standard deviation, from 3 replicates. Dashed lines show model output after fitting. Values indicate

the percentage of model points that fall within the range of the corresponding *in vitro* data point +/-

standard deviation. cfu: colony-forming units. Note that cfu per mL are shown on a log-scale.

The Hill equations fitted well to the effect of varying concentrations of antibiotics on the bacteria

(Figure 2, Figure S1). The corresponding parameter values are presented in Table 1. The antibiotic

effect curves for the DRPET1 are similar to the one for NE201KT7 for tetracycline and NE327 for

erythromycin (Figure S1), which was expected since DRPET1 contains both antibiotic-resistance genes

from NE201KT7 (*tetK*) and NE327 (*ermB*).

371   **Table 1: Model parameter values.** Parameters with no units are dimensionless. All estimates were

372   obtained by fitting the model to *in vitro* data, except those marked with a * which are assumed.

Name (unit)		Symbol	Estimate	Range for sensitivity analysis	Reference
Bacterial growth parameters	Carrying capacity (bacteria.mL^{-1})	N^{max}	2.76×10^9	-	[28]
	B_E growth rate (h^{-1})	μ^{max}_E	1.61	1.59 - 1.63	
	B_T growth rate (h^{-1})	μ^{max}_T	1.51	1.49 - 1.53	
	B_{ET} growth rate (h^{-1})	μ^{max}_{ET}	1.44	1.42 - 1.47	
Phage parameters	Phage adsorption rate (phage^{-1}.h^{-1})	β	2.3×10^{-10}	2.1×10^{-10} - 2.4×10^{-10}	
	Phage burst size (phage)	δ^{max}	76	70 - 83	
	Phage latent period (h)	τ	0.72	0.69 - 0.77	
	Transduction probability (phage^{-1})	α	1.0×10^{-8}	8.5×10^{-9} - 1.4×10^{-8}	
	Phage decay rate (h^{-1})	γ_P	0*	0 - 0.1	[49]
Antibiotic parameters	Erythromycin decay rate (h^{-1})	γ_E	0*	0 - 0.1	[50]
	Tetracycline decay rate (h^{-1})	γ_T	0*	0 - 0.1	
Effect of erythromycin on B_E	Max effect (h^{-1})	$\varepsilon^{max}_{E,E}$	1.10	-	This study
	Half maximal effective concentration (mg.L^{-1})	$EC50_{E,E}$	3.42	-	
	Hill coefficient	$H_{E,E}$	0.67	-	
Effect of erythromycin on B_T	Max effect (h^{-1})	$\varepsilon^{max}_{E,T}$	3.91	-	
	Half maximal effective concentration (mg.L^{-1})	$EC50_{E,T}$	9.26	-	
	Hill coefficient	$H_{E,T}$	0.22	-	
Effect of erythromycin	Max effect (h^{-1})	$\varepsilon^{max}_{E,ET}$	0.95	-	

on B_{ET}	Half maximal effective concentration (mg.L^{-1})	EC50$_{E,ET}$	2.47	-
	Hill coefficient	H$_{E,ET}$	0.79	-
Effect of tetracycline on B_E	Max effect (h^{-1})	$\varepsilon^{max}_{T,E}$	5.40	-
	Half maximal effective concentration (mg.L^{-1})	EC50$_{T,E}$	81.43	-
	Hill coefficient	H$_{T,E}$	0.30	-
Effect of tetracycline on B_T	Max effect (h^{-1})	$\varepsilon^{max}_{T,T}$	1.66	-
	Half maximal effective concentration (mg.L^{-1})	EC50$_{T,T}$	7.27	-
	Hill coefficient	H$_{T,T}$	2.41	-
Effect of tetracycline on B_{ET}	Max effect (h^{-1})	$\varepsilon^{max}_{T,ET}$	1.58	-
	Half maximal effective concentration (mg.L^{-1})	EC50$_{T,ET}$	4.42	-
	Hill coefficient	H$_{T,ET}$	1.70	-

Model-predicted antibacterial effect of concurrent erythromycin, tetracycline and bacteriophage presence

When simulating the dynamics of two single-resistant *S. aureus* strains in our mathematical model, as

expected, the presence of only one antibiotic at 1 mg/L (4 x MIC for susceptible strains) leads to a

decrease in the susceptible strain, while the resistant strain does not decrease (Figure 3a, top row).

On the other hand, the presence of both antibiotics, or of only phage without transduction, causes a

decrease in both bacterial strains (Figure 3a, top and middle rows). The presence of phage and one

antibiotic leads to a decrease in both bacterial strains, with the antibiotic-susceptible strain decreasing

faster (Figure 3a, middle row). Out of all the conditions shown here, the presence of phage and both

antibiotics leads to the fastest decrease in both bacterial strains (Figure 3b). For comparison, to

replicate the combined effect of 10^9 pfu/mL of phage, 1 mg/L of erythromycin, and 1 mg/L of

tetracycline, we would need 4.58 mg/L of erythromycin and 1.14 mg/L of tetracycline in the absence

of phage (Figure S2).

Transduction does not appear to affect the antibacterial activity of antibiotics and phage when phage

are either present alone, alongside erythromycin, or alongside tetracycline (Figure 3a, bottom row).

Double-resistant progeny bacteria (B_{ET}) appear, but only reach a maximum concentration of 30

cfu/mL, and do not remain higher than 1 cfu/mL for more than 8h (Figure 3a, bottom row). However,

when both erythromycin and tetracycline are present alongside phage capable of transduction, there

is a steady increase in the number of B_{ET} throughout 24h, reaching 80 cfu/mL after 24h (Figure 3a,

bottom row & Figure 3b). It is important to note here that when both antibiotics are present, phage

numbers do not increase throughout the 24h period, regardless of transduction ability (Figure 3a,

middle and bottom rows & Figure 3b, right).

Figure 3: a) Model-predicted dynamics with two single-resistant strains starting at carrying capacity (10⁹ colony-forming units (cfu)/mL), in the presence of no antibiotics (1st column), erythromycin only (2nd column), tetracycline only (3rd column), or both erythromycin and tetracycline (4th column), combined with either no phage (top row), phage incapable of transduction (middle row), or phage capable of generalised transduction (bottom row). The starting strains are either erythromycin-resistant (B_E) or tetracycline-resistant (B_T). Antibiotics and/or phage (P_L) are present at the start of the simulation, at concentrations of 1 mg/L (4 x MIC for susceptible strains) and 10⁹ plaque-

forming units (pfu)/mL respectively. Double-resistant bacteria (B_{ET}) can be initially generated by

generalised transduction only, and then by replication of existing B_{ET}. Dashed line indicates the

detection threshold of 1 cfu or pfu/mL. **b) Change in bacteria (single-resistant to erythromycin,**

single-resistant to tetracycline, or double-resistant) and phage numbers depending on the antibiotic

exposure, in the presence of phage capable of generalised transduction. Dashed line indicates the

detection threshold of 1 cfu or pfu/mL.

With a reduced starting concentration of either *S. aureus* strain to 10^6 cfu/mL, while the other remains

at 10^9 cfu/mL, the conclusions are similar as described above, regardless of which strain is in the

minority. Figure 4 shows the scenario where tetracycline-resistant bacteria are in minority, and Figure

S3 shows the scenario where erythromycin-resistant bacteria are in minority. When erythromycin,

tetracycline, and phage capable of generalised transduction are all present, B_{ET} are still generated,

although they do not reach a concentration higher than 1 cfu/mL within 24h, instead of 11h above

(Figure 4, Figure 3).

Figure 4: a) Model-predicted dynamics with one single-resistant strains starting at carrying capacity (10⁹ colony-forming units (cfu)/mL) and the second in minority (10⁶ cfu/mL), in the presence of no antibiotics (1st column), erythromycin only (2nd column), tetracycline only (3rd column), or both erythromycin and tetracycline (4th column), combined with either no phage (top row), phage incapable of transduction (middle row), or phage capable of generalised transduction (bottom row). Erythromycin-resistant bacteria (B_E) are initially present at a concentration of 10^9 cfu/mL, and tetracycline-resistant bacteria (B_T) at 10^6 cfu/mL. Antibiotics and/or phage (P_L) are present at the start

of the simulation, at concentrations of 1 mg/L (4 x MIC for susceptible strains) and 10^9 plaque-forming

units (pfu)/mL respectively. Double-resistant bacteria (B_{ET}) can initially be generated by generalised

transduction only, and then by replication of existing B_{ET}. Dashed line indicates the detection threshold

of 1 cfu or pfu/mL. **b) Change in bacteria (single-resistant to erythromycin, single-resistant to**

tetracycline, or double-resistant) and phage numbers depending on the antibiotic exposure, in the

presence of phage capable of generalised transduction. Dashed line indicates the detection threshold

of 1 cfu or pfu/mL.

Effect of variation in antibiotic and phage timing and concentration on bacterial populations

Under conditions where both erythromycin and tetracycline are present, the timing and concentration

of antibiotics and phage capable of generalised transduction drives substantial variation in dynamics,

with failure to clear all bacteria after 48h being a possibility (Figure 5a-b, top row).

With a phage concentration of 10^8 cfu/mL and for any antibiotics concentration between 0.2 and 2.2

mg/L (Figure 5a), the optimal conditions to clear bacteria within 48h are when phage are initially

present, and antibiotics are introduced at least 5h later (Figure 5a, top row). However, this

systematically leads to B_{ET} appearance, with a peak concentration between 10 and 100 cfu/mL (Figure

5a, middle row), and a presence time (hours when cfu/mL > 1) of up to 10h (Figure 5a, bottom row).

Interestingly, if antibiotics are introduced exactly 5h after phage, the peak concentration of B_{ET} is only

between 1 and 10 (Figure 5a, middle row). The presence of antibiotics at the same time or shortly

before phage leads to failure to clear bacteria within 48h (up to between 10^8 and 10^{10} cfu/mL

remaining after 48h - top row), and substantial B_{ET} appearance (maximum concentration up to

between 10^8 and 10^{10} cfu/mL - middle row; presence time up to between 30 and 40h- bottom row).

Regardless of timings, the addition of at least 2.2 mg/L of antibiotics guarantees that less than 100

bacteria remain after 48h (top row), and no detectable B_{ET} if the antibiotics are added at least 2h

before the phage (bottom row).

When keeping the antibiotic concentrations at 1 mg/L, but varying the phage concentration, the

impact of the delay between phage and antibiotic presence on the bacterial population is strongly

dependent on phage concentration (Figure 5b). We again see that bacteria are not cleared within 48h

if antibiotics are present at the same time as or shortly before a concentration of phage between 10^5

and 10^9 pfu/mL (Figure 5b, top row). However, we now note that this also occurs if antibiotics are

introduced after a concentration of phage between 10^5 and 10^8 pfu/mL (Figure 5b, top row).

Regardless of timing, the presence of a phage concentration of more than 10^9 pfu/mL guarantees

bacterial clearance within 48h (top row), but leads to B_{ET} if the antibiotics are introduced after the

phage (maximum concentration between 10 and 100 cfu/mL - middle row; presence time between 1

and 10h - bottom row).

Figure 5: a-b) Varying timing (x-axis) and dose of antibiotic and phage (y-axis) affects total bacterial

count after 48h (top), maximum concentration of double-resistant bacteria (B$_{ET}$) (middle), and time

when the concentration of B$_{ET}$ is greater than 1 colony-forming unit (cfu) per mL (bottom). a) Adding

10^8 plaque-forming units (pfu) per mL of phage, and between 0.2 and 2.2 mg/L of both erythromycin

and tetracycline. **b)** Adding 1 mg/L of both erythromycin and tetracycline, and between 10^5 and 10^{10}

pfu/mL of phage. The x-axis indicates the time when antibiotics were added, relative to when phage

were added. For example, the value "4" indicates that phage were present at the start of the

simulation, and antibiotics were introduced 4h later. The segments with black borders correspond to

the dynamics shown in c). **c) Phage and bacteria dynamics over 48h for 4 conditions taken from panel**

b. In all 4 conditions, indicated by the black rectangles, phage are initially present at a concentration

of 10^8 pfu/mL, while erythromycin and tetracycline are both introduced at concentrations of 1 mg/L

after either 0h, 3h, 5h or 15h, stated on the plots, with the timing indicated by the vertical dashed

lines. Horizontal dotted lines indicate bacteria remaining after 48h (corresponding to the top row of

a-b) and maximum double-resistant bacteria (B_{ET}) concentration (middle row of a-b). Solid line

indicates the detection threshold of 1 cfu or pfu/mL. The concentrations of single-resistant bacteria

(B_E, blue, and B_T, green) overlap and cannot be distinguished.

To investigate the dynamics behind these results, we selected four conditions from Figure 5b

(indicated by the 4 black rectangles) with varying antibiotic addition times and plotted the underlying

phage and bacteria dynamics over 48h for each (Figure 5c). In all four conditions the starting

concentrations are 10^8 pfu/mL for phage and 1 mg/L for antibiotics, but antibiotics are introduced

either 0h, 3h, 5h or 15h after phage. These plots show that if phage are increasing, they stop

immediately following antibiotic addition (Figure 5c, 3 and 5h delay). If antibiotics are added too soon

after phage (Figure 5c, 0 and 3h delay), phage do not reach a high enough number to exert a sufficient

killing pressure on bacteria. In that case B_{ET}, which are not substantially affected by either antibiotic,

replicate faster than they are killed by phage. After more than 35h, the B_{ET} population reaches a

sufficiently high number such that the phage population increases, and the resulting pressure is

enough to lead to a net negative bacterial growth rate. If antibiotics are added 15h or later after phage

(Figure 5b, Figure 5c), B_{ET} generation will not change, as during this period they will have already arisen

by transduction and been removed by phage predation. The presence of antibiotics 5h after phage is

optimal to ensure the lowest maximum number of B_{ET} (Figure 5c - compare horizontal dotted lines).

This timing allows phage to initially increase to a concentration of almost 5 x 10^9, sufficiently high to

exert a strong killing pressure on bacteria, while the added effect of antibiotics prevents further B_{ET}

generation by decreasing the single-resistant strains.

These results also apply to a scenario where one of the two bacterial strains is in the minority (starting

concentration of 10^6 instead of 10^9 cfu/mL), regardless of which strain is in the minority (Figures S4 &

S5). However, B_{ET} peak, presence time, and total bacteria remaining after 48h decrease faster with a

higher dose of antibiotic or phage, or with a higher delay between phage and antibiotics, suggesting

that phage and antibiotics are able to exert a greater killing pressure and generate fewer B_{ET} when one

strain is in the minority.

Effect of variation in phage and bacteria parameters on multidrug-resistance evolution

Our results above rely on parameters estimated from phage and bacteria interactions *in vitro*, but

these may vary depending on the bacteria, phage, and environment. We can explore these different

conditions using our model to quantify the dynamics under varying values for parameters governing

phage-bacteria interactions (see Table 1 for the ranges used) to determine whether our results would

hold.

We first examine the impact of varying the transduction probability on our results (corresponding to

the probability that a transducing phage carrying an AMR gene is released instead of a lytic phage

during bacterial burst) as this is a vital and yet poorly quantified parameter. When 10^9 cfu/mL of each

single-resistant strain are simultaneously exposed to 10^9 pfu/mL of phage and 1 mg/L of erythromycin

and tetracycline, varying the transduction probability between 10^{-10} and 10^{-6} leads to a similar log-fold

increase in double-resistant bacteria numbers (B_{ET}, Figure 6a). Decreasing the probability only delays

the appearance of B_{ET} in the model, and does not prevent it. However, a probability lower than 10^{-9}

may prevent the appearance of B_{ET} in reality, since the single-resistant strains become undetectable

(< 1 cfu/mL) before the B_{ET} become detectable. If antibiotics are added more than 10h after phage, B_{ET}

will have already started declining due to phage predation, hence the antibiotics only contribute to

further increasing the decline in bacterial numbers (Figure 5, Figure 6b). Under these conditions, a

transduction probability lower than 10^{-9} is necessary to prevent B_{ET} from increasing past the detection

threshold (1 cfu/mL) (Figure 6b). If antibiotics are introduced 10h before phage, the resulting decline

in single-resistant bacteria prevents any B_{ET} from reaching a detectable level before single-resistant

bacteria are eradicated (< 1 cfu/mL), even with the highest transduction probability of 10^{-6} (Figure 6c).

Looking at how changes in other phage and bacteria parameters may affect our results, partial rank

correlation shows that an increase in phage predation either through an increase in phage adsorption

rate (β) or phage burst size (δ^{max}) correlates with a decrease in maximum B_{ET} detected over 48h (Figure

6d, blue). For example, the correlation coefficient of -0.65 between β and maximum B_{ET} implies that a

100% increase in adsorption rate is correlated with a 65% decrease in maximum double-resistant

bacteria detected over 48h. An increase in phage burst size is also correlated with a decrease in

bacteria remaining after 48h. However, adsorption rate is not substantially correlated with bacteria

remaining after 48h (Figure 6d, red). Finally, an increase in latent period (τ), equivalent to a decrease

in predation since phage will take longer before lysing the bacteria, is correlated with an increase in

maximum B_{ET}, but is not substantially correlated with bacteria remaining after 48h.

Although we were not able to parameterise our model for antibiotic decay, our 24h time-kill curves

suggest that antibiotics no longer decrease bacterial numbers after 24h (Figure S6). Phage may also

be affected by decay in the environment. Hence, we included these parameters in our partial rank

correlation analysis. An increase in phage or antibiotic decay (γ_P, γ_E, γ_T) is correlated with an increase

in maximum B_{ET} (Figure 6d, blue), but unexpectedly with a decrease in bacteria remaining after 48h

(Figure 6d, red). This is explained by the fact that such an increase leads to a weakened killing pressure

on B_{ET}, which are able to increase faster. This translates to a shorter time before the phage population

is able to increase again due to enough bacteria being available for predation, and thus a shorter time

before B_{ET} start decreasing due to phage killing (Figure S7). This also explains why an increase in the

transduction probability (α) is correlated with a decrease in bacteria remaining after 48h, as it leads

to a faster increase in B_{ET}, and thus a shorter time before the phage are able to increase and eradicate

the B_{ET} population. Finally, single-resistant bacterial growth rates (μ^{max}_E, μ^{max}_T) are not substantially

correlated with either maximum B_{ET} or remaining bacteria, while double-resistant growth rate (μ^{max}_{ET})

is only weakly positively correlated with maximum B_{ET}, and negatively with remaining bacteria.

Figure 6: Sensitivity of phage-bacteria dynamics to changes in model parameters. Effect of varying

the transduction probability between 10^{-10} and 10^{-6} when a) antibiotics and phage are present at the

start of the simulation, b) phage are present at the start, antibiotics are introduced 10h later, and c)

antibiotics are present at the start, phage are added 10h later. Transduction probability is defined as

the probability that a transducing phage carrying an AMR gene is released instead of a lytic phage

during bacterial burst. The dashed lines for single-resistant bacteria overlap and cannot be

distinguished. Vertical dashed lines indicate timing of addition of antibiotics or phage. cfu/mL: colony-

forming units per mL. **d) Partial rank correlation between model parameters, and remaining bacteria**

after 48h (pink) or maximum double-resistant bacteria (B_{ET}) concentration (blue). Information on the

parameter ranges investigated can be found in Table 1. β: adsorption rate, δ^{max}: burst size, τ: latent

period, α: transduction probability, γ_P: phage decay, γ_E: erythromycin decay, γ_T: tetracycline decay,

μ^{max}_E: B_E growth rate, μ^{max}_T: B_T growth rate, μ^{max}_{ET}: B_{ET} growth rate.

DISCUSSION

Summary of results

In this work, we reconcile the existing literature suggesting either that phage and antibiotics can synergistically kill bacteria, or that antibiotics reduce the efficacy of phage predation, whilst also considering the synergy of phage and antibiotics on antimicrobial resistance (AMR) evolution. We showed that although phage replication is limited in the presence of antibiotics, which negatively affect bacterial growth, phage are still able to exert a strong killing pressure on bacteria to complement the action of antibiotics. Under such conditions, the concentration and timing of antibiotics and phage are essential: phage introduced after antibiotics at a low concentration may not be able to replicate and hence not contribute to killing the bacterial population. Phage and antibiotics can act synergistically to drive AMR evolution when phage generate multidrug-resistant bacteria by transduction, and antibiotics act as a selection pressure. This is again exacerbated by the reduction in phage replication caused by antibiotics: a low concentration of phage capable of transduction introduced after antibiotics will not exert a strong killing pressure on bacteria, and instead generate multidrug-resistant bacteria at a background rate which are then selected for by the antibiotics.

Optimal conditions to clear bacteria and minimise AMR evolution

The best conditions to guarantee bacterial eradication within 48h whilst minimising the risk of multidrug-resistant bacteria evolution in our system are when antibiotics are present before phage, and either when erythromycin and tetracycline are present at a concentration of at least 2 mg/L (8 x MIC for susceptible strains), or phage at a concentration of at least 10^{10} pfu/mL (Figure 5a-b). Multidrug-resistant bacteria generation is also restricted if phage are introduced at least 10h after antibiotics (Figure 5a-b). This may be particularly relevant in the context of antibacterial treatment,

587 further discussed below. The worst outcome in which antibiotics and phage fail to rapidly clear all

588 bacteria and instead act synergistically to drive AMR evolution is a combination of a low antibiotic

589 dose (< 1 mg/L) with a low phage dose (< 10^9 pfu/ml), introduced around the same time (Figure 5a-b).

590 Unfortunately, this may correspond to values seen in natural environments where phage are

591 commonly found and antibiotics are residually present due to pollution [2,3].

Importance of transduction in the environment and during phage therapy

Our results highlight the necessity to better understand the role of transduction in AMR spread and

evolution, and not assume by default that it is too rare to be relevant compared to other mechanisms

of horizontal gene transfer such as conjugation and transformation. We found that multidrug-

resistance evolution remained possible even at the lowest probability we considered here of a

transducing phage carrying an AMR gene being released instead of a lytic phage during burst (1

transducing phage per 10^9 lytic phage). Additionally, in this work we assumed that transduction rates

are constant, but previous research has shown that sub-MIC antibiotic exposure can lead to an

increase in transducing phage, but not lytic ones [51]. Hence, our results may still underestimate the

relative impact of transduction versus predation by phage when antibiotics are present. However, our

findings are encouraging for conditions under which AMR evolution is limited, since even with a high

risk of transduction (1 transducing phage per 10^6 lytic phage), if phage are only introduced to an

environment more than 10h after antibiotics, multidrug-resistant bacteria do not reach detectable

levels before single-resistant strains are eradicated (Figure 6c).

To the best of our knowledge, our model is the first to consider the potential consequences of

transduction in a system where concentrations of bacteria, phage, and antibiotic are similar to those

we may see during phage therapy. Previously our work only considered bacteria and phage capable

of generalised transduction in the absence of antibiotics [28], whilst others have explored bacteria

phage and antibiotics without including generalised transduction [6–8,10]. We echo the conclusions

from these previous studies which highlighted that the timing of antibiotics and phage introduction

during phage therapy can affect the rate at which bacteria are cleared, but extend these to show that

timing may also impact the risk for multidrug-resistant bacteria to be generated. Here, we suggest

that, although both sequential treatments can ultimately lead to bacterial eradication, the timing leads

to a trade-off between a slower clearing rate of bacteria (if antibiotics are added before phage), and

a higher risk of multidrug-resistance evolution (if phage are added before antibiotics). Future studies

and clinical trials of phage therapy should investigate varying timings of phage and antibiotics, instead

of only investigating their simultaneous application, and consider the risk of transduction during

treatment.

In any case, our ability to measure the impact of transduction as a driver of AMR evolution *in vivo* is

currently limited since individuals are not routinely screened for phage. A first step to measure this

despite the limitation may be to investigate evidence for within-patient changes in the resistance

profile of *S. aureus* isolates, as these would likely be caused by transduction [23]. In the case of

antibiotic treatment, the natural presence of phage capable of transduction may explain instances of

treatment failure, if these generate multidrug-resistant strains which are then selected for by the

antibiotics. Future studies monitoring therapeutic outcomes of antibacterial treatment in patients

where phage are also detected will be essential to better understand how our findings translate to *in*

vivo settings.

630   Limitations

The major limitation of our work is the deterministic nature of our model. While it does not account

for stochastic events which would play a large role when bacterial numbers are low, the deterministic

model is useful for analysis purposes, as it represents the average scenario we would observe. In

reality, we would likely see either bacterial clearance or unexpected increases at low numbers of

bacteria. We only generated model predictions for up to 48h, as our parameter values were obtained

using data from experiments over 24h. Beyond this time, bacteria and phage may be affected

negatively due to resource depletion, depending on the environment.

Our model does not include some dynamics which may be present *in vivo*, as we do not currently have

robust data available to parameterise these features, and would instead have had to rely on

assumptions or previously estimated parameter values from different settings. Notably, we have not

included the effect of the immune system, which may limit the number of multidrug-resistant bacteria

generated as it could suppress both bacteria and phage populations *in vivo* [52,53]. If the model was

extended to include the immune system, it would also have to consider potential detrimental effects

of large doses of phage and antibiotics, which would restrict these concentrations to prevent side

effects *in vivo* [54,55].

In addition, we assume that all the bacteria in our environment are equally susceptible to phage

infection, and have not considered the possibility for any further evolution. This includes adaptation

mutation allowing multidrug-resistant bacteria to overcome any fitness cost, as well as resistance to

phage. In our own previous *in vitro* experiments using these *S. aureus* strains and phage, we did not

detect evidence of bacterial resistance to phage appearing within 24h [28]. However, such resistance

may arise over longer periods of time [56,57]. The threat represent by phage resistance evolution is

clear for multidrug-resistant bacteria, as these would then no longer be affected by either antibiotics

nor phage. On the other hand, if single antibiotic-resistant strains developed resistance to phage in

our system, they would still be removed by the other antibiotic they are susceptible to, but may then

be more likely to benefit from transduction and become multidrug-resistant, depending on the phage

resistance mechanism. If resistance occurs by targeting and neutralising phage DNA in the cell (e.g.

CRISPR-Cas [57]), bacterial DNA injected by a transducing phage would not be targeted, hence

protecting the resistant bacteria from phage lytic infection whilst still allowing transduction. However,

if resistance occurs by preventing of phage binding (e.g. surface receptor modification [57], although

there is limited evidence of this type of phage resistance in *S. aureus* [58]), then the bacteria would no

longer benefit from transduction. In any case, the risk of phage resistance could be mitigated by rapid

killing of bacteria, before phage resistance arises, highlighting the importance of measuring how fast

bacterial eradication occurs.

Although we varied the concentration of antibiotics in our results, we have consistently added

erythromycin and tetracycline in equal amounts. Our model would allow us to change this, yet we

have chosen not to for simplicity and because the antibacterial effect curves look similar for these two

antibiotics in our setting (Figure S1). However, for other antibiotics it may be necessary to revisit this

assumption and investigate concentrations which may better reflect those to which bacteria are

exposed to in the environment or during antibacterial treatment.

Generalisability

Our model is extensively parameterised using data from a single phage and three *S. aureus* strains,

making it a robust tool to study the dynamics of these organisms, as it relies on a minimum number

of assumptions [28]. However, the parameters we have estimated (adsorption rate, burst size, latent

period and transduction probability) will likely vary depending on the phage, bacteria, and

environment studied. Our sensitivity analysis shows that the model outputs are reasonable with

alternative parameter values, predicting for example that phage with a higher predation capacity

(higher adsorption rate or burst size, or lower latent period) would clear more bacteria within 48h,

and reduce the maximum number of multidrug-resistant bacteria generated. This model has been

developed as part of an interdisciplinary project alongside *in vitro* experiments, hence it could be easily

re-parameterised using data for other strains of bacteria and phage showing similar dynamics of lysis

and generalised transduction. The structure of the model is generalisable to other systems of

generalised transducing phage and bacteria, as it captures the relevant biological characteristics of

phage predation and generalised transduction [28].

The unique dynamics of phage, bacteria, and antibiotics

We suggest that transduction and the effect of antibiotics should be considered in the context of the previously described unique dynamics of phage and bacteria. These imply that phage must first reach a certain concentration (previously referred to as "inundation threshold") before they can offset bacterial growth and decrease the bacterial population, and bacteria must first reach a certain concentration ("proliferation threshold") before the phage population can increase [10]. Generalised transduction and antibiotics affect the size of the bacterial population, and therefore how phage interact with bacteria to clear them and generate multidrug-resistant bacteria. Thus, multidrug-resistant bacteria are able to increase in our model if phage are initially present at a concentration lower than the inundation threshold (Figure 5c). This also explains our counterintuitive observation that higher decay rates may lead to less bacteria remaining after 48h (Figure 6d), as this would allow bacteria to reach the proliferation threshold sooner, and therefore allow phage to increase up to the inundation threshold sooner (Figure S7). Importantly, our results similarly suggest that higher decay rates for antibiotics present alongside phage would reduce bacteria remaining after 48h, at the cost of a higher peak concentration of multidrug-resistant bacteria, since this decay would allow bacteria to reach the proliferation threshold sooner (Figure 6c, Figure S7). This knowledge may be further useful in the context of phage therapy, to select the antibiotics that will be given alongside phage [50].

Conclusions

Our results demonstrate the complex synergy between phage and antibiotics to kill bacteria and drive the evolution of AMR. We suggest this synergy leads to a trade-off between a slower clearing rate of bacteria (if antibiotics are added before phage), and a higher risk of multidrug-resistance evolution (if phage are added before antibiotics), further exacerbated by low concentrations of either phage or antibiotics. Interdisciplinary frameworks such as ours combining *in vitro* data and mathematical models are key to understanding both fundamental AMR evolution, and new interventions like phage

708   therapy or screening for phage in patients. Our conclusions form hypotheses to guide future

709   experimental and clinical work, notably for studies of phage therapy which should consider the risk

710   for multidrug-resistance evolution by transduction, and investigate varying timings and

711   concentrations of phage and antibiotics instead of only their simultaneous use.

ACKNOWLEDGMENTS

The authors would like to thank Caroline Memmi for helpful discussions on antibiotic bioavailability and *in vivo* antibiotic concentrations, as well as Joseph Standing and his research group for advice regarding antibiotic pharmacodynamics modelling.

FUNDING

Q.J.L was supported by a studentship from the Medical Research Council Intercollegiate Doctoral Training Program (MR/N013638/1). G.M.K was supported by a Skills Development Fellowship from the Medical Research Council (MR/P014658/1). G.M.K and J.A.L were supported by a Research Grant from the Medical Research Council (MR/P028322/1). The funders played no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

1.   Clokie MRJ, Millard AD, Letarov AV, Heaphy S. Phages in nature. Bacteriophage. 2011;1: 31–45. doi:10.4161/bact.1.1.14942

2.   Batinovic S, Wassef F, Knowler SA, Rice DTF, Stanton CR, Rose J, et al. Bacteriophages in Natural and Artificial Environments. Pathogens. 2019;8: 100. doi:10.3390/pathogens8030100

3.   Larsson DGJ. Pollution from drug manufacturing: review and perspectives. Philos Trans R Soc B Biol Sci. 2014;369: 20130571. doi:10.1098/rstb.2013.0571

4.   Brives C, Pourraz J. Phage therapy as a potential solution in the fight against AMR: obstacles and possible futures. Palgrave Commun. 2020;6: 1–11. doi:10.1057/s41599-020-0478-4

5.   Li X, He Y, Wang Z, Wei J, Hu T, Si J, et al. A combination therapy of Phages and Antibiotics: Two is better than one. Int J Biol Sci. 2021;17: 3573–3582. doi:10.7150/ijbs.60551

6.   Berryhill BA, Huseby DL, McCall IC, Hughes D, Levin BR. Evaluating the potential efficacy and limitations of a phage for joint antibiotic and phage therapy of Staphylococcus aureus infections. Proc Natl Acad Sci. 2021;118. doi:10.1073/pnas.2008007118

7.   Lopes A, Pereira C, Almeida A. Sequential Combined Effect of Phages and Antibiotics on the
Inactivation   of   Escherichia   coli.   Microorganisms.   2018;6:   125.
doi:10.3390/microorganisms6040125

8.   Torres-Barceló C, Arias-Sánchez FI, Vasse M, Ramsayer J, Kaltz O, Hochberg ME. A Window of
Opportunity to Control the Bacterial Pathogen Pseudomonas aeruginosa Combining Antibiotics
and Phages. PLoS ONE. 2014;9: e106628. doi:10.1371/journal.pone.0106628

9.   Abedon ST. Phage-Antibiotic Combination Treatments: Antagonistic Impacts of Antibiotics on
the   Pharmacodynamics   of   Phage   Therapy?   Antibiotics.   2019;8:   182.
doi:10.3390/antibiotics8040182

10. Payne RJH, Jansen VAA. Understanding Bacteriophage Therapy as a Density-dependent Kinetic
Process. J Theor Biol. 2001;208: 37–48. doi:10.1006/jtbi.2000.2198

11. Wang L, Tkhilaishvili T, Bernal Andres B, Trampuz A, Gonzalez Moreno M. Bacteriophage–
antibiotic combinations against ciprofloxacin/ceftriaxone-resistant Escherichia coli in vitro and
in an experimental Galleria mellonella model. Int J Antimicrob Agents. 2020;56: 106200.
doi:10.1016/j.ijantimicag.2020.106200

12. Kumaran D, Taha M, Yi Q, Ramirez-Arcos S, Diallo J-S, Carli A, et al. Does Treatment Order
Matter? Investigating the Ability of Bacteriophage to Augment Antibiotic Activity against
Staphylococcus aureus Biofilms. Front Microbiol. 2018;9: 127. doi:10.3389/fmicb.2018.00127

13. Chaudhry WN, Concepción-Acevedo J, Park T, Andleeb S, Bull JJ, Levin BR. Synergy and Order
Effects of Antibiotics and Phages in Killing Pseudomonas aeruginosa Biofilms. PLoS ONE.
2017;12: e0168615. doi:10.1371/journal.pone.0168615

14. von Wintersdorff CJH, Penders J, van Niekerk JM, Mills ND, Majumder S, van Alphen LB, et al.
Dissemination of Antimicrobial Resistance in Microbial Ecosystems through Horizontal Gene
Transfer. Front Microbiol. 2016;7. doi:10.3389/fmicb.2016.00173

15. Fernández L, Rodríguez A, García P. Phage or foe: an insight into the impact of viral predation on
microbial communities. ISME J. 2018;12: 1171–1179. doi:10.1038/s41396-018-0049-5

16. Balcázar JL. How do bacteriophages promote antibiotic resistance in the environment? Clin
Microbiol Infect. 2018;24: 447–449. doi:10.1016/j.cmi.2017.10.010

17. Haaber J, Penadés JR, Ingmer H. Transfer of Antibiotic Resistance in Staphylococcus aureus.
Trends Microbiol. 2017;25: 893–905. doi:10.1016/j.tim.2017.05.011

18. Leclerc QJ, Lindsay JA, Knight GM. Mathematical modelling to study the horizontal transfer of
antimicrobial resistance genes in bacteria: current state of the field and recommendations. J R
Soc Interface. 2019;16: 20190260. doi:10.1098/rsif.2019.0260

19. Verheust C, Pauwels K, Mahillon J, Helinski DR, Herman P. Contained use of Bacteriophages: Risk
Assessment   and   Biosafety   Recommendations.   Appl   Biosaf.   2010;15:   32–44.
doi:10.1177/153567601001500106

20. Jassim SAA, Limoges RG. Natural solution to antibiotic resistance: bacteriophages 'The Living
Drugs.' World J Microbiol Biotechnol. 2014;30: 2153–2170. doi:10.1007/s11274-014-1655-7

21.   Chen J, Quiles-Puchalt N, Chiang YN, Bacigalupe R, Fillol-Salom A, Chee MSJ, et al. Genome
hypermobility   by   lateral   transduction.   Science.   2018;362:   207–212.
doi:10.1126/science.aat5867

22.   den Heijer CDJ, van Bijnen EME, Paget WJ, Pringle M, Goossens H, Bruggeman CA, et al.
Prevalence and resistance of commensal Staphylococcus aureus, including meticillin-resistant S
aureus, in nine European countries: a cross-sectional study. Lancet Infect Dis. 2013;13: 409–415.
doi:10.1016/S1473-3099(13)70036-7

23.   Lindsay JA. Staphylococcus aureus genomics and the impact of horizontal gene transfer. Int J
Med Microbiol. 2014;304: 103–109. doi:10.1016/j.ijmm.2013.11.010

24.   Stanczak-Mrozek KI, Manne A, Knight GM, Gould K, Witney AA, Lindsay JA. Within-host diversity
of   MRSA   antimicrobial   resistances.   J   Antimicrob   Chemother.   2015;70:   2191–2198.
doi:10.1093/jac/dkv119

25.   Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery,
research, and development of new antibiotics: the WHO priority list of antibiotic-resistant
bacteria   and   tuberculosis.   Lancet   Infect   Dis.   2018;18:   318–327.   doi:10.1016/S1473-
3099(17)30753-3

26.   Rice LB. Federal Funding for the Study of Antimicrobial Resistance in Nosocomial Pathogens: No
ESKAPE. J Infect Dis. 2008;197: 1079–1081. doi:10.1086/533452

27.   Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al. Attributable
deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in
the EU and the European Economic Area in 2015: a population-level modelling analysis. Lancet
Infect Dis. 2019;19: 56–66. doi:10.1016/S1473-3099(18)30605-4

28.   Leclerc QJ, Wildfire J, Gupta A, Lindsay JA, Knight GM. Growth-Dependent Predation and
Generalized Transduction of Antimicrobial Resistance by Bacteriophage. mSystems. 2022;0:
e00135-22. doi:10.1128/msystems.00135-22

29.   McCarthy AJ, Loeffler A, Witney AA, Gould KA, Lloyd DH, Lindsay JA. Extensive Horizontal Gene
Transfer during Staphylococcus aureus Co-colonization In Vivo. Genome Biol Evol. 2014;6: 2697–
2708. doi:10.1093/gbe/evu214

30.   Fish R, Kutter E, Wheat G, Blasdel B, Kutateladze M, Kuhl S. Bacteriophage treatment of
intransigent diabetic toe ulcers: a case series. J Wound Care. 2016;25: S27–S33.
doi:10.12968/jowc.2016.25.Sup7.S27

31.   Lehman SM, Mearns G, Rankin D, Cole RA, Smrekar F, Branston SD, et al. Design and Preclinical
Development of a Phage Product for the Treatment of Antibiotic-Resistant Staphylococcus
aureus Infections. Viruses. 2019;11: 88. doi:10.3390/v11010088

32.   Ngassam-Tchamba C, Duprez JN, Fergestad M, De Visscher A, L'Abee-Lund T, De Vliegher S, et
al. In vitro and in vivo assessment of phage therapy against Staphylococcus aureus causing
bovine mastitis. J Glob Antimicrob Resist. 2020;22: 762–770. doi:10.1016/j.jgar.2020.06.020

33.   Ooi ML, Drilling AJ, Morales S, Fong S, Moraitis S, Macias-Valle L, et al. Safety and Tolerability of
Bacteriophage Therapy for Chronic Rhinosinusitis Due to Staphylococcus aureus. JAMA
Otolaryngol Neck Surg. 2019;145: 723–729. doi:10.1001/jamaoto.2019.1191

34.   Petrovic Fabijan A, Lin RCY, Ho J, Maddocks S, Ben Zakour NL, Iredell JR. Safety of bacteriophage
therapy in severe Staphylococcus aureus infection. Nat Microbiol. 2020;5: 465–472.
doi:10.1038/s41564-019-0634-z

35.   Raj JRM, Karunasagar I. Phages amid antimicrobial resistance. Crit Rev Microbiol. 2019;45: 701–
711. doi:10.1080/1040841X.2019.1691973

36.   Hassan AY, Lin JT, Ricker N, Anany H. The Age of Phage: Friend or Foe in the New Dawn of
Therapeutic   and   Biocontrol   Applications?   Pharmaceuticals.   2021;14:   199.
doi:10.3390/ph14030199

37.   Rodríguez-Rubio L, Jofre J, Muniesa M. Is Genetic Mobilization Considered When Using
Bacteriophages   in   Antimicrobial   Therapy?   Antibiotics.   2017;6:   32.
doi:10.3390/antibiotics6040032

38.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A Genetic Resource for Rapid
and Comprehensive Phenotype Screening of Nonessential Staphylococcus aureus Genes. mBio.
2013;4. doi:10.1128/mBio.00537-12

39.   Mouton JW, Vinks AA. Pharmacokinetic/Pharmacodynamic Modelling of Antibacterials In Vitro
and In Vivo Using Bacterial Growth and Kill Kinetics. Clin Pharmacokinet. 2005;44: 201–210.
doi:10.2165/00003088-200544020-00005

40.   Wayne P, Clinical and Laboratory Standards Institute. CSLI document M07-A9. 9th ed. Methods
for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved
Standard. 9th ed. CSLI; 2012.

41.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing; 2020. Available: https://www.R-project.org/

42.   Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. Pharmacodynamic Functions:
a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. Antimicrob Agents
Chemother. 2004;48: 3670–3676. doi:10.1128/AAC.48.10.3670-3676.2004

43.   Weld RJ, Butts C, Heinemann JA. Models of phage growth and their applicability to phage
therapy. J Theor Biol. 2004;227: 1–11.

44.   Schrag SJ, Mittler JE. Host-Parasite Coexistence: The Role of Spatial Refuges in Stabilizing
Bacteria-Phage Interactions. Am Nat. 1996;148: 348–377. doi:10.1086/285929

45.   Gelman A, Rubin DB. Inference from Iterative Simulation Using Multiple Sequences. Stat Sci.
1992;7: 457–472.

46.   Gefen O, Fridman O, Ronin I, Balaban NQ. Direct observation of single stationary-phase bacteria
reveals a surprisingly long period of constant protein production activity. Proc Natl Acad Sci.
2014;111: 556–561. doi:10.1073/pnas.1314114111

47.   Krasniqi S, Matzneller P, Kinzig M, Sörgel F, Hüttner S, Lackner E, et al. Blood, Tissue, and
Intracellular Concentrations of Erythromycin and Its Metabolite Anhydroerythromycin during
and   after   Therapy.   Antimicrob   Agents   Chemother.   2012;56:   1059–1064.
doi:10.1128/AAC.05490-11

48.   Somogyi AA, Bochner F, Hetzel D, Williams DB. Evaluation of the Intestinal Absorption of
853   Erythromycin in Man: Absolute Bioavailability and Comparison with Enteric Coated
854   Erythromycin. Pharm Res. 1995;12: 149–154. doi:10.1023/A:1016215510223

855   49.   Heldal M, Bratbak G. Production and decay of viruses in aquatic environments. Mar Ecol Prog
856   Ser. 1991;72: 205–212.

857   50.   Levin BR, Udekwu KI. Population Dynamics of Antibiotic Treatment: a Mathematical Model and
858   Hypotheses for Time-Kill and Continuous-Culture Experiments. Antimicrob Agents Chemother.
859   2010;54: 3414–3426. doi:10.1128/AAC.00381-10

860   51.   Stanczak-Mrozek KI, Laing KG, Lindsay JA. Resistance gene transfer: induction of transducing
861   phage by sub-inhibitory concentrations of antimicrobials is not correlated to induction of lytic
862   phage. J Antimicrob Chemother. 2017;72: 1624–1631. doi:10.1093/jac/dkx056

863   52.   Banuelos S, Gulbudak H, Horn MA, Huang Q, Nandi A, Ryu H, et al. Investigating the Impact of
864   Combination Phage and Antibiotic Therapy: A Modeling Study. In: Segal R, Shtylla B, Sindi S,
865   editors. Using Mathematics to Understand Biological Complexity: From Cells to Populations.
866   Cham: Springer International Publishing; 2021. pp. 111–134. doi:10.1007/978-3-030-57129-0_6

867   53.   Rodriguez-Gonzalez RA, Leung CY, Chan BK, Turner PE, Weitz JS. Quantitative Models of Phage-
868   Antibiotic Combination Therapy. mSystems. 2020;5: e00756-19. doi:10.1128/mSystems.00756-
869   19

870   54.   Liu D, Van Belleghem JD, de Vries CR, Burgener E, Chen Q, Manasherob R, et al. The Safety and
871   Toxicity of Phage Therapy: A Review of Animal and Clinical Studies. Viruses. 2021;13: 1268.
872   doi:10.3390/v13071268

873   55.   Imani S, Buscher H, Marriott D, Gentili S, Sandaradura I. Too much of a good thing: a
874   retrospective study of β-lactam concentration–toxicity relationships. J Antimicrob Chemother.
875   2017;72: 2891–2897. doi:10.1093/jac/dkx209

876   56.   Koskella B, Brockhurst MA. Bacteria–phage coevolution as a driver of ecological and evolutionary
877   processes in microbial communities. FEMS Microbiol Rev. 2014;38: 916–931. doi:10.1111/1574-
878   6976.12072

879   57.   Hampton HG, Watson BNJ, Fineran PC. The arms race between bacteria and their phage foes.
880   Nature. 2020;577: 327–336. doi:10.1038/s41586-019-1894-8

881   58.   Dalen R van, Peschel A, Sorge NM van. Wall Teichoic Acid in Staphylococcus aureus Host
882   Interaction. Trends Microbiol. 2020;28: 985–998. doi:10.1016/j.tim.2020.05.017