Nematode histone H2A variant evolution reveals diverse histories of retention and

loss and evidence for conserved core-like variants

Swadha Singh[1], Diana Chu[2*], Scott Roy[1,2*]

[1]Quantitative & Systems Biology, University of California, Merced, California, USA.

[2]Department of Biology, San Francisco State University, California, USA.

[*]Corresponding authors: Emails; sroy5@ucmerced.edu; chud@sfsu.edu

ABSTRACT

Histone variants are paralogs that replace canonical histones in nucleosomes, often

imparting novel functions. Despite their importance, how histone variants arise and evolve

is poorly understood. Reconstruction of histone protein evolution is challenging due to high

amino acid conservation and large differences in evolutionary rates across gene lineages

and sites. Here we combined amino acid sequences and intron position data from 108

nematode genomes to trace the evolutionary histories of the three H2A variants found in

Caenorhabditis elegans: the ancient H2A.Z[HTZ-1], the sperm-specific HTAS-1, and HIS-35,

which differs from canonical H2A by a single glycine-to-alanine C-terminal change. We

find disparate evolutionary histories. Although the H2A.Z[HTZ-1] protein is highly conserved,

its gene exhibits recurrent intron gain and loss. This pattern suggests that it is intron

presence, rather than specific intron sequences or positions, that may be important to

H2A.Z functionality. In contrast, for HTAS-1 and HIS-35, we find variant-specific intron

positions that are conserved across species. HIS-35 arose in the ancestor of *Caenorhabditis*

and its sister group, including the genus *Diploscapter*, while the sperm-specific variant HTAS-1 arose more recently in the ancestor of a subset of *Caenorhabditis* species. HIS-35 exhibits gene retention in some descendent lineages but also recurrent gene loss in others, suggesting that histone variant use or functionality is highly flexible in this case. We also find that the single amino acid differentiating HIS-35 from core H2A is ancestral and common across canonical *Caenorhabditis* H2A sequences and identify one nematode species that bear identical HIS-35 and canonical H2A proteins, findings that are not predicted from the hypothesis that HIS-35 has a distinct function. Instead, we speculate that HIS-35 enables H2A expression across the cell cycle or in distinct tissues; genes encoding such partially-redundant functions may be advantageous yet relatively replaceable over evolutionary times, consistent with the patchwork pattern of retention and loss of both genes. Our study shows the evolutionary trajectory for histone H2A variants with distinct functions and the utility of intron positions for reconstructing the evolutionary history of gene families, particularly those undergoing idiosyncratic sequence evolution.

INTRODUCTION

All characterized eukaryotic cells compact their DNA into nucleosomes, which consist of an octameric histone complex comprised of two copies each of the core histone proteins H2A, H2B, H3, and H4, and which wrap around ~147 bps of DNA (1-4). In many species, genes encoding canonical core histones (also referred to as 'replication-dependent histones') do not contain introns and are organized into multiple gene copies and in a tandem-repeat structure, which facilitates their rapid, coordinated expression (4-7). The

expression of canonical histones is tightly coupled with the S-phase of the cell cycle because of the critical need for the large bulk of histone proteins required to package and compact the newly synthesized DNA (8-10).

Histones have evolved variant forms that further regulate chromatin compaction and affect processes like transcription, DNA repair, and development (11-20). Such histone variants are often considered to be a part of the 'histone code' and control distinct sets of genes during specific times and tissues in both animals and plants (11, 12, 21-23). Among all the core histones, H2A is the fastest evolving histone, showing the most diversity in the variant types and expression patterns (24, 25). For example, the highly conserved H2A.Z is present in every eukaryotic species and ubiquitously expressed. It has been shown in different species to play a variety of roles including transcriptional regulation, heterochromatin boundaries, DNA repair, DNA replication, and dosage compensation (26-31). In contrast, H2A.X has arisen independently in several lineages, with functions in DNA damage response and transcription (18, 31, 32), while macroH2A, which is involved in X inactivation and stress response, is present only in some lineages, making its origins unclear (3, 4, 24, 33-36). Finally, short histone variants, such as H2A.B, H2A.P, H2A.L, and H2A.Q, are expressed in testis (except H2A.B which is also expressed in the brain cells) (25, 37-40). In short, a great wealth of histone H2A variants are observed, ranging from those shared across all eukaryotes to others that are species-specific (3, 4, 18, 19, 35). H2A variants also have distinct functions in a broad range of processes and can be ubiquitously expressed or expressed only in certain tissues (11, 25, 41, 42). However, with

the exception of the well-studied ancient variant H2A.Z, how histone H2A variants have

arisen and evolved remains unknown.

Clues to how histone variants differ may stem from the distinct gene structures and

expression patterns compared to canonical histones. Unlike canonical histone genes,

variant histone gene expression is not restricted to S phase expression and can be tissue-

specific. Most histone variants are typically found in a single copy in the genome and

contain introns in their pre-mRNAs (5-7, 43, 44). Notably, the roles of these introns in

variant function remain poorly studied, including whether the introns themselves impart

novel functionality. For instance, sequences found within introns of other genes can

regulate gene expression (43, 45-48). Introns can also allow for alternative splicing, which

is common in animals and plays important role in development and disease (49-52). Introns

have been used in other studies to determine the evolutionary trajectory of other protein

families (53-57). Thus, the presence of introns in histone variants may provide useful clues

to exploring the histone evolution and function across closely related species.

Core histone functions are expected to be highly conserved across eukaryotes, given

their central roles in ensuring DNA packaging and protection (4, 61). Thus, observed

protein changes in core histones are expected to be largely neutral with respect to protein

function. On the other hand, amino acid differences between variant histones and core

histones are thought to generally lead to functional differences (3, 4, 62). Indeed, protein

sequence differences between well-studied variants and their core homologs have been

shown to affect chromatin structure and function. The histone H3.3 variants, for example,

which differ from each other by 4-5 amino acids, play distinct roles in transcriptional

activation, chromatin remodeling, heterochromatin formation, and development (14, 16,

63-66). On the other hand, the functional significance of protein differences between

variant histones and their core counterparts in some cases is unclear. For instance, the

protein sequence of *C. elegans* HIS-35 differs by a single 'A' instead of a 'G' at the 124th

position from the S-phase *C. elegans* H2A, the functional significance of which is

unknown.

In this study, we focus on the evolutionary histories of the three very different H2A

variants found in *Caenorhabditis elegans:* H2A.Z^{HTZ-1}, HIS-35, and HTAS-1 (Figure 1).

H2A.Z^{HTZ-1} is an ortholog of the evolutionarily conserved variant, H2A.Z. HIS-35

intriguingly differs from core H2A by a single amino acid difference. HTAS-1 shows a

greater divergence from core H2A, particularly at the highly divergent C and N termini,

and appears to be expressed only in sperm and has only been reported in *C. elegans* to date

(67). The presence of a discrete number of H2A variants with distinct functions and/or

expression patterns in combination with the availability of sequence data across a large

number of nematode species allows a unique opportunity to track the evolutionary

trajectory of this histone variant family.

RESULTS

Sequence-based phylogenetic methods fail to reconstruct H2A gene family evolution

Taking advantage of vast sequencing (68-70) efforts across nematode species, we

used BLAST searches across 108 reported nematode genomes to identify all annotated

copies of H2A and H2A-related gene variants across the diversity of nematodes. After

filtering and collapsing identical proteins, we were left with 408 unique sequences. We

then used standard phylogenetic methods to reconstruct the evolutionary history of these

sequences (Figure S1, intronless (thus putative core or canonical) H2As are highlighted in

yellow, HTZ-1 homologs are highlighted in pink, putative HIS-35 homologs are

highlighted in blue, and putative HTAS-1 homologs are highlighted in gray). However,

scrutiny of the recovered phylogenetic tree revealed several bizarre findings. For instance,

core H2A proteins were grouped as separate clades that included very deeply-diverged

nematode sequences; on the other hand, species-specific variants were often found grouped

far from core proteins from the same or related species. Perhaps the clearest case can be

shown by performing phylogenetic reconstruction on core H2A orthologs and on the HIS-

35 orthologs identified based on intron positions (described in detail below). Here, we

expect distinct clades of H2A and HIS-35 sequences, yet we recover no such separate

clades (Figure S2, HIS-35 is highlighted).

Some of these anomalies are as expected by errors in phylogenetic reconstruction

due to model misspecification (the phenomenon in which differences between the assumed

model of sequence evolution and the actual evolutionary process led to errors in

phylogenetic reconstruction) (71). The possibility of model misspecification is increased

in the case of histone genes by several factors: extreme differences in rates across sites

(some sites show conservation across eukaryotes, others are variable within

Caenorhabditis); large relative differences in rates across gene lineages (the extremely

slow evolution of core proteins, but substantially more change in some variants); and the

generally small number of total sites given the short length of histone proteins.

Intron position and phase distinguish the three H2A variants of *C. elegans*.

As an alternative approach, we used another potential source of phylogenetic

information: the position of the spliceosomal introns that disrupt nuclear genes, including

variant histones. Intron positions can be conserved over very long times in orthologous

genes (53-56, 58, 59). Whereas early work considered the possibility of intron 'sliding',

in which an intron would migrate a few base pairs along a gene, recent work has shown

that intron sliding is a very rare occurrence, and thus that intron positions are very often

conserved over very long times (54, 59). For example, in *Theileria* apicomplexans, 99.7%

of intron positions are conserved between *T. parva* and *T. annulata*, diverging roughly 82

million years ago (53). In mammals, 99.9% of intron positions are conserved between

humans and dogs, diverging around 100 million years ago (55).

We began our study of intron-exon structures in H2A variants by obtaining intron-

exon structures for all H2A gene family members and performing alignments to determine

intron position sharing across genes. We first aligned the three main *C. elegans* H2A

variants (Figure 1), HIS-35, HTAS-1, and H2A.Z^{HTZ-1}, each of which has a single intron

position (highlighted boxes in Figure 1). Scrutiny of the intron positions showed that the

three genes contain introns at different positions, differing in the codon that they interrupt

or and/or in the phase at which they disrupt the codon. Interestingly, the intron position in

HIS-35 falls very near to that found in H2A.Z^{HTZ-1}. However, these introns are unlikely to

represent a shared intron, for two reasons. First, intron positions rarely slide between

phases (54, 59). Second, HIS-35's near identity to H2A (see below) strongly suggests that

it more likely evolved from intronless H2A by gene duplication and intron gain, and not

from H2A.Z^{HTZ-1}. Thus, the differences in the intron positions between the three *C. elegans*

variants suggest that the three variants evolved independently as different duplicates of

core H2A, rather than from each other.

Intron position and sequence evidence indicates the origin of HTAS-1 within

***Caenorhabditis* and subsequent retention and loss**

The *C. elegans* sperm-specific H2A variant, HTAS-1, contains a single intron between

codons 26 and 27 ("phase 0"; Figure 1, highlighted with yellow). Alignment across all the

H2A variants of 108 nematodes revealed 16 genes that share an intron at the exact

homologous position as *C. elegans* HTAS-1 (Figure 2) from species falling within a single

subclade of 22 species (represented by the red hash mark on the tree branch in Figure 2)

within the *Caenorhabditis* genus, suggesting an origin of this intron position within the

common ancestor of these species. Scrutiny of the sequence gene tree (Figure S1,

highlighted in gray, IP_152) revealed that this same set of genes (i.e., those sharing the

intron) appears together as a clade, reinforcing the notion that the genes sharing the HTAS-

1 intron position represent a set of orthologs. We next sought to identify potential HTAS-

1 orthologs in additional *Caenorhabditis* species, both the 6/22 within the HTAS-1-

containing subclade that lack intron-containing HTAS-1 candidates (marked by purple

hash on the tree branch in Figure 2) as well as the 10 species outside this subclade. No

genes from any such species were found grouping with the putative HTAS-1 ortholog

clade; thus, overall, we couldn't find any other genes that are candidates for HTAS-1

orthologs. In total, the data is consistent with the origin of HTAS-1 and its gene-specific

intron occurring in the same ancestor of a subset of *Caenorhabditis* species.

As noted above, in six species within the HTAS-1-containing clade of

Caenorhabditis (*C. wallacei*, *C. brenneri* and *C.inopinata* (*sp34*), *C. kamaaina*, *C.*

panamensis (*sp28*), *and C. japonica*), we could not find annotated genes with either an

intron position at the HTAS-1-specific position (Figure 2, marked with purple hashes on

the tree branch) or other genes that grouped with the putative HTAS-1 orthologs (Figure

S1). However, we were unable to distinguish whether these absences represent true

biological loss or failure of gene annotation, which is particularly possible given that

histone genes' short lengths increase the possibility they would be missed by standard

annotation methods. Indeed, we identified one species harboring a likely HTAS-1

sequence for which only the second exon was annotated (Figure S1), however, TBLASTN

searches against the genome failed to confidently detect this gene. This is likely because,

given the much higher rate of evolution of HTAS-1, HTAS-1 from one species often shows

a greater degree of core H2A from another species than to its HTAS-1 ortholog;

consequently, TBLASTN searches using HTAS-1 from one species typically give many

hits, with the likely HTAS-1 not being within the top dozen hits, making it difficult to

distinguish gene absence from the failure of annotation. In total, then, the data is consistent

with a single origin of HTAS-1 and its characteristic intron position within the ancestor of

a subset of studied *Caenorhabditis* species, but we cannot determine with certainty whether

HTAS-1 has been lost in up to six independent lineages (Figure 2).

Intron position conservation suggests HIS-35 originated in the *Caenorhabditis-*

***Diploscapter* ancestor and has undergone subsequent gene retention and loss**

The variant HIS-35 of *C. elegans* has a phase zero intron between the 50th and the 51st

codon (Figure 1). Alignment across all H2A variants revealed 20 species with genes that

share this intron position (Figure 3, marked with a plus sign). In all cases, these genes

show very high sequence similarity with H2A at the protein level. These genes are from

species falling in the clades of *Caenorhabditis* and its sister genus *Diploscapter*,

representing 20/32 species within this clade, suggesting an origin of this intron position

within the common ancestor of *Diploscapter* and *Caenorhabditis* (Figure 3).

We sought to determine whether the 12 descendent species without a clear candidate

HIS-35 gene represent true losses or merely failures of gene annotation. TBLASTN and

BLASTN searches failed to recover likely HIS-35 orthologs absent from the annotation.

In contrast to the case of HTAS-1 above, we are confident in the ability of these methods

to detect HIS-35 orthologs, since we were able to detect the annotated HIS-35 ortholog by

identical methods (i.e., the positive control). For instance, using *C. japonica* HIS-35 as a

query, BLASTN searches revealed the (annotated) putative ortholog in *C. macrosperma*,

but did not reveal unannotated candidate orthologs in *C. nouroguensis, C. waitukubuli, C.*

panamensis, or *C. becei*, which are equally closely related to *C. japonica*. In addition,

whereas species lacking HTAS-1 appear to be randomly scattered across the HTAS-1-

containing clade, as expected by random gene annotation failures, species lacking HIS-35

within the HIS-35-containing clade group in subclades. This is as expected by true

biological loss and not by random annotation failures. Thus, we conclude that HIS-35 has

been lost some 5 times independently in different *Caenorhabditis* lineages.

Conservative protein evolution of HIS-35 and evidence for occasional gene conversion

with core H2A

219   HIS-35 provides a particularly interesting example for histone protein evolution. In

220 *C. elegans*, the protein sequence of HIS-35 differs by just one amino acid from the S-phase

221 H2A despite the substantial evolutionary time. Namely, HIS-35 has an "A", while H2A

222 has a "G" at position 124 of the amino acid sequence. If an "A" at this position is an

223 overriding change in the HIS-35 variant, then we could expect this change to encode a

224 different function from the canonical H2A.

225   However, when we looked at position 124 the canonical H2A sequences of all the

Caenorhabditis species we actually found out that the "A" is ancestral to canonical H2A

(Figure 3). This conservation of the "A" at position 124 suggests that HIS-35 likely has not

diverged in function from the ancestral H2A despite this variant being retained across many

Caenorhabditis species. We also found that the predicted protein sequences of HIS-35 and

H2A of species *C. kamaaina* are exactly the same (Figure 3). These findings are not as

expected if canonical H2A and HIS-35 proteins have different functions, but are as

expected if the proteins are functionally redundant.

The multiple copies of core histone genes are known to undergo so-called concerted

evolution, with sequences being transferred between gene copies by recombination (72-

75). We, therefore, wondered whether concerted evolution plays a role in the identical

protein sequence changes observed in the H2A and HIS-35 paralogs of some species.

Under concerted evolution, the two sequences undergoing concerted evolution are

homogenized (one overwrites the other). Consequently, the prediction is that such events

should lead the interconverting partners to the group together on a phylogenetic tree. To

search for evidence of concerted evolution, we reconstructed separate phylogenetic trees

of exon-1 and exon-2 for all H2A and HIS-35 sequences. Most of the reconstructed tree

largely reflected the species tree, suggesting against the possibility of widespread concerted

evolution (Figures S3 and S4). However, we did observe the grouping of the two gene

sequences for *Caenorhabditis sp21*, consistent with the concerted evolution of HIS-35 and

H2A in only this species. Concerted evolution of these genes is consistent with a lack of

functional differentiation, though admittedly the low rate of such events weakens the

strength of this argument.

The intron loss and gain in HTZ-1 show a dynamic evolutionary history

The ubiquitously expressed *C. elegans* H2A variant H2A.Z^{HTZ-1} is the ortholog of

H2A.Z, which is evolutionary conserved across all the eukaryotes (3, 76). *C. elegans*

H2A.Z^{HTZ-1} has an intron that splits the 57th codon at position 2 (Figure 1). Within the

alignment across all H2A variants, we searched for genes that share the single intron

position of *C. elegans* H2A.Z^{HTZ-1}, revealing 30 genes share this position (Figure 4, marked

with a plus sign; Figure S1). These are putative H2A.Z^{HTZ-1} orthologs; consistent with this

hypothesis, they grouped together on the tree. Unexpectedly, we found that 22 of these 30

putative H2A.Z^{HTZ-1} genes have two (or more in *Diploscapter*) introns in their genes, one

at position 57.2 and the other at position 111.1. Both introns have been repeatedly

individually lost in different lineages (including in the lineage leading to the single-intron

C. elegans H2A.Z^{HTZ-1} gene). Species including *C. elegans*, *C. tropicallis*, *C. sp32*, *C. afra*,

C. guadeloupensis, *C. virilis* have lost the second H2A.Z intron which is at position 111.1,

whereas *C. casteli* and *C. angaria* have lost their first intron. These results are consistent

with general results in protein-coding genes, wherein intron loss is common across the

Caenorhabditis phylogeny (77, 78). Nonetheless, the finding that general trends of intron

loss may equally apply to histone variant genes is important in understanding the functional

implications of variant introns.

Intron presence is a conspicuous difference between core and variant histones,

raising the question of the functional significance of variant histone introns: specifically,

are variant histone introns important for differentiating the specific functions of those

variants from their core paralogs? The finding of recurrent loss of introns from a variant

gene suggests that the specific positions or sequences of introns in a particular histone

variant gene may not be of particular importance. Nonetheless, it is of note that all observed

H2A.Z orthologs contain at least one intron, suggesting that the presence of at least one

intron, at whatever position, could be important for efficient expression of variant histone

genes, consistent with their expression through canonical gene expression pathways, in

which intron presence often promotes gene expression.

Discussion

Introns as sources of phylogenetic information

This study is among the first to leverage the information present in intron positions

to decipher the evolutionary history of histone variants (60). Previous studies have shown

intron position conservation among widely diverged eukaryotic species (53-55, 57-60). For

instance, intron positions are highly conserved between humans, mice, and fish (57). Thus,

intron positions contain a record of evolutionary history that can facilitate insights into

gene history. The utility of introns here is different for the different variants and allows

different questions to be addressed. The clearest case comes from HIS-35, in which nearly

complete evolutionary conservation of H2A and HIS-35 sequences leads to a lack of

phylogenetic signal within proteins. At the other end of the spectrum lies HTAS-1, which

shows much more rapid sequence evolution. However, here, the discrepancy between rates

between HTAS-1 and H2A, and likely between different sites in the two proteins (i.e., the

N and C termini are highly constrained in H2A but fast-evolving for HTAS-1) makes it

impossible to define a single evolutionary model across the gene family. As expected by

general long branch attraction consideration, this leads to fast-evolving HTAS-1

incorrectly grouping far away from *Caenorhabditis* core H2A sequences. This generally

undermines our confidence in sequence-based phylogenetic reconstruction of HTAS-1

genes. Ironically, intron-defined HTAS-1 orthologs do group as a clade, indicating that

sequence-based phylogenetic reconstruction was likely successful for this group; however,

without the intron position information, our general lack of confidence in the methods'

success for HTAS-1 would lead us to question this finding. Thus, for this case, even though

the sequence-based phylogenetic methods apparently correctly identified the HTAS-1

clade, the orthologous information from intron positions was necessary for us to be

confident in the sequence-based phylogeny. Importantly, intron positions were

indispensable in distinguishing the origins of HTAS-1. Because HTAS-1 arose in an

ancestor containing two genes encoding nearly identical proteins (H2A and HIS-35), it is

very difficult to determine whether H2A evolved from the preexisting variant HIS-35 or

de novo from H2A. The fact that all candidate HTAS-1 genes lack the HIS-35 intron

strongly suggests *de novo* evolution from H2A, not HIS-35.

Evidence from phylogenetic analysis for functional significance of recently-evolved histone

variants.

In addition to tracing the origins and subsequent history of gene loss and retention,

our results provide insights into the possible functions of histone H2A variants. For

example, HIS-35 differs by a single amino acid from the S-phase H2A ('A' instead of 'G'

at the 124th position). Given that characterized histone variants are thought to largely

represent functionally distinct proteins, one hypothesis is that this single difference

functionally differentiates HIS-35 protein from H2A. One way that a single amino acid

change could have outsize effects is through altering the landscape of posttranslational

modifications, which are key to histone function. For instance, this is the case with H3

variants, H3.1 and H3.2 which differ from one another by single amino acid and show

distinct patterns of expression and post-translational modifications (64, 65).

Given it is possible for a small amino acid difference to change function, we

approached the single H2A/HIS-35 difference with the hypothesis that the single difference

could lead to functionally different proteins. However, when we looked at the H2A

sequences of all the *Caenorhabditis* species, we found an 'A' at position 124 to be

ancestral. Considering the presence of an A at position 124 in other canonical H2As,

suggests the variant HIS-35 might have the same function as the canonical H2A. Lack of

functional differentiation is also consistent with the similarity of A and G amino acids. This

hypothesis is also supported by the case of *C. kamaanina*, in which the encoded HIS-35

and H2A protein sequences are exactly the same. While these results do not disprove the

hypothesis that H2A and HIS-35 encode proteins with important functional differences

(except in the case of *C. kamaanina*), we propose instead that HIS-35 protein's functional

importance lies in allowing for the expression of a protein with overlapping or redundant

functions to canonical H2A that is not restricted to S-phase, as is the case with canonical

H2A. This could allow expression in differentiated cells that do not undergo mitosis and

enable tissue-specific expression. Such a potential semi-redundancy could help to explain

the ambivalent phylogenetic pattern, in which retention of HIS-35 in most species suggests

functional importance whereas loss in 5 independent lineages suggests conditional

expendability. Interestingly, a similar pattern of lineage-specific loss has been observed

for H2B variants, which has also been interpreted as encoding functionally important but

partially redundant functions (79).

Function in reproduction for the sperm-specific variant HTAS-1

Sperm-specific proteins show generally elevated rates of evolution, consistent with

strong selection on sperm functions because of sperm competition (67, 80). The current

data show that this is decisively the case for the sperm-specific variant HTAS-1. We show

that *C. elegans* HTAS-1's greater divergence from core H2A is not because of differences

in an evolutionary age, but very much despite it: HTAS-1 is most divergent variant protein

despite being the most recent to diverge from core H2A, having since evolved at a rate

many times higher than any other H2A paralog. This high rate of evolution strongly

suggests that HTAS-1 may be adapted to play roles important for reproduction in some

species.

However, our data also have somewhat ambiguous implications for HTAS-1

function. On the one hand, though HTAS-1 has been maintained for long periods of time

in many lineages it may have also been lost in multiple independent lineages, as found for

HIS-35. While rapid evolution of HTAS-1 made it impossible to exclude the possibility

that apparent HTAS-1 losses actually represent gene annotation failures, the fact that

careful scrutiny revealed no gene annotation failures for HIS-35 weighs against the

possibility of annotation failure, suggesting real loss of HTAS-1.

On the other hand, the much larger degree of protein sequence difference between

HTAS-1 and H2A would seem to decrease the probability that HTAS-1 protein is

functionally identical to H2A protein particularly given the extended C & N terminus of

HTAS-1 which has previously been reported to play a vital role in DNA compaction,

chromosome segregation, and fertility (67). Moreover, the particular chromatin constraints

of sperm production raise the possibility that HTAS-1 proteins could encode distinct

functions relative to H2A proteins, for instance by leading to greater sperm DNA

compaction; however, it is also possible that a distinct H2A paralog is simply necessary to

ensure expression of H2A proteins well after germline mitosis in later stages of

spermatogenesis that undergo transcription, DNA recombination and repair, and

division. Thus, more study of the functional significance of HTAS-1 homologs in different

species is clearly needed to distinguish between these possibilities.

Concluding remarks

To summarize, these results show exceptions to previously reported patterns,

challenging sometimes implicit assumptions about non-core histones. First, whereas

protein sequence differences between core and variant histone paralogs are often assumed

to reflect differences in protein function, here we show that the variant protein HIS-35 is

likely to have a redundant function with core H2A despite the sequence difference. Second,

while all *C. elegans* H2A variants have a single intron, our observation of multi-intron variants and of recurrent intron loss, suggests that specific introns may not have crucial roles in the expression of histone variants. Instead, the role of introns in variant histones may simply lie in introns' general roles in promoting expression. Third, the combination of conservation and loss of variant histones points to potentially lineage-specific, partially redundant, or easily replaced roles of some histone variants. Future studies should explore the generality of these patterns across other lineages of eukaryotes. In addition to our specific findings for histone variant biology, these results highlight that introns can be useful in the reconstruction of the histories of complex gene families.

MATERIAL AND METHODS

Data source

Genomic sequences and gene feature format files of 108 nematode species were obtained from WormBase and the *Caenorhabditis* database (68, 69).

Data mining and processing

All the known genes of 108 nematode species with characterized exon-intron structures were fetched from their genomes using their respective 'gene feature format' file. We then noted the positions of the introns in the header of their respective genes and translated the *Caenorhabditis* gene sequences.

To identify the homologs of H2A and their variants, BLASTP, version 2.9.0+, was

performed using standard parameters while treating the translated gene sequences (of 108

nematode species) as the database and H2A and variant (H2A.Z^{HTZ-1}, HTAS-1, HIS-35)

protein sequences as the query (81). Using a maximum e-value of 1e-10, 8003 hits were

retrieved which were the homologs of H2A and H2A-variant genes. We then removed

dubious genes encoding proteins more than 200 amino acids long, because histone proteins

are generally shorter. We collapsed the genes whose introns align at the position and have

an identical protein sequence. After filtering off the genes we were left with 408 distinct

entries.

Previous studies have shown the intron position conservation among widely diverged

eukaryotic species (53-55, 57-60). Therefore, to assess the intron position conservation

among the putative H2A variant genes, we performed a Multiple Sequence Alignment

(MSA) using the default parameters of CLUSTALW (82). We mapped the intron positions

of each gene onto the corresponding protein CLUSTALW alignment, allowing us to

identify as potential H2A.Z^{HTZ-1}, HTAS-1, HIS-35 orthologs those genes with intron

positions matching *C. elegans* intron positions.

Phylogenetic Analysis

Multiple sequence alignment of 408 H2A variants homologs was performed using

default parameters of MUSCLE to generate a phylogenetic tree (Figure S1) using IQtree,

which does an automatic selection of the model by doing a model fit test and likelihood

scoring (83, 84). VT+R9, the variable time method, was selected by IQtree for our

data. The tree didn't yield a clear phylogenetic signal for HIS-35 or H2A.Z^{HTZ-1}, with

homologs exhibiting the *C. elegans* HIS-35 or H2A.Z^{HTZ-1} intron positions are scattered

over the tree (Figure S1; IP 234 and IP 233 respectively). However, when we took a closer

look at the HIS-35 characteristic intron-containing genes, we see that genes containing an

intron at the *C. elegans* HIS-35 intron position (Figure S1, IP 234) are restricted to most

species of *Caenorhabditis* and its sister genus *Diploscapter*. A clear clade of species was

seen which had HTAS-1 characteristic intron position (Figure S1, IP 152). For consistency

in Figures 2-4, we show the portion of the phylogenetic tree containing *Caenorhabditis*

species and *Disploscapter*.

421   Confirmation of H2A variant losses

We found a loss of H2A.Z^{HTZ-1}, HTAS-1, HIS-35 characteristic introns in a few

lineages (marked by a minus sign in Figures 2, 3, and 4). To know whether this is a real

loss or reflected errors in gene annotation, tblastn searches were performed across the

genome of these species. This manual curation led to the variant's characteristic intron

splice sites being identified by eye in a few species due to alignment gaps at the exact intron

position, indicating that these species truly contain the variant and that failure to initially

identify the variant is due to a failure of the annotation to include these genes. We included

a material and method flowchart as the Figure S4.

Acknowledgments

We thank our lab members for their helpful discussions. This work was supported by NSF:1616878; NSF STC DBI-1548297; NSF MCB RUI 1817611; NIH NICHD R03HD093990.

References

1. Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. Crystal structure of the nucleosome core particle at 2.8 Å resolution. Nature. 1997;389(6648):251-60.

2. Luger K, Dechassa ML, Tremethick DJ. New insights into nucleosome and chromatin structure: an ordered state or a disordered affair? Nature reviews Molecular cell biology. 2012;13(7):436-47.

3. Talbert PB, Henikoff S. Histone variants - ancient wrap artists of the epigenome. Nature reviews. Molecular cell biology. 2010 Apr;11(4):264-75.

4. Malik HS, Henikoff S. Phylogenomics of the nucleosome. Nature structural & molecular biology. 2003;10(11):882-91.

5. Mei Q, Huang J, Chen W, Tang J, Xu C, Yu Q, et al. Regulation of DNA replication-coupled histone gene expression. Oncotarget. 2017 -10-16;8(55):95005.

6. Pandey NB, Chodchoy N, Liu T, Marzluff WF. Introns in histone genes alter the distribution of 3′ ends. Nucleic Acids Res. 1990;18(11):3161-70.

7. Romeo V, Schümperli D. Cycling in the nucleus: regulation of RNA 3′ processing and

nuclear organization of replication-dependent histone genes. Curr Opin Cell Biol.

2016;40:23-31.

8. Biterge B, Schneider R. Histone variants: key players of chromatin. Cell Tissue Res.

2014;356(3):457-66.

9. Henikoff S, Smith MM. Histone variants and epigenetics. Cold Spring Harbor

perspectives in biology. 2015 Jan 05,;7(1):a019364.

10. Talbert PB, Ahmad K, Almouzni G, Ausió J, Berger F, Bhalla PL, et al. A unified

phylogeny-based nomenclature for histone variants. Epigenetics & chromatin.

2012;5(1):1-19.

11. Martire S, Banaszynski LA. The roles of histone variants in fine-tuning chromatin

organization and function. Nature reviews. Molecular cell biology. 2020 Sep;21(9):522-

41.

12. Weber CM, Henikoff S. Histone variants: dynamic punctuation in transcription. Genes

Dev. 2014;28(7):672-82.

13. Talbert PB, Henikoff S. Environmental responses mediated by histone variants. Trends

Cell Biol. 2014;24(11):642-50.

14. Filipescu D, Szenker E, Almouzni G. Developmental roles of histone H3 variants and

their chaperones. Trends in Genetics. 2013;29(11):630-40.

15. Maze I, Noh K, Soshnev AA, Allis CD. Every amino acid matters: essential

contributions of histone variants to mammalian development and disease. Nature Reviews

Genetics. 2014;15(4):259-71.

16. Otero S, Desvoyes B, Gutierrez C. Histone H3 dynamics in plant cell cycle and

development. Cytogenetic and genome research. 2014;143(1-3):114-24.

17. Volle C, Dalal Y. Histone variants: the tricksters of the chromatin world. Curr Opin

Genet Dev. 2014;25:8-14.

18. Herchenröther A, Wunderlich TM, Lan J, Hake SB. Spotlight on histone H2A variants:

From B to X to Z. Seminars in Cell & Developmental Biology; Elsevier; 2022.

19. Talbert PB, Henikoff S. Histone variants at a glance. J Cell Sci. 2021;134(6):jcs244749.

20. Phillips EO, Gunjan A. Histone variants: The unsung guardians of the genome. DNA

repair. 2022;112:103301.

21. Deal RB, Henikoff S. Histone variants and modifications in plant gene regulation. Curr

Opin Plant Biol. 2011;14(2):116-22.

22. Lei B, Berger F. H2A variants in Arabidopsis: versatile regulators of genome activity.

Plant Communications. 2020;1(1):100015.

23. Tessarz P, Kouzarides T. Histone core modifications regulating nucleosome structure

and dynamics. Nature reviews Molecular cell biology. 2014;15(11):703-8.

24. Bönisch C, Hake SB. Histone H2A variants in nucleosomes and chromatin: more or

less stable? Nucleic acids research. 2012 Nov;40(21):10719-41.

25. Molaro A, Young JM, Malik HS. Evolutionary origins and diversification of testis-

specific short histone H2A variants in mammals. Genome Res. 2018;28(4):460-73.

26. Faast R, Thonglairoam V, Schulz TC, Beall J, Wells JR, Taylor H, et al. Histone variant

H2A. Z is required for early mammalian development. Current Biology.

2001;11(15):1183-7.

27. Meneghini MD, Wu M, Madhani HD. Conserved histone variant H2A. Z protects

euchromatin from the ectopic spread of silent heterochromatin. Cell. 2003;112(5):725-36.

28. Fan JY, Rangasamy D, Luger K, Tremethick DJ. H2A. Z alters the nucleosome surface

to promote HP1α-mediated chromatin fiber folding. Mol Cell. 2004;16(4):655-61.

29. Colino-Sanguino Y, Clark SJ, Valdes-Mora F. The H2A. Z-nuclesome code in

mammals: emerging functions. Trends in Genetics. 2021.

30. Petty EL, Collette KS, Cohen AJ, Snyder MJ, Csankovszki G. Restricting dosage

compensation complex binding to the X chromosomes by H2A. Z/HTZ-1. PLoS genetics.

2009;5(10):e1000699.

31. Long J, Carter B, Johnson ET, Ogas J. Contribution of the histone variant H2A. Z to

expression of responsive genes in plants. Seminars in Cell & Developmental Biology;

Elsevier; 2022.

32. Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded

breaks induce histone H2AX phosphorylation on serine 139. J Biol Chem.

1998;273(10):5858-68.

33. Rivera-Casas C, Gonzalez-Romero R, Cheema MS, Ausió J, Eirin-Lopez JM. The

characterization of macroH2A beyond vertebrates supports an ancestral origin and

conserved role for histone variants in chromatin. Epigenetics. 2016;11(6):415-25.

34. Sun Z, Bernstein E. Histone variant macroH2A: from chromatin deposition to

molecular function. Essays Biochem. 2019;63(1):59-74.

35. Osakabe A, Molaro A. Histone renegades: Unusual H2A histone variants in plants and

animals. Seminars in Cell & Developmental Biology; Elsevier; 2022.

36. Guberovic I, Farkas M, Corujo D, Buschbeck M. Evolution, structure and function of

divergent macroH2A1 splice isoforms. Seminars in Cell & Developmental Biology;

Elsevier; 2022.

37. Contrepois K, Coudereau C, Benayoun BA, Schuler N, Roux P, Bischof O, et al.

Histone variant H2A. J accumulates in senescent cells and promotes inflammatory gene

expression. Nature communications. 2017;8(1):1-18.

38. Shaytan AK, Landsman D, Panchenko AR. Nucleosome adaptability conferred by

sequence and structural variations in histone H2A–H2B dimers. Curr Opin Struct Biol.

2015;32:48-57.

39. Jiang X, Soboleva TA, Tremethick DJ. Short histone H2A variants: small in stature but

not in function. Cells. 2020;9(4):867.

40. Hoghoughi N, Barral S, Vargas A, Rousseaux S, Khochbin S. Histone variants:

essential actors in male genome programming. The Journal of Biochemistry.

2018;163(2):97-103.

41. Millar CB. Organizing the genome with H2A histone variants. Biochem J.

2013;449(3):567-79.

42. Buschbeck M, Hake SB. Variants of core histones and their roles in cell fate decisions,

development and cancer. Nature reviews Molecular cell biology. 2017;18(5):299-314.

43. Wolffe AP. Histone Genes. In: Brenner's Online Encyclopedia of Genetics, Four-

Volume Set. Elsevier Inc; 2001. p. 948-52.

44. Marzluff WF, Wagner EJ, Duronio RJ. Metabolism and regulation of canonical histone

mRNAs: life without a poly (A) tail. Nature Reviews Genetics. 2008;9(11):843-54.

45. Le Hir H, Nott A, Moore MJ. How introns influence and enhance eukaryotic gene

expression. Trends Biochem Sci. 2003;28(4):215-20.

46. Dixon RJ, Eperon IC, Samani NJ. Complementary intron sequence motifs associated

with human exon repetition: a role for intragenic, inter-transcript interactions in gene

expression. Bioinformatics. 2007;23(2):150-5.

47. Rose AB. Intron-mediated regulation of gene expression. Nuclear pre-mRNA

processing in plants. 2008:277-90.

48. Casas-Mollano JA, Lao NT, Kavanagh TA. Intron-regulated expression of SUVH3, an

Arabidopsis Su (var) 3-9 homologue. J Exp Bot. 2006;57(12):3301-11.

49. Luco RF, Allo M, Schor IE, Kornblihtt AR, Misteli T. Epigenetics in alternative pre-

mRNA splicing. Cell. 2011;144(1):16-26.

50. Armstrong JF, Pritchard-Jones K, Bickmore WA, Hastie ND, Bard JB. The expression

of the Wilms' tumour gene, WT1, in the developing mammalian embryo. Mech Dev.

1993;40(1-2):85-97.

51. Cieply B, Carstens RP. Functional roles of alternative splicing factors in human disease.

Wiley Interdisciplinary Reviews: RNA. 2015;6(3):311-26.

52. Faustino NA, Cooper TA. Pre-mRNA splicing and human disease. Genes Dev.

2003;17(4):419-37.

53. Roy SW, Fedorov A, Gilbert W. Large-Scale Comparison of Intron Positions in

Mammalian Genes Shows Intron Loss but No Gain. Proceedings of the National Academy

of Sciences - PNAS. 2003 Jun 10,;100(12):7158-62.

54. Sêton Bocco S, Csűrös M. Splice Sites Seldom Slide: Intron Evolution in Oomycetes.

Genome biology and evolution. 2016 Aug 25,;8(8):2340-50.

55. Rogozin IB, Wolf YI, Sorokin AV, Mirkin BG, Koonin EV. Remarkable interkingdom

conservation of intron positions and massive, lineage-specific intron loss and gain in

eukaryotic evolution. Current Biology. 2003;13(17):1512-7.

56. Coulombe-Huntington J, Majewski J. Characterization of intron loss events in

mammals. Genome Res. 2007;17(1):23-32.

57. Irimia M, Roy SW. Spliceosomal introns as tools for genomic and evolutionary

analysis. Nucleic acids research. 2008 Mar;36(5):1703-12.

58. Sakharkar MK, Tan TW, de Souza SJ. Generation of a database containing discordant

intron positions in eukaryotic genes (MIDB). Bioinformatics. 2001;17(8):671-5.

59. Rogozin IB, Lyons-Weiler J, Koonin EV. Intron sliding in conserved gene families.

Trends in Genetics. 2000;16(10):430-2.

60. Van Daal A, White EM, Elgin SC, Gorovsky MA. Conservation of intron position

indicates separation of major and variant H2As is an early event in the evolution of

eukaryotes. J Mol Evol. 1990;30(5):449-55.

61. Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. Chromosomal DNA and

its packaging in the chromatin fiber. In: Molecular Biology of the Cell. 4th edition. Garland

Science; 2002.

62. Draizen EJ, Shaytan AK, Mariño-Ramírez L, Talbert PB, Landsman D, Panchenko AR.

HistoneDB 2.0: a histone database with variants—an integrated resource to explore

histones and their variants. Database. 2016;2016.

63. Ahmad K, Henikoff S. Histone H3 variants specify modes of chromatin assembly.

Proceedings of the National Academy of Sciences. 2002;99(suppl 4):16477-84.

64. Hake SB, Allis CD. Histone H3 variants and their potential role in indexing mammalian

genomes: the "H3 barcode hypothesis". Proceedings of the National Academy of Sciences.

2006;103(17):6428-35.

65. Hake SB, Garcia BA, Duncan EM, Kauer M, Dellaire G, Shabanowitz J, et al.

Expression patterns and post-translational modifications associated with mammalian

histone H3 variants. J Biol Chem. 2006;281(1):559-68.

66. Santenard A, Ziegler-Birling C, Koch M, Tora L, Bannister AJ, Torres-Padilla M.

Heterochromatin formation in the mouse embryo requires critical residues of the histone

variant H3. 3. Nat Cell Biol. 2010;12(9):853-62.

67. Chu DS, Liu H, Wu TF, Ralston EJ, Nix P, Meyer BJ, et al. Sperm chromatin

proteomics identifies evolutionarily conserved fertility factors. Nature (London). 2006 Sep

07,;443(7107):101-5.

68. Davis P, Zarowiecki M, Arnaboldi V, Becerra A, Cain S, Chan J, et al. WormBase in

2022—data, processes, and tools for analyzing Caenorhabditis elegans. Genetics.

2022;220(4):iyac003.

69. Stevens L, Félix M, Beltran T, Braendle C, Caurcel C, Fausett S, et al. Comparative

genomics of 10 new Caenorhabditis species. Evolution Letters. 2019;3(2):217-36.

70. Carlton PM, Davis RE, Ahmed S. Nematode chromosomes. Genetics.

2022;221(1):iyac014.

71. Kelchner SA, Thomas MA. Model use in phylogenetics: nine key questions. Trends in Ecology & Evolution. 2007;22(2):87-94.

72. Nei M, Rooney AP. CONCERTED AND BIRTH-AND-DEATH EVOLUTION OF MULTIGENE FAMILIES. Annual review of genetics. 2005 Dec 15,;39(1):121-52.

73. Liao D. Concerted evolution. Nature Encyclopedia of the human genome. 2003;1:938-42.

74. Baldo AM, Les DH, Strausbaugh LD. Potentials and limitations of histone repeat sequences for phylogenetic reconstruction of Sophophora. Mol Biol Evol. 1999;16(11):1511-20.

75. Fitch DH, Strausbaugh LD. Low codon bias and high rates of synonymous substitution in Drosophila hydei and D. melanogaster histone genes. Mol Biol Evol. 1993;10(2):397-413.

76. Meneghini MD, Wu M, Madhani HD. Conserved histone variant H2A. Z protects euchromatin from the ectopic spread of silent heterochromatin. Cell. 2003;112(5):725-36.

77. Irimia M, Rukov JL, Penny D, Garcia-Fernandez J, Vinther J, Roy SW. Widespread evolutionary conservation of alternatively spliced exons in Caenorhabditis. Mol Biol Evol. 2008;25(2):375-82.

78. Kiontke K, Gavin NP, Raynes Y, Roehrig C, Piano F, Fitch DH. Caenorhabditis phylogeny predicts convergence of hermaphroditism and extensive intron loss. Proceedings of the National Academy of Sciences. 2004;101(24):9003-8.

79. Raman P, Rominger MC, Young JM, Molaro A, Tsukiyama T, Malik HS. Novel classes

and evolutionary turnover of histone H2B variants in the mammalian germline. Mol Biol

Evol. 2022;39(2):msac019.

80. Kasimatis KR, Phillips PC. Rapid gene family evolution of a nematode sperm protein

despite sequence hyper-conservation. G3: Genes, Genomes, Genetics. 2018;8(1):353-62.

81. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic

acids research. 1997 Sep;25(17):3389-402.

82. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of

progressive multiple sequence alignment through sequence weighting, position-specific

gap penalties and weight matrix choice. Nucleic acids research. 1994 Nov 11,;22(22):4673-

80.

83. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic acids research. 2004 Mar 01,;32(5):1792-7.

84. Nguyen L, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Molecular

biology and evolution. 2015 Jan;32(1):268-74.

```
HTAS-1    MARLKQRPNRILNTS--TKTSSAKKKKKRISRSTRSGLTFPVGRIHRKLRETTRGKQRIS          58
HTZ-1     ----------MAGGKGKAGKDSGKSKSKVVSRSARAGLQFPVGRIHRFLKQRTTSSGRVG          50
HIS-35    -----------MSGRGKGGKA--KTGGKAKSRSSRAGLQFPVGRLHRILRKGN-YAQRVG          46
H2A       -----------MSGRGKGGKA--KTGGKAKSRSSRAGLQFPVGRLHRILRKGN-YAQRVG          46
                      .          .    *.   *   ***:*:** *****:** *:: .     *:.

HTAS-1    AGASVFMAATLEYLTTELMEMSAIAANESKKSRVTPRHLHLAIYGDQETAQLLDKVTLPQ          118
HTZ-1     ATAAVYSAAILEYLTAEVLELAGNASKDLKVKRITPRHLHLAIRGDEELDTLIK-ATIAG          109
HIS-35    AGAPVYLAAVLEYLAAEVLELAGNAARDNKKTRIAPRHLQLAVRNDEELNKLLAGVTIAQ          106
H2A       AGAPVYLAAVLEYLAAEVLELAGNAARDNKKTRIAPRHLQLAVRNDEELNKLLAGVTIAQ          106
          * * *: ** ****::*::*::. *:.: * .*::****:**: .*:*   *:   .*:

HTAS-1    GGVTPMPIHPSLLPKKKAKEDDKENNS-----     145
HTZ-1     GGVIPH-IHRYLMNKKGAPVPGKPGAPGQGPQ     140
HIS-35    GGVLPN-IQAVLLPKKTAGDKE----------     127
H2A       GGVLPN-IQAVLLPKKTGGDKE----------     127
          *** *  *:  *: **  .
```

Figure 1. Sequence and intron position comparison of three H2A variants of *C. elegans*. *C. elegans* core histone H2A and its three variant paralogs contain different sequences and intron positions. The variants differ either in intron positions or the phases in which they interrupt the codon. HTAS-1 (highlighted with a yellow) has a phase 0 intron between the 26th and 27th amino acid; HIS-35 (highlighted with pink) has a phase 0 intron between 50th and 51st amino acid. HTZ-1 (highlighted with blue) has a phase 2 intron splitting the 57th amino acid.

Figure 2: Intron position and sequence evidence indicate the origin of HTAS-1 within *Caenorhabditis* and subsequent loss. On the left is the previously reconstructed species tree of *Caenorhabditis* and *Diploscapter*. The likely origin of HTAS-1 is indicated with a dark pink bar. HTAS-1 presence in a species is indicated by the plus sign. The question mark denotes species where HTAS-1 homologs could not be identified.. The species where HTAS-1 was not incorporated are indicated as an asterisk. On the right is the multiple sequence alignment of HTAS-1 proteins showing the aligned intron positions (highlighted with yellow). For consistency, we show the portion of the phylogenetic tree containing the *Caenorhabditis* species and *Disploscapter*.

HIS-35

H2A

Intron Position 51.0

Figure 3: Comparison of the variant HIS-35 (left) with canonical H2A (right) sequences across *Caenorhabditis* species. Inferred sequence changes relative to the reconstructed ancestral sequence are highlighted with different colors (pink, red, yellow, blue, green), with identical changes colored identical colors across the figure (i.e., the L-to-R change at position 35th is colored purple on both protein alignments).

Figure 4: The dynamic history of intron loss and gain in HTZ-1. On the left is the previously reconstructed species tree of *Caenorhabditis*. HTZ-1 characteristic intron presence and absence in a species is indicated by the plus and minus sign. Yellow hash marks on the tree branch depict the loss of intron-1 whereas the blue hash mark on the branch suggests the loss of intron 2 in those lineages.