An initial assessment of the sustainability of waterbird harvest in the United Kingdom

Matthew B. Ellis (Corresponding author)

British Association for Shooting & Conservation

Marford Mill,

Rossett,

Wrexham, UK

LL12 0HL

matt.ellis@basc.org.uk

ORCID 0000-0003-2215-3190

Tom C. Cameron

School of Life Sciences

University of Essex

Wivenhoe Park

Colchester, UK

CO4 3SQ

tcameron@essex.ac.uk

ORCID 0000-0002-5875-1494

Abstract

1. There is a need to assess the sustainability of current harvest levels of wild birds in the United Kingdom (UK), and more widely, across Europe. Yet data on population and harvest size, as well as population demographics of many species are limited.

2. We used a Demographic Invariant Method (DIM) to estimate Potential Excess Growth (PEG) for populations of waterbirds wintering in the UK and calculated a Sustainable Harvest Index (SHI) for each species or population. We compared this with population trends and conservation classifications (e.g. Birds of Conservation Concern) to assess the sustainability of harvests and the utility of these classifications.

3. Our approach found evidence for potential overharvest in the United Kingdom of mallard *Anas platyrhynchos*, Eurasian teal *Anas crecca*, gadwall *Mareca strepera*, Canada geese *Branta canadensis* and both British and Icelandic greylag geese *Anser anser*. Whether DIM methods predict overharvest for ducks is highly dependent on estimates of maximum population growth rates inferring PEG. We found estimates of maximum population growth to be variable across a range of different methods.

4. We found no relationship between sustainable harvest indices and short-term or long-term wintering trends in harvested species, or conservation classification under the UK's Birds of Conservation Concern framework.

Policy Implications: Our results suggest that UK based harvest is unlikely to be a major determinant of population trends for overwintering waterbirds in the UK, but harvest rates for some species require further investigation. Additionally, the lack of a relationship between conservation classifications and SHI strongly suggests that these are not an appropriate tool for making decisions about harvest management. Our assessment provides the basis for a framework to make evidence-based decisions on sustainable harvest levels in the UK in the face of incomplete data.

Investment in data collection for European waterbird population size and harvest is

long overdue.

Keywords

Waterbirds, harvest, hunting management, potential excess growth, demographic invariant

method

Word count

6766

Introduction

Hunting of waterbirds in the UK with shotguns has occurred since at least the 1600s (Helme,

1614; Markham, 1621). Contemporary recreational hunting of waterbirds in the UK targets

nine species of duck, four species of goose and three waders. Coot *Fulica atra* and moorhen

Gallinula chloropus are also huntable but are rarely pursued and there are no modern

estimates of the harvest of these species. The total annual harvest of ducks, geese and

waders in the UK is estimated at approximately 1.4 million (Aebischer, 2019; PACEC, 2014).

These same species, hunted in winter in the UK, breed across Northern, Eastern Europe

and Russia and are also hunted by nations in the breeding and wintering range and across

their migration routes.

The British Association for Shooting and Conservation (BASC) is the largest

representative body for shooting sports in the UK. BASC has approximately 8,000

wildfowling members, which is likely to represent the majority of self-identified wildfowlers

hunting ducks and geese on UK coasts and estuaries. Of these wildfowlers approximately

25% hunt at least once per season, and they harvest an average of 8 birds each per season

(Crown Estate, 2021; Ellis, 2014). This leads to an estimated coastal harvest of 16,000

waterbirds. The approximately 1.4 million remaining harvested waterbirds are harvested

inland over a broad range of habitats, including artificially fed sites. Released mallard *Anas*

platyrhynchos are a significant proportion of the inland waterbird harvest with current release

estimates of approximately 2.6 million (Madden, 2021), and harvest estimates of

approximately 950,000 birds per year (Aebischer, 2019). A significant proportion of other

wild duck and goose species are also harvested in these inland areas due to the creation

and provision of appropriate habitat, as well as supplementary feeding.

The rarity, localisation and population trends of birds in the UK has been assessed at

regular intervals under the Birds of Conservation Concern (BoCC) methodology since 1996

(Gibbons et al., 1996). Species are assigned the highest priority rating (Red>Amber) for any

of the criteria they exceed such that, for example, a species which exceeded the Amber

breeding range decline criterion and the Red breeding population decline criterion would be

classified as Red-listed. Species which do not exceed any of the assessment criteria are

placed on the Green list. A species with an increasing or stable UK population can be

classed as Amber due to, for example, the UK holding an internationally important proportion

of the European population– e.g. Pink-footed goose *Anser brachyrhynchus*. This approach is

different to the better-known International Union for the Conservation of Nature (IUCN) Red

List which only reviews the extinction risk for species (IUCN, 2020). This can lead to

disparities due to differences in the scale of assessment. For example, the Eurasian

woodcock *Scolopax rusticola* is classed as least conservation concern by the IUCN due to a

very large and stable global population and large range but has Red status on the UK BoCC

due to long and short term declines in UK breeding range.

Increasingly there have been calls to prohibit or restrict the hunting of species that

are Red or Amber listed on BoCC (Thomas, 2021; Welsh Parliament, 2020). Similar calls

drawing attention to the BoCC status of hunted birds have been made by social media

influencers such as Chris Packham (Common, 2016; UK Government and Parliament,

2016). Currently all huntable ducks and geese in the UK, with the exception of tufted duck

Aythya fuligula and Canada goose *Branta canadensis*, are Red or Amber listed using BoCC,

most commonly due to the UK holding >20% of the European wintering population, rather

than due to declines in breeding or wintering populations. Apriori then, BoCC may not be the

most appropriate metric to judge conservation risks from additional winter mortality, but this

has not yet been tested. Likewise, there have been calls to prohibit or restrict harvest in

nationally and internationally protected sites. Typically, on these sites, harvest data is robust,

and harvests are low (approximately 1% of the estimated UK harvest), compared to

unprotected inland sites where regulatory authorities currently have limited room to act.

These regulatory disparities lead to conflicts and this calls for an assessment of the

sustainability of the UK harvest of such highly mobile species, that are constantly responding

to food availability, recreational disturbance and hunting pressure with shifting range size

and range location, at an appropriate spatial scale (Beatty et al., 2014; O'Connell et al.,

2007).

In the absence of regular and systematic monitoring of geographic population

structures or harvest returns for huntable species in the UK it has historically been difficult to

assess the likely sustainability of current harvest levels. Particularly so for data poor ducks

compared to more regularly monitored European goose populations (Holopainen et al.,

2018; Madsen et al., 2015). However, the development of methods allowing for estimates of

harvest levels in the UK (Aebischer, 2019), and the development of analytical tools to assess

the sustainability of harvest with incomplete demographic data now allows for initial

assessments to be made even in the face of uncertainty (Eraud et al., 2021). The UK is a

party to the Agreement on the Conservation of African-Eurasian Migratory Waterbirds

(AEWA) which is developing mechanisms to ensure flyway-level harvest sustainability and in

the long term will be a more appropriate level to make these decisions (Madsen et al. 2015).

However, in the absence of these mechanisms we present an initial assessment of the

sustainability of winter hunting mortality of UK waterbirds using best-available data, whilst

recognising the likely sources of error and identifying areas where further research would be

particularly valuable.

Materials and Methods

We collated mean and confidence intervals for harvest estimates, population size estimates,

and demographic data, and also population trends on huntable ducks, geese and waders in

the UK. This included nine species of ducks (mallard *Anas platyrhynchos*; Eurasian teal

Anas crecca; Eurasian wigeon *Mareca penelope*; gadwall *Mareca strepera*; northern pintail

Anas acuta; northern shoveler *Spatula clyptea*; tufted duck *Aythya fuligula*; common pochard

Aythya ferina; and common goldeneye *Bucephala clangula*), three waders (Eurasian

woodcock *Scolopax rusticola*; common snipe *Gallinago gallinago*; and golden plover

Pluvialis apricaria) and three goose species comprising a total of four goose populations

(British (resident) and Icelandic (migratory) greylag goose A*nser anser*; Greenland/Icelandic

pink-footed goose *Anser brachyrhynchus*; and, resident Canada goose *Branta canadensis*).

These data were used to estimate a Sustainable Hunting Index using a stochastic simulation

approach using the "popharvest" R package (Eraud et al. 2021).

Estimating harvest sustainability

A Sustainable Hunting Index (SHI + 95% confidence interval (CI)) was calculated for each

species using the potential excess growth (PEG) function in the popharvest R package

(Eraud et al., 2021). This index is calculated as the ratio of harvest to PEG such that any

value in excess of 1 indicates overharvest and values under 1 are increasingly less likely to

be unsustainable as they approach zero. This is not unlike a surplus production model

approach that is commonly applied to data poor or initial assessments for harvest of other

vertebrate systems such as fisheries (Jennings et al., 2008).

The PEG for each species was calculated using a Demographic Invariant Method

(DIM) which allows estimation of maximal potential population growth rates (λ_{max}) with

relatively few demographic parameters including age at first reproduction, life-history

strategy categorisation and adult survival (Ernaud et al., 2021; Johnson et al., 2012). This

stochastic simulation approach allows us to best account for the uncertainty in data sources

The choice of life history strategy ("long" versus "short") and the use of adult survival

estimates from either ringing studies or estimated from species mass can have significant

effects on the estimate of SHI. Given the uncertainty around these variables and the

significant effects each can have on the estimated outputs we ran four models per species

varying life history strategy ("long" versus "short") and adult survival ("reported" versus

"estimated") whilst holding other variables the same and present the results averaged over

these four approaches. In each model we ran 10,000 Monte Carol simulations per species

with both population and harvest estimates drawn from a uniform range equal to the 95%

confidence intervals of published estimates (full details on data sources given in

supplementary online materials). The age at first reproduction is poorly understood for most

species so was taken to be fixed at 1 for all duck species, 2 for all waders and 3 for all geese

(Robinson, 2005). Distributions for drawing simulation parameters from were guided by

those recommended by the package authors (Eraud et al., 2021).

Models using "reported" adult survival use the average adult survival reported from

ring returns. However, this can underestimate maximal survival rates under optimal

conditions for harvested species and bias low the estimated sustainable harvest index. In

models with "estimated" adult survival we provided body mass information to the model and

allow it to draw on an estimate of adult survival in optimum conditions (Johnson 2012) from

estimates of how adult survival responds to variation in body mass across many bird taxa

(Ricklefs, 2000).

The popharvest package includes a safety factor (F_s) which can be set from 0-1 to

limit harvest to only a proportion of the PEG (with a recommended maximum of $F_s = 0.5$).

Based on published recommendations (Dillingham & Fletcher, 2008), we set $F_s = 0.5$ for all

species listed as least concern by IUCN. Pochard is listed as Vulnerable under IUCN and

therefore we assigned an $F_s = 0.1$. Setting F_s to a recommended maximum of 0.5 does not

well represent the sustainability of a harvest where hunters are being encouraged to

maximise their harvest, for example on non-native geese or geese with expanding

populations, e.g. Canada or pink-footed goose. We shall examine this issue later in the

discussion. As well as calculating a confidence interval for SHI the package also calculates

the probability that the current level of harvest is unsustainable. Estimated λ_{max} using

published survival estimates from ringing studies and the inbuilt body mass predicted adult

survival were compared with other published assessments for Canada goose and mallard to

make sure our DIM approaches were producing reasonable estimates (Table S1).

We tested for relationships between SHI and Birds of Conservation Concern

category using a Welch's one-way ANOVA which allows for unequal variance and between

SHI and wintering population trend (short and long term) using multiple linear regression. All

analyses were conducted in R version 4.1.2 (R Core Team, 2021).

Results

The estimated potential excess growth (PEG), sustainable hunt index (SHI) and probability

of unsustainable harvest averaged across the four models can be found in Table 1

(individual model estimates presented in Tables S2-S5). The current harvest for mallard,

teal, gadwall, Canada goose and both British and Icelandic greylag goose were predicted to

be unsustainable against an objective of harvesting 50% or less of maximum annual

productivity. Harvest levels were up to eight times greater than half the potential excess

growth for each population (i.e.when F_s is set to 0.5). The highest predicted SHI for duck

species was for Eurasian teal at 1.071 (95% CI 0.686-1.487), followed by mallard, gadwall,

shoveler and wigeon in decreasing order (Table 1). The probability of an unsustainable

harvest of wintering UK ducks, that is harvest greater than 50% of the maximum potential

surplus production of current winter population sizes, were generally low , even for h mallard

(0.502) and teal (0.514) which was indecisive for the model average. For pochard, where the

threshold for sustainability was set to only 10% of the maximum surplus growth due to recent

breeding population declines in continental Europe as well as poor UK breeding and harvest

data, SHI was 0.358 (95% CI 0.102 – 0.634) with a low probability of unsustainable harvest

(0.002). For the four wader populations considered none had SHIs or 95% CIs exceeding 1

and the probability of an unsustainable harvest was low (Table 1). However, the upper 95%

confidence interval approached 1 for both resident (0.989) and migratory (0.880) woodcock.

The choice of life history strategy (e.g. short vs long) and survival estimates (e.g.

reported vs DIM estimate) had a large effect on the magnitude of mean SHI estimates, but

had little effect on the overall species order (Figure 1 and Tables S2-S5).

There was no significant relationship between the average SHI and the BoCC5

category across all species ($F_{(2,4.569)}$ = 2.990, p = 0.148), and there was also no significant

relationship between average SHI and short-term (β = 0.011, p = 0.303) or long-term

wintering trends (β = 0.000, p = 0.843)(Figure 2).

Figure 1: Mean (and 95% confidence interval) sustainable harvest index (SHI) for selected

huntable waterbirds in the United Kingdom modelled using survival rates estimated from

mass and (A) a long life history or (B) a short life history, and survival rates reported in the

literature and a (C) long life history or (D) short life history. SHI is based on an F_s of 0.5 for

all species and the dotted line at SHI=1 represents the threshold for overharvest. Species

were selected to represent those huntable waterbirds which are most well studied.

Figure 2: Mean and 95% confidence intervals for the sustainable harvest index (SHI) for

huntable waterbirds in the United Kingdom along with their short-term (2008-2018) and long-

term (1970 – 2018) wintering population trends. The dotted line at SHI = 1 represents the

threshold for overharvest such that species falling to the right of the line are likely to be

experiencing overharvest. SHI is calculated based on an F_s of 0.5 for all species other than

pochard which is set to 0.1. Species are colour coded according to their classification under

Birds of Conservation Concern 5 (BoCC5).

Discussion

Discussion

In this study we used data on waterbird population sizes and harvest levels to estimate

sustainable harvest indices (SHI). For the majority of ducks and all waders hunted in the UK

we found no evidence of unsustainable harvests, but we recommend further scrutiny for

mallard, teal and gadwall where the mean SHI or upper 95% CI exceeded 1. For this

assessment we used a moderating factor (F_s) of 0.5 for all ducks and waders other than

pochard. If a socio-ecological management plan for duck or wader harvest in the UK and

Europe aimed for higher or lower harvests than 50% of the surplus production, then the SHI

assessment would change.

Our estimates of SHI for wild geese suggest that Canada geese and Icelandic and

British greylag geese may be subject to unsustainable harvest levels in the UK. This remains

the case for Canada goose and both greylag goose populations if a higher moderating factor

is chosen (F_s=0.9 where harvest target is 90% of maximum surplus growth). However,

British greylag goose, Canada goose and indeed gadwall experience strongly increasing

population trends in recent years (Frost et al., 2019). Although there is some evidence of

levelling off of these trend increases in the most recent years (Frost et al., 2021), particularly

for Canada goose which could indicate that the curent harvest is finally having an effect on

stabilising or reducing its numbers. In this context an unsustainable harvest of Canada

goose may be desirable but woud suggest mansgement to reduce that harvest may be

required at some point. Our SHI estimate for the migratory Icelandic greylag goose

population is high with a high probability of unsustainable harvest. The Icelandic greylag

goose population trend is declining, and this species is subject to significant additional

hunting mortality outside of the UK. However, there are difficulties in accurately assessing

the proportion of the harvest to assign to the resident and migratory populations and

between the two main nations that hunt this population (UK and Iceland), which can clearly

have a significant impact on the calculation of the SHI. For example, the mean annual

harvest of greylag geese in Iceland was 43,000 per annum from 2015/16 to 2019/20

(Statistics Iceland, 2022), and we estimated the harvest for the UK as 22,500, giving a total

estimated harvest across the flyway of 65,500 from a total estimated population of 60,000

(Brides et al., 2021). But this does suggest action to improve estimates and limit the total

harvest of this population would be warranted, at least as an adaptive experiment to test for

linkage between harvest and population recovery.

The high SHIs for mallard and teal indicate that these species may be subject to

unsustainably high levels of harvest and this appears to accord with slowing of growth rates

in teal populations and ongoing declines in mallard. The mean SHIs for these species were

close to or exceeded 1 in three out of the four models we ran (Tables S2-S5). Our apriori

best candidate model for mallard and teal uses a short life history and DIM estimates adult

survival (Table S5), with maximum adult survival values that are reasonable from our review

of the literature (Table S1) also predicts mean SHI values above 1. This result highlights a

need to improve population demographic data and reduce harvest, as harvest reduction can

help us understand how the populations respond to harvest control and therefore better

inform and manage the sustainability of future hunting opportunities.

We found no relationship between SHI and short-term or long-term wintering

population trends. We suggest that this indicates that hunting of waterbirds in the UK is not a

primary driver of population trends. This is to be expected as waterfowl harvests tend to be a

function of waterfowl productivity in a given year (Fox et al., 2016; Holopainen, Christensen,

et al., 2018). This means hunters tend to harvest more ducks in years when surplus

production is higher and moderate harvests tend not to have long term effects on waterfowl

population sizes. That is not to say that the sum of harvests across the flyway would give

similar results, and we encourage further work using these tools but on international scales.

Likewise further scrutiny of trends and over what period they apply may be useful as long-

term trends for the total Eurasian woodcock population size (e.g. 25yr), were stated as

stable in 2015 but are now stated as having strong declines (Birdlife 2015 vs 2021).

Immediate steps would be to build further realism into this initial model assessment of

sustainability by incorporating the demographic breakdown of the hunting bag as affecting

juvenile or adult survival and attempting the model with estimates of population sizes and

hunting mortality at larger spatial scales across the UK, mainland Europe and western

Russia.

The lack of a relationship between SHI and BoCC5 categories suggests that

classification as Red or Amber listed by this scheme is not an appropriate tool to use as an

aid to harvest management decisions. This is not unexpected as BoCC5 classification

depends on a variety of factors including breeding and wintering localisation and is not solely

dependent on international population trends or demographic sensitivities of the species

concerned. For example, pink-footed geese are amber listed for non-breeding localisation

(greater than 50% of the UK non-breeding population at ten or fewer sites) and non-breeding

international importance (UK holds at least 20% of the European non-breeding

population)(Stanbury et al., 2021), despite significant long and short term population

increases (Burns et al., 2020). However, this does not undermine the value of BoCC as a

useful tool for the initial prioritisation of species of potential conservation concern at

appropriate spatial scales; particularly for making habitat availability and restoration

decisions.

In estimating the potential excess growth or annual recruitment in a bird population

with which to assess the sustainability of harvest, a safety feature has been included so that

the threshold with which to judge overharvesting is a proportion of the excess growth (Eraud

et al., 2021; Johnson et al., 2012; Niel & Lebreton, 2005). This is wise due to well-

recognised risks in using surplus growth approaches in data poor systems (Jennings et al.,

2008). The sequential use of this safety feature in frameworks designed for initial

assessments of the sustainability of wild bird harvests has led to a clear and well described

framework where the proportion of the excess growth that should not be exceeded by

harvest can decline with data availability, population size, and declining conservation status

(Dillingham & Fletcher, 2008; Eraud et al., 2021). But this initial assessment method does

not take into account any specific harvest management objectives that may be equally

important in utilitarian conservation systems such as legal recreational fishing or hunting,

where the objective may be to maximise harvest opportunities for some species. This may

be due to the species being non-native (e.g. Canada or Egyptian goose *Alopchen*

aegyptiaca and Mandarin duck *Aix galericulata* in the UK), causing damage (e.g. pink-footed

goose) or due to its relatively high abundance and productivity (e.g. Eurasian teal) making it

a candidate for safe, sustainable regulated harvesting of wild food resources. Here we have

chosen a "safety" net of 0.5 to apply to excess growth of all species (except pochard) as they

are of Least Concern conservation status according to IUCN (IUCN, 2022).

Harvesting up to 50% of the surplus production of highly productive species such as

waterfowl is not unreasonable, where, in North America for example, 5.9 million Mallards

were harvested from a breeding population of 14 million in 1999 (Wilkins & Cooch, 1999).

Assuming the mean DIM estimated maximum population growth rate with a slow life-history

of 1.765 for mallard, that represents 55% of the estimated surplus production on mallard in

that year. Similar estimates of large harvest of 30+% of the annual surplus production of

mallards in North America are common and sustained (e.g. U.S. Fish and Wildlife Service

2018). However, it is potentially informative to examine the effect of altering the level of

precaution applied to the estimates of SHI. For example, reducing the F_s for Eurasian

wigeon from 0.5 (reported in Table 1) to 0.1 changes the mean SHI estimate from 0.413 to

2.064. Likewise, these models can be used to estimate the harvest required to achieve an

SHI value ≥1 for pink-footed goose if the aim is to reduce or stabilise population growth and

expansion, though the Prescribed Take Level (PTL) function of the popharvest package is

better suited to this task.

We have completed this initial assessment of the sustainability of waterbird harvest in

the UK using the best available data at this time and applying all potential estimates of

maximum potential growth. However, we recognise that there are multiple sources of

uncertainty in the data and parameter estimates. Population estimates of overwintering

waterbirds in the UK are well documented and widely accepted, but there are known issues

that influence reliability for this exercise. For example, the latest population estimates are

scored for reliability out of 3 (3 = least reliable), and greylag geese, Canada geese, teal and

goldeneye are scored 2 and mallard, snipe, woodcock and golden plover are scored 3 (Frost

et al., 2019). Wetland bird survey (WebS) counts are made by an extensive network of

volunteers overseen by the British Trust for Ornithology (BTO), and for many sites data

extends back to 1947. However, coverage is incomplete and 39% of WeBS sites are

currently vacant, of which 19.5% are "high" or "very high" priority (BTO, 2022). Such gaps

undoubtedly contribute to the low reliability of some population estimates. Furthermore,

changes to the methods used to make population estimates can have significant impacts.

For example, the latest teal population estimates doubled from previous estimates as a

result of new calculation methods (Musgrove et al. 2011; Méndez et al. 2015).

Overall, UK population estimates are likely to be underestimates due to count

methods and the combined effects of hunting mortality and migration – this will result in

underestimates of the potential excess growth and lead to a perceived greater proportion of

the population being harvested. Depending on the management objectives (maximising

population versus maximising sustainable harvest) this may be more or less important for

decision makers.

Waterbird survival estimates are reported variously from the UK, North America or

across Europe, but generally from ringing recaptures. The DIM approach requires good

estimates of maximal adult survival under optimal conditions (Eraud et al., 2021).

Underestimates of survival rates will bias estimates of population growth rates, potentially

overestimating excess growth and hence underestimating the sustainable harvest index. We

reviewed different approaches to estimate the maximum population growth for two of the

most well studied hunted species to see if the DIM method produced "sensible" estimates

(Table S1). We assumed Canada geese had a "long" life history and mallard "short"

(equivalent to Table S2 and S3 respectively). The DIM estimate of maximum population

growth using reported survival of 1.225 for Canada goose seems reasonable in light of

studies across its range over varying time periods (1.07-1.74). However, for mallard, the DIM

estimated maximum population growth using reported survival of 1.980 appears high relative

to realised maximum population growth observations of 1.20-1.59 (Table S1). However, for

both species taking published estimates of nesting attempts, nest success and brood

survival can lead to productivity estimates more in line with the highest values e.g 1.58-2.10

for mallard (Table S1). As all realised population growth estimates from time series counts

include hunting mortality, natural mortality and variation caused by environmental conditions

and also suffer survey design error and biases, this may explain this discrepancy where

observed trends are never estimating maximum population growth potential. Further work is

required but given this review we feel the best estimates for Geese likely come from Table

S2 and S4, and best SHI estimates for ducks and waders likely come from Table S5.

The accuracy of current UK waterbird harvest estimates is unknown (Aebischer,

2019) and developing a harvest recording system for the UK should remain a priority, not

least to assist with management decisions at the flyway-scale (Madsen et al., 2015). The

prospects for improving harvest data quality through voluntary systems seem low in the UK,

especially in light of the diversity of hunters and hunter organisations, low trust in regulatory

agencies by hunters and low levels of participation in other voluntary schemes (Ellis, 2020).

However, it would seem to be of high policy importance to hunters and their organisations to

be able to demonstrate the sustainability of their hunting activities. Appropriate policy

intervention may be welcome in this area provided it is applied to all participants, so that

harvest data is collected from inland and private land areas as well as public coastal areas,

areas of significantly higher and lower hunting pressure respectively.

Making decisions with incomplete data is difficult. We have used the best available

data to make this preliminary assessment of the sustainability of harvest levels in the UK, but

clearly further work is needed, including a structured assessment of the harvest at the

flyway-scale. Our assessment provides a first step towards an informed discussion of

harvest levels in the UK and whilst improved data collection will be useful, a collaborative

approach to management based on the principles of Structured Decision Making and

Adaptive Management could provide a transparent and trusted system to initiate with

existing data that is "good enough" (Holopainen et al., 2018; Johnson et al., 2018; Williams &

Brown, 2014). This approach is beginning to be developed in Europe by AEWA through

mechanisms such as the European Goose Management Platform. This removes the need to

rely on simplistic and inappropriate traffic light systems (such as Birds of Conservation

Concern) which invariably result in poorly supported and simplistic outcomes (prohibiting

versus status-quo). This is especially the case as we have shown their use to be unrelated

to hunting mortality across all species. We would instead recommend a move towards a

system which explicitly values recreational hunting, such as how the UK values recreational

fishing, and allows for an ecologically sustainable approach which is socially acceptable,

evidence based and able to initiate with incomplete data.

Authors' Contributions

ME and TC conceived the idea, ME designed the methodology, collected and analysed the

data, and wrote the first draft of the manuscript. TC contributed critically to drafts of the

manuscript, validated the analyses and produced the supplementary information.

Conflict of interest

MBE is employed by and is a member of BASC, the UK's largest representative body for

shooting sports. TCC is a member of BASC, the British Trust for Ornithology and Essex

Wildlife Trust.

Data availability statement

Data will be hosted on the DataDryad Digital Repository [DOI to be added]

References

Aebischer, N. J. (2019). Fifty-year trends in UK hunting bags of birds and mammals, and

calibrated estimation of national bag size, using GWCT's National Gamebag Census.

European Journal of Wildlife Research, *65*(4), 64. https://doi.org/10.1007/s10344-019-

1299-x

Balmer, D. E., Gillings, S., Caffrey, B. J., Swann, R. L., Downie, I. S., & Fuller, R. J. (2013).

Bird Atlas 2007-11: the breeding and wintering birds of Britain and Ireland. BTO books.

Thetford, UK.

Beatty, W. S., Webb, E. B., Kesler, D. C., Raedeke, A. H., Naylor, L. W., & Humburg, D. D.

(2014). Landscape effects on mallard habitat selection at multiple spatial scales during

the non-breeding period. *Landscape Ecology*, *29*(6), 989–1000.

https://doi.org/10.1007/s10980-014-0035-x

BirdLife International. (2021). European Red List of Birds. Publications Office of the

European Union. Luxembourg.

Brides, K., Mitchell, C., & Auhage, S. N. V. (2021). *Status and distribution of Icelandic-*

breeding geese: results of the 2020 international census. Wildfowl and Wetlands Trust.

Slimbride, UK.

BTO. (2022). *Vacant sites*. https://app.bto.org/websonline/sites/vacant/vacant-sites.jsp date

accessed: 07/02/2022

Burns, F., Eaton, M. A., Balmer, D. E., Banks, A., Caldow, R., Donelan, J. L., Douse, A.,

Duigan, C., Foster, S., Frost, T., Grice, P. V., Hall, C., Hanmer, H. J., Harris, S. J.,

Johnstone, I., Lindley, P., McCulloch, N., Noble, D. G., Risely, K., … Wotton, S. (2020).

The State of the UK's Birds 2020. RSPB, BTO, WWT, DAERA, JNCC, NatureScot, NE

and NRW. Sandy, UK.

Common, J. (2016). *New campaign sparks controversy over the shooting of wading birds*.

Conservation Jobs. https://www.conservationjobs.co.uk/articles/new-campaign-sparks-

controversy-over-the-shooting-of-wading-birds/ date accessed: 02/03/2022.

Crown Estate. (2021). *Crown Estate foreshore sporting rights leases to wildfowling clubs*

season summaries from 2007/08 to 2020/21 England, Wales and Northern Ireland.

https://www.thecrownestate.co.uk/media/3934/wildfowl-returns-season-summary.pdf

date accessed: 07/02/2022.

Dillingham, P. W., & Fletcher, D. (2008). Estimating the ability of birds to sustain additional

human-caused mortalities using a simple decision rule and allometric relationships.

Biological Conservation, *141*(7), 1783–1792.

https://doi.org/10.1016/j.biocon.2008.04.022

Ellis, M. B. (2014). A Profile of Waterfowl Hunting and Hunters in the UK. *Human*

Dimensions of Wildlife, *19*(4), 391–392. https://doi.org/10.1080/10871209.2014.908251

Ellis, M. B. (2020). Waterfowl hunters' motivations and barriers to participation in a citizen

science project. *Wildfowl*, *70*, 228–241.

Eraud, C., Devaux, T., Villers, A., Johnson, F. A., & Francesiaz, C. (2021). popharvest: An R

package to assess the sustainability of harvesting regimes of bird populations. *Ecology*

and Evolution, *11*(23), 16562–16571. https://doi.org/10.1002/ece3.8212

Fox, A. D., Clausen, K. K., Dalby, L., Christensen, T. K., & Sunde, P. (2016). Between-year

variations in sex/age ratio bias in hunter wings of Eurasian Wigeon (Anas penelope)

compared to field samples. *Ornis Fennica*, *93*(1), 26–30.

Frederiksen, M. (2002). Indirect estimation of the number of migratory Greylag and pink-

footed geese shot in Britain. *Wildfowl*, *53*, 27–34.

Frost, T. M., Austin, G., Hearn, R., Mcavoy, S., Robinson, A., Stroud, D., Woodward, I.,

Wotton, S., & Allen, R. (2019). Population estimates of wintering waterbirds in Great

Britain. *British Birds*, *112*(March 2019), 130–145.

Frost, T. M., Calbrade, N. A., Birtles, G. A., Hall, C., Robinson, A. E., Wotton, S. R., Balmer,

D. E., & Austin, G. E. (2021). *Waterbirds in the UK 2019/20: The Wetland Bird Survey*.

BTO/RSPB/JNCC. Thetford, UK.

Gibbons, D. W., Reid, J. B., & Chapman, R. A. (1996). Bird species of conservation concern

in the United Kingdom, Channel Islands and Isle of Man: revising the Red Data List.

RSPB Conservation Review, *10*, 7–18.

Helme, J. (1614). *A jewell for gentrie. Being an exact dictionary, or true method, to make any*

man understand all the art, secrets, and worthy knowledges belonging to hawking,

hunting, fowling and fishing. Together with all the true measures for winding of the

horne. Snodham, T. London, UK

Heward, C. J., Hoodless, A. N., Conway, G. J., Aebischer, N. J., Gillings, S., & Fuller, R. J.

(2015). Current status and recent trend of the Eurasian Woodcock Scolopax rusticola

as a breeding bird in Britain. *Bird Study*, *62*(4), 535–551.

https://doi.org/10.1080/00063657.2015.1092497

Holopainen, S., Arzel, C., Elmberg, J., Fox, A. D., Guillemain, M., Gunnarsson, G., Nummi,

P., Sjöberg, K., Väänänen, V.-M., Alhainen, M., & Pöysä, H. (2018). Sustainable

management of migratory European ducks: finding model species. *Wildlife Biology*,

2018(1). https://doi.org/10.2981/wlb.00336

Holopainen, S., Christensen, T. K., Pöysä, H., Väänänen, V. M., Rintala, J., & Fox, A. D.

(2018). Associations between duck harvest, hunting wing ratios and measures of

reproductive output in Northern Europe. *European Journal of Wildlife Research*, *64*(6).

https://doi.org/10.1007/s10344-018-1227-5

IUCN. (2020). IUCN Red List 2017-2020 report. IUCN. Gland, Switzerland.

IUCN. (2022). *The IUCN Red List of Threatened Species. Version 2021-3.*

https://www.iucnredlist.org date accessed 07/02/2022.

Jennings, S., Kaiser, M. J., & Reynolds, J. D. (2008). *Marine fisheries ecology*. Wiley-

Blackwell. Oxford, UK.

Johnson, F. A., Alhainen, M., Fox, A. D., Madsen, J., & Guillemain, M. (2018). Making do

with less: must sparse data preclude informed harvest strategies for European

waterbirds? *Ecological Applications*, *28*(2), 427–441. https://doi.org/10.1002/eap.1659

Johnson, F. A., Walters, M. A. H., & Boomer, G. S. (2012). Allowable levels of take for the

trade in Nearctic songbirds. *Ecological Applications*, *22*(4), 1114–1130.

https://doi.org/10.1890/11-1164.1

Madden, J. R. (2021). How many gamebirds are released in the UK each year? *European*

Journal of Wildlife Research, *67*(4), 1–14. https://doi.org/10.1007/s10344-021-01508-z

Madsen, J., Guillemain, M., Nagy, S., Defos du Rau, P. ., Modain-Monval, J.-Y., Griffin, C.,

Williams, J. H., Bunnefeld, N., Czajkowski, A., Hearn, R., Grauer, A., Alhainen, M., &

Middleton, A. (2015). *Towards sustainable Management of Huntable Migratory*

Waterbirds in Europe: A report by the Waterbird Harvest Specialist Group of Wetlands

International. Wetlands International. Wageningen, Netherlands.

Markham, G. (1621). *Hunger's prevention: Or, the whole art of fowling by water and land*.

Francis Grove. London, UK.

Méndez, V., Austin, G. E., Musgrove, A. J., Ross-Smith, V. H., Hearn, R. D., Stroud, D. A.,

Wotton, S. R., & Holt, C. A. (2015). Use of environmental stratification to derive non-

breeding population estimates of dispersed waterbirds in Great Britain. *Journal for*

Nature Conservation, *28*, 56–66. https://doi.org/10.1016/j.jnc.2015.09.001

Musgrove, A. J., Austin, G. E., Hearn, R. D., Holt, C. A., Stroud, D. A., & Wotton, S. R.

(2011). Overwinter population estimates of British waterbirds. *British Birds*, *104*(364–

397).

Niel, C., & Lebreton, J. D. (2005). Using demographic invariants to detect overharvested bird

populations from incomplete data. *Conservation Biology*, *19*(3), 826–835.

https://doi.org/10.1111/j.1523-1739.2005.00310.x

O'Connell, M., Ward, R., & Onoufriou, C. (2007). Integrating multi-scale data to model the

relationship between food resources, waterbird distribution and human activities in

freshwater systems: preliminary findings. *Ibis*, *149*, 65–72.

https://doi.org/10.1111/j.1474-919X.2007.00659.x

PACEC. (2014). *The Value of Shooting: The economic, environmental and social benefits of*

shooting sports to the UK. PACEC. Cambridge, UK.

R Core Team. (2021). *R: A language and environment for statistical computing. R*

Foundation for Statistical Computing. R Foundation for Statistical Computing. Geneva,

Switzerland.

Ricklefs, R. E. (2000) Intrinsic aging-related mortality in birds. *Journal of Avian Biology*, *31*,

103-111. https://doi.org/10.1034/j.1600-048X.2000.210201.x

Robinson, R. A. (2005). *BirdFacts: profiles of birds occurring in Britain & Ireland*.

http://www.bto.org/birdfacts date accessed: 07/02/2022.

Stanbury, A., Eaton, M., Aebischer, N., Balmer, D., Brown, A., Douse, A., Lindley, P.,

Mcculloch, N., Noble, D., & Win, I. (2021). The status of our bird populations: the fifth

Birds of Conservation Concern in the United Kingdom, Channel Islands and Isle of Man

and second IUCN Red List assessment of extinction risk for Great Britain. *British Birds*,

114(December 2021), 723–747.

Statistics Iceland. (2022). *Online database: https://statice.is* date accessed: 07/02/2022

Thomas, J. (2021). *Unfair game: the birds that can still be shot in the UK*. Bird Guides.

        https://www.birdguides.com/articles/conservation/unfair-game-the-birds-that-can-still-

        be-shot-in-the-uk/ date accessed: 07/02/2022.

U.S. Fish and Wildlife Service. (2018). *Waterfowl population status, 2018*. U.S. Department

        of the Interior. Washington D.C., USA.

UK Government and Parliament. (2016). *Introduce a moratorium on the hunting of critically*

        declining wading birds. https://petition.parliament.uk/archived/petitions/167410 date

        accessed: 07/02/2022.

Welsh Parliament. (2020). *Ban the shooting of critically endangered birds...give them the*

        protection they so desperately need! Petitions.

        https://petitions.senedd.wales/petitions/244712 date accessed: 07/02/2022.

Wilkins, K. A., & Cooch, E. G. (1999). *Waterfowl population status, 1999*. U.S. Fish & Wildlife

        Service, Department of the Interior. Washington D.C., USA.

Williams, B. K., & Brown, E. D. (2014). Adaptive management: From more talk to real action.

        Environmental Management, *53*(2), 465–479. https://doi.org/10.1007/s00267-013-0205-

Woodward, I. D., Massimino, D., Hammond, M. J., Barber, L., Barimore, C., Harris, S. J.,

        Leech, D. I., Noble, D. G., Walker, R. H., Baillie, S. R., & Robinson, R. A. (2020).

        BirdTrends 2020: trends in numbers, breeding success and survival for UK breeding

        birds. BTO Research Report 732. BTO. Thetford, UK.

570 Table 1: The estimated mean λ_{max}, potential excess growth (PEG; reported to two significant figures), sustainable hunt index (SHI) and the
571 probability of current harvest levels being unsustainable with a safety factor (F_s) = 0.5 for huntable waterbird species populations overwintering
572 in the United Kingdom, except for pochard where F_s = 0.1. The presented figures are an average of four models using either a "long" or "short"
573 life history strategy and either the "reported" or "estimated" adult survival rates.
574

Species	Estimated λ_{max} (95% CI)	Mean Potential Excess Growth (95% CI)	Mean Sustainable Hunt Index (95% CI)	Probability of unsustainable harvest
Mallard	1.626 (1.590 - 1.661)	1,000,000 (960,000 – 1,100,000)	1.054 (0.793 – 1.326)	0.502
Teal	1.714 (1.632 - 1.789)	160,000 (130,000 – 170,000)	1.071 (0.686 – 1.487)	0.514
Wigeon	1.690 (1.630 - 1.749)	160,000 (130,000 – 180,000)	0.413 (0.167 – 0.685)	0.016
Gadwall	1.573 (1.422 - 1.721)	8,900 (6,400 – 11,00)	0.773 (0.303 – 1.391)	0.249
Pintail	1.609 (1.484 - 1.727)	6,100 (4,500 – 7,700)	0.222 (0.045 – 0.432)	0.000
Shoveler	1.668 (1.604 - 1.727)	6,500 (5,500 – 7,500)	0.426 (0.164 – 0.712)	0.020
Tufted duck	1.587 (1.560 - 1.614)	41,000 (39,000 – 43,000)	0.155 (0.076 – 0.235)	0.000
Pochard	1.616 (1.584 - 1.648)	1,800 (1,600 – 2,000)	0.358 (0.102 – 0.634)	0.002

Goldeneye	1.538	5,600	0.201	0.000
	(1.446 - 1.632)	(4,500 – 6,800)	(0.028 – 0.396)	
Canada goose	1.215	17,000	2.001	0.867
	(1.210 - 1.220)	(17,000 – 19,000)	(0.879 – 3.147)	
Greylag goose (Icelandic)	1.184	8,400	2.761	1.000
	(1.143 - 1.222)	(6,400 – 10,000)	(2.173 – 3.658)	
Greylag goose (British)	1.184	13,000	4.149	0.954
	(1.143 - 1.221)	(9,900 – 16,000)	(0.877 – 8.052)	
Pink-footed goose	1.188	48,000	0.441	0.006
	(1.176 - 1.199)	(36,000 – 57,000)	(0.072 – 0.866)	
Woodcock (Resident)	1.375	31,000	0.662	0.079
	(1.334 - 1.411)	(21,000 – 42,000)	(0.437 – 0.989)	
Woodcock (Migratory)	1.374	230,000	0.592	0.038
	(1.334 - 1.411)	(160,000 – 310,000)	(0.392 – 0.880)	
Common Snipe	1.419	230,000	0.365	0.000
	(1.354 - 1.470)	(180,000 – 280,000)	(0.137 – 0.643)	
Golden plover	1.646	130,000	0.018	0.000
	(1.520 - 1.769)	(100,000 – 160,000)	(0.003 – 0.035)	