An initial assessment of the sustainability of waterbird harvest in the United Kingdom

Matthew B. Ellis (Corresponding author)

British Association for Shooting & Conservation

Marford Mill,

Rossett,

Wrexham, UK

LL12 0HL

matt.ellis@basc.org.uk

ORCID 0000-0003-2215-3190

Tom C. Cameron

School of Life Sciences

University of Essex

Wivenhoe Park

Colchester, UK

CO4 3SQ

tcameron@essex.ac.uk

ORCID 0000-0002-5875-1494

Abstract

1. There is a need to assess the sustainability of current harvest levels of wild birds in the United Kingdom (UK), and more widely, across Europe. Yet data on population and harvest size, as well as population demographics of many species are limited.

2. We used a Demographic Invariant Method (DIM) to estimate Potential Excess Growth (PEG) for populations of waterbirds wintering in the UK and calculated a Sustainable Harvest Index (SHI) for each species or population. We compared this with population trends and conservation classifications (e.g. Birds of Conservation Concern) to assess the sustainability of harvests and the utility of these classifications.

3. Our approach found no evidence for overharvest of any duck or wader species, except potentially for gadwall *Mareca strepera*. We estimated that Canada geese *Branta canadensis* and both British and Icelandic greylag geese *Anser anser* are potentially overharvested in the UK. With the exception of Icelandic greylag geese, all of the goose populations predicted to be experiencing overharvest show strong population growth. This conflicting result raises concern about current population size estimates, harvest estimates or both.

4. We found no relationship between sustainable harvest indices and short-term or long-term wintering trends in harvested species, or conservation classification under the UK's Birds of Conservation Concern framework.

Policy Implications: Our results suggest that UK based harvest is unlikely to be a major determinant of population trends for overwintering waterbirds in the UK. Additionally, the lack of a relationship between conservation classifications and SHI strongly suggests that these are not an appropriate tool for making decisions about harvest management. Our assessment provides the basis for a framework to make evidence-based decisions on sustainable harvest levels in the UK in the face of

incomplete data. Investment in data collection for European waterbird population size

and harvest is long overdue.

Keywords

Waterbirds, harvest, hunting management, potential excess growth, demographic invariant

method

Word count

6996

Introduction

Hunting of waterbirds in the UK with shotguns has occurred since at least the 1600s (Helme,

1614; Markham, 1621). Contemporary recreational hunting of waterbirds in the UK targets

nine species of duck, four species of goose and three waders. Coot *Fulica atra* and moorhen

Gallinula chloropus are also huntable but are rarely pursued and there are no modern

estimates of the harvest of these species. The total annual harvest of ducks, geese and

waders in the UK is estimated at approximately 1.4 million (Aebischer, 2019; PACEC, 2014).

These same species, hunted in winter in the UK, breed across Northern, Eastern Europe

and Russia and are also hunted by nations in the breeding range and across their migration

routes.

The British Association for Shooting and Conservation (BASC) is the largest

representative body for shooting sports in the UK. BASC has approximately 8,000

wildfowling members, which is likely to represent the majority of self-identified wildfowlers

hunting ducks and geese on UK coasts and estuaries. Of these wildfowlers approximately

25% hunt at least once per season, and they harvest an average of 8 birds each per season

(Crown Estate, 2021; Ellis, 2014). This leads to an estimated coastal harvest of 16,000

waterbirds. The approximately 1.4 million remaining harvested waterbirds are harvested

inland on fields and a broad range of wetlands, often on artificially fed sites. Released

mallard *Anas platyrhynchos* are a significant proportion of the inland waterbird harvest with

current release estimates of approximately 2.6 million (Madden, 2021), and harvest

estimates of approximately 950,000 birds per year (Aebischer, 2019). Therefore, a

significant proportion of other wild duck and goose species are also harvested in these

inland areas due to the creation and provision of appropriate habitat, as well as

supplementary feeding.

The conservation status of birds in the UK has been assessed under the Birds of

Conservation Concern (BoCC) methodology since 1996 (Gibbons et al., 1996). This

assessment is carried out at regular intervals (approximately every six years) by a

consortium of Non-Governmental Organisations (NGOs) and statutory agencies and

assesses the broad conservation status of birds in terms of rarity, localisation and population

trends. Species are assessed against the published criteria and assigned the highest priority

rating (Red>Amber) for any of the criteria they satisfy such that, for example, a species

which exceeded the Amber breeding range decline criterion and the Red breeding

population decline criterion would be classified as Red-listed. Any species which do not

satisfy any of the assessment criteria are placed on the Green list. A species with an

increasing or stable UK population could be classed as Amber due to, for example, the UK

holding an internationally important proportion of the European population– e.g. Pink-footed

goose *Anser brachyrhynchus*. This approach is different to the better known International

Union for the Conservation of Nature (IUCN) Red List which only reviews the extinction risk

for species (IUCN, 2020). This can lead to disparities due to differences in the scale of

assessment. For example, a species such as the Eurasian woodcock *Scolopax rusticola* can

be classed as being of least conservation concern by the IUCN due to a very large and

stable global population and large range but has Red status in the UK due to long and short

term declines in UK breeding range.

Increasingly there have been calls to prohibit or restrict the hunting of species that

are Red or Amber listed on BoCC (Thomas, 2021; Welsh Parliament, 2020). Similar calls

drawing attention to the BoCC status of hunted birds have been made by social media

influencers such as Chris Packham (Common, 2016; UK Government and Parliament,

2016). Currently all huntable ducks and geese in the UK, with the exception of tufted duck

Aythya fuligula and Canada goose *Branta canadensis*, are Red or Amber listed using BoCC,

and the most common reason for listing was due to the UK holding >20% of the European

wintering population, rather than due to declines in breeding or wintering populations. Apriori

then, BoCC may not be the most appropriate metric to judge conservation risks from

additional winter mortality, but this has not yet been tested. Likewise, there have been calls

to prohibit or restrict the harvest in protected areas where statutory agencies have regulatory

roles, for example in coastal hunting areas over nationally and internationally protected sites.

Typically, on these sites, harvest data is robust, and harvests are low (1% of the estimated

UK harvest), compared to inland areas which account for 99.9% of the UK waterbird harvest

but where regulatory authorities currently have limited room to act. These regulatory

disparities lead to conflicts and this calls for an assessment of the sustainability of the UK

harvest of such highly mobile species, that are constantly responding to food availability,

recreational disturbance and hunting pressure with shifting range size and range location, at

an appropriate spatial scale (Beatty et al., 2014; O'Connell et al., 2007).

In the absence of regular and systematic monitoring of breeding range population

sizes, geographic populations structures or harvest returns for huntable species in the UK it

has historically been difficult to assess the likely sustainability of current harvest levels.

Particularly so for data poor ducks compared to more regularly monitored European goose

populations (Holopainen, Arzel, et al., 2018; Madsen et al., 2015). However, the

development of methods allowing for estimates of harvest levels in the UK (Aebischer,

2019), and the development of analytical tools to assess the sustainability of harvest with

incomplete demographic data now allows for initial assessments to be made even in the face

of uncertainty (Eraud et al., 2021). The UK is a party to the Agreement on the Conservation

of African-Eurasian Migratory Waterbirds (AEWA) which is developing mechanisms to

ensure flyway-level harvest sustainability and in the long term will be a more appropriate

level to make these decisions (Elmberg et al. 2006; Madsen et al. 2015). However, in the

absence of these mechanisms we present an initial assessment of the sustainability of

winter hunting mortality of UK waterbirds using best-available data, whilst recognising the

likely sources of error and identifying areas where further research would be particularly

valuable.

Materials and Methods

We collated harvest estimates, population estimates, population trends and demographic

data on huntable ducks, geese and waders in the UK. This included nine species of ducks

(mallard *Anas platyrhynchos*; Eurasian teal *Anas crecca*; Eurasian wigeon *Mareca penelope*;

gadwall *Mareca strepera*; northern pintail *Anas acuta*; northern shoveler *Spatula clyptea*;

tufted duck *Aythya fuligula*; common pochard *Aythya ferina*; and common goldeneye

Bucephala clangula), three waders (Eurasian woodcock *Scolopax rusticola*; common snipe

Gallinago gallinago; and golden plover *Pluvialis apricaria*) and three goose species

comprising a total of four goose populations (British (resident) and Icelandic (migratory)

greylag goose A*nser anser*; Greenland/Icelandic pink-footed goose *Anser brachyrhynchus*;

and, resident Canada goose *Branta canadensis*).

Estimating harvest sustainability

A Sustainable Hunting Index (SHI + 95% confidence interval (CI)) was calculated for each

species using the potential excess growth (PEG) function in the popharvest R package

(Eraud et al., 2021). This index is calculated as the ratio of harvest to PEG such that any

value in excess of 1 indicates overharvest and values under 1 are increasingly less likely to

be unsustainable as they approach zero (Niel and Lebreton 2005). This is not unlike a

surplus production model approach that is commonly applied to data poor or initial

assessments for harvest of other vertebrate systems, i.e. fisheries (Jennings et al., 2008).

The PEG for each species was calculated using a Demographic Invariant Method (DIM)

which allows estimation of maximal potential population growth rates (λ_{max}) with relatively

few demographic parameters including age at first reproduction and life-history strategy

categorisation (Johnson et al., 2012). Estimated λ_{max} were compared with other published

assessments in the most common species geographic range and differential growth in BTO

breeding and wintering abundance indexes to make sure our DIM approach was producing

reasonable estimates (Table S1).

In order to account for uncertainty in reported life history characteristics we ran

10,000 Monte Carlo simulations per species with population estimates drawn from a uniform

range equal to the 95% confidence intervals calculated from the Wetland Bird Survey

(WeBS) indices for 2012/13 – 2016/17. Adult survival was drawn from a beta distribution with

a standard deviation equal to the calculated 95% confidence intervals. Harvest was drawn

from a uniform range equal to the 95% confidence intervals of published estimates

(Aebischer, 2019). Distributions for drawing simulation parameters from were guided by

those recommended by the package authors (Eraud et al., 2021) and references therein.

Age at first reproduction was taken to be fixed at 1 for all duck species, 2 for all waders and

3 for all geese (Robinson, 2005). All ducks and waders were assumed to have a "short" life

history strategy and geese a "long" life history strategy. The statistical package includes a

safety factor (F_s) which can be set from 0-1 to limit harvest to only a proportion of the PEG

(with a recommended maximum of $F_s = 0.5$) and it is recommended to set this at $F_s = 0.5$ for

species listed as least concern by IUCN (Dillingham & Fletcher, 2008). We therefore set $F_s =$

0.5 for all species except pochard which is listed as Vulnerable under IUCN and therefore

we assigned an $F_s = 0.1$. However, this does not well represent the sustainability of a

harvest where hunters are being encouraged to maximise their harvest, for example on non-

native geese or geese with expanding populations, e.g. Canada or pink footed goose. We

shall examine this issue later in the discussion. As well as calculating an SHI the package

also calculates the probability that the current level of harvest is unsustainable.

We tested for relationships between SHI and Birds of Conservation Concern

category using a Welch's one-way ANOVA, which allows for unequal variance and between

SHI and wintering population trend (short and long term) using multiple linear regression. All

analyses were conducted in R version 4.1.2 (R Core Team, 2021).

Data sources

Wintering bird populations for the UK were obtained from the latest published estimates

(Frost et al., 2019). The estimate for mallard was increased by 2.6 million birds to account

for the annual release of captive-raised birds for shooting purposes (Madden, 2021). We

estimated a standard deviation for the population estimates by averaging the Wetland Bird

Survey (WeBS; Frost et al. 2021) index for each species for the winters 2012/13 – 2016/17,

which corresponded to the period used to estimate duck populations. We treated this mean

index as equal to the estimate from Frost et al. (2019) and then calculated population

estimates for each year based on their WeBS indices and calculated the standard deviation

of these estimates. Population estimates for greylag goose, pink-footed goose, golden

plover, snipe and woodcock were based on single years, but the same time frame was

applied to standard deviation estimates in order to account for any interannual differences.

The latest (2016) harvest estimates and 95% confidence intervals were taken from

Aebischer (2019).

We modelled the resident and migratory populations of greylag geese and woodcock

separately. The resident woodcock population was estimated as three times the number of

breeding males. This was subtracted from the total estimated overwintering woodcock

population to provide an estimate of the migratory woodcock population. Resident woodcock

harvest was estimated as equal to the proportion of breeding birds in the overwinter

population (13%) The remaining 87% of total harvest was assigned to migratory woodcock.

We recognise that further mortality of UK breeding woodcock will occur outside the UK in

southern Europe, but we have no estimate of this mortality and can only consider the

contribution of UK hunters to the harvest of birds that winter in the UK in this initial

assessment. Population estimates for Icelandic and British greylag geese overwintering in

the UK are reported separately and no adjustment is needed. An estimated harvest of

migratory Icelandic greylag geese in the UK (Frederiksen, 2002) was deducted from the total

UK greylag goose harvest to provide an estimate of the harvest of British greylag geese.

However, it should be noted that the estimate was from 1996-2000 and no new estimates

have been made.

Short-term (2008-2018) and long-term (1970-2018) wintering population trends were

taken from Burns et al. (2020), except for common snipe and Eurasian woodcock. Common

snipe trends were taken from Woodward et al. (2020), with caution advised due to the small

sample size. Resident woodcock long-term and short-term population trends were estimated

at -29% for both on the basis of reported declines in breeding woodcock (Balmer et al.,

2013; Heward et al., 2015). Migratory woodcock short-term and long-term trends were

estimated at -11% and -22% on the basis of a 4-18% decline from 2008-2018 and an 11-

33% decline from 1980-2018 (BirdLife International, 2021). Bird population status in the UK

(Red/Amber/Green) was taken from Birds of Conservation Concern 5 (BoCC5; Stanbury et

al. 2021).

Estimates of adult survival were taken from Robinson (2005) with reported standard

errors multiplied by 1.96 to give an approximate 95% confidence interval. The average

standard error for all reported waterbirds (0.03) was used for species when no standard error

was reported (e.g. wigeon, shoveler, greylag goose, woodcock and golden plover). We used

the same survival estimates for both resident and migratory populations of woodcock and

greylag geese. These adult survival estimates include mortality from hunting and so are

likely to underestimate the maximum achievable survival rates under optimal conditions and

are therefore conservative. Age at first reproduction (alpha) was also taken from Robinson

(2005).

Results

The estimated potential excess growth (PEG), sustainable hunt index (SHI) and probability

of unsustainable harvest can be found in Table 1. The current harvest for Canada goose and

both British and Icelandic greylag goose were predicted to be unsustainable with harvest

levels two, three and four times greater than potential excess growth for each population

respectively when F_s is set to 0.5. The current harvest of Pink-footed goose has an SHI

value of 0.44 with a low probability of an unsustainable harvest (Table 1). The highest

predicted SHI for duck species was for Eurasian teal at 0.61, followed by mallard, gadwall,

shoveler and wigeon in decreasing order (Table 1). As well as no duck mean SHI

approaching a value of 1, the 95% CIs did not overlap a value of 1 for all ducks other than

gadwall. The probability of an unsustainable harvest of wintering UK ducks, that is harvest

greater than 50% of the maximum potential surplus production of current winter population

sizes, were all very low (0-0.04). For pochard, where the threshold for sustainability was set

to only 10% of the maximum surplus growth due to recent breeding population declines in

continental Europe as well as poor UK breeding and harvest data, SHI was 0.21 with a low

probability of unsustainable harvest. For the four wader populations considered none had

SHIs or 95% CIs exceeding 1 and the probability of an unsustainable harvest was low (Table

1).

There was no significant relationship between the average SHI and the BoCC5 category

across all species ($F_{(2,4.569)}$ = 1.017, p = 0.431), and there was also no significant relationship

between average SHI and short-term (β = -0.004, p = 0.678) or long-term wintering trends (β

= 0.009, p = 0.061).

261
262

Figure 1: Mean and 95% confidence intervals for the sustainable harvest index (SHI) for

huntable waterbirds in the United Kingdom along with their short-term (2008-2018) and long-

term (1970 – 2018) wintering population trends. The dotted line at SHI = 1 represents the

threshold for overharvest such that species falling to the right of the line are likely to be

experiencing overharvest. SHI is calculated based on an F_s of 0.5 for all species other than

pochard which is set to 0.1. Species are colour coded according to their classification under

Birds of Conservation Concern 5 (BoCC5).

Discussion

In this study we used data on waterbird population sizes and harvest levels to estimate

sustainable harvest indices (SHI). For the majority of ducks and all waders hunted in the UK

we found no evidence of unsustainable harvests, but we recommend further scrutiny for

gadwall. For this assessment we used a moderating factor (F_s) of 0.5 for all ducks and

waders other than pochard. If a socio-ecological management plan for duck or wader

harvest in the UK and Europe aimed for higher or lower harvests than 50% of the surplus

production, then the SHI assessment would change.

Our estimates of SHI for wild geese suggest that Canada geese and Icelandic and

British greylag geese may be subject to unsustainable harvest levels in the UK. This remains

the case for Canada goose and both greylag goose populations if a higher moderating factor

is chosen (F_s=0.9 where harvest target is 90% of maximum surplus growth). However,

British greylag goose, Canada goose and indeed gadwall experience strongly increasing

population trends in recent years (Frost et al., 2019). Although there is some evidence of

levelling off of these trend increases in the most recent years (Frost et al., 2021). Our review

of observed maximum population growth rates of the most common hunted waterbird

species across their range gives us confidence that our DIM approach is not underestimating

λ_{max}, e.g. maximum realised population growth of Canada goose in North America and the

UK is c8-10% per year (Table S1). This suggests that there are likely to be errors in the

assessment of the UK population size or hunting mortality. This uncertainty is reflected in the

large confidence intervals of the SHI for these three populations.

Our SHI estimate for the migratory Icelandic greylag goose population is high with a

high probability of unsustainable harvest. The Icelandic greylag goose population trend is

declining, and this species is subject to significant additional hunting mortality outside of the

UK. However, there are difficulties in accurately assessing the proportion of the harvest to

assign to the resident and migratory populations and between the two main nations that hunt

this population, which can clearly have a significant impact on the calculation of the SHI. For

example, the mean annual harvest of greylag geese in Iceland was 43,000 per annum from

2015/16 to 2019/20 (Statistics Iceland, 2022), and we estimated the harvest for the UK as

22,500, giving a total estimated harvest of 65,500 from a total estimated population of

60,000 (Brides et al., 2021). But this does suggest action to improve estimates or limit the

total harvest of this population would be warranted, at least as an adaptive experiment to

seek a small positive change in population growth, e.g. adaptive management.

We found no relationship between SHI and short-term or long-term wintering

population trends. We suggest that this indicates that hunting of waterbirds in the UK is not a

primary driver of population trends. This is to be expected as waterfowl harvests tend to be a

function of waterfowl productivity in a given year (Fox et al., 2016; Holopainen, Christensen,

et al., 2018).This means hunters tend to harvest more ducks in years when surplus

production is higher and moderate harvests tend not to have long term effects on waterfowl

population sizes. That is not to say that the sum of harvests across the flyway would give

similar results, and we encourage further work using these tools but on international scales.

Likewise further scrutiny of trends and over what period they apply may be useful as long-

term trends for the total Eurasian woodcock population size (e.g. 25yr), were stated as

stable in 2015 but are now stated as having strong declines (Birdlife 2015 vs 2021).

Immediate steps would be to build further realism into this initial model assessment of

sustainability by incorporating the demographic breakdown of the hunting bag as affecting

juvenile or adult survival and attempting the model with estimates of population sizes and

hunting mortality at larger spatial scales across the UK, mainland Europe and western

Russia.

The lack of a relationship between SHI and BoCC5 categories suggests that

classification as Red or Amber listed by this scheme is not an appropriate tool to use as an

aid to harvest management decisions. This is not unexpected as BoCC5 classification

depends on a variety of factors including breeding and wintering localisation and is not solely

dependent on international population trends or demographic sensitivities of the species

concerned. For example, pink-footed geese are amber listed for non-breeding localisation

(greater than 50% of the UK non-breeding population at ten or fewer sites) and non-breeding

international importance (UK holds at least 20% of the European non-breeding

population)(Stanbury et al., 2021), despite significant long and short term population

increases (Burns et al., 2020). However, this does not undermine the value of BoCC as a

useful tool for the initial prioritisation of species of potential conservation concern at

appropriate spatial scales; particularly for making habitat availability and restoration

decisions.

In estimating the potential excess growth or annual recruitment in a bird population

with which to assess the sustainability of harvest, a safety feature has been included so that

the threshold with which to judge overharvesting is a proportion of the excess growth (Eraud

et al., 2021; Johnson et al., 2012; Niel & Lebreton, 2005). This is wise due to well

recognised risks in using surplus growth approaches in data poor systems (Jennings et al.,

2008). The sequential use of this safety feature in frameworks designed for initial

assessments of the sustainability of wild bird harvests has led to a clear and well described

framework where the proportion of the excess growth that should not be exceeded by

harvest can decline with data availability, population size, and declining conservation status

(Dillingham & Fletcher, 2008; Eraud et al., 2021). But this initial assessment method does

not take into account any specific harvest management objectives that may be equally

important in utilitarian conservation systems such as legal recreational fishing or hunting,

where the objective may be to maximise harvest opportunities for some species. This may

be due to the species being non-native (e.g. Canada or Egyptian goose *Alopchen*

aegyptiaca and Mandarin duck *Aix galericulata* in the UK), causing damage (e.g. pink-footed

goose) or due to its relatively high abundance and productivity (e.g. Eurasian teal) it is a

candidate for safe, sustainable regulated harvesting of wild food resources. Here we have

chosen a "safety" net of 0.5 to apply to excess growth of all species (except pochard) as they

are of Least Concern conservation status according to IUCN (IUCN, 2022). Harvesting up to

50% of the surplus production of highly productive species such as waterfowl is not

unreasonable, where, in North America for example, 5.9 million Mallards were harvested

from a breeding population of 14 million in 1999 (Wilkins & Cooch, 1999). Assuming the DIM

estimated maximum population growth rate used here of 1.979 for mallard, 50% of the

surplus production would be 6.85 million ducks. Similar estimates of large harvest of 30+%

of the annual surplus production of mallards in North America are common and sustained

(e.g. U.S. Fish and Wildlife Service 2018). However, it is potentially informative to examine

the effect of altering the level of precaution applied to the estimates of SHI. For example,

reducing the F_s for Eurasian wigeon from 0.5 (reported in table 1) to 0.1 changes the mean

SHI estimate from 0.222 to 1.127. Likewise, these models can be used to estimate the

harvest required to achieve an SHI value ≥1 for pink-footed goose if the aim is to reduce or

stabilise population growth and expansion, though the Prescribed Take Level (PTL) function

of the popharvest package is better suited to this task.

We have completed this initial assessment of the sustainability of waterbird harvest in

the UK using the best available data at this time. However, we recognise that there are

multiple sources of uncertainty in our data. Population estimates of overwintering waterbirds

in the UK are well documented and widely accepted, but there are known issues that

influence reliability for this exercise. The latest population estimates are scored for reliability

out of 3, where 3 is less reliable, and greylag geese, Canada geese, teal and goldeneye are

scored 2 and mallard, snipe, woodcock and golden plover are scored 3 (Frost et al., 2019).

Any underestimate of the population will result in underestimates of the potential excess

growth, and changes to the methods used to make population estimates can have significant

impacts. For example, the latest teal population estimates doubled from previous estimates

as a result of new calculation methods (Musgrove et al. 2011; Méndez et al. 2015).

Population estimates for waterbirds in the UK are essentially the average of peak counts at

all counted sites across five years, with correction factors for missing data and extrapolation

factors for species known to be undercounted (Musgrove et al., 2011). Counts are made by

an extensive network of volunteers overseen by the British Trust for Ornithology (BTO), and

for many sites data extends back to 1947. However, coverage is incomplete and 39% of

WeBS sites are currently vacant, of which 19.5% are "high" or "very high" priority (BTO,

2022). Such gaps undoubtedly have an impact on the accuracy of population estimates.

These are likely our best estimates of the wintering population of hunted birds but do not

solve the problem of an accurate assessment of the total Flyway or northern European

breeding population of waterfowl as would be required for more realistic predictions of

waterfowl productivity (Elmberg et al., 2006; Holopainen, Arzel, et al., 2018).

Waterbird survival estimates are reported variously from the UK, or across Europe,

but generally from ringing recaptures. The DIM approach is sensitive to adult survival and

requires good estimates of maximal survival under optimal conditions (Eraud et al., 2021).

Our survival estimates are known underestimates as they are invariably measured in the

presence of hunting mortality. Underestimates of survival rates will bias estimates of

population growth rates, potentially overestimating excess growth and hence

underestimating the sustainable harvest index. To test whether the DIM approach is

generating sensible estimates of maximum population growth for waterbirds we reviewed

estimates for two of the most well studied hunted species; Canada goose and mallard (Table

S1). The DIM estimate of maximum population growth of 1.22 for Canada goose seems

reasonable in light of studies across its range over varying time periods (1.07-1.74).

However, for mallard, the DIM estimated maximum population growth of 1.979 appears high

relative to realised maximum population growth observations of 1.2-1.59 (Table S1). As all

realised population growth estimates include hunting mortality, variation caused by

environmental conditions and suffer survey design error and biases, this may explain this

discrepancy, but it warrants further investigation.

The accuracy of current UK waterbird harvest estimates is unknown (Aebischer,

2019). Developing a harvest recording system for the UK should remain a priority, not least

to assist with management decisions at the flyway-scale (Madsen et al., 2015). However, at

the UK-scale, the value of perfect harvest information is likely to be low so long as

population estimates remain at their current level of precision. The prospects for improving

harvest data quality through voluntary systems seem low in the UK, especially in light of the

diversity of hunters and hunter organisations, low trust in regulatory agencies by hunters and

low levels of participation in other voluntary schemes (Ellis, 2020). However, it would seem

to be of high policy importance to hunters and their organisations to be able to demonstrate

the sustainability of their hunting activities. Appropriate policy intervention may be welcome

in this area provided it is applied to all participants, so that harvest data is collected from

inland and private land areas as well as public coastal areas, areas of significantly higher

and lower hunting pressure respectively.

Making decisions with incomplete data is difficult. We have used the best available

data to make this preliminary assessment of the sustainability of harvest levels in the UK, but

clearly further work is needed, including a structured assessment of the harvest at the

flyway-scale and identification of key areas which could improve the accuracy of these initial

sustainability assessments. Our assessment provides a first step towards an informed

discussion of harvest levels in the UK and whilst improved data collection will be useful, a

collaborative approach to management based on the principles of Structured Decision

Making and Adaptive Management could provide a transparent and trusted system to initiate

with existing data that is "good enough" (Holopainen, Arzel, et al., 2018; Johnson et al.,

2018; Williams & Brown, 2014). This approach is beginning to be developed in Europe by

AEWA through mechanisms such as the European Goose Management Platform. This

removes the need to rely on simplistic and inappropriate traffic light systems (such as Birds

of Conservation Concern) which invariably result in poorly supported and simplistic

outcomes (prohibiting versus continue allowing hunting). This is especially the case as we

have shown their use to be unrelated to hunting mortality across all species. We would

instead recommend a move towards a system which explicitly values recreational hunting,

as the UK values recreational fishing, and allows for an ecologically sustainable approach

which is socially acceptable, evidence based and able to initiate with incomplete data.

Authors' Contributions

ME and TC conceived the idea, ME designed the methodology, collected and analysed the

data, and wrote the first draft of the manuscript. TC contributed critically to drafts of the

manuscript, validated the analyses and produced the supplementary information

Conflict of interest

MBE is employed by and is a member of BASC, the UK's largest representative body for

shooting sports. TCC is a member of BASC, the British Trust for Ornithology and Essex

Wildlife Trust.

Data availability statement

Data will be hosted on the DataDryad Digital Repository [DOI to be added]

References

Aebischer, N. J. (2019). Fifty-year trends in UK hunting bags of birds and mammals, and
    calibrated estimation of national bag size, using GWCT's National Gamebag Census.
    European Journal of Wildlife Research, *65*(4), 64. https://doi.org/10.1007/s10344-019-
    1299-x

Balmer, D. E., Gillings, S., Caffrey, B. J., Swann, R. L., Downie, I. S., & Fuller, R. J. (2013).
    Bird Atlas 2007-11: the breeding and wintering birds of Britain and Ireland. BTO books.
    Thetford, UK.

Beatty, W. S., Webb, E. B., Kesler, D. C., Raedeke, A. H., Naylor, L. W., & Humburg, D. D.
    (2014). Landscape effects on mallard habitat selection at multiple spatial scales during
    the non-breeding period. *Landscape Ecology*, *29*(6), 989–1000.
    https://doi.org/10.1007/s10980-014-0035-x

BirdLife International. (2021). European Red List of Birds. Publications Office of the
    European Union. Luxembourg.

Brides, K., Mitchell, C., & Auhage, S. N. V. (2021). *Status and distribution of Icelandic-
    breeding geese: results of the 2020 international census*. Wildfowl and Wetlands Trust.
    Slimbride, UK.

BTO. (2022). *Vacant sites*. https://app.bto.org/websonline/sites/vacant/vacant-sites.jsp date
    accessed: 07/02/2022

Burns, F., Eaton, M. A., Balmer, D. E., Banks, A., Caldow, R., Donelan, J. L., Douse, A.,

Duigan, C., Foster, S., Frost, T., Grice, P. V., Hall, C., Hanmer, H. J., Harris, S. J.,

Johnstone, I., Lindley, P., McCulloch, N., Noble, D. G., Risely, K., … Wotton, S. (2020).

The State of the UK's Birds 2020. RSPB, BTO, WWT, DAERA, JNCC, NatureScot, NE

and NRW. Sandy, UK.

Common, J. (2016). *New campaign sparks controversy over the shooting of wading birds*.

Conservation Jobs. https://www.conservationjobs.co.uk/articles/new-campaign-sparks-

controversy-over-the-shooting-of-wading-birds/ date accessed: 02/03/2022.

Crown Estate. (2021). *Crown Estate foreshore sporting rights leases to wildfowling clubs*

season summaries from 2007/08 to 2020/21 England, Wales and Northern Ireland.

https://www.thecrownestate.co.uk/media/3934/wildfowl-returns-season-summary.pdf

date accessed: 07/02/2022.

Dillingham, P. W., & Fletcher, D. (2008). Estimating the ability of birds to sustain additional

human-caused mortalities using a simple decision rule and allometric relationships.

Biological Conservation, *141*(7), 1783–1792.

https://doi.org/10.1016/j.biocon.2008.04.022

Ellis, M. B. (2014). A Profile of Waterfowl Hunting and Hunters in the UK. *Human*

Dimensions of Wildlife, *19*(4), 391–392. https://doi.org/10.1080/10871209.2014.908251

Ellis, M. B. (2020). Waterfowl hunters' motivations and barriers to participation in a citizen

science project. *Wildfowl*, *70*, 228–241.

Elmberg, J., Nummi, P., Pöysä, H., Sjöberg, K., Gunnarsson, G., Clausen, P., Guillemain,

M., Rodrigues, D., & Väänänen, V.-M. (2006). The scientific basis for new and

sustainable management of migratory European ducks. *Wildlife Biology*, *12*(2), 121–

127. https://doi.org/10.2981/0909-6396(2006)12[121:TSBFNA]2.0.CO;2

Eraud, C., Devaux, T., Villers, A., Johnson, F. A., & Francesiaz, C. (2021). popharvest: An R

package to assess the sustainability of harvesting regimes of bird populations. *Ecology*

and Evolution, *11*(23), 16562–16571. https://doi.org/10.1002/ece3.8212

Fox, A. D., Clausen, K. K., Dalby, L., Christensen, T. K., & Sunde, P. (2016). Between-year

variations in sex/age ratio bias in hunter wings of Eurasian Wigeon (Anas penelope)

compared to field samples. *Ornis Fennica*, *93*(1), 26–30.

Frederiksen, M. (2002). Indirect estimation of the number of migratory Greylag and pink-

footed geese shot in Britain. *Wildfowl*, *53*, 27–34.

Frost, T. M., Austin, G., Hearn, R., Mcavoy, S., Robinson, A., Stroud, D., Woodward, I.,

Wotton, S., & Allen, R. (2019). Population estimates of wintering waterbirds in Great

Britain. *British Birds*, *112*(March 2019), 130–145.

Frost, T. M., Calbrade, N. A., Birtles, G. A., Hall, C., Robinson, A. E., Wotton, S. R., Balmer,

D. E., & Austin, G. E. (2021). *Waterbirds in the UK 2019/20: The Wetland Bird Survey*.

BTO/RSPB/JNCC. Thetford, UK.

Gibbons, D. W., Reid, J. B., & Chapman, R. A. (1996). Bird species of conservation concern

in the United Kingdom, Channel Islands and Isle of Man: revising the Red Data List.

RSPB Conservation Review, *10*, 7–18.

Helme, J. (1614). *A jewell for gentrie. Being an exact dictionary, or true method, to make any*

man understand all the art, secrets, and worthy knowledges belonging to hawking,

hunting, fowling and fishing. Together with all the true measures for winding of the

horne. Snodham, T. London, UK

Heward, C. J., Hoodless, A. N., Conway, G. J., Aebischer, N. J., Gillings, S., & Fuller, R. J.

(2015). Current status and recent trend of the Eurasian Woodcock Scolopax rusticola

as a breeding bird in Britain. *Bird Study*, *62*(4), 535–551.

https://doi.org/10.1080/00063657.2015.1092497

Holopainen, S., Arzel, C., Elmberg, J., Fox, A. D., Guillemain, M., Gunnarsson, G., Nummi,

P., Sjöberg, K., Väänänen, V.-M., Alhainen, M., & Pöysä, H. (2018). Sustainable

management of migratory European ducks: finding model species. *Wildlife Biology*,

2018(1). https://doi.org/10.2981/wlb.00336

Holopainen, S., Christensen, T. K., Pöysä, H., Väänänen, V. M., Rintala, J., & Fox, A. D.

(2018). Associations between duck harvest, hunting wing ratios and measures of

reproductive output in Northern Europe. *European Journal of Wildlife Research*, *64*(6).

https://doi.org/10.1007/s10344-018-1227-5

IUCN. (2020). IUCN Red List 2017-2020 report. IUCN. Gland, Switzerland.

IUCN. (2022). *The IUCN Red List of Threatened Species. Version 2021-3.*

https://www.iucnredlist.org date accessed 07/02/2022.

Jennings, S., Kaiser, M. J., & Reynolds, J. D. (2008). *Marine fisheries ecology*. Wiley-

Blackwell. Oxford, UK.

Johnson, F. A., Alhainen, M., Fox, A. D., Madsen, J., & Guillemain, M. (2018). Making do

with less: must sparse data preclude informed harvest strategies for European

waterbirds? *Ecological Applications*, *28*(2), 427–441. https://doi.org/10.1002/eap.1659

Johnson, F. A., Walters, M. A. H., & Boomer, G. S. (2012). Allowable levels of take for the

trade in Nearctic songbirds. *Ecological Applications*, *22*(4), 1114–1130.

https://doi.org/10.1890/11-1164.1

Madden, J. R. (2021). How many gamebirds are released in the UK each year? *European*

Journal of Wildlife Research, *67*(4), 1–14. https://doi.org/10.1007/s10344-021-01508-z

Madsen, J., Guillemain, M., Nagy, S., Defos du Rau, P. ., Modain-Monval, J.-Y., Griffin, C.,

Williams, J. H., Bunnefeld, N., Czajkowski, A., Hearn, R., Grauer, A., Alhainen, M., &

Middleton, A. (2015). *Towards sustainable Management of Huntable Migratory*

Waterbirds in Europe: A report by the Waterbird Harvest Specialist Group of Wetlands

International. Wetlands International. Wageningen, Netherlands.

Markham, G. (1621). *Hunger's prevention: Or, the whole art of fowling by water and land*.

Francis Grove. London, UK.

Méndez, V., Austin, G. E., Musgrove, A. J., Ross-Smith, V. H., Hearn, R. D., Stroud, D. A.,

Wotton, S. R., & Holt, C. A. (2015). Use of environmental stratification to derive non-

breeding population estimates of dispersed waterbirds in Great Britain. *Journal for*

Nature Conservation, *28*, 56–66. https://doi.org/10.1016/j.jnc.2015.09.001

Musgrove, A. J., Austin, G. E., Hearn, R. D., Holt, C. A., Stroud, D. A., & Wotton, S. R.

(2011). Overwinter population estimates of British waterbirds. *British Birds*, *104*(364–

397).

Niel, C., & Lebreton, J. D. (2005). Using demographic invariants to detect overharvested bird

populations from incomplete data. *Conservation Biology*, *19*(3), 826–835.

https://doi.org/10.1111/j.1523-1739.2005.00310.x

O'Connell, M., Ward, R., & Onoufriou, C. (2007). Integrating multi-scale data to model the

relationship between food resources, waterbird distribution and human activities in

freshwater systems: preliminary findings. *Ibis*, *149*, 65–72.

https://doi.org/10.1111/j.1474-919X.2007.00659.x

PACEC. (2014). *The Value of Shooting: The economic, environmental and social benefits of*

shooting sports to the UK. PACEC. Cambridge, UK.

R Core Team. (2021). *R: A language and environment for statistical computing. R*

Foundation for Statistical Computing. R Foundation for Statistical Computing. Geneva,

Switzerland.

Robinson, R. A. (2005). *BirdFacts: profiles of birds occurring in Britain & Ireland*.

http://www.bto.org/birdfacts date accessed: 07/02/2022.

Stanbury, A., Eaton, M., Aebischer, N., Balmer, D., Brown, A., Douse, A., Lindley, P.,

Mcculloch, N., Noble, D., & Win, I. (2021). The status of our bird populations: the fifth

Birds of Conservation Concern in the United Kingdom, Channel Islands and Isle of Man

and second IUCN Red List assessment of extinction risk for Great Britain. *British Birds*,

114(December 2021), 723–747.

Statistics Iceland. (2022). *Online database: https://statice.is* date accessed: 07/02/2022

Thomas, J. (2021). *Unfair game: the birds that can still be shot in the UK*. Bird Guides.

https://www.birdguides.com/articles/conservation/unfair-game-the-birds-that-can-still-

be-shot-in-the-uk/ date accessed: 07/02/2022.

U.S. Fish and Wildlife Service. (2018). *Waterfowl population status, 2018*. U.S. Department

of the Interior. Washington D.C., USA.

UK Government and Parliament. (2016). *Introduce a moratorium on the hunting of critically*

declining wading birds. https://petition.parliament.uk/archived/petitions/167410 date

accessed: 07/02/2022.

Welsh Parliament. (2020). *Ban the shooting of critically endangered birds...give them the*

protection they so desperately need! Petitions.

https://petitions.senedd.wales/petitions/244712 date accessed: 07/02/2022.

Wilkins, K. A., & Cooch, E. G. (1999). *Waterfowl population status, 1999.* U.S. Fish & Wildlife

Service, Department of the Interior. Washington D.C., USA.

Williams, B. K., & Brown, E. D. (2014). Adaptive management: From more talk to real action.

Environmental Management, 53(2), 465–479. https://doi.org/10.1007/s00267-013-0205-

7

Woodward, I. D., Massimino, D., Hammond, M. J., Barber, L., Barimore, C., Harris, S. J.,

Leech, D. I., Noble, D. G., Walker, R. H., Baillie, S. R., & Robinson, R. A. (2020).

BirdTrends 2020: trends in numbers, breeding success and survival for UK breeding

birds. BTO Research Report 732. BTO. Thetford, UK.

Table 1: estimated mean λ_{max}, potential excess growth (PEG; reported to two significant figures), sustainable hunt index (SHI) and the
probability of current harvest levels being unsustainable with a safety factor (F_s) = 0.5 for huntable waterbird species populations overwintering
in the United Kingdom, except for pochard where F_s = 0.1. The mean potential excess growth is also displayed as a proportion of the population
estimate.
598

Species	Estimated λ_{max} (95% CI)	Mean Potential Excess Growth (95% CI)	PEG as proportion of population	Mean Sustainable Hunt Index (95% CI)	Probability of unsustainable harvest
Mallard	1.979 (1.886 – 2.068)	1,600,0000 (1,500,000 – 1,800,000)	0.49	0.596 (0.441 – 0.764)	0.000
Teal	2.112 (1.900 – 2.302)	240,000 (190,000 – 290,000)	0.56	0.610 (0.376 – 0.894)	0.004
Wigeon	2.224 (1.902 – 2.307)	250,000 (210,00 – 300,000)	0.56	0.222 (0.089 – 0.375)	0.000
Gadwall	1.818 (1.447 – 2.197)	13,000 (6900 – 19,000)	0.41	0.522 (0.184 – 1.099)	0.043
Pintail	1.916 (1.603 – 2.213)	9200 (5900 – 13,000)	0.46	0.136 (0.027 – 0.286)	0.000
Shoveler	2.047 (1.880 – 2.204)	10,000 (8300 – 12,000)	0.52	0.241 (0.091 – 0.414)	0.000
Tufted duck	1.853 (1.784 – 1.920)	60,000 (55,000 – 65,000)	0.43	0.098 (0.048 – 0.151)	0.000
Pochard	1.947 1.866 – 2.028)	2700 (2400 – 3100)	0.09	0.209 (0.058 – 0.372)	0.000

Goldeneye	1.473 (1.334 – 1.232)	5000 (3500 – 6500)	0.24	0.218 (0.030 – 0.454)	0.000
Canada goose	1.225 (1.218 – 1.232)	19,000 (17,000 – 20,000)	0.11	1.826 (0.811 – 2.881)	0.885
Greylag goose (Icelandic)	1.185 (1.133 – 1.231)	8400 (6000 – 11,000)	0.09	2.732 (2.047 – 3.793)	1.000
Greylag goose (British)	1.185 (1.133 – 1.231)	13,000 (9300 – 16,000)	0.09	4.101 (0.851 – 8.123)	0.952
Pink-footed goose	1.188 (1.173 – 1.202)	48,000 (39,000 – 57,000)	0.09	0.436 (0.073 – 0.859)	0.000
Woodcock (Resident)	1.354 (1.310 – 1.393)	36,000 (25,000 – 49,000)	0.22	0.540 (0.354 – 0.812)	0.00
Woodcock (Migratory)	1.354 (1.310 – 1.393)	270,000 (180,000 – 370,000)	0.22	0.482 (0.317 – 0.722)	0.00
Common Snipe	1.393 (1.327 – 1.446)	220,000 (170,000 – 270,000)	0.25	0.294 (0.109 – 0.524)	0.000
Golden plover	1.517 (1.404 – 1.626)	110,000 (80,000 – 130,000)	0.41	0.020 (0.003 – 0.041)	0.000