Engineering bacterial blight-resistant plants through CRISPR/Cas9-targeted editing of the *MeSWEET10a* promoter in cassava

Yajie Wang [a,1], Mengting Geng[b,1], Ranran Pan[d,1], Tong Zhang[b], Xiaohua Lu[b], Xinghou Zhen[b], Yannian Che[b], Ruimei Li[a,c],

Jiao Liu[a,c], Yinhua Chen[b,*], Jianchun Guo[a,c*], Yuan Yao[a,c*]

[a] *Key Laboratory of Biology and Genetic Resources of Tropical Crops, Ministry of Agriculture, Institute of Tropical Bioscience and Biotechnology, Chinese*

  Academy of Tropical Agricultural Sciences & Key Laboratory for Biology and Genetic Resources of Tropical Crops of Hainan Province, Hainan Institute for

  Tropical Agricultural Resources, Haikou 571101, PR China

[b] *Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresource, College of Tropical Crops, Hainan University, Haikou 570228, China*

[c] *Hainan Yazhou Bay Seed Lab, Sanya 572025, PR China*

[d] *College of Agriculture & Biotechnology, Zhejiang University, Hangzhou 310058, PR China*

* Corresponding author.

  E-mail address: yhchen@hainanu.edu.cn (Y. Chen); guojianchun@itbb.org.cn (J. Guo); yaoyuan@itbb.org.cn (Y.Yao);

[1] These authors contributed equally to this work.

Abstract: Cassava starch is a widely used raw materials for industrial production. However, cassava bacterial blight (CBB) caused by *Xanthomonas axonopodis* pv. *manihotis* (*Xam*) results in severe yield losses and is the most destructive bacterial disease in all worldwide cassava planting regions. This study showed that editing of the promoter in the disease-susceptibility gene *MeSWEET10a* of SC8 cassava confers resistance to CBB. All mutated cassava lines had normal morphological and yield-related traits as the wild type. The results lay a research foundation for breeding cassava resistant to bacterial blight.

Keywords: Cassava; Cassava bacterial blight; CRISPR/Cas9; *MeSWEET10a* promoter

Highlights:

MeSWEET10a gene in SC8 cultivar is hijacked by TALE20 from *Xam11* strain.

Editing of the *MeSWEET10a* promoter in SC8 cultivar confers resistance to CBB.

All the mutated cassava lines had similar yield-related traits compared to wild-type.

1. Introduction

Starch is one of the most important polymers that has been extensively used in industry. Cassava is a major available source of commercially produced starches in the market. It is extensively cultivated throughout the tropical and subtropical regions.

However, cassava bacterial blight (CBB) caused by *Xanthomonas axonopodis* pv. *manihotis* (*Xam*) results in severe yield losses and

is the most destructive bacterial disease in all worldwide cassava planting regions (McCallum et al., 2017). The vast majority of

cassava cultivars are not resistant to CBB. *Xam* pathogen infection secretes transcription activator-like effectors (TALEs) into

cassava host cells through the type III secretion system (Boch and Bonas, 2010). The repeat variable diresidues (RVDs) in TALEs

recognize the effector-binding elements (EBEs) on the host susceptibility (S) gene promoters and activate S gene expression which

facilitates bacterial growth and disease symptom formation. *Xam668* is a highly virulent *Xam* strain to cassava which contains five

transcription activator-like effectors (TALEs) including $TALE20_{Xam668}$ (Bart et al., 2012). The repeat variable diresidues (RVDs) in

$TALE20_{Xam668}$ recognize the effector-binding element (EBE) on the *MeSWEET10a* promoter from cassava 60444 cultivar and

activate *MeSWEET10a* gene expression which facilitates sucrose transport from the interior of plant cells to the apoplasts and

provide carbon sources for the bacteria (Cohn et al., 2014). Thus, editing the EBE in the *MeSWEET10a* promoter may potentially

confer resistance to CBB in cassava cultivars.

Xam11 is a highly pathogenic subspecies from Hainan, China that infects the Chinese local planting cassava SC8 cultivar

(*Manihot esculenta* Crantz South China 8) with marked pathogenic symptoms. In this study, the CRISPR/Cas9 system was used to

edit the EBE_{TALE20} of the *MeSWEET10a* promoter in SC8 cultivar. The EBE_{TALE20} mutant plants showed enhanced resistance to

Xam11, and did not show any significant differences in major yield-related traits compared to wild-type plants.

2. Materials and Methods

2.1. Plant Materials

For cassava transformation, the SC8 cultivar was used as the recipient. For pathogen inoculation, SC8 and CRISPR/Cas9

mutant plants were cultivated in a greenhouse under a 16-h/8-h light/dark cycle at 23-28□. For agronomic trait (weight and num-

ber of tuber root per plant, dry matter ratio, starch content) evaluation, all of the tested mutant plants and WT were planted in the

15-cm round pots containing 10% Perlite, 80% soil.

2.2. Dual LUC Reporter Gene Assay

Genomic DNA extraction from SC8 leaves was performed according to the Plant DNA Kit. The promoter sequence of

MeSWEET10a from SC8 was amplified by using the specific primers (F: 5'-AACTTTTAGAATGAGCCCTTG-3'; R:

5'-TTCTCCGGCTATAGTAGAGAC-3'), then inserted into pGreenII0800-LUC vector, named as pGreen

II0800-pMeSWEET10a-LUC vector. The EBE_{TALE20} from the *MeSWEET10a* promoter was deleted and generated

pSWEET10a-EBE, then inserted into pGreenII0800-LUC, named as pGreenII0800-pMeSWEET10a- EBE_{TALE20}-LUC vector.

Full-length CDS of $TALE20_{xam11}$ was inserted into pGreen II62-SK vector to generate pGreen II62-SK-$TALE20_{Xam11}$ vector. The

vectors of pGreen II62-SK or pGreen II62-SK-TALE20$_{Xam11}$ were used as the effectors, and pGreen II0800-pMeSWEET10a-LUC

or pGreen II0800-pMeSWEET10a-EBE$_{TALE20}$-LUC were used as the reporters. The effector and reporter vectors were transformed

into tobacco leaves as previously described (Liu et al., 2019b). The activity of Firefly luciferase and the reference Renilla luciferase

were quantified using the Dual-Luciferase Reporter Assay System (Promega), according to the manufacturer's instructions. Three

biological repeats were analyzed.

2.3. Editing of MeSWEET10a EBE$_{TALE20}$ by CRISPR/Cas9 System

In our previous study, the target sgRNA of *MeSWEET10a* EBE$_{TALE20}$ has been designed and constructed the vector of

pCAMBIA1301-Cas9-EBE-sgRNA, and the editing function of target sgRNA has been validated in cassava FEC by transient

transformation (Zhang et al., 2021). In our current research, the friable embryogenic callus (FEC) of SC8 cassava was infected by

Agrobacterium LBA4404 harboring editing vector of pCAMBIA1301-Cas9-EBE-sgRNA. The transgenic positive lines were

produced according to the protocol described by Nyaboga et al. (2015) (Nyaboga et al., 2015). The transformation conditions of

cassava SC8 were *Agrobacterium* strain LBA4404 cell infection (density OD$_{600}$ = 0.65), 250 µM acetosyringone induction, and

agro-cultivation with wet FEC for 3 days in dark. To analyze the genotypes of the CRISPR/Cas9 mutants, genomic DNA from the

transgenic positive lines were extracted, and analyzed using the Hi-TOM program for high-throughput mutations (Liu et al.,

2019a).

2.4. Plant Inoculation Xam11 and Symptom Observation

Xanthomonas strain *Xam11* was resuspended in 10 mM MgCl$_2$. For watersoaking assays and qRT-PCR analyses, the leaves

of SC8 and mutant lines (#1, #2, #3, #4) were injected by *Xam11* via a 1-mL needleless syringe. 5 µL of the bacterial suspension

at OD$_{600}$ = 0.1 density was injected to the back side of leaves. Three biological repeats were analyzed. The expression of

MeSWEET10a gene was measured at 0, 0.5, 1, 3, and 7 dpi time points. Symptom development after *Xam11* inoculation was pho-

tographed at 3, 7, 14, and 40 dpi. The leaf disease areas after *Xam11* inoculation were measured at 3, 7, and 14 dpi. Bacterial

numbers of *Xam11* in the SC8 and mutant lines were counted according to the method of Li et al. (2018)(Li et al., 2018).

2.5. qRT-PCR

To analyze *MeSWEET10a* expressions in SC8 cassava and mutant lines, total RNAs isolated from the leaves of SC8 and

mutant lines (#1, #2, #3, #4) using RNAplant Plus reagent (TianGen, Beijing, China) following the manufacturer's instructions

were reversed by using the reverse transcriptase kit (TaKaRa, Dalian, China). The qRT-PCR was performed with the reversed

cDNAs as substrates and the *MeSWEET10a* specific primers (F: 5'-TCCTCACCTTGACTGCGCTG-3'; R:

5'-AGCACCATCTGGACAATCCCA-3') by using the SYBR Premix Taq Kit (TaKaRa, Dalian, China) in the ABI7500 Re-

al-Time PCR System (Applied Biosystems, USA). The expression of cassava tubulin gene (Phytozome name: Manes.08G061700)

was used as an internal standard by using the primers (F: 5'-GTGGAGGAACTGGTTCTGGA-3'; R:

5'-TGCACTCATCTGCATTCTCC-3'). The $2^{-\Delta\Delta Ct}$ method was used for relative quantification (Livak and Schmittgen, 2001).

Expression data were collected from three biological repeats.

2.6. Statistical Analyses

89   All the data were presented as mean ± *SD* from at least three independent experiments with three replicates. Statistical anal-

90 yses were conducted by using Student's t-tests.

3. Results and discussion

3.1. MeSWEET10a Gene in SC8 Cultivar is Hijacked by TALE20$_{Xam11}$ from Xam11 Strain

TALE20 was cloned based on the genome sequencing information of *Xam11*. TALE20$_{Xam11}$ (TALE20 in *Xam11*) has 20 RVD

with variable amino acids at positions 12 and 13 in each repeat, and EBE$_{TALE20}$ is targeted by TALE20$_{Xam11}$ and is located in the -92

to -112 bp region before the *MeSWEET10a* start codon (Fig. 1A). The dual luciferase reporter assay demonstrated that TALE20$_{Xam11}$

activated the *MeSWEET10a* promoter from SC8 cultivar and initiated downstream expression of firefly luciferase; however,

TALE20$_{Xam11}$ could not activate the *MeSWEET10a* promoter lacking the EBE$_{TALE20}$ region (Fig. 1B). This result suggests that the

transcription activator-like effector TALE20$_{Xam11}$ of *Xam11* strain regulates the expression of *MeSWEET10a* genes by binding to the

EBE$_{TALE20}$ region of SC8 cultivar.

101   (**Color print**)

Fig. 1. Schematic representation of RVD in TALE20$_{Xam11}$ recognize the EBE$_{Xam11}$ on the *MeSWEET10a* promoter from cultivar cassava SC8 (**A**).

The C-terminus of TALE20$_{Xam11}$ has two nuclear localization signals (NLS) and an acidic activation domain (ADD). TALE20$_{Xam11}$ directly acti-

vates *MeSWEET10a* expression through the EBE region (**B**). SK: pGreen II62-SK vector (control vector); TALE20: pGreen

II62-SK-TALE20$_{Xam11}$ vector; pSWEET10a: pGreen II0800-pMeSWEET10a-LUC vector; pSWEET10a-EBE: pGreen

II0800-pMeSWEET10a-EBE$_{TALE20}$-LUC vector (deletion of EBE$_{TALE20}$ from the *MeSWEET10a* promoter). Data are displayed from three bio-

logical repeats (Student's *t*-test, **$P < 0.01$).

3.2. CRISPR/Cas9-Mediated Editing of EBE$_{TALE20}$ in the Promoter of MeSWEET10a

Inhibition of S gene expression by genetic engineering is an attractive strategy to improve plant disease resistance (Zaidi et al.,

2018). Disruption of EBEs in promoters of three major target rice *SWEET* genes by CRISPR-Cas9 avoids recognition by TALEs

from *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) and improves bacterial blight resistance (Xu et al., 2019). In this study, the effect of

genetically modifying the EBE in the *MeSWEET10a* promoter on CBB resistance after *Xam11* infection was investigated in SC8

cassava. The pCAMBIA1301-Cas9-EBE-sgRNA plasmid was constructed to target EBE$_{TALE20}$ of the *MeSWEET10a* promoter and

transformed into the SC8 cassava genome by *Agrobacterium*-mediated fragile embryonic callus (EFC) transformation. In total, 57

independent transgenic lines were regenerated. The EBE$_{TALE20}$ regions from all the transgenic lines were subjected to high

throughput tracking of mutation (Hi-TOM) sequences, the results showed that Cas9 activity efficiency was 98.2%. Three types of

mutations were obtained among the transgenic lines: homozygous mono-allelic (19.30%), homozygous bi-allelic (61.40%), and

heterozygous (17.54%) (Fig. 2A). The EBE$_{TALE20}$ regions in the four mutant types of transgenic cassava lines (#1, #2, #3, #4) and

wild type SC8 cassava (WT) were shown in Fig. 2B. These studies showed that the gene editing plasmid could efficiently edit the

EBE region of *MeSWEET10a* promoter in SC8 cassava and obtained high proportion of homozygous mutant lines. Up to now, there

were five genes (*nCBP-1*, *nCBP-2*, *GBSS*, *PTST1*, and *MePDS*) have been mutated in cassava by CRISPR/Cas9 technology;

however, the homozygous mutations in these genes were very low (Odipio et al., 2017; Bull et al., 2018; Gomez et al., 2019).

(**Color print**)

Fig. 2. CRISPR/Cas9-induced mutation types and frequency (**A**). Hi-TOM sequencing of the EBE$_{TALE20}$ regions of wild type (WT) non-transgenic

SC8 cassava compared with the engineered lines #1, #2, #3, and #4 (**B**). The EBE$_{TALE20}$ region of the TALE20$_{Xam11}$ target is marked in a red box in

WT; the sgRNA sequences are underlined; the protospacer adjacent motif (PAM) sites are marked in blue text; the hyphens are missing bases, and

the red letter is the inserted base.

3.3. The EBE_{TALE20} Mutant Shows Enhanced Resistance to CBB

The effects of EBE$_{TALE20}$ modification on *MeSWEET10a* expression in the mutant cassava lines were further investigated after inoculation with *Xam11*. The relative expression levels of *MeSWEET10a* in the mutant lines showed no obvious difference from that of the wild type at 0 and 0.5 day post-inoculation (dpi); however, markedly lower levels were observed in the mutant lines compared with that of the wild type at 1–7 dpi (Fig. 3), which showed that EBE$_{TALE20}$ mutations repressed *MeSWEET10a* expression after *Xam11* infection. The wild type had obvious water-soaked symptoms at 3 dpi, however, which were not observed in the mutant lines (#1, #2, #3, #4). At 7 dpi, the wild type had obvious pustule symptoms, while the mutant lines showed slightly water-soaked symptoms; at 14 dpi, the injection sites in the wild type were withered and turned yellow, while in the mutant lines , the injection sites started to wither and the plaque expanded slowly; at 40 dpi, the whole leaves in the wild type were withered and become yellow, whereas the leaves in the mutant lines only withered near the plaques (Fig. 4A). Diseased lesions in the mutant lines were significantly smaller at 3 dpi, 7 dpi, and 14 dpi than that in the wild type (Fig. 4B), it suggested that EBE$_{TALE20}$ mutations limited the expansion of CBB symptoms in cassava SC8. The bacterial growth assay showed that *Xam11* populations in the four mutant lines were significantly lesser than that in the wild type at 3 dpi and 7 dpi (Fig. 4C). These results revealed that EBE$_{TALE20}$ modification of *MeSWEET10a* in the four edited cassava lines had strong and stable resistance to CBB.

(**Color print**)

Fig. 3. Time course of *MeSWEET10a* expression in mutant lines and WT after *Xam11* inoculation. Three biological repeats were performed for each data point. Statistical tests were two-sided using Student's t-test compared to WT (**$p < 0.01$).

 (**Color print**)

Fig. 4. Symptom development after *Xam11* inoculation (**A**). Scale bar, 5 mm. Time course of leaf disease areas (**B**) and bacterial numbers (**C**) after *Xam11* inoculation. At least 10 leaves were harvested for the diseased areas with the bacterial number assay performed using three biological replicates. Statistical tests were two-sided using Student's *t*-test compared to WT (**$p < 0.01$).

3.4. The EBE$_{TALE20}$ Mutant Shows Yield-related Traits Similar to the Wild Type

Mutation of *Mildew resistance locus O* (*Mlo*) through CRISPR-Cas9 has conferred PM resistance in wheat (Wang et al., 2014). In addition, mutations of other S genes such as *eIF4E* and *MPK* enhances disease resistance (Gomez et al., 2019; Xie and Yang, 2013). However, these mutations often affect plant fitness with reduced growth, fertility, yield, and abiotic stress tolerance (Zaidi et al., 2018). Recently, Veley et al. (2022) has been reported that directed methylation to EBE$_{TALE20}$ region of *MeSWEET10a* promoter of TME419 in 60444 cassava cultivar, the plants displayed an increased resistance to *Xam668* strain, meanwhile, they maintained normal growth and development (Veley et al., 2022). In our results, there were no significant differences in tuber root traits and starch content between the EBE$_{TALE20}$ mutated lines and the wild type plants when grown under normal conditions (Fig. 5). This indicated that the EBE$_{TALE20}$ mutation in the *MeSWEET10a* promoter of SC8 cultivar has no effect on yield-related traits in cassava.

(A)

Lines	WT	#1	#2	#3	#4
Tuber root (g)	52.7±28.3a	46.1±15.1a	62.2±15.6a	60.6±10.7a	52.4±22.5a
Number of tuber roots /plant	3.6±2.1a	3.3±1.5a	4.3±1.5a	3.3±1.5a	3.3±0.6a
Dry matter ratio (%)	30.0±4.3a	32.2±3.2a	32.2±2.7a	31.9±6.5a	29.1±1.1a
Starch content (% dry wt)	65.7±11.6a	66.6±9.1a	65.3±11.5a	63.1±11.4a	65.9±5.0a

(**Color print**)

Fig. 5. Tuber root phenotypic characterizations **(A)**. Scale bar, 1 cm. Yield-related traits shown as the mean ± *SD* from five plants of each line **(B)**. The values marked with the same letter (a) represented no significant difference ($p < 0.01$).

4. Conclusions

In conclusion, editing of the promoter in the disease-susceptibility gene *MeSWEET10a* in SC8 cassava confers resistance to CBB. All the mutated cassava lines had normal morphological and yield-related traits as the wild type. The results lay a research foundation for breeding Chinese local planting cassava SC8 cultivar resistant to bacterial blight by CRISPR/Cas9 technology.

CRediT authorship contribution statement

Yajie Wang, **Mengting Geng** and **Ranran Pan:** Investigation, Conceptualization, Visualization, Writing – original draft. **Tong Zhang, Xiaohua Lu, Xinghou Zhen, Yannian Che, Ruimei Li** and **Jiao Liu:** Investigation, Validation. **Yinhua Chen**, **Jianchun Guo** and **Yao Yuan:** Writing - review & editing, Funding acquisition, Project administration, Supervision, Validation.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This research was supported by the National Key Research and Development Program of China: 2019YFD1001105; the Hainan Provincial Natural Science Foundation of China: 320RC492; the Major Science and Technology plan of Hainan Province: ZDKJ2021012; Hainan Yazhou Bay Seed Lab: B21HJ0303 and the the earmarked fund for CARS: CARS-11.

References

Bart, R., Cohn, M., Kassen, A., McCallum, E.J., Shybut, M., Petriello, A., et al., 2012. High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved effectors to target for durable resistance. Proc. Natl. Acad. Sci. 109, E1972-E1979.

Boch, J., Bonas, U., 2010. *Xanthomonas* AvrBs3 family-type III effectors: discovery and function. Annu. Rev. Phytopathol. 48, 419-436.

Bull, S.E., Seung, D., Chanez, C., Mehta, D., Kuon, J.E., Truernit, E., et al., 2018. Accelerated ex situ breeding of *GBSS-* and *PTST1*-edited cassava for modified starch. Sci. Adv. 4, eaat6086.

Cohn, M., Bart, R.S., Shybut, M., Dahlbeck, D., Gomez, M., Morbitzer, R., et al., 2014. *Xanthomonas axonopodis* virulence is promoted by a transcription activator-like effector–mediated induction of a SWEET sugar transporter in cassava. Mol. Plant Microbe Interact. 27, 1186-1198.

Gomez, M.A., Lin, Z.D., Moll, T., Chauhan, R.D., Hayden, L., Renninger, K., et al., 2019. Simultaneous CRISPR/Cas9-mediated editing of cassava eIF4E isoforms nCBP-1 and nCBP-2 reduces cassava brown streak disease symptom severity and incidence. Plant Biotechnol. J. 17, 421-434.

Li, X., Liu, W., Li, B., Liu, G., Wei, Y., He, C., et al., 2018. Identification and functional analysis of cassava DELLA proteins in plant disease resistance against cassava bacterial blight. Plant Physiol. Bioch. 124, 70-76.

Liu, Q., Wang, C., Jiao, X., Zhang, H., Song, L., Li, Y., et al., 2019a. Hi-TOM: a platform for high-throughput tracking of mutations induced by CRISPR/Cas systems. Sci. China Life Sci. 62, 1-7.

Liu, Y., Yang, T., Lin, Z., Gu, B., Xing, C., Zhao, L., et al., 2019b. A WRKY transcription factor PbrWRKY53 from Pyrus betulaefolia is involved in drought tolerance and AsA accumulation. Plant Biotechnol. J. 17, 1770-1787.

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods 25, 402-408.

Nyaboga, E.N., Njiru, J.M., Tripathi, L., 2015. Factors influencing somatic embryogenesis, regeneration, and *Agrobacterium*-mediated transformation of cassava (*Manihot esculenta* Crantz) cultivar TME14. Front. Plant Sci. 6, 411.

Odipio, J., Alicai, T., Ingelbrecht, I., Nusinow, D.A., Bart, R., Taylor, N.J., 2017. Efficient CRISPR/Cas9 genome editing of phytoene desaturase in cassava. Front. Plant Sci. 8, 1780.

Veley, K.M., Elliott, K., Jensen, G., Zhong, Z., Feng, S., Yoder, M., et al., 2022. Improving disease resistance in plants by editing the epigenome. BioRxiv.

Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C., et al., 2014. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. Nat. Biotechnol. 32, 947-951.

McCallum, E.J., Anjanappa, R.B., Gruissem, W., 2017. Tackling agriculturally relevant diseases in the staple crop cassava (*Manihot esculenta*). Curr. Opin. Plant Biol. 38, 50-58.

Xie, K., Yang, Y., 2013. RNA-guided genome editing in plants using a CRISPR-Cas system. Mol. Plant 6, 1975-1983.

Xu, Z., Xu, X., Gong, Q., Li, Z., Li, Y., Wang, S., et al., 2019. Engineering broad-spectrum bacterial blight resistance by simultaneously disrupting variable TALE-binding elements of multiple susceptibility genes in rice. Mol. Plant 12,1434-1446.

Zaidi, S.S.E.A., Mukhtar, M.S., Mansoor, S., 2018. Genome editing: targeting susceptibility genes for plant disease resistance. Trends Biotechnol. 36, 898-906.

Zhang, T., Wang, Y., Guo, W., Li, R., Liu, J., Guo, J., et al., 2021. Construction and verification of editing vector for EBE region of MeSWEET10a promoter in cassava. Chin. J. Trop. Crops 42, 3120.