Title: Engineering bacterial blight-resistant plants through CRISPR/Cas9-targeted editing of the *MeSWEET10a* promoter in cassava

Yajie Wang[1,2,4], Mengting Geng[2,4], Ranran Pan[3,4], Tong Zhang[2], Xinghou Zhen[2], Yannian Che[2], Ruimei Li[1], Jiao Liu[1], Yinhua Chen*[2], Jianchun Guo*[1], Yuan Yao*[1]

[1]Key Laboratory of Biology and Genetic Resources of Tropical Crops, Ministry of Agriculture, Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, China

[2]Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresource, Hainan University, Haikou, China

[3]College of Agriculture & Biotechnology, Zhejiang University, Hangzhou, China

[4]These authors contributed equally to the article.

*Correspondence: Yinhua Chen (yhchen@hainu.edu.cn), **Jianchun Guo** (guojianchun@itbb.org.cn), **Yuan Yao** (yaoyuan@itbb.org.cn)

Abstract:

Editing of the promoter in the disease-susceptibility gene *MeSWEET10a* of SC8 cassava confers resistance to cassava bacterial blight (CBB). All mutated cassava lines had normal morphological and yield-related traits as the wild type. The results lay a research foundation for breeding cassava resistant to bacterial blight.

Dear Editor,

Cassava (*Manihot esculenta* Crantz) is a major food crop in tropical and

subtropical regions that provides food security for 800 million people. However,

cassava bacterial blight (CBB) caused by *Xanthomonas axonopodis* pv. *manihotis*

(*Xam*) results in severe yield losses and is the most destructive bacterial disease in all

worldwide cassava planting regions (McCallum et al., 2017). The vast majority of

cassava cultivars are not resistant to CBB. *Xam* pathogen infection secretes

transcription activator-like effectors (TALEs) into cassava host cells through the type

III secretion system (Boch and Bonas, 2010). The repeat variable diresidues (RVDs)

in TALEs recognize the effector-binding elements (EBEs) on the host susceptibility

(S) gene promoters and activate S gene expression which facilitates bacterial growth

and disease symptom formation.

Inhibition of S gene expression by genetic engineering is an attractive strategy to

improve plant disease resistance (Zaidi et al., 2018). *Mildew resistance locus O* (*Mlo*)

is one of the best-known S genes and encodes a conserved membrane-associated

protein. Mlo is required for the penetration of powdery mildew (PM) fungi into host

epidermal cells. Mutation of *Mlo* through CRISPR-Cas9 has conferred PM resistance

in wheat (Wang et al., 2014). In addition, mutations of other S genes such as *eIF4E*

and *MPK* enhances disease resistance (Gomez et al., 2019; Xie and Yang, 2013).

However, these mutations often affect plant fitness with reduced growth, fertility,

yield, and abiotic stress tolerance (Zaidi et al., 2018). Genetic engineering of the core

S gene promoter elements recognized by pathogenic TALEs improves disease

resistance without affecting plant characteristics in citrus and rice (Peng et al., 2017;

Xu et al., 2019). *CsLOB1* is a citrus canker S gene and plays a key role in promoting

pathogen growth and blistering pustule formation. Deletion of the EBE$_{PthA4}$ sequence

from the *CsLOB1* gene promoter by CRISPR/Cas9 has conferred high resistance to

citrus canker (Peng et al., 2017). Sugars will eventually be exported transporters

(*SWEET*) genes are the other best-known S genes encoding sugar transporters. Some

SWEET genes can be hijacked by pathogenic bacteria to induce sucrose transport from

the interior of plant cells to the apoplasts and provide carbon sources for the bacteria.

Disruption of EBEs in promoters of three major target rice *SWEET* genes by

CRISPR-Cas9 avoids recognition by TALEs from *Xanthomonas oryzae* pv. *oryzae*

(*Xoo*) and improves bacterial blight resistance (Xu et al., 2019).

Generally, *Xam* contains 1-5 TALEs which determine its virulence (Bart et al.,

2012). TALE20 contributes toward most of the aggressiveness and virulence of *Xam*

in cassava (Cohn et al., 2014) by targeting a *SWEET* gene. *MeSWEET10a* expression

rapidly increases after infection with Xam_{TALE20} (*Xam* containing TALE20) which

induces sucrose translocation to the infection site and facilitates the multiplication and

spread of *Xam* in susceptible cassava varieties, but not in resistant varieties (Mora et

al., 2019). *Xam668* is a highly virulent *Xam* strain to cassava which contains five

TALEs including TALE20 (Bart et al., 2012). TALE20 disruption in *Xam668*

(*Xam668ΔTAL20*) reduces cassava water-soaking symptom development, while

reintroducing TALE20 into the plasmid-cured strain fully complemented the

symptom defect phenotype (Cohn et al., 2014). $TALE20_{Xam668}$ has 20 RVDs, and

directly targets the EBE in the *MeSWEET10a* promoter (Cohn et al., 2014). Thus,

editing the EBE in the *MeSWEET10a* promoter may potentially confer resistance to

CBB in cassava (Lin et al., 2019).

Xam11 is a highly pathogenic subspecies from Hainan, China that infects the

cassava SC8 cultivar with marked pathogenic symptoms (Figure S1). In this study, the

effect of genetically modifying the EBE in the *MeSWEET10a* promoter on CBB

resistance in cassava after *Xam11* infection was investigated. *TALE20* was cloned

based on the genome sequencing information of *Xam11*. $TALE20_{Xam11}$ (TALE20 in

Xam11) has 20 RVD with variable amino acids at positions 12 and 13 in each repeat

(Figure S2). The promoter sequence of *MeSWEET10a* in SC8 cassava was cloned,

which is identical to the reported sequence in Mora et al. (2019). EBE_{TALE20} is

targeted by $TALE20_{Xam11}$ and is located in the -92 to -112 bp region before the

MeSWEET10a start codon (Figure S3). The dual luciferase reporter assay

demonstrated that TALE20$_{Xam11}$ activated the *MeSWEET10a* promoter and initiated

downstream expression of firefly luciferase; however, TALE20$_{Xam11}$ could not activate

the *MeSWEET10a* promoter lacking the EBE$_{TALE20}$ region (Figure 1A).

CRISPR-P   v2.0   was   used   to   design   the   sgRNA

(TGGGCGAGAAGCGTTTATATA**GG**) targeting EBE$_{TALE20}$ based on the sequence

characteristics   of   the   *MeSWEET10a*   promoter   in   SC8   cassava.   The

pCAMBIA1301-Cas9-EBE-sgRNA plasmid was constructed and transformed into the

SC8 cassava genome by *Agrobacterium*-mediated fragile embryonic callus (EFC)

transformation. In total, 57 independent transgenic lines were regenerated after

hygromycin (20 μg/mL) selection and confirmation by PCR analysis for the *Cas9*

gene (Figure S4). The EBE$_{TALE20}$ regions from all the transgenic lines subjected to

Sanger sequencing and high throughput tracking of mutation (Hi-TOM) sequencing

(Figure S5, Table S1) showed that Cas9 activity efficiency was 98.2%. Three types of

mutations were obtained among the transgenic lines: homozygous mono-allelic

(19.30%), homozygous bi-allelic (61.40%), and heterozygous (17.54%) (Table S1).

The EBE$_{TALE20}$ regions in the four mutant types of transgenic cassava (#1, #2, #3, #4)

and wild type SC8 cassava (WT) were shown in Figure 1B. The effects of EBE$_{TALE20}$

modification   on   *MeSWEET10a*   expression   in   the   mutant   cassava   lines   were

investigated   after   inoculation   with   *Xam11*.   The   relative   expression   level   of

MeSWEET10a in mutant lines showed no obvious difference from that of the wild

type at 0 and 0.5 day post-inoculation (dpi); however, markedly lower levels were

observed in mutant lines compared with that of the wild type at 1−7 dpi (Figure 1C).

This showed that EBE$_{TALE20}$ mutations repressed *MeSWEET10a* expression after

Xam11 infection. Wild type had obvious water-soaked symptoms at 3 dpi which were

not observed in mutant lines (#1, #2, #3, #4). At 7 dpi, the wild type had obvious

pustule symptoms while the mutant lines showed slightly water-soaked symptoms; at

14 dpi, the wild type injection sites were withered and turned yellow, while the

mutant injection sites started to wither and the plaque expanded slowly; at 40 dpi,

whole leaves of the wild type were withered and become yellow whereas the leaves of

the mutant lines only withered near the plaques (Figure 1D). Diseased lesions in

mutant lines were significantly smaller at 3 dpi, 7 dpi, and 14 dpi (Figure 1E)

suggesting that EBE_{TALE20} mutations limited the expansion of CBB symptoms in

cassava. The bacterial growth assay showed that *Xam11* populations in the four

mutant lines were significantly smaller than that in the wild type at 3 dpi and 7 dpi

(Figure 1F). Callose is deposited on the cassava cell wall during the early stages of

Xam pathogen attack which restricts pathogen growth (Wei et al., 2021). The leaves

of the mutant cassava lines displayed significantly more callose depositions at 3 dpi

than that in the wild type (Figure 1G). These results revealed that EBE_{TALE20}

modification of *MeSWEET10a* in the four edited cassava lines had strong and stable

resistance to CBB.

There were no significant differences in tuber root traits and starch content

between the EBE_{TALE20} mutated cassava lines and the wild type plants when grown

under normal conditions (Figure 1H and 1I). This indicated that the EBE_{TALE20}

mutation in the *MeSWEET10a* promoter has no effect on yield-related traits in

cassava.

In conclusion, editing of the promoter in the disease-susceptibility gene

MeSWEET10a of SC8 cassava confers resistance to CBB. All mutated cassava lines

had normal morphological and yield-related traits as the wild type. The results lay a

research foundation for breeding cassava resistant to bacterial blight.

SUPPLEMENTAL INFORMATION

Supplemental Information is available at *Molecular Plant* Online.

FUNDING

This work was supported by grants from the National Key R&D Program of China

(2019YFD1001105), the Hainan Provincial Natural Science Foundation of China

(320RC492), and the Earmarked Fund for China Agriculture Research System

(CARS-11-HNGJC, CARS-11-HNCYH).

AUTHOR CONTRIBUTIONS

Y.Y., J.G., and Y.C. designed the experiments. Y.W., M.G., R.P., X.Z., R.L., and J.L.

performed the experiments. T.Z. and Y.C. conducted the cassava transformation. Y.Y.,

Y.W., M.G., and R.P. analyzed the data and wrote the manuscript. J.G. and Y.C.

revised the manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGEMENTS

The authors declare no conflict of interest.

REFERENCES

Bart, R., Cohn, M., Kassen, A., McCallum, E.J., Shybut, M., Petriello, A.,

Krasileva, K., Dahlbeck, D., Medina, C., Alicai, T., et al. (2012). High-throughput

genomic sequencing of cassava bacterial blight strains identifies conserved effectors

to target for durable resistance. Proceedings of the National Academy of Sciences of

the United States of America **109**:E1972-E1979.

https://doi.org/10.1073/pnas.1208003109.

Boch, J., and Bonas, U. (2010). *Xanthomonas* AvrBs3 family-type III effectors:

discovery and function. Annual Review of Phytopathology **48**:419-436.

https://doi.org/10.1146/annurev-phyto-080508-081936.

Cohn, M., Bart, R.S., Shybut, M., Dahlbeck, D., Gomez, M., Morbitzer, R.,

Hou, B.-H., Frommer, W.B., Lahaye, T., and Staskawicz, B.J. (2014).

Xanthomonas axonopodis virulence is promoted by a transcription activator-like

effector–mediated induction of a SWEET sugar transporter in cassava. Molecular

Plant-Microbe   Interactions   **27**:1186-1198.

https://doi.org/10.1094/MPMI-06-14-0161-R.

Gomez, M.A., Lin, Z.D., Moll, T., Chauhan, R.D., Hayden, L., Renninger, K.,

Beyene, G., Taylor, N.J., Carrington, J.C., and Staskawicz, B.J. (2019).

Simultaneous CRISPR/Cas9‐mediated editing of cassava eIF 4E isoforms nCBP‐1

and nCBP‐2 reduces cassava brown streak disease symptom severity and incidence.

Plant Biotechnology Journal **17**:421-434. https://doi.org/10.1111/pbi.12987.

Lin, Z.J.D., Taylor, N.J., and Bart, R. (2019). Engineering disease-resistant

cassava.   Cold   Spring   Harbor   Perspectives   in   Biology   **11**:a034595.

https://doi.org/10.1101/cshperspect.a034595.

McCallum, E.J., Anjanappa, R.B., and Gruissem, W. (2017). Tackling

agriculturally relevant diseases in the staple crop cassava (*Manihot esculenta*).

Current Opinion in Plant Biology **38**:50-58. https://doi.org/10.1016/j.pbi.2017.04.008.

Mora, R.E., Rodriguez, M.A., Gayosso, L.Y., and López, C.E. (2019). Using

in vitro plants to study the cassava response to *Xanthomonas phaseoli* pv. *manihotis*

infection.   Tropical   Plant   Pathology   **44**:423-429.

https://doi.org/10.1007/s40858-019-00296-x.

Peng, A., Chen, S., Lei, T., Xu, L., He, Y., Wu, L., Yao, L., and Zou, X. (2017).

Engineering canker‐resistant plants through CRISPR/Cas9‐targeted editing of the

susceptibility gene CsLOB1 promoter in citrus. Plant Biotechnology Journal

15:1509-1519. https://doi.org/10.1111/pbi.12733.

Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C., and Qiu, J.-L.

(2014). Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers

heritable resistance to powdery mildew. Nature Biotechnology **32**:947-951.

https://doi.org/10.1038/nbt.2969.

Wei, Y., Zeng, H., Liu, W., Cheng, X., Zhu, B., Guo, J., and Shi, H. (2021).

Autophagy‐related genes serve as heat shock protein 90 co-chaperones in disease

resistance against cassava bacterial blight. The Plant Journal*: for Cell And Molecular

Biology **107**:925-937. https://doi.org/10.1111/tpj.15355.

Xie, K., and Yang, Y. (2013). RNA-guided genome editing in plants using a

CRISPR–Cas   system.   Molecular   Plant   **6**:1975-1983.

https://doi.org/10.1093/mp/sst119.

Xu, Z., Xu, X., Gong, Q., Li, Z., Li, Y., Wang, S., Yang, Y., Ma, W., Liu, L.,

Zhu, B., et al. (2019). Engineering broad-spectrum bacterial blight resistance by

simultaneously disrupting variable TALE-binding elements of multiple susceptibility

genes   in   rice.   Molecular   Plant   **12**:1434-1446.

https://doi.org/10.1016/j.molp.2019.08.006.

Zaidi, S.S.-e.-A., Mukhtar, M.S., and Mansoor, S. (2018). Genome editing:

targeting susceptibility genes for plant disease resistance. Trends in Biotechnology

36:898-906. https://doi.org/10.1016/j.tibtech.2018.04.005.

Figures

Figure 1. Engineering CBB resistant plants through CRISPR/Cas9-targeted editing of the *MeSWEET10a* promoter in cassava. (A) TALE20$_{Xam11}$ directly activates *MeSWEET10a* expression through the EBE region. *MeSWEET10a* promoter-luciferase fusion and 35S-driven TALE20$_{Xam11}$ constructs were delivered into *Nicotiana benthamiana* cells through infection by *Agrobacterium tumefaciens*. Luciferase activity was quantified after 1 dpi. SK: pGreen II62-SK vector (control vector); TAL20: pGreen II62-SK-TAL20$_{Xam11}$ vector; pSWEET10a: pGreen II0800-pMeSWEET10a-LUC vector; pSWEET10a-EBE: pGreen II0800-pMeSWEET10a-EBE$_{TAL20}$-LUC vector (deletion of EBE$_{TAL20}$ from the *MeSWEET10a* promoter). Data are displayed as mean technical replicates ± standard deviation. Experiments were repeated three times with similar results (Student's t-test, ***P < 0.005). (B) Hi-TOM sequencing of the EBE$_{TALE20}$ regions of wild type (WT) non-transgenic SC8 cassava compared with the engineered lines #1, #2, #3, and #4. The EBE$_{TALE20}$ region of the TALE20$_{Xam11}$ target is marked in a red box in WT; the sgRNA sequences are underlined; the protospacer adjacent motif (PAM) sites are

marked in blue text; the hyphens are missing bases, and the red letter is the inserted

base. **(C)** Time course of *MeSWEET10a* expression in mutant lines and WT after

Xam11 inoculation. *MeSWEET10a* transcript levels in leaves determined by qPCR at

0, 0.5, 1, 3, and 7 dpi. Three biological repeats were performed for each data point

(Student's t-test, ***P < 0.005). **(D)** Symptom development after *Xam11* inoculation.

Scale bar, 5 mm. **(E)** and **(F)** Time course of leaf disease areas and bacterial numbers

after *Xam11* inoculation. At least 10 leaves were harvested for the diseased areas with

the bacterial number assay performed using three biological replicates (Student's

t-test, *P < 0.05, ***P < 0.005). **(G)** Callose depositions in leaves. Scale bar, 50 μm.

(H) Tuber root phenotypic characterizations. Scale bar, 1 cm. **(I)** Yield-related traits

shown as the mean ± SD from five plants of each line. *P* values were calculated using

Student's t-test. The values marked with the same letter (a) represented no significant

difference.

Supplemental Figures

uninfected cassava   infected cassava

235 **Figure S1.** ***Xam11* infects cassava cultivar SC8 with marked pathogenic**

236 **symptoms.** Red arrows indicate the region affected in the infected cassava.

MDPIRPRTPSPAHELLAGPQPDRVQPQPTADRGGAPPAGSPLDGLPARRTMSRTRLPSPPAPLPAFSAGSFSD
LLRQFDPSLLDTSLFNSMSAFGAPHTEAASGEGDEVQSGLRAADDPQATVQVAVTAARPPRAKPAPRRRA
AHTSDASPAGQVDLCTLGYSQQQQEKIKLKARSTVAQHHEALIGHGFTRAHIVALSQHPAALGTVAVKYQ
AMIAALPEATHEDIVGGGKQWSGARALEALLTVSGELRGPPLQLDTGQLLKIAKRGGVTAVEAVHAWRN
ALTGAPLN

20 Repeat variable diresidues (RVD)

LTPDQVVAIAS**NI**GGKQALETVQRLLPVLCEQHG
LTLDQVVAIAS**NG**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**NI**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**NN**GGKQALETVQRLLPVLCEQHG
LTRAQVVAIAS**NG**GGKQALETVQRLLPVLRQAHG
LTPAQVVAIAS**HD**GGKQALETVQQLLPVLCEQHG
LTPAQVVAIAS**NN**GGKQALETVQRLLPVLRQAHG
LTPDQVVAIAS**HD**GGKPALETVQRLLPVLCEQHG
LTPDQVVAIAS**NN**GGKPALETVQRLLPVLCEQHG
LTRAQVVAIAS**NG**GGKQALETVQRLLPVLRQAHG
LTPAQVVAIAS**HD**GGKQALETVQQLLPVLCEQHG
LTPAQVVAIAS**NS**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**HD**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**NN**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**HD**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**HD**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**HD**GGKQALETVQRLLPVLRQAHG
LTPDQVVAIAS**NI**GGKQALETVQRLLPVLCEQHG
LTPDQVVAIAS**NG**GGKQALETVQRLLPVLCEQHG
LTPAQVVAIAS**NG**GGRPALE

SIFAQLSRPDQALAALTNDHLVALACLGGRPALEAVKKGLPHAPTLIKRTNRRLPERTSHRVADHAQVARVL
GFFQCHSHPAQAFDEAMTQFGMSRHGLLQLFRRAGVTELEAHSGTLPPASQRWHRILQASGMKRAEPSG
ASAQTPDQASLHAFADALERELDAPSPIDRAGQALASSS**RKRSRS**ESSVTGSFAQQAVEVRVPEQRDALHF
LPLSW**GVKRPR**TRIGGGLPDPGTPMDADLAASSTVMW**E**QDADPFAGAADDFPAFNEEEMAWLMELFPQ

Nuclear localization Signals (NLS)   **Acidic activation domain (AAD)**

239 **Figure S2. Analysis of TALE20 protein sequence from *Xam11*.** TALE20*Xam11* has

240 20 RVD with variable amino acids at positions 12 and 13 in each repeat (highlighted

241 in red).

```
AACTTTTAGAATGAGCCCTTGAAATGAAACCTCTAAGCTTGTAGGTTCTAATAGGGT
TACTTAATTAAATGATAGACCTAACCCTTATAATTTGATTATGACTTTTTATAATAG
AGATGAACGAAACCACAAAATTCTCCAGTTACCCAATAAACGTGGGAATGAGAGTGT
GTGGTTAGGAATCCATGTGCATGGCAAGAGAGAAAATCACAGGACTTTCTTTCAAAT
TAATTCTTGGTTAATCATGATTACATATTAATCACCTTGTCTTTTACGTGCAATCTT
CAAGTTTTTCAACCATTATTAATTACGTTGTGGACGTGGACAACTTGGAAGAAATCG
TAAAGGGCCGTCAACAAGCATTCACAAGAGCTGTTATCAGCGAGGGGGAATGAGACA
ACGGCGGCTGATGTGTTATGGTGGTAAACTGGGAAATACTAGGCACGGTTTCTTGTA
TGGGAAAAACAGCATATCCTCTGCCATGATGCTGAGCTCGTTTCTCCACCGTTTGAT
TGAATGTTCTGATTTGTTTCATCATCATCAAATGCGATTACACATCAGAAAGCTTGT
TGATGGGATTCTGTAACAAATGAAGTAAGCTTCAAAGCCTTTTGGATCCCTACTATG
GGAATGATCCAAGTGGGTTAACTACTATTAGTTAGTTGACCTCATCCACTGCGCTAT
AGGCTCATATAGGTCCAGTTAACTCGATCTCTGAAATTAAATTTCTATTTTTTAAAT
GAAATTAAGTCAAATGCATTATACGTACCCTTCCTCAACACCCTATATTATATTAAC
AAAAATCTTTATATTGCATTGCATCCAGTAAATGATAAAAAAAAATCAGGGTATA
ATTTTCTTAATCCATGCATGATAATCTGATTCATTTCATGTATAAATTTAAATATAA
TTTGTTGAGAAAGAGCAGGAGTCCTCTGGAACTAAAATATGAGTTGTAACCCTTAAA
ACCACTTGGTGGGTGACACATCAGAAAGAACAAAGTGGGACCCTTTGAATTTTGTGC
ACGTACTAGGAATGGTACCTACAAGTAGGGCTACTTTTGTATAGTGAGAGTTTACCT
GAAGAAAATTAACCATAGATGGATAAATATAGGGCATTAGGTTTAGAAAGCTGTTC
GGATCATTTTAACGTAAAAGCACTTAGTAGGACCAACTTAAAAGTAACCATAAAT
TAATATTGTTTTCTCTTTCTTCAAAAAAAATAATAAAAGAAACAAGGCCACTGTTAC
ATTGACATATTTTATTCACTTTAATCATGCATGCAACTTGACTTCATTCCGTTCCCT
GGATTCCTCCCCTATATAAACGCTTCTCGCCCATCCATCATTGCACAACATAGC
TAGAGTTTCCTCTTGAGAAAGAGAGTTTCCTCTGCACAAGGGAAAGAGAGTCTCTAC
TATAGCCGGAGAAATGGCCTTGCACTTGTCATTGGACTTCGTTTTTGGCGTTTTAGG
TACTA
```

Figure S5. Sanger sequencing peak map of target fragments in mutant lines #1, #2, #3, #4 and wild type (WT). Red boxes represent the protospacer adjacent motif (PAM), and underlines represent the target sequences.

257