Title: Engineering bacterial blight-resistant plants through

CRISPR/Cas9-targeted editing of the *MeSWEET10a* promoter in cassava

Yajie Wang[1,2,†], Mengting Geng[2,†], Ranran Pan[3,†], Tong Zhang[2], Xinghou Zhen[2],

Yannian Che[2], Ruimei Li[1], Jiao Liu[1], Yinhua Chen[2,*], Jianchun Guo[1,*], Yuan Yao[1,*]

[1]Key Laboratory of Biology and Genetic Resources of Tropical Crops, Ministry of

Agriculture, Institute of Tropical Bioscience and Biotechnology, Chinese Academy of

Tropical Agricultural Sciences, Haikou, China

[2]Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresource, College

of Tropical Crops, Hainan University, Haikou, China

[3]College of Agriculture & Biotechnology, Zhejiang University, Hangzhou, China

[*]Correspondence (Tel +86-898-66989251; fax +86-898-66987152; email

yaoyuan@itbb.org.cn (YY);Tel +86-898-66986031; fax +86-898-66987152; email

guojianchun@itbb.org.cn (JG); Tel +86-898-66279046; fax +86-898-66279046; email

yhchen@hainanu.edu.cn (YC))

[†] These authors contributed equally to the article.

Keywords: Cassava bacterial blight, Cassava, CRISPR/Cas9, *MeSWEET10a*

promoter.

Cassava is a major food crop in tropical and subtropical regions that provides food

security for 800 million people. However, cassava bacterial blight (CBB) caused by

Xanthomonas axonopodis pv. *manihotis* (*Xam*) results in severe yield losses and is the

most destructive bacterial disease in all worldwide cassava planting regions

(McCallum *et al.*, 2017). The vast majority of cassava cultivars are not resistant to

CBB. *Xam668* is a highly virulent *Xam* strain to cassava which contains five

transcription activator-like effectors (TALEs) including TALE20$_{Xam668}$ (Bart *et al.*,

2012). The repeat variable diresidues (RVDs) in TALE20$_{Xam668}$ recognize the

effector-binding element (EBE) on the *MeSWEET10a* promoter from cassava 60444

cultivar and activate *MeSWEET10a* gene expression which facilitates sucrose

transport from the interior of plant cells to the apoplasts and provide carbon sources

for the bacteria (Cohn *et al.*, 2014). Thus, editing the EBE in the *MeSWEET10a*

promoter may potentially confer resistance to CBB in cassava.

Xam11 is a highly pathogenic subspecies from Hainan, China that infects the

Chinese local planting cassava SC8 cultivar with marked pathogenic symptoms

(Figure 1A). *TALE20$_{Xam11}$* was cloned based on the genome sequencing information of

Xam11. TALE20$_{Xam11}$ has 20 RVD, and EBE$_{TALE20}$ is targeted by TALE20$_{Xam11}$ and is

located in the -92 to -112 bp region before the *MeSWEET10a* start codon (Figure 1B).

The dual luciferase reporter assay demonstrated that TALE20$_{Xam11}$ activated the

MeSWEET10a promoter from SC8 cultivar and initiated downstream expression of

firefly luciferase; however, TALE20$_{Xam11}$ could not activate the *MeSWEET10a*

promoter lacking the EBE$_{TALE20}$ region (Figure 1C).

CRISPR-P v2.0 was used to design the sgRNA targeting EBE$_{TALE20}$ of the

MeSWEET10a promoter in SC8 cassava. The pCAMBIA1301-Cas9-EBE-sgRNA

plasmid was constructed and transformed into the SC8 cassava genome by

Agrobacterium-mediated fragile embryonic callus (EFC) transformation. In total, 57

independent transgenic lines were regenerated. The EBE$_{TALE20}$ regions from all the

transgenic lines subjected to high throughput tracking of mutation (Hi-TOM)

sequencing showed that Cas9 activity efficiency was 98.2%. Three types of mutations

were obtained among the transgenic lines: homozygous mono-allelic (19.30%),

homozygous bi-allelic (61.40%), and heterozygous (17.54%) (Figure 1D). The

EBE_{TALE20} regions in the four mutant types of transgenic cassava (#1, #2, #3, #4) and

wild type SC8 cassava (WT) were shown in Figure 1E. The effects of EBE_{TALE20}

modification on *MeSWEET10a* expression in the mutant cassava lines were

investigated after inoculation with *Xam11*. The relative expression level of

MeSWEET10a in mutant lines showed no obvious difference from that of the wild

type at 0 and 0.5 day post-inoculation (dpi); however, markedly lower levels were

observed in mutant lines compared with that of the wild type at 1–7 dpi (Figure 1F).

This showed that EBE_{TALE20} mutations repressed *MeSWEET10a* expression after

Xam11 infection. Wild type had obvious water-soaked symptoms at 3 dpi which were

not observed in mutant lines (#1, #2, #3, #4). At 7 dpi, the wild type had obvious

pustule symptoms while the mutant lines showed slightly water-soaked symptoms; at

14 dpi, the wild type injection sites were withered and turned yellow, while the

mutant injection sites started to wither and the plaque expanded slowly; at 40 dpi,

whole leaves of the wild type were withered and become yellow whereas the leaves of

the mutant lines only withered near the plaques (Figure 1G). Diseased lesions in

mutant lines were significantly smaller at 3 dpi, 7 dpi, and 14 dpi (Figure 1H)

suggesting that EBE_{TALE20} mutations limited the expansion of CBB symptoms in

cassava. The bacterial growth assay showed that *Xam11* populations in the four

mutant lines were significantly smaller than that in the wild type at 3 dpi and 7 dpi

(Figure 1I). Callose is deposited on the cassava cell wall during the early stages of

Xam pathogen attack which restricts pathogen growth (Wei *et al.*, 2021). The leaves

of the mutant cassava lines displayed significantly more callose depositions at 3 dpi

than that in the wild type (Figure 1J). These results revealed that EBE_{TALE20}

modification of *MeSWEET10a* in the four edited cassava lines had strong and stable

resistance to CBB.

There were no significant differences in tuber root traits and starch content between the EBE$_{TALE20}$ mutated cassava lines and the wild type plants when grown under normal conditions (Figure 1K and 1L). This indicated that the EBE$_{TALE20}$ mutation in the *MeSWEET10a* promoter has no effect on yield-related traits in cassava.

In conclusion, editing of the promoter in the disease-susceptibility gene *MeSWEET10a* of SC8 cassava confers resistance to CBB. All mutated cassava lines had normal morphological and yield-related traits as the wild type. The results lay a research foundation for breeding cassava resistant to bacterial blight.

Acknowledgements

This work was supported by grants from the the National Key Research and Development Program of China (2019YFD1001105), the Hainan Provincial Natural Science Foundation of China (320RC492), the Major Science and Technology plan of Hainan Province (ZDKJ2021012), Hainan Yazhou Bay Seed Lab (B21HJ0303) and the Earmarked Fund for China Agriculture Research System (CARS-11-HNGJC, CARS-11-HNCYH).

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

Y.Y., J.G. and Y.C. designed the experiments. Y.W., M.G., R.P., T.Z., X.Z., Y.C., R.L. and J.L. performed the experiments. Y.W., M.G. and R.P. wrote the manuscript. Y.Y., J.G. and Y.C. revised the manuscript.

References

Bart, R., Cohn, M., Kassen, A., McCallum, E.J., Shybut, M., Petriello, A., Krasileva, K. *et al.* (2012) High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved effectors to target for durable resistance. *Proc. Natl Acad. Sci. USA,* **109**, E1972-E1979.

Cohn, M., Bart, R.S., Shybut, M., Dahlbeck, D., Gomez, M., Morbitzer, R., Hou,

B.-H. *et al.* (2014) *Xanthomonas axonopodis* virulence is promoted by a

transcription activator-like effector–mediated induction of a SWEET sugar

transporter in cassava. *Mol. Plant Microbe Interact.* **27**, 1186-1198.

McCallum, E.J., Anjanappa, R.B., and Gruissem, W. (2017) Tackling agriculturally

relevant diseases in the staple crop cassava (*Manihot esculenta*). *Curr. Opin.*

Plant Biol. **38**, 50-58.

Wei, Y., Zeng, H., Liu, W., Cheng, X., Zhu, B., Guo, J. and Shi, H. (2021)

Autophagy-related genes serve as heat shock protein 90 co-chaperones in disease

resistance against cassava bacterial blight. *Plant J.* **107**, 925-937.

Figures

Figure 1. Engineering CBB resistant plants through CRISPR/Cas9-targeted editing of the *MeSWEET10a* promoter in cassava. (A) *Xam11* infects the cultivar cassava SC8 with marked pathogenic symptoms. **(B)** Schematic representation of RVD in TALE20$_{Xam11}$ recognize the EBE$_{Xam11}$ on the *MeSWEET10a* promoter from cultivar cassava SC8. The C-terminus of TALE20$_{Xam11}$ has two nuclear localization signals (NLS) and an acidic activation domain (ADD). **(C)** TALE20$_{Xam11}$ directly activates *MeSWEET10a* expression through the EBE region. *MeSWEET10a* promoter-luciferase fusion and 35S-driven TALE20$_{Xam11}$ constructs were delivered into *Nicotiana benthamiana* cells through infection by *Agrobacterium tumefaciens*. Luciferase activity was quantified after 1 dpi. SK: pGreen II62-SK vector (control vector); TAL20: pGreen II62-SK-TAL20$_{Xam11}$ vector; pSWEET10a: pGreen II0800-pMeSWEET10a-LUC vector; pSWEET10a-EBE: pGreen

II0800-pMeSWEET10a-EBE$_{TAL20}$-LUC vector (deletion of EBE$_{TAL20}$ from the

MeSWEET10a promoter). Data are displayed as mean technical replicates ± standard

deviation. Experiments were repeated three times with similar results (Student's t-test,

***P < 0.005). **(D)** CRISPR/Cas9-induced mutation types and frequency. **(E)**

Hi-TOM sequencing of the EBE$_{TALE20}$ regions of wild type (WT) non-transgenic SC8

cassava compared with the engineered lines #1, #2, #3, and #4. The EBE$_{TALE20}$ region

of the TALE20$_{Xam11}$ target is marked in a red box in WT; the sgRNA sequences are

underlined; the protospacer adjacent motif (PAM) sites are marked in blue text; the

hyphens are missing bases, and the red letter is the inserted base. **(F)** Time course of

MeSWEET10a expression in mutant lines and WT after *Xam11* inoculation.

MeSWEET10a transcript levels in leaves determined by qPCR at 0, 0.5, 1, 3, and 7 dpi.

Three biological repeats were performed for each data point (Student's t-test, ***P <

0.005). **(G)** Symptom development after *Xam11* inoculation. Scale bar, 5 mm. **(H)**

and **(I)** Time course of leaf disease areas and bacterial numbers after *Xam11*

inoculation. At least 10 leaves were harvested for the diseased areas with the bacterial

number assay performed using three biological replicates (Student's t-test, *P < 0.05,

***P < 0.005). **(J)** Callose depositions in leaves. Scale bar, 50 μm. **(K)** Tuber root

phenotypic characterizations. Scale bar, 1 cm. **(L)** Yield-related traits shown as the

mean ± SD from five plants of each line. *P* values were calculated using Student's

t-test. The values marked with the same letter (a) represented no significant

difference.

151