1 **Parametric effects of light acting via multiple photoreceptors contribute to**

2 **circadian entrainment in *Drosophila melanogaster***

4 Lakshman Abhilash, Orie Thomas Shafer[#]

5 The Advanced Science Research Center, The City University of New York; The Graduate Center
6 at the City University of New York
7

9 *Running title: Parametric entrainment in Drosophila*

10
11
12

13 [#]Correspondence: oshafer@gc.cuny.edu

14
15

Abstract

Circadian rhythms in physiology and behavior have near 24-hour periodicities that must adjust to the exact 24-hour geophysical cycles on earth to ensure adaptive daily timing. Such adjustment is called entrainment. One major mode of entrainment is via the continuous modulation of circadian period by the prolonged presence of light. Although *Drosophila melanogaster* is a prominent insect model of chronobiology, there is little evidence for such continuous effects of light in the species. In this study, we demonstrate that prolonged light exposure at specific times of the day shapes the daily timing of activity in flies. We also establish that continuous blue- and UV-blocked light lengthens the circadian period of *Drosophila* and provide evidence that this is produced by the combined action of multiple photoreceptors which, includes the cell autonomous photoreceptor *cryptochrome*. Finally, we introduce ramped light cycles as an entrainment paradigm that produces light entrainment that lacks the large light-driven startle responses typically displayed by flies and requires multiple days for entrainment to shifted cycles. These features are reminiscent of entrainment in mammalian models systems and make possible new experimental approaches to understanding the mechanisms underlying entrainment in the fly.

Keywords: parametric entrainment, continuous entrainment, tonic entrainment, Drosophila, circadian, locomotor activity, Aschoff's rule, free-running period, constant light, skeleton photoperiods

Introduction

Circadian rhythms are ubiquitous endogenous rhythms in behavior, physiology and metabolism. Such rhythms "free-run" in the absence of cyclic environments at periodicities close to but significantly different from 24-h. To support optimal daily timing on a rotating planet, these rhythms must be coordinated with the exact 24-h rhythms of the earth's geophysical cycles, which include daily changes in light and temperature. Such coordination, called entrainment, enables organisms to restrict their daily activities and other biological processes to appropriate times of the day and to anticipate predictable daily changes in the environment [1–4]. Poor entrainment contributes significantly to many health problems in modern societies, making entrainment a central problem to circadian biology and general well-being [4].

Organisms are thought to entrain to cyclic environments using two distinct processes [1,5]. In the first, referred to as non-parametric, environmental transitions instantaneously reset the phase of the circadian clock, much like watches being reset upon the switch to daylight saving time [1,6]. In the second, referred to as parametric, the continuous action of light serves to speed-up or slow-down the angular velocity of internal clocks in a time-of-day dependent manner [1,5,7]. Real world entrainment likely involves both parametric and non-parametric processes [5] and understanding both processes is critical if we are to understand the entrainment of the circadian system and how clocks operate under both natural and artificial light environments.

Both parametric and non-parametric effects of light have been incorporated into unified models of circadian entrainment, including for *Drosophila,* an organism that offers a powerful model system for understanding the neuronal and molecular basis of light entrainment [8–10]. However, the

fly's circadian system does not maintain robust endogenous circadian rhythms under constant light

(LL), making the mechanisms of parametric entrainment difficult to address. This is because of

light and CRYPTOCHROME (CRY; blue light photoreceptor) mediated degradation of

TIMELESS (TIM), a core component of the molecular circadian clock ([11], reviewed in [12]).

Indeed, very few studies have examined the parametric effects of light on the fly clock. Free-

running rhythms have been observed for *Drosophila*, both in *cryptochrome* loss of function

mutations, which maintain rhythms in LL [13–16] and in small numbers of wild-type flies [17,18].

These studies suggest that LL conditions lengthen the clock's endogenous period relative to

constant darkness (DD). However, the sensitivity of the fly clock to light means that LL conditions

most often cause arrhythmicity or internal desynchronization in wild-type strains [15] and this has

limited the utility of the fly as a model of parametric light entrainment. For this reason, we have

sought to develop approaches that allow us to use the powerful fly model to understand the

parametric effects of light.

We therefore set out to identify light regimes that better support an examination of parametric light

effects and their contribution to circadian entrainment in *Drosophila*. Using various photoperiodic

regimes (see Methods) we show that the timing of locomotor activity under standard laboratory

conditions is strongly shaped by long durations of light at specific times of the day. This suggested

that speed of the clock regulating these rhythms must be modulated by time-of-day dependent

continuous effects of light. In our search for LL conditions that support free-running rhythms in

the fly, we find that LL regimes in which UV and blue wavelengths have been blocked support

robust free-running rhythms and produce lengthened circadian periods (slower running clocks).

We also show that external photoreceptors and the blue light circadian photoreceptor

cryptochrome differentially govern the parametric effects of light under these conditions. Finally,

we define light regimes that entrain behavioral rhythms in a manner that fails to induce the

confounding startle responses associated with widely used LD12:12 cycles (12-h of light and 12-

h of darkness) and that, as in mammalian systems, such regimes require multiple cycles for

entrainment to shifted light cycles. These results reveal important features of entrainment in

Drosophila and establish a useful new approach to understanding light resetting in this important

model organism.

Methods

Fly stocks

The following fly lines were used in this study: *Canton-S* (*CS*; Bloomington Drosophila Stock Center (BDSC) stock number: 64349), white-eye (*w[1118]*) [19], yellow-white (*yw*; BDSC stock number: 1495), *yw;;cry[OUT]* [20], *w;;glass[60j]* [13], *+;;norpA[7]* (BDSC stock number: 5685), *+;;per[01]* (BDSC stock number: 80928) and *+;;clk[JRK]* (BDSC stock number: 80927). All these genotypes were maintained in bottles with corn syrup soy media (Archon Scientific; Durham, NC). Three to five-day old male flies were collected from bottles for loading into *Drosophila* Activity Monitors (DAM; Trikinetics, Waltham, MA). Sample sizes and replication for each experiment and the environmental conditions experienced before behavioral experiments began are described separately for each experiment below.

Entrainment of locomotor activity rhythms under skeleton photoperiods

Fly lines, recording conditions and light conditions:

CS, *w[1118]* and *yw* males were used in these experiments. Flies were maintained under white light LD12:12 before being loaded into locomotor tubes for one cycle after being loaded into the DAM systems before being transferred to the different skeleton photoperiod conditions. ~32 flies of each genotype were collected under CO_2 anesthesia and gently placed into 5-mm glass capillary tubes containing sucrose-agar media, which was sealed with paraffin wax at the media containing end and plugged with a short length of white yarn. Fly-containing tubes were then loaded into DAM monitors and secured with rubber bands woven between tubes. Infrared beam crossings were recorded every minute for the duration of the experiment. Our symmetric skeleton photoperiod (SPP; Fig. 1A), consisted of 30-min pulses starting and ending at dawn (Zeitgeber Time (ZT) 00)

and dusk (ZT 12) of the LD cycle used during rearing, respectively. Additionally, we used two

asymmetric skeleton photoperiods (aSPP), the first of which consisted of a dawn light pulse lasting

6-h with a 30-min pulse ending at dusk (aSPP-1; Fig. 1A), and the second of which consisted of a

dusk light pulse lasting 6-h and a 30-min pulse starting at dawn (aSSP-2; Fig. 1A). All these

conditions employed the standard broad spectrum white light produced by the incubators without

the use of any wavelength filters. These incubators were programmed to run at a constant 25 °C.

The light intensity for all these experiments was 400-500 lux.

Statistical analyses:

The first step in the analysis of these data was to identify flies that were entrained versus flies that

were free-running. Owing to the strong startle responses associated with step-function cycles,

especially under the skeleton photoperiod conditions, it was not possible to use raw data to perform

periodogram analyses and estimate period values. This is because startles to transitions (see Fig.

1A) cause a spike in activity at recurring 24-h intervals which produce strong 24-h signals in

standard periodogram analyses, even when the non-startled component of activity appear to be

free-running. Therefore, we used the Savitzky-Golay (SG) filter with a filter order of 2 and filter

frame length of 251-min to smooth the raw data, to minimize the contributions of startle to the

periodogram results. We then used the χ^2 periodogram on the filtered data and estimated period

values. Flies that had period values between 23.67-h and 24.33-h were considered entrained, and

the ones outside this range were considered free-running. These values were chosen because

they're the two closest points to 24-h (on either side) in our χ^2 periodogram analysis; this enabled

us to include flies that entrain but have periodicities that may be slightly wavering around 24-h.

Profiles of activity were averaged across only the flies that were categorized as entrained. Phases

of evening peaks of activity were estimated by applying the SG filter on raw data of entrained flies

only. For this purpose, the filter order was kept at two, but the filter frame length was reduced to

81-min so that small nuances of peaks could be identified. For each genotype, the effect of light

environment on phases of evening peaks of activity were assessed by Kruskal-Wallis tests

followed by Bonferroni corrections to account for multiple comparisons from the *agricolae*

package for R [21]. All the analyses were carried out using custom R scripts.

Assessment of free-running period under constant light

Fly lines, recording conditions and the constant light paradigm:

CS, w^{1118}, *yw*, cry^{OUT}, $glass^{[60j]}$ and $norpA^{[7]}$ males were used in these experiments. ~32 flies of

each genotype were collected and loaded into locomotor tubes as described above. Infrared beam

crossings were recorded every minute for at least 10 days under constant darkness (DD) or constant

light (LL) in Percival incubators (Model: LED-30HL1, Geneva Scientific, Fontana, WI) at 25 °C.

Data from at least two independent runs are reported. See Suppl. Table 1 for sample sizes and

number of replicate runs.

Given that wildtype *Drosophila* are arrhythmic under constant white light, we sought to establish

light conditions under which wild-type flies maintain robust free-running rhythms. It is well

established that the arrhythmicity of flies under white LL is driven by the Cryptochrome (CRY)

mediated TIMELESS (TIM) degradation [22,23]. Furthermore, CRY is maximally sensitive to

blue and UV wavelengths of light [24]. We therefore used multiple light filters which had varying

degrees of overlap with the CRY action spectrum (Fig. 2A). We used constant light of green

(Filter number: 139; 8-12 umol/m2/sec), orange (Filter number: 105), true yellow (Filter number:

101; 8-12 umol/m2/sec) and blue and UV-blocked (Filter number: 821) wavelengths by means of

using specific filters (Lee Filters, UK; http://www.leefilters.com/). We also used constant red light

using the in-built red LED in our Percival incubators (Fig. 2A). In all cases, we used light

intensities between 6-12 $\mu mol/m^2/sec$. For examining the dose dependent effect of blue and UV-

blocked light we used three intensities, i.e., 14-18 (~16) $\mu W/cm^2$, 28-32 (~30) $\mu W/cm^2$ and 47-53

(~50) $\mu W/cm^2$ at 600nm. We used the S120C sensor and the PM100D console from Thorlabs

(Newton, NJ; https://www.thorlabs.com/) to measure the single wavelength light intensities.

Comparisons of period for various genotypes reported in Figs. 3B and 3C (and Suppl. Fig. 1) under

blue and UV-blocked light were done after assaying flies under intensities of 30-45 $\mu W/cm^2$ at

600nm.

Statistical analyses:

Free-running period values and corresponding rhythmic power were estimated using the χ^2

periodogram (implemented using in-house custom R scripts). Period and power values were

pooled from all independent replicate runs and were compared using Wilcoxon's tests for two

independent samples (also implemented using custom R scripts). Dose response analyses were

carried out using functions from the *drda* package [25] on R using logistic models. The logistic

model is a form of non-linear (sigmoidal) growth curve that was fit with our observed data using

the default four-parameter function. An ANOVA on the fitted model then tests the model against

a null hypothesis of a flat line representing the absence of light intensity dependent changes in the

free-running period.

Entrainment to ramped light cycles

Fly lines, recording conditions and light conditions:

CS, w[1118], yw, cry[OUT], glass[60j], norpA[7], per[01] and *clk[JRK]* males were used in these experiments.

Flies were reared under ramped yellow light cycles in Percival incubators (Model: LED-30HL1,

Geneva Scientific, Fontana, WI) at a constant temperature of 25 °C. ~32 males of each genotype

were collected for each run and loaded into locomotor tubes as described above. Ramping light

cycles were generated such that there was no darkness at any point in the cycle. We programmed

gradual light increases from ~10 $\mu W/cm^2$ to 40-50 $\mu W/cm^2$ (measured at 600nm) (ZT00-ZT12)

followed by gradual light decreases back to ~10 $\mu W/cm^2$ during the next 12-h to complete the

cycle. We settled on these light cycles based on our desire to avoid distinct jumps in light intensity

from complete darkness, which produce large startle responses, and to reduce the possibility of

any stark transition in light intensity that that could induce non-parametric clock resetting. We

recorded beam crossings, as described above, under the ramped light cycles for nine consecutive

days, at the end of which flies were kept under constant light starting from the nadir of the ramped

cycle (~10 $\mu W/cm^2$ intensity) to assess phase control. If entrained, flies must have a similar phase

of locomotor activity on the first day under constant conditions as they did during entrainment.

This phenomenon, referred to as phase-control, is a pre-requisite for a rhythm to be considered

entrained [2].

Statistical analyses:

Owing to the novelty of this light regime, it was necessary to assess if flies could entrain to sit at

all. For each fly, we plotted an actogram over nine ramp cycles, computed a wavelet power

spectrum to get daily estimates of period and plotted the standard χ^2 periodogram. The wavelet

power spectrum was computed using the *WaveletComp* package on R [26] and is appropriate for

use with circadian time series [27,28]. Individual flies were considered "entrained" only if visual

inspection of the actogram showed that timing of activity had stabilized by the sixth ramping cycle,

if the wavelet power spectrum showed stable period values between 23.5-24.5-h post the sixth

cycle, and/or if the χ^2 periodogram showed peaks in the same range. A fly was categorized as

"transiently entrained" or "relatively coordinated" if there were obvious period changes during the

time-course (for instance, a period that shifted from period of < 24-h to > 24-h) in the actograms,

if there was no period stability established by the sixth cycle based on the wavelet power spectrum,

if there were multiple peaks in the χ^2 periodogram, or if flies appeared entrained for the first few

cycles and then appeared to free-run. Flies were deemed "free-running" if the actograms showed

activity bouts drifting in a single direction and if the χ^2 periodogram showed peaks at period values

less than 23.5 or greater than 24.5-h. Flies were considered "arrhythmic" if there was no

significant peak in the χ^2 periodogram or if no detectable period in the wavelet power spectrum.

For loss-of-function clock mutants, flies were considered rhythmic if there was a significant peak

in the χ^2 periodogram or if there was a detectable period in the wavelet power spectrum. All these

analyses were carried out using custom R scripts.

Only the flies categorized as entrained (or rhythmic, in case of loss-of-function clock mutants)

were used to construct the average profiles of activity and to assess phase control under constant

dim light. We estimated phases of activity peaks using the SG filter for the entrained flies under

cyclic conditions and on the first day of constant dim light. We used a filter order of one and filter

frame length of 41 on activity data binned in 15-min intervals to smooth the data before identifying

peaks. Phase control was assessed by *V*-tests implemented through the *CircStats* package for R

[29]. *V*-tests are Rayleigh's tests of uniformity on circular data that have an alternate hypothesis

of a known mean angle [30]. Difference in phases of peaks between *CS* and *per^{01}* flies was tested

using a Wilcoxon's test, also implemented on R.

Results

Pulses of light at dawn and dusk are insufficient to mirror the entrained waveform of locomotor activity under standard LD12:12 conditions

One major prediction of the non-parametric (a.k.a., instantaneous) model of entrainment is that light presented at dawn and dusk are the most critical signals for entrainment, and that light during the rest of the day is largely superfluous [5,6]. If this were true of *D. melanogaster* locomotor rhythms, we would expect the locomotor activity waveform of flies be highly similar under LD12:12 and symmetric skeleton photoperiods (SPP), in which only two 30-min light pulses are provided to flies in an otherwise dark environment (see Methods; Fig. 1A). Although such experiments have been done on *Drosophila* eclosion rhythms [31,32], ours is the first, to the best of our knowledge, to report results of such experiments on the locomotor activity rhythm in *D. melanogaster*. All wild-type and genetic background control strains we examined readily entrained to SPP displaying no differences from LD12:12 flies in the percentage of entrained flies (Suppl. Table 2).

Averaged locomotor activity profiles of *CS*, w^{1118} and *yw* flies indicate that there were distinct differences in waveforms between LD12:12 and SPP (Fig. 1A). We quantified the phases of evening activity peaks from smoothed data (see Methods) and found that median phases in all three genotypes were significantly advanced under SPP compared to LD12:12 (See Methods for details of statistical analysis; Fig. 1B). These results clearly show that skeleton photoperiods do not recapitulate entrainment under LD12:12 and that light in the middle of the day has significant effects on the phase of evening activity.

Long durations of light only during the second half of the day shape the timing of locomotor

activity under standard LD cycles

If parametric effects contribute to entrainment under step function LD cycles, it is expected that

increased durations of skeleton light exposure will have measurable effects on the phase of

entrainment in a manner that depends on the timing of the increased light [7,32]. The parametric

model of entrainment posits that long durations of light delivered at dawn will accelerate the clock,

whereas light delivered at dusk will have a decelerating effect on the clock [7,33]. The prediction,

therefore, is that long durations of light around dawn would lead to reduced free-running period,

thereby leading to an advanced phase of entrainment and that long durations of light during dusk

would lengthen free-running period and consequently delay the phase of entrainment [34]. In

contrast, if non-parametric effects dominate entrainment at transitions, activity profiles under

skeleton light cycles should be largely unaffected by the length of the skeleton light periods. In

order to test the extent to which the fly is entrained to LD cycles via parametric or non-parametric

effects, we subjected our flies to two asymmetric skeleton photoperiods (aSPP), one with 6-h of

light during the first half of the day and a 30-min pulse at dusk (30-min; aSPP-1; Fig. 1A), and one

with a 30-min dawn pulse and 6-h of light during the second half of the day (aSPP-2; Fig. 1A). As

for SPP, we first sought to determine the extent to which wild-type and genetic background strains

entrain to aSPP regimes. We found that there was no significant reduction in the number of flies

entraining to either of the asymmetric skeleton photoperiods for any genotypes (Suppl. Table 2).

We next compared the phases of the evening peak of activity under our two aSPP conditions to

those seen under LD12:12 and found that, for wildtype *CS* flies, there was a significant advance

in the timing of evening activity when the long-duration of light was delivered around dawn (aSSP-

1; Figs. 1A and 1B). Though the evening peak appeared to be delayed when long duration light

was delivered around dusk, the differences in the median phases of individual flies did not reach

significance (aSPP-2; Figs. 1A and 1B). Similar effects of long-duration light around dawn were

observed for w^{1118} and yw flies (aSPP-1; Fig. 1) and the evening peak was significantly delayed in

these strains when long duration light was delivered around dusk (aSSP-2; Fig. 1). The difference

in phases of evening peak of yw flies under LD12:12 and aSPP-2, although significant was only

~15-min and is therefore barely apparent in the profiles (Fig. 1).

Taken together, these results reveal that long durations of light at the end of the day are sufficient

for flies to achieve a waveform similar to that displayed under LD12:12. Therefore, entrained

waveforms of locomotor activity under standard laboratory LD conditions are strongly shaped by

time-of-day dependent continuous effects of light.

Wildtype and background strains have longer free-running periods under constant blue- and

UV-blocked light

While the number of discrete transitions between light and dark are the same across the different

skeleton photoperiod regimes, the timings of these transitions are different (see Fig. 1A). This

may therefore appear as a confound in our experimental design. However, instantaneous effects

of light on the clock around mid-day (which is where our transitions occur) are almost non-existent

[2,18,35,36]. Thus, our results described above are consistent with the speed of internal clocks

being modulated by long durations, and continuous effects, of light. The most direct method

available for measuring such light effects is the demonstration that free-running periods of a

circadian rhythm differ between constant light and constant dark conditions. Given the fact that

the presence of the blue light circadian photoreceptor CRY renders the *Drosophila* clock

arrhythmic under white LL, we assayed locomotor activity in flies under various light conditions

with differing degrees of overlap with the CRY action spectrum (Fig. 2A). We assayed *CS* flies

under constant green, orange, yellow, blue- and UV-blocked and red lights and found that flies

were arrhythmic under green and yellow lights (see example actograms in Fig. 2B). While many

flies were arrhythmic under orange light, some flies displayed complex rhythms (Fig. 2B) in which

spontaneous changes in free-running period are apparent. Under constant red light, flies were

rhythmic and had periodicities similar to those measured under constant darkness (a little less than

24-h; Fig. 2B). In contrast, we found that when the blue- and UV-components of white light are

blocked, flies are robustly rhythmic and are characterized by periodicities longer than those

measured under constant darkness (Fig. 2B; Suppl. Fig. 1A; Suppl. Table 1). Thus, the presence

and speed of endogenous rhythms appear to depend on the extent of overlap between the spectrum

of constant light used and the action spectrum of CRY. There is considerably larger overlap in

case of green and yellow light (Fig. 2A), conditions under which arrhythmicity was observed. In

case of orange light, there is relatively little overlap, and this was associated with of complex

rhythms in a subset of flies. The complete lack of overlap in case of red light does not seem to

affect the presence or speed of the clock. However, the blue- and UV-blocked light, which has

minimal but non-zero overlap with the CRY absorption spectrum, supports robust but slower

rhythms. Taken together, these results suggest that CRY may play a role in the regulation of clock

speed during prolonged illumination when light conditions do not drive the CRY mediated loss of

circadian timekeeping.

Before examining the role of CRY on rhythm speed, we performed dose response analyses to

examine effects of light intensity on the free-running period of the clock driving rhythms in

locomotor activity. We used *CS*, w^{1118} and *yw* under three intensities of constant blue- and UV-

blocked light. Fitting the logistic model (see Methods) indicated that there was significant light

intensity dependent change in speed of the locomotor activity rhythm in all three fly strains (Fig.

3A). However, the extent of change in rhythm speed depended on the strain used (Figs. 3A-C).

While *CS* flies displayed a period lengthening of ~1-h, the w^{1118} and *yw* strains displayed

lengthening of ~3 to ~3.5-h (Figs. 3A-C). We also examined the strength of these rhythms under

constant light and constant darkness (see Methods) and found that w^{1118} and *yw* flies had

significantly lower rhythmic strengths under LL conditions (Suppl. Fig. 1A; Suppl. Table 1). w^{1118}

and *yw* strains lack normal screening pigment in the eyes due to a loss of function mutation in the

white gene. Our results therefore suggest that external photoreceptors may play a role in light

dependent changes in rhythm strength and speed, owing to the *white* gene's role in photoreception

e.g., [37].

Continuous effects of light are dependent on both external photoreceptors and *cryptochrome*

The *Drosophila* circadian system has two major photoreceptive systems, one mediated by the deep

brain, cell autonomous blue-light photoreceptor *cryptochrome (cry),* and the other consisting of

external photoreceptors, i.e, the eyes, ocelli and the HB eyelets [38]. Previous work by others has

established that external photoreceptors and CRY cooperate to entrain the fly's circadian clock to

12:12LD cycles, e.g., [39]. Our results also suggest that both CRY and external photoreceptors

likely contribute to light dependent changes in endogenous period. We therefore expanded our

analyses to fly strains that have mutations in these light input pathways. The loss-of-function

cry^{OUT} mutation lacks functional *cry* [20] and the *glass^{[60j]}* mutant lacks all external photoreceptors

due to the loss of a transcription factor required for their development [13]. Previous work has

shown that *cry/glass* double mutants are circadian blind to environmental light, suggesting that,

taken together, *cry* and external photoreceptors account for most if not all light input to the

circadian clock [13]. Further, studies have also shown that *cry* may have the ability to integrate

dim light information over long durations and that *cry* mutants continue to display weak resetting

responses to light [35,40].

Therefore, to further examine the extent to which external photoreceptors and CRY contribute to

the parametric effects of light, we examined the effects of constant light on mutants lacking normal

photoreception/transduction. We found that loss-of-function *cry^{OUT}* mutants do not increase their

free-running period under continuous light; rather, they showed a significant *reduction* in free-

running period compared to their values under constant darkness, although this difference was

quite small. This indicates that CRY is sensitive to long wavelengths light under prolonged

illumination, and that it contributes to parametric light effects under a broader range of

wavelengths than expected from its action spectra [24] (Figs. 3B and C; Suppl. Table 1).

Remarkably, loss-of-function *cryptochrome* mutants show twice as much advance in the phase of

their evening peaks of activity under SPP compared to their background controls (compare Fig. 1

and Suppl. Fig. 2). This is also consistent with the idea that *cryptochrome* tends to lengthen period

under long durations of light.

We also found that a large proportion of *glass^{[60j]}* mutants were arrhythmic under constant light,

but that the free-running period of rhythmic flies was longer under constant light by ~1.75-h (Figs.

3B and 3C; Suppl. Table 1).   In addition, we found that all *norpA* mutants, which lack the

phopholipase-C signaling that mediates rhodopsin-based phototransduction, were rhythmic and

their free-running periods were longer under constant light by ~3.5-h (Figs. 3A and 3C; Suppl.

Table 1). Thus, in contrast to cry^{OUT} mutants, which showed a very small reduction in period

under constant light, flies lacking external photoreceptors were characterized by large parametric

increases in period.

When we compared the rhythmic strength of these mutants under constant light and darkness, we

found that while cry^{OUT} displayed comparable rhythmic strength under the two regimes (Suppl.

Fig. 1B; Suppl. Table 1), both *glass[60j]* and *norpA* both displayed significantly weaker rhythms

under constant light (Suppl. Fig. 1B; Table 1). Thus, the loss of *cry* function drastically decreases

the parametric effects of light, while the loss of external photoreceptor function increases them

and weakens endogenous rhythms under constant light conditions.

These results suggest that parametric effects of light modulate clock speed in *Drosophila* and

suggest that these are likely to contribute to light entrainment. Surprisingly, *cryptochrome* appears

to be required for such lengthening of free-running periods, even under light conditions lacking

blue and UV light. Furthermore, the increased extent of period lengthening by constant light in

norpA mutants compared to wildtype *CS* flies suggests that phospholipase-C dependent

phototransduction normally acts in opposition to *cry* with regard to the parametric effects of light,

a result that supports previous work supporting antagonistic roles for *cry* and external

photoreceptors in the adjustment of the evening peak of activity under step function LD cycles

[39] and suggesting that parametric light effects make significant contributions to the timing of the

entrained evening peak of activity.

Ramped blue- and UV-blocked light cycles have significant experimental utility for

understanding entrainment

Although our skeleton photoperiod experiments revealed that entrained waveforms of locomotor

activity under LD12:12 are significantly shaped by parametric effects of light, the large startle

responses to the discrete transitions between darkness and light which can be seen clearly in the

activity profiles (Fig. 1), may contribute to the daily timing of activity in ways that may obscure

the effects of light on the clock. For example, arousal signals caused by the masking effects of

light likely contribute significantly to the timing of major peaks of activity. Thus, such startle

responses pose a challenge to understanding the true contribution of the entrained circadian system

to these waveforms. Moreover, waveforms of loss-of-function clock and light input mutants look

remarkably normal under step function LD cycles in that they reliably produce two daily peaks of

activity at dawn and dusk [13,41]. To characterize the timing of entrained circadian outputs most

accurately there is therefore a need for a light entrainment paradigm to which wildtype flies readily

entrain but that do not induce strong diurnal rhythms in loss-of-function clock mutants.

Furthermore, light regimes that better reveal entrainment deficits in flies lacking normal light input

pathways would be of great utility to the field. Experiments conducted under such light regimes,

although structurally very different than widely used step-function LD cycles, could reveal

neuronal and molecular underpinnings of circadian entrainment that step-function LD cycles will

not.

To this end, we asked if ramping light cycles in which light intensity gradually changes from a low

but non-zero value to a high value and then returns to the low value over 24-h, might provide novel

experimental utility. We argue that such a regime would be useful because (i) there are no discrete

jumps between states of darkness and light and therefore no cause for large startle responses, (ii)

ramping of the intensity represents a much more challenging light condition for entrainment than

step function LD cycles, as it requires the circadian system to constantly monitor a slowly changing

environment, and (iii) there is no opportunity under such a light cycle for the discrete phase-shifts

of the clock. If so, ramping light cycles would offer a unique regime of entrainment biased toward

the parametric effects of light, making it a useful complement to the widely employed LD12:12

cycle. Light ramps might, therefore, make possible new approaches to understanding the

mechanisms underlying this important form of circadian entrainment.

To examine the potential utility of ramping light cycles, we placed our wild-type, genetic

background, and several mutant strains under ramping light cycles that were filtered for UV and

blue wavelengths (see Methods) and characterized individual flies as either (i) entrained, (ii)

displaying relative coordination (transient or unstable entrainment), (iii) free-running, or (iv)

arrhythmic (Fig. 4). We found that while close to 80% of wildtype *CS* flies entrained to yellow

ramps, only ~50% of w^{1118} and *yw* flies entrain. Interestingly, ~80% of cry^{OUT} and ~97% of *norpA*

mutants displayed free-running rhythms under these conditions (Fig. 4; Suppl. Table 3) indicating

the presence of a strong circadian oscillation that had failed to entrain to the light cycle. In the

case of $glass^{[60j]}$ mutants, ~50% of the flies entrained and ~40% of flies are arrhythmic under these

ramping light cycles (Fig. 4; Suppl. Table 3).

We examined the locomotor activity profiles of entrained wildtype *CS*, *w^1118* and *yw* flies under

these conditions, and found one major peak of activity that was approximately centered around the

second half of the declining phase of light intensity (Fig. 5A). There is also a smaller peak of

activity especially visible in *w^1118* flies, which commences at the beginning of the inclining phase

of light intensity (Fig. 5A). Interestingly, while the median phase of the dominant peak of activity

for *CS* and *w^1118* flies was ~ZT19 (with ZT00 here defined as the start of the inclining light phase),

the median phase of the dominant peak in *yw* flies was ~ZT17 (Fig. 7B), a difference of ~2-h in

the phase of entrained activity peaks that is not apparent under standard LD12:12 entrainment

conditions (e.g., Fig. 3C and D). This surprisingly large difference between strains under light

ramps but not under LD12:12 lends strong support to our notion that such light ramp cycles will

allow the experimentalist to detect the effects of genetic and neuronal manipulations on

entrainment that would not be apparent using standard LD12:12 cycles, thanks, most likely, to the

absence of startle/arousal mediated effects on the timing of behavior and to the biasing of

entrainment toward parametric effects of light.

A critical property of entrainment is phase control, which means that once allowed to free-run

following entrainment to a cyclic environment, organisms must show similar phases of rhythm on

the first day of constant conditions as it did during stable entrainment [2]. In the absence of phase

control, one cannot conclude that the organism was really entrained during their exposure to any

particular environmental cycle (add same references as above in methods). We estimated phase

control in the wild-type and background control strains and found that phases of the dominant peak

of *CS* flies on the first day under constant conditions is significantly clustered around the median

phase of the same flies under entrainment (Fig. 5B). We found similar significant clustering in

case of the w^{1118} (Fig. 5B) and *yw* strains (Fig. 5B), thereby indicating that these flies had truly

entrained to the previous light ramps rather than being driven directly by changes in environmental

light.

To further establish the extent to which our ramped light cycles provide experimental conditions

that reduce the masking effects of light on behavior, we examined loss-of-function clock mutants

under these conditions. We found that only of ~35% of per^{01} loss-of-function mutants displayed

some form of rhythmicity, with most flies displaying arrhythmicity (~65%; Fig. 5C; Suppl. Table

4). We note that the residual rhythmicity was much weaker than what is observed under LD12:12,

a condition in which the majority of per^{01} mutant display rhythms. In the per^{01} flies that were

rhythmic, there was a single peak of activity very close to ZT00 or the time of the day when light

intensity was at a minimum (Fig. 5D). We quantified their phases and found that the median phase

of activity was ~ZT22.29. We think that such residual rhythmicity in per^{01} flies is some form of

bright light avoidance behavior/masking. We found that this phase of peak is significantly delayed

compared to the phase of their background control *CS* flies (Wilcoxon's test: $W = 36$, $p = 9.3e$-

09), which is not typically the case under standard LD12:12 conditions, wherein evening peaks

coincide with dusk. For clk^{JRK} mutants, which bear a loss-of-function mutation of the *clock* gene,

all flies were arrhythmic under light ramps (Fig. 5C), which is in stark contrast to the relatively

nocturnal activity rhythms displayed by this mutant under LD12:12 [42]. These results suggest

that light ramp cycles will offer significantly increased experimental utility for the examination of

light entrainment in *Drosophila*.

Discussion

Real world entrainment is mediated by a combined contribution of parametric and non-parametric

effects of light. Parametric effects of light have been modeled in the past and have successfully

explained entrainment under various photoperiodic conditions in *Neurospora*, *Drosophila* and

mammals [8–10], unlike the non-parametric model which has limited ability to predict entrainment

under different photoperiods [5,33]. In fact, the non-parametric model alone also has limited utility

in predicting entrainment in humans (see [8] and references therein, and [43]). Therefore, a holistic

study of entrainment requires the understanding of both parametric and non-parametric processes,

and our study is a significant first step in that direction for *Drosophila*.

Antagonistic effects of CRY and norpA in parametric light entrainment

Our results show that wildtype flies have longer free-running periods under constant light. The

amount of this lengthening appears to be strongly dependent on both *cry* and *norpA* mediated

phototransduction. Not only does the loss of CRY prevent constant light from lengthening free-

running period, but it also appears to modestly reduce it. On the other hand, the loss of *norpA*

function results in larger LL-induced increases in period compared to wildtype controls (see Fig.

3 and Suppl. Table 1). These clearly indicate an antagonistic interaction between *cryptochrome*

and *norpA* under constant light. The idea of such an antagonism is also supported by studies in

flies showing opposite effects of *cry* and visual photoreception on the evening peak of locomotor

activity and their effects on clock oscillations under long days [39].

Such an antagonistic relationship can be hypothesized to be of adaptive value in the context of

parametric entrainment, wherein organisms must modulate the speed of their clocks in a time-of-

day dependent manner. Such antagonism can explain the entrainment of locomotor rhythms to

skeleton photoperiods. *CS* flies displayed a significantly advanced phase of evening peak activity

under the symmetrical skeleton photoperiod (SPP) relative to LD12:12, consistent with the idea

that the absence of a full 12-h photoperiod leads to a faster running clock, thereby advancing

evening peak phase under SPP. Similar results in the w^{1118} and *yw* background strains lend further

support to this idea.

The significant evening peak advances displayed by wild-type and background strains under SPP

indicates that the two brief light pulses do not comprise the minimum required light schedule to

recapitulate LD12:12 behavior. We therefore extended our analyses to asymmetric SPPs to ask if

longer durations of light are required at specific times of day for activity to be timed in manner

similar to that observed under standard LD12:12. In this way we tested the prediction made by

the parametric model of entrainment that long durations of light contiguous with dawn would have

a net accelerating effect on the clock while long durations of light contiguous with dusk would

have a net decelerating effect on the clock. Indeed, this is exactly what we found: wild-type and

genetic background flies had significantly advanced phase under long dawn skeletons and a

tendency towards increased similarity to LD12:12 under long dusk skeletons. To the best of our

knowledge, this is the first study to demonstrate that long durations of light at specific times of the

day are required for entrainment phases like those observed under LD12:12 in the locomotor

activity rhythm of flies.   Similar experiments have been performed to examine the light

requirement schedules of the eclosion rhythm for *D. melanogaster* flies that were artificially

selected for morning and evening adult emergence [32]. This study found that for genotypes that

had shorter than 24-h periods under constant darkness and a larger delay/advance ratio of their

phase response curves, regimes that had long durations of light contiguous with dusk comprised

the minimum light requirement schedule to mirror behavior under LD12:12 [32], providing further

support for time-of-day specific parametric light effects.

Cryptochrome mediates the parametric effects of light in D. melanogaster

The most direct effect of light on the molecular circadian clock in *D. melanogaster* is through the

deep brain blue light photoreceptor CRY, which has long been thought to mediate the non-

parametric effects of light the circadian clock by driving the rapid degradation of TIM upon

exposure to light, thereby causing phase dependent advances or delays in the molecular circadian

oscillator (e.g., [11]). However, the construction of light phase response curves (plots of time-of-

day dependent phase-shifts in response to brief pulses of light) for the cry^{baby} mutant, revealed that

parametric light effects were still present, though reduced, in the absence of CRY function [35].

The presence of CRY is also thought to explain why dipteran clocks cannot maintain their rhythms

under constant light conditions: CRY action under continuous presence of light prevents sufficient

build-up of TIM for the molecular clock to run [44–46]. Given that CRY has long been thought

to be sensitive specifically to UV/blue wavelengths, we assumed that our blue- and UV-blocked

regime would "work-around" CRY. Indeed, the strong endogenous timekeeping displayed by

wild-type flies under this regime was reminiscent of the rhythms displayed by loss of function *cry*

mutants under the continuous white light conditions that render normal flies arrhythmic [44]. This

suggested that CRY was not engaged under blue- and UV-blocked light. However, the lack of

parametric effects in cry^{OUT} flies (Fig. 3) revealed, surprisingly, that non-UV/blue wavelengths

likely do engage CRY and contribute to the parametric effects of light.

Owing to the clear absence of period lengthening under blue- and UV-blocked light in loss-of-

function *cryptochrome* mutants, we hypothesize that CRY activation and CRY mediated TIM

degradation may be at the heart of period lengthening in this regime. This is because the action

spectrum of CRY suggests that it is likely to be weakly sensitive to wavelengths of light longer

than 500nm (Fig. 2A) [24]. Furthermore, flies display clear behavioral preferences for specific

wavelengths of light throughout the circadian cycle, avoiding blue light and seeking out green and

red shifted wavelengths during the day [47]. We suspect that, in real world settings, flies are likely

to be exposed, for significant durations, to light that is significantly more yellow/red shifted than

the white light used in the laboratory. Therefore, under natural and freely behaving conditions,

CRY mediated TIM degradation is expected to be significantly lower than that seen in the lab.

This would allow CRY to mediate the parametric effect of light by controlling TIM turnover in a

manner that depends on light intensity and duration, thereby adjusting PERIOD accumulation and

nuclear entry to modulate the clock's speed. This, we argue, is mechanistically similar to the

effects of a hypomorphic *cry* mutation under bright white LL, which produces lengthened free-

running periods [48]. Future work examining molecular clock cycling under a range of light

conditions will be necessary to test this hypothesis.

On the utility of light ramps to study entrainment

The vastly simpler nervous system and highly conserved molecular clock in *Drosophila* has made

it a superlative model system to understand mechanisms of entrainment that are relevant to

entrainment in animals generally. However, one aspect of light entrainment in flies has limited its

relevance to light resetting in mammals: whereas mammals including humans require several

cycles to entrain to a shifted (i.e., jet lagged) LD cycle (reviewed in [2]), flies adjust to even large

shifts in 12:12 LD cycles almost instantaneously upon encountering a shifted light transition

(reviewed in [38]), and even flies lacking major components of light input pathway re-entrain

fairly rapidly (reviewed in [38]). For these reasons, new experimental light regimes are necessary

if *Drosophila* is to be used to address entrainment in a manner with the greatest possible relevance

to animals generally.

We wonder if this apparent instantaneous re-entrainment to shifted LD cycles is the result of the

large startle responses produced by the light/dark cycles under which flies are typically studied.

Standard step-function LD cycles (and our various skeleton photoperiods – Fig. 1 and Suppl. Fig.

2) promote significant startle responses at light transitions, and these, rather than light's effect on

the clock, may be driving the rapid re-entrainment seen in flies. We propose the blue- and UV-

blocked ramping light cycles as one such regime, as they fail to produce the large startles

associated with step-function LD cycles and require multiple cycles for entrainment (i.e., they

produce transients; Fig. 4). Furthermore, unlike standard LD12:12 conditions, blue and UV-

blocked light ramps reveal significant entrainment defects in flies lacking light input pathways and

fail to drive strong daily rhythms in loss-of-function clock mutants. This regime, therefore, allows

us to explore aspects of entrainment in flies that are highly relevant but difficult to study using

step-function LD cycles.

Conclusion

In summary, we have demonstrated that parametric effects have a strong influence on entrainment

under the long-employed step function LD12:12 and that, as predicted by parametric models of

entrainment, there are clear time-of-day dependent parametric effects of light. We have also

established light regimes that demonstrate significant effects of continuous light on the *Drosophila*

circadian clock and reveal an unexpected role for CRY in the mediation of these effects, at

wavelengths beyond those associated with its established role as a blue light circadian

photoreceptor. Interestingly, the apparent antagonism of CRY and phototransduction from

external photoreceptors explains the patterns of entrainment displayed under various skeleton

photoperiod regimes. Finally, our work establishes experimental light conditions that will offer

great utility for future research aimed at understanding the parametric effects of light on the clock

and the ways in which specific genes and neural pathways contribute to circadian light

entrainment.

Acknowledgments

This work was supported by a grant from the National Institute of Neurological Disorders and Stroke (R01NS077933) and start-up funds provided by the State of New York. We would like to thank Hannah Pettibone, Matthew Ciolkowski and Lukasz Widziszewsky for technical support and Rob Veline and Budha Chowdhury for comments on the manuscript.

References

[1]   Pittendrigh CS. Circadian Systems: Entrainment. Biological Rhythms, Boston, MA: Springer US; 1981, p. 95–124. https://doi.org/10.1007/978-1-4615-6552-9_7.

[2]   Moore-Ede MC, Sulzman FM, Fuller CA (Charles A. The clocks that time us : physiology of the circadian timing system. Harvard University Press; 1982.

[3]   Vaze KM, Sharma VK. On the Adaptive Significance of Circadian Clocks for Their Owners. Chronobiol Int 2013;30:413–33. https://doi.org/10.3109/07420528.2012.754457.

[4]   Roenneberg T, Merrow M. The circadian clock and human health. Current Biology 2016;26:R432–43. https://doi.org/10.1016/j.cub.2016.04.011.

[5]   Daan S. Colin Pittendrigh, Jürgen Aschoff, and the Natural Entrainment of Circadian Systems. J Biol Rhythms 2000;15:195–207. https://doi.org/10.1177/074873040001500301.

[6]   Pittendrigh CS, Daan S. A functional analysis of circadian pacemakers in nocturnal rodents IV. Entrainment: Pacemaker as clock. Journal of Comparative Physiology ? A 1976;106:291–331. https://doi.org/10.1007/BF01417859.

[7]   Swade RH. Circadian rhythms in fluctuating light cycles: Toward a new model of entrainment. J Theor Biol 1969;24:227–39. https://doi.org/10.1016/S0022-5193(69)80048-2.

[8]   Rémi J, Merrow M, Roenneberg T. A Circadian Surface of Entrainment: Varying T, τ, and Photoperiod in Neurospora crassa. J Biol Rhythms 2010;25:318–28. https://doi.org/10.1177/0748730410379081.

[9]   Beersma DGM, Daan S, Hut RA. Accuracy of Circadian Entrainment under Fluctuating Light Conditions: Contributions of Phase and Period Responses. J Biol Rhythms 1999;14:320–9. https://doi.org/10.1177/074873099129000740.

[10]   Abhilash L, Sharma VK. Mechanisms of photic entrainment of activity/rest rhythms in populations of *Drosophila* selected for divergent timing of eclosion. Chronobiol Int 2020:1–16. https://doi.org/10.1080/07420528.2020.1727917.

[11]   Hunter-Ensor M, Ousley A, Sehgal A. Regulation of the Drosophila protein timeless suggests a mechanism for resetting the circadian clock by light. Cell 1996;84. https://doi.org/10.1016/S0092-8674(00)81046-6.

643   [12]   Hardin PE. Molecular genetic analysis of circadian timekeeping in Drosophila. Adv Genet,

644   vol. 74, Academic Press Inc.; 2011, p. 141–73. https://doi.org/10.1016/B978-0-12-387690-

645   4.00005-2.

646   [13]   Helfrich-Förster C, Winter C, Hofbauer A, Hall JC, Stanewsky R. The circadian clock of

647   fruit flies is blind after elimination of all known photoreceptors. Neuron 2001;30.

648   https://doi.org/10.1016/S0896-6273(01)00277-X.

649   [14]   Yoshii T, Funada Y, Ibuki-Ishibashi T, Matsumoto A, Tanimura T, Tomioka K. Drosophila

650   cryb mutation reveals two circadian clocks that drive locomotor rhythm and have different

651   responsiveness to light. J Insect Physiol 2004;50.

652   https://doi.org/10.1016/j.jinsphys.2004.02.011.

653   [15]   Rieger D, Shafer OT, Tomioka K, Helfrich-Förster C. Functional analysis of circadian

654   pacemaker neurons in Drosophila melanogaster. Journal of Neuroscience 2006;26.

655   https://doi.org/10.1523/JNEUROSCI.1234-05.2006.

656   [16]   Dolezelova E, Dolezel D, Hall JC. Rhythm defects caused by newly engineered null

657   mutations in Drosophila's cryptochrome gene. Genetics 2007;177:329–45.

658   https://doi.org/10.1534/GENETICS.107.076513.

659   [17]   Konopka RJ, Pittendrigh C, Orr D. Reciprocal behaviour associated with altered

660   homeostasis and photosensitivity of Drosophila clock mutants. J Neurogenet 2007;21:243–

661   52. https://doi.org/10.1080/01677060701695391.

662   [18]   Nikhil KL, Vaze KM, Karatgi R, Sharma VK. Circadian clock properties of fruit flies

663   Drosophila melanogaster exhibiting early and late emergence chronotypes. Chronobiol Int

664   2016;33:22–38. https://doi.org/10.3109/07420528.2015.1108981.

665   [19]   Barber AF, Erion R, Holmes TC, Sehgal A. Circadian and feeding cues integrate to drive

666   rhythms of physiology in Drosophila insulin-producing cells. Genes Dev 2016;30.

667   https://doi.org/10.1101/gad.288258.116.

668   [20]   Yoshii T, Todo T, Wülbeck C, Stanewsky R, Helfrich-Förster C. Cryptochrome is present

669   in the compound eyes and a subset of Drosophila's clock neurons. Journal of Comparative

670   Neurology 2008;508. https://doi.org/10.1002/cne.21702.

[21]   de Mendiburu Felipe. agricolae: Statistical Procedures for Agricultural Research 2021.

[22]   Stanewsky R, Kaneko M, Emery P, Beretta B, Wager-Smith K, Kay SA, et al. The cryb
Mutation Identifies Cryptochrome as a Circadian Photoreceptor in Drosophila. Cell
1998;95:681–92. https://doi.org/10.1016/S0092-8674(00)81638-4.

[23]   Emery P, Stanewsky R, Hall JC, Rosbash M. A unique circadian-rhythm photoreceptor.
Nature 2000;404:456–7. https://doi.org/10.1038/35006558.

[24]   VanVickle-Chavez SJ, van Gelder RN. Action spectrum of Drosophila cryptochrome. J Biol
Chem 2007;282:10561–6. https://doi.org/10.1074/JBC.M609314200.

[25]   Malyutina A, Tang J, Pessia A. drda: An R package for dose-response data analysis.
BioRxiv 2021.

[26]   Roesch A, Schmidbauer H. WaveletComp: Computational Wavelet Analysis 2018.

[27]   Leise TL. Analysis of Nonstationary Time Series for Biological Rhythms Research:
Http://DxDoiOrg/101177/0748730417709105   2017;32:187–94.
https://doi.org/10.1177/0748730417709105.

[28]   Leise TL. Wavelet analysis of circadian and ultradian behavioral rhythms. J Circadian
Rhythms 2013;11:5. https://doi.org/10.1186/1740-3391-11-5.

[29]   Lund U, Agostinelli C. CircStats: Circular Statistics, from Topics in Circular Statistics
(2001) 2018.

[30]   Zar JH. Biostatistical analysis. Prentice Hall; 1999.

[31]   Pittendrigh CS, Minis DH. The Entrainment of Circadian Oscillations by Light and Their
Role as Photoperiodic Clocks. Am Nat 1964;98. https://doi.org/10.1086/282327.

[32]   Vaze KM, Nikhil KL, Abhilash L, Sharma VK. Early-and late-emerging Drosophila
melanogaster fruit flies differ in their sensitivity to light during morning and evening.
Chronobiol Int 2012;29:674–82. https://doi.org/10.3109/07420528.2012.680557.

[33]   Roenneberg T, Hut R, Daan S, Merrow M. Entrainment Concepts Revisited. J Biol Rhythms
2010;25:329–39. https://doi.org/10.1177/0748730410379082.

[34]   Roenneberg T, Dragovic Z, Merrow M. Demasking biological oscillators: properties and
principles of entrainment exemplified by the Neurospora circadian clock. Proc Natl Acad
Sci U S A 2005;102:7742–7. https://doi.org/10.1073/pnas.0501884102.

[35]   Kistenpfennig C, Hirsh J, Yoshii T, Helfrich-Förster C. Phase-shifting the fruit fly clock
without   cryptochrome.   J   Biol   Rhythms   2012;27.
https://doi.org/10.1177/0748730411434390.

[36] Dollish HK, Kaladchibachi S, Negelspach DC, Fernandez FX. The Drosophila circadian
phase response curve to light: Conservation across seasonally relevant photoperiods and
anchorage   to   sunset.   Physiol   Behav   2022;245.
https://doi.org/10.1016/j.physbeh.2021.113691.

[37] Lee SJ, Montell C. Suppression of constant-light-induced blindness but not retinal
degeneration by inhibition of the rhodopsin degradation pathway. Current Biology 2004;14.
https://doi.org/10.1016/j.cub.2004.11.054.

[38] Helfrich-Förster C. The Drosophila Clock System. Biological Timekeeping: Clocks,
Rhythms   and   Behaviour,   New   Delhi: Springer   India;   2017,   p.   133–76.
https://doi.org/10.1007/978-81-322-3688-7_6.

[39] Kistenpfennig C, Nakayama M, Nihara R, Tomioka K, Helfrich-Förster C, Yoshii T. A Tug-
of-War between Cryptochrome and the Visual System Allows the Adaptation of Evening
Activity to Long Photoperiods in Drosophila melanogaster. J Biol Rhythms 2018;33.
https://doi.org/10.1177/0748730417738612.

[40] Vinayak P, Coupar J, Hughes SE, Fozdar P, Kilby J, Garren E, et al. Exquisite Light
Sensitivity of Drosophila melanogaster Cryptochrome. PLoS Genet 2013;9.
https://doi.org/10.1371/journal.pgen.1003615.

[41] Wheeler DA, Hamblen-Coyle MJ, Dushay MS, Hall JC. Behavior in Light-Dark Cycles of
Drosophila Mutants That Are Arrhythmic, Blind, or Both. J Biol Rhythms 1993;8.
https://doi.org/10.1177/074873049300800106.

[42] Kim EY, Bae K, Ng FS, Glossop NRJ, Hardin PE, Edery I. Drosophila clock protein is
under posttranscriptional control and influences light-induced activity. Neuron 2002;34.
https://doi.org/10.1016/S0896-6273(02)00639-6.

[43] Roenneberg T, Kuehnle T, Juda M, Kantermann T, Allebrandt K v., Gordijn MCM, et al.
Epidemiology   of   the   human   circadian   clock.   Sleep   Med   Rev   2007;11:429–38.
https://doi.org/10.1016/j.smrv.2007.07.005.

[44] Emery P, Stanewsky R, Helfrich-Förster C, Emery-Le M, Hall JC, Rosbash M. Drosophila
CRY   Is   a   Deep   Brain   Circadian   Photoreceptor.   Neuron   2000;26:493–504.
https://doi.org/10.1016/S0896-6273(00)81181-2.

[45]   Ingram KK, Kutowoi A, Wurm Y, Shoemaker DW, Meier R, Bloch G. The Molecular
733   Clockwork of the Fire Ant Solenopsis invicta. PLoS One 2012;7.
734   https://doi.org/10.1371/journal.pone.0045715.

735   [46]   Rubin EB, Shemesh Y, Cohen M, Elgavish S, Robertson HM, Bloch G. Molecular and
736   phylogenetic analyses reveal mammalian-like clockwork in the honey bee (Apis mellifera)
737   and shed new light on the molecular evolution of the circadian clock. Genome Res 2006;16.
738   https://doi.org/10.1101/gr.5094806.

739   [47]   Lazopulo S, Lazopulo A, Baker JD, Syed S. Daytime colour preference in Drosophila
740   depends on the circadian clock and TRP channels. Nature 2019;574.
741   https://doi.org/10.1038/s41586-019-1571-y.

742   [48]   Busza A, Emery-Le M, Rosbash M, Emery P. Roles of the two Drosophila
743   CRYPTOCHROME structural domains in circadian photoreception. Science (1979)
744   2004;304:1503–6.
745   https://doi.org/10.1126/SCIENCE.1096973/SUPPL_FILE/BUSZA.SOM.PDF.

 Figure Legends

Figure 1: Entrainment to white light skeleton photoperiods. (A) Mean (±SEM) locomotor activity

profiles of *CS* (top), *w^{1118}* (middle) and *yw* (bottom) flies under symmetric skeleton photoperiods

(left) and two asymmetric skeleton photoperiods (middle and right). The photoperiodic regimes

are shown on top of the profile plots. Black shaded regions indicate darkness and white regions

indicate light phases (400-500 lux) of the 24-h cycle. Note that the profiles under LD12:12 are

replotted across panels to facilitate pair-wise comparisons. (B) Phases of the evening peak of

activity for the three strains across the different photoperiodic conditions are shown. Boxplots that

share the same letter are not statistically significantly different from each other. Statistical

comparison of phases across all four photoperiodic conditions were done using Kruskal-Wallis

tests. *CS*: $\chi_3^2 = 95.05$, $p = 0$; *w^{1118}*: $\chi_3^2 = 123.47$, $p = 0$; *yw*: $\chi_3^2 = 113.58$, $p = 0$. Profiles and phases

of evening peak of activity are pooled from two independent replicate runs for each genotype.

Figure 2: Locomotor activity rhythms under various constant light conditions. (A) Spectral

composition of broad-spectrum white light overlayed with spectral composition of lights when

various filters were used, and with the action spectrum of CRYPTOCHROME (CRY). Data for

the spectral composition were extracted from the Lee Filters website (see Methods) and for the

action spectrum of CRY were taken from VanVickle-Chavez and Van Gelder, 2007 [24]. (B)

Example actograms of *CS* flies under constant darkness and constant light of various wavelength

compositions. Red lines on the actograms are eye-fit lines through the offset of locomotor activity.

Figure 3: Free-running locomotor activity rhythms of flies under constant blue- and UV-blocked

light. (A) Dose-dependent effect of constant light on the free-running period of *CS* (left), *w^{1118}*

(middle) and *yw* (right) flies. The lack of data point for highest intensity of light in *yw* flies is

because flies were arrhythmic under this condition. The yellow dashed line is dose response curve

fitted through a logistic model (see Methods). ANOVAs on the fitted logistic model revealed a

statistically significant effect of intensity on the free-running period for all three genotypes (*CS*: *p*

< 2e-16; *w^1118^*: *p* < 2e-16; *yw*: *p* < 2e-16). (B) Representative actograms of *Canton-S* (*CS*), *w^1118^*,

yw, *yw;;cry^OUT^*, *w;;glass^[60j]^* and *+;;norpA^[7]^* flies under constant darkness (top) and constant blue-

and UV-blocked light (bottom) of 30-45 $\mu W/cm^2$. (C) Frequency distributions of free-running

periods of all genotypes under constant dark and constant light conditions. The dots and dashed

vertical lines represent the median values of free-running periods under each regime. Statistical

comparisons are based on Wilcoxon's tests (*CS*: $W = 2676.5$, $p = 6.036e-08$; *w^1118^*: $W = 502$, $p =$

2.2e-16; *yw*: $W = 618.5$, $p = 1.583e-08$; *yw;;cry^OUT^*: $W = 7391$, $p = 2.233e-08$; *w;;glass^[60j]^*: $W =$

305, $p = 1.092e-05$; *+;;norpA^[7]^*: $W = 10$, $p = 2.2e-16$). Values are pooled from at least two

independent runs (see Suppl. Table 1).

Figure 4: Example actograms from all six genotypes illustrating entrainment, relative

coordination, free-running and/or arrhythmicity. The red dashed lines are drawn to trace the offset

of locomotor activity in the top three rows. The percentage values on top of each actogram are the

percentage of all flies showing that behavior. The shading on the actograms represents the ramping

up and down of light, starting at Zeitgeber Time-00.

Figure 5: Entrainment to ramped blue- and UV-blocked light cycles. (A) Mean (±SEM)

locomotor activity profiles of entrained *CS* (left), *w^1118^* (middle) and *yw* (right) flies under ramped

light cycles (top) and on the first day under constant low intensity light (bottom). (B) Polar plots

showing the phases of the predominant peak of locomotor activity for the three genotypes (*CS* –

left; *w¹¹¹⁸* – middle; *yw* – right) under entrainment and on the first day post entrainment under

constant conditions. Phase control was statistically examined using V-tests (see Methods). *CS*: *p*

= 7.67e-17; *w¹¹¹⁸*: *p* = 1.45e-10; *yw*: *p* = 7.38e-10. (C) Representative actograms showing rhythmic

and arrhythmic flies under ramped yellow cycles for loss-of-function clock mutants. The red

dashed line indicates the offset of locomotor activity. The shading on the actograms represents

the ramping up and down of light, starting at Zeitgeber Time (ZT) 00. (D) Mean (±SEM)

locomotor activity profile (left) and polar plot depicting phases of the peak of activity of *per⁰¹* flies

under ramped light cycles. The black dashed line in the top row of panel (A) and (D-left) is a

schematic representation of light ramping from a lower minimum value (~10 $\mu W/cm^2$) at ZT00 to

the maximum value (40-50 $\mu W/cm^2$) at ZT12 and back to the minimum value by ZT24. In all the

polar plots individual dots represent a single fly and the line from the center points to the mean

phase across all flies. The distance of these lines from the center is an estimate of the across-fly

variation in phase. Values close to 1 (gray dashed circle) have lower variances.

Supplementary Figure and Table Legends

Suppl. Table 1: Details of experiments performed to assay free-running locomotor activity rhythms in various genotypes of flies under constant darkness and constant light (also see Fig. 3).

Suppl. Table 2: Percentage of flies entraining to each of the four environmental conditions and the total number of flies based on which these percentages were computed. There was no statistically significant reduction in the percentage of entrainment for any of these strains under any photoperiodic regime. Statistical comparisons were made using the Fisher Exact Test.

Suppl. Table 3: Percentage of flies that belong to one of the four entrainment/rhythm categories under ramped blue- and UV-blocked light cycles, along with the total number of flies based on which these percentages were calculated.

Suppl. Table 4: Percentage of flies that are rhythmic and arrhythmic for loss-of-function clock mutants under ramped blue- and UV-blocked light cycles, along with the total number of flies based on which these percentages were calculated.

Suppl. Figure 1: Power of circadian locomotor rhythms under constant darkness (DD) and constant light (LL) conditions for background controls and photoreceptor mutants. (A) Power of the rhythm for *CS* (left), w^{1118} (middle) and *yw* (right). Statistical comparisons are based on Wilcoxon's tests (*CS*: $W = 3749$, $p = 0.0071$; w^{1118}: $W = 7579$, $p = 1.525e\text{-}12$; *yw*: $W = 3196$, $p = 4.799e\text{-}12$). Values of power are pooled from at least two independent runs (see Suppl. Table 1). $* < 0.05$, $** < 0.01$, $*** < 0.001$, and NS (Not Significant). (B) Power of the rhythm for three

photoreceptor mutants. Statistical comparisons are based on Wilcoxon's tests ($yw;;cry^{OUT}$: $W =$

5289, $p = 0.76$; $w;;glass^{[60j]}$: $W = 1315$, $p < 0.001$; $+;;norpA^{[7]}$: $W = 2365$, $p < 0.001$). Values of

rhythm power are pooled from at least two independent runs (see Suppl. Table 1). * < 0.05, ** $<$

0.01, *** < 0.001, and NS (Not Significant).

Suppl. Figure 2: Entrainment to white light skeleton photoperiods. (A) Mean (±SEM) locomotor

activity profiles of $yw;;cry^{OUT}$ flies under symmetric skeleton photoperiods (left) and two

asymmetric skeleton photoperiods (middle and right). The photoperiodic regimes are shown on

top of the profile plots. Black shaded regions indicate darkness and white regions indicate light

phases (400-500 lux) of the 24-h cycle. Note that the profiles under LD12:12 are replotted across

panels to facilitate pair-wise comparisons. (B) Phases of the evening peak of activity across the

different photoperiodic conditions are shown. Boxplots that share the same letter are not

statistically significantly different from each other. Statistical comparison of phases across all four

photoperiodic conditions were done using Kruskal-Wallis tests. $\chi^2_3 = 63.46$, $p = 1.07e-13$. Profiles

and phases of evening peak of activity are pooled from two independent replicate runs for each

genotype.

Figure 1

A

White light   CRY action spectrum   Zone of CRY activation

Green   Orange   Yellow   Blue+UV blocked   Red

Normalized intensity

Wavelength (nm)

log$_{10}$ % sensitivity of CRY

B

Representative *CS* actograms

LL

DD   Green   Orange   Yellow   Blue+UV blocked   Red

Days

Time

Figure 2

Figure 3

Figure 4

Figure 5