Validation of carbon isotopologue distribution measurements by GC-MS and application to [13]C-metabolic flux analysis of plant tricarboxylic acid cycle

Younès Dellero[1, 2, 4*], Solenne Berardocco[1, 2], Cécilia Berges[3, 4], Olivier Filangi[1, 2, 4], Alain
Bouchereau[1, 2, 4]

[1]Institute for Genetics, Environment and Plant Protection (IGEPP), National Research Institute for
Agriculture, Food and Environment (INRAE), Université Rennes, Institut Agro , Le Rheu, France

[2]Plateau de Profilage Métabolique et de Métabolomique (P2M2), Institute for Genetics, Environment
and Plant Protection, Biopolymers Interactions Assemblies, French National Research, Le Rheu,
France

[3]TBI, Université de Toulouse, National center for Scientific Research (CNRS), National Institute for
Research for Agriculture, food and Environment (INRAE), National Institute of Applied Sciences
(INSA), Toulouse, France;

[4]MetaToul-MetaboHUB, National infrastructure of metabolomics and fluxomics, Toulouse, France

* Correspondence:
Corresponding Author
younes.dellero@inrae.fr

Keywords: Isotope, gas chromatography-mass spectrometry, trimethylsilyl (TMS) derivatives,
accuracy, regulation, phosphoenolpyruvate carboxylase, decarboxylation. (Min.5-Max. 8)

Abstract

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of [13]C-Metabolic Flux Analysis at
isotopically non-stationary steady-state. This approach requires a high level of accuracy for the
measurement of Carbon Isotopologue Distribution in plant metabolites, which has still not been
accurately evaluated at the isotopoloque level for GC-MS. Thus, uncertainties remain about the
accuracy of the CID measured and subsequently, the metabolic fluxes associated. Moreover, a
significant part of our knowledge about the functioning and the regulation of plant TCA cycle is solely
based on [13]C-labeling experiments analyzed with Nuclear Magnetic Resonnance. But this method is
not very sensitive at low [13]C abundance and can hardly fully resolve isotopologue dynamics inherent
to the TCA cycle compared to MS. Here, we developed a workflow to validate the measurements of
CIDs from plant metabolites with GC-MS by producing tailor-made *E. coli* standard extracts harboring
a predictable binomial CID for all metabolites. Most of our TMS-derivatives mass fragments were
validated with these standards and at natural isotope abundance in plant matrices. Then, we applied
this validated MS method to investigate the light/dark regulation of plant TCA cycle metabolic fluxes
by incorporating U-[13]C-pyruvate to *Brassica napus* leaf discs. We took advantage of pathway-specific
isotopologues/isotopomers to show that the TCA cycle has a nearly four-fold higher contribution to
Citrate and Malate production than phosphoenolpyruvate carboxylase (PEPc). Overall, the dynamic of
[13]C-enrichment in Citrate, Glutamate, Succinate and Malate ruled out the hypothesis of the stored
citrate contribution to TCA cycle and highlithed the major contribution of the forward plant TCA cycle
to malate biosynthesis in both dark and light conditions. Interestingly, the light-dependent [13]C-

incorporation into Glycine and Serine showed that nearly 5% of TCA cycle-derived CO_2 was actively
reassimilated by the photosynthesis. We also observed an increase of PEPc contribution to malate
production in the light while there was a higher involvement of PEPc-derived carbons to citrate
production in dark condition. The results suggested an important role for Malate import into
mitochondria in sustaining TCA cycle in dark conditions.

1   INTRODUCTION

Metabolic fluxes within central metabolism are among the most important determinants of plant cell
physiology, supporting central energetical functions and allowing them to quickly adapt to
environmental changes by controlling metabolite turnover (Clark et al., 2020). Given that most of
fluxes supported by a metabolic network cannot be measured directly, Metabolic Flux Analysis (MFA)
technics were developed to infer information about metabolic fluxes from isotopic labeling
experiments (Ratcliffe and Shachar-Hill, 2006).^{13}C-MFA relies on the tracing of ^{13}C throughout a
metabolic network in a relative metabolic steady-state to model and/or compare metabolic fluxes, *i.e.*
that the metabolic fluxes are stable across the experiment because the associated metabolite pool sizes
are relatively constant over the time of labeling except for the source (consumption) and sink
(production) metabolites (Allen and Young, 2019;Antoniewicz, 2021). ^{13}C-MFA is usually performed
at isotopic steady-state for heterotrophic tissues for practical reasons, and also because they can reach
this isotopic state (Hay et al., 2014). Conversely, ^{13}C-MFA in phototrophic tissues can only be
performed at isotopically non-stationary steady-state (INST-MFA), thus requiring a higher accuracy in
the measurement of ^{13}C enrichment within metabolites (Cheah and Young, 2018). Over the last
decades, ^{13}C-MFA has been applied successfully to microorganisms to highlight new metabolic routes
and identify metabolic bottlenecks for the purpose of metabolic engineering (Cheah et al.,
2020;Antoniewicz, 2021;Treves et al., 2022). In plants, ^{13}C-INST-MFA has been scarcely applied to
date, due to the complexity in modelling accurately metabolic fluxes in a multicompartmental system
(Heise et al., 2015;Kruger and Ratcliffe, 2021;Wieloch, 2021). However, accurate measurement of
^{13}C-enrichment is a prerequisite for precise evaluation of isotopic labeling dynamics and accurate
modelling of metabolic fluxes. Indeed, small errors in mass isotopologue distribution can propagate to
large errors in estimated fluxes (Antoniewicz et al., 2006).

Two major analytical methods are currently employed to analysis the Carbon Isotope Distribution
(CID) within plant metabolites during ^{13}C-isotope tracing experiments: Nuclear Magnetic Resonance
(NMR) and Mass Spectrometry (MS). NMR analysis of the ^{13}C enrichment is generally performed
with ^{13}C-decoupling ^{1}H sequences and provides information respectively at isotopomer level (by
integrating ^{13}C-^{13}C interactions) (Kruger et al., 2007;Masakapalli et al., 2010;Abadie et al., 2017).
Although this method is quantitative for the ^{13}C at the isotopomer level and allows to significantly
separate carbon atoms of different chemical structures, it has a low sensitivity at natural ^{13}C abundance
(large amounts of fresh material required), thereby affecting the accuracy for ^{13}C enrichment
measurements regarding metabolites found in small quantities (Dellero et al., 2016;Abadie and
Tcherkez, 2021). In addition, this method is time-consuming due to the longer relaxation time for ^{13}C
nucleus and the need to accumulate many spectrums to get a good signal (Abadie and Tcherkez, 2021).
As a consequence, most of the ^{13}C-tracing isotope experiments recently performed in plants and
analyzed by ^{13}C-NMR have only one time-point (Abadie et al., 2016;Abadie et al., 2018;Abadie and
Tcherkez, 2019a;b), which questions the estimation of metabolic fluxes therein (metabolic/isotopic
steady-state and treatment of inactive metabolite pools). To date, the accurate estimation of metabolic
fluxes in phototrophic tissues requires to determine the isotopic and metabolic state of the system by
coupling transient kinetics of ^{13}C-labeling with absolute quantification of metabolite pools (Kruger and
Ratcliffe, 2015;Cheah and Young, 2018;Clark et al., 2020;Antoniewicz, 2021). Some methodological

improvements were recently made with highly sensitive [1]H-derived sequences to improve the
resolution of the [13]C atoms at the isotopomer level (Sinnaeve et al., 2018), and its future use for [13]C-
INST-MFA in plants will provide interesting perspectives. Conversely, mass spectrometry has been
preferentially used to analysis the Carbon Isotope Distribution (CID) within plant metabolites during
time-course [13]C-isotope tracing experiments (Szecowka et al., 2013;Ma et al., 2014;Medeiros et al.,
2018;Treves et al., 2022). This method is more sensitive at natural [13]C abundance and can be
implemented at high throughput. However, MS only gives information at the isotopologue level and
an absolute quantification of the metabolite content is generally required to subsequently infers
metabolic fluxes. In addition, the analysis of CID for amino acids by LC-MS revealed some bias and
errors in the measurement of CID (Heuillet et al., 2018), thus highlighting the need to evaluate
measurement accuracy with GC-MS fragments for subsequent [13]C-MFA in plants.

Among plant metabolites, organic and amino acids are major players for plant primary metabolism.
They are actively involved in crucial physiological processes such as photorespiration, *de novo*
nitrogen assimilation and mitochondrial respiration (Xu et al., 2012;Zhang and Fernie, 2018;Dellero,
2020;Timm, 2020). Most of the carbon backbones for amino acid biosynthesis arise from organic acids
of the tricarboxylic acid (TCA) cycle (Galili et al., 2016). Interestingly, an important part of our
knowledge about the functioning and regulation of plant TCA cycle is almost exclusively based on
[13]C-labeling studies that used NMR. These previous works suggested that the plant TCA cycle is not
cyclic in the light, with two branches acting separately (Tcherkez et al., 2009;Sweetlove et al., 2010):
i) a first branch stimulated by the anaplerotic activity of phosphoenolpyruvate carboxylase (PEPc)
leading to the production of Oxaloacetate-malate-fumarate ; ii) a second branch stimulated by the
pyruvate dehydrogenase activity leading to the production of citrate-isocitrate-2oxoglutarate-
glutamate. Based on the NMR analysis of combined [13]CO_2 and [15]NH_4[15]NO_3 labeling experiments in
oilseed rape leaves, it was proposed that a high proportion of non-labeled citrate (vacuolar pool
probably) was involved in the synthesis of 2-oxoglutarate under light conditions to sustain *de novo*
nitrogen assimilation through glutamate (Gauthier et al., 2010). Under this scenario, the citrate pool
would be replenished during dark conditions from glycolysis-derived [13]C-pyruvate, leading to a higher
[13]C enrichment of the citrate when compared to light conditions. Thus, the light and dark environmental
conditions are expected to introduce important variations of metabolic fluxes within the plant TCA
cycle and its associated PEPc anaplerotic pathway. However, such conclusions and hypotheses remain
to be carefully investigated with mass spectrometry and a [13]C-MFA framework, *i.e.* with time-course
[13]C-isotope tracing experiments and pool size quantification.

Currently, the routine analysis of CID on plants is often assessed for organic and amino acids by GC-
MS using TMS-derivatives (Szecowka et al., 2013;Araujo et al., 2014;Lima et al., 2018;Treves et al.,
2022). This TMS-based method notably allowed to follow the fate of sucrose breakdown within guard
cells in connection with TCA cycle (Medeiros et al., 2018). More recently, a [13]C-positionnal
isotopomer approach was also proposed for glutamate and malate to investigate the fate of PEPc-
derived carbon from [13]C-labeling experiments (Lima et al., 2021). The comparison of two
derivatization methods suggested that TBDMS-derivatives were potentially less prone to errors for
amino acid analysis than TMS-derivatives (Antoniewicz et al., 2007;Young et al., 2014). However, the
suitability of a TMS-derivative molecular ion in each of these studies was only performed using U-
[13]C-metabolites, which do not evaluate the entire CID but only the carbon distribution between the
unlabeled isotopologue and the fully labeled isotopologue (M0 and M5 for a 5-carbon metabolite).
Unfortunately, isotopic effects can occur at all stages of the analytical procedures and depend on the
tested metabolite and its isotopic content. For instance, the present methodologies used have not
assessed the impact of each isotopologue to the final isotopologue distribution, which can lead to
important bias during interpretation of the results. Thus, the reliability of CID measurements by GC-

MS has to be evaluated for each individual isotopologue of all the compounds of interest to
subsequently ensure a reliable use for monitoring isotopic labeling dynamics. This is even more
important for plants, given that plant metabolites show strong differences of ^{13}C-enrichment dynamics
between amino and organic acids (Szecowka et al., 2013). For this purpose, the recent development of
an *E. coli* ^{13}C biological standard with controlled patterns of CID for organic and amino acids may
help to address this issue (Millard et al., 2014). Indeed, the recent use of these reference samples
allowed to identify important analytical biases in the CID measurement of Leucine and Asparagine in
a concentration-dependent manner by LC-MS (Heuillet et al., 2018). Interestingly, this validated LC-
MS method successfully measured nitrogen isotope distribution within plant amino acids and led to
the accurate estimation of metabolic fluxes from ^{15}N-isotope transient labeling experiments in oilseed
rape leaves at a metabolic steady-state using the ScalaFlux approach (Dellero et al., 2020b;Heuillet et
al., 2020;Millard et al., 2020).

Here, we proposed a workflow to assess the measurement accuracy of CID from organic and amino
acids by GC-MS for applications in plants. To do so, we built up an analytical method maximizing
information on carbon backbones and analyzed two specific *E. coli* ^{13}C biological standards with
controlled patterns of CID for organic and amino acids (^{13}C-PT). We also analyzed chemical standards
and plant samples at natural ^{13}C abundance to identify potential bias arising from matrix and analytical
systems. Next, we applied our MS method to investigate the light/dark regulation of metabolic fluxes
within plant TCA cycle by performing U-^{13}C-pyruvate labeling kinetic experiments on oilseed rape
leaves under light or dark conditions. Using ^{13}C-MFA principles and some specific isotopologues, we
calculated the fractional contribution of phosphoenolpyruvate carboxylase (PEPc) and TCA cycle to
malate and citrate production and proposed a simplified carbon flux map with relative metabolic fluxes.
Our results notably revealed that under non-limiting pyruvate conditions, PEPc activity remained
nearly four-fold lower than the TCA cycle activity independently of the light/dark stimulation. In
addition, we showed that pyruvate dehydrogenase and TCA cycle-derived CO_2 contributed to around
5% of net CO_2 assimilation under light conditions.

2   MATERIALS AND METHODS

Organisms and culture

The strain Escherichia coli K-12 MG1655 was grown on minimal medium containing 5 mM
KH_2PO_4,10mM Na_2HPO_4, 9 mM NaCl, 40 mM NH_4Cl, 0.8 mM $MgSO_4$, 0.1 mM $CaCl_2$, 0.3 mM
thiamine, and 45 mM of the ^{13}C-labeled Acetate mixture as the sole carbon source. Acetate and
thiamine were sterilized by filtration, whereas the other compounds were autoclaved separately. Batch
cultures were carried out in 1-L baffled flasks with 500 mL of medium sparged with synthetic gas (80%
N_2/20%O_2 containing <1 ppm CO_2, Air Liquide). Cultivation was performed in a Multifors Bioreactor
(Infors HT, Bottmingen-Basel, Switzerland) at 37 °C and 200 rpm of orbital shaking. The pH was
maintained at 7.0 by the addition of 2 M HCl. Cell growth was monitored by measuring optical density
at 600 nm with a Genesys 6 spectrophotometer (Thermo, Carlsbad, CA, USA). Amount of biomass in
each sample was calculated using a coefficient of 0.37 g of cell dry weight per OD unit. Arabidopsis
thaliana seeds were sterilized for 10 min with absolute ethanol containing 7.5% NaClO and rinsed two
times with absolute ethanol. Seeds were sowed on ½ MS media containing 1% Agar and incubated to
the dark at 4°C for 48-72 hours to break seed dormancy. Seedlings were grown at ambient air under a
day/night cycle with 16 h light/8h dark at a light intensity of 120 µmol photons.m^{-2}.s^{-1} and 20°C
day/18°C night. Seedlings were harvested after 2-3 weeks by instant sampling with liquid nitrogen and
lyophilized for 96 h before to be stored at -80°C until analysis. 2-month old *Brassica napus* plants were

obtained from seeds exactly as described before (Dellero et al., 2020a). The leaf L9 was used for all
experiments (natural abundance and U-^{13}C-pyruvate incorporation).

U-^{13}C-pyruvate labeling experiments

Four hours after the beginning of the light period, each leaf was cut at the basis of the petiole and then
66 leaf discs (0.8 cm^2) were randomly punched with a cork-borer in both laminas of each leaf. Leaf
discs were first floated in a 10 mM MES-KOH (pH 6.5) buffer for 15 min in order to process the L9
leaf for four plants. In this experiment, one leaf of a plant corresponded to a biological replicate. Then
for each time point, six leaf discs from the same leaf were transferred in a well of a 6-well microplate
filled with 4 mL/well of a buffer containing 10 mM MES-KOH (pH 6.5) and 10 mM U-^{13}C-Pyruvate
(^{13}C isotopic purity of 99%) (Eurisotop, France). All incubations were performed at 20 °C with an
orbital shaking of 70 rpm under either a continuous light of 120 µmol photons m^{-2} s^{-1} or under dark
conditions using multiple opaque plastic bags, similarly to the previous experimental set-up used for
dark-induced senescence (Dellero et al., 2020a). The timepoint 0 was the same for both conditions as
it corresponded the beginning of the experiment. For each time point (0, 0.5, 1, 2, 4, 6 hours of
incubation), the leaf discs were harvested and rinsed three times (10 sec) with a neutral buffer (i.e.,
without U-^{13}C-Pyruvate) and dried for 10 s on a clean tissue before being frozen in liquid nitrogen and
stored at -80 ∘C. The timepoint 0 of the experiment was incubated 2 sec with the labeled buffer before
being rinsed in order to evaluate the fraction of ^{13}C-pyruvate detected in our analysis that was not
incorporated inside leaf discs (leading to a mean ^{13}C enrichment close to 35% for pyruvate at T0 in our
study).

Analysis of ^{13}C-labeled Acetate by NMR

An equimolar mixture of all ^{13}C-isotopomers of acetate was produced from ^{12}C-Acetate, 1-^{13}C-Acetate,
2-^{13}C-Acetate and U-^{13}C-Acetate (13C-purity 99%, Eurisotop, France; **Figure 1**). Quantitative 1H 1D-
NMR was used to control the isotopic composi-tion of the ^{13}C-labeled substrate before use. NMR
spectra were recorded with an Avance 500 MHz spectrometer (Bruker, Rheinstetten, Germany) at 298
K, using a 30° pulse and a relaxation delay of 20 s. The proportion of each isotopic form was quanti-
fied by fitting using a mixed Gaussian−Lorentzian model (Millard et al., 2017).

Metabolite extraction and quantification

^{13}C-PT full metabolites and ^{13}C-PT protein hydrolysates extracts were obtained following procedures
described previously (Heuillet et al., 2018). Briefly, for the ^{13}C-PT full metabolites extract, 1 mL of *E.*
coli broth at OD600 nm = 1.7 was micro-filtered and the pellet was rapidly quenched at -20 °C into a
mixture of acetonitrile/methanol/H$_2$0 (2:2:1) with 0.1% formic acid. After a centrifugation step (7000g,
−20 °C, 15 min), the supernatant was evaporated and resuspended in 250 µL of Milli-Q water, to be
stored at −80 °C until analysis. For the ^{13}C-PT protein hydrolysates extract, the pellet obtained with
the cellular extract was subjected to an acid hydrolysis with 6N HCl at 110 °C overnight. Samples were
evaporated and rinsed 3 times with Milli-Q water. After a centrifugation step (10 min, 12 000g), the
supernatant was stored at −80 °C. Polar metabolites of freeze-dried Arabidopsis seedlings and *B. napus*
leaves were extracted with a MeOH/Chloroform/H$_2$O mixture following procedures described
previously (Dellero et al., 2020b). The quantification of amino acid content was performed by HPLC-
UV exactly as described previously (Dellero et al., 2020b). The quantification of organic acids was
performed by GC-FID exactly as described previously (Bianchetti et al., 2021).

Determination of Carbon Isotopologue Distribution by GC-MS

Dried aliquots of samples were derivatized on a MultiPurpose Sampler Gerstel and analyzed by GC-
MS (GC-2010 Plus Shimadzu coupled to a triple quadrupole mass spectrometer GCMS-TQ8040
Shimadzu). Polar metabolites were first derivatized for 1 h at 40°C with 50 µL of pyridine containing
20 g.l^{-1} methoxyamine chloride and then were incubated with 50 µL of N-Methyl-N-(trimethylsilyl)-
trifluoroacetamide (MSTFA) for 0.5 h at 40°C. 1 µL was injected with a split mode 1/20 at 260°C on
a TG-5MS column (26098-1420, Thermo Scientific). Separation was performed in a Helium gas-
stream at 1 mL.min^{-1} using the following temperature ramp : 4 min at 70°C, +10°C.min^{-1} until 198°C,
2 min at 198°C, +1°C.min^{-1} until 202°C, +15°C.min^{-1} until 268°C, 3 min at 268°C, +1°C.min^{-1} until
272°C,+10°C.min^{-1} until 310°C and 7 min à 310°C. The transfer line was maintained at 300°C and the
ion source at 200°C. Ionization was performed by electron impact at 70 eV and the MS acquisition was
carried out after a solvent delay of 5 min in full scan mode at 7 spectra.s$_{-1}$ over the m/z range 45-700.
MS Spectra were processed with the software "GCMS Solutions" supplied by Shimadzu to adjust
retention times between each run using a mixture of alkanes (C7 to C40, Sigma-Aldrich). Peak identity
was established according to their retention time and mass fragmentation spectra with respect to
authentic standards derivatized (**Table S1**), and by reference to mass spectra of derivatives in the US
National Institute of Standards and Technology database and in the Golm metabolome database. All
mass isotopologue raw areas were subjected to correction for the contribution of naturally occurring
isotopes of all elements except the carbon arising from the metabolite backbone using the IsoCor v2.1.4
package of Python (https://github.com/MetaSys-LISBP/IsoCor) in low resolution mode (Millard et al.,
2019). In addition, unlabeled chemical standards and Arabidopsis and *B. napus* leaves were
additionally corrected for the contribution of naturally occurring isotopes of the carbon arising from
the metabolite backbone (giving a residual molecular ^{13}C enrichment of 0 for molecules labeled at
natural ^{13}C abundance). ^{13}C-PT samples were not concerned by this additional correction since they
held fully controlled ^{12}C/^{13}C patterns at all positions. A correction method dedicated to IsoCor was
specifically created for the TMS-derivative fragment described in this study (**Table 1**), *i.e.* by
considering the contribution of all atoms added by the derivatization on the isotopic measurement of
the monitored fragments (chemical separation of all atoms arising from the metabolite and the
derivative). All files required to run this method on IsoCor are available in the supplementary material
of this article (**Supplementary files**) and all information will be updated indefinitely in the following
INRAE dataverse (Dellero and Filangi, 2021). The method is publicly available and can also be run
directly from GC-MS source files using the following Galaxy workflow (https://galaxy.genouest.org/,
section "Metabolomics", subsections "Conversion GCMS PostRun Analysis to IsoCor" and "Isotope
Correction for mass spectrometry labeling experiments"). For this purpose, each raw GC-MS dataset
should contain a column "Name" for the fragments considered and a column "Area". The name of each
fragment must be written exactly as specified in the "Metabolite.dat" (an example file is provided in
the **supplementary material**). Fractional Carbon Isotopologue Distribution and mean ^{13}C enrichment
were directly obtained from IsoCor outputs (isotopologue_fraction and mean_enrichment) while CIDs
were converted to binomial distributions by multiplying with 2^n, where n is the number of carbon atoms
(Millard et al., 2014). The accuracy of measurements was assessed from the closeness of experimental
and predicted values for CIDs and mean ^{13}C enrichments at either ^{13}C-PT abundance or at natural ^{13}C
abundance. CID accuracy was formally defined as the mean of absolute differences between measured
and predicted values for each isotopologue (Heuillet et al., 2018).

Pathway fractional contribution

Pathway fractional contribution (FC) can be applied to a metabolic network in a metabolic and isotopic
steady-state to compare the relative contribution of the considered pathway to the production of a
metabolite, as long as these pathways produced specific isotopologues that can be monitored (Buescher
et al., 2015). Here, FC was calculated from PEPc and TCA cycle-specific mass

isotopomers/isotopologues for Malate and Citrate from the timepoint 4 and 6 hours (metabolic steady-
state and assumed similar pseudo-isotopic steady-state). In the following equations, *mi* refers to the
isotopologue enrichment calculated with respect to the CID (sum of all *mi* for a given metabolite equal
to 1). As all these calculations are relative, it is also possible to use corrected areas of mass
isotopologues (outputs from IsoCor). For Malate_2TMS_C1C4, M1 isotopologue measurement
reflected both PEPc and TCA cycle isotopomers. However, the ratio between M1 and M3
isotopologues of Malate_2TMS_C1C4 produced by the TCA cycle (denoted *X* in the following
equations) can also be observed in Succinate_2TMS_C1C4 and Glutamate_2TMS_C2C5, thereby
reflecting the M0 and M2 from mitochondrial active pool of Acetyl-CoA, *i.e.*:

$$Succinate C1C4 \left(\frac{m3}{m1 + m3} \right) = Glutamate C2C5 \left(\frac{m3}{m1 + m3} \right) = Acetate \left(\frac{m2}{m0 + m2} \right) = X$$

From this parameter, we calculated the FC of PEPc and TCA cycle to Malate and Citrate production
as follows:

$$FC\ Malate_{PEPc} = Malate C1C4 \left(\frac{m1 - \left(\left(\frac{1 - X}{X} \right) \times m3 \right)}{m1 + m2 + m3} \right)$$

$$FC\ Malate_{TCA\ cycle} = Malate C1C4 \left(\frac{m2 + \frac{m3}{X}}{m1 + m2 + m3} \right)$$

$$FC\ Citrate_{PEPc} = Citrate C1C6 \left(\frac{m1 + m3}{m1 + m2 + 2 \times m3 + m4} \right)$$

$$FC\ Citrate_{TCA\ cycle} = Citrate C1C6 \left(\frac{m2 + m3 + m4}{m1 + m2 + 2 \times m3 + m4} \right)$$

It should be noted that *m3* was used twice in the FC Citrate calculations because this isotopologue is
produced by both pathways. Finally, the FC of pyruvate dehydrogenase (PDH) and TCA cycle-derived
CO_2 to photosynthesis was calculated using *X* as a proxy for the $^{13}C/^{12}C$ ratio of this CO_2 fraction in
light conditions. Considering the isotopic and metabolic steady-state for these metabolites, the mean
^{13}C enrichment of *X* should be observed in Serine if this fraction of CO_2 accounted for 100% of CO_2
fixed by photosynthesis, *i.e.*:

$$FC\ photosynthesis_{PDH\ and\ TCA\ cycle\ CO_2} = \frac{mean\ ^{13}C\ enrichment\ (Serine C1C3)}{X}$$

Mean ^{13}C enrichment of Serine C1C3 was recalculated from M1 isotopologue of Serine_2TMS_C1C2
and Serine_2TMS_C2C3 assuming that Serine was only labeled in the C2 of these fragments based on
the validated hypothesis (**Figure 7B**). Statistical comparisons were performed using a Student-test (2
groups) or an ANOVA followed by a post-hoc Tukey-HSD test (3 or more groups) with a level of p-
value <0.05 and Rstudio (RStudio Team, 2016).

3   RESULTS

The proposed workflow (**Figure 1**) consisted in three steps: 1) the build-up of a method to identify and
select abundant molecular ions deriving from TMS-derivatives with a large number of carbons; 2) the
production of two reference [13]C-labeled samples with a controlled pattern for CID using *E. coli* as a
metabolic factory; 3) the validation of TMS-derivatives mass fragments for CID measurements with
our GC-MS method for applications in plants using unlabeled and [13]C-labeled plant samples.

**Selection of GC-MS mass fragments maximizing information on carbon back-bones of organic
and amino acids**

We combined a literature survey analyzing common pathway of TMS-derivative mass fragmentations
with the injection of chemical standards in our analytical system to select some abundant fragments
maximizing the information on carbon backbones. Their chemical identity was further confirmed by
the observed CID pattern with our two reference [13]C-labeled samples with a controlled pattern for CID.
Derivatives with only one TMS group were not considered in this study due to the low yield of these
fragments after fragmentation and the occurrence of important contaminations for low m/z. Overall,
TMS-derivatives of most α-amino acids were characterized by an α-cleavage of the carboxylic function
in the form of a TMS-CO_2 (m/z 117) (Harvey and Vouros, 2020), thus leading to the abundant
formation of fragments containing all others atoms of the α-amino acids except the C1 (**Table 1**). Only
Ornithine(4TMS) could be detected without any mass loss (**Table 1**, (Harvey and Vouros, 2020)).
Aromatic amino acids showed an additional α-cleavage of the amine function leading to the additional
production of an abundant m/z 218 fragment containing the C1-C2 for Phenylalanine, Tryptophan and
Tyrosine (**Table 1**, (Lai and Fiehn, 2018;Harvey and Vouros, 2020)). Only tryptophan did not proceed
through an α-cleavage of the carboxylic function compared to Phenylalanine and Tyrosine, but the
other fragment containing the C3 to C11 atoms resulting from the α-cleavage of the amine function
could be abundantly detected. Serine(3TMS) and Valine(2TMS) showed also some specific abundant
molecular ions in addition to the α-cleavage of the carboxylic function (**Table 1**,(Harvey and Vouros,
2020)). Arginine, Histidine and Cysteine could not be detected by GC-MS using TMS derivatization
and were not further considered in our study. Beside this, β-Alanine, an important β -amino acid from
plants, showed a classical abundant fragment resulting from the loss of methyl from a TMS group,
which is known to be very low in α-amino acids (**Table 1**, (Lai and Fiehn, 2018)). Interestingly,
fragmentation of the TMS-derivatives of organic acids related to the TriCarboxylic Acid cycle
proceeded through an α-cleavage within a TMS group leading to the loss of a methyl radical and
producing a [M-15] fragment containing all carbon atoms. However, this fragment was very weak for
Citrate and Malate. The highest fragments maximizing information on carbon backbones were m/z 273
(C1 to C5) for Citrate(4TMS) and m/z 233 (C2 to C4) for Malate(3TMS). The identity of the carbons
hold by these fragments has been recently assigned using position-specific [13]C-labeled standards
prepared by *in vitro* enzymatic reactions (Okahashi et al., 2019).

**Evaluation of CID measurements by GC-MS using reference [13]C-PT samples and plant samples
at natural isotope abundance**

Next, we produced reference [13]C-labeled samples using *E. coli*, denoted here [13]C-PT. This strategy
relies on the use of an accurate equimolar mixture of all [13]C isotopomers of acetate to biologically
produce metabolites with fully controlled and predictable [13]C-labeling patterns (Millard et al., 2014).
Thus, molecular [13]C enrichment of each carbon-containing metabolite will be theoretically equal to
50% and all isotopomers will be present in equimolar concentrations, leading to a binomial distribution
of isotopologues following Pascal's triangle. For this purpose, we grew *E. coli* in a minimal media with
this acetate mixture as the sole carbon source of the media (**Figure 1**) and checked its isotopic content
by NMR (molecular [13]C enrichment expected at 49.9 % ± 0.8). From cell harvest, two specific

reference [13]C-PT samples were produced during the extraction: i) a [13]C-PT full metabolite extract
containing all polar metabolites for the evaluation of CID for organic acids; ii) a [13]C-PT protein
hydrolysates extract containing proteogenic amino acids in high concentrations for the evaluation of
CID for amino acids. This strategy allowed to maximize the amount of amino acids that can be weakly
abundant as soluble pools when *E. coli* is grown in a minimal media. It should be mentioned that the
hydrolysis process deamidated Asparagine into Aspartate and Glutamine into Glutamate, thus these
amino acids could not be evaluated (Kambhampati et al., 2019). In addition, non-proteogenic amino
acids from the **Table 1** could not be evaluated with this extract (Ornithine, β-Alanine).

To facilitate comparison of CIDs between the different amino acids, we converted the measured [13]C-
enrichments among isotopologues of each fragment into binomial distribution, thus reflecting the
number of isotopomers considered for each isotopologue (**Figure 2** and Material and Methods).
Overall, all the considered amino acids were detected in the [13]C-PT protein hydrolysates except
Tryptophan (**Figures 2A, S1A**). We found very a high level of correlation between the measured CID
(blue bars) and the expected CID (black bars) for numerous amino acid fragments considered in our
method: α-Alanine_2TMS_C2C3,   Aspartate_C2C4   either   in   2TMS   or   3TMS,
Glutamate_3TMS_C2C5, Glycine_3TMS_C2, Isoleucine_2TMS_C2C6, Leucine_2TMS_C2C6,
Phenylalanine_2TMS_C1C2 and C2C9, Serine_2TMS_C2C3, Serine_3TMS_C1C2 and C2C3 and
Tyrosine_2TMS_C1C2. CID measurements for Methionine_2TMS_C2C5 showed a high variability
coupled to a lower estimation of M3 and M4 isotopologues at the expense of M1 and M2 (**Figure 2A**).
However, this amino acid was in a very low abundance in the [13]C-PT protein hydrolysates compared
to the others amino acids (**Figure S1A**). Lysine and Tyrosine_2TMS_C2C9 had also important
variability for their CID measurements, leading to misestimations of their M3 isotopologues. The most
important and significant errors for CID measurements of amino acids concerned Proline_2TMS
(overestimation of M4), Threonine_3TMS (overestimation of M3 at the expense of M0 and M1),
Valine_2TMS_C1C2 (overestimation of M2) and Valine_2TMS_C2C5 (overestimation of M3 at the
expense of M1 and M2). Considering the recent establishment of a [13]C-positional isotopomer approach
for Glutamate to analysis its C1 (Lima et al., 2021), we also analyzed the CID for the two low abundant
fragments Glutamate_3TMS_C1 (CO_2-TMS, M0 at m/z 117) and Glutamate_3TMS_C1C5 fragment
(CH3 loss, M0 at m/z 348). However, they both showed large and significant misestimations of their
CIDs, thus questioning the accuracy of this method (**Figure S1B**). The analysis of organic acids from
[13]C-PT full metabolites extract showed very good CID for Citrate_2TMS_C1C5, Malate_3TMS_C2C4
and Fumarate_2TMS_C1C4 (**Figure 2B**). Citrate_4TMS_C1C6 and Malate_3TMS_C1C4 showed
also relatively good CID but they had a higher variability compared to their respective C1C5 and C2C4
fragments. There was a very high overestimation of M0 in Succinate_2TMS_C1C4 at the expense of
M2, M3 and M4 isotopologues, However, these misestimations have been previously attributed to
contaminations inherent in the [13]C-PT full metabolites extract (Millard et al., 2014), thus preventing
the analytical validation for this fragment. For Pyruvate_MEOXTMS_C1C3, the variability was too
high to assess the CID measurements, perhaps reflecting its weak derivatization yield and low
abundance in [13]C-PT full metabolites extract (**Figure S1A**). To further assess the distribution of
measurements errors among isotopologues, we analyzed the CID accuracy, using threshold values of
±0.05 (**Figure 3A**). Overall, the bias of measurement for each isotopologue of each metabolite was
relatively low except for Methionine, Valine, Threonine, Succinate and Pyruvate. Interestingly, these
very high levels of accuracy for CID measurements were translated into very high levels of accuracy
for mean [13]C enrichment measurements, leading to very small shifts between expected (0.499 ±0.08)
and measured values (below 1% of error), except for Methionine, Valine, Threonine, Succinate and
Pyruvate   (**Figure   3B**).   However,   Malate_3TMS_C1C4,   Lysine_2TMS_C2C6   and
Serine_2TMS_C2C3 showed important variability for mean [13]C enrichment measurements although
no strong biases in their CID measurements were previously observed. Conversely, mean [13]C

enrichment of Phenylalanine_2TMS_C1C2 and proline_2TMS_C2C5 were slightly overestimated and
reflected the overestimation of their respective isotopologues M2 and M4. To ensure that the tested
fragments are suitable for isotopic analyses of metabolites from plant samples, we analyzed their
accuracy at natural ^{13}C abundance. For this purpose, we produced two types of plant samples (seedlings
from Arabidopsis and leaves from *Brassica napus*) and we measured the CID for each fragment of our
method to calculate the residual molecular ^{13}C enrichment (**Figure 3C**). This variable corresponded to
the ^{13}C enrichment remaining after removing the contribution of natural abundance for all isotopes (a
value of 0 is expected if no contamination occurs). Among the metabolites that were detected in the
plant samples, most of the fragments for organic and amino acids showed relatively high accuracy at
natural ^{13}C abundance, with most of the mean values of residual molecular ^{13}C enrichment ranging
between and 0.05 and 3% (**Figure 3C**). Interestingly, some discrepancies were observed between
Arabidopsis seedlings and *B. napus* leaves, notably for the full carbon backbone fragments
Citrate_4TMS_C1C6 and Malate_3TMS_C1C4, which were more accurately measured in *B. napus* at
low enrichment levels (critical values close to 5% for Arabidopsis). Conversely, Arabidopsis seedlings
showed slightly higher accuracies than *B. napus* leaves for AlphaAlanine_2TMS_C2C3,
Aspartate_2TMS_C2C4, Fumarate_2TMS_C1C4 and Serine_3TMS fragments. Phenylalanine_2TMS
fragments showed residual molecular ^{13}C enrichments close to 5% but only in plants samples.
Surprisingly, Proline_2TMS_C2C5, Threonine_3TMS_C2C4 and Valine_2TMS_C2C5 showed
relatively important residual molecular ^{13}C enrichments (nearly 10%) in both chemical standards and
unlabeled plant extracts, suggesting the occurrence of analytical bias. A detailed survey showed that
the isotopic cluster of Proline_2TMS_C2C5 was contaminated in the M4 isotopologue (m/z 146) by a
mass fragment deriving from near elution of Isoleucine_2TMS, and correlated with previous
measurements of CIDs ((**Figures 2, 3D**). Similarly, the isotopic cluster of Valine_2TMS_C2C5 was
contaminated in the M3 isotopologue (m/z 147) while Threonine_3TMS_C2C4 showed a surprising
very high abundance of the M1 isotopologue (40% of the CID) in both chemical standards and
unlabeled plant extracts.

Assessment of the light/dark regulation of plant TCA cycle using a ^{13}C-MFA approach

To further validate our MS method, we applied it to ^{13}C-labeled plant samples. For this purpose, we
performed transient kinetics of U-^{13}C-pyruvate incorporation to *B. napus* leaf discs subjected to either
light or dark conditions for up to 6 hours. The aim of this experiment was to obtain an overview of the
light/dark regulation of metabolic fluxes associated to TCA cycle and PEPc using MS methodology
and isotopologue-based approaches. We used U-^{13}C-pyruvate as a probe since the mitochondrial
Pyruvate DeHydrogenase (PDH) will be able to produce simultaneously $^{13}C_2$-Acetyl-CoA molecules
suitable for the investigation of the TCA cycle and $^{13}CO_2$ molecules suitable for the investigation of
PEPc activity through the combined action of carbonic anhydrase (H$^{13}CO_3^-$).

Following our experiments, the analysis of mean ^{13}C enrichment for all conditions revealed a
significant increase for α-Alanine, Aspartate, Citrate, Glutamate, Glycine, Malate, Pyruvate, Serine
and Succinate (**Figure 4A**). There was a very high biological variability and small ^{13}C enrichment for
Fumarate_2TMS_C1C4. Surprisingly, these enrichment dynamics remained relatively equivalent
between light and dark conditions for all the fragments considered, except for Glycine and Serine,
which were almost exclusively enriched in light conditions. Considering the importance of the light-
dependent photorespiratory pathway for glycine and serine production in leaves, this result clearly
showed that a significant part of $^{13}CO_2$ arising from PDH and TCA cycle decarboxylations (isocitrate
dehydrogenase and 2-oxoglutarate dehydrogenase) was reassimilated by photosynthesis. Overall,
Glycine_3TMS_C2, Serine_3TMS_C1C2 and Serine_3TMS_C2C3 reached a pseudo-isotopic steady-
state after 2 hours of illumination. Other fragments showed a relatively linear evolution of their mean

^{13}C enrichment, except for Pyruvate_MEOXTMS_C1C3 which rapidly reached an isotopic steady-
state after 0.5 hours only. Interestingly, labeling data from Pyruvate_MEOXTMS_C1C3 and
Succinate_2TMS_C1C4 fragments had a very good resolution (weak analytical and biological
variability) compared to previous evaluation with ^{13}C-PT full metabolites extract. Next, we performed
quantification of metabolite pool sizes along the experiment to determine whether the considered
metabolic network was in a metabolic steady-state (**Figure 4B**). For this purpose, we used GC-FID for
organic acids and HPLC-UV for amino acids using AccQTag methodology as they have a very high
level of stability for absolute quantification. Indeed, TMS-derivatives of amino acids are highly
unstable along a GC-MS run, leading to drastic differences of response when quantifying them with
external standards curves (Data not shown). In addition, the reliable absolute quantification of
metabolites using their CID masses can only be achieved by using metabolite-specific isotopic
standards (not our case in the present study) (Heuillet et al., 2020;Evers et al., 2021). While metabolite
pool sizes were relatively different between them, they all reached a remarkable pseudo steady-state
starting at 0-0.5 hours, except for Succinate and Serine where this state was reached at 2 hours.
Nevertheless, the range of pool size variation for these two metabolites was below +/- 2 µmol.g^{-1}DW.
Glycine was only quantified in light conditions (not detected in "dark condition" samples with HPLC-
UV). To summarize, we have found some similar ^{13}C enrichment dynamics for TCA cycle derived
metabolites in both light and dark conditions combined to similar metabolite pool sizes at a metabolic
steady-state.

Next, we focused our analysis on the isotopologue dynamics of these metabolites in order to depict the
considered metabolic network and compare metabolic fluxes associated to TCA cycle and PEPc
(**Figure 5**). Mean ^{13}C enrichment of Citrate, Glutamate and Succinate was essentially driven by an
increase of the M2 isotopologue for light and dark conditions, thus reflecting the incorporation of M2
Acetyl-CoA arising from U-^{13}C-pyruvate. However, there was also an increase of M1 and M3
isotopologues for Glutamate and Succinate after 2 hours while M1, M3 and M4 isotopologues of
Citrate_4TMS_C1C6 and Citrate_2TMS_C1C5 were increased starting from 4 hours. M1 and M4 of
Citrate_4TMS_C1C6 were essentially increased in dark conditions compared to light conditions after
4 hours. There was a very high variability for Fumarate_2TMS_C2C4 CID in both conditions, which
prevented its use for a reliable isotopic investigation. Interestingly, Malate_3TMS_C1C4 fragment
showed a higher enrichment of M1 compared to M2 in the light condition while M2 remains higher
than M1 in the dark condition. This shift may reflect an increase of the PEPc activity under light
condition (M1 arising from the reassimilation of the mitochondrial $^{13}CO_2$ decarboxylations). Since
there was no difference of mean ^{13}C enrichment between light and dark conditions for this fragment,
this result reinforced the importance of analyzing specifically CIDs for the interpretation of labeling
data. In addition, the CID of Malate_3TMS_C2C4 showed a similar dynamic for both light and dark
conditions, hence ruling out the use of this fragment to disentangle PEPc and TCA cycle contribution
to Malate production. Aspartate_3TMS_C2C4 showed a very high variability for the light condition,
but the CID dynamic in the dark condition was similar to the CID dynamic of Malate_3TMS_C2C4.
Thus, both fragments seemed to mimic Oxaloacetate labeling dynamics. Pyruvate_MEOXTMS_C1C3
was only enriched with the M3 isotopologue (nearly 80% M3/20% M0), thereby suggesting that the
activity of the malic enzyme, converting malate to pyruvate, was negligible in our metabolic network.

From these observations, we build a simplified metabolic network for TCA cycle composed of two
"turns" to explain the production of M1 and M3 in both Succinate_2TMS_C1C4 and
Glutamate_2TMS_C2C5 (**Figure 6A**). In this network, the second "turn" reflected the commitment of
a molecule that has already experienced all reactions of the TCA cycle to another round of reactions
(Orange and blue labels to compare acetyl-CoA-derived labeling from 1st and 2nd "turn"). Based on
previous works (Abadie and Tcherkez, 2019a), the reverse reaction of the fumarase was also considered

in the network, although it has no effect on the M1 isotopologue monitored by mass spectrometry (C4
vs C1 and C4 isotopomers of M1 isotopologue). The involvement of a light-dependent stored ^{12}C-
Citrate pool was not considered here since it only provided M0 isotopologues. From this metabolic
network, we proposed to evaluate PEPc-dependent and TCA cycle-dependent contributions to Citrate
production through M1, M2, M3 and M4 isotopologues and to Malate production through M1, M2 and
M3 isotopologues. These contributions were only calculated using timepoints 4 and 6 hours, because
all the fragments to consider were mostly visible at these moments. The main idea of this pathway
fractional contribution (FC) to a given metabolite production rely on the use of pathway-specific mass
isotopomer ratios for the metabolite, as long as the metabolic network is in a metabolic and isotopic
pseudo steady-state. Here, we observed a rapid metabolic steady-state and a relatively negligible
evolution of the relative proportions between the considered isotopologues (comparison of M1, M2
and M3 only for Malate for example) between 4 and 6 hours, *i.e.* the relative contribution of PEPc and
TCA cycle to either Malate and Citrate production remains in the same order of magnitude between
T4 and T6 hours. An important issue to address for calculating FC concerned the M1 from
Malate_2TMS_C1C4, which monitored at the same time PEPc and TCA cycle dependent isotopomers.
To deal with, we took advantage of Succinate_2TMS_C1C4 and Glutamate_2TMS_C1C4 fragments
which both harbored M1 and M3 isotopologues that are the precursor of TCA cycle-dependent M1 and
M3 of Malate_2TMS_C1C4 (**Figure 5**). Analysis of the M1/(M1+M3) ratio for these two fragments
showed a remarkable stability between dark and light conditions, thus reflecting a stable M2
enrichment of 40% for the active pool of acetyl-CoA (**Figure 6 B** and Equations). Using this ratio, we
calculated the FC of PEPc and TCA cycle for malate and citrate respectively (See Equations). We
found that TCA cycle had a significantly four-fold higher contribution to either Malate or Citrate
production compared to PEPc in all conditions (around 80% versus 20%). Interestingly, there was a
1.6-fold increase of the FC of PEPc to Malate production from dark to light conditions (14% versus
23%), although it remained not statistically significant (high variability, t-test p-value =0.08).
Conversely, PEPc contribution to Citrate production was significantly 1.5-fold higher in dark condition
than in light condition (26% versus 17%). Thus, the results suggested a higher commitment of PEPc-
derived carbons to the forward part of the TCA cycle in the dark, while commitment of PEPc-derived
carbons to the reverse part of the TCA cycle (Malate and perhaps Fumarate) was reduced at the same
time.

Finally, we explored the isotopologue distribution of Glycine and Serine, some amino acids which
were previously identified as a proxy for photosynthetic reassimilation of PDH and TCA cycle-derived
$^{13}CO_2$ (**Figure 7A**). Our results showed that the increase of mean ^{13}C enrichment for the three
considered fragments in light condition arisen from an increase of their M1 isotopologue. Given that
the light-dependent photorespiratory enzymes glycine decarboxylase and serine:hydroxymethyl
aminotransferase convert 2 glycine into 1 serine, we wanted to know whether the fragments
Serine_2TMS_C1C2 and Serine_2TMS_C2C3 were labeled in the same or different carbons, thus
revealing if the C1 of Glycine was also labeled (**Figure 7B**). By reviewing all the possible scenario
with different combinations of labeled glycine, we showed that the CID associated to our fragments
(isotopologue occurrence) could only be produced with one molecule of glycine labeled at the C2 and
an unlabeled molecule of glycine, leading to the production of a molecule of Serine labeled only at the
C2 and an unlabeled CO_2 molecule. Since we have shown that Glycine and Serine were both at the
isotopic and metabolic steady-state between 2 and 6 hours, we used Serine fragments to evaluate the
contribution of PDH and TCA cycle-derived CO_2 to the photosynthetic activity. For this purpose, the
steady-state enrichment of Acetyl-CoA calculated before was used as a proxy of the $^{12}C/^{13}C$ ratio of
the CO_2 produced by pyruvate dehydrogenase since a significant proportion of PDH-derived ^{12}C could
still contribute to photosynthesis (See equations), and given its major contributor for light respiration
(CO_2 emissions) (Tcherkez et al., 2005). After recalculating the mean ^{13}C-enrichment of SerineC1C3

from both Serine fragments, we showed that the light-dependent assimilation of PDH and TCA cycle
decarboxylations represented around 5% of photosynthesis (**Figure 7C**).

4   DISCUSSION

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of ^{13}C-INST-MFA. This approach
requires a high level of accuracy for CID measurements to avoid propagation of small measurement
errors to large errors in estimated fluxes (Antoniewicz et al., 2006). Considering the need to perform
high throughput measurements at isotopically non-stationary steady-state, MS has been preferentially
selected by the field in the recent years to measure CID in organic and amino acids. However, the
suitability of TMS-derivative molecular ions for these metabolites has never been evaluated at the
isotopologue level, *i.e.* by monitoring the impact of each isotopologue to the final CID measurement.
Conversely, a significant part of our knowledge about the functioning and the regulation of plant TCA
cycle is based on ^{13}C-labeling experiments with NMR detection of the ^{13}C signal. Thus, the use of a
validated MS methodology to investigate ^{13}C-labeling dynamics within plant TCA cycle will definitely
help to counterbalance our vision of the plant TCA cycle.

A robust methodology to evaluate CID measurements and identify analytical and matrix biases

In this study, we proposed a workflow to assess the measurement accuracy of CID from TMS-
derivatives of organic and amino acids by GC-MS for applications in plants, based on tailor-made *E.*
coli ^{13}C biological standards with controlled patterns of CID (^{13}C-PT) (**Figure 1**). From the TMS-
derivatives (selected to maximize information on carbon backbones), we showed that most of them
had a remarkable accuracy, leading to very small bias in the measurement of CIDs (below 5%) and
mean ^{13}C enrichments (below 1%). We further validated their suitability for application in plants by
monitoring very low residual molecular ^{13}C-enrichments in Arabidopsis seedlings and *Brassica napus*
leaves, although these accuracies could be variable between the two plant matrices
(Citrate_4TMS_C1C6, Malate_3TMS_C1C4, α-Alanine_2TMS_C2C3, Aspartate_2TMS_C2C4 and
Serine_3TMS). However, among amino acid TMS-derivatives, we identified three important biases at
the isotopologue level. The first bias concerned the contamination of the isotopic cluster of
Proline_2TMS_C2C5 at the m/z 146 (**Figure 3D**). A previous study reported this contaminant using
chemical standards at natural ^{13}C abundance, but no explanation was provided for this bias
(Antoniewicz et al., 2006). Here, we showed that the m/z 146 arisen from the co-elution with
Isoleucine, *i.e.* a separation by 0.02 seconds in our chromatograms, equivalent to 0.33 °C with the ramp
time for temperature during the elution (+10°C.min^{-1}). Nevertheless, this contamination only led to a
slight overestimation of the M4 isotopologue (^{13}C-PT samples), which did not strongly impact the
accuracy measurement for mean ^{13}C enrichment (**Figure 3B**). Thus, this fragment may still be suitable
for ^{13}C-MFA in plants, as long as the ^{13}C-labeling strategy for the metabolic network is adapted to
prevents the production of M4 isotopologues. The second bias identified concerned the contamination
of the isotopic cluster of Valine_2TMS_C2C5 at the m/z 147, which was observed in both chemical
standards and plant matrices. Interestingly, when a siliconium ion (loss of a methyl in a TMS group)
operates a cyclization by attacking another TMS group of the same molecule, it produces an ion
[(CH$_3$)$_3$Si-O-Si(CH$_3$)$_2$]$^+$ of the same mass 147 (Harvey and Vouros, 2020). We also detected a
significant abundance of an m/z 246 in the fragmentation spectra of Valine_2TMS, which seems to
support this hypothesis for the contamination (**Figure S2**). The third bias concerned the very high level
of M1 isotopologue for Threonine_3TMS_C2C4 detected in both chemical standards and plant
matrices. In its fragmentation spectra, we found an additional M1 isotopologue contamination for the
peak m/z 291, an important fragment for the identification of Threonine_3TMS by GC-MS (**Figure**

S2, NIST database). Given that we are working at a nominal mass resolution, we cannot determine
whether this contamination is due to: i) a co-elution of a contaminant chemically similar to Threonine;
ii) a strong isotopic effect during the derivatization. The use of GC coupled to high-resolution MS may
help to resolve this analytical issue (instrument not available at present in our platform). Besides this,
we observed a high level of M0 contamination for CID measurement of Succinate using the ^{13}C-PT
full metabolites extract, that has been reported previously (Millard et al., 2014), but prevented its
accurate evaluation with our workflow. Nevertheless, there was a small biological and analytical
variability for Succinate_2TMS_C1C4 in our ^{13}C-labeling experiments and we measured similar ratios
for M1 and M3 with Glutamate_2TMS_C2C5 that was already validated by our method (**Figure 6B**).
Since M1 and M3 were both impacted by the M0 contamination in ^{13}C-PT samples, our results suggest
that this fragment can be reliably used for ^{13}C-MFA.

Using our validated method, we provided important information about the role of to the activity of the
TCA cycle that contrasted with some proposed flux modes for TCA cycle in illuminated leaves
(Tcherkez et al., 2009;Gauthier et al., 2010;Sweetlove et al., 2010;Tcherkez et al., 2017;da Fonseca-
Pereira et al., 2021).

Stored Citrate is unlikely to contribute to TCA cycle activity

The hypothesis about the role of night-stored citrate molecules relies on two pioneer works. In the first
work, after 50 min of $^{13}CO_2$ incorporation in *Xanthium stomarium* leaves, ^{13}C signal was analyzed by
NMR and led the authors to propose "transmission coefficients" for metabolic reactions based on^{13}C
redistribution towards metabolites normalized to the $^{13}CO_2$ assimilated (Tcherkez et al., 2009). To
ensure the fitting with their experimental data, the authors had to introduce the contribution of an
important pool of unlabeled Citrate. The second work used the following experimental set-up ($^{13}CO_2$
6h light/$^{12}CO_2$ 6h dark/$^{12}CO_2$ 6h light, with $^{15}NH_4$$^{15}NO_3$ available all the time) with *B. napus* leaves
and a petiole-feeding assay with (Gauthier et al., 2010). Analysis of the results by NMR showed a weak
^{13}C-enrichment for Citrate, Glutamate, Glutamine and Malate at the end of the first light period and 30
min after the beginning of the dark period compared to the end of the experiment. Combined to a higher
occurrence of ^{12}C-^{15}N interactions compared to ^{13}C-^{15}N interactions in Glutamate after 6h of light with
$^{13}CO_2$, the results were interpreted as evidences for the contribution of night-stored vacuolar Citrate
and Malate molecules to TCA cycle activity in the subsequent light period. Here, using U-^{13}C-pyruvate
labeling experiment on *B. napus* with a leaf disc-feeding assay, we provided evidences for the absence
of Stored Citrate contribution to the TCA cycle in a light/dark dependent manner in our conditions
(**Figure 8**). Notably, we showed that the M2/(M0+M2) ratio of the Acetyl-CoA active pool was close
to 0.4 after 4 to 6 hours in our experiments, thereby showing that most of M0 isotopologues from
metabolite active pools arisen from M0 Acetyl-CoA and M0 OAA and not Citrate (**Figure 6B**). Given
the important production of PEPc-derived molecules with non-labeled PEP in our experiments, ^{12}C-
Pyruvate must be produced by the glycolysis and dilute the mitochondrial U-^{13}C-Pyruvate pool (nearly
20% of M0 detected in Pyruvate). It is also possible that thioredoxin-mediated activation of the ATP
citrate lyase activity in light conditions leads to the conversion of stored ^{12}C-Citrate pool into ^{12}C-OAA
and ^{12}C-Acetyl-CoA (Sweetlove et al., 2010;da Fonseca-Pereira et al., 2021). However, we showed
that the dynamic of ^{13}C-labeling for Citrate, Glutamate and Succinate was not diluted by light
conditions compared to dark conditions while the system was in a metabolic steady-state (**Figure 4**).
The remobilization of ^{12}C-stored Citrate in the light was expected to introduce a ^{12}C isotopic dilution
to Glutamate and Succinate compared to the dark, if this contribution was relatively important
(Gauthier et al., 2010), but this was not observed here. Interestingly, the ^{13}C-enrichment of Acetyl-
CoA was calculated from ^{13}C-labeled fragments of either Succinate or Glutamate (independent of ^{12}C-
molecules), thus this measurement cannot be biased with the contribution of stored ^{12}C-Citrate. Besides

this, we did not observe any significant modification of the Citrate pool during the 6 hours of our
experiment while a strong decrease in light conditions and a strong increase in dark conditions were
expected with the hypothesis of the stored citrate remobilization (**Figure 4B**). Since the previous
pioneer experiments performed with $^{13}CO_2$ and not U-^{13}C-pyruvate and used different methods for the
detection of ^{13}C, one might think that it is difficult to reconcile them. However, a ^{13}C-MFA performed
in Arabidopsis rosette leaves with $^{13}CO_2$ showed that Pyruvate was rapidly labeled and reached a
pseudo-isotopic state at nearly 40% of mean ^{13}C enrichment after 1 hour, similarly to the ^{13}C-
enrichment of Acetyl-CoA we calculated between 4 and 6 hours (Szecowka et al., 2013). In addition,
another ^{13}C-MFA performed in tobacco mature leaves showed that after 5 hours of $^{13}CO_2$
incorporation, only 20% of the starch produced was ^{13}C-labeled while ADP-Glucose pool was ^{13}C-
labeled by up to 88% (Chu et al., 2022). Thus, a high proportion of non-labeled carbon is still expected
to enter the glycolysis and the TCA cycle during the first hours of $^{13}CO_2$ incorporation, without
necessarily implying the contribution of stored Citrate. Given the similar dynamic of ^{13}C-enrichment
for Glutamate in both light and dark conditions in our experiments, the higher commitment of old ^{12}C
than photosynthetic ^{13}C towards Glutamate biosynthesis previously observed in the light seems to be
essentially attributed to both: i) an isotopic dilution of newly assimilated ^{13}C with pre-existing ^{12}C
metabolic pools; ii) a relatively slow turn-over of TCA cycle metabolic intermediates independently
of the light/dark condition. In addition, Glutamate is produced from 2-Oxoglutarate and Glutamine,
thus stored ^{12}C carbons are also incorporated at this step from Glutamine. Overall, our present findings
confirmed that stored Citrate is unlikely to contribute to plant TCA cycle in either light or dark
conditions (**Figure 8**). Since most of the Citrate and Malate pools are found in the vacuole
independently of the leaf developmental stage (Szecowka et al., 2013;Destailleur et al., 2021), their
remobilization may be expected only under stress conditions.

The forward cyclic flux mode of the plant TCA cycle majorly contributes to Malate biosynthesis

To date, non-cyclic nature of the TCA cycle in illuminated leaves has been proposed based on few
studies that used NMR to interpret ^{13}C signals in terms of ^{13}C-^{13}C interactions (Sweetlove et al.,
2010;Abadie et al., 2017). One branch is supposed to be fed essentially by PEPc (OAA, Malate and
Fumarate) while the other branch is only fed by (Citrate, 2-Oxoglutarate and Succinate) and the
conversion of Succinate to Fumarate is almost non-existent. In addition, it has also been proposed that
80% of Glutamate molecules in the light arisen from PEPc-derived carbons (Abadie et al., 2017).
However, one evidence is based on the absence of ^{13}C enrichment detected in the Succinate compared
to Fumarate with a 2-hours $^{13}C_1$-pyruvate incorporation while both metabolites showed similar ^{13}C-
enrichments with a 2-hours U-^{13}C-Pyruvate (Tcherkez et al., 2009). Unfortunately, the reassimilation
of $^{13}CO_2$ from $^{13}C_1$-Pyruvate decarboxylation by PEPc cannot be metabolically transmitted from
Malate to Succinate since the C1 and the C4 of the OAA backbone are lost during Citrate to Succinate
conversion (**Figure 6A**) (Alves et al., 2015). Another evidence is based on "transmission coefficient"
reflecting the ^{13}C-redistribution towards TCA cycle metabolites after 50 min of $^{13}CO_2$ labeling
(Tcherkez et al., 2009). The authors suggested that PEPc provided carbons essentially to OAA, Malate
and Fumarate, while stored Citrate supplied most of the carbon precursors for Glutamate (lower PDH
activity compared to PEPc). To assess the flux modes of the TCA Cycle, we took advantage of
pathway-specific mass isotopologues/isotopomers and a ^{13}C-MFA approach to investigate the TCA
cycle and PEPc metabolic fluxes after a U-^{13}C-Pyruvate incorporation in *B. napus* leaf discs. First, we
showed that Glutamate and Succinate rapidly harbored M2 istopologues, corresponding to *de novo*
incorporation of M2 Acetyl-CoA from the TCA Cycle, followed by an increase of M1 and M3
isotopologues over the time, corresponding to M2 isotopologues that had experienced a complete turn
of the forward cyclic mode of the TCA cycle (**Figure 5**). A brief comparison shows that the
(M1+M3)/M2 isotopologues ratio is close to 0.5 after 6 hours of labeling for both Glutamate and

Succinate (**Figure 5**). Since M2 isotopologues can be produced by both M0 OAA fixing M2 Acetyl-
CoA and M2 OAA fixing M0 acetyl-CoA, the commitment of M2 isotopologue to the forward TCA
cycle is necessarily higher than this ratio expected. Using a ^{13}C-MFA approach combined to pathway
specific MS fragments and the indirect measurement of M2/(M0+M2) ratio for the Acetyl-CoA active
pool, we showed that the forward part of the TCA cycle contributed to nearly 80% of both Malate and
Citrate production compared to PEPc (4:1 activity ratio), while malic enzyme activity was not
significantly detected (**Figure 6**). Interestingly, this modelling approach integrate the M0 istopologues
into the metabolic pools without the need to measure them, *i.e.* a potential contribution of any
additional ^{12}C pool can still be compatible. However, this pathway fractional contribution assumes that
most of the CO_2 is produced by the PDH, and its conversion to HCO_3^- are not rate-limiting for PEPc
activity. While the second hypothesis may not be relevant given that CO_2 was actively reassimilated
by photosynthesis in our light conditions (**Figure 7**), the second hypothesis can be questioned given
the metabolic network (**Figure 6**). Indeed, isocitrate and 2-oxoglutarate dehydrogenases necessarily
decarboxylate more $^{12}CO_2$ than $^{13}CO_2$ compared to the PDH, which may imbalance the reassimilation
of $^{13}CO_2$ by PEPc if the first part of the TCA cycle operates at a high rate. Assuming an equivalent
metabolic flux from PDH to 2oxoglutarate dehydrogenase, the overall mitochondrial $^{12}CO_2/^{13}CO_2$ ratio
should not exceed a 2:1 dilution compared to the $^{12}CO_2/^{13}CO_2$ produced by the PDH, based on the
isotopologue distribution of Citrate, Glutamate and Succinate and their associated CO_2 losses. Thus,
the forward and cyclic mode of the TCA cycle will still majorly contribute to Malate and Citrate
compared to PEPc (2:1 ratio with this scenario) (**Figure 8**). Interestingly, a ^{13}C-MFA performed at
isotopic and metabolic steady-state in heterotrophic suspension cells culture under non-limiting
glucose concentrations showed nearly similar ratio of 2:1 for forward TCA cycle/PEPc metabolic
fluxes, with a negligible malic enzyme activity (10-fold less than PEPc) (Masakapalli et al., 2010).
Considering Arabidopsis leaves, a ^{13}C-MFA was performed recently using a 1-hour kinetic of $^{13}CO_2$
incorporation and suggested that PEPc activity was 1.5 fold higher than the TCA cycle flux, with
fumarase and malate dehydrogenase being inactive (value below 0.1 compared to PEPc at 1.2 µmol.g$^-$
^{1} FW.h^{-1}) (Xu et al., 2021). However, such conclusions were drawn with questionable modelling, *i.e.*
M1, M2 and M3 isotopologues of Citrate were already present at high levels when starting the
experiment and metabolic flux variations were only recorded as the production of M4 isotopologue
(unrealistic metabolic network compared to other work (Szecowka et al., 2013)). Based on ^{13}C-
positionnaly accumulated in Malate after a $^{13}CO_2$ incorporation, another study proposed an important
flux for PEPc at high photorespiration rates in sunflower (Abadie and Tcherkez, 2019a), a condition
which may happen in our experiments using leaf disc floating on a solution. However, the conclusion
for PEPc flux were also drawn with questionable modeling approach: i) calculations were made without
the dynamic of ^{13}C-enrichment for both the substrate and the product (mandatory (Buescher et al.,
2015;Millard et al., 2020) ; ii) the estimated metabolic fluxes associated to the use of HCO_3^- or PEP
were largely different while both metabolites are the substrates of the same reaction (same metabolic
flux); iii) the metabolic steady-state of the system was not considered (accumulation of Malate is
unlikely to occur). To date, the accurate estimation of absolute PEPc metabolic flux remains
challenging in plants, as it is difficult to routinely get access to ^{13}C-enrichments for HCO_3^- and PEP by
either NMR or GC-MS. Here, we showed that the anaplerotic contribution of PEPc to the mitochondrial
metabolism remains lower than the contribution of the TCA cycle, perhaps reflecting the constraints
imposed by PEPc-derived Malate transport from cytosol to mitochondria.

Besides this, the forward cyclic mode of the TCA cycle has to be reconnected to the relative ^{13}C content
found in Glutamate and Aspartate, which can represent important escapes for ^{13}C that will drive the
relative differences of metabolic flux for the biosynthesis of Glutamate and Succinate, and Malate and
OAA respectively (branching points) (**Figure 5**). However, a significant proportion of Glutamate and
Aspartate may come back to the TCA cycle. In astrocyte cells, U-^{13}C-Glutamate labeling

incorporations showed that Glutamate M5 was rapidly metabolized into Glutamate M3 after one turn
of the TCA cycle (Nissen et al., 2015) thereby proposing a similar role for plant mitochondrial
Glutamate dehydrogenase (Terce-Laforgue et al., 2013). The potential role of GABA (Gamma-
aminobutyric acid) shunt has been proposed to reroute Glutamate towards Succinate (Tcherkez et al.,
2009). Finally, mitochondrial transminases such as Alanine:2-oxoglutarate and Aspartate:2-
oxoglutarate could also participle in this process. Since Glutamate is necessarily produced from ^{13}C-2-
Oxoglutarate and ^{12}C-Glutamine molecules in our metabolic network, Glutamate rerouting pathways
are supposed to introduce a 50% isotopic dilution between ^{12}C and ^{13}C molecules for mitochondrial
active pools of either Succinate or 2-Oxoglutarate. But it is not possible for instance to have access to
the size of the mitochondrial active pool for both metabolites and by extension their exact ^{13}C-
enrichments without a complete isotopic steady-state of the plant TCA cycle. Given the fractional
contributions of PEPc and TCA cycle calculated in this work for Malate and Citrate, a higher metabolic
flux for PDH (^{13}C-Acetyl-CoA) than 2-Oxoglutarate dehydrogenase (^{13}C-Succinate) is expected and
necessarily constraints the PEPc flux to Malate (**Figure 8**). To which extent Glutamate and Aspartate
molecules escape the TCA cycle remains to be determined. Overall, this forward cyclic flux mode of
the TCA cycle in both light and dark conditions strongly supports its key roles in *de novo* nitrogen
assimilation and mitochondrial ATP production (through Succinate dehydrogenase activity)
(Sweetlove et al., 2010).

Light/dark regulation of PEPc and TCA cycle metabolic fluxes: the potential roles of glycolysis
and Malate import into mitochondria

We showed that Malate_2TMS_C1C4 fragment experienced a strong shift in its M1/M2 isotopologues
in light conditions compared to dark conditions, which was associated to a relatively higher
contribution of PEPc to Malate production in light condition than in dark condition. Such shift could
be attributed to both increased PEPc activity and TCA cycle inhibition in the light compared to dark
conditions. Interestingly, PEPc activity is positively regulated by its light-dependent thioredoxin-
mediated phosphorylation in C_3 and C_4 plants (Caburatan and Park, 2021). In the light, the lower
respiration rate measured compared to dark conditions has been partly attributed to an inhibition of
PDH and TCA cycle-derived decarboxylations (Tcherkez et al., 2005;Xu et al., 2021). In our
experiments, it was not possible to compare absolute fluxes of TCA cycle in light and dark conditions,
due to the relative nature of the fractional contribution measurements (total of 100% in each condition).
However, we found very similar dynamics of ^{13}C-enrichment for TCA cycle metabolites combined to
similar metabolic pool sizes in a metabolic steady-state (**Figure 5**). Thus, variations of TCA cycle
metabolic fluxes between light and dark conditions are expected to be very low in our experimental
set-up, and subsequently the variation of light/dark respiration. It is tempting to think that our
experimental set-up badly reflects plant physiology and its light/dark associated regulation (leaf discs
floating on a solution). However, by integrating a ^{13}C-positionnal isotopomer approach to the
photorespiratory metabolism, we showed that PDH and TCA-cycle derived CO_2 were actively
reassimilated by the photosynthesis, leading to a contribution of around 5% to net CO_2 assimilation
(**Figure 7**). Under the light irradiance of around 120 µmol photons.m^{-2}.s^{-1}, net CO_2 uptake of *B. napus*
leaves at a similar developmental stage (L11-L7) is close to 5 µmol CO_2. m^{-2}.s^{-1} (Dellero et al., 2021),
leading to a reassimilation rate of 0.25 µmol CO_2. m^{-2}.s^{-1}. Since the dark respiration rate for these
leaves is close to 1.25 µmol CO_2. m^{-2}.s^{-1} (Dellero et al., 2021), and given the proposed inhibition of
day respiration by nearly 50% in plants (Tcherkez et al., 2005), this means that the current
measurement/estimation of day respiration rate through CO_2 mass balance at the leaf level (Tcherkez
et al., 2017;Berghuijs et al., 2019) is still underestimated by nearly 40% (photosynthetic reassimilation
not taken into account because not evaluated). Therefore, the inhibition of day respiration at the
metabolic flux level is likely be lower than measured by gas-exchanges (**Figure 8**). For instance, the

very similar dynamics of ^{13}C-enrichment observed in our experiments for light and dark conditions can
be explained by two major hypotheses: i) a relatively higher control of TCA cycle activity by glycolysis
rather than by pyruvate dehydrogenase, given that starch degradation and glycolysis are essentially
inhibited by light conditions (Plaxton and Podestá, 2006;Thalmann and Santelia, 2017); ii) a specific
activation of pyruvate dehydrogenase in both conditions, given that we incorporated a strong amount
of U-^{13}C-pyruvate into leaf discs (Tovar-Mendez et al., 2003). Nevertheless, the shift in the fractional
contribution of PEPc to malate production between dark and light conditions still confirms the
robustness of our method to infer reliable information about the light/dark regulation of metabolic
fluxes. Besides this, we observed another unexpected but interesting result: the higher contribution of
PEPc derived-carbons to Citrate production in the dark condition compared to the light condition. Since
we have shown that PEPc contribution to Malate production was reduced in dark compared to light
conditions, this result can only be explained by a higher transport of OAA/Malate from the cytosol to
the mitochondria in dark conditions (**Figure 8**). This idea is supported by the fact that PEPc is majorly
localized to the cytosol (Caburatan and Park, 2021) and the recent identification of a mitochondrial
malate importer in Arabidopsis with a decisive impact on mitochondrial respiration (Lee et al., 2021).
Whether a higher malate import into mitochondria allows to increase the absolute metabolic flux of the
TCA cycle in dark conditions in our experimental set-up remains to be elucidated.

Conclusions

Overall, we proposed a robust workflow to evaluate the accuracy of CID measurements by GC-MS for
organic and amino acids and successfully applied our validated method to the investigation of plant
TCA cycle and PEPc metabolic fluxes. Notably, our results do not support the hypothesis of the stored
citrate contribution to TCA cycle and demonstrated that the forward and cyclic flux mode of the TCA
cycle largely contributed to Malate and Citrate production compared to PEPc (4:1 ratio). Thus, the
activity of the succinate dehydrogenase in both dark and light conditions has been largely
underestimated until now. Reassimilation of PDH and TCA-cycle derived CO_2 by photosynthesis
represented nearly 5% of the net CO_2 assimilation rate in our conditions, and this should be considered
when day respiration rate is measured by CO_2 mass balance at the leaf level. Although the regulation
of the PEPc activity by light was still operating, the higher involvement of PEPc-derived carbons to
citrate production in dark conditions proposed an important role for Malate import into mitochondria
in sustaining TCA cycle in dark conditions. The next stage of this work should be focused on the
accurate modeling of this metabolic network to infer absolute metabolic fluxes. This perspective
sounds stimulating given the development of multiple modeling approaches over the last decade but
still remains highly challenging (Kruger and Ratcliffe, 2021).

5   Conflict of Interest

794 The authors declare that the research was conducted in the absence of any commercial or financial
795 relationships that could be construed as a potential conflict of interest.

6   Author Contributions

797 Conceptualization, Y.D., A.B.; methodology, Y.D..; software, Y.D., O.F.; validation, Y.D., S.B. and
798 C.B.; formal analysis, Y.D.; investigation, Y.D., S.B. and C.B.; data curation, Y.D.; writing-original
799 draft preparation, Y.D.; writing-review and editing, Y.D., A.B.; visualization, Y.D.; supervision,
800 Y.D.; project administration, Y.D.; funding acquisition, Y.D., A.B. All authors have read and agreed
801 to the published version of the manuscript.

7   Funding

803 This research was supported by the IB2021_TRICYCLE project funded by the INRAE BAP division
804 and by the METABOHUB project funded by the program Investments for the Future of the French
805 national agency for research (ANR grant number 11-INBS-0010).

8   Acknowledgments

P2M2 (Metabolomics & Metabolic Profiling Platform, Le Rheu, France, www6.inrae.fr/p2m2/) and its
staff members are gratefully acknowledged for access to mass spectrometry facilities and for
supporting the methodological developments required for this study. We thank MetaToul
(Metabolomics & Fluxomics Facitilies, Toulouse, France, www.metatoul.fr) for technical support and
access to NMR facilities. We are grateful to Nathalie Marnet and Mathieu Aubert for technical
assistance in the quantification of amino acid content and Pierre Millard and Floriant Bellvert for
fruitful scientific discussions.

9   Supplementary Material

815 Supplementary Material is attached to the preprint.

10   References

Abadie, C., Bathellier, C., and Tcherkez, G. (2018). Carbon allocation to major metabolites in
illuminated leaves is not just proportional to photosynthesis when gaseous conditions (CO2
and O2) vary. *New Phytol* 218**,** 94-106.

Abadie, C., Boex-Fontvieille, E.R., Carroll, A.J., and Tcherkez, G. (2016). In vivo stoichiometry of
photorespiratory metabolism. *Nat Plants* 2**,** 15220.

Abadie, C., Lothier, J., Boex-Fontvieille, E., Carroll, A., and Tcherkez, G. (2017). Direct assessment
of the metabolic origin of carbon atoms in glutamate from illuminated leaves using (13) C-
NMR. *New Phytol* 216**,** 1079-1089.

Abadie, C., and Tcherkez, G. (2019a). In vivo phosphoenolpyruvate carboxylase activity is
controlled by CO2 and O2 mole fractions and represents a major flux at high photorespiration
rates. *New Phytol* 221**,** 1843-1852.

Abadie, C., and Tcherkez, G. (2019b). Plant sulphur metabolism is stimulated by photorespiration.
Commun Biol 2**,** 379.

Abadie, C., and Tcherkez, G. (2021). (13)C Isotope Labelling to Follow the Flux of Photorespiratory
Intermediates. *Plants (Basel)* 10.

Allen, D.K., and Young, J.D. (2019). Tracing metabolic flux through time and space with isotope
labeling experiments. *Curr Opin Biotechnol* 64**,** 92-100.

Alves, T.C., Pongratz, R.L., Zhao, X., Yarborough, O., Sereda, S., Shirihai, O., Cline, G.W., Mason,
G., and Kibbey, R.G. (2015). Integrated, Step-Wise, Mass-Isotopomeric Flux Analysis of the
TCA Cycle. *Cell Metab* 22**,** 936-947.

Antoniewicz, M.R. (2021). A guide to metabolic flux analysis in metabolic engineering: Methods,
tools and applications. *Metab Eng* 63**,** 2-12.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2006). Determination of confidence
intervals of metabolic fluxes estimated from stable isotope measurements. *Metab Eng* 8**,** 324-
337.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2007). Accurate Assessment of Amino
Acid Mass Isotopomer Distributions for Metabolic Flux Analysis. *Anal Chem* 79**,** 6.

Araujo, W.L., Tohge, T., Nunes Nesi, A., Obata, T., and Fernie, A. (2014). "Analysis of Kinetic
Labeling of Amino Acids and Organic Acids by GC-MS," in *Plant Metabolic Flux Analysis:*
Methods and Protocols, ed. M.I.M. Biology.).

Berghuijs, H.N.C., Yin, X., Ho, Q.T., Retta, M.A., Nicolai, B.M., and Struik, P.C. (2019). Using a
reaction-diffusion model to estimate day respiration and reassimilation of (photo)respired
CO2 in leaves. *New Phytol* 223**,** 619-631.

Bianchetti, G., Baron, C., Carrillo, A., Berardocco, S., Marnet, N., Wagner, M.H., Demilly, D.,
Ducournau, S., Manzanares-Dauleux, M.J., Caherec, F.L., Buitink, J., and Nesi, N. (2021).
Dataset for the metabolic and physiological characterization of seeds from oilseed rape
(Brassica napus L.) plants grown under single or combined effects of drought and clubroot
pathogen Plasmodiophora brassicae. *Data Brief* 37**,** 107247.

Buescher, J.M., Antoniewicz, M.R., Boros, L.G., Burgess, S.C., Brunengraber, H., Clish, C.B.,
Deberardinis, R.J., Feron, O., Frezza, C., Ghesquiere, B., Gottlieb, E., Hiller, K., Jones, R.G.,
Kamphorst, J.J., Kibbey, R.G., Kimmelman, A.C., Locasale, J.W., Lunt, S.Y., Maddocks,
O.D., Malloy, C., Metallo, C.M., Meuillet, E.J., Munger, J., Noh, K., Rabinowitz, J.D.,
Ralser, M., Sauer, U., Stephanopoulos, G., St-Pierre, J., Tennant, D.A., Wittmann, C., Vander
Heiden, M.G., Vazquez, A., Vousden, K., Young, J.D., Zamboni, N., and Fendt, S.M. (2015).
A roadmap for interpreting (13)C metabolite labeling patterns from cells. *Curr Opin*
Biotechnol 34**,** 189-201.

Caburatan, L., and Park, J. (2021). Differential Expression, Tissue-Specific Distribution, and
Posttranslational Controls of Phosphoenolpyruvate Carboxylase. *Plants (Basel)* 10.

Cheah, Y.E., Xu, Y., Sacco, S.A., Babele, P.K., Zheng, A.O., Johnson, C.H., and Young, J.D. (2020).
Systematic identification and elimination of flux bottlenecks in the aldehyde production
pathway of Synechococcus elongatus PCC 7942. *Metab Eng* 60**,** 56-65.

Cheah, Y.E., and Young, J.D. (2018). Isotopically nonstationary metabolic flux analysis (INST-
MFA): putting theory into practice. *Curr Opin Biotechnol* 54**,** 80-87.

Chu, K.L., Koley, S., Jenkins, L.M., Bailey, S.R., Kambhampati, S., Foley, K., Arp, J.J., Morley,
S.A., Czymmek, K.J., Bates, P.D., and Allen, D.K. (2022). Metabolic flux analysis of the
non-transitory starch tradeoff for lipid production in mature tobacco leaves. *Metab Eng* 69**,**
231-248.

Clark, T.J., Guo, L., Morgan, J., and Schwender, J. (2020). Modeling Plant Metabolism: From
Network Reconstruction to Mechanistic Models. *Annu Rev Plant Biol*.

Da Fonseca-Pereira, P., Souza, P.V.L., Fernie, A.R., Timm, S., Daloso, D.M., and Araujo, W.L.
(2021). Thioredoxin-mediated regulation of (photo)respiration and central metabolism. *J Exp*
Bot 72**,** 5987-6002.

Dellero, Y. (2020). Manipulating amino acid metabolism to improve crop nitrogen use efficiency for
a sustainable agriculture. *Frontiers in Plant Science* 11**,** 1857

Dellero, Y., Clouet, V., Marnet, N., Pellizzaro, A., Dechaumet, S., Niogret, M.F., and Bouchereau,
A. (2020a). Leaf status and environmental signals jointly regulate proline metabolism in
winter oilseed rape. *J Exp Bot* 71**,** 2098-2111.

Dellero, Y., and Filangi, O. (2021). "Corrective method dedicated to Isocor for calculating carbon
isotopologue distribution from GCMS runs". V1 ed.: Portail Data INRAE).

Dellero, Y., Heuillet, M., Marnet, N., Bellvert, F., Millard, P., and Bouchereau, A. (2020b).
Sink/Source Balance of Leaves Influences Amino Acid Pools and Their Associated Metabolic
Fluxes in Winter Oilseed Rape (Brassica napus L.). *Metabolites* 10**,** 16.

Dellero, Y., Jossier, M., Bouchereau, A., Hodges, M., and Leport, L. (2021). Leaf Phenological
Stages of Winter Oilseed Rape (Brassica napus L.) Have Conserved Photosynthetic
Efficiencies but Contrasted Intrinsic Water Use Efficiencies at High Light Intensities.
Frontiers in Plant Science 12.

Dellero, Y., Jossier, M., Glab, N., Oury, C., Tcherkez, G., and Hodges, M. (2016). Decreased
glycolate oxidase activity leads to altered carbon allocation and leaf senescence after a
transfer from high CO2 to ambient air in Arabidopsis thaliana. *J Exp Bot* 67**,** 3149-3163.

Destailleur, A., Poucet, T., Cabasson, C., Alonso, A.P., Cocuron, J.C., Larbat, R., Vercambre, G.,
Colombie, S., Petriacq, P., Andrieu, M.H., Beauvoit, B., Gibon, Y., and Dieuaide-Noubhani,
M. (2021). The Evolution of Leaf Function during Development Is Reflected in Profound
Changes in the Metabolic Composition of the Vacuole. *Metabolites* 11.

Evers, B., Gerding, A., Boer, T., Heiner-Fokkema, M.R., Jalving, M., Wahl, S.A., Reijngoud, D.J.,
and Bakker, B.M. (2021). Simultaneous Quantification of the Concentration and Carbon
Isotopologue Distribution of Polar Metabolites in a Single Analysis by Gas Chromatography
and Mass Spectrometry. *Anal Chem* 93**,** 8248-8256.

Galili, G., Amir, R., and Fernie, A.R. (2016). The Regulation of Essential Amino Acid Synthesis and
Accumulation in Plants. *Annu Rev Plant Biol* 67**,** 153-178.

Gauthier, P.P., Bligny, R., Gout, E., Mahe, A., Nogues, S., Hodges, M., and Tcherkez, G.G. (2010).
In folio isotopic tracing demonstrates that nitrogen assimilation into glutamate is mostly
independent from current CO2 assimilation in illuminated leaves of Brassica napus. *New*
Phytol 185**,** 988-999.

Harvey, D.J., and Vouros, P. (2020). Mass Spectrometric Fragmentation of Trimethylsilyl and
Related Alkylsilyl Derivatives. *Mass Spectrom Rev* 39**,** 105-211.

Hay, J.O., Shi, H., Heinzel, N., Hebbelmann, I., Rolletschek, H., and Schwender, J. (2014).
Integration of a constraint-based metabolic model of Brassica napus developing seeds with
(13)C-metabolic flux analysis. *Front Plant Sci* 5**,** 724.

Heise, R., Fernie, A.R., Stitt, M., and Nikoloski, Z. (2015). Pool size measurements facilitate the
determination of fluxes at branching points in non-stationary metabolic flux analysis: the case
of Arabidopsis thaliana. *Front Plant Sci* 6**,** 386.

Heuillet, M., Bellvert, F., Cahoreau, E., Letisse, F., Millard, P., and Portais, J.C. (2018).
Methodology for the Validation of Isotopic Analyses by Mass Spectrometry in Stable-Isotope
Labeling Experiments. *Anal Chem* 90**,** 1852-1860.

Heuillet, M., Millard, P., Cisse, M.Y., Linares, L.K., Letisse, F., Manie, S., Le Cam, L., Portais, J.C.,
and Bellvert, F. (2020). Simultaneous Measurement of Metabolite Concentration and Isotope
Incorporation by Mass Spectrometry. *Anal Chem*.

Kambhampati, S., Li, J., Evans, B.S., and Allen, D.K. (2019). Accurate and efficient amino acid
analysis for protein quantification using hydrophilic interaction chromatography coupled
tandem mass spectrometry. *Plant Methods* 15**,** 46.

Kruger, N.J., Huddleston, J.E., Le Lay, P., Brown, N.D., and Ratcliffe, R.G. (2007). Network flux
analysis: impact of 13C-substrates on metabolism in Arabidopsis thaliana cell suspension
cultures. *Phytochemistry* 68**,** 2176-2188.

Kruger, N.J., and Ratcliffe, R.G. (2015). Fluxes through plant metabolic networks: measurements,
predictions, insights and challenges. *Biochem J* 465**,** 27-38.

Kruger, N.J., and Ratcliffe, R.G. (2021). Whither metabolic flux analysis in plants? *Journal of*
Experimental Botany.

Lai, Z., and Fiehn, O. (2018). Mass spectral fragmentation of trimethylsilylated small molecules.
Mass Spectrom Rev 37**,** 245-257.

Lee, C.P., Elsasser, M., Fuchs, P., Fenske, R., Schwarzlander, M., and Millar, A.H. (2021). The
versatility of plant organic acid metabolism in leaves is underpinned by mitochondrial malate-
citrate exchange. *Plant Cell* 33**,** 3700-3720.

Lima, V.F., Erban, A., Daubermann, A.G., Freire, F.B.S., Porto, N.P., Candido-Sobrinho, S.A.,
Medeiros, D.B., Schwarzlander, M., Fernie, A.R., Dos Anjos, L., Kopka, J., and Daloso,
D.M. (2021). Establishment of a GC-MS-based (13) C-positional isotopomer approach
suitable for investigating metabolic fluxes in plant primary metabolism. *Plant J*.

Lima, V.F., Perez Souza, L., Williams, T.C., Fernie, A., and Daloso, D.M. (2018). "Gas
Chromatography–Mass Spectrometry-Based 13C-Labeling Studies in Plant Metabolomics,"
in *Plant Metabolomics: Methods and Protocol,* ed. M.I.M. Biology.).

Ma, F., Jazmin, L.J., Young, J.D., and Allen, D.K. (2014). Isotopically nonstationary 13C flux
analysis of changes in Arabidopsis thaliana leaf metabolism due to high light acclimation.
Proc Natl Acad Sci U S A 111**,** 16967-16972.

Masakapalli, S.K., Le Lay, P., Huddleston, J.E., Pollock, N.L., Kruger, N.J., and Ratcliffe, R.G.
(2010). Subcellular flux analysis of central metabolism in a heterotrophic Arabidopsis cell
suspension using steady-state stable isotope labeling. *Plant Physiol* 152**,** 602-619.

Medeiros, D.B., Perez Souza, L., Antunes, W.C., Araujo, W.L., Daloso, D.M., and Fernie, A.R.
(2018). Sucrose breakdown within guard cells provides substrates for glycolysis and
glutamine biosynthesis during light-induced stomatal opening. *Plant J* 94**,** 583-594.

Millard, P., Cahoreau, E., Heuillet, M., Portais, J.C., and Lippens, G. (2017). (15)N-NMR-Based
Approach for Amino Acids-Based (13)C-Metabolic Flux Analysis of Metabolism. *Anal Chem*
89**,** 2101-2106.

Millard, P., Delepine, B., Guionnet, M., Heuillet, M., Bellvert, F., and Letisse, F. (2019). IsoCor:
isotope correction for high-resolution MS labeling experiments. *Bioinformatics* 35**,** 4484-
4487.

Millard, P., Massou, S., Portais, J.C., and Letisse, F. (2014). Isotopic studies of metabolic systems by
mass spectrometry: using Pascal's triangle to produce biological standards with fully
controlled labeling patterns. *Anal Chem* 86**,** 10288-10295.

Millard, P., Schmitt, U., Kiefer, P., Vorholt, J.A., Heux, S., and Portais, J.-C. (2020). ScalaFlux: a
scalable approach to quantify fluxes in metabolic subnetworks. *Plos Computational Biology*
16**,** e1007799.

Nissen, J.D., Pajecka, K., Stridh, M.H., Skytt, D.M., and Waagepetersen, H.S. (2015). Dysfunctional
TCA-Cycle Metabolism in Glutamate Dehydrogenase Deficient Astrocytes. *Glia* 63**,** 2313-
2326.

Okahashi, N., Kawana, S., Iida, J., Shimizu, H., and Matsuda, F. (2019). Fragmentation of
Dicarboxylic and Tricarboxylic Acids in the Krebs Cycle Using GC-EI-MS and GC-EI-
MS/MS. *Mass Spectrom (Tokyo)* 8**,** A0073.

Plaxton, W.C., and Podestá, F.E. (2006). The Functional Organization and Control of Plant
Respiration. *Critical Reviews in Plant Sciences* 25**,** 159-198.

Ratcliffe, R.G., and Shachar-Hill, Y. (2006). Measuring multiple fluxes through plant metabolic
networks. *Plant J* 45**,** 490-511.

Rstudio Team (2016). RStudio: Integrated Development for R.

.

Sinnaeve, D., Dinclaux, M., Cahoreau, E., Millard, P., Portais, J.C., Letisse, F., and Lippens, G.
(2018). Improved Isotopic Profiling by Pure Shift Heteronuclear 2D J-Resolved NMR
Spectroscopy. *Anal Chem* 90**,** 4025-4031.

Sweetlove, L.J., Beard, K.F., Nunes-Nesi, A., Fernie, A.R., and Ratcliffe, R.G. (2010). Not just a
circle: flux modes in the plant TCA cycle. *Trends Plant Sci* 15**,** 462-470.

Szecowka, M., Heise, R., Tohge, T., Nunes-Nesi, A., Vosloh, D., Huege, J., Feil, R., Lunn, J.,
Nikoloski, Z., Stitt, M., Fernie, A.R., and Arrivault, S. (2013). Metabolic fluxes in an
illuminated Arabidopsis rosette. *Plant Cell* 25**,** 694-714.

Tcherkez, G., Cornic, G., Bligny, R., Gout, E., and Ghashghaie, J. (2005). In vivo respiratory
metabolism of illuminated leaves. *Plant Physiol* 138**,** 1596-1606.

Tcherkez, G., Gauthier, P., Buckley, T.N., Busch, F.A., Barbour, M.M., Bruhn, D., Heskel, M.A.,
Gong, X.Y., Crous, K.Y., Griffin, K., Way, D., Turnbull, M., Adams, M.A., Atkin, O.K.,
Farquhar, G.D., and Cornic, G. (2017). Leaf day respiration: low CO2 flux but high
significance for metabolism and carbon balance. *New Phytol* 216**,** 986-1001.

Tcherkez, G., Mahe, A., Gauthier, P., Mauve, C., Gout, E., Bligny, R., Cornic, G., and Hodges, M.
(2009). In folio respiratory fluxomics revealed by 13C isotopic labeling and H/D isotope
effects highlight the noncyclic nature of the tricarboxylic acid "cycle" in illuminated leaves.
Plant Physiol 151**,** 620-630.

Terce-Laforgue, T., Bedu, M., Dargel-Grafin, C., Dubois, F., Gibon, Y., Restivo, F.M., and Hirel, B.
(2013). Resolving the role of plant glutamate dehydrogenase: II. Physiological
characterization of plants overexpressing the two enzyme subunits individually or
simultaneously. *Plant Cell Physiol* 54**,** 1635-1647.

Thalmann, M., and Santelia, D. (2017). Starch as a determinant of plant fitness under abiotic stress.
New Phytol 214**,** 943-951.

Timm, S. (2020). The impact of photorespiration on plant primary metabolism through metabolic and
redox regulation. *Biochemical Society Transactions* 48**,** 2495-2504.

Tovar-Mendez, A., Miernyk, J.A., and Randall, D.D. (2003). Regulation of pyruvate dehydrogenase
complex activity in plant cells. *Eur J Biochem* 270**,** 1043-1049.

Treves, H., Kuken, A., Arrivault, S., Ishihara, H., Hoppe, I., Erban, A., Hohne, M., Moraes, T.A.,
Kopka, J., Szymanski, J., Nikoloski, Z., and Stitt, M. (2022). Carbon flux through
photosynthesis and central carbon metabolism show distinct patterns between algae, C3 and
C4 plants. *Nat Plants* 8**,** 78-91.

Wieloch, T. (2021). The next phase in the development of 13C isotopically nonstationary metabolic
flux analysis. *J Exp Bot*.

Xu, G., Fan, X., and Miller, A.J. (2012). Plant nitrogen assimilation and use efficiency. *Annu Rev
Plant Biol* 63**,** 153-182.

Xu, Y., Fu, X., Sharkey, T.D., Shachar-Hill, Y., and Walker, B.J. (2021). The metabolic origins of
non-photorespiratory CO2 release during photosynthesis: A metabolic flux analysis. *Plant
Physiol*.

Young, J.D., Allen, D.K., and Morgan, J.A. (2014). "Isotopomer measurement techniques in
metabolic flux analysis II: mass spectrometry," in *Methods Mol Biol*.), 85-108.

Zhang, Y., and Fernie, A.R. (2018). On the role of the tricarboxylic acid cycle in plant productivity. *J
Integr Plant Biol* 60**,** 1199-1216.

1022

1023

1024

1025

Table 1. Molecular ions selected for Carbon Isotopologue Distribution analysis of organic and amino acid TMS-derivatives by GC-MS.

Compounds	TMS derivative		Loss		Molecular ion		
	Formula	mass	Formula	mass	Formula	m/z	Carbon skeleton
Organic acids							
Citrate(4TMS)	C18H40O7Si4	480	CH3	15	C17H37O7Si4	465	1-2-3-4-5-6
Citrate(4TMS)	C18H40O7Si4	480	TMS-CO2, TMS-OH	207	C11H21O4Si2	273	1-2-3-4-5
Fumarate(2TMS)	C10H20O4Si2	260	CH3	15	C9H17O4Si2	245	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	CH3	15	C12H27O5Si3	335	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	TMS-CO2	117	C9H21O3Si2	233	2-3-4
Pyruvate(MEOX)(TMS)	C7H15NO3Si	189	CH3	15	C6H12NO3Si	174	1-2-3
Succinate(2TMS)	C10H22O4Si2	262	CH3	15	C9H19O4Si2	247	1-2-3-4
Amino acids							
Asparagine(2TMS)	C10H24N2O3Si2	276	TMS-CO2	117	C6H15N2OSi	159	2-3-4
Asparagine(3TMS)	C13H32N2O3Si3	348	TMS-CO2	117	C9H23N2OSi2	231	2-3-4
Aspartate(2TMS)	C10H23NO4Si2	277	TMS-CO2	117	C6H14NO2Si	160	2-3-4
Aspartate(3TMS)	C13H31NO4Si3	349	TMS-CO2	117	C9H22NO2Si2	232	2-3-4
Glutamate(3TMS)	C14H33NO4Si3	363	TMS-CO2	117	C10H24NO2Si2	246	2-3-4-5
Glutamine(3TMS)	C14H34N2O3Si3	362	TMS-CO2	117	C10H25N2OSi2	245	2-3-4-5
Glycine(3TMS)	C11H29NO2Si3	291	TMS-CO2	117	C7H20NSi2	174	2
Isoleucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Leucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Lysine(4TMS)	C18H46N2O2Si4	434	TMS-CO2	117	C14H37N2Si3	317	2-3-4-5-6
Methionine(2TMS)	C11H27NO2SSi2	293	TMS-CO2	117	C7H18NSSi	176	2-3-4-5
Ornithine(4TMS)	C17H44N2O2Si4	420	none		none		1-2-3-4-5
Phenylalanine(2TMS)	C15H27NO2Si2	309	TMS-CO2	117	C11H18NSi	192	2-3-4-5-6-7-8-9
Phenylalanine(2TMS)	C15H27NO2Si2	309	C7H7	91	C8H20NO2Si2	218	1-2
Proline(2TMS)	C11H25NO2Si2	259	TMS-CO2	117	C7H16NSi	142	2-3-4-5
Serine(2TMS)	C9H23NO3Si2	249	TMS-CO2	117	C5H14NOSi	132	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CO2	117	C8H22NOSi2	204	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CH2O	103	C8H20NO2Si2	218	1-2
Threonine(3TMS)	C13H33NO3Si3	335	TMS-CO2	117	C9H24NOSi2	218	2-3-4
Tryptophan(2TMS)	C17H28N2O2Si2	348	2TMS-C2H2NO2	218	C9H8N	130	3-4-5-6-7-8-9-10-11
Tryptophan(2TMS)	C17H28N2O2Si2	348	C9H8N	130	C8H20NO2Si2	218	1-2
Tryptophan(3TMS)	C20H36N2O2Si3	420	2TMS-C2H2NO2	218	C12H16NSi	202	3-4-5-6-7-8-9-10-11
Tryptophan(3TMS)	C20H36N2O2Si3	420	C12H16NSi	202	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	C10H15OSi	179	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	TMS-CO2	117	C14H26NOSi2	280	2-3-4-5-6-7-8-9
Valine(2TMS)	C11H27NO2Si2	261	TMS-CO2	117	C7H18NSi	144	2-3-4-5
Valine(2TMS)	C11H27NO2Si2	261	C3H7	43	C9H20NO2Si2	218	1-2
α-Alanine(2TMS)	C9H23NO2Si2	233	TMS-CO2	117	C5H14NSi	116	2-3
β-Alanine(3TMS)	C12H31NO2Si3	305	CH3	15	C11H28NO2Si3	290	1-2-3

1028

1029

Figure legends

Figure 1. Workflow used in this study.

Figure 2. **Evaluation of Carbon Isotopologue Distribution (CID) measurements for organic and**
amino acids by GC-MS using ^{13}C-PT reference samples. **A**, Evaluation of amino acids using ^{13}C-
PT protein hydrolysates. B, Evaluation of organic acids using ^{13}C-PT full metabolites extract. Black
bars correspond to expected CIDs and blue bars to the measured CIDs. The results are presented as the
mean ± SD of four independent biological replicates.

Figure 3. Evaluation of the measurement accuracy for CID and mean ^{13}C enrichment at ^{13}C-PT
and ^{13}C natural abundance in plants. A, CID accuracy and **B**, Mean ^{13}C enrichment for the selected
TMS-derivatives of organic and amino acids using ^{13}C-PT samples. **C**, Residual mean ^{13}C enrichment
measured in chemical standards, Arabidopsis seedlings and *Brassica napus* leaves after correction for
isotope natural abundance. **D**, Isotopologue-specific biases identified at ^{13}C natural abundance for
Proline, Valine and Threonine fragments. The results are presented as the mean ± SD of four
independent biological replicates or independent preparations for commercial chemical standards.

Figure 4. **Dynamics of U-^{13}C-pyruvate incorporation in *B. napus* leaves exposed to light or dark**
conditions. **A**, Mean fractional ^{13}C enrichment and **B**, Metabolite pool sizes. Mean fractional ^{13}C
enrichment were monitored with GC-MS, while absolute quantification of organic and amino acids
was performed respectively with GC-FID and HPLC-UV. The results are presented as the mean ± SD
of four independent biological replicates.

Figure 5. **Fractional ^{13}C enrichment at the isotopologue level for TCA-cycle derived metabolites.**
A, Light conditions and B, Dark conditions. The M0 isotopologue was not represented to facilitate the
visualization of low isotopologue enrichments. The results are presented as the mean ± SD of four
independent biological replicates.

Figure 6. **Fractional contribution of PEPc and TCA cycle to Malate and Citrate production in**
light and dark conditions. A, Network reconstruction at the isotopomer level based on the
isotopologues detected. **B**, Fractional contribution of PEPc and TCA cycle to Malate and Citrate
production based on pathway-specific mass isotopologues/isotopomers. In the proposed network, the
second "turn" reflected the commitment of a molecule that has already experienced all reactions of the
TCA cycle to another round of reactions (Orange and blue labels to compare acetyl-CoA-derived
labeling from 1st and 2nd "turn"). Note that M0 istopologues of all metabolites will still continue to
produce M0 istopologues at a significant propotion after a real turn of the TCA cycle, given the ^{13}C-
enrichment for Acetyl-CoA active pool. For these calculations, only the timepoint T4 and T6 hours
were used and pooled. The results are presented as the mean ± SD of four independent biological
replicates. Statistical comparison of Acetyl-CoA enrichment for light and dark conditions was
performed with a Student-test, while fractional contributions were compared with an ANOVA
followed by a post-hoc Tukey-HSD test (letters indicate groups that were separated with a p-value
<0.05).

Figure 7. **Fractional contribution of PDH and TCA-cycle derived decarboxylations to**
photosynthesis. A, Fractional ^{13}C enrichment at the isotopologue level for Glycine and Serine. **B**,
Working hypotheses to explain the photorespiratory-dependent labeling pattern of Glycine and Serine.
C, Estimation of the fractional contribution of PDH and TCA-cycle CO_2 to photosynthesis with Serine
(expressed in %). For these calculations, only the timepoint T4 and T6 hours were used and pooled
(metabolic and isotopic steady-state requirements, assumption with ^{13}C/^{12}C of CO_2 released). The

results are presented as the mean ± SD of four independent biological replicates. Statistical comparison
of the two Serine fragments was performed with a Student-test.

Figure 8. **Proposed metabolic fluxes for plant TCA cycle and PEPc in light and dark conditions**
based on this study. The contribution of stored citrate to is unlikely to occur based on our results,
while the forward TCA cycle has a higher contribution to both malate and citrate production than PEPc.
Questions remains about the exact proportion of Glutamate and Aspartate carbon backbones escaping
the TCA cycle. Despite a reduction of PEPc activity in dark conditions compared to light conditions,
PEPc-derived carbons show a higher commitment to Citrate production in dark conditions due to a
higher rate of Malate import into mitochondria. Reassimilation of PDH and TCA-cycle derived CO_2
by photosynthesis may represent up to 40% of the TCA cycle metabolic flux. See Discussion for
detailed explanations. TCA, Tricarboxylic acids; PEPc, Phosphoenolpyruvate carboxylase; OAA,
Oxaloacetate; GDH, Glutamate dehydrogenase; GABA, Gamma-aminobutyric acid; ALAAT,
Alanine:2-oxoglutarate aminotransferase; ASAT, Aspartate:2-oxoglutarate aminotransferase.

Identification of TMS-derivative mass fragments maximizing carbon backbones

Production of reference labelled samples

Validation of TMS-derivative mass fragments for CID measurements

A CID accuracy

B Fractional Mean ^{13}C enrichment

C Bias at natural ^{13}C abundance

D
Proline_2TMS_C2C5
m/z = 146
Valine_2TMS_C2C5
Threonine_3TMS_C2C4

A Mean fractional ¹³C enrichment

B Metabolite pool sizes

A

Glycine_3TMS_C2

Glycine_3TMS_C2

Serine_3TMS_C1C2

Serine_3TMS_C1C2

Serine_3TMS_C2C3

Serine_3TMS_C2C3

Fractional ^{13}C enrichment

Time (hours)

isotopologue
1
2

B

C

Contribution of PDH and TCA cycle derived CO_2 to photosynthesis

t-test p-value = 0.07