Validation of carbon isotopologue distribution measurements by GC-MS and application to [13]C-metabolic flux analysis of the tricarboxylic acid cycle in *Brassica napus* leaves

Younès Dellero[1, 2, 4*], Solenne Berardocco[1, 2], Cécilia Berges[3, 4], Olivier Filangi[1, 2, 4], Alain
Bouchereau[1, 2, 4]

[1]Institute for Genetics, Environment and Plant Protection (IGEPP), National Research Institute for
Agriculture, Food and Environment (INRAE), Université Rennes, Institut Agro , Le Rheu, France

[2]Metabolic profiling and metabolomics platform, Institute for Genetics, Environment and Plant
Protection, Biopolymers Interactions Assemblies, French National Research, Le Rheu, France

[3]TBI, Université de Toulouse, National center for Scientific Research (CNRS), National Institute for
Research for Agriculture, food and Environment (INRAE), National Institute of Applied Sciences
(INSA), Toulouse, France;

[4]MetaToul-MetaboHUB, National infrastructure of metabolomics and fluxomics, Toulouse, France

* Correspondence:
Corresponding Author
younes.dellero@inrae.fr

Keywords: Isotope, gas chromatography-mass spectrometry, trimethylsilyl (TMS) derivatives,
accuracy, regulation, phosphoenolpyruvate carboxylase, decarboxylation, TCA cycle. (Min.5-
Max. 8)

Abstract

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of [13]C-Metabolic Flux Analysis at
isotopically non-stationary steady-state. This approach requires a high level of accuracy for the
measurement of Carbon Isotopologue Distribution in plant metabolites. But this accuracy has still not
been evaluated at the isotopologue level for GC-MS, leading to uncertainties for the metabolic fluxes
calculated based on these fragments. Here, we developed a workflow to validate the measurements of
CIDs from plant metabolites with GC-MS by producing tailor-made *E. coli* standard extracts harboring
a predictable binomial CID for some organic and amino acids. Overall, most of our TMS-derivatives
mass fragments were validated with these standards and at natural isotope abundance in plant matrices.
Then, we applied this validated MS method to investigate the light/dark regulation of plant TCA cycle
by incorporating U-[13]C-pyruvate to *Brassica napus* leaf discs. We took advantage of pathway-specific
isotopologues/isotopomers observed between two and six hours of labeling to show that the TCA cycle
can operate in a cyclic manner under both light and dark conditions. Interestingly, this forward cyclic
flux mode has a nearly four-fold higher contribution for pyruvate-to-citrate and pyruvate-to-malate
fluxes than the phosphoenolpyruvate carboxylase (PEPc) flux reassimilating carbon derived from some
mitochondrial enzymes. The contribution of stored citrate to the mitochondrial TCA cycle activity was
also questioned based on dynamics of [13]C-enrichment in citrate, glutamate and succinate and variations
of citrate total amounts under light and dark conditions. Interestingly, there was a light-dependent [13]C-
incorporation into glycine and serine showing that decarboxylations from pyruvate dehydrogenase

complex and TCA cycle enzymes were actively reassimilated and could represent up to 5% to net
photosynthesis.

1   INTRODUCTION

Metabolic fluxes within central metabolism are among the most important determinants of plant cell
physiology, supporting central energetical functions and allowing them to quickly adapt to
environmental changes by controlling metabolite turnover (Clark et al., 2020). They can be estimated
from ^{13}C-Metabolic Flux Analysis (MFA), which relies on the tracing of ^{13}C throughout a metabolic
network in a relative metabolic steady-state to model and/or compare metabolic fluxes (Ratcliffe and
Shachar-Hill, 2006;Allen and Young, 2019;Antoniewicz, 2021). In phototrophic tissues, ^{13}C-MFA can
only be performed at isotopically non-stationary steady-state (INST-MFA) and requires a high level of
accuracy for the measurement of ^{13}C enrichment within metabolites (Cheah and Young, 2018;Cheah
et al., 2020;Antoniewicz, 2021;Treves et al., 2022). Since small errors in mass isotopologue
distribution can propagate to large errors in estimated fluxes (Antoniewicz et al., 2006), an accurate
measurement of ^{13}C-enrichment still remains an important prerequisite to explore plant metabolism
dynamics through isotopic labeling approaches. Two major analytical methods are currently employed
to analyze the Carbon Isotope Distribution (CID) within plant metabolites: Nuclear Magnetic
Resonance (NMR) and Mass Spectrometry (MS). These methods can provide information either at the
isotopologue level, *i.e.* there is a separation of labeled molecules according to their numbers of ^{13}C
atoms for a given metabolite, or at the isotopomer level, *i.e.* there is a separation of labeled molecules
according to the position of ^{13}C atoms for a given metabolite (Des Rosiers et al., 2004;Ratcliffe and
Shachar-Hill, 2006). While NMR analysis of the ^{13}C enrichment can provide positional labeling
information, it has not been widely used in plants due to the need of large amounts of fresh material to
overcome its low sensitivity at natural ^{13}C abundance, thus preventing the possibility to perform time-
course ^{13}C-isotope tracing experiments (Kruger et al., 2007;Masakapalli et al., 2010;Dellero et al.,
2016;Abadie et al., 2017;Abadie and Tcherkez, 2021). Conversely, mass spectrometry has been
preferentially used to analyze the Carbon Isotope Distribution (CID) within plant metabolites during
time-course ^{13}C-isotope tracing experiments (Szecowka et al., 2013;Ma et al., 2014;Medeiros et al.,
2018;Treves et al., 2022). This method is highly sensitive at natural ^{13}C abundance but essentially gives
information at the isotopologue level, given that MS fragments can hardly cover the complete set of
all possible isotopomers for a given metabolite (Lima et al., 2021). Interestingly, the recent analysis of
CID for amino acids by LC-MS revealed some biases and errors in the measurement of CID (Heuillet
et al., 2018), thus highlighting the need to also evaluate measurement accuracy of MS methods for
subsequent use in ^{13}C-MFA of plants.

Among plant metabolites, organic and amino acids are major players for plant primary metabolism.
They are actively involved in crucial physiological processes such as photorespiration, *de novo*
nitrogen assimilation and mitochondrial respiration (Xu et al., 2012;Zhang and Fernie, 2018;Dellero,
2020;Timm, 2020). Interestingly, most of the carbon backbones for amino acid biosynthesis come from
organic acids of the tricarboxylic acid (TCA) cycle, raising this cycle as a central hub for plant primary
metabolism (Araujo et al., 2012;Galili et al., 2016;Tcherkez et al., 2017). The functioning and
regulation of the plant TCA cycle has been described based on flux balance models (Sweetlove et al.,
2010;Araujo et al., 2012;Cheung et al., 2014) but also based on ^{13}C-labeling studies with different cell
types/experimental setups: attached leaves (Xu et al., 2021;Xu et al., 2022), detached leaves (Tcherkez
et al., 2009;Gauthier et al., 2010;Zhang et al., 2021), leaf discs (Le et al., 2021), guard cell cultures
(Daloso et al., 2017), heterotrophic cell cultures (Masakapalli et al., 2010;Masakapalli et al., 2014) and
isolated mitochondria (Le et al., 2021;Le et al., 2022). Overall, most of these works proposed that the
plant TCA cycle is not cyclic in the light, with two branches acting separately: i) a first branch

stimulated by the anaplerotic activity of phosphoenolpyruvate carboxylase (PEPc) leading to the
production of oxaloacetate, malate and fumarate; ii) a second branch stimulated by the entry of citrate
leading to the production of isocitrate, 2-oxoglutarate, glutamate and sometimes succinate. These
conclusions were notably supported by the inhibition of the pyruvate dehydrogenase complex (PDC)
and the mitochondrial respiration in the light while nitrogen assimilation was relatively high compared
to dark conditions and required important amounts of 2-oxoglutarate (Tcherkez et al., 2005;Gauthier
et al., 2010). In addition, the photorespiratory activity of the mitochondrial glycine decarboxylase
complex could largely compete with the second branch of the TCA cycle for NADH production,
thereby sustaining most of the mitochondrial energetic production in the light, independently of the
activity of TCA cycle enzymes (Timm and Hagemann, 2020). Besides this, the requirement to support
continued export of sugar and amino acids from the leaf during the night and to meet cellular
maintenance costs was also invoked to explain the light-dependent bi-directional functioning of the
TCA cycle (Cheung et al., 2014). Regarding the metabolic origin of citrate for the second branch in
the light, it can come from: i) TCA cycle *de novo* biosynthesis using acetyl-CoA derived from either
glycolysis and alanine transamination; ii) a stored citrate pool in a light/dark dependent-manner; iii)
the peroxisomal glyoxylate cycle (Gauthier et al., 2010;Sweetlove et al., 2010;Cheung et al.,
2014;Dong et al., 2018;Le et al., 2021). Based on the NMR analysis of combined $^{13}CO_2$ and
$^{15}NH_4^{15}NO_3$ labeling experiments in *Brassica napus* leaves, it was proposed that a high proportion of
stored and non-labeled citrate (vacuolar pool probably) was involved in the synthesis of 2-oxoglutarate
under light conditions to sustain *de novo* nitrogen assimilation through glutamate (Gauthier et al.,
2010). Under this scenario, the citrate pool would be replenished during dark conditions from
glycolysis-derived ^{13}C-pyruvate, leading to a higher ^{13}C enrichment of the citrate after a light/dark
cycle. Thus, the light and dark environmental conditions are expected to introduce important variations
of metabolic fluxes within the plant TCA cycle (cyclic versus non-cyclic) and its associated PEPc
anaplerotic pathway in *B. napus* leaves. Nevertheless, these variations have still not been carefully
investigated with time-course ^{13}C-isotope tracing experiments and pool size quantification/estimation
(^{13}C-MFA concepts), thus casting doubt on previous conclusions drawn in *B. napus* from NMR-based
experiments (one time-point, no pool size quantification/estimation). In addition, recent $^{13}CO_2$ labeling
experiments coupled to ^{13}C-INST-MFA succeeded to model leaf primary metabolism (including TCA
cycle) without the need of a stored ^{12}C Citrate pool in *Camelina sativa* and Arabidopsis leaves (Xu et
al., 2021;Xu et al., 2022).

To date, the routine analysis of CID for plant organic and amino acids has been often assessed by GC-
MS using TMS-derivatives (Szecowka et al., 2013;Araujo et al., 2014;Lima et al., 2018;Medeiros et
al., 2018;Lima et al., 2021;Treves et al., 2022) although previous comparisons of two derivatization
methods for GC-MS suggested that TBDMS-derivatives were potentially less prone to errors for amino
acid analysis than TMS-derivatives (Antoniewicz et al., 2007;Young et al., 2014). However, the
suitability of molecular ions from TMS and TBDMS derivatives was only performed using U-^{13}C-
metabolites in each of these studies, which did not evaluate the entire CID but only the carbon
distribution between the unlabeled isotopologue and the fully labeled isotopologue (M0 and M5 for a
5-carbon metabolite for example). Given that isotopic effects and analytical biases may occur for any
isotopologue of each tested metabolite, it is essential to validate MS measurements for each
isotopologue. This validation step is even more important for plants, given that plant metabolites show
strong differences of ^{13}C-enrichment dynamics between amino and organic acids (Szecowka et al.,
2013). For this purpose, the recent development of an *E. coli* ^{13}C biological standard with controlled
patterns of CID for organic and amino acids may help to address this issue (Millard et al., 2014).
Indeed, the recent use of these reference samples allowed to identify important analytical biases in the
CID measurement of Leucine and Asparagine in a concentration-dependent manner by LC-MS
(Heuillet et al., 2018). Interestingly, this validated LC-MS method successfully measured nitrogen

isotope distribution within plant amino acids and led to the estimation of metabolic fluxes from ^{15}N-
isotope transient labeling experiments in *B. napus* leaves at a metabolic steady-state using the
ScalaFlux approach (Dellero et al., 2020b;Heuillet et al., 2020;Millard et al., 2020).

Here, we proposed a workflow to assess the measurement accuracy of CID from organic and amino
acids by GC-MS for applications in plants. This methodology was based on chemical standards, plant
samples at natural ^{13}C abundance and two specific *E. coli* ^{13}C reference samples (^{13}C-PT), produced to
specifically harbor controlled patterns of CID for organic and amino acids (Pascal's Triangle
coefficients). Next, we applied our validated MS method to investigate the light/dark regulation of
metabolic fluxes within plant TCA cycle by performing U-^{13}C-pyruvate labeling kinetic experiments
on *B. napus* leaf discs under light or dark conditions. Using ^{13}C-MFA principles and some specific
isotopologues, we questioned the contribution of stored citrate to the TCA cycle activity and showed
that the TCA cycle partially works as a cycle under light conditions in *B. napus*. We also analyzed the
fractional contribution of phosphoenolpyruvate carboxylase (PEPc) and TCA cycle for pyruvate-to-
malate and for pyruvate-to-citrate fluxes.

2   MATERIALS AND METHODS

Organisms and culture

The strain *Escherichia coli* K-12 MG1655 was grown on minimal medium containing 5 mM
KH_2PO_4,10mM Na_2HPO_4, 9 mM NaCl, 40 mM NH_4Cl, 0.8 mM $MgSO_4$, 0.1 mM $CaCl_2$, 0.3 mM
thiamine, and 45 mM of the ^{13}C-labeled acetate mixture as the sole carbon source. Acetate and thiamine
were sterilized by filtration, whereas the other compounds were autoclaved separately. Batch cultures
were carried out in 1-L baffled flasks with 500 mL of medium sparged with synthetic gas (80%
N_2/20% O_2 containing <1 ppm CO_2, Air Liquide). Cultivation was performed in a Multifors Bioreactor
(Infors HT, Bottmingen-Basel, Switzerland) at 37 °C and 200 rpm of orbital shaking. The pH was
maintained at 7.0 by the addition of 2 M HCl. Cell growth was monitored by measuring optical density
at 600 nm with a Genesys 6 spectrophotometer (Thermo, Carlsbad, CA, USA). Amount of biomass in
each sample was calculated using a coefficient of 0.37 g of cell dry weight per OD unit. Arabidopsis
thaliana seeds were sterilized for 10 min with absolute ethanol containing 7.5% NaClO and rinsed two
times with absolute ethanol. Seeds were sowed on ½ MS media containing 1% Agar and incubated to
the dark at 4°C for 48-72 hours to break seed dormancy. Seedlings were grown at ambient air under a
day/night cycle with 16 h light/8h dark at a light intensity of 120 µmol photons.m^{-2}.s^{-1} and 20°C
day/18°C night. Seedlings were harvested after 2-3 weeks by instant sampling with liquid nitrogen and
lyophilized for 96 h before to be stored at -80°C until analysis. Two-month old *Brassica napus* plants
were obtained from seeds exactly as described before (Dellero et al., 2020a). The leaf L9 was used for
all experiments (natural ^{13}C abundance and U-^{13}C-pyruvate incorporation).

U-^{13}C-pyruvate labeling experiments

Four hours after the beginning of the light period, each leaf was cut at the basis of the petiole and then
66 leaf discs (0.8 cm^2) were randomly punched with a cork-borer in both laminas of each leaf. Leaf
discs were first floated in a 10 mM MES-KOH (pH 6.5) buffer for 15 min in order to process the L9
leaf for four plants (see details for leaf annotation in (Dellero et al., 2020a)). In this experiment, one
leaf of a plant corresponded to a biological replicate. Then for each time point, six leaf discs from the
same leaf were transferred in a well of a 6-well microplate filled with 4 mL/well of a buffer containing
10 mM MES-KOH (pH 6.5) and 10 mM U-^{13}C-pyruvate (^{13}C isotopic purity of 99%; sodium salt from
Cambridge Isotope Laboratories (Eurisotop, France)). All incubations were performed at 20 °C with

an orbital shaking of 70 rpm under either a continuous light of 120 µmol photons m^{-2} s^{-1} or under dark
conditions using multiple opaque plastic bags, similarly to the previous experimental set-up used for
dark-induced senescence (Dellero et al., 2020a). The timepoint 0 was the same for both conditions as
it corresponded the beginning of the experiment. For each time point (0, 0.5, 1, 2, 4, 6 hours of
incubation), the leaf discs were harvested and rinsed three times (10 sec) with a neutral buffer (i.e.,
without U-^{13}C-pyruvate) and dried for 10 s on a clean tissue before being frozen in liquid nitrogen and
stored at -80 °C. The timepoint 0 of the experiment was incubated for 2 sec with the labeled buffer
before being directly rinsed. This was done to evaluate the residual amount of ^{13}C$_3$-pyruvate detected
in our analysis that was not truly incorporated into leaf discs (a mean ^{13}C enrichment close to 35% for
pyruvate at T0 was observed in our study). This amount reflected probably biases in the washing
procedure and/or an artificial fixation of ^{13}C$_3$-pyruvate to the leaf disc surface.

Analysis of ^{13}C-labeled Acetate by NMR

An equimolar mixture of all ^{13}C-isotopomers of acetate was produced from ^{12}C-acetate, 1-^{13}C-acetate,
2-^{13}C-acetate and U-^{13}C-acetate (^{13}C isotopic purity of 99%, sodium salts from Cambridge Isotope
Laboratories (Eurisotop, France); **Figure 1**). Quantitative 1H 1D- NMR was used to control the
isotopic composition of the ^{13}C-labeled substrate before use. NMR spectra were recorded with an
Avance 500 MHz spectrometer (Bruker, Rheinstetten, Germany) at 298 K, using a 30° pulse and a
relaxation delay of 20 s. The proportion of each isotopic form was quantified by fitting using a mixed
Gaussian−Lorentzian model (Millard et al., 2017). We obtained a molecular ^{13}C enrichment of 49.9 %
± 0.8.

Metabolite extraction and quantification

^{13}C-PT full metabolites and ^{13}C-PT protein hydrolysates samples were obtained following procedures
described previously (Heuillet et al., 2018). Briefly, for the ^{13}C-PT full metabolites extract, 1 mL of *E.*
coli broth at OD600 nm = 1.7 was micro-filtered and the pellet was rapidly quenched at -20 °C into a
mixture of acetonitrile/methanol/H$_2$0 (2:2:1) with 0.1% formic acid. After a centrifugation step (7000g,
−20 °C, 15 min), the supernatant was evaporated and resuspended in 250 µL of Milli-Q water, to be
stored at −80 °C until analysis. For the ^{13}C-PT protein hydrolysates extract, the pellet obtained with
the cellular extract was subjected to an acid hydrolysis with 6N HCl at 110 °C overnight. Samples were
evaporated and rinsed 3 times with Milli-Q water. After a centrifugation step (10 min, 12 000g), the
supernatant was stored at −80 °C. Polar metabolites of freeze-dried Arabidopsis seedlings and *B. napus*
leaves were extracted with a MeOH/chloroform/H$_2$O mixture following procedures described
previously (Dellero et al., 2020b). The quantification of amino acid content was performed by HPLC-
UV exactly as described previously (Dellero et al., 2020b). The quantification of organic acids was
performed by GC-FID exactly as described previously (Bianchetti et al., 2021).

Determination of Carbon Isotopologue Distribution by GC-MS

Dried aliquots of samples were derivatized on a MultiPurpose Sampler Gerstel and analyzed by GC-
MS (GC-2010 Plus Shimadzu coupled to a triple quadrupole mass spectrometer GCMS-TQ8040
Shimadzu). Polar metabolites were first derivatized for 1 h at 40°C with 50 µL of pyridine containing
20 g.l^{-1} methoxyamine chloride and then were incubated with 50 µL of N-Methyl-N-(trimethylsilyl)-
trifluoroacetamide (MSTFA) for 0.5 h at 40°C. 1 µL was injected with a split mode 1/20 at 260°C on
a TG-5MS column (26098-1420, Thermo Scientific). Separation was performed in a Helium gas-
stream at 1 mL.min^{-1} using the following temperature ramp : 4 min at 70°C, +10°C.min^{-1} until 198°C,
2 min at 198°C, +1°C.min^{-1} until 202°C, +15°C.min^{-1} until 268°C, 3 min at 268°C, +1°C.min^{-1} until

272°C,+10°C.min^{-1} until 310°C and 7 min à 310°C. The transfer line was maintained at 300°C and the
ion source at 200°C. Ionization was performed by electron impact at 70 eV and the MS acquisition was
carried out after a solvent delay of 5 min in full scan mode at 7 spectra.s$_{-1}$ over the m/z range 45-700.
MS Spectra were processed with the software "GCMS Solutions" supplied by Shimadzu to adjust
retention times between each run using a mixture of alkanes (C7 to C40, Sigma-Aldrich). Peak identity
was established according to their retention time and mass fragmentation spectra with respect to
authentic standards derivatized (**Table S1**), and by reference to mass spectra of derivatives in the US
National Institute of Standards and Technology database and in the Golm metabolome database. All
mass isotopologue raw areas were subjected to correction for the contribution of naturally occurring
isotopes of all elements except the carbon arising from the metabolite backbone using the IsoCor v2.1.4
package of Python (https://github.com/MetaSys-LISBP/IsoCor) in low resolution mode (Millard et al.,
2019). In addition, unlabeled chemical standards, Arabidopsis seedlings and *B. napus* leaves were
additionally corrected for the contribution of naturally occurring isotopes of the carbon arising from
the metabolite backbone (giving a residual molecular ^{13}C enrichment of 0 for molecules labeled at
natural ^{13}C abundance). ^{13}C-PT samples were not concerned by this additional correction since they
held fully controlled ^{12}C/^{13}C patterns at all positions. A correction method dedicated to IsoCor was
specifically created for the TMS-derivative fragment described in this study (**Table 1**), *i.e.* by
considering the contribution of all atoms added by the derivatization on the isotopic measurement of
the monitored fragments (chemical separation of all atoms arising from the metabolite and the
derivative). All files required to run this method on IsoCor are available in the supplementary material
of this article (**Supplementary files**) and all information will be updated indefinitely in the following
INRAE dataverse (Dellero and Filangi, 2021). The method is publicly available and can also be run
directly from GC-MS source files using the following Galaxy workflow (https://galaxy.genouest.org/,
section "Metabolomics", subsections "Conversion GCMS PostRun Analysis to IsoCor" and "Isotope
Correction for mass spectrometry labeling experiments"). For this purpose, each raw GC-MS dataset
should contain a column "Name" for the fragments considered and a column "Area". The name of each
fragment must be written exactly as specified in the "Metabolite.dat" (an example file is provided in
the **supplementary material**). Fractional Carbon Isotopologue Distribution and mean ^{13}C enrichment
were directly obtained from IsoCor outputs (isotopologue_fraction and mean_enrichment) while CIDs
were rescaled to Pascal's Triangle coefficients by multiplying with 2^n, where n is the number of carbon
atoms (Millard et al., 2014). The accuracy of measurements was assessed from the closeness of
experimental and predicted values for CIDs and mean ^{13}C enrichments at either ^{13}C-PT abundance or
at natural ^{13}C abundance. CID accuracy was formally defined as the mean of absolute differences
between measured and predicted values for each isotopologue (Heuillet et al., 2018).

Pathway fractional contribution

Pathway fractional contribution (FC) can be applied to a metabolic network in a metabolic and isotopic
steady-state to compare the relative contribution of the considered pathway to a metabolite, as long as
these pathways produced specific isotopologues for this metabolite (Buescher et al., 2015). Basically,
it consists in the comparison (ratios) of mass isotopologues derived from each pathway in the final
metabolite considered to determine their relative contributions (for example: isotopologues from
pathway A/(isotopologues from pathway A and pathway B)). Here, FC was calculated starting from
U-^{13}C-pyruvate and considering malate and citrate as final metabolites. We considered two major
fluxes for the biosynthesis of malate and citrate from pyruvate: the forward mode of the TCA cycle,
reflecting the successive action of multiple enzyme from this cycle, and the PEPc flux, but only
associated to the reassimilation of TCA cycle and PDC-derived CO_2. Indeed, PEPc could also use
$^{12}CO_2$ arising independently from the decarboxylation of U-^{12}C-pyruvate, leading to an unknow
fraction of unlabeled malate derived from PEPc and entering the TCA cycle. To apply this

methodology, we selected the timepoint 4 and 6 hours because there was a metabolic steady-state and
an isotopic pseudo steady-state for isotopologue ratios. Malate biosynthesis from $^{13}C_3$-pyruvate
produced M1, M2 and M3 isotopologues in the fragment Malate_3TMS_C1C4, with M1 belonging to
both PEPc and TCA cycle fluxes, while M2 and M3 were exclusively attributed to the TCA cycle (See
result section). Interestingly, TCA cycle produced both M1 and M3 at the same time in the fragment
Malate_3TMS_C1C4. This M3/(M1+M3) ratio was also observed in other fragments
(Succinate_2TMS_C1C4 and Glutamate_2TMS_C2C5), relatively stable, and corresponded to the
same metabolite precursors, thereby reflecting the M0 and M2 from mitochondrial active pool of
acetyl-CoA (See result section). Thus, this ratio was used to calculate the proportion of M1 from
Malate_3TMS_C1C4 fragment belonging exclusively to the TCA cycle based on the M3 fragment
produced only by the contribution of the TCA cycle. In the following equations, *mi* refers to the
isotopologue enrichment calculated with respect to the CID (sum of all *mi* for a given metabolite equal
to 1). As all these calculations are relative, it is also possible to use corrected areas of mass
isotopologues (outputs from IsoCor).

278
$$Succinate C1C4 \left(\frac{m3}{m1+m3}\right) = Glutamate C2C5 \left(\frac{m3}{m1+m3}\right) = Acetate \left(\frac{m2}{m0+m2}\right) = X$$

279
$$FC\ Malate_{PEPc} = Malate C1C4 \left(\frac{m1 - \left(\left(\frac{1-X}{X}\right) \times m3\right)}{m1+m2+m3}\right)$$

280
$$FC\ Malate_{TCA\ cycle} = Malate C1C4 \left(\frac{m2+\frac{m3}{X}}{m1+m2+m3}\right)$$

281 Citrate biosynthesis from U-^{13}C-pyruvate produced M1, M2, M3 and M4 isotopologues in the fragment
282 Citrate_4TMS_C1C6, with M1 belonging exclusively to PEPc, M3 belonging to both PEPc and TCA
283 cycle and M2 and M4 belonging exclusively to TCA cycle (See result section). Therefore, FC were
284 calculated as follow:

285
$$FC\ Citrate_{PEPc} = Citrate C1C6 \left(\frac{m1+m3}{m1+m2+2\times m3+m4}\right)$$

286
$$FC\ Citrate_{TCA\ cycle} = Citrate C1C6 \left(\frac{m2+m3+m4}{m1+m2+2\times m3+m4}\right)$$

Finally, the FC of TCA cycle and PDC-derived CO_2 to photosynthesis was calculated using serine
fragments and the ratio "X" as a proxy for the $^{13}C/^{12}C$ ratio of this CO_2 fraction. We did not consider
the contribution of plastidial PDC here since a recent ^{13}C-MFA of tobacco leaves showed that this flux
was relatively negligible compared to the mitochondrial PDC flux (Chu et al., 2022). Considering the
isotopic and metabolic steady-state for serine fragments, the mean ^{13}C enrichment of X should be
observed in serine if the mitochondrial fraction of CO_2 accounted for 100% of the CO_2 fixed by
photosynthesis. Thus, FC was calculated as follow:

294
$$FC\ photosynthesis_{PDH\ and\ TCA\ cycle\ CO_2} = \frac{mean\ ^{13}C\ enrichment\ (Serine C1C3)}{X}$$

295 For this, mean ^{13}C enrichment of serine was recalculated from M1 isotopologue of either
296 Serine_2TMS_C1C2 or Serine_2TMS_C2C3, since serine was only labeled in the C2 of these
297 fragments based on the hypothesis validated by our results.

Statistical analysis

299 Statistical comparisons were performed using either confidence intervals (comparison of a group of
300 value against one reference value), a Student-test (comparison of 2 groups of values) or an ANOVA
301 followed by a post-hoc Tukey-HSD test (comparison of 3 or more groups of values) with a level of p-
302 value <0.05 by using the publicly available Rstudio software and dedicated packages (RStudio Team,
303 2016).

3   RESULTS

Evaluation of CID measurements by GC-MS

The proposed workflow (**Figure 1**) consisted in three steps: 1) the build-up of a method to identify and
select abundant molecular ions deriving from TMS-derivatives with a large number of carbons; 2) the
production of two reference ^{13}C-labeled samples with a controlled pattern for CID (^{13}C-PT) using *E.*
coli as a metabolic factory; 3) the validation of TMS-derivatives mass fragments for CID measurements
with our GC-MS method for applications in plants using ^{13}C-PT reference samples and unlabeled and
^{13}C-labeled plant samples. First, we combined a literature survey analyzing common pathway of TMS-
derivative mass fragmentations with the injection of chemical standards in our analytical system (Lai
and Fiehn, 2018;Okahashi et al., 2019;Harvey and Vouros, 2020). We selected the most abundant
fragments maximizing the information on carbon backbones (**Table 1**). Their chemical identity was
further confirmed by the observed CID pattern with our two reference ^{13}C-PT samples. Next, we
produced reference ^{13}C-labeled samples with *E. coli* (^{13}C-PT samples) using an accurate equimolar
mixture of all ^{13}C isotopomers of acetate as previously described (Millard et al., 2014). This strategy
produced metabolites with fully controlled and predictable ^{13}C-labeling patterns, *i.e.* a mean ^{13}C
enrichment close of 50% (49.9 % ± 0.8 in our study based on NMR analysis) and a CID following
coefficients from Pascal's triangle (binomial distribution) (Millard et al., 2014). From *E. coli* cell
harvest, two specific ^{13}C-PT samples were produced during the extraction: i) a ^{13}C-PT full metabolite
extract containing all polar metabolites for the evaluation of CID for organic acids; ii) a ^{13}C-PT protein
hydrolysates extract containing proteogenic amino acids in high concentrations for the evaluation of
CID for amino acids. While this strategy allowed to maximize the amount of soluble proteogenic amino
acids detectable, it prevented the evaluation of glutamine and asparagine due to the hydrolysis process
(Kambhampati et al., 2019).To facilitate comparison of CIDs between the different amino acids, we
rescaled the measured ^{13}C-enrichments among isotopologues of each fragment into binomial
distribution, thus reflecting the average number of isotopomers constituting each isotopologue (**Figure**
2 and Material and Methods). In the following figures, GC-MS fragments will be denoted according
to the considered metabolite, its TMS-derivative analyzed and the metabolite carbon backbone (for
example, Glutamate_3TMS_C2C5 means that the fragment contains the carbon C2-C3-C4-C5
(**Table1**)).

Overall, all the considered amino acids were detected in the ^{13}C-PT protein hydrolysates except
tryptophan (**Figures 2A, S1A**). We found very a high level of correlation between the measured CID
(blue bars) and the expected CID (black bars) for most of the amino acid fragments considered in our
method. CID measurements for Methionine_2TMS_C2C5, Lysine_4TMS_C2C6 and
Tyrosine_2TMS_C2C9 showed a high variability, leading to small misestimations for some

isotopologues (**Figure 2A**). For methionine, this could be a consequence of its very low abundance in
the ^{13}C-PT protein hydrolysates compared to the others amino acids (**Figure S1A**). The most important
and significant errors for CID measurements of amino acids concerned Proline_2TMS_C2C5
(overestimation of M4), Threonine_3TMS_C2C4 (overestimation of M3 at the expense of M0 and
M1), Valine_2TMS_C1C2 (overestimation of M2) and Valine_2TMS_C2C5 (overestimation of M3 at
the expense of M1 and M2). Considering the recent establishment of a ^{13}C-positional isotopomer
approach for glutamate to analyze its C1 carbon in connection with PEPc activity (Lima et al., 2021),
we also analyzed the CID for the two low abundant fragments Glutamate_3TMS_C1 (CO_2-TMS, M0
at m/z 117) and Glutamate_3TMS_C1C5 fragment (CH3 loss, M0 at m/z 348) (**Figure S1B**). However,
they both showed large and significant misestimations of their CIDs, thus questioning the accuracy of
this isotopomer-based method. The analysis of organic acids from ^{13}C-PT full metabolites extract
showed   relatively   good   CID   for   Citrate_2TMS_C1C5,   Malate_3TMS_C2C4   and
Fumarate_2TMS_C1C4 (**Figure 2B**). Citrate_4TMS_C1C6 and Malate_3TMS_C1C4 showed also
relatively good CID but they had a higher variability compared to their respective C1C5 and C2C4
fragments. There was a very high overestimation of M0 in Succinate_2TMS_C1C4, a contamination
already reported for this ^{13}C-PT full metabolites extract (Millard et al., 2014), which prevented its
validation by the workflow. For Pyruvate_MEOXTMS_C1C3, the variability was too high to assess
the CID measurements, perhaps due to its weak derivatization yield and low abundance in the^{13}C-PT
full metabolites extract (**Figure S1A**). To further assess the distribution of measurement errors among
isotopologues, we analyzed the CID accuracy, using threshold values of ±0.05 (**Figure 3A**). Overall,
the bias of measurement for each isotopologue of each metabolite was relatively low except for
methionine, valine, threonine, succinate and pyruvate. Interestingly, these very high levels of accuracy
for CID measurements were converted into very high levels of accuracy for fractional mean ^{13}C
enrichment measurements, leading to very small shifts between expected (0.499 ±0.08) and measured
values (below 1% of error), except again for methionine, valine, threonine, succinate and pyruvate
(**Figure 3B**). Nevertheless, Malate_3TMS_C1C4, Lysine_2TMS_C2C6 and Serine_2TMS_C2C3
showed important variability for mean ^{13}C enrichment measurements although no strong biases in their
CID measurements were previously observed. Conversely, fractional mean ^{13}C enrichment of
Phenylalanine_2TMS_C1C2 and Proline_2TMS_C2C5 were slightly overestimated and reflected the
overestimation of their respective isotopologues M2 and M4.

Then, we evaluated the suitability of our GC-MS method for application in plants. First, we analyzed
measurement accuracy at natural ^{13}C abundance using either seedlings from Arabidopsis or leaves from
B. napus. To do so, we measured the fractional residual molecular ^{13}C enrichment, *i.e* the fractional
^{13}C enrichment remaining after removing the contribution of natural abundance for all isotopes (a value
of 0 is expected if no contamination occurs) (**Figure 3C**). Among the metabolites detected in the plant
samples, most of the fragments for organic and amino acids showed relatively high accuracies: mean
values ranged between and 0.005 and 0.03 (**Figure 3C**). Some discrepancies were observed between
Arabidopsis seedlings and *B. napus* leaves for Citrate_4TMS_C1C6 and Malate_3TMS_C1C4 (higher
accuracy   for   *B.   napus*),   and   for   AlphaAlanine_2TMS_C2C3,   Aspartate_2TMS_C2C4,
Fumarate_2TMS_C1C4 and Serine_3TMS fragments (higher accuracy for Arabidopsis). Residual
molecular ^{13}C enrichments for Phenylalanine_2TMS fragments were close to 0.05 but only in plants
samples.   The   major   errors   at   natural ^{13}C   abundance   concerned   Proline_2TMS_C2C5,
Threonine_3TMS_C2C4 and Valine_2TMS_C2C5 and were observed in both chemical standards and
unlabeled plant extracts, confirming the occurrence of systematic matrix-independent analytical bias.
A detailed survey showed that the isotopic cluster of Proline_2TMS_C2C5 was contaminated in the
M4 isotopologue (m/z 146) by a mass fragment deriving from near elution of Isoleucine_2TMS, and
correlated with the M4 overestimation previously observed in ^{13}C-PT protein hydrolysate sample
(**Figures 2, 3D**). Similarly, the isotopic cluster of Valine_2TMS_C2C5 was contaminated in the M3

isotopologue (m/z 147) while Threonine_3TMS_C2C4 showed a surprising very high abundance of
the M1 isotopologue (40% of the CID).

388 U-^{13}C-pyruvate incorporations into leaf discs and qualification of metabolic and isotopic states

We used our validated MS method to explore the light/dark regulation of TCA cycle and PEPc fluxes
in *B. napus* leaves, with a special emphasize on day/night citrate metabolism and the cyclic/non-cyclic
modes of the TCA cycle. For this purpose, we selected U-^{13}C-pyruvate as a metabolic probe since the
mitochondrial Pyruvate Dehydrogenase complex (PDC) will be able to produce simultaneously ^{13}C$_2$-
acetyl-CoA molecules suitable for the investigation of the TCA cycle and ^{13}CO$_2$ molecules suitable for
the investigation of PEPc activity through the combined action of carbonic anhydrase (H^{13}CO$_3^-$).

Following transient kinetics of 10 mM U-^{13}C-pyruvate incorporation in either light or dark conditions
for up to 6 hours, the analysis of mean ^{13}C enrichment for all conditions revealed a significant increase
for fragments of alphaalanine, aspartate, citrate, glutamate, glycine, malate, pyruvate, serine and
succinate along time (**Figure 4A**, **Table S2**; **Table S4** for statistical analysis). There was a very high
biological variability and small ^{13}C enrichment for fumarate fragment. Overall, most of these
enrichment dynamics remained relatively equivalent between light and dark conditions for all the
fragments considered, except for glycine and serine, which were almost exclusively enriched in light
conditions. These results were surprising since the commitment of an unlabeled stored ^{12}C-citrate
during light conditions was expected to introduce an isotopic dilution compared to dark conditions,
that could have been reflected in citrate, glutamate and succinate notably. Considering glycine and
serine, the importance of the light-dependent photorespiratory pathway for their production in leaves
clearly suggested that a significant part of ^{13}CO$_2$ released was reassimilated by photosynthesis. A
statistical analysis revealed that Glycine_3TMS_C2, Serine_3TMS_C1C2 and Serine_3TMS_C2C3
reached an isotopic pseudo steady-state starting after 2 hours of illumination (**Table S4**). Other
fragments showed a relatively linear evolution of their mean ^{13}C enrichment, except for
Pyruvate_MEOXTMS_C1C3 which rapidly reached an isotopic steady-state after 0.5 hours only.
Interestingly, labeling data from Pyruvate_MEOXTMS_C1C3 and Succinate_2TMS_C1C4 fragments
had a very good resolution (weak analytical and biological variability) compared to previous results
obtained with ^{13}C-PT full metabolites extract.

Next, we performed quantification of total pool sizes for these metabolites to give an approximate
indication whether the leaf metabolism was in a metabolic pseudo steady-state (plant cells are
extensively compartmented) (**Figure 4B, Table S3**). This was done by GC-FID for organic acids and
UPLC-UV for amino acids using AccQTag methodology. Indeed, the response of TMS-derivatives
for amino acids is too unstable along a GC-MS run (**Figure S3**) and absolute quantification of
metabolites using their CID masses have to be achieved according to internal isotopic standards for
each metabolite (not our case in the present study) (Heuillet et al., 2020;Evers et al., 2021). Overall,
total metabolite pool sizes were relatively different between each other's for any given time but there
was no significant difference between light and dark conditions, except for Aspartate. Glycine was
only quantified in light conditions (not detected in "dark condition" samples with UPLC-UV).
Interestingly, a statistical analysis showed that all the total metabolite pool sizes reached a significant
pseudo steady-state starting after 1 hour (values belonging to similar statistical groups in **Table S5**).
To summarize, we found some similar ^{13}C enrichment dynamics for TCA cycle derived metabolites in
both light and dark conditions combined to similar total metabolite pool sizes at a metabolic pseudo
steady-state.

429 Metabolic network considerations and dynamics of CIDs for TCA cycle-derived metabolites

According to our ^{13}C-labeling strategy and the metabolites detected with our MS method, we
constructed a metabolic network containing all the possible isotopomers that will be produced from
two "turns" of the TCA cycle and the PEPc activity, including the reassimilation of TCA cycle and
PDC-derived CO_2 (**Figure 5A**). In this network, the second "turn" reflected the commitment of a
molecule that has already experienced all reactions of the TCA cycle to another round of reactions
(Orange and blue labels to compare acetyl-CoA-derived labeling from 1st and 2nd "turn"). Based on
previous works (Abadie and Tcherkez, 2019), the reverse reaction of the fumarase was also considered
in the network, although it had no effect on the M1 isotopologue monitored by mass spectrometry (C4
and C1 isotopomers are both detected as M1 isotopologues in our study). The involvement of a light-
dependent stored ^{12}C-citrate pool was not represented here since it only provided M0 isotopologues
(no additional isotopologues to explain). The activity of the malic enzyme, converting malate to
pyruvate, was also not considered here since it may have been inhibited due to the high incorporation
of pyruvate. Overall, the PEPc-dependent reassimilation of TCA cycle and PDC-derived CO_2 was
expected to introduce M1 isotopologues for Malate_3TMS_C1C4 and M1 and M3 isotopologues for
Citrate_4TMS_C1C6. Conversely, the non-cyclic mode of the TCA cycle should introduce M2
isotopologues to Citrate_4TMS_C1C6, Glutamate_2TMS_C2C5 and Succinate_2TMS_C1C4
(incorporation of M2 acetyl-CoA arising from U-^{13}C-pyruvate) while the cyclic mode should introduce
M2 isotopologue to malate, followed by M1, M3 and M4 isotopologues for all these metabolites if the
M2 isotopologues were committed again to the TCA cycle (second "turn").

Analysis of the dynamic of CIDs showed that mean ^{13}C enrichment of citrate, glutamate and succinate
was essentially driven by an increase of the M2 isotopologue for light and dark conditions, thus
reflecting the conventional non-cyclic mode of the TCA cycle in both conditions (**Figure 5B,C**).
However, there was also an increase of M1 and M3 isotopologues for Glutamate_2TMS_C2C5 and
Succinate_2TMS_C1C4 starting after 2 hours while M1, M3 and M4 isotopologues of
Citrate_4TMS_C1C6 and Citrate_2TMS_C1C5 were increased starting from 4 hours (**Figure 5B,C**).
Therefore, the cyclic mode of the TCA cycle was necessarily operating during both light and dark
conditions. Nevertheless, M1 and M4 of Citrate_4TMS_C1C6 were essentially increased in dark
conditions compared to light conditions after 4 hours, perhaps reflecting a higher turn-over for the TCA
cycle in dark conditions. There was a very high variability for Fumarate_2TMS_C2C4 CID in both
conditions, which prevented its use for a reliable isotopic investigation. Interestingly,
Malate_3TMS_C1C4 fragment showed a higher enrichment of M1 compared to M2 in the light
condition while M2 remains higher than M1 in the dark condition. This shift may reflect an increase
of the PEPc activity under light condition. Since there was no difference of mean ^{13}C enrichment
between light and dark conditions for this fragment (**Figure 4A**), this result reinforced the importance
of analyzing specifically CIDs for the interpretation of labeling data from TCA cycle. In addition, the
CID of Malate_3TMS_C2C4 showed a similar dynamic for both light and dark conditions, hence ruling
out the use of this fragment to disentangle PEPc and TCA cycle contribution to malate production
(**Figure 5B,C**). Aspartate_3TMS_C2C4 showed a very high variability for the light condition, but the
CID dynamic in the dark condition was similar to the CID dynamic of Malate_3TMS_C2C4. Thus,
both fragments seemed to mimic oxaloacetate labeling dynamics. Pyruvate_MEOXTMS_C1C3 was
only enriched with the M3 isotopologue (nearly 80% M3/20% M0), thereby confirming a negligible
activity of the malic enzyme in our metabolic network.

Pathway fractional contribution for TCA cycle and PEPc-dependent reassimilation of TCA cycle
and PDC-derived CO_2 to either malate and citrate biosynthesis

We took advantage of the pathway-specific mass isotopologues produced respectively by the two
"turns" of the TCA cycle and PEPc to calculate pathway fractional contributions (FC) for pyruvate-to-

malate and pyruvate-to-citrate (**Figure 6A**). As PEPc activity could also be driven by other sources of
CO_2 (photorespiration and ambient air) diluting the final $^{13}C/^{12}C$ ratio for HCO_3^-, our results presented
here only reflected the PEPc activity devoted to the reassimilation of TCA cycle and PDC-derived
$^{13}CO_2$. We used only the timepoints 4 and 6 hours, because all the fragments considered for these FC
were mostly visible at these moments. In addition, pathway FC can only be applied if the metabolic
network is at both metabolic and isotopic pseudo steady-states. Here, we observed a rapid metabolic
pseudo steady-state (**Figure 4B**, **Table S5**) and a relatively negligible evolution of the ratios for the
considered isotopologues between 4 and 6 hours (similar evolutions of M1, M2 and M3 isotopologues
for malate between 4 and 6 hours for example). Thus, the relative contribution of PEPc and TCA cycle
to either malate and citrate remained in the same order of magnitude between 4 and 6 hours (**Figure
5B,C**). An important issue to address for calculating FC concerned the M1 from Malate_3TMS_C1C4,
which monitored at the same time PEPc and TCA cycle dependent isotopomers (**Figure 5A**). To deal
with, we took advantage of Succinate_2TMS_C1C4 and Glutamate_2TMS_C2C5 fragments which
both harbored M1 and M3 isotopologues that are the precursors of TCA cycle-dependent M1 and M3
of Malate_3TMS_C1C4 (**Figures 5A and 6A**). Analysis of the M3/(M1+M3) ratio for these two
fragments showed a remarkable stability between dark and light conditions, thus reflecting a stable M2
enrichment of 40% for the active pool of acetyl-CoA (**Figure 6B** and Equations). This result also
confirmed that the isotopologue ratios considered here were at an isotopic pseudo steady-state. Using
this ratio, we calculated the FC of PEPc and TCA cycle for pyruvate-to-malate and pyruvate-to-citrate
respectively (**See equations**). We found that TCA cycle had a significant four-fold higher contribution
for pyruvate-to-malate or pyruvate-to-citrate compared to PEPc in all conditions (around 80% versus
20%). Thus, PEPc-dependent reassimilation of TCA cycle and PDC-derived CO_2 was relatively low
compared to the forward cyclic mode of the TCA cycle in *B. napus* leaf discs under either light or dark
conditions. Interestingly, there was a 1.6-fold increase of the FC of PEPc for pyruvate-to-malate from
dark to light conditions (14% versus 23%), although it remained not statistically significant (high
variability, t-test p-value =0.08). Conversely, PEPc contribution for pyruvate-to-citrate was
significantly 1.5-fold higher in dark condition than in light condition (26% versus 17%). These last
results suggested a possible light/dark regulation of PEPc activity and a higher commitment of PEPc-
derived carbons to the forward part of the TCA cycle in the dark.

Contribution of TCA cycle and PDC-derived CO_2 to photosynthesis

To finish, we explored the isotopologue distribution of glycine and serine, some amino acids which
were previously identified as a proxy for photosynthetic reassimilation of PDC and TCA cycle-derived
$^{13}CO_2$ (**Figure 7A**). We did not consider the contribution of plastidial PDC here since a recent ^{13}C-
MFA of tobacco leaves showed that this flux was relatively negligible compared to the mitochondrial
PDC flux (Chu et al., 2022). Our results showed that the increase of mean ^{13}C enrichment for the three
considered fragments in light condition came from an increase of their M1 isotopologue. Given that
the light-dependent photorespiratory enzymes glycine decarboxylase and serine:hydroxymethyl
aminotransferase convert 2 glycine into 1 serine, we wanted to know whether the fragments
Serine_2TMS_C1C2 and Serine_2TMS_C2C3 were labeled in the same or different carbons, thus
revealing if the C1 of glycine was also labeled (**Figure 7B**). By reviewing all the possible scenarios
with different combinations of labeled glycine, we showed that the CID associated to our fragments
(isotopologue occurrence) could only be produced with one molecule of glycine labeled at the C2 and
an unlabeled molecule of glycine, leading to the production of a molecule of serine labeled only at the
C2 and an unlabeled CO_2 molecule. Since we have shown that glycine and serine were both at the
isotopic and metabolic steady-state between 2 and 6 hours (**Figure 4**, **Table S4**, **Table S5**), we used
serine fragments to evaluate the contribution of mitochondrial decarboxylations to the photosynthetic
activity. For this purpose, the steady-state enrichment of acetyl-CoA calculated before was used as a

proxy of the $^{13}C/^{12}C$ ratio of the TCA cycle and PDC-derived CO_2 (**See equations**) given its
preponderant contribution for light respiration (CO_2 emissions) (Tcherkez et al., 2005). After
recalculating the mean ^{13}C-enrichment of serine from both serine fragments (C1-C2 or C2-C3), we
showed that the light-dependent assimilation of TCA cycle and PDC-derived decarboxylations could
represent around 5% of net photosynthesis (**Figure 7C**).

4   DISCUSSION

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of ^{13}C-INST-MFA. This approach
requires a high level of accuracy for CID measurements to avoid propagation of small measurement
errors to large errors in estimated fluxes (Antoniewicz et al., 2006). Considering the need to perform
high throughput measurements at isotopically non-stationary steady-state, MS has been preferentially
selected by the field in the recent years to measure CID in organic and amino acids. However, the
suitability of TMS-derivative molecular ions for these metabolites has never been evaluated up to the
isotopologue level, *i.e.* by monitoring the measurement accuracy for each possible mass isotopologue
and its impact on the final CID. The use of a validated MS methodology will help to routinely
investigate ^{13}C-labeling dynamics within plant TCA cycle and strengthen our knowledge about its
functioning.

A robust methodology to evaluate CID measurements and identify analytical and matrix biases

In this study, we proposed a workflow to assess the measurement accuracy of CID from TMS-
derivatives of organic and amino acids by GC-MS for applications in plants, based on tailor-made *E.*
coli ^{13}C biological standards with controlled patterns of CID (^{13}C-PT) and plant samples at natural ^{13}C-
abundance (**Figure 1**). Overall, most of the TMS-derivatives selected (Table 1) were validated,
harboring remarkable accuracies leading to very small biases in the measurement of CIDs (below 0.05)
and fractional mean ^{13}C enrichments (below 0.01). Nevertheless, we identified three important biases
at the isotopologue level. The first bias concerned the contamination of the isotopic cluster of
Proline_2TMS_C2C5 at the m/z 146 (**Figure 3D**), which was previously reported with chemical
standards (Antoniewicz et al., 2006). Here, we showed that the contamination at m/z 146 could occur
from the co-elution with isoleucine: separation by 0.02 s in our chromatograms, *i.e* 0.33 °C along the
temperature gradient (+10°C.min^{-1}). Interestingly, this contamination had a weak impact on the mean
^{13}C enrichment measured (**Figure 3B**), suggesting that this fragment may still be used, as long as the
^{13}C-labeling strategy can prevent the production of M4 isotopologues. The second bias identified
concerned the contamination of the isotopic cluster of Valine_2TMS_C2C5 at the m/z 147. During the
fragmentation process, a siliconium ion (loss of a methyl in a TMS group) can operate a cyclization by
attacking another TMS group of the same molecule, thereby producing an ion $[(CH^3)^3Si-O-Si(CH^3)^2]^+$
of the same mass 147 (Harvey and Vouros, 2020). We also detected a significant abundance of an m/z
246 (siliconium ion) in the fragmentation spectra of Valine_2TMS, which seems to support this
hypothesis for the contamination (**Figure S2**). The third bias concerned the very high level of M1
isotopologue detected for Threonine_3TMS_C2C4. In its fragmentation spectra, we also found a M1
isotopologue contamination for the peak m/z 291, an important fragment for the identification of
Threonine_3TMS by GC-MS (**Figure S2**, NIST database). Thus, this contamination could be due to:
i) a co-elution of a contaminant chemically similar to threonine; ii) a strong analytical isotopic effect.
The use of GC coupled to high-resolution MS may help to resolve this analytical issue. Besides this,
we could not evaluate Succinate_2TMS_C1C4 with our workflow, due to a previously reported *E. coli*-
dependent contamination of ^{13}C-PT samples (Millard et al., 2014). However, this fragment showed
small biological and analytical variabilities in our ^{13}C-labeling experiments and provided

M3/(M1+M3) ratios similar to those calculated with Glutamate_2TMS_C2C5 (**Figure 6B**), hence
partially validating its use for ^{13}C-MFA in plants. Using our validated MS method, we provided
relevant information about the activity of the TCA cycle and citrate metabolism in *B. napus* leaves that
counterbalanced some conclusions from previous works (Tcherkez et al., 2009;Gauthier et al.,
2010;Sweetlove et al., 2010;Tcherkez et al., 2017;da Fonseca-Pereira et al., 2021).

Does a stored citrate pool really contribute to the mitochondrial TCA cycle activity in *Brassica*
***napus* leaves in a light/dark-dependent manner ?**

The hypothesis about the role of night-stored citrate molecules was initially proposed based on ^{13}CO$_2$
labeling incorporation in detached leaves of *Xanthium stumarium* in order to fit a model based on
"transmission coefficients" (Tcherkez et al., 2009). A second work combined ^{13}CO$_2$ with ^{5}NH$_4$^{15}NO$_3$
labeling experiments in *B. napus* leaves (petiole-feeding assay) to demonstrate that newly assimilated
^{15}N was mostly fixed on "old" ^{12}C carbon and that mean ^{13}C enrichment of citrate was higher after a
light/dark cycle (Gauthier et al., 2010). However, it was not clear to which extent this contribution
interfered with the mitochondrial pool of citrate and no quantification of the absolute amount of citrate
was done. In addition, a flux balance analysis using a diel model suggested that this stored citrate
contribution was explained by the need to support continued export of sugar and amino acids from the
leaf during the night and to meet cellular maintenance costs (Cheung et al., 2014). However, previous
NMR-based works used both detached leaves to propose the contribution of the stored citrate, which
is an experimental setup that necessarily stops sink/source relationships (sugar and amino acids
exports) (Gauthier et al., 2010). Here, using *B. napus* leaf discs and a U-^{13}C-pyruvate labeling
incorporation, we showed that the M2/(M0+M2) ratio of the acetyl-CoA active pool was close to 0.4
after 4 to 6 hours in our experiments. This clearly suggested that most of M0 isotopologues from
mitochondrial active pools of TCA cycle metabolites came from M0 acetyl-CoA and M0 oxaloacetate
and not a stored pool of ^{12}C-citrate (**Figures 6B,8**). Given the important production of PEPc-derived
molecules with non-labeled PEP in our experiments, ^{12}C-pyruvate necessarily arisen from glycolysis
thereby diluting the mitochondrial U-^{13}C-pyruvate pool (nearly 20% of M0 still detected for the
pyruvate pool at the leaf level). Nevertheless, the concept of stored citrate could also rely on the light-
dependent activation of ATP citrate lyase, by converting stored ^{12}C-citrate pool into ^{12}C-oxaloacetate
and ^{12}C-acetyl-CoA (Sweetlove et al., 2010;da Fonseca-Pereira et al., 2021). In our experiments the
^{13}C-enrichment of acetyl-CoA was indirectly estimated from ^{13}C-labeled fragments of either succinate
or glutamate, *i.e.* this measurement could not capture a potential contribution of stored ^{12}C-citrate to
^{12}C-acetyl-CoA. But if such contribution was relatively important in the light and negligible in the dark
(concept of the stored citrate contribution), it would necessarily introduce a ^{12}C-isotopic dilution for
mean ^{13}C enrichment of glutamate and succinate in the light compared to the dark (Gauthier et al.,
2010). Surprisingly, this isotopic dilution was not observed here while the system was in a metabolic
steady-state, thus ruling out a possible bypass with ATP citrate lyase (**Figure 4**). Besides this, we did
not observe any significant modification of the citrate pool during the 6 hours of our experiment
(**Figure 4B, Table S4**) while a strong decrease in light conditions and a strong increase in dark
conditions were expected with the concept of the stored citrate contribution (Cheung et al., 2014). One
may question the use of pyruvate instead of CO$_2$ (previous works) since pyruvate incorporation may
have artificially increased PEPc activity, leading to an important import of malate into mitochondria
that stimulate the export of citrate rather than its import through the dicarboxilate carrier 2 (DIC2) (Lee
et al., 2021). This could in turn partially explain why the proportion of molecules that experienced a
second "turn" of the TCA cycle was relatively weak for citrate, glutamate and succinate compared to
malate in our experiments. Nevertheless, it seems more plausible that stored citrate contribution to
glutamate biosynthesis was fully decoupled from mitochondrial pools to explain our results (**Figure 8**)

(Sweetlove et al., 2010). Overall, our results casted serious doubt about the contribution of stored
citrate to the mitochondrial TCA cycle in *B. napus* leaves.

The TCA cycle partially works in a cyclic mode under light conditions in *B. napus* leaves

The non-cyclic nature of the TCA cycle in illuminated leaves has been experimentally demonstrated
from works that used a 1-hour incorporation of $^{13}CO_2$ to either detached or attached leaves of different
plant species (Tcherkez et al., 2009;Xu et al., 2021;Xu et al., 2022). Only one work used a 2-hour
incorporation of $^{13}C_1$-pyruvate and $^{13}C_3$-pyruvate (Tcherkez et al., 2009), but the major argument for
the non-cyclic nature was questionable given that the reassimilation of $^{13}CO_2$ arising from $^{13}C_1$-
pyruvate decarboxylation by PEPc cannot be metabolically transmitted from malate to succinate since
the C1 and the C4 of the oxaloacetate backbone are lost during citrate to succinate conversion (**Figure**
6A) (Alves et al., 2015;Lima et al., 2021). Overall, the two INST-MFA approaches showed that PEPc
flux and the citrate-to-2-oxoglutarate conversion experienced both relatively similar rates (ratio close
to 1), while the flux from 2-oxoglutarate to succinate and from oxaloacetate to malate were both equal
to 0 in Arabidopsis and *Camelina sativa* (Xu et al., 2021;Xu et al., 2022). Here, after 2 hours of labeling
with U-^{13}C-pyruvate, we observed a significant production of TCA cycle-derived isotopologues in
citrate, glutamate, succinate and malate fragments followed by an increase of other isotopologues from
2 to 6 hours that corresponded to labeled molecules experiencing a second "turn" of the TCA cycle.
Our results provided for the first time in plants a unique ^{13}C-labeling signature demonstrating that the
TCA cycle can work in a cyclic mode under light conditions in *B. napus* leaves (**Figures 5,6,8**).
Therefore, how can we explain discrepancies compared to previous works ? Perhaps, previous works
may have missed this cyclic mode due to a shorter time of ^{13}C incorporation, but also due to the ^{13}C
labeling strategy. Indeed, pyruvate can be more rapidly metabolized through the TCA cycle compared
to CO_2. In addition, $^{13}CO_2$ could be quickly assimilated by the PEPc thanks to the anhydrase carbonic
while the $^{13}CO_2$ fixed by the RuBiSCO will require more time to be transmitted to PEP and pyruvate
(See the dynamics of CIDs for trioses phosphates, alanine, citrate and aspartate therein (Xu et al.,
2021)). In this scenario, ^{13}C enrichment of citrate and glutamate will be artificially higher than for
succinate (C1 and the C4 of the oxaloacetate backbone are lost during citrate to succinate conversion
(**Figure 5A**)). Conversely, it is also important to question our labeling strategy (10 mM U-^{13}C-pyruvate
incorporations) since it could artificially increase TCA cycle turn-over compared to experiments with
$^{13}CO_2$ by eliminating the glycolytic regulation step, leading for example to the rapid apparition of M1
and M3 isotopologues for Glutamate_2TMS_C2C5 and Succinate_2TMS_C1C4. A recent
incorporation of 15 mM U-^{13}C-pyruvate or 15 mM U-^{13}C-glucose into Arabidopsis with a petiole-
feeding assay during 4 hours in the light identified important ^{13}C enrichments in citrate, glutamate and
succinate but the CIDs were not shown (Zhang et al., 2021). However, another experiment
(Arabidopsis leaf discs, 20 mM U-^{13}C-pyruvate, up to 8 hours in the dark) showed similar results but
only observed M2 isotopologues for citrate, glutamate and succinate and not the M1 and M3
isotopologues that were observed in our experiments (complete dataset available on request from the
authors) (Le et al., 2021). Against these observations for Arabidopsis, it is evident that the important
commitment of TCA cycle-derived malate towards another "turn" of the TCA cycle is specific to *B.*
napus (**Figure 8**). Therefore, part of the discrepancies between our results and previous works may
also be explained by the use of different plant species. Since the non-cyclic nature of the TCA cycle in
the light was mostly explained by the need to sustain *de novo* nitrogen assimilation (Gauthier et al.,
2010), the analysis of 2-oxoglutarate molecule escaping the mitochondrial TCA cycle should stimulate
particular attention in ^{13}C labeling experiments in order to compare the rates of cyclic and non-cyclic
flux modes (**Figure 8**). For this purpose, the potential role of gamma-aminobutyric acid shunt and
glutamate dehydrogenase in rerouting glutamate towards either 2-oxoglutarate or succinate may have

to be considered regarding some important studies (Tcherkez et al., 2009;Terce-Laforgue et al.,
2013;Nissen et al., 2015).

Light/dark regulation of PEPc and TCA cycle fluxes in *Brassica napus* leaves

PEPc activity is positively regulated by its light-dependent thioredoxin-mediated phosphorylation in
C_3 and C_4 plants (Caburatan and Park, 2021). Consequently, PEPc is expected to be more active in the
light than in the dark, as recently suggested in guard cell cultures based on the differential ^{13}C
enrichment found for the C1 of glutamate (Lima et al., 2021). Conversely, TCA cycle is inhibited in
light conditions compared to dark conditions, notably due to the inhibition of the PDC and the
inhibition of TCA cycle dehydrogenases by the photorespiratory activity of the mitochondrial GDC
(Tcherkez et al., 2005;Timm and Hagemann, 2020). Here, we took advantage of pathway specific
isotopologue/isotopomer to calculate fractional contribution (FC) of TCA cycle and PEPc for pyruvate-
to-malate and pyruvate-to-citrate. We showed that the contribution of PEPc-dependent reassimilation
of TCA cycle and PDC-derived CO_2 was relatively low compared to the TCA cycle for malate (4:1
ratio) but was positively regulated by the light condition (1.5-fold increase), reflecting the strong shift
for M1/M2 isotopologues of Malate_3TMS_C1C4 in light conditions compared to dark conditions
(**Figures 6,8**). Considering the relative nature of pathway FC, this shift could be attributed to both
increased PEPc activity and TCA cycle inhibition in the light. However, in our experiments, we found
very similar dynamics of ^{13}C-enrichment for TCA cycle metabolites in both dark and light conditions
combined to similar metabolic pool sizes in a metabolic steady-state (**Figure 4**). Thus, variations of
TCA cycle metabolic fluxes between light and dark conditions are expected to be very low. This may
be attributed to the artificial incorporation of pyruvate that eliminated the PDC-dependent regulation
of carbon entry for plant TCA cycle. Nevertheless, the importance of this regulation for carbon entry
towards the TCA cycle can also be questioned. A previous work based on $^{13}CO_2$ decarboxylations with
either ^{13}C-pyruvate or ^{13}C-glucose showed that PDC was inhibited by only 27% in the light compared
to the dark, while glycolysis could be inhibited by up to 95% (Tcherkez et al., 2005). Recently, a U-
^{13}C-pyruvate incorporation into Arabidopsis leaves during 4 hours in the light showed that the *pdk*
mutant (constant activation of the PDC) had higher ^{13}C-enrichments for citrate, glutamate and succinate
compared to the control but these differences were not observed when using U-^{13}C-glucose (Zhang et
al., 2021). Based on these observations, we think that pyruvate incorporation essentially eliminated the
light/dark regulation of glycolysis in our experiments. Thus, our ^{13}C-labeling strategy can only be used
to probe light/dark regulation of PEPc activity, through the reassimilation of TCA cycle and PDC-
derived CO_2. To which extent this PEPc-dependent reassimilation is close to the total PEPc flux
remains an important question to address for a complete comparison and TCA cycle and PEPc fluxes.
For instance, the two ^{13}C-INST-MFA performed in plant leaves with $^{13}CO_2$ obtained a 1:1 ratio with
only non-cyclic flux mode for the TCA cycle. A^{13}C-MFA performed at isotopic and metabolic steady-
state in heterotrophic suspension cells culture incubated with U-^{13}C-glucose showed a similar ratio of
2:1 for forward TCA cycle/PEPc metabolic fluxes (ana1 versus tca2, tca3, tca4, tca5), with a negligible
malic enzyme activity (10-fold less than PEPc) (Masakapalli et al., 2010). Whether the total PEPc flux
is higher than the cyclic mode of the TCA cycle for both malate and citrate in *B. napus* leaves will have
to be addressed using complementary ^{13}C-labeling strategy, such U-^{13}C-glucose or U-^{13}C-sucrose, or
with a full $^{13}CO_2$-based INST-MFA with longer periods of labeling compared to previous works.
Besides this, we observed an unexpected higher contribution of PEPc derived-carbons to citrate in the
dark condition compared to the light condition. Since we have shown that PEPc contribution to malate
was reduced in the dark compared to the light, this result could reflect a higher transport of
oxaloacetate/malate from the cytosol to the mitochondria in dark conditions. This idea is supported by
the fact that PEPc is majorly localized to the cytosol (Caburatan and Park, 2021) and the recent
identification of a mitochondrial malate importer in Arabidopsis with an important impact on

mitochondrial respiration in the dark (Lee et al., 2021). The import/export of malate and citrate in
mitochondria should also be considered in future ^{13}C-MFA of the TCA cycle.

Conclusions

Overall, we proposed a robust workflow to evaluate the accuracy of CID measurements by GC-MS for
organic and amino acids and successfully applied our validated method to the investigation of TCA
cycle and PEPc metabolic fluxes in *Brassica napus* leaves using a ^{13}C-MFA concept. Our results did
not support the hypothesis of the stored citrate contribution to mitochondrial TCA cycle in a light/dark-
dependent manner and demonstrated that the forward cyclic flux mode of the TCA cycle can operate
under light conditions, leading to nonnegligible commitment of *de novo* biosynthesized TCA cycle
intermediates to another "turn" of the cycle (**Figure 8**). We also showed that pathway fractional
contribution could be used to probe the light/dark regulation of PEPc-dependent reassimilation of
mitochondrial decarboxylations. Notably, the reassimilation of PDC and TCA-cycle derived CO_2 by
photosynthesis represented nearly 5% of the net CO_2 assimilation rate in our conditions. The next stage
of this work could be focused on the accurate modeling of this metabolic network to infer absolute
metabolic fluxes. This perspective sounds stimulating given the development of multiple modeling
approaches over the last decade but still remains highly challenging (Kruger and Ratcliffe, 2021).

5   Conflict of Interest

724 The authors declare that the research was conducted in the absence of any commercial or financial
725 relationships that could be construed as a potential conflict of interest.

6   Author Contributions

727 Conceptualization, Y.D., A.B.; methodology, Y.D..; software, Y.D., O.F.; validation, Y.D., S.B. and
728 C.B.; formal analysis, Y.D.; investigation, Y.D., S.B. and C.B.; data curation, Y.D.; writing-original
729 draft preparation, Y.D.; writing-review and editing, Y.D., A.B.; visualization, Y.D.; supervision,
730 Y.D.; project administration, Y.D.; funding acquisition, Y.D., A.B. All authors have read and agreed
731 to the published version of the manuscript.

7   Funding

733 This research was supported by the IB2021_TRICYCLE project funded by the INRAE BAP division
734 and by the METABOHUB project funded by the program Investments for the Future of the French
735 national agency for research (ANR grant number 11-INBS-0010).

8   Acknowledgments

P2M2 (Metabolic Profiling & Metabolomics Platform, Le Rheu, France, www6.inrae.fr/p2m2/) and its
staff members are gratefully acknowledged for access to mass spectrometry facilities and for
supporting the methodological developments required for this study. We thank MetaToul
(Metabolomics & Fluxomics Facitilies, Toulouse, France, www.metatoul.fr) for technical support and
access to NMR facilities. We are grateful to Nathalie Marnet and Mathieu Aubert for technical
assistance in the quantification of amino acid content and Pierre Millard and Floriant Bellvert for
fruitful scientific discussions.

9   Supplementary Material

745   Supplementary Material is available upon request (issues with uploading on the biorxiv server).

746   **10   References**

747   Abadie, C., Lothier, J., Boex-Fontvieille, E., Carroll, A., and Tcherkez, G. (2017). Direct assessment
748   of the metabolic origin of carbon atoms in glutamate from illuminated leaves using (13) C-
749   NMR. *New Phytol* 216**,** 1079-1089.

750   Abadie, C., and Tcherkez, G. (2019). In vivo phosphoenolpyruvate carboxylase activity is controlled
751   by CO2 and O2 mole fractions and represents a major flux at high photorespiration rates. *New*
752   *Phytol* 221**,** 1843-1852.

753   Abadie, C., and Tcherkez, G. (2021). (13)C Isotope Labelling to Follow the Flux of Photorespiratory
754   Intermediates. *Plants (Basel)* 10.

755   Allen, D.K., and Young, J.D. (2019). Tracing metabolic flux through time and space with isotope
756   labeling experiments. *Curr Opin Biotechnol* 64**,** 92-100.

757   Alves, T.C., Pongratz, R.L., Zhao, X., Yarborough, O., Sereda, S., Shirihai, O., Cline, G.W., Mason,
758   G., and Kibbey, R.G. (2015). Integrated, Step-Wise, Mass-Isotopomeric Flux Analysis of the
759   TCA Cycle. *Cell Metab* 22**,** 936-947.

760   Antoniewicz, M.R. (2021). A guide to metabolic flux analysis in metabolic engineering: Methods,
761   tools and applications. *Metab Eng* 63**,** 2-12.

762   Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2006). Determination of confidence
763   intervals of metabolic fluxes estimated from stable isotope measurements. *Metab Eng* 8**,** 324-
764   337.

765   Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2007). Accurate Assessment of Amino
766   Acid Mass Isotopomer Distributions for Metabolic Flux Analysis. *Anal Chem* 79**,** 6.

767   Araujo, W.L., Nunes-Nesi, A., Nikoloski, Z., Sweetlove, L.J., and Fernie, A.R. (2012). Metabolic
768   control and regulation of the tricarboxylic acid cycle in photosynthetic and heterotrophic plant
769   tissues. *Plant Cell Environ* 35**,** 1-21.

770   Araujo, W.L., Tohge, T., Nunes Nesi, A., Obata, T., and Fernie, A. (2014). "Analysis of Kinetic
771   Labeling of Amino Acids and Organic Acids by GC-MS," in *Plant Metabolic Flux Analysis:*
772   *Methods and Protocols,* ed. M.I.M. Biology.).

773   Bianchetti, G., Baron, C., Carrillo, A., Berardocco, S., Marnet, N., Wagner, M.H., Demilly, D.,
774   Ducournau, S., Manzanares-Dauleux, M.J., Caherec, F.L., Buitink, J., and Nesi, N. (2021).
775   Dataset for the metabolic and physiological characterization of seeds from oilseed rape
776   (Brassica napus L.) plants grown under single or combined effects of drought and clubroot
777   pathogen Plasmodiophora brassicae. *Data Brief* 37**,** 107247.

778   Buescher, J.M., Antoniewicz, M.R., Boros, L.G., Burgess, S.C., Brunengraber, H., Clish, C.B.,
779   Deberardinis, R.J., Feron, O., Frezza, C., Ghesquiere, B., Gottlieb, E., Hiller, K., Jones, R.G.,
780   Kamphorst, J.J., Kibbey, R.G., Kimmelman, A.C., Locasale, J.W., Lunt, S.Y., Maddocks,
781   O.D., Malloy, C., Metallo, C.M., Meuillet, E.J., Munger, J., Noh, K., Rabinowitz, J.D.,
782   Ralser, M., Sauer, U., Stephanopoulos, G., St-Pierre, J., Tennant, D.A., Wittmann, C., Vander
783   Heiden, M.G., Vazquez, A., Vousden, K., Young, J.D., Zamboni, N., and Fendt, S.M. (2015).
784   A roadmap for interpreting (13)C metabolite labeling patterns from cells. *Curr Opin*
785   *Biotechnol* 34**,** 189-201.

Caburatan, L., and Park, J. (2021). Differential Expression, Tissue-Specific Distribution, and
Posttranslational Controls of Phosphoenolpyruvate Carboxylase. *Plants (Basel)* 10.

Cheah, Y.E., Xu, Y., Sacco, S.A., Babele, P.K., Zheng, A.O., Johnson, C.H., and Young, J.D. (2020).
Systematic identification and elimination of flux bottlenecks in the aldehyde production
pathway of Synechococcus elongatus PCC 7942. *Metab Eng* 60**,** 56-65.

Cheah, Y.E., and Young, J.D. (2018). Isotopically nonstationary metabolic flux analysis (INST-
MFA): putting theory into practice. *Curr Opin Biotechnol* 54**,** 80-87.

Cheung, C.Y., Poolman, M.G., Fell, D.A., Ratcliffe, R.G., and Sweetlove, L.J. (2014). A Diel Flux
Balance Model Captures Interactions between Light and Dark Metabolism during Day-Night
Cycles in C3 and Crassulacean Acid Metabolism Leaves. *Plant Physiol* 165**,** 917-929.

Chu, K.L., Koley, S., Jenkins, L.M., Bailey, S.R., Kambhampati, S., Foley, K., Arp, J.J., Morley,
S.A., Czymmek, K.J., Bates, P.D., and Allen, D.K. (2022). Metabolic flux analysis of the
non-transitory starch tradeoff for lipid production in mature tobacco leaves. *Metab Eng* 69**,**
231-248.

Clark, T.J., Guo, L., Morgan, J., and Schwender, J. (2020). Modeling Plant Metabolism: From
Network Reconstruction to Mechanistic Models. *Annu Rev Plant Biol*.

Da Fonseca-Pereira, P., Souza, P.V.L., Fernie, A.R., Timm, S., Daloso, D.M., and Araujo, W.L.
(2021). Thioredoxin-mediated regulation of (photo)respiration and central metabolism. *J Exp*
Bot 72**,** 5987-6002.

Daloso, D.M., Medeiros, D.B., Dos Anjos, L., Yoshida, T., Araujo, W.L., and Fernie, A.R. (2017).
Metabolism within the specialized guard cells of plants. *New Phytol* 216**,** 1018-1033.

Dellero, Y. (2020). Manipulating amino acid metabolism to improve crop nitrogen use efficiency for
a sustainable agriculture. *Frontiers in Plant Science* 11**,** 1857

Dellero, Y., Clouet, V., Marnet, N., Pellizzaro, A., Dechaumet, S., Niogret, M.F., and Bouchereau,
A. (2020a). Leaf status and environmental signals jointly regulate proline metabolism in
winter oilseed rape. *J Exp Bot* 71**,** 2098-2111.

Dellero, Y., and Filangi, O. (2021). "Corrective method dedicated to Isocor for calculating carbon
isotopologue distribution from GCMS runs". V1 ed.: Portail Data INRAE).

Dellero, Y., Heuillet, M., Marnet, N., Bellvert, F., Millard, P., and Bouchereau, A. (2020b).
Sink/Source Balance of Leaves Influences Amino Acid Pools and Their Associated Metabolic
Fluxes in Winter Oilseed Rape (Brassica napus L.). *Metabolites* 10**,** 16.

Dellero, Y., Jossier, M., Glab, N., Oury, C., Tcherkez, G., and Hodges, M. (2016). Decreased
glycolate oxidase activity leads to altered carbon allocation and leaf senescence after a
transfer from high CO2 to ambient air in Arabidopsis thaliana. *J Exp Bot* 67**,** 3149-3163.

Des Rosiers, C., Lloyd, S., Comte, B., and Chatham, J.C. (2004). A critical perspective of the use of
(13)C-isotopomer analysis by GCMS and NMR as applied to cardiac metabolism. *Metab Eng*
6**,** 44-58.

Dong, H., Bai, L., Zhang, Y., Zhang, G., Mao, Y., Min, L., Xiang, F., Qian, D., Zhu, X., and Song,
C.P. (2018). Modulation of Guard Cell Turgor and Drought Tolerance by a Peroxisomal
Acetate-Malate Shunt. *Mol Plant* 11**,** 1278-1291.

Evers, B., Gerding, A., Boer, T., Heiner-Fokkema, M.R., Jalving, M., Wahl, S.A., Reijngoud, D.J.,
and Bakker, B.M. (2021). Simultaneous Quantification of the Concentration and Carbon

828   Isotopologue Distribution of Polar Metabolites in a Single Analysis by Gas Chromatography
829   and Mass Spectrometry. *Anal Chem* 93**,** 8248-8256.

830 Galili, G., Amir, R., and Fernie, A.R. (2016). The Regulation of Essential Amino Acid Synthesis and
831   Accumulation in Plants. *Annu Rev Plant Biol* 67**,** 153-178.

832 Gauthier, P.P., Bligny, R., Gout, E., Mahe, A., Nogues, S., Hodges, M., and Tcherkez, G.G. (2010).
833   In folio isotopic tracing demonstrates that nitrogen assimilation into glutamate is mostly
834   independent from current CO2 assimilation in illuminated leaves of Brassica napus. *New*
835   *Phytol* 185**,** 988-999.

836 Harvey, D.J., and Vouros, P. (2020). Mass Spectrometric Fragmentation of Trimethylsilyl and
837   Related Alkylsilyl Derivatives. *Mass Spectrom Rev* 39**,** 105-211.

838 Heuillet, M., Bellvert, F., Cahoreau, E., Letisse, F., Millard, P., and Portais, J.C. (2018).
839   Methodology for the Validation of Isotopic Analyses by Mass Spectrometry in Stable-Isotope
840   Labeling Experiments. *Anal Chem* 90**,** 1852-1860.

841 Heuillet, M., Millard, P., Cisse, M.Y., Linares, L.K., Letisse, F., Manie, S., Le Cam, L., Portais, J.C.,
842   and Bellvert, F. (2020). Simultaneous Measurement of Metabolite Concentration and Isotope
843   Incorporation by Mass Spectrometry. *Anal Chem*.

844 Kambhampati, S., Li, J., Evans, B.S., and Allen, D.K. (2019). Accurate and efficient amino acid
845   analysis for protein quantification using hydrophilic interaction chromatography coupled
846   tandem mass spectrometry. *Plant Methods* 15**,** 46.

847 Kruger, N.J., Huddleston, J.E., Le Lay, P., Brown, N.D., and Ratcliffe, R.G. (2007). Network flux
848   analysis: impact of 13C-substrates on metabolism in Arabidopsis thaliana cell suspension
849   cultures. *Phytochemistry* 68**,** 2176-2188.

850 Kruger, N.J., and Ratcliffe, R.G. (2021). Whither metabolic flux analysis in plants? *Journal of*
851   *Experimental Botany*.

852 Lai, Z., and Fiehn, O. (2018). Mass spectral fragmentation of trimethylsilylated small molecules.
853   *Mass Spectrom Rev* 37**,** 245-257.

854 Le, X.H., Lee, C.P., and Millar, A.H. (2021). The mitochondrial pyruvate carrier (MPC) complex
855   mediates one of three pyruvate-supplying pathways that sustain Arabidopsis respiratory
856   metabolism. *Plant Cell* 33**,** 2776-2793.

857 Le, X.H., Lee, C.P., Monachello, D., and Millar, A.H. (2022). Metabolic evidence for distinct
858   pyruvate pools inside plant mitochondria. *Nat Plants*.

859 Lee, C.P., Elsasser, M., Fuchs, P., Fenske, R., Schwarzlander, M., and Millar, A.H. (2021). The
860   versatility of plant organic acid metabolism in leaves is underpinned by mitochondrial malate-
861   citrate exchange. *Plant Cell* 33**,** 3700-3720.

862 Lima, V.F., Erban, A., Daubermann, A.G., Freire, F.B.S., Porto, N.P., Candido-Sobrinho, S.A.,
863   Medeiros, D.B., Schwarzlander, M., Fernie, A.R., Dos Anjos, L., Kopka, J., and Daloso,
864   D.M. (2021). Establishment of a GC-MS-based (13) C-positional isotopomer approach
865   suitable for investigating metabolic fluxes in plant primary metabolism. *Plant J*.

866 Lima, V.F., Perez Souza, L., Williams, T.C., Fernie, A., and Daloso, D.M. (2018). "Gas
867   Chromatography–Mass Spectrometry-Based 13C-Labeling Studies in Plant Metabolomics,"
868   in *Plant Metabolomics: Methods and Protocol,* ed. M.I.M. Biology.).

Ma, F., Jazmin, L.J., Young, J.D., and Allen, D.K. (2014). Isotopically nonstationary 13C flux
analysis of changes in Arabidopsis thaliana leaf metabolism due to high light acclimation.
Proc Natl Acad Sci U S A 111**,** 16967-16972.

Masakapalli, S.K., Bryant, F.M., Kruger, N.J., and Ratcliffe, R.G. (2014). The metabolic flux
phenotype of heterotrophic Arabidopsis cells reveals a flexible balance between the cytosolic
and plastidic contributions to carbohydrate oxidation in response to phosphate limitation.
Plant J 78**,** 964-977.

Masakapalli, S.K., Le Lay, P., Huddleston, J.E., Pollock, N.L., Kruger, N.J., and Ratcliffe, R.G.
(2010). Subcellular flux analysis of central metabolism in a heterotrophic Arabidopsis cell
suspension using steady-state stable isotope labeling. *Plant Physiol* 152**,** 602-619.

Medeiros, D.B., Perez Souza, L., Antunes, W.C., Araujo, W.L., Daloso, D.M., and Fernie, A.R.
(2018). Sucrose breakdown within guard cells provides substrates for glycolysis and
glutamine biosynthesis during light-induced stomatal opening. *Plant J* 94**,** 583-594.

Millard, P., Cahoreau, E., Heuillet, M., Portais, J.C., and Lippens, G. (2017). (15)N-NMR-Based
Approach for Amino Acids-Based (13)C-Metabolic Flux Analysis of Metabolism. *Anal Chem*
89**,** 2101-2106.

Millard, P., Delepine, B., Guionnet, M., Heuillet, M., Bellvert, F., and Letisse, F. (2019). IsoCor:
isotope correction for high-resolution MS labeling experiments. *Bioinformatics* 35**,** 4484-
4487.

Millard, P., Massou, S., Portais, J.C., and Letisse, F. (2014). Isotopic studies of metabolic systems by
mass spectrometry: using Pascal's triangle to produce biological standards with fully
controlled labeling patterns. *Anal Chem* 86**,** 10288-10295.

Millard, P., Schmitt, U., Kiefer, P., Vorholt, J.A., Heux, S., and Portais, J.-C. (2020). ScalaFlux: a
scalable approach to quantify fluxes in metabolic subnetworks. *Plos Computational Biology*
16**,** e1007799.

Nissen, J.D., Pajecka, K., Stridh, M.H., Skytt, D.M., and Waagepetersen, H.S. (2015). Dysfunctional
TCA-Cycle Metabolism in Glutamate Dehydrogenase Deficient Astrocytes. *Glia* 63**,** 2313-
2326.

Okahashi, N., Kawana, S., Iida, J., Shimizu, H., and Matsuda, F. (2019). Fragmentation of
Dicarboxylic and Tricarboxylic Acids in the Krebs Cycle Using GC-EI-MS and GC-EI-
MS/MS. *Mass Spectrom (Tokyo)* 8**,** A0073.

Ratcliffe, R.G., and Shachar-Hill, Y. (2006). Measuring multiple fluxes through plant metabolic
networks. *Plant J* 45**,** 490-511.

Rstudio Team (2016). RStudio: Integrated Development for R.

.

Sweetlove, L.J., Beard, K.F., Nunes-Nesi, A., Fernie, A.R., and Ratcliffe, R.G. (2010). Not just a
circle: flux modes in the plant TCA cycle. *Trends Plant Sci* 15**,** 462-470.

Szecowka, M., Heise, R., Tohge, T., Nunes-Nesi, A., Vosloh, D., Huege, J., Feil, R., Lunn, J.,
Nikoloski, Z., Stitt, M., Fernie, A.R., and Arrivault, S. (2013). Metabolic fluxes in an
illuminated Arabidopsis rosette. *Plant Cell* 25**,** 694-714.

Tcherkez, G., Cornic, G., Bligny, R., Gout, E., and Ghashghaie, J. (2005). In vivo respiratory
metabolism of illuminated leaves. *Plant Physiol* 138**,** 1596-1606.

Tcherkez, G., Gauthier, P., Buckley, T.N., Busch, F.A., Barbour, M.M., Bruhn, D., Heskel, M.A.,
Gong, X.Y., Crous, K.Y., Griffin, K., Way, D., Turnbull, M., Adams, M.A., Atkin, O.K.,
Farquhar, G.D., and Cornic, G. (2017). Leaf day respiration: low CO2 flux but high
significance for metabolism and carbon balance. *New Phytol* 216**,** 986-1001.

Tcherkez, G., Mahe, A., Gauthier, P., Mauve, C., Gout, E., Bligny, R., Cornic, G., and Hodges, M.
(2009). In folio respiratory fluxomics revealed by 13C isotopic labeling and H/D isotope
effects highlight the noncyclic nature of the tricarboxylic acid "cycle" in illuminated leaves.
Plant Physiol 151**,** 620-630.

Terce-Laforgue, T., Bedu, M., Dargel-Grafin, C., Dubois, F., Gibon, Y., Restivo, F.M., and Hirel, B.
(2013). Resolving the role of plant glutamate dehydrogenase: II. Physiological
characterization of plants overexpressing the two enzyme subunits individually or
simultaneously. *Plant Cell Physiol* 54**,** 1635-1647.

Timm, S. (2020). The impact of photorespiration on plant primary metabolism through metabolic and
redox regulation. *Biochemical Society Transactions* 48**,** 2495-2504.

Timm, S., and Hagemann, M. (2020). Photorespiration - how is it regulated and regulates overall
plant metabolism? *J Exp Bot*.

Treves, H., Kuken, A., Arrivault, S., Ishihara, H., Hoppe, I., Erban, A., Hohne, M., Moraes, T.A.,
Kopka, J., Szymanski, J., Nikoloski, Z., and Stitt, M. (2022). Carbon flux through
photosynthesis and central carbon metabolism show distinct patterns between algae, C3 and
C4 plants. *Nat Plants* 8**,** 78-91.

Xu, G., Fan, X., and Miller, A.J. (2012). Plant nitrogen assimilation and use efficiency. *Annu Rev
Plant Biol* 63**,** 153-182.

Xu, Y., Fu, X., Sharkey, T.D., Shachar-Hill, Y., and Walker, B.J. (2021). The metabolic origins of
non-photorespiratory CO2 release during photosynthesis: A metabolic flux analysis. *Plant
Physiol*.

Xu, Y., Wieloch, T., Kaste, J.a.M., Shachar-Hill, Y., and Sharkey, T.D. (2022). Reimport of carbon
from cytosolic and vacuolar sugar pools into the Calvin–Benson cycle explains
photosynthesis labeling anomalies. *Proceedings of the National Academy of Sciences* 119.

Young, J.D., Allen, D.K., and Morgan, J.A. (2014). "Isotopomer measurement techniques in
metabolic flux analysis II: mass spectrometry," in *Methods Mol Biol*.), 85-108.

Zhang, Y., and Fernie, A.R. (2018). On the role of the tricarboxylic acid cycle in plant productivity. *J
Integr Plant Biol* 60**,** 1199-1216.

Zhang, Y., Giese, J., Kerbler, S.M., Siemiatkowska, B., Perez De Souza, L., Alpers, J., Medeiros,
D.B., Hincha, D.K., Daloso, D.M., Stitt, M., Finkemeier, I., and Fernie, A.R. (2021). Two
mitochondrial phosphatases, PP2c63 and Sal2, are required for posttranslational regulation of
the TCA cycle in Arabidopsis. *Mol Plant* 14**,** 1104-1118.

Table 1. Molecular ions selected for Carbon Isotopologue Distribution analysis of organic and amino
acid TMS-derivatives by GC-MS.

Compounds	TMS derivative		Loss		Molecular ion		
	Formula	mass	Formula	mass	Formula	m/z	Carbon skeleton
Organic acids							
Citrate(4TMS)	C18H40O7Si4	480	CH3	15	C17H37O7Si4	465	1-2-3-4-5-6
Citrate(4TMS)	C18H40O7Si4	480	TMS-CO2, TMS-OH	207	C11H21O4Si2	273	1-2-3-4-5
Fumarate(2TMS)	C10H20O4Si2	260	CH3	15	C9H17O4Si2	245	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	CH3	15	C12H27O5Si3	335	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	TMS-CO2	117	C9H21O3Si2	233	2-3-4
Pyruvate(MEOX)(TMS)	C7H15NO3Si	189	CH3	15	C6H12NO3Si	174	1-2-3
Succinate(2TMS)	C10H22O4Si2	262	CH3	15	C9H19O4Si2	247	1-2-3-4
Amino acids							
Asparagine(2TMS)	C10H24N2O3Si2	276	TMS-CO2	117	C6H15N2OSi	159	2-3-4
Asparagine(3TMS)	C13H32N2O3Si3	348	TMS-CO2	117	C9H23N2OSi2	231	2-3-4
Aspartate(2TMS)	C10H23NO4Si2	277	TMS-CO2	117	C6H14NO2Si	160	2-3-4
Aspartate(3TMS)	C13H31NO4Si3	349	TMS-CO2	117	C9H22NO2Si2	232	2-3-4
Glutamate(3TMS)	C14H33NO4Si3	363	TMS-CO2	117	C10H24NO2Si2	246	2-3-4-5
Glutamine(3TMS)	C14H34N2O3Si3	362	TMS-CO2	117	C10H25N2OSi2	245	2-3-4-5
Glycine(3TMS)	C11H29NO2Si3	291	TMS-CO2	117	C7H20NSi2	174	2
Isoleucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Leucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Lysine(4TMS)	C18H46N2O2Si4	434	TMS-CO2	117	C14H37N2Si3	317	2-3-4-5-6
Methionine(2TMS)	C11H27NO2SSi2	293	TMS-CO2	117	C7H18NSSi	176	2-3-4-5
Ornithine(4TMS)	C17H44N2O2Si4	420	none		none		1-2-3-4-5
Phenylalanine(2TMS)	C15H27NO2Si2	309	TMS-CO2	117	C11H18NSi	192	2-3-4-5-6-7-8-9
Phenylalanine(2TMS)	C15H27NO2Si2	309	C7H7	91	C8H20NO2Si2	218	1-2
Proline(2TMS)	C11H25NO2Si2	259	TMS-CO2	117	C7H16NSi	142	2-3-4-5
Serine(2TMS)	C9H23NO3Si2	249	TMS-CO2	117	C5H14NOSi	132	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CO2	117	C8H22NOSi2	204	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CH2O	103	C8H20NO2Si2	218	1-2
Threonine(3TMS)	C13H33NO3Si3	335	TMS-CO2	117	C9H24NOSi2	218	2-3-4
Tryptophan(2TMS)	C17H28N2O2Si2	348	2TMS-C2H2NO2	218	C9H8N	130	3-4-5-6-7-8-9-10-11
Tryptophan(2TMS)	C17H28N2O2Si2	348	C9H8N	130	C8H20NO2Si2	218	1-2
Tryptophan(3TMS)	C20H36N2O2Si3	420	2TMS-C2H2NO2	218	C12H16NSi	202	3-4-5-6-7-8-9-10-11
Tryptophan(3TMS)	C20H36N2O2Si3	420	C12H16NSi	202	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	C10H15OSi	179	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	TMS-CO2	117	C14H26NOSi2	280	2-3-4-5-6-7-8-9
Valine(2TMS)	C11H27NO2Si2	261	TMS-CO2	117	C7H18NSi	144	2-3-4-5
Valine(2TMS)	C11H27NO2Si2	261	C3H7	43	C9H20NO2Si2	218	1-2
α-Alanine(2TMS)	C9H23NO2Si2	233	TMS-CO2	117	C5H14NSi	116	2-3
β-Alanine(3TMS)	C12H31NO2Si3	305	CH3	15	C11H28NO2Si3	290	1-2-3

Figure legends

Figure 1. Workflow used in this study. A GC-MS method was built with abundant molecular ions
deriving from TMS-derivatives of organic and amino acids that contained a large number of
informative carbons. Then this method was evaluated by using: i) chemical standards and tailor-made
^{13}C-standards that had a predictable CID using *E. coli* as a metabolic factory (^{13}C-PT); ii) unlabeled
and ^{13}C-labeled plant samples.

Figure 2. **Evaluation of Carbon Isotopologue Distribution (CID) measurements for organic and**
amino acids by GC-MS using ^{13}C-PT reference samples. **A**, Evaluation of amino acids using ^{13}C-
PT protein hydrolysates. **B**, Evaluation of organic acids using ^{13}C-PT full metabolites extracts. Black
bars correspond to expected CIDs and blue bars to the measured CIDs. The results are presented as the
mean ± SD of four independent biological replicates. Statistical differences between predicted and
measured CID for each isotopologue of each fragment are denoted with asterisks (*) and were
established by considering the 95% confidence intervals of the measured CID. GC-MS fragments are
denoted according to the considered metabolite, its MEOX/TMS-derivatives analyzed and the
metabolite carbon backbone (for example, Glutamate_3TMS_C2C5 means that the fragment comes
from glutamate(3TMS) and contains the carbon C2-C3-C4-C5 (**Table 1**)).

Figure 3. Evaluation of the measurement accuracy for CID and mean ^{13}C enrichment at ^{13}C-PT
and ^{13}C natural abundance in plants. A, CID accuracy and **B**, Mean ^{13}C enrichment for the selected
TMS-derivatives of organic and amino acids using ^{13}C-PT samples. **C**, Residual mean ^{13}C enrichment
measured in chemical standards, Arabidopsis seedlings and *B. napus* leaves after correction for isotope
natural abundance. **D**, Isotopologue-specific biases identified at ^{13}C natural abundance for proline,
valine and threonine fragments. The results are presented as the mean ± SD of four independent
biological replicates or independent preparations for commercial chemical standards.

Figure 4. **Dynamics of U-^{13}C-pyruvate incorporation in *Brassica napus* leaf discs exposed to light**
or dark conditions. **A**, Mean fractional ^{13}C enrichment and **B**, Metabolite pool sizes. Mean fractional
^{13}C enrichment were monitored with GC-MS, while absolute quantification of organic and amino acids
was performed respectively with GC-FID and HPLC-UV. The results are presented as the mean ± SD
of four independent biological replicates. Statistical differences between dark and light conditions are
denoted with asterisks according to a Student test (*, p-value <0.05; **, p-value <0.01; ***, p-value
<0.001). The complete dataset is available in **Table S2, S3**. Supplementary statistics (ANOVA and
post-hoc Tukey-HSD tests) are available in **Table S4, S5**.

Figure 5. **Fractional ^{13}C enrichment at the isotopologue level for TCA-cycle derived metabolites.**
A, Metabolic network considered based on the ^{13}C-labeling strategy. **B**, Light conditions and **C**, Dark
conditions. The M0 isotopologue was not represented in the graphs to facilitate the visualization of
low-enriched isotopologues. The results are presented as the mean ± SD of four independent biological
replicates. The complete dataset is available in **Table S2**. In the proposed network, the second "turn"
reflected the commitment of a molecule that has already experienced all reactions of the TCA cycle to
another round of reactions (Orange and blue labels to compare acetyl-CoA-derived labeling from 1st
and 2nd "turn"). Note that M0 istopologues of all metabolites will still continue to produce M0
istopologues at an important rate after a "turn" of the TCA cycle, given the ^{13}C-enrichment for acetyl-
CoA active pool. PEPc, Phosphoenolpyruvate carboxylase; PDC, Pyruvate dehydrogenase complex,
CS, Citrate synthase; ACO, Aconitase; ICDH, Isocitrate dehydrogenase; OGDH, 2-oxoglutarate
dehydrogenase; SDH, Succinate dehydrogenase complex; FUM, Fumarase; MDH, Malate
dehydrogenase.

Figure 6. **Fractional contribution of TCA cycle and PEPc-dependent reassimilation of**
mitochondrial CO_2 for pyruvate to malate and pyruvate to citrate in light and dark conditions.
A, Pathway-specific mass isotopologues considered for the calculations. **B**, Fractional contribution of
PEPc and TCA cycle to malate and citrate based on pathway-specific mass isotopologues/isotopomers.
For these calculations, only the timepoint T4 and T6 hours were used and pooled (metabolic steady-
state and isotopic steady-state for isotopologue ratios). The results are presented as the mean ± SD of
four independent biological replicates. Statistical comparison of acetyl-CoA enrichment for light and
dark conditions was performed with a Student-test, while fractional contributions were compared with
an ANOVA followed by a post-hoc Tukey-HSD test (different letters indicate groups that were
separated with a p-value <0.05).

Figure 7. **Fractional contribution of PDC and TCA-cycle derived decarboxylations to**
photosynthesis. A, Fractional [13]C enrichment at the isotopologue level for glycine and serine. **B**,
Working hypotheses to explain the photorespiratory-dependent labeling pattern of glycine and serine.
C, Estimation of the fractional contribution of PDC (pyruvate dehydrogenase complex) and TCA-cycle
CO_2 to photosynthesis with Serine (expressed in %). For these calculations, only the timepoint T4 and
T6 hours were used and pooled (metabolic and isotopic steady-state, assumption with [13]C/[12]C of CO_2
released). The results are presented as the mean ± SD of four independent biological replicates.
Statistical comparison of the two serine fragments was performed with a Student-test.

Figure 8. **Proposed flux modes for TCA cycle in *B. napus* leaves under light conditions**. The
contribution of stored citrate is unlikely to interact with mitochondrial TCA cycle. The mitochondrial
TCA cycle operate in both cyclic and non-cyclic flux modes. To which extent 2-oxoglutarate molecules
escape the cycle and imbalance this cyclic/non-cyclic ratio remain to be determined.

1018

Identification of TMS-derivative mass fragments maximizing carbon backbones

Production of reference ¹³C labeled samples with controlled CIDs (¹³C-PT)

Validation of TMS-derivative mass fragments for CID measurements

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

A

Mean fractional ¹³C enrichment

B

Metabolite pool sizes

1034

1035

1036

1037

1038

1039

1040

A

B

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066