Validation of carbon isotopologue distribution measurements by GC-MS and application to [13]C-metabolic flux analysis of the tricarboxylic acid cycle in *Brassica napus* leaves

Younès Dellero[1, 2, 4*], Solenne Berardocco[1, 2], Cécilia Berges[3, 4], Olivier Filangi[1, 2, 4], Alain
Bouchereau[1, 2, 4]

[1]Institute for Genetics, Environment and Plant Protection (IGEPP), National Research Institute for
Agriculture, Food and Environment (INRAE), Université Rennes, Institut Agro , Le Rheu, France

[2]Metabolic profiling and metabolomics platform, Institute for Genetics, Environment and Plant
Protection, Biopolymers Interactions Assemblies, French National Research, Le Rheu, France

[3]TBI, Université de Toulouse, National center for Scientific Research (CNRS), National Institute for
Research for Agriculture, food and Environment (INRAE), National Institute of Applied Sciences
(INSA), Toulouse, France;

[4]MetaToul-MetaboHUB, National infrastructure of metabolomics and fluxomics, Toulouse, France

* Correspondence:
Corresponding Author
younes.dellero@inrae.fr

Keywords: Isotope, gas chromatography-mass spectrometry, trimethylsilyl (TMS) derivatives,
accuracy, regulation, phosphoenolpyruvate carboxylase, decarboxylation, TCA cycle. (Min.5-
Max. 8)

Abstract

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of [13]C-Metabolic Flux Analysis at
isotopically non-stationary steady-state. This approach requires a high level of accuracy for the
measurement of Carbon Isotopologue Distribution in plant metabolites, which has still not been
accurately evaluated at the isotopologue level for GC-MS. Thus, uncertainties remain about the
accuracy of the CID measured and subsequently, the metabolic fluxes associated. Here, we developed
a workflow to validate the measurements of CIDs from plant metabolites with GC-MS by producing
tailor-made *E. coli* standard extracts harboring a predictable binomial CID for some organic and amino
acids. Overall, most of our TMS-derivatives mass fragments were validated with these standards and
at natural isotope abundance in plant matrices. Then, we applied this validated MS method to
investigate the light/dark regulation of plant TCA cycle by incorporating U-[13]C-pyruvate to *Brassica*
napus leaf discs. We took advantage of pathway-specific isotopologues/isotopomers to show that the
TCA cycle can operate in a cyclic manner at high rates in *Brassica napus* under both light and dark
conditions. Notably, this forward cyclic flux mode has a nearly four-fold higher contribution to Citrate
and Malate production than phosphoenolpyruvate carboxylase (PEPc) in our experimental setup. The
light condition did not introduce an isotopic dilution to the dynamic of [13]C-enrichment in Citrate,
Glutamate and Succinate compared to dark conditions and there was no evidence for significant
variations of citrate pools in either light or dark conditions. Our results suggested that the contribution
of stored citrate to the TCA cycle activity was unlikely to occur in *Brassica napus*. Interestingly, there
was a light-dependent [13]C-incorporation into Glycine and Serine which revealed that TCA cycle-

derived CO_2 was actively reassimilated and contributed by up to 5% to net photosynthesis. We also
observed an increase of PEPc contribution to malate production in the light while there was a higher
involvement of PEPc-derived carbons to citrate production in dark condition. This suggested an
important role for Malate import into mitochondria in sustaining TCA cycle in dark conditions.

1   INTRODUCTION

Metabolic fluxes within central metabolism are among the most important determinants of plant cell
physiology, supporting central energetical functions and allowing them to quickly adapt to
environmental changes by controlling metabolite turnover (Clark et al., 2020). Given that most of
fluxes supported by a metabolic network cannot be measured directly, Metabolic Flux Analysis (MFA)
technics were developed to infer information about metabolic fluxes from isotopic labeling
experiments (Ratcliffe and Shachar-Hill, 2006).[13]C-MFA relies on the tracing of [13]C throughout a
metabolic network in a relative metabolic steady-state to model and/or compare metabolic fluxes, *i.e.*
that the metabolic fluxes are stable across the experiment because the associated metabolite pool sizes
are relatively constant over the time of labeling except for source (consumption) and sink (production)
metabolites (Allen and Young, 2019;Antoniewicz, 2021). [13]C-MFA is usually performed at isotopic
steady-state for heterotrophic tissues for practical reasons, and also because they can reach this isotopic
state (Hay et al., 2014). Conversely, [13]C-MFA in phototrophic tissues can only be performed at
isotopically non-stationary steady-state (INST-MFA), thus requiring a higher accuracy in the
measurement of [13]C enrichment within metabolites (Cheah and Young, 2018). Over the last decades,
[13]C-MFA has been applied successfully to microorganisms to highlight new metabolic routes and
identify metabolic bottlenecks for the purpose of metabolic engineering (Cheah et al.,
2020;Antoniewicz, 2021;Treves et al., 2022). In plants, [13]C-INST-MFA has been scarcely applied to
date, due to the complexity in modelling accurately metabolic fluxes in a multicompartmental system
(Heise et al., 2015;Kruger and Ratcliffe, 2021;Wieloch, 2021). However, accurate measurement of
[13]C-enrichment is a prerequisite for precise evaluation of isotopic labeling dynamics and accurate
modelling of metabolic fluxes. Indeed, small errors in mass isotopologue distribution can propagate to
large errors in estimated fluxes (Antoniewicz et al., 2006).

Two major analytical methods are currently employed to analyze the Carbon Isotope Distribution
(CID) within plant metabolites during [13]C-isotope tracing experiments: Nuclear Magnetic Resonance
(NMR) and Mass Spectrometry (MS). NMR analysis of the [13]C enrichment is generally performed
with [13]C-decoupling [1]H sequences and provides information at isotopomer level (by integrating [13]C-
[13]C interactions) (Kruger et al., 2007;Masakapalli et al., 2010;Abadie et al., 2017). Although this
method is quantitative for the [13]C and allows to significantly separate carbon atoms of different
chemical structures, it has a low sensitivity at natural [13]C abundance (large amounts of fresh material
required), thereby affecting the accuracy for [13]C enrichment measurements regarding metabolites
found in small quantities (Dellero et al., 2016;Abadie and Tcherkez, 2021). This method is also time-
consuming due to the longer relaxation time for [13]C nucleus and the need to accumulate many
spectrums to get a good signal (Abadie and Tcherkez, 2021). As a consequence, most of the [13]C-tracing
isotope experiments recently performed in plants and analyzed by [13]C-NMR have only one time-point
(Abadie et al., 2016;Abadie et al., 2018;Abadie and Tcherkez, 2019a;b), which questions the estimation
of metabolic fluxes therein (metabolic/isotopic steady-state and treatment of metabolically inactive
pools). To date, the accurate estimation of metabolic fluxes in phototrophic tissues requires to
determine the isotopic and metabolic state of the system by coupling transient kinetics of [13]C-labeling
with absolute quantification of metabolite pools (Kruger and Ratcliffe, 2015;Cheah and Young,
2018;Clark et al., 2020;Antoniewicz, 2021). Some methodological improvements were recently made
with highly sensitive [1]H-derived sequences to improve the resolution of the [13]C atoms at the isotopomer

level (Sinnaeve et al., 2018), and its future use for [13]C-INST-MFA in plants will provide interesting
perspectives. Conversely, mass spectrometry has been preferentially used to analyze the Carbon
Isotope Distribution (CID) within plant metabolites during time-course [13]C-isotope tracing
experiments (Szecowka et al., 2013;Ma et al., 2014;Medeiros et al., 2018;Treves et al., 2022). This
method is more sensitive at natural [13]C abundance and can be implemented at high throughput.
However, MS essentially gives information at the isotopologue level since isotopomer approaches are
extremely limited by the availability of multiple MS fragments to cover the complete set of all possible
isotopomers. An absolute quantification of the metabolite content is also required in general to
subsequently infer metabolic fluxes. In addition, the analysis of CID for amino acids by LC-MS
revealed some biases and errors in the measurement of CID (Heuillet et al., 2018), thus highlighting
the need to evaluate measurement accuracy with GC-MS fragments for subsequent use in [13]C-MFA of
plants.

Among plant metabolites, organic and amino acids are major players for plant primary metabolism.
They are actively involved in crucial physiological processes such as photorespiration, *de novo*
nitrogen assimilation and mitochondrial respiration (Xu et al., 2012;Zhang and Fernie, 2018;Dellero,
2020;Timm, 2020). Most of the carbon backbones for amino acid biosynthesis come from organic
acids of the tricarboxylic acid (TCA) cycle (Galili et al., 2016). The functioning and regulation of the
plant TCA cycle has been described based on flux balance models (Sweetlove et al., 2010;Araujo et
al., 2012;Cheung et al., 2014) but also based on [13]C-labeling studies with different cells/experimental
setups: attached leaves (Xu et al., 2021), detached leaves (Tcherkez et al., 2009;Gauthier et al.,
2010;Zhang et al., 2021), leaf discs (Le et al., 2021), guard cell cultures (Daloso et al., 2017),
heterotrophic cell cultures (Masakapalli et al., 2010). Overall, most of these works proposed that the
plant TCA cycle is not cyclic in the light, with two branches acting separately: i) a first branch
stimulated by the anaplerotic activity of phosphoenolpyruvate carboxylase (PEPc) leading to the
production of Oxaloacetate, malate and fumarate; ii) a second branch stimulated by the entry of Citrate
leading to the production of isocitrate, 2-oxoglutarate, glutamate and sometimes succinate. These
conclusions were notably supported by the inhibition of the pyruvate dehydrogenase (PDH) complex
and the mitochondrial respiration in the light while nitrogen assimilation was relatively high compared
to dark conditions and required important amounts of 2-oxoglutarate (Tcherkez et al., 2005;Gauthier
et al., 2010). However, the requirement to support continued export of sugar and amino acids from the
leaf during the night and to meet cellular maintenance costs was also invoked (Cheung et al., 2014).
Regarding the metabolic origin of citrate for the second branch in the light, it can come from: i) TCA
cycle *de novo* biosynthesis using Acetyl-CoA derived from either glycolysis and alanine
transamination; ii) a stored citrate pool in a light/dark dependent-manner; iii) the peroxisomal
glyoxylate cycle (Gauthier et al., 2010;Sweetlove et al., 2010;Cheung et al., 2014;Dong et al., 2018;Le
et al., 2021). Based on the NMR analysis of combined [13]CO$_2$ and [15]NH$_4$[15]NO$_3$ labeling experiments in
Brassica napus leaves, it was proposed that a high proportion of non-labeled citrate (vacuolar pool
probably) was involved in the synthesis of 2-oxoglutarate under light conditions to sustain *de novo*
nitrogen assimilation through glutamate (Gauthier et al., 2010). Under this scenario, the citrate pool
would be replenished during dark conditions from glycolysis-derived [13]C-pyruvate, leading to a higher
[13]C enrichment of the citrate after a light/dark cycle. Thus, the light and dark environmental conditions
are expected to introduce important variations of metabolic fluxes within the plant TCA cycle (cyclic
versus non-cyclic) and its associated PEPc anaplerotic pathway. Nevertheless, these variations remain
to be carefully investigated in *B. napus* leaves with a [13]C-MFA framework, *i.e.* with time-course [13]C-
isotope tracing experiments and pool size quantification.

Currently, the routine analysis of CID on plants is often assessed for organic and amino acids by GC-
MS using TMS-derivatives (Szecowka et al., 2013;Araujo et al., 2014;Lima et al., 2018;Treves et al.,

2022). This TMS-based method notably allowed to follow the fate of sucrose breakdown within guard
cells in connection with the TCA cycle (Medeiros et al., 2018). More recently, a ^{13}C-positionnal
isotopomer approach was also proposed for glutamate and malate to investigate the fate of PEPc-
derived carbon from ^{13}C-labeling experiments (Lima et al., 2021). Previous comparisons of two
derivatization methods for GC-MS suggested that TBDMS-derivatives were potentially less prone to
errors for amino acid analysis than TMS-derivatives (Antoniewicz et al., 2007;Young et al., 2014).
However, the suitability of molecular ions from TMS and TBDMS derivatives was only performed
using U-^{13}C-metabolites in each of these studies, which did not evaluate the entire CID but only the
carbon distribution between the unlabeled isotopologue and the fully labeled isotopologue (M0 and
M5 for a 5-carbon metabolite for example). Unfortunately, isotopic effects can occur at all stages of
the analytical procedures and depend on the tested metabolite and its isotopic content. For instance, the
present methodologies used have not assessed the impact of each isotopologue to the final isotopologue
distribution, which can lead to important biases during the interpretation of the results. Thus, the
reliability of CID measurements by GC-MS has to be evaluated for each individual isotopologue of all
the compounds of interest to subsequently ensure a reliable use for monitoring isotopic labeling
dynamics. This is even more important for plants, given that plant metabolites show strong differences
of ^{13}C-enrichment dynamics between amino and organic acids (Szecowka et al., 2013). For this
purpose, the recent development of an *E. coli* ^{13}C biological standard with controlled patterns of CID
for organic and amino acids may help to address this issue (Millard et al., 2014). Indeed, the recent use
of these reference samples allowed to identify important analytical biases in the CID measurement of
Leucine and Asparagine in a concentration-dependent manner by LC-MS (Heuillet et al., 2018).
Interestingly, this validated LC-MS method successfully measured nitrogen isotope distribution within
plant amino acids and led to the accurate estimation of metabolic fluxes from ^{15}N-isotope transient
labeling experiments in *B. napus* leaves at a metabolic steady-state using the ScalaFlux approach
(Dellero et al., 2020b;Heuillet et al., 2020;Millard et al., 2020).

Here, we proposed a workflow to assess the measurement accuracy of CID from organic and amino
acids by GC-MS for applications in plants. To do so, we built up an analytical method maximizing
information on carbon backbones and analyzed two specific *E. coli* ^{13}C biological standards, produced
to specifically harbor controlled patterns of CID for organic and amino acids (^{13}C-PT). We also
analyzed chemical standards and plant samples at natural ^{13}C abundance to identify potential biases
arising from matrix and analytical systems. Next, we applied our MS method to investigate the
light/dark regulation of metabolic fluxes within plant TCA cycle by performing U-^{13}C-pyruvate
labeling kinetic experiments on *B. napus* leaf discs under light or dark conditions. Using ^{13}C-MFA
principles and some specific isotopologues, we calculated the fractional contribution of
phosphoenolpyruvate carboxylase (PEPc) and TCA cycle to malate and citrate production and
proposed a simplified carbon flux map with relative metabolic fluxes. Our results notably revealed that
this TCA cycle/PEPc ratio was close to 4 for Malate and highlighted the cyclic nature of the TCA cycle
in *B. napus* leaves. This ratio could be substantially regulated by light/dark conditions and mostly
reflected the light-dependent activation of PEPc. In addition, we showed that pyruvate dehydrogenase
and TCA cycle-derived CO_2 contributed to around 5% of net CO_2 assimilation under light conditions.

2 MATERIALS AND METHODS

Organisms and culture

The strain Escherichia coli K-12 MG1655 was grown on minimal medium containing 5 mM
KH$_2$PO$_4$,10mM Na$_2$HPO$_4$, 9 mM NaCl, 40 mM NH$_4$Cl, 0.8 mM MgSO$_4$, 0.1 mM CaCl$_2$, 0.3 mM
thiamine, and 45 mM of the ^{13}C-labeled Acetate mixture as the sole carbon source. Acetate and

thiamine were sterilized by filtration, whereas the other compounds were autoclaved separately. Batch
cultures were carried out in 1-L baffled flasks with 500 mL of medium sparged with synthetic gas (80%
N_2/20%O_2 containing <1 ppm CO_2, Air Liquide). Cultivation was performed in a Multifors Bioreactor
(Infors HT, Bottmingen-Basel, Switzerland) at 37 °C and 200 rpm of orbital shaking. The pH was
maintained at 7.0 by the addition of 2 M HCl. Cell growth was monitored by measuring optical density
at 600 nm with a Genesys 6 spectrophotometer (Thermo, Carlsbad, CA, USA). Amount of biomass in
each sample was calculated using a coefficient of 0.37 g of cell dry weight per OD unit. Arabidopsis
thaliana seeds were sterilized for 10 min with absolute ethanol containing 7.5% NaClO and rinsed two
times with absolute ethanol. Seeds were sowed on ½ MS media containing 1% Agar and incubated to
the dark at 4°C for 48-72 hours to break seed dormancy. Seedlings were grown at ambient air under a
day/night cycle with 16 h light/8h dark at a light intensity of 120 µmol photons.m^{-2}.s^{-1} and 20°C
day/18°C night. Seedlings were harvested after 2-3 weeks by instant sampling with liquid nitrogen and
lyophilized for 96 h before to be stored at -80°C until analysis. Two-month old *Brassica napus* plants
were obtained from seeds exactly as described before (Dellero et al., 2020a). The leaf L9 was used for
all experiments (natural ^{13}C abundance and U-^{13}C-pyruvate incorporation).

U-^{13}C-pyruvate labeling experiments

Four hours after the beginning of the light period, each leaf was cut at the basis of the petiole and then
66 leaf discs (0.8 cm^2) were randomly punched with a cork-borer in both laminas of each leaf. Leaf
discs were first floated in a 10 mM MES-KOH (pH 6.5) buffer for 15 min in order to process the L9
leaf for four plants (see details for leaf annotation in (Dellero et al., 2020a)). In this experiment, one
leaf of a plant corresponded to a biological replicate. Then for each time point, six leaf discs from the
same leaf were transferred in a well of a 6-well microplate filled with 4 mL/well of a buffer containing
10 mM MES-KOH (pH 6.5) and 10 mM U-^{13}C-Pyruvate (^{13}C isotopic purity of 99%; sodium salt from
Cambridge Isotope Laboratories (Eurisotop, France)). All incubations were performed at 20 °C with
an orbital shaking of 70 rpm under either a continuous light of 120 µmol photons m^{-2} s^{-1} or under dark
conditions using multiple opaque plastic bags, similarly to the previous experimental set-up used for
dark-induced senescence (Dellero et al., 2020a). The timepoint 0 was the same for both conditions as
it corresponded the beginning of the experiment. For each time point (0, 0.5, 1, 2, 4, 6 hours of
incubation), the leaf discs were harvested and rinsed three times (10 sec) with a neutral buffer (i.e.,
without U-^{13}C-Pyruvate) and dried for 10 s on a clean tissue before being frozen in liquid nitrogen and
stored at -80 ∘C. The timepoint 0 of the experiment was incubated 2 sec with the labeled buffer before
being rinsed in order to evaluate the fraction of ^{13}C$_3$-pyruvate detected in our analysis that was not
incorporated inside leaf discs (leading to a mean ^{13}C enrichment close to 35% for pyruvate at T0 in our
study).

Analysis of ^{13}C-labeled Acetate by NMR

An equimolar mixture of all ^{13}C-isotopomers of acetate was produced from ^{12}C-Acetate, 1-^{13}C-Acetate,
2-^{13}C-Acetate and U-^{13}C-Acetate (^{13}C isotopic purity of 99%, sodium salts from Cambridge Isotope
Laboratories (Eurisotop, France); **Figure 1**). Quantitative 1H 1D- NMR was used to control the
isotopic composition of the ^{13}C-labeled substrate before use. NMR spectra were recorded with an
Avance 500 MHz spectrometer (Bruker, Rheinstetten, Germany) at 298 K, using a 30° pulse and a
relaxation delay of 20 s. The proportion of each isotopic form was quantified by fitting using a mixed
Gaussian−Lorentzian model (Millard et al., 2017).

Metabolite extraction and quantification

^{13}C-PT full metabolites and ^{13}C-PT protein hydrolysates extracts were obtained following procedures
described previously (Heuillet et al., 2018). Briefly, for the ^{13}C-PT full metabolites extract, 1 mL of *E.*
coli broth at OD600 nm = 1.7 was micro-filtered and the pellet was rapidly quenched at -20 °C into a
mixture of acetonitrile/methanol/H$_2$0 (2:2:1) with 0.1% formic acid. After a centrifugation step (7000g,
−20 °C, 15 min), the supernatant was evaporated and resuspended in 250 µL of Milli-Q water, to be
stored at −80 °C until analysis. For the ^{13}C-PT protein hydrolysates extract, the pellet obtained with
the cellular extract was subjected to an acid hydrolysis with 6N HCl at 110 °C overnight. Samples were
evaporated and rinsed 3 times with Milli-Q water. After a centrifugation step (10 min, 12 000g), the
supernatant was stored at −80 °C. Polar metabolites of freeze-dried Arabidopsis seedlings and *B. napus*
leaves were extracted with a MeOH/Chloroform/H$_2$O mixture following procedures described
previously (Dellero et al., 2020b). The quantification of amino acid content was performed by HPLC-
UV exactly as described previously (Dellero et al., 2020b). The quantification of organic acids was
performed by GC-FID exactly as described previously (Bianchetti et al., 2021).

Determination of Carbon Isotopologue Distribution by GC-MS

Dried aliquots of samples were derivatized on a MultiPurpose Sampler Gerstel and analyzed by GC-
MS (GC-2010 Plus Shimadzu coupled to a triple quadrupole mass spectrometer GCMS-TQ8040
Shimadzu). Polar metabolites were first derivatized for 1 h at 40°C with 50 µL of pyridine containing
20 g.l^{-1} methoxyamine chloride and then were incubated with 50 µL of N-Methyl-N-(trimethylsilyl)-
trifluoroacetamide (MSTFA) for 0.5 h at 40°C. 1 µL was injected with a split mode 1/20 at 260°C on
a TG-5MS column (26098-1420, Thermo Scientific). Separation was performed in a Helium gas-
stream at 1 mL.min^{-1} using the following temperature ramp : 4 min at 70°C, +10°C.min^{-1} until 198°C,
2 min at 198°C, +1°C.min^{-1} until 202°C, +15°C.min^{-1} until 268°C, 3 min at 268°C, +1°C.min^{-1} until
272°C,+10°C.min^{-1} until 310°C and 7 min à 310°C. The transfer line was maintained at 300°C and the
ion source at 200°C. Ionization was performed by electron impact at 70 eV and the MS acquisition was
carried out after a solvent delay of 5 min in full scan mode at 7 spectra.s$_{-1}$ over the m/z range 45-700.
MS Spectra were processed with the software "GCMS Solutions" supplied by Shimadzu to adjust
retention times between each run using a mixture of alkanes (C7 to C40, Sigma-Aldrich). Peak identity
was established according to their retention time and mass fragmentation spectra with respect to
authentic standards derivatized (**Table S1**), and by reference to mass spectra of derivatives in the US
National Institute of Standards and Technology database and in the Golm metabolome database. All
mass isotopologue raw areas were subjected to correction for the contribution of naturally occurring
isotopes of all elements except the carbon arising from the metabolite backbone using the IsoCor v2.1.4
package of Python (https://github.com/MetaSys-LISBP/IsoCor) in low resolution mode (Millard et al.,
2019). In addition, unlabeled chemical standards and Arabidopsis and *B. napus* leaves were
additionally corrected for the contribution of naturally occurring isotopes of the carbon arising from
the metabolite backbone (giving a residual molecular ^{13}C enrichment of 0 for molecules labeled at
natural ^{13}C abundance). ^{13}C-PT samples were not concerned by this additional correction since they
held fully controlled ^{12}C/^{13}C patterns at all positions. A correction method dedicated to IsoCor was
specifically created for the TMS-derivative fragment described in this study (**Table 1**), *i.e.* by
considering the contribution of all atoms added by the derivatization on the isotopic measurement of
the monitored fragments (chemical separation of all atoms arising from the metabolite and the
derivative). All files required to run this method on IsoCor are available in the supplementary material
of this article (**Supplementary files**) and all information will be updated indefinitely in the following
INRAE dataverse (Dellero and Filangi, 2021). The method is publicly available and can also be run
directly from GC-MS source files using the following Galaxy workflow (https://galaxy.genouest.org/,
section "Metabolomics", subsections "Conversion GCMS PostRun Analysis to IsoCor" and "Isotope
Correction for mass spectrometry labeling experiments"). For this purpose, each raw GC-MS dataset

should contain a column "Name" for the fragments considered and a column "Area". The name of each
fragment must be written exactly as specified in the "Metabolite.dat" (an example file is provided in
the **supplementary material**). Fractional Carbon Isotopologue Distribution and mean ^{13}C enrichment
were directly obtained from IsoCor outputs (isotopologue_fraction and mean_enrichment) while CIDs
were converted to binomial distributions by multiplying with 2^n, where n is the number of carbon atoms
(Millard et al., 2014). The accuracy of measurements was assessed from the closeness of experimental
and predicted values for CIDs and mean ^{13}C enrichments at either ^{13}C-PT abundance or at natural ^{13}C
abundance. CID accuracy was formally defined as the mean of absolute differences between measured
and predicted values for each isotopologue (Heuillet et al., 2018).

Pathway fractional contribution

Pathway fractional contribution (FC) can be applied to a metabolic network in a metabolic and isotopic
steady-state to compare the relative contribution of the considered pathway to the production of a
metabolite, as long as these pathways produced specific isotopologues that can be monitored (Buescher
et al., 2015). Here, FC was calculated from PEPc and TCA cycle-specific mass
isotopomers/isotopologues for Malate and Citrate from the timepoint 4 and 6 hours (metabolic steady-
state and pseudo-isotopic steady-state for isotopologue ratios). In the following equations, *mi* refers to
the isotopologue enrichment calculated with respect to the CID (sum of all *mi* for a given metabolite
equal to 1). As all these calculations are relative, it is also possible to use corrected areas of mass
isotopologues (outputs from IsoCor). For Malate_3TMS_C1C4 (corresponding to the GC-MS
fragment from Malate(3TMS) metabolite than contained the C1-C2-C3-C4 carbons of Malate
(**Table1**)), M1 isotopologue measurement reflected both PEPc and TCA cycle isotopomers (**Figure
6**). However, the ratio between M1 and M3 isotopologues of Malate_3TMS_C1C4 produced by the
TCA cycle (denoted *X* in the following equations) can also be observed in Succinate_2TMS_C1C4 and
Glutamate_2TMS_C2C5, thereby reflecting the M0 and M2 from mitochondrial active pool of Acetyl-
CoA, *i.e.*:

$$Succinate C1C4\left(\frac{m3}{m1+m3}\right) = Glutamate C2C5\left(\frac{m3}{m1+m3}\right) = Acetate\left(\frac{m2}{m0+m2}\right) = X$$

From this parameter, we calculated the FC of PEPc and TCA cycle to Malate and Citrate production
as follows:

$$FC\ Malate_{PEPc} = Malate C1C4\left(\frac{m1-\left(\left(\frac{1-X}{X}\right)\times m3\right)}{m1+m2+m3}\right)$$

$$FC\ Malate_{TCA\ cycle} = Malate C1C4\left(\frac{m2+\frac{m3}{X}}{m1+m2+m3}\right)$$

$$FC\ Citrate_{PEPc} = Citrate C1C6\left(\frac{m1+m3}{m1+m2+2\times m3+m4}\right)$$

$$FC\ Citrate_{TCA\ cycle} = Citrate C1C6\left(\frac{m2+m3+m4}{m1+m2+2\times m3+m4}\right)$$

It should be noted that *m3* was used twice in the FC Citrate calculations because this isotopologue is
produced by both pathways. Finally, the FC of pyruvate dehydrogenase (PDH) and TCA cycle-derived
CO_2 to photosynthesis was calculated using X as a proxy for the $^{13}C/^{12}C$ ratio of this CO_2 fraction in
light conditions. Considering the isotopic and metabolic steady-state for these metabolites, the mean
^{13}C enrichment of X should be observed in Serine if this fraction of CO_2 accounted for 100% of CO_2
fixed by photosynthesis, *i.e.:*

304
$$FC\ photosynthesis_{PDH\ and\ TCA\ cycle\ CO_2} = \frac{mean\ ^{13}C\ enrichment\ (SerineC1C3)}{X}$$

305 Mean ^{13}C enrichment of Serine C1C3 was recalculated from M1 isotopologue of Serine_2TMS_C1C2
306 and Serine_2TMS_C2C3 assuming that Serine was only labeled in the C2 of these fragments based on
307 the validated hypothesis (**Figure 7B**).

Statistical analysis

309 Statistical comparisons were performed using either confidence intervals (comparison of a group of
310 value against one reference value), a Student-test (comparison of 2 groups of values) or an ANOVA
311 followed by a post-hoc Tukey-HSD test (comparison of 3 or more groups of values) with a level of p-
312 value <0.05 by using the publicly available Rstudio software (RStudio Team, 2016).

3 RESULTS

The proposed workflow (**Figure 1**) consisted in three steps: 1) the build-up of a method to identify and
select abundant molecular ions deriving from TMS-derivatives with a large number of carbons; 2) the
production of two reference ^{13}C-labeled samples with a controlled pattern for CID (^{13}C-PT) using *E.*
coli as a metabolic factory; 3) the validation of TMS-derivatives mass fragments for CID measurements
with our GC-MS method for applications in plants using ^{13}C-PT reference samples and unlabeled and
^{13}C-labeled plant samples.

**Selection of GC-MS mass fragments maximizing information on carbon back-bones of organic
and amino acids**

We combined a literature survey analyzing common pathway of TMS-derivative mass fragmentations
with the injection of chemical standards in our analytical system to select some abundant fragments
maximizing the information on carbon backbones. Their chemical identity was further confirmed by
the observed CID pattern with our two reference ^{13}C-PT samples. Derivatives with only one TMS
group were not considered in this study due to the low yield of these fragments after fragmentation and
the occurrence of important contaminations for low m/z. Overall, TMS-derivatives of most α-amino
acids were characterized by an α-cleavage of the carboxylic function in the form of a TMS-CO_2 (m/z
117) (Harvey and Vouros, 2020), thus leading to the abundant formation of fragments containing all
others atoms of the α-amino acids except the C1 (**Table 1**). Only Ornithine(4TMS) could be detected
without any mass loss (**Table 1**, (Harvey and Vouros, 2020)). Aromatic amino acids showed an
additional α-cleavage of the amine function leading to the additional production of an abundant m/z
218 fragment containing the C1-C2 for Phenylalanine, Tryptophan and Tyrosine (**Table 1**, (Lai and
Fiehn, 2018;Harvey and Vouros, 2020)). Only tryptophan did not proceed through an α-cleavage of
the carboxylic function compared to Phenylalanine and Tyrosine, but the other fragment containing
the C3 to C11 atoms resulting from the α-cleavage of the amine function could be abundantly detected.
Serine(3TMS) and Valine(2TMS) showed also some specific abundant molecular ions in addition to
the α-cleavage of the carboxylic function (**Table 1**,(Harvey and Vouros, 2020)). Arginine, Histidine

and Cysteine could not be detected by GC-MS using TMS derivatization and were not further
considered in our study. Beside this, β-Alanine, an important β -amino acid from plants, showed a
classical abundant fragment resulting from the loss of methyl from a TMS group, which is known to
be very low in α-amino acids (**Table 1**, (Lai and Fiehn, 2018)). Interestingly, fragmentation of the
TMS-derivatives of organic acids related to the TriCarboxylic Acid cycle proceeded through an α-
cleavage within a TMS group leading to the loss of a methyl radical and producing a [M-15] fragment
containing all carbon atoms. However, this fragment was very weak for Citrate and Malate. The highest
fragments maximizing information on carbon backbones were m/z 273 (C1 to C5) for Citrate(4TMS)
and m/z 233 (C2 to C4) for Malate(3TMS). The identity of the carbons hold by these fragments has
been recently assigned using position-specific [13]C-labeled standards prepared by *in vitro* enzymatic
reactions (Okahashi et al., 2019).

Evaluation of CID measurements by GC-MS using reference [13]C-PT samples and plant samples
at natural isotope abundance

Next, we produced reference [13]C-labeled samples using *E. coli*, denoted here [13]C-PT. This strategy
relies on the use of an accurate equimolar mixture of all [13]C isotopomers of acetate to biologically
produce metabolites with fully controlled and predictable [13]C-labeling patterns (Millard et al., 2014).
Thus, molecular [13]C enrichment of each carbon-containing metabolite will be theoretically equal to
50% and all isotopomers will be present in equimolar concentrations, leading to a binomial distribution
of isotopologues following Pascal's triangle. For this purpose, we grew *E. coli* in a minimal media with
this acetate mixture as the sole carbon source of the media (**Figure 1**) and checked its isotopic content
by NMR (molecular [13]C enrichment expected at 49.9 % ± 0.8). From cell harvest, two specific
reference [13]C-PT samples were produced during the extraction: i) a [13]C-PT full metabolite extract
containing all polar metabolites for the evaluation of CID for organic acids; ii) a [13]C-PT protein
hydrolysates extract containing proteogenic amino acids in high concentrations for the evaluation of
CID for amino acids. This strategy allowed to maximize the amount of amino acids that can be weakly
abundant as soluble pools when *E. coli* is grown in a minimal media. It should be mentioned that the
hydrolysis process deamidated Asparagine into Aspartate and Glutamine into Glutamate, thus these
amino acids could not be evaluated (Kambhampati et al., 2019). In addition, non-proteogenic amino
acids from the **Table 1** could not be evaluated with this extract (Ornithine, β-Alanine).

To facilitate comparison of CIDs between the different amino acids, we converted the measured [13]C-
enrichments among isotopologues of each fragment into binomial distribution, thus reflecting the
number of isotopomers considered for each isotopologue (**Figure 2** and Material and Methods). In the
following figures, GC-MS fragments will be denoted according to the considered metabolite, its TMS-
derivative analyzed and the metabolite carbon backbone (for example, Glutamate_3TMS_C2C5 means
that the fragment contains the carbon C2-C3-C4-C5 (**Table1**)). Overall, all the considered amino acids
were detected in the [13]C-PT protein hydrolysates except Tryptophan (**Figures 2A, S1A**). We found
very a high level of correlation between the measured CID (blue bars) and the expected CID (black
bars) for numerous amino acid fragments considered in our method: α-Alanine_2TMS_C2C3,
Aspartate_C2C4 either in 2TMS or 3TMS, Glutamate_3TMS_C2C5, Glycine_3TMS_C2,
Isoleucine_2TMS_C2C6, Leucine_2TMS_C2C6, Phenylalanine_2TMS_C1C2 and C2C9,
Serine_2TMS_C2C3, Serine_3TMS_C1C2 and C2C3 and Tyrosine_2TMS_C1C2. CID
measurements for Methionine_2TMS_C2C5 showed a high variability coupled to a lower estimation
of M3 and M4 isotopologues at the expense of M1 and M2 (**Figure 2A**). However, this amino acid
was in a very low abundance in the [13]C-PT protein hydrolysates compared to the others amino acids
(**Figure S1A**). Lysine and Tyrosine_2TMS_C2C9 had also important variability for their CID
measurements, leading to misestimations of their M3 isotopologues. The most important and

significant errors for CID measurements of amino acids concerned Proline_2TMS (overestimation of
M4), Threonine_3TMS (overestimation of M3 at the expense of M0 and M1), Valine_2TMS_C1C2
(overestimation of M2) and Valine_2TMS_C2C5 (overestimation of M3 at the expense of M1 and
M2). Considering the recent establishment of a ^{13}C-positional isotopomer approach for Glutamate to
analysis its C1 carbon (Lima et al., 2021), we also analyzed the CID for the two low abundant fragments
Glutamate_3TMS_C1 (CO_2-TMS, M0 at m/z 117) and Glutamate_3TMS_C1C5 fragment (CH3 loss,
M0 at m/z 348). However, they both showed large and significant misestimations of their CIDs, thus
questioning the accuracy of this isotopomer-based method (**Figure S1B**). The analysis of organic acids
from ^{13}C-PT full metabolites extract showed relatively good CID for Citrate_2TMS_C1C5,
Malate_3TMS_C2C4 and Fumarate_2TMS_C1C4 (**Figure 2B**). Citrate_4TMS_C1C6 and
Malate_3TMS_C1C4 showed also relatively good CID but they had a higher variability compared to
their respective C1C5 and C2C4 fragments. There was a very high overestimation of M0 in
Succinate_2TMS_C1C4, which seemed to have an impact on M2, M3 and M4 isotopologues,
However, these misestimations for Succinate have been previously attributed to contaminations
inherent in the ^{13}C-PT full metabolites extract used here (Millard et al., 2014), thus preventing the
analytical validation for this fragment. For Pyruvate_MEOXTMS_C1C3, the variability was too high
to assess the CID measurements, perhaps reflecting its weak derivatization yield and low abundance
in ^{13}C-PT full metabolites extract (**Figure S1A**). To further assess the distribution of measurement
errors among isotopologues, we analyzed the CID accuracy, using threshold values of ±0.05 (**Figure
3A**). Overall, the bias of measurement for each isotopologue of each metabolite was relatively low
except for Methionine, Valine, Threonine, Succinate and Pyruvate. Interestingly, these very high levels
of accuracy for CID measurements were converted into very high levels of accuracy for mean ^{13}C
enrichment measurements, leading to very small shifts between expected (0.499 ±0.08) and measured
values (below 1% of error), except for Methionine, Valine, Threonine, Succinate and Pyruvate (**Figure
3B**). However, Malate_3TMS_C1C4, Lysine_2TMS_C2C6 and Serine_2TMS_C2C3 showed
important variability for mean ^{13}C enrichment measurements although no strong biases in their CID
measurements were previously observed. Conversely, mean ^{13}C enrichment of
Phenylalanine_2TMS_C1C2 and proline_2TMS_C2C5 were slightly overestimated and reflected the
overestimation of their respective isotopologues M2 and M4. To ensure that the tested fragments were
suitable for isotopic analyses of metabolites from plant samples, we analyzed their accuracy at natural
^{13}C abundance. For this purpose, we produced two types of plant samples (seedlings from Arabidopsis
and leaves from *B. napus*) and we measured the CID for each fragment of our method to calculate the
residual molecular ^{13}C enrichment (**Figure 3C**).

This variable corresponded to the ^{13}C enrichment remaining after removing the contribution of natural
abundance for all isotopes (a value of 0 is expected if no contamination occurs). Among the metabolites
that were detected in the plant samples, most of the fragments for organic and amino acids showed
relatively high accuracy at natural ^{13}C abundance, with most of the mean values of residual molecular
^{13}C enrichment ranging between and 0.05 and 3% (**Figure 3C**). Interestingly, some discrepancies were
observed between Arabidopsis seedlings and *B. napus* leaves, notably for the full carbon backbone
fragments Citrate_4TMS_C1C6 and Malate_3TMS_C1C4, which were more accurately measured in
B. napus at low enrichment levels (critical values close to 5% for Arabidopsis). Conversely,
Arabidopsis seedlings showed slightly higher accuracies than *B. napus* leaves for
AlphaAlanine_2TMS_C2C3, Aspartate_2TMS_C2C4, Fumarate_2TMS_C1C4 and Serine_3TMS
fragments. Phenylalanine_2TMS fragments showed residual molecular ^{13}C enrichments close to 5%
but only in plants samples. Surprisingly, Proline_2TMS_C2C5, Threonine_3TMS_C2C4 and
Valine_2TMS_C2C5 showed relatively important residual molecular ^{13}C enrichments (nearly 10%) in
both chemical standards and unlabeled plant extracts, suggesting the occurrence of analytical bias. A
detailed survey showed that the isotopic cluster of Proline_2TMS_C2C5 was contaminated in the M4

isotopologue (m/z 146) by a mass fragment deriving from near elution of Isoleucine_2TMS, and
correlated with previous measurements of CIDs (**Figures 2, 3D**). Similarly, the isotopic cluster of
Valine_2TMS_C2C5 was contaminated in the M3 isotopologue (m/z 147) while
Threonine_3TMS_C2C4 showed a surprising very high abundance of the M1 isotopologue (40% of
the CID) in both chemical standards and unlabeled plant extracts.

Assessment of the light/dark regulation of TCA cycle and PEPc fluxes in *Brassica napus* leaves
using a ^{13}C-MFA approach

448   To further validate our MS method, we applied it to ^{13}C-labeled plant samples. For this purpose, we
performed transient kinetics of U-^{13}C-pyruvate incorporation to *B. napus* leaf discs subjected to either
light or dark conditions for up to 6 hours. The aim of this experiment was to obtain an overview of the
light/dark regulation of metabolic fluxes associated to TCA cycle and PEPc using MS methodology
and isotopologue-based approaches. We used U-^{13}C-pyruvate as a probe since the mitochondrial
Pyruvate DeHydrogenase (PDH) will be able to produce simultaneously ^{13}C$_2$-Acetyl-CoA molecules
suitable for the investigation of the TCA cycle and ^{13}CO$_2$ molecules suitable for the investigation of
PEPc activity through the combined action of carbonic anhydrase (H^{13}CO$_3^-$).

Following our experiments, the analysis of mean ^{13}C enrichment for all conditions revealed a
significant increase for α-Alanine, Aspartate, Citrate, Glutamate, Glycine, Malate, Pyruvate, Serine
and Succinate along the U-^{13}C-pyruvate incorporation (**Figure 4A**, **Table S2**; **Table S4** for statistical
analysis). There was a very high biological variability and small ^{13}C enrichment for
Fumarate_2TMS_C1C4. Overall, most of these enrichment dynamics remained relatively equivalent
between light and dark conditions for all the fragments considered, except for Glycine and Serine,
which were almost exclusively enriched in light conditions. These results were surprising since
previous experiments using a 4-hour incorporation of 15 mM U-^{13}C-pyruvate into Arabidopsis leaves
showed a higher mean ^{13}C enrichment for Citrate, Glutamate and succinate in the *pdk* mutant (light-
dependent activator of the PDH complex) compared to the control (Zhang et al., 2021). Considering
Glycine and Serine, the importance of the light-dependent photorespiratory pathway for their
production in leaves clearly suggested that a significant part of ^{13}CO$_2$ arising from PDH and TCA cycle
decarboxylations (isocitrate dehydrogenase and 2-oxoglutarate dehydrogenase) was reassimilated by
photosynthesis. A statistical analysis revealed that Glycine_3TMS_C2, Serine_3TMS_C1C2 and
Serine_3TMS_C2C3 reached a pseudo-isotopic steady-state starting after 2 hours of illumination
(**Table S4**). Other fragments showed a relatively linear evolution of their mean ^{13}C enrichment, except
for Pyruvate_MEOXTMS_C1C3 which rapidly reached an isotopic steady-state after 0.5 hours only.
Interestingly, labeling data from Pyruvate_MEOXTMS_C1C3 and Succinate_2TMS_C1C4 fragments
had a very good resolution (weak analytical and biological variability) compared to previous evaluation
with ^{13}C-PT full metabolites extract. Next, we performed quantification of metabolite pool sizes along
the experiment to determine whether the considered metabolic network was in a metabolic steady-state
(**Figure 4B, Table S3**). For this purpose, we used GC-FID for organic acids and UPLC-UV for amino
acids using AccQTag methodology as they have a very high level of stability for absolute
quantification. Indeed, TMS-derivatives of amino acids are highly unstable along a GC-MS run,
leading to drastic differences of response when quantifying them with external standards curves (Data
not shown). In addition, the reliable absolute quantification of metabolites using their CID masses can
only be achieved by using internal isotopic standards for each metabolite considered (not our case in
the present study) (Heuillet et al., 2020;Evers et al., 2021). Overall, metabolite pool sizes were
relatively different for each of them but there was no significant difference between light and dark
conditions, except for Aspartate. Glycine was only quantified in light conditions (not detected in "dark
condition" samples with UPLC-UV). Interestingly, a statistical analysis showed that all the metabolite

pool sizes reached a significant pseudo steady-state starting after 1 hour (similar statistical groups in
Table S5). To summarize, we have found some similar [13]C enrichment dynamics for TCA cycle
derived metabolites in both light and dark conditions combined to similar metabolite pool sizes at a
metabolic steady-state.

Next, we focused our analysis on the isotopologue dynamics of these metabolites in order to depict the
considered metabolic network and compare metabolic fluxes associated to TCA cycle and PEPc
(**Figure 5**). Mean [13]C enrichment of Citrate, Glutamate and Succinate was essentially driven by an
increase of the M2 isotopologue for light and dark conditions, thus reflecting the incorporation of M2
Acetyl-CoA arising from U-[13]C-pyruvate. However, there was also an increase of M1 and M3
isotopologues for Glutamate and Succinate after 2 hours while M1, M3 and M4 isotopologues of
Citrate_4TMS_C1C6 and Citrate_2TMS_C1C5 were increased starting from 4 hours. M1 and M4 of
Citrate_4TMS_C1C6 were essentially increased in dark conditions compared to light conditions after
4 hours. There was a very high variability for Fumarate_2TMS_C2C4 CID in both conditions, which
prevented its use for a reliable isotopic investigation. Interestingly, Malate_3TMS_C1C4 fragment
showed a higher enrichment of M1 compared to M2 in the light condition while M2 remains higher
than M1 in the dark condition. This shift may reflect an increase of the PEPc activity under light
condition (M1 arising from the reassimilation of the mitochondrial [13]CO$_2$ decarboxylations). Since
there was no difference of mean [13]C enrichment between light and dark conditions for this fragment
(**Figure 4A**), this result reinforced the importance of analyzing specifically CIDs for the interpretation
of labeling data. In addition, the CID of Malate_3TMS_C2C4 showed a similar dynamic for both light
and dark conditions, hence ruling out the use of this fragment to disentangle PEPc and TCA cycle
contribution to Malate production (**Figure 5**). Aspartate_3TMS_C2C4 showed a very high variability
for the light condition, but the CID dynamic in the dark condition was similar to the CID dynamic of
Malate_3TMS_C2C4. Thus, both fragments seemed to mimic Oxaloacetate labeling dynamics.
Pyruvate_MEOXTMS_C1C3 was only enriched with the M3 isotopologue (nearly 80% M3/20% M0),
thereby suggesting that the activity of the malic enzyme, converting malate to pyruvate, was negligible
in our metabolic network.

From these observations, we built a simplified metabolic network for TCA cycle composed of two
"turns" to explain the production of M1 and M3 in both Succinate_2TMS_C1C4 and
Glutamate_2TMS_C2C5 (**Figure 6A**). In this network, the second "turn" reflected the commitment of
a molecule that has already experienced all reactions of the TCA cycle to another round of reactions
(Orange and blue labels to compare acetyl-CoA-derived labeling from 1[st] and 2[nd] "turn"). Based on
previous works (Abadie and Tcherkez, 2019a), the reverse reaction of the fumarase was also considered
in the network, although it has no effect on the M1 isotopologue monitored by mass spectrometry (C4
and C1 isotopomers are still detected as M1 isotopologues in our study). The involvement of a light-
dependent stored [12]C-Citrate pool was not represented here since it only provided M0 isotopologues
(no additional isotopologues to explain). From this metabolic network, we proposed to evaluate PEPc-
dependent and TCA cycle-dependent contributions to Citrate production through M1, M2, M3 and M4
isotopologues and to Malate production through M1, M2 and M3 isotopologues. These contributions
were only calculated using timepoints 4 and 6 hours, because all the fragments to consider for these
contributions were mostly visible at these moments. The main idea of this pathway fractional
contribution (FC) to a given metabolite production rely on the use of pathway-specific mass isotopomer
ratios for the metabolite considered, as long as the metabolic network is in a metabolic and isotopic
pseudo steady-state. Here, we observed a rapid metabolic steady-state (**Figure 4B**, **Table S5**) and a
relatively negligible evolution of the ratios for the considered isotopologues between 4 and 6 hours
(comparison of ratios for M1, M2 and M3 isotopologues only for Malate for example). In this context,
the relative contribution of PEPc and TCA cycle to either Malate and Citrate production remained in

the same order of magnitude between T4 and T6 hours (**Figure 5**). An important issue to address for
calculating FC concerned the M1 from Malate_3TMS_C1C4, which monitored at the same time PEPc
and TCA cycle dependent isotopomers. To deal with, we took advantage of Succinate_2TMS_C1C4
and Glutamate_2TMS_C1C4 fragments which both harbored M1 and M3 isotopologues that are the
precursor of TCA cycle-dependent M1 and M3 of Malate_3TMS_C1C4 (**Figure 5**). Analysis of the
M1/(M1+M3) ratio for these two fragments showed a remarkable stability between dark and light
conditions, thus reflecting a stable M2 enrichment of 40% for the active pool of acetyl-CoA (**Figure 6**
B and Equations). This result also showed that the isotopologue ratios considered here were at a pseudo
isotopic steady-state. Using this ratio, we calculated the FC of PEPc and TCA cycle for malate and
citrate respectively (**See equations**). We found that TCA cycle had a significant four-fold higher
contribution to either Malate or Citrate production compared to PEPc in all conditions (around 80%
versus 20%). Interestingly, there was a 1.6-fold increase of the FC of PEPc to Malate production from
dark to light conditions (14% versus 23%), although it remained not statistically significant (high
variability, t-test p-value =0.08). Conversely, PEPc contribution to Citrate production was significantly
1.5-fold higher in dark condition than in light condition (26% versus 17%). Thus, the results suggested
a higher commitment of PEPc-derived carbons to the forward part of the TCA cycle in the dark, while
commitment of PEPc-derived carbons to the reverse part of the TCA cycle (Malate and perhaps
Fumarate) was reduced at the same time.

Finally, we explored the isotopologue distribution of Glycine and Serine, some amino acids which
were previously identified as a proxy for photosynthetic reassimilation of PDH and TCA cycle-derived
$^{13}CO_2$ (**Figure 7A**). Our results showed that the increase of mean ^{13}C enrichment for the three
considered fragments in light condition came from an increase of their M1 isotopologue. Given that
the light-dependent photorespiratory enzymes glycine decarboxylase and serine:hydroxymethyl
aminotransferase convert 2 glycine into 1 serine, we wanted to know whether the fragments
Serine_2TMS_C1C2 and Serine_2TMS_C2C3 were labeled in the same or different carbons, thus
revealing if the C1 of Glycine was also labeled (**Figure 7B**). By reviewing all the possible scenarios
with different combinations of labeled glycine, we showed that the CID associated to our fragments
(isotopologue occurrence) could only be produced with one molecule of glycine labeled at the C2 and
an unlabeled molecule of glycine, leading to the production of a molecule of Serine labeled only at the
C2 and an unlabeled CO_2 molecule. Since we have shown that Glycine and Serine were both at the
isotopic and metabolic steady-state between 2 and 6 hours (**Figure 4**, **Table S4**, **Table S5**), we used
Serine fragments to evaluate the contribution of PDH and TCA cycle-derived CO_2 to the photosynthetic
activity. For this purpose, the steady-state enrichment of Acetyl-CoA calculated before was used as a
proxy of the $^{12}C/^{13}C$ ratio of the CO_2 produced by pyruvate dehydrogenase since a significant
proportion of PDH-derived ^{12}C could still contribute to photosynthesis (**See equations**), and given its
presumable contribution for light respiration (CO_2 emissions) (Tcherkez et al., 2005). After
recalculating the mean ^{13}C-enrichment of SerineC1C3 from both Serine fragments (C1-C2 or C2-C3),
we showed that the light-dependent assimilation of PDH and TCA cycle decarboxylations could
represent around 5% of net photosynthesis (**Figure 7C**).

4   DISCUSSION

The estimation of metabolic fluxes in photosynthetic organisms represents an important challenge that
has gained interest over the last decade with the development of ^{13}C-INST-MFA. This approach
requires a high level of accuracy for CID measurements to avoid propagation of small measurement
errors to large errors in estimated fluxes (Antoniewicz et al., 2006). Considering the need to perform
high throughput measurements at isotopically non-stationary steady-state, MS has been preferentially
selected by the field in the recent years to measure CID in organic and amino acids. However, the

suitability of TMS-derivative molecular ions for these metabolites has never been evaluated at the
isotopologue level, *i.e.* by monitoring the impact of each isotopologue to the final CID measurement.
The use of a validated MS methodology to investigate ^{13}C-labeling dynamics within plant TCA cycle
may help to strengthen our knowledge about its functioning.

A robust methodology to evaluate CID measurements and identify analytical and matrix biases

In this study, we proposed a workflow to assess the measurement accuracy of CID from TMS-
derivatives of organic and amino acids by GC-MS for applications in plants, based on tailor-made *E.*
coli ^{13}C biological standards with controlled patterns of CID (^{13}C-PT) (**Figure 1**). From the TMS-
derivatives (selected to maximize information on carbon backbones), we showed that most of them
had a remarkable accuracy, leading to very small biases in the measurement of CIDs (below 5%) and
mean ^{13}C enrichments (below 1%). We further validated their suitability for application in plants by
monitoring very low residual molecular ^{13}C-enrichments in Arabidopsis seedlings and *B. napus* leaves,
although these accuracies could be variable between the two plant matrices (Citrate_4TMS_C1C6,
Malate_3TMS_C1C4, α-Alanine_2TMS_C2C3, Aspartate_2TMS_C2C4 and Serine_3TMS).
However, among amino acid TMS-derivatives, we identified three important biases at the isotopologue
level. The first bias concerned the contamination of the isotopic cluster of Proline_2TMS_C2C5 at the
m/z 146 (**Figure 3D**). A previous study reported this contaminant using chemical standards at natural
^{13}C abundance, but no explanation was provided for this bias (Antoniewicz et al., 2006). Here, we
showed that the m/z 146 came from the co-elution with Isoleucine, *i.e.* a separation by 0.02 seconds in
our chromatograms, equivalent to 0.33 °C with the ramp time for temperature during the elution
(+10°C.min^{-1}). Nevertheless, this contamination only led to a slight overestimation of the M4
isotopologue (^{13}C-PT samples) (**Figure 2A**), which did not strongly impact the accuracy measurement
for mean ^{13}C enrichment (**Figure 3B**). Thus, this fragment may still be suitable for ^{13}C-MFA in plants,
as long as the ^{13}C-labeling strategy for the metabolic network is adapted to prevent the production of
M4 isotopologues. The second bias identified concerned the contamination of the isotopic cluster of
Valine_2TMS_C2C5 at the m/z 147, which was observed in both chemical standards and plant
matrices. Interestingly, when a siliconium ion (loss of a methyl in a TMS group) operates a cyclization
by attacking another TMS group of the same molecule, it produces an ion $[(CH_3)^3Si\text{-}O\text{-}Si(CH_3)^2]^+$ of
the same mass 147 (Harvey and Vouros, 2020). We also detected a significant abundance of an m/z
246 in the fragmentation spectra of Valine_2TMS, which seems to support this hypothesis for the
contamination (**Figure S2**). The third bias concerned the very high level of M1 isotopologue for
Threonine_3TMS_C2C4 detected in both chemical standards and plant matrices. In its fragmentation
spectra, we found an additional M1 isotopologue contamination for the peak m/z 291, an important
fragment for the identification of Threonine_3TMS by GC-MS (**Figure S2**, NIST database). Given
that we are working at a nominal mass resolution, we cannot determine whether this contamination is
due to: i) a co-elution of a contaminant chemically similar to Threonine; ii) a strong isotopic effect
during the derivatization. The use of GC coupled to high-resolution MS may help to resolve this
analytical issue (instrument not available at present in our platform). Besides this, we observed a high
level of M0 contamination for CID measurement of Succinate using the ^{13}C-PT full metabolites extract,
that has been reported previously (Millard et al., 2014), but prevented its accurate evaluation with our
workflow. Nevertheless, there was a small biological and analytical variability for
Succinate_2TMS_C1C4 in our ^{13}C-labeling experiments and we measured similar ratios for M1 and
M3 with Glutamate_2TMS_C2C5 that was already validated by our method (**Figure 6B**). Since M1
and M3 were both impacted by the M0 contamination in ^{13}C-PT samples, our results suggest that this
fragment can be reliably used for ^{13}C-MFA in plants.

Using our validated method, we provided important information about the activity of the TCA cycle
that contrasted with some proposed flux modes for TCA cycle in illuminated leaves or during a
light/dark cycle (Tcherkez et al., 2009;Gauthier et al., 2010;Sweetlove et al., 2010;Tcherkez et al.,
2017;da Fonseca-Pereira et al., 2021).

Stored Citrate is unlikely to contribute to TCA cycle activity in *Brassica napus* leaves

The hypothesis about the role of night-stored citrate molecules relies on few works to date. In the first
work, after 50 min of $^{13}CO_2$ incorporation in *Xanthium stumarium* leaves, ^{13}C redistribution towards
metabolites were used to propose "transmission coefficients" that required the contribution of an
important pool of unlabeled Citrate to work (Tcherkez et al., 2009). The second work used the
following experimental set-up ($^{13}CO_2$ 6h light/$^{12}CO_2$ 6h dark/$^{12}CO_2$ 6h light, with $^{15}NH_4^{15}NO_3$
available all the time) with *B. napus* leaves and a petiole-feeding assay (Gauthier et al., 2010). Analysis
of the results by NMR showed a weak ^{13}C-enrichment for Citrate, Glutamate, Glutamine and Malate
at the end of the first light period and 30 min after the beginning of the dark period compared to the
end of the experiment. Combined to a higher occurrence of ^{12}C-^{15}N interactions compared to ^{13}C-^{15}N
interactions in Glutamate after 6h of light with $^{13}CO_2$, the results were interpreted as evidences for the
contribution of night-stored vacuolar Citrate and Malate molecules to TCA cycle activity in the
subsequent light period. However, the citrate pool size was not monitored and the ^{13}C enrichment of
TCA cycle metabolites at the end of the dark period and/or at the beginning of the second light period
remained unknown. Nevertheless, these results were then partly supported by a flux balance analysis
using a diel model which explained this stored citrate contribution by the need to support continued
export of sugar and amino acids from the leaf during the night and to meet cellular maintenance costs
(Cheung et al., 2014). Recently, this concept has been reinforced by the subcellular distribution of
Citrate in plant leaves, which is essentially present in the vacuoles (Destailleur et al., 2021). However,
previous NMR works used detached leaves to propose the contribution of the stored citrate, which is
an experimental setup that necessarily stops sink/source relationships (sugar and amino acids exports)
(Gauthier et al., 2010). Thus, the light/dark regulation of the stored citrate contribution should be
observed in both detached leaves and leaf discs. Here, using *B. napus* leaf discs and a U-^{13}C-pyruvate
labeling incorporation, we provided evidences for the absence of stored Citrate contribution to the TCA
cycle in a light/dark dependent manner (**Figure 8**). First, we showed that the M2/(M0+M2) ratio of the
Acetyl-CoA active pool was close to 0.4 after 4 to 6 hours in our experiments, thereby showing that
most of M0 isotopologues from active pools of TCA cycle metabolites came from M0 Acetyl-CoA and
M0 OAA and not a stored pool of ^{12}C-Citrate (**Figure 6B**). Given the important production of PEPc-
derived molecules with non-labeled PEP in our experiments, ^{12}C-Pyruvate may have been produced
by the glycolysis during our kinetic, thereby diluting the mitochondrial U-^{13}C-Pyruvate pool (nearly
20% of M0 still detected for the Pyruvate poll at the leaf level). However, it is also possible that the
thioredoxin-mediated activation of the ATP citrate lyase activity in light conditions leads to the
conversion of stored ^{12}C-Citrate pool into ^{12}C-OAA and ^{12}C-Acetyl-CoA (Sweetlove et al., 2010;da
Fonseca-Pereira et al., 2021). In our experiments, the ^{13}C-enrichment of Acetyl-CoA was calculated
from ^{13}C-labeled fragments of either Succinate or Glutamate (independent of their ^{12}C-molecules (M0
isotopologues)), thus this measurement can only be biased if there is a contribution of stored ^{12}C-Citrate
converted to ^{12}C-Acetyl-CoA. However, if such contribution was relatively important in the light and
negligible in the dark (concept of the stored citrate contribution), it would necessarily introduce a ^{12}C
isotopic dilution for mean ^{13}C enrichment of Glutamate and Succinate in the light compared to the dark
(Gauthier et al., 2010), and this was not observed here while the system was in a metabolic steady-state
(**Figure 4**). Second, we did not observe any significant modification of the Citrate pool during the 6
hours of our experiment (**Figure 4B, Table S4**) while a strong decrease in light conditions and a strong
increase in dark conditions were expected with the hypothesis of the stored citrate remobilization

(Cheung et al., 2014). Altogether, our results showed that the contribution of stored citrate to the TCA
cycle activity is unlikely to occur in *B. napus* leaves (**Figure 8**).

**The forward cyclic flux mode of the TCA cycle has a higher contribution to Malate biosynthesis
compared to PEPc in *Brassica napus* leaves**

The non-cyclic nature of the TCA cycle has been previously proposed in illuminated leaves from works
that used different experimental setup/labeling strategies. This non-cyclic nature in the light was mostly
explained by the need to sustain *de novo* nitrogen assimilation and was supported by higher PEPc
activities compared to dark conditions (Gauthier et al., 2010;Caburatan and Park, 2021). A first
experiment was based on detached leaves of *Xanthium stumarium* and ^{13}C labeling with either ^{13}C$_1$-
pyruvate or U-^{13}C-Pyruvate (Tcherkez et al., 2009). The authors did not observe a significant ^{13}C
enrichment in the Succinate compared to Fumarate with a 2-hour ^{13}C$_1$-pyruvate incorporation while
both metabolites showed similar ^{13}C-enrichments with a 2-hour U-^{13}C-Pyruvate incorporation.
However, the reassimilation of ^{13}CO$_2$ from ^{13}C$_1$-Pyruvate decarboxylation by PEPc cannot be
metabolically transmitted from Malate to Succinate since the C1 and the C4 of the OAA backbone are
lost during Citrate to Succinate conversion (**Figure 6A**) (Alves et al., 2015;Lima et al., 2021). Three
other experiments were based on 1-hour ^{13}CO$_2$ labeling with either detached or attached leaves from
Xanthium stumarium, Arabidopsis or *Camelina sativa* (Tcherkez et al., 2009;Xu et al., 2021;Xu et al.,
2022). The two INST-MFA approaches notably predicted that PEPc and the pyruvate-to-2-
oxoglutarate conversion experienced both relatively similar metabolic fluxes (ratio close to 1), while
the flux from 2-oxoglutarate to succinate and from OAA to malate were both equal to 0. Thus, these
previous works never identified the relative contribution of PEPc and TCA cycle to Malate
biosynthesis. Perhaps, this can be partly attributed to the ^{13}C labeling strategy: ^{13}CO$_2$ must be quickly
assimilated by the PEPc thanks to the anhydrase carbonic while the ^{13}CO$_2$ fixed by the RuBiSCO will
require more time to be transmitted to PEP and pyruvate (See the dynamics of CIDs for Trioses
phosphates, Alanine, Citrate and Aspartate therein (Xu et al., 2021)). Consequently, ^{13}C enrichment
could be artificially higher in Citrate and Glutamate compared to Succinate with this ^{13}CO$_2$ labeling
strategy (C1 and the C4 of the OAA backbone are lost during Citrate to Succinate conversion (**Figure
6A**)).

Here, we assessed the flux modes of the TCA Cycle by taking advantage of pathway-specific mass
isotopologues/isotopomers and a ^{13}C-MFA approach after a U-^{13}C-Pyruvate incorporation in *B. napus*
leaf discs. This approach was notably based on the simultaneous application of ^{13}C labeling thanks to
the PDH that produced ^{13}C$_2$-AcetylCoA for the TCA cycle and ^{13}CO$_2$ for the PEPc. Interestingly, we
showed that Glutamate and Succinate rapidly harbored M2 istopologues, corresponding to *de novo*
incorporation of M2 Acetyl-CoA from the TCA Cycle, followed by an increase of M1 and M3
isotopologues that were clearly visible after only 2 hours. These isotopologues corresponded to M2
isotopologues that had experienced a complete turn of the forward cyclic mode of the TCA cycle. They
provided for the first time in plants a unique ^{13}C-labeling signature following U-^{13}C-pyruvate
incorporation that demonstrated the cyclic nature of the plant TCA cycle in *B. napus* leaves (**Figure
5**). From these results, we combined a ^{13}C-MFA approach with pathway specific MS fragments and
the indirect measurement of M2/(M0+M2) ratio for the Acetyl-CoA active pool to show that the
forward cyclic mode of the TCA cycle contributed to nearly 80% of both Malate and Citrate production
compared to PEPc (4:1 activity ratio) in both light and dark conditions with our experimental setup,
while malic enzyme activity was not significantly detected (**Figure 6**). One may question the use of
pyruvate for ^{13}C labeling strategies, since it may introduce an artificial increase of TCA cycle turn-
over compared to experiments with ^{13}CO$_2$. This could ultimately lead to important ^{13}C enrichments for
Succinate and the rapid apparition of M1 and M3 isotopologues for Glutamate and Succinate. A recent

incorporation of 15 mM U-^{13}C-pyruvate or 15 mM U-^{13}C-glucose into Arabidopsis with a petiole
feeding assay during 4 hours in the light identified important ^{13}C enrichments in Citrate, Glutamate
and Succinate (Zhang et al., 2021). However, another experiment (Arabidopsis leaf discs, 20 mM U-
^{13}C-pyruvate, up to 8 hours in the dark) showed similar results but only observed M2 isotopologues
for Citrate, Glutamate and Succinate and not M1 and M3 isotopologues which were expected for a
cyclic mode (complete dataset available on request from the authors) (Le et al., 2021). Thus, the use
of pyruvate does not seem to artificially increase the turn-over of the TCA cycle in Arabidopsis, *i.e.*
metabolites experiencing multiples turns of the TCA cycle. Therefore, the forward cyclic mode of the
TCA cycle has definitely a higher contribution to Malate biosynthesis compared to PEPc in *B. napus*
leaves. Nevertheless, this TCA cycle/PEPc ratio (4:1) may have been slightly overestimated in our
study since this pathway fractional contribution assumes that most of the CO_2 is produced by the PDH,
and its conversion to HCO_3^- are not rate-limiting for PEPc activity. While the second hypothesis may
not be so constraining given that CO_2 was actively reassimilated by photosynthesis in our light
conditions (**Figure 7**), the second hypothesis can be questioned given the metabolic network (**Figure
6**). Indeed, isocitrate and 2-oxoglutarate dehydrogenases necessarily decarboxylate more $^{12}CO_2$ than
$^{13}CO_2$ compared to the PDH (see mean ^{13}C enrichment of metabolically active pool of Acetyl-CoA,
Figure 6B). This could imbalance the reassimilation of $^{13}CO_2$ by PEPc if the first part of the TCA
cycle operates at a rate close to the PEPc activity. Assuming an equivalent metabolic flux from PDH
to 2-oxoglutarate dehydrogenase based on previous works with $^{13}CO_2$ (Xu et al., 2021;Xu et al., 2022),
the overall mitochondrial $^{12}CO_2/^{13}CO_2$ ratio should not exceed a 2:1 dilution compared to the
$^{12}CO_2/^{13}CO_2$ produced by the PDH (see CID of Citrate, Glutamate and Succinate and their associated
CO_2 losses). Thus, the forward cyclic mode of the TCA cycle will still majorly contribute to Malate
and Citrate production compared to PEPc in our experimental setup with this scenario (a 2:1 ratio is
expected) (**Figure 8**). Interestingly, a ^{13}C-MFA performed at isotopic and metabolic steady-state in
heterotrophic suspension cells culture incubated with U-^{13}C-glucose showed a similar ratio of 2:1 for
forward TCA cycle/PEPc metabolic fluxes (ana1 versus tca2, tca3, tca4, tca5), with a negligible malic
enzyme activity (10-fold less than PEPc) (Masakapalli et al., 2010).

Besides this, the forward cyclic mode of the TCA cycle has to be reconnected to the relative ^{13}C content
found in Glutamate and Aspartate in our study. This can represent important escapes for ^{13}C, thereby
driving the relative differences of metabolic flux for the biosynthesis of Glutamate and Succinate, and
Malate and OAA respectively (branching points) (**Figure 5**). Nevertheless, a significant proportion of
Glutamate and Aspartate may come back to the TCA cycle. In astrocyte cells, U-^{13}C-Glutamate
labeling incorporations showed that Glutamate M5 was rapidly metabolized into Glutamate M3 after
one turn of the TCA cycle (Nissen et al., 2015) thereby proposing a similar role for plant mitochondrial
Glutamate dehydrogenase (Terce-Laforgue et al., 2013). The potential role of GABA (Gamma-
aminobutyric acid) shunt has been proposed to reroute Glutamate towards Succinate (Tcherkez et al.,
2009). Finally, mitochondrial transaminases such as Alanine:2-oxoglutarate and Aspartate:2-
oxoglutarate could also participle in this process. Indeed, Alanine:2-oxoglutarate aminotransferase is
one of the three pathway supplying pyruvate to mitochondria and was notably solicited in the *mpc1*
mutants to compensate for the loss of mitochondrial pyruvate import (Le et al., 2021). Regarding
Glutamate, its production from ^{13}C-2-Oxoglutarate and ^{12}C-Glutamine molecules in our metabolic
network will necessarily introduce a 50% isotopic dilution between ^{12}C and ^{13}C molecules. Thus, its
reassimilation will also introduce this 50% isotopic dilution to mitochondrial active pools of either
Succinate or 2-Oxoglutarate. For instance, it is not possible to have access to the size of the
mitochondrial active pool for both metabolites and by extension their exact ^{13}C-enrichments without a
complete isotopic steady-state of the plant TCA cycle and perhaps subcellular fractionation or very fast
mitochondrial purification by centrifugation (Lee et al., 2019;Destailleur et al., 2021). To which extent
Glutamate and Aspartate molecules escape the TCA cycle remains to be determined.

Light/dark regulation of PEPc and TCA cycle contributions to Malate and Citrate biosynthesis
in *Brassica napus* leaves

PEPc activity is positively regulated by its light-dependent thioredoxin-mediated phosphorylation in
C_3 and C_4 plants (Caburatan and Park, 2021). Thus, PEPc is expected to be more active in the light
than in the dark, as recently suggested in guard cell cultures based on the differential ^{13}C enrichment
found for the C1 of Glutamate (Lima et al., 2021). Here, we focused our analysis on both Malate and
Citrate to compare PEPc and TCA cycle contribution to their biosynthesis. We notably showed that
Malate_3TMS_C1C4 fragment experienced a strong shift in its M1/M2 isotopologues in light
conditions compared to dark conditions (**Figure 5**). This was associated to a relatively higher
contribution of PEPc to Malate production in light conditions compared to dark conditions (**Figure**
6B). However, such shift could be attributed to both increased PEPc activity and TCA cycle inhibition
in the light. In our experiments, we found very similar dynamics of ^{13}C-enrichment for TCA cycle
metabolites in both dark and light conditions combined to similar metabolic pool sizes in a metabolic
steady-state (**Figure 4**). Thus, variations of TCA cycle metabolic fluxes between light and dark
conditions are expected to be very low in our experimental set-up. Since we used 10 mM U-^{13}C-
pyruvate for labeling experiments, perhaps this has activated the PDH thereby eliminating one of the
major bottlenecks regulating the entry of carbon for plant TCA cycle. A previous work based on $^{13}CO_2$
decarboxylations with either ^{13}C-pyruvate or ^{13}C-glucose showed that PDH was inhibited by up to 27%
in the light compared to the dark, while glycolysis could be inhibited by up to 95% (Tcherkez et al.,
2005). However, these inhibitions were necessarily underestimated by the measurement method which
did not considered the reassimilation of $^{13}CO_2$ by the RuBiSCO. Here, by integrating a ^{13}C-positionnal
isotopomer approach to the photorespiratory metabolism, we showed that PDH and TCA-cycle derived
CO_2 were actively reassimilated by the RuBiSCO, leading to a contribution of around 5% to net CO_2
assimilation (**Figure 7**). Under the light irradiance of around 120 µmol photons.m^{-2}.s^{-1} (same light
intensity of our experimental setup), net CO_2 uptake of *B. napus* leaves at a similar developmental
stage (L11 and L7 leaves) is close to 5 µmol CO_2. m^{-2}.s^{-1} (Dellero et al., 2021), leading to a
reassimilation rate of 0.25 µmol CO_2. m^{-2}.s^{-1}. Since the dark respiration rate for these leaves is close to
1.25 µmol CO_2. m^{-2}.s^{-1} (Dellero et al., 2021), and given the proposed inhibition of mitochondrial
respiration by nearly 50% in the light (Tcherkez et al., 2005), this will give a day respiration rate of
0.625 µmol CO_2. m^{-2}.s^{-1}, but without considering the reassimilation rate of PDH and TCA-derived
decarboxylations. Therefore, the current measurement/estimation of day respiration rate through CO_2
mass balance at the leaf level (Tcherkez et al., 2017;Berghuijs et al., 2019) must be underestimated by
nearly 40% (0.875 versus 0.625). Thus, the inhibition of day respiration at the metabolic flux level is
likely be lower than measured by gas-exchanges and could only represent 30% in our experiments
(1.25 in the dark versus 0.875 in the light) (**Figure 8**). Besides this, the *pdk* mutant responsible for the
activation of the PDH in the light was recently used for ^{13}C labeling experiments. After a U-^{13}C-
pyruvate incorporation into Arabidopsis leaves during 4 hours in the light, the *pdk* mutant had higher
^{13}C-enrichments for Citrate, Glutamate and Succinate compared to the control (Zhang et al., 2021).
However, there were no differences when using U-^{13}C-Glucose. This result questioned the regulatory
power of the PDH for the activity of the TCA cycle in the light, especially when carbon arising from
glycolysis is not limiting. Based on these observations, we think that the similar dynamics of ^{13}C-
enrichment observed in our experiments for light and dark conditions are essentially explained by the
elimination of the light/dark regulation of glycolysis. Since the inhibition of PDH and TCA cycle
decarboxylations in the light (50%) is supposed to be close to 30% only at the metabolic flux level, the
fractional contributions calculated before have necessarily overestimated the contribution of the TCA
cycle in light conditions by up to 30% compared to dark conditions. Nevertheless, the forward cyclic
flux mode of the TCA cycle will still have a major contribution to Malate and Citrate biosynthesis
compared to PEPc (ratio 4:1 would be close to 3:1 after correction). Therefore, the shift in M1/M2

isotopologues of Malate_3TMS_C1C4 fragment in light conditions compared to dark conditions
essentially captured the light/dark regulation of PEPc.

Besides this, we observed another unexpected but interesting result: the higher contribution of PEPc
derived-carbons to Citrate production in the dark condition compared to the light condition. Since we
have shown that PEPc contribution to Malate production was reduced in the dark compared to the light,
this result can only be explained by a higher transport of OAA/Malate from the cytosol to the
mitochondria in dark conditions (**Figure 8**). This idea is supported by the fact that PEPc is majorly
localized to the cytosol (Caburatan and Park, 2021) and the recent identification of a mitochondrial
malate importer in Arabidopsis with an important impact on mitochondrial respiration in the dark (Lee
et al., 2021). Whether a higher malate import into mitochondria allows to increase the absolute
metabolic flux of the TCA cycle in dark conditions in our experimental set-up remains to be elucidated.

Conclusions

Overall, we proposed a robust workflow to evaluate the accuracy of CID measurements by GC-MS for
organic and amino acids and successfully applied our validated method to the investigation of TCA
cycle and PEPc metabolic fluxes in *Brassica napus* leaves. Notably, our results did not support the
hypothesis of the stored citrate contribution to TCA cycle in a light/dark-dependent manner and
demonstrated that the forward cyclic flux mode of the TCA cycle largely contributed to Malate and
Citrate production compared to PEPc (4:1 ratio) in *B. napus* leaves. Thus, the activity of the succinate
dehydrogenase in both dark and light conditions has been largely underestimated until now.
Reassimilation of PDH and TCA-cycle derived CO_2 by photosynthesis represented nearly 5% of the
net CO_2 assimilation rate in our conditions, and this should be considered when day respiration rate is
measured by CO_2 mass balance at the leaf level (underestimation by nearly 40%). Although the
regulation of the PEPc activity by light was still operating, the higher involvement of PEPc-derived
carbons to citrate production in dark conditions proposed an important role for Malate import into
mitochondria in sustaining TCA cycle in dark conditions. The next stage of this work will be focused
on the accurate modeling of this metabolic network to infer absolute metabolic fluxes. This perspective
sounds stimulating given the development of multiple modeling approaches over the last decade but
still remains highly challenging (Kruger and Ratcliffe, 2021).

5   Conflict of Interest

835   The authors declare that the research was conducted in the absence of any commercial or financial
836   relationships that could be construed as a potential conflict of interest.

6   Author Contributions

838   Conceptualization, Y.D., A.B.; methodology, Y.D..; software, Y.D., O.F.; validation, Y.D., S.B. and
839   C.B.; formal analysis, Y.D.; investigation, Y.D., S.B. and C.B.; data curation, Y.D.; writing-original
840   draft preparation, Y.D.; writing-review and editing, Y.D., A.B.; visualization, Y.D.; supervision,
841   Y.D.; project administration, Y.D.; funding acquisition, Y.D., A.B. All authors have read and agreed
842   to the published version of the manuscript.

7   Funding

This research was supported by the IB2021_TRICYCLE project funded by the INRAE BAP division
and by the METABOHUB project funded by the program Investments for the Future of the French
national agency for research (ANR grant number 11-INBS-0010).

8 Acknowledgments

P2M2 (Metabolic Profiling & Metabolomics Platform, Le Rheu, France, www6.inrae.fr/p2m2/) and its
staff members are gratefully acknowledged for access to mass spectrometry facilities and for
supporting the methodological developments required for this study. We thank MetaToul
(Metabolomics & Fluxomics Facitilies, Toulouse, France, www.metatoul.fr) for technical support and
access to NMR facilities. We are grateful to Nathalie Marnet and Mathieu Aubert for technical
assistance in the quantification of amino acid content and Pierre Millard and Floriant Bellvert for
fruitful scientific discussions.

9 Supplementary Material

Supplementary Material available with the files deposited.

10 References

Abadie, C., Bathellier, C., and Tcherkez, G. (2018). Carbon allocation to major metabolites in
illuminated leaves is not just proportional to photosynthesis when gaseous conditions (CO_2
and O_2) vary. *New Phytol* 218**,** 94-106.

Abadie, C., Boex-Fontvieille, E.R., Carroll, A.J., and Tcherkez, G. (2016). In vivo stoichiometry of
photorespiratory metabolism. *Nat Plants* 2**,** 15220.

Abadie, C., Lothier, J., Boex-Fontvieille, E., Carroll, A., and Tcherkez, G. (2017). Direct assessment
of the metabolic origin of carbon atoms in glutamate from illuminated leaves using (13) C-
NMR. *New Phytol* 216**,** 1079-1089.

Abadie, C., and Tcherkez, G. (2019a). In vivo phosphoenolpyruvate carboxylase activity is
controlled by CO_2 and O_2 mole fractions and represents a major flux at high photorespiration
rates. *New Phytol* 221**,** 1843-1852.

Abadie, C., and Tcherkez, G. (2019b). Plant sulphur metabolism is stimulated by photorespiration.
Commun Biol 2**,** 379.

Abadie, C., and Tcherkez, G. (2021). (13)C Isotope Labelling to Follow the Flux of Photorespiratory
Intermediates. *Plants (Basel)* 10.

Allen, D.K., and Young, J.D. (2019). Tracing metabolic flux through time and space with isotope
labeling experiments. *Curr Opin Biotechnol* 64**,** 92-100.

Alves, T.C., Pongratz, R.L., Zhao, X., Yarborough, O., Sereda, S., Shirihai, O., Cline, G.W., Mason,
G., and Kibbey, R.G. (2015). Integrated, Step-Wise, Mass-Isotopomeric Flux Analysis of the
TCA Cycle. *Cell Metab* 22**,** 936-947.

Antoniewicz, M.R. (2021). A guide to metabolic flux analysis in metabolic engineering: Methods,
tools and applications. *Metab Eng* 63**,** 2-12.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2006). Determination of confidence
intervals of metabolic fluxes estimated from stable isotope measurements. *Metab Eng* 8**,** 324-
337.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2007). Accurate Assessment of Amino
Acid Mass Isotopomer Distributions for Metabolic Flux Analysis. *Anal Chem* 79**,** 6.

Araujo, W.L., Nunes-Nesi, A., Nikoloski, Z., Sweetlove, L.J., and Fernie, A.R. (2012). Metabolic
control and regulation of the tricarboxylic acid cycle in photosynthetic and heterotrophic plant
tissues. *Plant Cell Environ* 35**,** 1-21.

Araujo, W.L., Tohge, T., Nunes Nesi, A., Obata, T., and Fernie, A. (2014). "Analysis of Kinetic
Labeling of Amino Acids and Organic Acids by GC-MS," in *Plant Metabolic Flux Analysis:*
Methods and Protocols, ed. M.I.M. Biology.).

Berghuijs, H.N.C., Yin, X., Ho, Q.T., Retta, M.A., Nicolai, B.M., and Struik, P.C. (2019). Using a
reaction-diffusion model to estimate day respiration and reassimilation of (photo)respired
CO2 in leaves. *New Phytol* 223**,** 619-631.

Bianchetti, G., Baron, C., Carrillo, A., Berardocco, S., Marnet, N., Wagner, M.H., Demilly, D.,
Ducournau, S., Manzanares-Dauleux, M.J., Caherec, F.L., Buitink, J., and Nesi, N. (2021).
Dataset for the metabolic and physiological characterization of seeds from oilseed rape
(Brassica napus L.) plants grown under single or combined effects of drought and clubroot
pathogen Plasmodiophora brassicae. *Data Brief* 37**,** 107247.

Buescher, J.M., Antoniewicz, M.R., Boros, L.G., Burgess, S.C., Brunengraber, H., Clish, C.B.,
Deberardinis, R.J., Feron, O., Frezza, C., Ghesquiere, B., Gottlieb, E., Hiller, K., Jones, R.G.,
Kamphorst, J.J., Kibbey, R.G., Kimmelman, A.C., Locasale, J.W., Lunt, S.Y., Maddocks,
O.D., Malloy, C., Metallo, C.M., Meuillet, E.J., Munger, J., Noh, K., Rabinowitz, J.D.,
Ralser, M., Sauer, U., Stephanopoulos, G., St-Pierre, J., Tennant, D.A., Wittmann, C., Vander
Heiden, M.G., Vazquez, A., Vousden, K., Young, J.D., Zamboni, N., and Fendt, S.M. (2015).
A roadmap for interpreting (13)C metabolite labeling patterns from cells. *Curr Opin*
Biotechnol 34**,** 189-201.

Caburatan, L., and Park, J. (2021). Differential Expression, Tissue-Specific Distribution, and
Posttranslational Controls of Phosphoenolpyruvate Carboxylase. *Plants (Basel)* 10.

Cheah, Y.E., Xu, Y., Sacco, S.A., Babele, P.K., Zheng, A.O., Johnson, C.H., and Young, J.D. (2020).
Systematic identification and elimination of flux bottlenecks in the aldehyde production
pathway of Synechococcus elongatus PCC 7942. *Metab Eng* 60**,** 56-65.

Cheah, Y.E., and Young, J.D. (2018). Isotopically nonstationary metabolic flux analysis (INST-
MFA): putting theory into practice. *Curr Opin Biotechnol* 54**,** 80-87.

Cheung, C.Y., Poolman, M.G., Fell, D.A., Ratcliffe, R.G., and Sweetlove, L.J. (2014). A Diel Flux
Balance Model Captures Interactions between Light and Dark Metabolism during Day-Night
Cycles in C3 and Crassulacean Acid Metabolism Leaves. *Plant Physiol* 165**,** 917-929.

Clark, T.J., Guo, L., Morgan, J., and Schwender, J. (2020). Modeling Plant Metabolism: From
Network Reconstruction to Mechanistic Models. *Annu Rev Plant Biol*.

Da Fonseca-Pereira, P., Souza, P.V.L., Fernie, A.R., Timm, S., Daloso, D.M., and Araujo, W.L.
(2021). Thioredoxin-mediated regulation of (photo)respiration and central metabolism. *J Exp*
Bot 72**,** 5987-6002.

Daloso, D.M., Medeiros, D.B., Dos Anjos, L., Yoshida, T., Araujo, W.L., and Fernie, A.R. (2017).
Metabolism within the specialized guard cells of plants. *New Phytol* 216**,** 1018-1033.

Dellero, Y. (2020). Manipulating amino acid metabolism to improve crop nitrogen use efficiency for
a sustainable agriculture. *Frontiers in Plant Science* 11**,** 1857

Dellero, Y., Clouet, V., Marnet, N., Pellizzaro, A., Dechaumet, S., Niogret, M.F., and Bouchereau,
A. (2020a). Leaf status and environmental signals jointly regulate proline metabolism in
winter oilseed rape. *J Exp Bot* 71**,** 2098-2111.

Dellero, Y., and Filangi, O. (2021). "Corrective method dedicated to Isocor for calculating carbon
isotopologue distribution from GCMS runs". V1 ed.: Portail Data INRAE).

Dellero, Y., Heuillet, M., Marnet, N., Bellvert, F., Millard, P., and Bouchereau, A. (2020b).
Sink/Source Balance of Leaves Influences Amino Acid Pools and Their Associated Metabolic
Fluxes in Winter Oilseed Rape (Brassica napus L.). *Metabolites* 10**,** 16.

Dellero, Y., Jossier, M., Bouchereau, A., Hodges, M., and Leport, L. (2021). Leaf Phenological
Stages of Winter Oilseed Rape (Brassica napus L.) Have Conserved Photosynthetic
Efficiencies but Contrasted Intrinsic Water Use Efficiencies at High Light Intensities.
Frontiers in Plant Science 12.

Dellero, Y., Jossier, M., Glab, N., Oury, C., Tcherkez, G., and Hodges, M. (2016). Decreased
glycolate oxidase activity leads to altered carbon allocation and leaf senescence after a
transfer from high CO2 to ambient air in Arabidopsis thaliana. *J Exp Bot* 67**,** 3149-3163.

Destailleur, A., Poucet, T., Cabasson, C., Alonso, A.P., Cocuron, J.C., Larbat, R., Vercambre, G.,
Colombie, S., Petriacq, P., Andrieu, M.H., Beauvoit, B., Gibon, Y., and Dieuaide-Noubhani,
M. (2021). The Evolution of Leaf Function during Development Is Reflected in Profound
Changes in the Metabolic Composition of the Vacuole. *Metabolites* 11.

Dong, H., Bai, L., Zhang, Y., Zhang, G., Mao, Y., Min, L., Xiang, F., Qian, D., Zhu, X., and Song,
C.P. (2018). Modulation of Guard Cell Turgor and Drought Tolerance by a Peroxisomal
Acetate-Malate Shunt. *Mol Plant* 11**,** 1278-1291.

Evers, B., Gerding, A., Boer, T., Heiner-Fokkema, M.R., Jalving, M., Wahl, S.A., Reijngoud, D.J.,
and Bakker, B.M. (2021). Simultaneous Quantification of the Concentration and Carbon
Isotopologue Distribution of Polar Metabolites in a Single Analysis by Gas Chromatography
and Mass Spectrometry. *Anal Chem* 93**,** 8248-8256.

Galili, G., Amir, R., and Fernie, A.R. (2016). The Regulation of Essential Amino Acid Synthesis and
Accumulation in Plants. *Annu Rev Plant Biol* 67**,** 153-178.

Gauthier, P.P., Bligny, R., Gout, E., Mahe, A., Nogues, S., Hodges, M., and Tcherkez, G.G. (2010).
In folio isotopic tracing demonstrates that nitrogen assimilation into glutamate is mostly
independent from current CO2 assimilation in illuminated leaves of Brassica napus. *New
Phytol* 185**,** 988-999.

Harvey, D.J., and Vouros, P. (2020). Mass Spectrometric Fragmentation of Trimethylsilyl and
Related Alkylsilyl Derivatives. *Mass Spectrom Rev* 39**,** 105-211.

Hay, J.O., Shi, H., Heinzel, N., Hebbelmann, I., Rolletschek, H., and Schwender, J. (2014).
Integration of a constraint-based metabolic model of Brassica napus developing seeds with
(13)C-metabolic flux analysis. *Front Plant Sci* 5**,** 724.

Heise, R., Fernie, A.R., Stitt, M., and Nikoloski, Z. (2015). Pool size measurements facilitate the
determination of fluxes at branching points in non-stationary metabolic flux analysis: the case
of Arabidopsis thaliana. *Front Plant Sci* 6**,** 386.

Heuillet, M., Bellvert, F., Cahoreau, E., Letisse, F., Millard, P., and Portais, J.C. (2018).
Methodology for the Validation of Isotopic Analyses by Mass Spectrometry in Stable-Isotope
Labeling Experiments. *Anal Chem* 90**,** 1852-1860.

Heuillet, M., Millard, P., Cisse, M.Y., Linares, L.K., Letisse, F., Manie, S., Le Cam, L., Portais, J.C.,
and Bellvert, F. (2020). Simultaneous Measurement of Metabolite Concentration and Isotope
Incorporation by Mass Spectrometry. *Anal Chem*.

Kambhampati, S., Li, J., Evans, B.S., and Allen, D.K. (2019). Accurate and efficient amino acid
analysis for protein quantification using hydrophilic interaction chromatography coupled
tandem mass spectrometry. *Plant Methods* 15**,** 46.

Kruger, N.J., Huddleston, J.E., Le Lay, P., Brown, N.D., and Ratcliffe, R.G. (2007). Network flux
analysis: impact of 13C-substrates on metabolism in Arabidopsis thaliana cell suspension
cultures. *Phytochemistry* 68**,** 2176-2188.

Kruger, N.J., and Ratcliffe, R.G. (2015). Fluxes through plant metabolic networks: measurements,
predictions, insights and challenges. *Biochem J* 465**,** 27-38.

Kruger, N.J., and Ratcliffe, R.G. (2021). Whither metabolic flux analysis in plants? *Journal of*
Experimental Botany.

Lai, Z., and Fiehn, O. (2018). Mass spectral fragmentation of trimethylsilylated small molecules.
Mass Spectrom Rev 37**,** 245-257.

Le, X.H., Lee, C.P., and Millar, A.H. (2021). The mitochondrial pyruvate carrier (MPC) complex
mediates one of three pyruvate-supplying pathways that sustain Arabidopsis respiratory
metabolism. *Plant Cell* 33**,** 2776-2793.

Lee, C.P., Elsasser, M., Fuchs, P., Fenske, R., Schwarzlander, M., and Millar, A.H. (2021). The
versatility of plant organic acid metabolism in leaves is underpinned by mitochondrial malate-
citrate exchange. *Plant Cell* 33**,** 3700-3720.

Lee, W.D., Mukha, D., Aizenshtein, E., and Shlomi, T. (2019). Spatial-fluxomics provides a
subcellular-compartmentalized view of reductive glutamine metabolism in cancer cells. *Nat*
Commun 10**,** 1351.

Lima, V.F., Erban, A., Daubermann, A.G., Freire, F.B.S., Porto, N.P., Candido-Sobrinho, S.A.,
Medeiros, D.B., Schwarzlander, M., Fernie, A.R., Dos Anjos, L., Kopka, J., and Daloso,
D.M. (2021). Establishment of a GC-MS-based (13) C-positional isotopomer approach
suitable for investigating metabolic fluxes in plant primary metabolism. *Plant J*.

Lima, V.F., Perez Souza, L., Williams, T.C., Fernie, A., and Daloso, D.M. (2018). "Gas
Chromatography–Mass Spectrometry-Based 13C-Labeling Studies in Plant Metabolomics,"
in *Plant Metabolomics: Methods and Protocol,* ed. M.I.M. Biology.).

Ma, F., Jazmin, L.J., Young, J.D., and Allen, D.K. (2014). Isotopically nonstationary 13C flux
analysis of changes in Arabidopsis thaliana leaf metabolism due to high light acclimation.
Proc Natl Acad Sci U S A 111**,** 16967-16972.

Masakapalli, S.K., Le Lay, P., Huddleston, J.E., Pollock, N.L., Kruger, N.J., and Ratcliffe, R.G.
(2010). Subcellular flux analysis of central metabolism in a heterotrophic Arabidopsis cell
suspension using steady-state stable isotope labeling. *Plant Physiol* 152**,** 602-619.

Medeiros, D.B., Perez Souza, L., Antunes, W.C., Araujo, W.L., Daloso, D.M., and Fernie, A.R.
(2018). Sucrose breakdown within guard cells provides substrates for glycolysis and
glutamine biosynthesis during light-induced stomatal opening. *Plant J* 94**,** 583-594.

Millard, P., Cahoreau, E., Heuillet, M., Portais, J.C., and Lippens, G. (2017). (15)N-NMR-Based
Approach for Amino Acids-Based (13)C-Metabolic Flux Analysis of Metabolism. *Anal Chem*
89**,** 2101-2106.

Millard, P., Delepine, B., Guionnet, M., Heuillet, M., Bellvert, F., and Letisse, F. (2019). IsoCor:
isotope correction for high-resolution MS labeling experiments. *Bioinformatics* 35**,** 4484-
4487.

Millard, P., Massou, S., Portais, J.C., and Letisse, F. (2014). Isotopic studies of metabolic systems by
mass spectrometry: using Pascal's triangle to produce biological standards with fully
controlled labeling patterns. *Anal Chem* 86**,** 10288-10295.

Millard, P., Schmitt, U., Kiefer, P., Vorholt, J.A., Heux, S., and Portais, J.-C. (2020). ScalaFlux: a
scalable approach to quantify fluxes in metabolic subnetworks. *Plos Computational Biology*
16**,** e1007799.

Nissen, J.D., Pajecka, K., Stridh, M.H., Skytt, D.M., and Waagepetersen, H.S. (2015). Dysfunctional
TCA-Cycle Metabolism in Glutamate Dehydrogenase Deficient Astrocytes. *Glia* 63**,** 2313-
2326.

Okahashi, N., Kawana, S., Iida, J., Shimizu, H., and Matsuda, F. (2019). Fragmentation of
Dicarboxylic and Tricarboxylic Acids in the Krebs Cycle Using GC-EI-MS and GC-EI-
MS/MS. *Mass Spectrom (Tokyo)* 8**,** A0073.

Ratcliffe, R.G., and Shachar-Hill, Y. (2006). Measuring multiple fluxes through plant metabolic
networks. *Plant J* 45**,** 490-511.

Rstudio Team (2016). RStudio: Integrated Development for R.

.

Sinnaeve, D., Dinclaux, M., Cahoreau, E., Millard, P., Portais, J.C., Letisse, F., and Lippens, G.
(2018). Improved Isotopic Profiling by Pure Shift Heteronuclear 2D J-Resolved NMR
Spectroscopy. *Anal Chem* 90**,** 4025-4031.

Sweetlove, L.J., Beard, K.F., Nunes-Nesi, A., Fernie, A.R., and Ratcliffe, R.G. (2010). Not just a
circle: flux modes in the plant TCA cycle. *Trends Plant Sci* 15**,** 462-470.

Szecowka, M., Heise, R., Tohge, T., Nunes-Nesi, A., Vosloh, D., Huege, J., Feil, R., Lunn, J.,
Nikoloski, Z., Stitt, M., Fernie, A.R., and Arrivault, S. (2013). Metabolic fluxes in an
illuminated Arabidopsis rosette. *Plant Cell* 25**,** 694-714.

Tcherkez, G., Cornic, G., Bligny, R., Gout, E., and Ghashghaie, J. (2005). In vivo respiratory
metabolism of illuminated leaves. *Plant Physiol* 138**,** 1596-1606.

Tcherkez, G., Gauthier, P., Buckley, T.N., Busch, F.A., Barbour, M.M., Bruhn, D., Heskel, M.A.,
Gong, X.Y., Crous, K.Y., Griffin, K., Way, D., Turnbull, M., Adams, M.A., Atkin, O.K.,
Farquhar, G.D., and Cornic, G. (2017). Leaf day respiration: low CO2 flux but high
significance for metabolism and carbon balance. *New Phytol* 216**,** 986-1001.

Tcherkez, G., Mahe, A., Gauthier, P., Mauve, C., Gout, E., Bligny, R., Cornic, G., and Hodges, M.
(2009). In folio respiratory fluxomics revealed by 13C isotopic labeling and H/D isotope
effects highlight the noncyclic nature of the tricarboxylic acid "cycle" in illuminated leaves.
Plant Physiol 151**,** 620-630.

Terce-Laforgue, T., Bedu, M., Dargel-Grafin, C., Dubois, F., Gibon, Y., Restivo, F.M., and Hirel, B.
(2013). Resolving the role of plant glutamate dehydrogenase: II. Physiological
characterization of plants overexpressing the two enzyme subunits individually or
simultaneously. *Plant Cell Physiol* 54**,** 1635-1647.

Timm, S. (2020). The impact of photorespiration on plant primary metabolism through metabolic and
redox regulation. *Biochemical Society Transactions* 48**,** 2495-2504.

Treves, H., Kuken, A., Arrivault, S., Ishihara, H., Hoppe, I., Erban, A., Hohne, M., Moraes, T.A.,
Kopka, J., Szymanski, J., Nikoloski, Z., and Stitt, M. (2022). Carbon flux through
photosynthesis and central carbon metabolism show distinct patterns between algae, C3 and
C4 plants. *Nat Plants* 8**,** 78-91.

Wieloch, T. (2021). The next phase in the development of 13C isotopically nonstationary metabolic
flux analysis. *J Exp Bot*.

Xu, G., Fan, X., and Miller, A.J. (2012). Plant nitrogen assimilation and use efficiency. *Annu Rev
Plant Biol* 63**,** 153-182.

Xu, Y., Fu, X., Sharkey, T.D., Shachar-Hill, Y., and Walker, B.J. (2021). The metabolic origins of
non-photorespiratory CO2 release during photosynthesis: A metabolic flux analysis. *Plant
Physiol*.

Xu, Y., Wieloch, T., Kaste, J.a.M., Shachar-Hill, Y., and Sharkey, T.D. (2022). Reimport of carbon
from cytosolic and vacuolar sugar pools into the Calvin–Benson cycle explains
photosynthesis labeling anomalies. *Proceedings of the National Academy of Sciences* 119.

Young, J.D., Allen, D.K., and Morgan, J.A. (2014). "Isotopomer measurement techniques in
metabolic flux analysis II: mass spectrometry," in *Methods Mol Biol*.), 85-108.

Zhang, Y., and Fernie, A.R. (2018). On the role of the tricarboxylic acid cycle in plant productivity. *J
Integr Plant Biol* 60**,** 1199-1216.

Zhang, Y., Giese, J., Kerbler, S.M., Siemiatkowska, B., Perez De Souza, L., Alpers, J., Medeiros,
D.B., Hincha, D.K., Daloso, D.M., Stitt, M., Finkemeier, I., and Fernie, A.R. (2021). Two
mitochondrial phosphatases, PP2c63 and Sal2, are required for posttranslational regulation of
the TCA cycle in Arabidopsis. *Mol Plant* 14**,** 1104-1118.

1077

1078

1079

1080

1081 **Table 1.** Molecular ions selected for Carbon Isotopologue Distribution analysis of organic and amino
1082 acid TMS-derivatives by GC-MS.

Compounds	TMS derivative		Loss		Molecular ion		
	Formula	mass	Formula	mass	Formula	m/z	Carbon skeleton
Organic acids							
Citrate(4TMS)	C18H40O7Si4	480	CH3	15	C17H37O7Si4	465	1-2-3-4-5-6
Citrate(4TMS)	C18H40O7Si4	480	TMS-CO2, TMS-OH	207	C11H21O4Si2	273	1-2-3-4-5
Fumarate(2TMS)	C10H20O4Si2	260	CH3	15	C9H17O4Si2	245	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	CH3	15	C12H27O5Si3	335	1-2-3-4
Malate(3TMS)	C13H30O5Si3	350	TMS-CO2	117	C9H21O3Si2	233	2-3-4
Pyruvate(MEOX)(TMS)	C7H15NO3Si	189	CH3	15	C6H12NO3Si	174	1-2-3
Succinate(2TMS)	C10H22O4Si2	262	CH3	15	C9H19O4Si2	247	1-2-3-4
Amino acids							
Asparagine(2TMS)	C10H24N2O3Si2	276	TMS-CO2	117	C6H15N2OSi	159	2-3-4
Asparagine(3TMS)	C13H32N2O3Si3	348	TMS-CO2	117	C9H23N2OSi2	231	2-3-4
Aspartate(2TMS)	C10H23NO4Si2	277	TMS-CO2	117	C6H14NO2Si	160	2-3-4
Aspartate(3TMS)	C13H31NO4Si3	349	TMS-CO2	117	C9H22NO2Si2	232	2-3-4
Glutamate(3TMS)	C14H33NO4Si3	363	TMS-CO2	117	C10H24NO2Si2	246	2-3-4-5
Glutamine(3TMS)	C14H34N2O3Si3	362	TMS-CO2	117	C10H25N2OSi2	245	2-3-4-5
Glycine(3TMS)	C11H29NO2Si3	291	TMS-CO2	117	C7H20NSi2	174	2
Isoleucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Leucine(2TMS)	C12H29NO2Si2	275	TMS-CO2	117	C8H20NSi	158	2-3-4-5-6
Lysine(4TMS)	C18H46N2O2Si4	434	TMS-CO2	117	C14H37N2Si3	317	2-3-4-5-6
Methionine(2TMS)	C11H27NO2SSi2	293	TMS-CO2	117	C7H18NSSi	176	2-3-4-5
Ornithine(4TMS)	C17H44N2O2Si4	420	none		none		1-2-3-4-5
Phenylalanine(2TMS)	C15H27NO2Si2	309	TMS-CO2	117	C11H18NSi	192	2-3-4-5-6-7-8-9
Phenylalanine(2TMS)	C15H27NO2Si2	309	C7H7	91	C8H20NO2Si2	218	1-2
Proline(2TMS)	C11H25NO2Si2	259	TMS-CO2	117	C7H16NSi	142	2-3-4-5
Serine(2TMS)	C9H23NO3Si2	249	TMS-CO2	117	C5H14NOSi	132	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CO2	117	C8H22NOSi2	204	2-3
Serine(3TMS)	C12H31NO3Si3	321	TMS-CH2O	103	C8H20NO2Si2	218	1-2
Threonine(3TMS)	C13H33NO3Si3	335	TMS-CO2	117	C9H24NOSi2	218	2-3-4
Tryptophan(2TMS)	C17H28N2O2Si2	348	2TMS-C2H2NO2	218	C9H8N	130	3-4-5-6-7-8-9-10-11
Tryptophan(2TMS)	C17H28N2O2Si2	348	C9H8N	130	C8H20NO2Si2	218	1-2
Tryptophan(3TMS)	C20H36N2O2Si3	420	2TMS-C2H2NO2	218	C12H16NSi	202	3-4-5-6-7-8-9-10-11
Tryptophan(3TMS)	C20H36N2O2Si3	420	C12H16NSi	202	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	C10H15OSi	179	C8H20NO2Si2	218	1-2
Tyrosine(3TMS)	C18H35NO3Si3	397	TMS-CO2	117	C14H26NOSi2	280	2-3-4-5-6-7-8-9
Valine(2TMS)	C11H27NO2Si2	261	TMS-CO2	117	C7H18NSi	144	2-3-4-5
Valine(2TMS)	C11H27NO2Si2	261	C3H7	43	C9H20NO2Si2	218	1-2
α-Alanine(2TMS)	C9H23NO2Si2	233	TMS-CO2	117	C5H14NSi	116	2-3
β-Alanine(3TMS)	C12H31NO2Si3	305	CH3	15	C11H28NO2Si3	290	1-2-3

Figure legends

Figure 1. Workflow used in this study. A GC-MS method was built with abundant molecular ions
deriving from TMS-derivatives of organic and amino acids that contained a large number of
informative carbons. Then this method was evaluated by using: i) chemical standards and tailor-made
13C-standards that had a predictable CID y using *E. coli* as a metabolic factory (^{13}C-PT); ii) unlabeled
and ^{13}C-labeled plant samples.

Figure 2. **Evaluation of Carbon Isotopologue Distribution (CID) measurements for organic and**
amino acids by GC-MS using ^{13}C-PT reference samples. **A**, Evaluation of amino acids using ^{13}C-
PT protein hydrolysates. **B**, Evaluation of organic acids using ^{13}C-PT full metabolites extracts. Black
bars correspond to expected CIDs and blue bars to the measured CIDs. The results are presented as the
mean $\pm$ SD of four independent biological replicates. Statistical differences between predicted and
measured CID for each isotopologue of each fragment are denoted with asterisks (*) and were
established by considering the 95% confidence intervals of the measured CID. GC-MS fragments are
denoted according to the considered metabolite, its MEOX/TMS-derivatives analyzed and the
metabolite carbon backbone (for example, Glutamate_3TMS_C2C5 means that the fragment comes
from Glutamate(3TMS) and contains the carbon C2-C3-C4-C5 (**Table 1**)).

Figure 3. Evaluation of the measurement accuracy for CID and mean ^{13}C enrichment at ^{13}C-PT
and ^{13}C natural abundance in plants. A, CID accuracy and **B**, Mean ^{13}C enrichment for the selected
TMS-derivatives of organic and amino acids using ^{13}C-PT samples. **C**, Residual mean ^{13}C enrichment
measured in chemical standards, Arabidopsis seedlings and *B. napus* leaves after correction for isotope
natural abundance. **D**, Isotopologue-specific biases identified at ^{13}C natural abundance for Proline,
Valine and Threonine fragments. The results are presented as the mean $\pm$ SD of four independent
biological replicates or independent preparations for commercial chemical standards.

Figure 4. **Dynamics of U-^{13}C-pyruvate incorporation in *Brassica napus* leaf discs exposed to light**
or dark conditions. **A**, Mean fractional ^{13}C enrichment and **B**, Metabolite pool sizes. Mean fractional
^{13}C enrichment were monitored with GC-MS, while absolute quantification of organic and amino acids
was performed respectively with GC-FID and HPLC-UV. The results are presented as the mean $\pm$ SD
of four independent biological replicates. Statistical differences between dark and light conditions are
denoted with asterisks according to a Student test (*, p-value <0.05; **, p-value <0.01; ***, p-value
<0.001). The complete dataset is available in **Table S2, S3**.

Figure 5. **Fractional ^{13}C enrichment at the isotopologue level for TCA-cycle derived metabolites.**
A, Light conditions and **B**, Dark conditions. The M0 isotopologue was not represented to facilitate the
visualization of low-enriched isotopologues. The results are presented as the mean $\pm$ SD of four
independent biological replicates. The complete dataset is available in **Table S2**.

Figure 6. **Fractional contribution of PEPc and TCA cycle to Malate and Citrate production in**
light and dark conditions. A, Network reconstruction at the isotopomer level based on the
isotopologues detected. **B**, Fractional contribution of PEPc and TCA cycle to Malate and Citrate
production based on pathway-specific mass isotopologues/isotopomers. In the proposed network, the
second "turn" reflected the commitment of a molecule that has already experienced all reactions of the
TCA cycle to another round of reactions (Orange and blue labels to compare acetyl-CoA-derived
labeling from 1st and 2nd "turn"). Note that M0 isotopologues of all metabolites will still continue to
produce M0 istopologues at an important rate after a "turn" of the TCA cycle, given the ^{13}C-enrichment
for Acetyl-CoA active pool. For these calculations, only the timepoint T4 and T6 hours were used and

pooled (metabolic steady-state and isotopic steady-state for isotopologue ratios). The results are
presented as the mean ± SD of four independent biological replicates. Statistical comparison of Acetyl-
CoA enrichment for light and dark conditions was performed with a Student-test, while fractional
contributions were compared with an ANOVA followed by a post-hoc Tukey-HSD test (different
letters indicate groups that were separated with a p-value <0.05).

Figure 7. **Fractional contribution of PDH and TCA-cycle derived decarboxylations to**
photosynthesis. A, Fractional ^{13}C enrichment at the isotopologue level for Glycine and Serine. **B**,
Working hypotheses to explain the photorespiratory-dependent labeling pattern of Glycine and Serine.
C, Estimation of the fractional contribution of PDH (pyruvate dehydrogenase) and TCA-cycle CO_2 to
photosynthesis with Serine (expressed in %). For these calculations, only the timepoint T4 and T6
hours were used and pooled (metabolic and isotopic steady-state requirements, assumption with
$^{13}C/^{12}C$ of CO_2 released). The results are presented as the mean ± SD of four independent biological
replicates. Statistical comparison of the two Serine fragments was performed with a Student-test.

Figure 8. **Proposed metabolic fluxes for TCA cycle and PEPc in light and dark conditions in**
***Brassica napus* leaves based on this study.** The contribution of stored citrate is unlikely to occur
based on our results, while the forward cyclic mode of the TCA cycle has a higher contribution to both
malate and citrate production than the PEPc. Questions remains about the exact proportion of
Glutamate and Aspartate carbon backbones escaping the TCA cycle. Despite a reduction of PEPc
activity in dark conditions compared to light conditions, PEPc-derived carbons show a higher
commitment to Citrate production in dark conditions, thereby suggesting a higher rate of Malate import
into mitochondria. PDH and TCA-cycle derived CO_2 are reassimilated by photosynthesis and may
represent up to 30% of the TCA cycle metabolic flux. See Discussion for detailed explanations. TCA,
Tricarboxylic acids; PEPc, Phosphoenolpyruvate carboxylase; OAA, Oxaloacetate; GDH, Glutamate
dehydrogenase; GABA, Gamma-aminobutyric acid; ALAAT, Alanine:2-oxoglutarate
aminotransferase; ASAT, Aspartate:2-oxoglutarate aminotransferase.

Identification of TMS-derivative mass fragments maximizing carbon backbones

Production of reference ¹³C labelled samples with controlled CIDs (¹³C-PT)

Validation of TMS-derivative mass fragments for CID measurements

A CID accuracy

TMS-derivatives

Alphaalanine_2TMS_C2C3
Aspartate_2TMS_C2C4
Aspartate_3TMS_C2C4
Citrate_2TMS_C1C5
Citrate_4TMS_C1C6
Fumarate_2TMS_C1C4
Glutamate_3TMS_C2C5
Glycine_3TMS_C2
Isoleucine_2TMS_C2C6
Leucine_2TMS_C2C6
Lysine_4TMS_C2C6
Malate_3TMS_C1C4
Malate_3TMS_C2C4
Methionine_2TMS_C2C5

Phenylalanine_2TMS_C1C2
Phenylalanine_2TMS_C2C9
Proline_2TMS_C2C5
Pyruvate_MEOXTMS_C1C3
Serine_2TMS_C2C3
Serine_3TMS_C1C2
Serine_3TMS_C2C3
Succinate_2TMS_C1C4
Threonine_3TMS_C2C4
Tyrosine_3TMS_C1C2
Tyrosine_3TMS_C2C9
Valine_2TMS_C1C2
Valine_2TMS_C2C5

B Fractionnal Mean ^{13}C enrichment

C Bias at natural ^{13}C abundance

Standard
Arabidopsis seedlings
Brassica Napus leaves

D

Proline_2TMS_C2C5

Standard
Arabidopsis
Brassica napus

	M0	M1	M2	M3	**M4**
m/z	142	143	144	145	**146**

m/z = 146

Isoleucine_2TMS_xxx

Proline_2TMS_C2C5

Valine_2TMS_C2C5

	M0	M1	M2	**M3**	M4
m/z	144	145	146	**147**	148

Threonine_3TMS_C2C4

	M0	**M1**	M2	M3
m/z	218	**219**	220	221

A Mean fractional ^{13}C enrichment

B Metabolite pool sizes

A

Glycine_3TMS_C2

Glycine_3TMS_C2

Serine_3TMS_C1C2

Serine_3TMS_C1C2

Serine_3TMS_C2C3

Serine_3TMS_C2C3

Fractional ^{13}C enrichment

Time (hours)

isotopologue
1
2

B

C

Contribution of PDH and TCA cycle derived CO_2 to photosynthesis

Serine3C1C2   Serine3C2C3

t-test p-value = 0.07