Causal disconnectomics of motion perception: insights from TMS-induced BOLD responses

Short title: BOLD correlates of TMS perturbation in the visual system

Estelle RAFFIN[1,2,3], Roberto F. SALAMANCA-GIRON[1,2], Krystel R. HUXLIN[4], Olivier REYNAUD[5], Loan MATTERA[5], Roberto MARTUZZI[5], Friedhelm C. HUMMEL[1,2,6]

[1] Defitech Chair of Clinical Neuroengineering, Neuro-X Institute (INX), Ecole Polytechnique Fédérale de Lausanne (EPFL), Geneva, Switzerland.

[2] Defitech Chair for Clinical Neuroengineering, INX, EPFL Valais, Clinique Romande de Réadaptation, Sion, Switzerland.

[3] LPNC, CNRS UMR 5105, Université Grenoble Alpes, Saint-Martin-d'Heres, France.

[4] The Flaum Eye Institute and Center for Visual Science, University of Rochester, Rochester, NY, USA.

[5] Human Neuroscience Platform, Fondation Campus Biotech Genève (FCBG), Campus Biotech, Geneva, Switzerland.

[6] Department of Clinical Neurosciences, Geneva University Hospital (HUG), Geneva, Switzerland.

Corresponding Author: Estelle Raffin, LPNC, CNRS UMR 5105,Université Grenoble Alpes, Saint-Martin-d'Hères, France. Email: estelle.raffin@univ-grenoble-alpes.fr. https://orcid.org/0000-0003-3262-5251

Highlights:

- TMS-induced perturbation of the early visual areas (EVA) or the mediotemporal area (MT) area selectively impairs motion direction discrimination.
- The TMS perturbation is associated with a context-dependent local up-scaling of BOLD activity in both areas.
- The two visual areas display distinct topological networks' adaptation in response to TMS, reflecting different levels of network resilience to a focal lesion.
- TMS-fMRI coupling can be used to assess 'causal disconnectomics' and to precisely map how a local perturbation propagates to large-scale behavioural deficits.

Abstract:

Understanding how focal perturbations lead to large-scale network (re)organization is

essential for accurately predicting the behavioral consequences of brain lesions. In this

study, we applied a 'virtual lesion' approach by means of short bursts of 10 Hz

transcranial magnetic stimulation (TMS) over either early visual areas (EVA) or the

medio-temporal area (MT) in healthy participants, while acquiring concurrent functional

MRI. TMS delivered during the early stages of motion processing selectively impaired

direction discrimination at both sites, while global motion perception remained

unaffected. These behavioral effects were accompanied by a common local increase in

BOLD activity, but distinct patterns of network reorganization. Perturbation of EVA led

to more robust and efficient functional adaptation, suggesting greater resilience to focal

disruption. In contrast, behavioral impairments following MT stimulation were

associated with a less organized, more random network structure. Together, these

findings underscore the potential of TMS-fMRI coupling as a powerful approach for

mapping *causal disconnectomics*—the dynamic relationships between localized neural

disruption and widespread functional and behavioral outcomes providing a better

understanding of lesion-induced brain changes in neurological disorders such as stroke.

1. Introduction:

Brain networks dynamically adjust their spatiotemporal organization to changing contexts and ongoing demands (Telesford *et al.*, 2011; Papo *et al.*, 2014). Such adaptive networks can be characterized by their topological transformation, their state-dependency and/or their state transition. The adaptability of brain networks is crucial for maintaining an optimal balance between information segregation and integration, ensuring functional stability, efficient local processing, and communication through long-range connections. Understanding how flexible and distributed brain networks give rise to complex behavior is a central challenge in systems neuroscience. This adaptability may enhance the resilience of functional networks in response to progressive neurodegeneration or focal lesions (Bassett & Bullmore, 2009; Bullmore & Sporns, 2012; Fornito *et al.*, 2015). Today, a comprehensive understanding of the causal relationship between a focal lesion and brain network organization is still missing, leading to the inability to accurately predict resulting symptoms and recovery processes and develop targeted treatment strategies.

Directly testing the systematic association between a focal lesion and brain reorganization in humans is challenging, because of the difficulties in selectively altering activity *in vivo* and decipher the causal impact on brain networks. While advances in neuroimaging have provided detailed descriptions of functional and structural brain connectivity (Lim *et al.*, 2019; Ma *et al.*, 2022; Litwińczuk *et al.*, 2022; Blanco *et al.*, 2024), these correlational approaches offer limited insights into the causal mechanisms that underlie network dynamics and behavior. An emerging approach is to combine non-invasive brain stimulation (e.g., transcranial magnetic stimulation, TMS) with neuroimaging techniques (in particular functional magnetic resonance imaging (fMRI)) (Bergmann et al., 2021; Ozdemir et al., 2020; Vink et al., 2018, Raffin et al., in press). The

combination of TMS with fMRI provides a unique opportunity to temporarily perturb a

brain region and assess the changes in local brain activity and the resulting large-scale

network responses in vivo that may underlie changes in task performance (Beckers &

Hömberg, 1992; Ruff *et al.*, 2006; Bestmann S., 2008; Ricci *et al.*, 2012; Leitão *et al.*,

2015; Cocchi *et al.*, 2016). Several attempts have been made to combine fMRI with TMS

to capture the impact that TMS can have on patterns of functional connectivity in

specialized, large-scale, brain systems (Ruff *et al.*, 2008; Jung *et al.*, 2020). This holds

great promise for establishing *individual causal connectomics* (Glick *et al.*, 2024)— the

ability to map the local effects of targeted perturbations of brain nodes and how they

propagate across interconnected networks in individual participants.

Despite its conceptual appeal, key physiological mechanisms underlying TMS-fMRI

responses remain incompletely understood. First, the local neural response to TMS

reflects a complex interplay between intrinsic cortical excitability, synaptic connectivity,

ongoing oscillatory activity, and network state. Second, the associated local TMS-evoked

BOLD response and its physiological underpinnings remain to be fully elucidated (Rafiei

& Rahnev, 2022). However, earlier studies have shown that these local perturbations

can propagate through polysynaptic pathways, producing remote BOLD responses that

seem to reflect both direct anatomical projections and state-dependent functional

interactions (Ruff *et al.*, 2006; Bestmann S., 2008; Siebner *et al.*, 2022). The distinct

propagation patterns across brain regions, combined with individual variability, offer a

valuable   window   into   mapping   subject-specific   causal   connectivity—*causal*

connectomics.

Describing personalized causal connectomics is particularly relevant when considering

the application of TMS-fMRI to complex behavioral domains such as perceptual

decision-making and motion direction discrimination. Motion direction discrimination is

supported by a distributed network that integrates early visual areas (V1/V2) or EVA

(Moore *et al.*, 2001; Koivisto *et al.*, 2010), motion-sensitive temporal regions (MT)

(Newsome & Paré, 1988; Liu & Pack, 2017), the intraparietal sulcus involved in sensory

evidence accumulation (Chen *et al.*, 2017; Zhang *et al.*, 2022; Wongtrakun *et al.*, 2025),

and prefrontal circuits supporting decision-making and response selection (Kennerley &

Walton, 2011; Lin *et al.*, 2020). Impairments in this system can arise from dysfunctions

at multiple network levels, leading to behavioral deficits that may be difficult to

attribute to any single region. Disentangling whether such deficits arise from primary

sensory dysfunction, impaired integration, or altered top-down control requires tools

that can interrogate how information flows across the network. By applying TMS-fMRI

perturbations to key nodes within the motion discrimination network, here to the EVA

and MT, it becomes possible to experimentally probe how focal disruptions alter

distributed network dynamics and behavior. This approach may ultimately allow us to

characterize individual-specific network vulnerabilities that give rise to multidomain

symptoms, not only in healthy brain function but also in clinical populations where

sensory, cognitive, and motor symptoms often co-occur (Siegel *et al.*, 2016; Fleury *et al.*,

2022; Jimenez-Marin *et al.*, 2022). However, achieving such mechanistic understanding

requires further efforts to relate region-specific TMS-evoked local and network-level

responses to behavioral outcomes.

In this study, we aim to address these challenges by combining TMS-fMRI perturbation

mapping targeting EVA and MT with behavioral assessments of motion direction

discrimination, providing novel insights into the causal architecture of perceptual

decision-making networks. These two regions were chosen because they allowed us to

compare the whole brain consequences of a TMS perturbation affecting a primary

integrative brain region, acting as a gateway to other 'higher' visual areas, and in

contrast, a more functionally specialized brain region (Simoncelli & Heeger, 1998;

Grossberg *et al.*, 1999; Rust *et al.*, 2006; Mineault *et al.*, 2012). Simulation studies suggest that inhibitory rTMS targeting peripheral regions may exert more pronounced acute changes in network organization than stimulation of hub or integrative regions (Gollo *et al.*, 2017). Accordingly, we hypothesized that, as a primary integrative hub, EVA would display greater local and network-level homeostasis following TMS, thereby attenuating the propagation of perturbation effects and limiting both behavioral impairments and widespread network reconfiguration compared to MT (Das *et al.*, 2012; Gollo *et al.*, 2017; Tu *et al.*, 2021). Therefore, we anticipate that differences in TMS-induced behavioral impairments across behavioral state and across processing stages will be explained by distinct patterns of TMS signal propagation—shaped by the hierarchical network level of the targeted area—rather than by variations in local BOLD responses to TMS.

2. Methods

Participants

In total, thirty healthy subjects (seventeen males: mean age 28.8 years, range 19–39 years) were recruited in the experiment. All participants provided informed written consent prior the experiment and none of them met the MRI or TMS exclusion criteria (Rossi *et al.*, 2021). This study was approved by the local Swiss Ethics Committee (2017-01761) and performed in accordance with the Declaration of Helsinki.

Experimental Design and Task Procedure

Sixteen participants (nine males, mean age 26.5 years, range 19-32 years) performed

two TMS-fMRI sessions (except one drop-out for the second session). During the first

session, TMS was applied to the right EVA ($TMS_{(EVA)}$: mean($\pm$SD) MNI coordinate: 8(5);-

76(4);9(6)) using phosphenes perception when possible (in 9/16 participants) or the O2

position based on the 10-20 EEG system; during the second session, TMS was applied to

the functionally defined right MT ($TMS_{(MT)}$: mean MNI coordinates (SD): 46(4); -83(6);

11(5)) (Figure 1A). We chose this study design to maximize our chances to optimally

target the individual MT area through a dedicated MT localizer acquired during the first

session (see online supplementary materials for a description of the MT functional

localizer). For the two online TMS-fMRI sessions, TMS was applied during a *motion*

direction discrimination (MDD) Task and at *Rest*. These sessions had the same content

except the anatomical T1-MPRAGE sequence and the MT functional localizer sequence,

which were only performed during the first session. A short offline session prior the MT

session was needed to locate the individual MT cluster using a neuro-navigation system

(Localite GmbH, Bonn, Germany).

An offline control experiment was performed to control for TMS non-specific effects on

additional fifteen participants (seven males, mean age 28.9 years, range 22-39 years).

We repeated the *MDD task* with TMS using the same setup (neuronavigation and eye-

tracking systems) and the same task parameters (see below) but stimulating the right

motor cortex ($TMS_{(M1)}$, mean(SD) MNI coordinates: 53(7);-4(2);54(3), corresponding to

the individual first dorsal interossei muscle's motor hotspot of the participants), outside

the MRI scanner. Figure 1B summarizes the study timeline.

For the three sessions, the *MDD task* consisted in rating subjective perceptual

experience of visually presented motion stimuli and in performing a forced-choice

direction discrimination task on the same moving stimuli (Koivisto *et al.*, 2021). The

visual stimuli involved a group of static or moving dots coherently to the right or to the

left. During the familiarization phase outside the scanner, the duration of stimulus

presentation was individually defined (either 50, 67 or 84 ms, corresponding to 3, 4 or 5

frames) to ensure approximatively 70% accuracy in the *noTMS* trials. The stimulus

duration was then kept constant throughout the experiment (see Figure 1D for an

illustration of the task).

Just after the stimulus presentation, participants were asked to rate their perception

using a response box with their right hand using a 1 to 4 scale (see Figure 1's caption for

more details). In the main experimental conditions, TMS bursts were given to EVA when

visual signals first reach the target region 60 ms after the onset of the visual stimulus

($TMS_{(EVA)}$) (Lamme, 2001) and 30 ms after stimulus onset ($TMS_{(MT)}$) for MT to account for

the direct thalamic-extrastriate pathway (Beckers & Hömberg, 1992; Lamme &

Roelfsema, 2000; d'Alfonso *et al.*, 2002; Koivisto *et al.*, 2010). This ensured similar

perturbation of the feed-forward thalamic inputs to the two regions. Later TMS onsets

were also tested in half of the trials, 100ms for EVA and 130ms for MT selectively

interfering with late processing stages. The total duration of the TMS bursts was 200ms

to cover the full window of visual processing in the two regions. TMS intensity was set to

≈ 80% [range: 75 to 90%] maximal stimulator output (MSO), corresponding to dI/dt of

119–169 A/ µs. The resulting E-field magnitude at hotspot (99.9-th percentile)

corresponded to values between 73 and 112 V/m. This intensity was individually

adjusted prior the measurement to ensure phosphene sub-threshold stimulation and

progressively increased until participants report pain or discomfort. The intensity was

then chosen to get reliable BOLD signal while preserving participant's comfort. For the

M1 control experiment, the same EVA timings were used and TMS was given sub-

threshold to avoid motor evoked potentials elicitation.

Conditions randomly alternated between TMS versus noTMS trials, moving versus static

trials throughout the task, 50 trials each, giving a total of 300 trials. An inter-trial interval

(ITI) of approximatively 1.5 sec was used. The task was displayed in the scanner through

a 44cm x 27cm LCD monitor at a 2.5m distance via a mirror mounted on the head coil,

on a frame on top of the TMS-fMRI setup (Figure 1B). For the offline M1 control

experiment, the task was displayed on a mid-grey background LCD projector (1024 x

768, 144 Hz). To exclude bad performance due to TMS-induced blinking and to make

sure that participants were constantly looking at the fixation point, gaze and pupils'

movements were controlled in real time with an EyeLink 1000 Plus Eye Tracking System

(SR Research Ltd., Canada) in the scanner and in the behavioral room, sampling at a

frequency of 1000 Hz. Trials were aborted and redone if deviation exceeded 1° from the

fixation point. The duration of the *MDD Task-TMS* sequence was on average 16.5

minutes (SD = 4.30 minutes) depending on individual reaction times.

A *Rest-TMS* sequence was used to study the effect of TMS intensity. An event-related

design was used to map the effect of TMS bursts composed of three pulses at alpha

(10Hz) frequency. Three conditions were pseudorandomized and counterbalanced

across the run: high-intensity TMS (*HighTMS)*, low-intensity TMS *(LowTMS)* and no TMS

(noTMS), with 25 repetitions of each condition with an inter-trial interval (ITI) of 6

seconds (covering 3 repetition times). *HighTMS* intensity corresponded to the intensity

used during the MDD Task while *LowTMS* was set to ≈38 % [35 to 43%] MSO.

Participants were asked to look at a fixation cross throughout the acquisition. The

duration of the *Rest TMS* sequence was 9 minutes. Note that the results of the Rest-TMS

sequence are presented in the supplementary materials.

Figure 1.A: TMS targeting. *TMS targets were determined by the functional localizer of EVA*
and MT for precise TMS positioning (see online materials for further details about the
motion processing localizer). The two images on the right part show the T2 image of a
representative subject with the oil capsules placed on the TMS-MRI coil casing to monitor
the coil position located on the TMS coil for EVA and MT. These markers formed a "T-like
shape" indicating the center of the coil as well as the coil orientation as seen on a T2
HASTE image acquired with the two TMS-compatible MRI coil arrays (56 axial slices, slice
thickness = 0.8 mm, TR = 1500 ms, TE = 135 ms, FOV = 216 mm, flip angle = 140°). Coil
*positions were readjusted if needed. **B: MRI setup**. The MRI-TMS coil was positioned over*
the occipital area while the MR-receiving coil was positioned over the frontal lobe. Both
were maintained via a vacuum pillow to ensure the absence of movement or vibration and
*to ensure a comfortable position for the participant. **C: Study timeline**: Description of the*
*two experimental sessions; **D: One single trial of the motion discrimination task**. The*
visual stimuli involved a group of white dots in a 5˚ diameter circular aperture centered in
the middle of the screen. The dots were either static or moving coherently to the right or to
the left at a density of 2.6 dots per degree. Dots were displayed for 50, 67 or 84 ms
(corresponding to 3, 4 or 5 frames). This duration was individually adjusted prior the fMRI
measurement to ensure approximatively 70% accuracy in the noTMS trials. Just after the
stimulus presentation, participants were asked to subjectively rate their perception using
a response box on their right hand with a 1 to 4 scale, 1 being "I clearly saw the motion"
(Seeing), 2 being "I sensed the motion" (Sensing), 3 being "I saw something moving but
I'm unable to judge the direction" (No Direction) and 4 being "I didn't see any motion" (No
Motion) (Koivisto *et al.*, 2010; Grasso *et al.*, 2018). When the participants pressed the
three first options, they were then asked to judge whether the dots moved to the right or
to the left with a response box on their left hand. If participants pressed 4, the task moved
to the next trial. ***E:*** *Schematic of the BOLD sequence allowing artefact free combination of*
alpha TMS bursts.

Behavioral data analysis

Considering static stimuli, the mean motion perception accuracy, reflecting how well

participants could distinguish between moving and static stimuli, was extracted as the

percent of correct answer relative to the total number of trials belonging to the same

condition. Using moving stimuli, motion direction discrimination performances were

extracted as the percent of correct Left-Right judgments relative to the total number of

trials of that condition. For both scores, mixed ANOVA were built with factors Site (EVA,

MT and M1) and TMS (Early onset, Late onset and NoTMS). Violation of sphericity was

corrected using Greenhouse–Geisser corrections and Tukey's multiple comparisons tests

were applied for post hoc analyses when significant main effects or interactions were

found.

Magnetic Resonance Imaging & Transcranial Magnetic Stimulation

MRI images were acquired at the Magnetic Resonance Imaging facility of the Human

Neuroscience Platform of the Fondation Campus Biotech Genève (FCBG, Biotech

Campus, Geneva, Switzerland), using a Siemens Prisma 3 Tesla scanner (Siemens

Healthineers, Germany).

Standard MRI sequences acquisition

Anatomical images were acquired with a 64-channel head and neck coil using a 3D

MPRAGE sequence (TR/TE = 2300/2.96 ms, flip angle 9°, 192 sagittal slices, matrix = 256,

1x1x1 mm^3 resolution) covering the whole head. For retrospective distortion correction,

static field mapping was performed with a double-echo spoiled gradient echo sequence

(TR= 652 ms, TE1/TE2=4.92/7.38 ms, resolution 2.2x2.2x2.2 mm^3, whole head coverage),

generating two magnitude images and one image representing the phase difference

between the two echoes.

Combined TMS-fMRI sequences acquisition

For combined TMS-fMRI images, two dedicated coil arrays were used (Navarro de Lara

et al., 2017). This setup consisted of an ultra-slim 7-channel receive-only coil array,

which was placed between the subject's head and the TMS coil (MRI-B91, MagVenture,

Farum, Denmark) and connected to a MagPro XP stimulator (MagVenture, Farum,

Denmark). A second, receive-only MR coil was positioned over Cz in the EEG 10-20

system to allow a full coverage of the participant's brain.

The *Rest-TMS* and MDD *Task-TMS* sequences were acquired with a GE-EPI sequence

using the same parameters: 40 axial slices, slice thickness = 2.2 mm, in-plane resolution

= 2.2 mm, TR = 2000 ms, TE = 30 ms, FOV = 242 mm, flip angle = 67°, GRAPPA = 2,

Multiband Factor (MB) = 2. A gap was introduced between consecutive EPI volumes in

order to guaranty artefact free MR images after TMS stimulation (Navarro de Lara *et al.*,

2015). For both rest and task TMS sequence, a single repetition time (TR=2000 ms) was

therefore composed of 40 slices acquired during 1430 ms followed by a gap of 570 ms

before the next volume acquisition. The synchronization of TMS pulse was carried out

with an in-house script using Matlab (R2019).

Static field mapping was also performed with the TMS-MRI coils using the same double-

echo spoiled gradient echo sequence (TR = 652 ms, TE = 4.92 and 7.38 ms, slice

thickness: 2.2 mm, in-plane resolution = 2.2 mm, flip angle = 60°) that generates two

magnitude images and one image representing the phase difference between the two

echoes. Additional MRI sequences for coregistration of TMS-fMRI data are described in

the supplementary materials.

Preprocessing of fMRI images

Statistical Parametric Mapping software (SPM12, Wellcome Department of Imaging

Neuroscience, UK) was used for data pre-processing and analysis of the four datasets

(*Rest TMS* and *Task TMS* datasets, on both EVA and MT). Spatial distortions due to field

inhomogeneities were removed from all EPI images using the Fieldmap toolbox from

SPM12 (Andersson *et al.*, 2001). Subsequently, all time-series were motion-corrected to

the first volume using 6 degrees of freedom (3 rotations, 3 translations) and drifts in

signal time courses were corrected as well. To improve the normalization procedure,

two intermediate steps were added (see online materials for further details about the

additional sequences used for coregistration purposes). A first co-registration was

performed between the mean realigned and slice timing corrected image and the SSFP

sequence acquired with the same MR coil (and thus the same spatial coverage). The

resulting, co-registered image was once more co-registered to the SSFP sequence

acquired with the MR coil integrated into the scanner (i.e., the body coil, thus preserving

the contrast). The latter could then be easily co-registered to the high resolution T1

image acquired with the 64-channel head coil that covered the whole brain, and later

transformed into standard MNI space using a segmentation-based normalization

approach (Ashburner & Friston, 2005). Finally, the normalized images were spatially

smoothed using a Gaussian kernel (4 mm, Full-width half-maximal). The realignment

parameters estimated during spatial pre-processing for the *Rest TMS* and *Task TMS*

dataset were introduced in the design matrix as regressors of no interest in order to

prevent confounding activations related to minor head movements during scanning. The

experimental event-related designs were convolved with the canonical hemodynamic

response function, and the resulting models described in the following section were

estimated using a high-pass filter at 128 s to remove low-frequency artefacts.

fMRI Univariate Analysis

EVA and MT datasets were similarly analyzed using general linear models (GLM) to

calculate individual contrasts. For the *Task TMS* sequence, a design matrix with TMS

conditions (TMS_{early}, TMS_{late}, *noTMS*) and visual state (*Moving*, *Static*) was constructed

using all trials. To disentangle the effects of a TMS onsets and visual stimuli, we built

GLM with random effects at the second-level, by calculating a flexible factorial design, as

implemented in SPM12, with the within-subject factors *TMS condition* (TMS_{early}, TMS_{late})

and *Visual stimulus* (Moving and Static).

To explore the local effects of TMS across conditions, mean beta values were extracted

from the stimulated regions EVA and hMT+/V5 defined as a 8 mm radius sphere based

on the results of the group-level analysis of the motion processing localizer

(supplementary online materials) using the MarsBaR toolbox (Brett *et al.*, 2002)(right

EVA: 6, -83, 11 and right MT: 58, -66, -3). Repeated measure ANOVAs were built with

TMS (TMS_{early}, TMS_{late}, *noTMS*) and visual stimulus (*Moving*, *Static*) as within-subject

factors for each site of stimulation (EVA and MT).

For the *RestTMS* sequence, we defined a design matrix comprising three conditions

(*HighTMS*, *LowTMS* and *noTMS*). A second-level GLM with random effects was designed

using a flexible factorial design, with the within-subject factors *TMS intensity* (TMS_{High},

TMS_{Low} and *noTMS*). Beta values were extracted locally for each condition using the

same ROI as previsouly described.

The statistical significance threshold was set to a height threshold of $p < 0.001$

uncorrected, at the voxel level and to that of $p < 0.05$ at the cluster level after false-

discovery rate (FDR) correction.

fMRI Multivariate Analysis:

Independent Component Analysis (ICA) was used on the motion discrimination task

data to estimate independent spatiotemporal functional networks from the data and to

visualize networks specifically modulated by TMS. ICA uses fluctuations in BOLD signal to

split it into maximally independent spatial maps or components, each explaining a

unique part of variance from the 4D fMRI data. Then, all components have their specific

time course related to a coherent neural signal potentially associated with intrinsic brain

networks, artifacts, or both. The group ICA of fMRI Toolbox (GIFT version 3.0b; (Calhoun

et al., 2001)) was used. First, it concatenates the individual data followed by the

computation of subject-specific components and time course. Maximum Description

Length (MDL) and Akaike's criteria were applied to estimate the number of ICs in our

data. Using principal component analysis, individual data was reduced. Then, the

informax algorithm (Bell & Sejnowski, 1995) was applied for the group ICA and

estimated 16 components. In order to improve the IC's stability, the ICASSO was applied

and run 20 times (Himberg *et al.*, 2004). Over the 8 extracted network components, we

focused on the TMS specific networks defined by the overlap between the activated

clusters and the coordinates of the stimulation target. Based on these independent

components, we then examined how each condition (Moving or Static) covaries with the

TMS networks using the *temporal sorting* option in the GIFT toolbox. Temporal sorting

consists in regressing the fMRI specific time course for each individual against the design

matrix for the experimental conditions. The resulting betas weights represent the

degree to which network was recruited by the conditions. For a network, positive and

negative b weights indicate the level of network recruitment in each experimental

condition. Paired-sample t-tests were used to identify networks that were significantly

modulated by TMS at rest and during motion discrimination using $p < 0.05$.

Brain Networks generation and Graph Theoretical Analysis

In addition to ICA, we used the graph theoretical network analysis toolbox GRETNA to

image connectomics and construct the functional brain networks derived from the full

Task $TMS_{(EVA)}$ and $TMS_{(MT)}$ dataset comprising all the conditions of interest (e.g., noTMS,

TMS, motion or static stimuli) (Wang *et al.*, 2015). The whole brain was divided into 90

cortical and subcortical regions according to Automated Anatomical Labeling (Tzourio-

Mazoyer *et al.*, 2002), and the mean time series for each of the 90 regions was

extracted. Pearson's correlation coefficients for each pair of regions were calculated for

the mean time series of all of the 90 regions, and z-transformed using Fisher's Z

transformation. Subsequently, a positive binary undirected connection functional

network was constructed according to a range of selected threshold of the relation

matrix. As there is no defined standard for threshold selection in the construction of a

binary connection brain functional network, sparsity was used as a range of correlation

coefficient thresholds for correlation metrics. It was defined as the existing number of

edges in a graph divided by the maximum possible number of edges. In accord with

previous studies, we set both the sparsity and Pearson correlation threshold of the

functional network to range from 0.05 to 0.5 (in 0.05 steps), resulting in a more efficient

functional network than a random network with the number of artificial edges

minimized (Achard & Bullmore, 2007; Zhao *et al.*, 2017). Graph theoretical analysis was

applied to assess the topological and organizational properties focusing on 3 graph

theoretical metrics (Rubinov & Sporns, 2010). These metrics were: the clustering

coefficient (γ), which measures how much neighbors of a node are connected to each

other and is closely related to local efficiency; characteristic path length (λ), which is the

average number of edges needed to get from any node in the network to any other

node in the network and is inversely related to global efficiency; small worldness (δ),

which reflects whether the network balances global integration and local segregation for

efficient information processing (Latora & Marchiori, 2001). To compare graph

theoretical network properties between EVA and MT, integration of the topological

metric over all selected ranges of sparsity values was calculated using GRETNA. Paired t-

tests corrected for FDR (p < 0.05) were performed between the topological measures

of EVA and MT networks.

3. Results

3.1 Behavioural results

We used chronometric TMS-fMRI coupling in healthy participants to perturb and map

the effect of TMS over the right EVA (TMS$_{(EVA)}$) and over the right MT (TMS$_{(MT)}$) during a

motion direction discrimination task. An additional offline experiment targeted the right

primary motor cortex (TMS$_{(M1)}$) in another subgroup of healthy participants.

Figure 2A shows the mean and individual motion perception accuracy (+/- SEM),

reflecting how well participants could distinguish between moving and static stimuli

when a TMS burst was applied over EVA (top row), MT (middle row) and over the

control region M1 (bottom row). A mixed ANOVA revealed no main effect of TMS ($F_{(1,24)}$

= 2.9, p = 0.084, η^2 = 0.02) and no significant TMS by Site interaction ($F_{(4,48)}$ = 0.9, p =

0.452, η^2 = 0.031), suggesting that TMS has no consequence on motion perception in

these three regions. However, there was a Site effect ($F_{(2,24)}$ = 6.8, p = 0.004, η^2 = 0.16),

likely driven by the overall better performances in the M1 session. The same model

applied to the motion direction discrimination performances (Figure 2B) revealed a

significant TMS effect ($F_{(2,305)}$ = 22.012, p < 0.001, η^2 = 0.2) and Site effect ($F_{(2,30)}$ = 20.57,

p < 0.001, η^2 = 0.2). There was also a TMS by Site interaction ($F_{(4,60)}$ = 9.87, p < 0.001, η^2 =

0.124). When TMS was applied to EVA, performances were markedly affected by TMS$_{early}$

compared to noTMS ($t_{(15)}$ = 5.2, adjusted p < 0.001) and compared to TMS$_{late}$ ($t_{(15)}$ = 4.4,

adjusted p = 0.034). Although slightly lower, the difference between noTMS and TMS$_{late}$

was not significant (t$_{(15)}$ = 1.45, p = 0.35).

In contrast, when applied to MT, TMS$_{early}$ and TMS$_{late}$ negatively impacted performance

compared to noTMS (TMS$_{early}$: t$_{(15)}$ = 5.2, adjusted p < 0.001, TMS$_{late}$: t$_{(15)}$ = 5.2, p <

0.001). After correction for multiple comparisons, performance under TMS$_{early}$ and

TMS$_{late}$ were not different (t$_{(15)}$ = 2.2, p = 0.1). Performances were similar in the three

TMS conditions for M1 (all comparisons p>0.05) and baseline performances (noTMS

conditions) were not different across sites (all comparisons p>0.05).

We also examined the relationship between awareness and accuracy by taking into

account motion awareness ratings (see supplementary Figure S1). Without TMS, motion

direction discrimination increased (moved away from chance level) as subjective

awareness of motion increased. This indicates that without TMS, performance is tied to

conscious motion awareness. The rmANOVA with TMS and Rating as within subject

factors only revealed a significant Site by TMS condition interaction, confirming the

previous analysis (F$_{(4,12)}$ = 15.1, p < 0.011, η^2 = 0.09). There was no main effect of

Awareness nor interactions with any other factor (all p>0.05).

In summary, TMS bursts exerted negative effects on motion direction discrimination at

the early and late onsets in MT. TMS bursts were only efficient in disturbing motion

directions discrimination in the early onset for EVA. The two onsets did not affect

performances when targeting the control region M1.

453
Figure 2: *The upper row illustrates the individual data of motion perception under noTMS,*
TMS_early and TMS_late for TMS_(EVA) (A), TMS_(MT) (B) and TMS_(M1) (C); The bottom row illustrates the
individual data of motion direction under noTMS, TMS_early and TMS_late for TMS_(EVA) (A), TMS_(MT)
(B) and TMS_(M1) (C);

3.2 Whole-brain and local responses to TMS perturbation

Next, we analyzed local BOLD activity in response to TMS bursts over EVA and

MT applied during behavioral perturbation of motion direction discrimination. First,

Figure 3A (TMS_(EVA)) and Figure 3B (TMS_(MT)) report the results of the average effects of

TMS conditions. TMS_(EVA) induced significant clusters in the bilateral occipital cortex, the

right superior and middle temporal gyrus, in the bilateral anterior and middle cingulate

cortex, the left middle frontal gyrus, the bilateral caudate nucleus and in the right

posterior thalamus. When TMS was given over the right MT, the average effect of TMS

conditions returned significant clusters in the right occipital cortex, the right inferior and

frontal gyrus, the right intra-parietal sulcus and the right cerebellum (cruz VI). These

clusters overlapped with the TMS-induced activation at rest (see supplementary results

and Table S1).

We then examined the main effect of visual stimuli (difference between TMS

applied during a moving or static stimuli) returned significant clusters in the right middle

temporal gyrus and in the bilateral middle frontal gyrus the right superior frontal gyrus,

the posterior cingulate cortex, and the right occipital cortex when EVA was stimulated

(Figure 3C) and only in the right middle temporal gyrus, when MT was stimulated (Figure

3D). When stimulating the EVA, the main effect of TMS onsets revealed a significant

cluster in the right middle frontal gyrus and in the right superior frontal gyrus (Figure

3E). There was no significant cluster when stimulating MT (see Supplementary Table S2

and S3 for the MNI coordinates of the significant clusters of all effects).

The extracted beta values from the stimulated EVA and MT regions were

entered into an ANOVA with the factors TMS (noTMS, TMS_{early} and TMS_{late}) and Visual

stimuli (moving, static) and stimulation site (EVA, MT). This analysis revealed a

significant TMS effect ($F_{(2,22)}$ = 3.8, p = 0.038), mostly driven by an increase in beta

weights for the two TMS conditions compared to no TMS, while there was no difference

between the early and late TMS onsets. Finally, there was a significant TMS by Visual

stimuli interaction ($F_{(2,22)}$ = 6.1, p = 0.008) showing that the increase in activity only

occurred in the combination early TMS associated with moving stimuli (noTMS, Moving

vs early TMS, Moving: $t_{(12)}$ = -4.2, p = 0.003 and early TMS, Static vs early TMS, Moving:

$t_{(12)}$ = -3.6, p = 0.015, supplementary Table S4 for all posthoc comparisons). There was no

stimulation site effect, demonstrating that the local response to TMS were similar in

both regions.

Figure 3A: *Activation maps corresponding to the average TMS conditions for TMS(EVA)*; **B:** *Activation maps corresponding to the average TMS conditions for TMS(MT)*; **C:** *Main effect of visual stimuli for TMS(EVA)*; **D:** *Main effect if visual stimuli for TMS(MT)*; **E:** *Main effect of TMS onsets for TMS(EVA)*; **F:** *Main effect of TMS onsets for TMS(MT)*; **G:** *Local beta weights extracted for the EVA ROI:* **H:** *Local beta weights extracted for the MT ROI. P(FWE)) < 0.001 at the cluster level.*

3.3 Network changes in response to TMS perturbation

We further investigated whether and how long-range brain activity was modulated by

TMS during motion direction discrimination. We first applied an independent

component analysis to extract hidden spatiotemporal components contained within the

TMS(EVA) and TMS(MT) task data, returning a measure of functional connectivity. We then

applied temporal regression analysis on the components' time courses to compare

network activity associated with the different experimental conditions.

The ICA applied to the $TMS_{(EVA)}$ data during motion discrimination revealed 8

independent networks, among which 7 were considered as functional brain networks.

Five networks (IC1, IC3, IC4, IC7 and IC8) were labelled as TMS-related networks because

they overlapped with the stimulation site (Figure 4A and Supplementary Table S5 for the

MNI coordinates of the IC). IC1, IC7 and IC8 covered the bilateral EVA while IC3 only

encompassed the right EVA and IC4, the left EVA. Only IC1 and IC7 were significantly

modulated by TMS as revealed by one-way ANOVAs modeling the t scores obtained

from the temporal regression analysis on the components' time courses for moving and

static stimuli. IC1 was significantly down regulated by both TMS onsets when associated

with moving stimuli only (one-way ANOVA: $F_{(2,15)}$ = 3.6, p = 0.04, post hoc TMS_{early} vs

noTMS: $t_{(15)}$ = 1.9, p = 0.07, TMS_{late} vs noTMS: $t_{(15)}$ = 2.2, p = 0.04). IC7 showed a different

pattern of modulation reflecting a significant over-expression of the bilateral EVA only

during TMS_{early}, for moving ($F_{(2,15)}$ = 3.5, p = 0.045, post hoc TMS_{early} vs noTMS: $t_{(15)}$ = 2.4,

p = 0.03) and static stimuli ($F_{(2,15)}$ = 4.4, p = 0.03, post hoc TMS_{early} vs noTMS: $t_{(15)}$ = 2.3, p

= 0.035). All the other ANOVAs were not significant.

When TMS was given over MT, 7 components were considered as functional brain

networks. Among them, three networks (IC1, IC2 and IC8) were labelled as TMS-related

networks (see Supplementary Table S6 for the MNI coordinates of the IC). Interestingly,

these networks were not only restricted to the right MT. IC2 and 8 also included the IPS

and superior parietal cortex and the medial frontal gyrus. IC 1 activity was significantly

enhanced by both TMS onsets when stimuli were moving ($F_{(2,15)}$ = 3.6, p < 0.001 post hoc

TMS_{early} vs noTMS: $t_{(15)}$ = 2.7, p = 0.02, TMS_{late} vs noTMS: $t_{(15)}$ = 3.8, p = 0.002) while IC8

was significantly depressed during moving stimuli ($F_{(2,15)}$ = 7.4, p = 0.003, post hoc

TMS$_{early}$ vs noTMS: t$_{(15)}$ = 3.2, p = 0.007, TMS$_{late}$ vs noTMS: t$_{(15)}$ = 3.1, p = 0.008, Figure 4B).

All the other ANOVAs were not significant.

Figure 4A: *TMS$_{(EVA)}$-specific component networks that are modulated by TMS during motion*
discrimination and temporal regression analysis for the noTMS and TMS$_{(EVA)}$ conditions associated
with moving or static stimuli; ***B***: *TMS$_{(MT)}$-specific component networks that are modulated by TMS*
during motion discrimination and temporal regression analysis for the noTMS and TMS$_{(EVA)}$
conditions associated with moving or static stimuli; Note that the components are sorted based
on the percent of explained variance.

Finally, we used a graph theoretical analysis to assess the topological and organizational

properties of these networks, while being agnostic to the TMS condition and visual

stimulus. We first investigated small-world topological properties especially because

networks with high small-world properties are believed to play a role in protecting the

brain from local perturbation or damage. Both EVA and MT networks exhibited small-

world properties (λ≈1 and σ>1) among the selected sparsity values (see Methods

section). Interestingly, there was a significant difference in the small-world network

parameters (Sigma) between EVA networks and MT networks with higher Sigma values

for EVA networks at the sparsity values 0.3, 0.4, 0.45 and marginally at 0.35 (Figure 5A,

left panel and supplementary Table S7 for all paired t-tests), also considering the mean

across all sparsity values ($t_{(12)}$ = 2.4, p = 0.034) (Figure 5B, left panel). There was no

difference in normalized clustering coefficient (Gamma) ($t_{(14)}$ = 0.42, p = 0.68) (Figure 5A

& B, middle panels and supplementary Table S8 for all paired t--tests). Finally,

characteristic path length (Lambda) was significantly higher at sparsity thresholds of 0.2,

0.25 and marginally at 0.15 for MT networks (Figure 5A, right panel and supplementary

Table S9 for all paired t-tests). The difference between EVA and MT+/ networks was also

significant when averaging across scarcities ($t_{(14)}$ = 2.4, p = 0.03, Cohen's d: 0.75) (Figure

5B, right panel).

Figure 5A: *Small-worldness properties (left panel), normalized clustering coefficients (middle*
panel) and normalized characteristic path length (right panel) for TMS$_{(EVA)}$ and TMS$_{(MT)}$ along the
sparsity thresholds of 0.05-0.50; ***B:*** *Associated mean values of small-worldness properties (left*
panel), normalized clustering coefficients (middle panel) and normalized characteristic path
*length (right panel) for TMS$_{(EVA)}$ and TMS$_{(MT)}$ * p < 0.05.*

3.4 Predicting TMS effects on behaviour with BOLD activity

To explain the effect of TMS on motion direction discrimination, we performed a series of multiple linear regressions using for each of the four conditions (TMS$_{(EVA)}$early, TMS$_{(EVA)}$late, TMS$_{(MT)}$early, TMS$_{(MT)}$late), the associated local beta scores, the t scores derived from the temporal regression of the significant TMS-ICs and the three graph theory metrics as fixed effects. The significant model explained the drop in performance associated with TMS$_{(MT)}$early with Gamma values from the graph theory analysis as confirmed by the ANOVA on the model's parameters (F$_{(1,6)}$ = 6.6, p = 0.04), suggesting that TMS exerted a negative impact especially in participants showing lower cluster coefficient, implying a more random or decentralized network structure. All other models were not significant (see supplementary Table S10 for all the multiple linear regressions results).

4. Discussion

The goal of this study was to examine whether TMS perturbation elicits inter-individual differences in BOLD responses, potentially reflecting the magnitude of the induced behavioral impairment. We used short bursts of 10Hz TMS to disrupt either a low-level visual area (EVA) or a highly specialized, motion sensitive, visual area (MT) during concurrent fMRI. We first confirmed the causal involvement of both regions in motion direction discrimination, with more pronounced behavioral impairments following stimulation of MT. Our main findings were: (1) a local, context-dependent increase in BOLD activity common to both targeted areas; (2) distinct patterns of network reconfiguration; and (3) a relationship between TMS-induced performance decline after MT stimulation and a shift toward a more random network structure. These results suggest that TMS perturbation triggers area-specific topological adaptations, which may reflect differing levels of network resilience to focal disruption.

4.1. TMS-induced behavioural deterioration

In line with the concept of "virtual lesion" described in the literature (Pascual-Leone et al., 1999; Beynel et al., 2019), TMS bursts applied to either EVA and MT, resulted in a selective and transient deterioration of performance of a left/right motion direction discrimination task (except for the late TMS$_{(EVA)}$ condition). This was interpreted to reflect successful causal and selective disruption of brain functioning. Interestingly, participants remained able to dissociate moving from static stimuli. This dissociation is likely supported by the existence of independent substrates mediating unspecific motion processing versus global motion direction integration in EVA and MT (Simoncelli & Heeger, 1998). There is evidence showing that a sub-population of

neurons in MT focuses on the processing of local motion signals similar to that in EVA,

and some other neuronal populations rather process global motion (Born & Tootell,

1992). Our TMS perturbation might have specifically impacted on the neurons in charge

of motion directions decoding. The absence of significant impairment induced by the

late onsets of TMS$_{(EVA)}$ might be explained by different time delay of feedback loops in

the motion discrimination system (Foss & Milton, 2000; Joukes et al., 2014). Considering

now the relationship between motion awareness and motion direction discrimination

(please see supplementary Figure SX), we found that the direction of the moving trials

that were judged as "moving", were not correctly guessed when both areas were

stimulated. In contrast, "unaware" trials were unchanged. We can speculate that this

residual, unconscious ability of detecting changes in the dots' positions might be at least

partially supported by the activation of the fast, direct geniculo-extra-striate pathway

(Bridge et al., 2008; Ajina et al., 2015; Abed Rabbo et al., 2015).

4.2. Local TMS-induced activity

Our results demonstrate that short bursts of TMS can elicit local BOLD activation

beneath the coil in both EVA and MT, even at relatively low intensities—below the

phosphene threshold. While the presence of local BOLD activity elicitation in response

to TMS is still unresolved in the field (Rafiei & Rahnev, 2022), we specifically found a

local over-activation in the presence of moving visual stimuli for both regions. This

coupling between local BOLD increase and decreased performance suggests that the

additional activity may reflect the accumulation of task-irrelevant neural noise (Ruff et

al., 2009; Bancroft et al., 2014). Such a mechanism aligns with established models of

TMS-induced disruption, where stimulation is thought to activate neuronal populations

that are not optimally tuned to the task-relevant feature—in this case, motion direction

(Silvanto & Pascual-Leone, 2008). As a result, the effective signal-to-noise ratio within

the motion-sensitive network would be reduced, impairing accurate motion

discrimination. It also highlights the state-dependent nature of TMS effects: stimulation

during active motion direction processing leads to qualitatively different outcomes than

during static stimuli presentation, suggesting that the excitability and engagement of the

targeted network at the time of stimulation critically shape the neuronal consequences.

These findings reinforce the idea that TMS does not simply "excite" or "inhibit" local

tissue, but interacts dynamically with ongoing neural activity (Romero *et al.*, 2019;

Siebner *et al.*, 2022). This, together with other recent findings (Perera *et al.*, 2024; Luo *et*

al., 2025), might open the door to more tailored or context-sensitive brain stimulation

protocols.

3. TMS induced whole-brain activity

An important question is whether this "local noise injection" propagates in the

brain to destabilize larger functional networks. TMS-induced changes in BOLD activity

were observed not only locally in the stimulated regions, but also in remote areas

through functional coupling. For both EVA and MT stimulation, the spread of activity to

distal regions—such as the intraparietal sulcus (IPS), frontal eye fields (FEF), and medial

prefrontal cortex—replicates previous findings from TMS-fMRI studies (Ruff *et al.*, 2006;

Caparelli *et al.*, 2010; Leitão *et al.*, 2015). Notably, many of these remote regions are

task-relevant and are known to be engaged during motion perception and direction

discrimination (Offen *et al.*, 2010; Sani *et al.*, 2021), suggesting that the ongoing task

context may also shape the propagation of TMS-evoked activity through the actively

engaged network.

When TMS was applied during motion processing, additional BOLD activation

was observed in regions such as the medio-temporal cortex (including MT) and medial

frontal areas. Importantly, this increased activation coincided with a measurable decline

in behavioural performance on the motion direction discrimination task, as well as with

participants' subjective reports of increased task difficulty. These findings suggest that

TMS may disrupt efficient network integration during perceptual processing, potentially

by altering the balance of activity within task-relevant networks. The differential effect

induced by the early and late TMS onsets was only visible when stimulating EVA and

manifested as an increase in activity in the right FEF and in the medial prefrontal cortex

with the late onsets. These regions are involved in perceptual decision making and this

time could reflect a compensatory mechanism that maintain stable performance (Imani

et al., 2021).

4. TMS-induced network activity

To provide both a qualitative and quantitative framework for comparing

network activity across different TMS-induced perturbation, we combined group

Independent Component Analysis (ICA) with graph-theoretical metrics. ICA identified

multiple functional networks, some directly linked to TMS and others more domain-

general. Across both targeted regions (EVA and MT), the decomposition revealed

components with divergent responses to TMS. In the case of EVA, for example, IC7—

spatially overlapping with the site of stimulation—showed increased activation

specifically following early TMS onset. In contrast, IC1, located more rostrally, exhibited

a pattern of downregulation under the same stimulation condition. The opposing

responses may reflect a balance between excitatory and inhibitory processes, possibly

mediated by intra-areal or feedback projections (Shao & Burkhalter, 1996; Schwabe *et*

al., 2006). Furthermore, the specificity of IC7's response to early TMS onset raises

questions about the temporal dynamics of network susceptibility, with different

networks or nodes having critical windows of "vulnerability" or influence (Huang & Yu,

2017; Kottaram *et al.*, 2018). This dissociation suggests that ICA successfully captured

the complex, spatially and temporally differentiated dynamics of TMS-induced

responses. It highlights ICA's sensitivity not only to spatial localization but also to the

functional state of the network—whether it is task-relevant, engaged, or modulated by

TMS in a context-dependent manner. It further emphasizes that TMS effects are not

uniform within a target region but depend on the local functional architecture and

current cognitive or perceptual state.

Regarding network activity induced by $TMS_{(MT)}$, we found that TMS engaged

more spatially distributed networks, involving additional associative regions such as the

FEF, the posterior parietal cortex (PPC), and medial prefrontal cortex (mPFC). In

contrast, all but one of the independent components associated with $TMS_{(EVA)}$ were

largely confined to the stimulated visual area, indicating more localized network

engagement.

To quantitatively assess these differences, we applied graph-theoretical analyses

and observed distinct whole-brain network configurations depending on the stimulation

site. Specifically, networks following TMS over EVA exhibited higher small-worldness—a

topological property characterized by strong local clustering and short path lengths—

compared to those following TMS over MT. This network architecture is thought to

support efficient parallel information processing by maximizing information transfer at

minimal wiring cost (Achard & Bullmore, 2007). The clustered, locally efficient

organization of the $TMS_{(EVA)}$ network may promote functional resilience and flexible

redistribution of activity. This could help explain the relative preservation of behavioral

performance during late TMS onset, suggesting that the network maintains coherence

despite transient perturbations (Wu *et al.*, 2020). In contrast, the more dispersed

TMS$_{(MT)}$ network configuration likely reflects broader propagation of task-specific

interference across higher-order visual regions, potentially leading to a less efficient,

more vulnerable network state. Congruently, the variable that best accounted for the

TMS-induced performance impairments following TMS$_{(MT)}$ was the clustering coefficient.

This measure reflects the tendency of nodes within a network to form tightly

interconnected clusters. High clustering values indicate a well-organized, locally efficient

network structure, whereas lower values reflect a more random or fragmented

configuration, consistent with patterns observed in our ICA analyses. Our findings

showed that participants who exhibited greater performance declines had lower

clustering coefficients, suggesting that a more dispersed network organization is

detrimental to motion direction discrimination. It would be valuable in future work to

investigate whether these network disruptions extend to other visual subfunctions—or

even domain-general cognitive functions—in order to determine the broader functional

impact of focal perturbations on brain-wide network dynamics.

This result fits with our initial hypothesis which was that EVA, given its role as an

early-stage integrative cortical area, would maintain more stable synaptic and network

dynamics in response to TMS, resulting in greater resistance to externally induced

disruptions in both perceptual performance and functional connectivity compared to

MT. It is in line with a modelling study, showing that brain hubs, where activity is

integrated and further distributed (like EVA in our case) operate in a slower regime and

appear to be functionally less affected by focal perturbations (Gollo *et al.*, 2017). In

contrast, perturbations of peripheral or more associative areas (e.g., MT), might have

greater impact on acute network activity. Lesion studies also provided evidence that

damage to the homologue of MT in behaving cats impairs more relearning of motion

discrimination and learning transfer than lesions to EVA (Das *et al.*, 2012). To sum up,

the network analyses revealed differences between the two disrupted areas, not

observable with classical activation analyses. It highlights the relevance of graph-

theoretical metrics in capturing functionally meaningful changes in network topology. In

particular, the clustering coefficient appears to be a sensitive marker of TMS-induced

network reconfiguration and its behavioural consequences.

5. Limitations of the current work and open questions:

The aim of this study was to develop a causal connectomic framework for

understanding motion direction discrimination by selectively perturbing two key

network nodes using TMS. However, it is important to recognize certain methodological

limitations inherent to this approach. Unlike brain lesions, TMS-induced perturbations

are both temporally transient and spatially focal. More critically, the physiological

consequences of a lesion might differ fundamentally from those of TMS. Brain lesions

not only disrupt local neuronal activity but also provoke widespread, often nonspecific

responses—particularly in the hyperacute phase—such as excitotoxicity, inflammation,

and diaschisis (Lai *et al.*, 2014; Farooqui *et al.*, 2022). These processes can profoundly

alter interregional communication through mechanisms unrelated to normal synaptic

signaling.

In contrast, the type of TMS used in this study induces neuronal firing and inter-

areal communication, though likely mediated by different—and possibly more

superficial or excitatory—neuronal populations than those affected by a brain lesion.

This distinction is essential when interpreting the resulting BOLD signal changes. While

increases in BOLD activity are often taken to reflect compensatory engagement, they

could alternatively indicate maladaptive or inefficient network recruitment. Future work

using population tuning models or decoding approaches could help clarify the functional

relevance of these activations and determine whether they truly reflect adaptive

reorganization or instead a signature of disrupted processing.

Another intriguing aspect concerns the resting-state TMS-fMRI data (presented

in the supplementary results). These results helped confirm the spatial specificity of the

TMS effects, validating that stimulation targeted the intended network nodes even at

rest. However, we observed that the direction of BOLD signal changes at rest contrasted

with those seen during task performance. This apparent discrepancy has been reported

in prior studies and may reflect state-dependent effects of TMS—that is, the neural

response to TMS can differ markedly depending on whether the brain is at rest or

engaged in a task. Another contributing factor could be the experimental design itself:

we employed a block design during the resting-state acquisition and an event-related

design during the task-based sessions. These methodological differences could influence

the temporal integration of BOLD responses and the observed directionality of signal

change (Chee *et al.*, 2003; Tie *et al.*, 2009). Together, these findings underscore the

complexity of interpreting TMS-induced effects in fMRI data and highlight the need to

consider both brain state and experimental context when drawing causal inferences.

6. Conclusion

Despite growing interest and significant advances in computational modelling,

the causal mechanisms underlying neural network dynamics following a lesion remain

incompletely understood. In this study, we employed concurrent TMS-fMRI to probe the

neural mechanisms supporting functional reorganization across whole-brain networks,

using a targeted perturbation approach.

Our findings offer neuroimaging evidence for the context-dependent nature of
TMS effects, suggesting that such perturbations may preferentially engage specific
neuronal populations depending on task demands. Moreover, we show that inducing
focal, virtual lesions at different levels of the visual hierarchy leads to distinct patterns of
dysregulation across visual networks.

Complementing classical lesion studies (Das *et al.*, 2012), our results highlight
the utility of TMS-fMRI coupling as a powerful tool for investigating 'causal
disconnectomics'—the causal relationships between localized disruptions and large-
scale neural and behavioural consequences (Egger *et al.*, 2021). This approach holds
promise for precisely mapping how local perturbations propagate through brain
networks, especially when assessed during active cognitive processing. Ultimately, such
techniques may advance our understanding of brain resilience, compensation, and
vulnerability in the face of focal damage.

Acknowledgements:

We thank Holly Bridge for her insightful comments on the manuscript. We would like to

thank the MRI and neuromodulation facilities of the Human Neuroscience Platform of

the Fondation Campus Biotech Geneva, for technical advice. This study was supported

by the Bertarelli Foundation (Catalyst BC77O7 to FCH & ER), by the Swiss National

Science Foundation (PRIMA PR00P3_179867 to ER), and by the Defitech Foundation (to

FCH).

Conflict of interest statement: The authors declare no financial or non-

financial competing interests.

References

Abed Rabbo F, Koch G, Lefèvre C & Seizeur R (2015). Direct geniculo-extrastriate pathways: a review of the literature. *Surg Radiol Anat SRA* **37,** 891–899.

Achard S & Bullmore E (2007). Efficiency and cost of economical brain functional networks. *PLoS Comput Biol* **3,** e17.

Ajina S, Pestilli F, Rokem A, Kennard C & Bridge H (2015). Human blindsight is mediated by an intact geniculo-extrastriate pathway. *eLife* **4,** e08935.

d'Alfonso AA, van Honk J, Schutter DJ, Caffé AR, Postma A & de Haan EH (2002). Spatial and temporal characteristics of visual motion perception involving V5 visual cortex. *Neurol Res* **24,** 266–270.

Andersson JLR, Hutton C, Ashburner J, Turner R & Friston K (2001). Modeling Geometric Deformations in EPI Time Series. *NeuroImage* **13,** 903–919.

Ashburner J & Friston KJ (2005). Unified segmentation. *NeuroImage* **26,** 839–851.

Bancroft TD, Hogeveen J, Hockley WE & Servos P (2014). TMS-induced neural noise in sensory cortex interferes with short-term memory storage in prefrontal cortex. *Front Comput Neurosci*; DOI: 10.3389/fncom.2014.00023.

Bassett DS & Bullmore ET (2009). Human brain networks in health and disease. *Curr Opin Neurol* **22,** 340–347.

Beckers G & Hömberg V (1992). Cerebral visual motion blindness: transitory akinetopsia induced by transcranial magnetic stimulation of human area V5. *Proc Biol Sci* **249,** 173–178.

Bell AJ & Sejnowski TJ (1995). An information-maximization approach to blind separation and blind deconvolution. *Neural Comput* **7,** 1129–1159.

Bergmann TO, Varatheeswaran R, Hanlon CA, Madsen KH, Thielscher A & Siebner HR (2021). Concurrent TMS-fMRI for causal network perturbation and proof of target engagement. *NeuroImage* **237,** 118093.

Bestmann S. (2008). Mapping Causal Interregional Influences With Concurrent TMS-fMRI - PubMed. *Exp Brain Res*. Available at: https://pubmed.ncbi.nlm.nih.gov/18936922/ [Accessed March 23, 2020].

Beynel L, Appelbaum LG, Luber B, Crowell CA, Hilbig SA, Lim W, Nguyen D, Chrapliwy NA, Davis SW, Cabeza R, Lisanby SH & Deng Z-D (2019). Effects of online repetitive transcranial magnetic stimulation (rTMS) on cognitive processing: A meta-analysis and recommendations for future studies. *Neurosci Biobehav Rev* **107,** 47–58.

Blanco R, Preti MG, Koba C, Ville DVD & Crimi A (2024). Comparing structure–function relationships in brain networks using EEG and fNIRS. *Sci Rep* **14,** 28976.

Born RT & Tootell RBH (1992). Segregation of global and local motion processing in
primate middle temporal visual area. *Nature* **357,** 497–499.

Brett M, Anton JL, Valabrgue R & Poline J-B (2002). Region of interest analysis using an
SPM toolbox. Presented at the 8th International Conference on Functional
Mapping of the Human Brain, June 2-6, 2002, Sendai, Japan. *Neuroimage* **13,**
210–217.

Bridge H, Thomas O, Jbabdi S & Cowey A (2008). Changes in connectivity after visual
cortical brain damage underlie altered visual function. *Brain J Neurol* **131,** 1433–
1444.

Bullmore E & Sporns O (2012). The economy of brain network organization. *Nat Rev*
Neurosci **13,** 336–349.

Calhoun VD, Adali T, Pearlson GD & Pekar JJ (2001). A method for making group
inferences from functional MRI data using independent component analysis.
Hum Brain Mapp **14,** 140–151.

Caparelli EC, Backus W, Telang F, Wang G-J, Maloney T, Goldstein RZ, Anschel D & Henn
F (2010). Simultaneous TMS-fMRI of the Visual Cortex Reveals Functional
Network, Even in Absence of Phosphene Sensation. *Open Neuroimaging J* **4,**
100–110.

Chee MWL, Venkatraman V, Westphal C & Siong SC (2003). Comparison of block and
event-related fMRI designs in evaluating the word-frequency effect. *Hum Brain*
Mapp **18,** 186–193.

Chen N, Lu J, Shao H, Weng X & Fang F (2017). Neural mechanisms of motion perceptual
learning in noise. *Hum Brain Mapp* **38,** 6029–6042.

Cocchi L, Sale MV, L Gollo L, Bell PT, Nguyen VT, Zalesky A, Breakspear M & Mattingley
JB (2016). A hierarchy of timescales explains distinct effects of local inhibition of
primary visual cortex and frontal eye fields ed. Culham JC. *eLife* **5,** e15252.

Das A, DeMagistris M & Huxlin KR (2012). Different Properties of Visual Relearning after
Damage to Early Versus Higher-Level Visual Cortical Areas. *J Neurosci* **32,** 5414–
5425.

Egger P, Evangelista GG, Koch PJ, Park C-H, Levin-Gleba L, Girard G, Beanato E, Lee J,
Choirat C, Guggisberg AG, Kim Y-H & Hummel FC (2021). Disconnectomics of the
Rich Club Impacts Motor Recovery After Stroke. *Stroke*STROKEAHA120031541.

Farooqui M, Ortega-Gutierrez S, Hernandez K, Torres VO, Dajles A, Zevallos CB,
Quispe-Orozco D, Mendez-Ruiz A, Manzel K, Ten Eyck P, Tranel D, Karandikar NJ
& Ortega SB (2022). Hyperacute immune responses associate with immediate
neuropathology and motor dysfunction in large vessel occlusions. *Ann Clin*
Transl Neurol **10,** 276–291.

Fleury L et al. (2022). Toward individualized medicine in stroke-The TiMeS project:
Protocol of longitudinal, multi-modal, multi-domain study in stroke. *Front*
Neurol **13,** 939640.

Fornito A, Zalesky A & Breakspear M (2015). The connectomics of brain disorders. *Nat*
Rev Neurosci **16,** 159–172.

Foss J & Milton J (2000). Multistability in Recurrent Neural Loops Arising From Delay. *J*
Neurophysiol **84,** 975–985.

Glick C, Gajawelli N, Sun Y, Badami F, Saggar M & Etkin A (2024). Concurrent single-pulse
(sp) TMS/fMRI to reveal the causal connectome in healthy and patient
populations. *BioRxiv Prepr Serv Biol* 2024.09.25.614833.

Gollo LL, Roberts JA & Cocchi L (2017). Mapping how local perturbations influence
systems-level brain dynamics. *NeuroImage* **160,** 97–112.

Grasso PA, Làdavas E, Bertini C, Caltabiano S, Thut G & Morand S (2018). Decoupling of
Early V5 Motion Processing from Visual Awareness: A Matter of Velocity as
Revealed by Transcranial Magnetic Stimulation. *J Cogn Neurosci* **30,** 1517–1531.

Grossberg S, Mingolla E & Pack C (1999). A neural model of motion processing and visual
navigation by cortical area MST. *Cereb Cortex N Y N 1991* **9,** 878–895.

Himberg J, Hyvärinen A & Esposito F (2004). Validating the independent components of
neuroimaging time series via clustering and visualization. *NeuroImage* **22,** 1214–
1222.

Huang D-W & Yu Z-G (2017). Dynamic-Sensitive centrality of nodes in temporal
networks. *Sci Rep* **7,** 41454.

Imani E, Harati A, Pourreza H & Goudarzi MM (2021). Brain-behavior relationships in the
perceptual decision-making process through cognitive processing stages.
Neuropsychologia **155,** 107821.

Jimenez-Marin A, De Bruyn N, Gooijers J, Llera A, Meyer S, Alaerts K, Verheyden G,
Swinnen SP & Cortes JM (2022). Multimodal and multidomain lesion network
mapping enhances prediction of sensorimotor behavior in stroke patients. *Sci*
Rep **12,** 22400.

Joukes J, Hartmann TS & Krekelberg B (2014). Motion detection based on recurrent
network dynamics. *Front Syst Neurosci*; DOI: 10.3389/fnsys.2014.00239.

Jung J, Bungert A, Bowtell R & Jackson SR (2020). Modulating Brain Networks With
Transcranial Magnetic Stimulation Over the Primary Motor Cortex: A Concurrent
TMS/fMRI Study. *Front Hum Neurosci*; DOI: 10.3389/fnhum.2020.00031.

Kennerley SW & Walton ME (2011). Decision Making and Reward in Frontal Cortex.
Behav Neurosci **125,** 297–317.

Koivisto M, Leino K, Pekkarinen A, Karttunen J, Railo H & Hurme M (2021). Transcranial
magnetic stimulation (TMS)-induced Blindsight of Orientation is Degraded
Conscious Vision. *Neuroscience*; DOI: 10.1016/j.neuroscience.2021.08.025.

Koivisto M, Mäntylä T & Silvanto J (2010). The role of early visual cortex (V1/V2) in
conscious and unconscious visual perception. *NeuroImage* **51,** 828–834.

Kottaram A, Johnston L, Ganella E, Pantelis C, Kotagiri R & Zalesky A (2018).
Spatio-temporal dynamics of resting-state brain networks improve
single-subject prediction of schizophrenia diagnosis. *Hum Brain Mapp* **39,** 3663–
3681.

Lai TW, Zhang S & Wang YT (2014). Excitotoxicity and stroke: Identifying novel targets
for neuroprotection. *Prog Neurobiol* **115,** 157–188.

Lamme VA (2001). Blindsight: the role of feedforward and feedback corticocortical
connections. *Acta Psychol (Amst)* **107,** 209–228.

Lamme VA & Roelfsema PR (2000). The distinct modes of vision offered by feedforward
and recurrent processing. *Trends Neurosci* **23,** 571–579.

Latora V & Marchiori M (2001). Efficient behavior of small-world networks. *Phys Rev Lett*
87, 198701.

Leitão J, Thielscher A, Tünnerhoff J & Noppeney U (2015). Concurrent TMS-fMRI Reveals
Interactions between Dorsal and Ventral Attentional Systems. *J Neurosci* **35,**
11445–11457.

Lim S, Radicchi F, van den Heuvel MP & Sporns O (2019). Discordant attributes of
structural and functional brain connectivity in a two-layer multiplex network. *Sci*
Rep **9,** 2885.

Lin Z, Nie C, Zhang Y, Chen Y & Yang T (2020). Evidence accumulation for value
computation in the prefrontal cortex during decision making. *Proc Natl Acad Sci*
117, 30728–30737.

Litwińczuk MC, Muhlert N, Cloutman L, Trujillo-Barreto N & Woollams A (2022).
Combination of structural and functional connectivity explains unique variation
in specific domains of cognitive function. *NeuroImage* **262,** 119531.

Liu LD & Pack CC (2017). The Contribution of Area MT to Visual Motion Perception
Depends on Training. *Neuron* **95,** 436-446.e3.

Luo H, Ye X, Cai H-T, Wang M, Wang Y, Liu Q, Xu Y, Mao Z, Cai Y, Hong J, Zhang C, Wei P,
Lu Y, Liu Q, Xu J & Yuan T-F (2025). Frequency-specific and state-dependent
neural responses to brain stimulation. *Mol Psychiatry* **30,** 2880–2890.

Ma S, Huang T, Qu Y, Chen X, Zhang Y & Zhen Z (2022). An fMRI dataset for whole-body
somatotopic mapping in humans. *Sci Data* **9,** 515.

Mineault PJ, Khawaja FA, Butts DA & Pack CC (2012). Hierarchical processing of complex
motion along the primate dorsal visual pathway. *Proc Natl Acad Sci* **109,** E972–
E980.

Moore T, Rodman HR & Gross CG (2001). Direction of motion discrimination after early
lesions of striate cortex (V1) of the macaque monkey. *Proc Natl Acad Sci* **98,**
325–330.

Navarro de Lara LI, Tik M, Woletz M, Frass-Kriegl R, Moser E, Laistler E & Windischberger
C (2017). High-sensitivity TMS/fMRI of the Human Motor Cortex Using a
Dedicated Multichannel MR Coil. *NeuroImage* **150,** 262–269.

Navarro de Lara LI, Windischberger C, Kuehne A, Woletz M, Sieg J, Bestmann S,
Weiskopf N, Strasser B, Moser E & Laistler E (2015). A novel coil array for
combined TMS/fMRI experiments at 3 T. *Magn Reson Med* **74,** 1492–1501.

Newsome WT & Paré EB (1988). A selective impairment of motion perception following
lesions of the middle temporal visual area (MT). *J Neurosci Off J Soc Neurosci* **8,**
2201–2211.

Offen S, Gardner J, Schluppeck D & Heeger D (2010). Differential roles for frontal eye
fields (FEF) and intraparietal sulcus (IPS) in visual working memory and visual
attention. *J Vis* **10,** 28.

Ozdemir RA, Tadayon E, Boucher P, Momi D, Karakhanyan KA, Fox MD, Halko MA,
Pascual-Leone A, Shafi MM & Santarnecchi E (2020). Individualized perturbation
of the human connectome reveals reproducible biomarkers of network
dynamics relevant to cognition. *Proc Natl Acad Sci* **117,** 8115–8125.

Papo D, Buldú JM, Boccaletti S & Bullmore ET (2014). Complex network theory and the
brain. *Philos Trans R Soc B Biol Sci* **369,** 20130520.

Pascual-Leone A, Bartres-Faz D & Keenan JP (1999). Transcranial magnetic stimulation:
studying the brain-behaviour relationship by induction of "virtual lesions." *Philos*
Trans R Soc Lond B Biol Sci **354,** 1229–1238.

Perera ND, Wischnewski M, Alekseichuk I, Shirinpour S & Opitz A (2024). State-
Dependent Motor Cortex Stimulation Reveals Distinct Mechanisms for
Corticospinal Excitability and Cortical Responses. *eNeuro*; DOI:
10.1523/ENEURO.0450-24.2024.

Rafiei F & Rahnev D (2022). TMS Does Not Increase BOLD Activity at the Site of
Stimulation: A Review of All Concurrent TMS-fMRI Studies. *eNeuro*; DOI:
10.1523/ENEURO.0163-22.2022.

Ricci R, Salatino A, Li X, Funk A, Logan S, Mu Q, Johnson K, Bohning D & George M
(2012). Imaging the neural mechanisms of TMS neglect-like bias in healthy
volunteers with the interleaved TMS/fMRI technique: preliminary evidence.
Front Hum Neurosci; DOI: 10.3389/fnhum.2012.00326.

Romero MC, Davare M, Armendariz M & Janssen P (2019). Neural effects of transcranial
magnetic stimulation at the single-cell level. *Nat Commun*; DOI:
10.1038/s41467-019-10638-7.

Rossi S et al. (2021). Safety and recommendations for TMS use in healthy subjects and
patient populations, with updates on training, ethical and regulatory issues:
Expert Guidelines. *Clin Neurophysiol* **132,** 269–306.

Rubinov M & Sporns O (2010). Complex network measures of brain connectivity: uses
and interpretations. *NeuroImage* **52,** 1059–1069.

Ruff CC, Blankenburg F, Bjoertomt O, Bestmann S, Freeman E, Haynes J-D, Rees G,
Josephs O, Deichmann R & Driver J (2006). Concurrent TMS-fMRI and
Psychophysics Reveal Frontal Influences on Human Retinotopic Visual Cortex.
Curr Biol **16,** 1479–1488.

Ruff CC, Blankenburg F, Bjoertomt O, Bestmann S, Weiskopf N & Driver J (2008).
Hemispheric Differences in Frontal and Parietal Influences on Human Occipital
Cortex: Direct Confirmation with Concurrent TMS–fMRI. *J Cogn Neurosci* **21,**
1146–1161.

Ruff CC, Driver J & Bestmann S (2009). Combining TMS and fMRI. *Cortex J Devoted Study*
Nerv Syst Behav **45,** 1043–1049.

Rust NC, Mante V, Simoncelli EP & Movshon JA (2006). How MT cells analyze the motion
of visual patterns. *Nat Neurosci* **9,** 1421–1431.

Sani I, Stemmann H, Caron B, Bullock D, Stemmler T, Fahle M, Pestilli F & Freiwald WA
(2021). The human endogenous attentional control network includes a ventro-
temporal cortical node. *Nat Commun* **12,** 360.

Schwabe L, Obermayer K, Angelucci A & Bressloff PC (2006). The Role of Feedback in
Shaping the Extra-Classical Receptive Field of Cortical Neurons: A Recurrent
Network Model. *J Neurosci* **26,** 9117–9129.

Shao Z & Burkhalter A (1996). Different Balance of Excitation and Inhibition in Forward
and Feedback Circuits of Rat Visual Cortex. *J Neurosci* **16,** 7353–7365.

Siebner HR et al. (2022). Transcranial magnetic stimulation of the brain: What is
stimulated? – A consensus and critical position paper. *Clin Neurophysiol* **140,**
59–97.

Siegel JS, Ramsey LE, Snyder AZ, Metcalf NV, Chacko RV, Weinberger K, Baldassarre A,
Hacker CD, Shulman GL & Corbetta M (2016). Disruptions of network
connectivity predict impairment in multiple behavioral domains after stroke.
Proc Natl Acad Sci U S A **113,** E4367-4376.

Silvanto J & Pascual-Leone A (2008). State-Dependency of Transcranial Magnetic
Stimulation. *Brain Topogr* **21,** 1–10.

Simoncelli EP & Heeger DJ (1998). A model of neuronal responses in visual area MT.
Vision Res **38,** 743–761.

Telesford QK, Simpson SL, Burdette JH, Hayasaka S & Laurienti PJ (2011). The Brain as a
Complex System: Using Network Science as a Tool for Understanding the Brain.
Brain Connect **1,** 295–308.

Tie Y, Suarez RO, Whalen S, Radmanesh A, Norton IH & Golby AJ (2009). Comparison of
blocked and event-related fMRI designs for pre-surgical language mapping.
NeuroImage **47,** T107–T115.

Tu W, Ma Z & Zhang N (2021). Brain network reorganization after targeted attack at a
hub region. *NeuroImage* **237,** 118219.

Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N,
Mazoyer B & Joliot M (2002). Automated anatomical labeling of activations in
SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject
brain. *NeuroImage* **15,** 273–289.

Vink JJT, Mandija S, Petrov PI, van den Berg CAT, Sommer IEC & Neggers SFW (2018). A
novel concurrent TMS-fMRI method to reveal propagation patterns of prefrontal
magnetic brain stimulation. *Hum Brain Mapp* **39,** 4580–4592.

Wang J, Wang X, Xia M, Liao X, Evans A & He Y (2015). Corrigendum: GRETNA: a graph
theoretical network analysis toolbox for imaging connectomics. *Front Hum*
Neurosci. Available at:
https://www.frontiersin.org/articles/10.3389/fnhum.2015.00458 [Accessed July
23, 2022].

Wongtrakun J, Zhou S-H, O'Connell RG, Chong TT-J, Bellgrove MA & Coxon JP (2025). The
role of human intraparietal sulcus in evidence accumulation revealed by EEG
and model-informed fMRI. 2025.02.05.636628. Available at:
https://www.biorxiv.org/content/10.1101/2025.02.05.636628v2 [Accessed June
13, 2025].

Wu YK, Hengen KB, Turrigiano GG & Gjorgjieva J (2020). Homeostatic mechanisms
regulate distinct aspects of cortical circuit dynamics. *Proc Natl Acad Sci* **117,**
24514–24525.

Zhang Z, Yin C & Yang T (2022). Evidence accumulation occurs locally in the parietal
cortex. *Nat Commun* **13,** 4426.

Zhao X, Tian L, Yan J, Yue W, Yan H & Zhang D (2017). Abnormal Rich-Club Organization
Associated with Compromised Cognitive Function in Patients with Schizophrenia
and Their Unaffected Parents. *Neurosci Bull* **33,** 445–454.

1055

1056

1057