TITLE

Endothelial FOXC1 and FOXC2 promote intestinal regeneration after ischemia-

reperfusion injury

Can Tan,[1] Pieter R. Norden,[1] Ting Liu,[1] Naoto Ujiie,[1] Xiaocai Yan,[2] Kazushi Aoto,[3] Sagrario

Ortega,[4] Isabelle G. De Plaen,[2] and Tsutomu Kume[1]

[1]Feinberg Cardiovascular and Renal Research Institute, Department of Medicine, Feinberg

School of Medicine, Northwestern University, Chicago, Illinois, USA

[2]Department of Pediatrics, Feinberg School of Medicine, Northwestern University, Chicago,

Illinois, USA

[3]Department of Biochemistry, Hamamatsu University School of Medicine, Hamamatsu, Japan

[4]Mouse Genome Editing Unit, Biotechnology Program, Spanish National Cancer Research

Centre, Madrid, Spain

Correspondence to:

Tsutomu Kume, Feinberg Cardiovascular and Renal Research Institute, Department of Medicine,

Northwestern University School of Medicine, 300 E. Superior Street, Chicago, Illinois 60611,

USA

Phone: 312.503.0623; E-mail: t-kume@northwestern.edu

Abstract

Intestinal ischemia induces mucosal damage while simultaneously activating intestinal stem cells (ISCs), which subsequently regenerate the damaged intestinal epithelium. However, whether paracrine factors secreted from vascular endothelial cells (ECs) - blood and lymphatic ECs (BECs and LECs, respectively) – regulate ISC-mediated regeneration have yet to be elucidated. Here, we identify FOXC1 and FOXC2 as essential regulators of paracrine signaling in regeneration of the small intestine after ischemia-reperfusion (I/R) injury. EC- and LEC-specific deletions of *Foxc1*, *Foxc2*, or both in mice augment I/R-induced intestinal damage by causing defects in vascular regrowth, expression of the chemokine CXCL12 and the Wnt activator R-spondin 3 in BECs and LECs, respectively, and activation of Wnt signaling in ISCs. Treatment with CXCL12 and R-spondin 3 rescues the I/R-induced intestinal damage in EC- and LEC-*Foxc* mutant mice, respectively. This study provides evidence that FOXC1 and FOXC2 are required for intestinal regeneration by stimulating paracrine CXCL12 and Wnt signaling.

Introduction

Tissue regeneration and repair is essential for maintaining physiological homeostasis and relies on the precise control of molecular networks that regulate, or are regulated by, the vasculature. Endothelial cells (ECs) present in the blood and lymphatic vessels are crucial participants in the vascular-dependent processes that restore damaged tissue because they control the secretion of paracrine factors from both the vessels themselves and nearby cells [1]. However, the fundamental mechanisms by which the vascular system regulates tissue regeneration and repair are poorly understood. Thus, an adequate understanding of the biological processes that contribute to EC-dependent tissue repair, as well as their roles in the pathogenesis and potential treatment of vascular disease or injury, is crucially dependent on a thorough characterization of how blood/lymphatic vessels control tissue regeneration.

Intestinal ischemia is a life-threatening vascular emergency and can be caused by thrombus formation in the mesenteric vasculature, embolisms that arise as a consequence of cardiopulmonary disease, and disease- or shock-induced declines in perfusion, as well as when blood flow is interrupted by intestinal transplantation [2, 3]. Moreover, impairments in intestinal microvasculature development contribute to the pathogenesis of neonatal necrotizing enterocolitis (NEC), which is the most common life-threatening gastrointestinal emergency in neonatal patients [4]. Both homeostasis and repair of the small intestine are mediated via intestinal stem cells (ISCs) [5, 6]. Active ISCs express the specific marker leucine-rich repeat-containing G protein-coupled receptor 5 (Lgr5) and are located at the base of the crypts of the small intestine where they vigorously proliferate to continuously regenerate the intestinal epithelium. Ischemia/reperfusion (I/R) injury, as well as radiation injury and stresses such as acute

inflammation, induce apoptosis in proliferating Lgr5$^+$ ISCs [7, 8], while ISC regeneration after

injury is restored largely by dedifferentiation of crypt cells [9]. However, the mechanisms that

coordinate the role of intestinal ECs in intestinal regeneration and repair, and whether blood and

lymphatic ECs (BECs and LECs) in the ISC niche regulate the regenerative activity of ISCs as

well as the preservation of the ISC niche after injury, have yet to be fully elucidated.

In adult mice, the proliferation of active ISCs is controlled, in part, by Wnt/β-catenin signaling [10,

11], and canonical Wnt/β-catenin signaling is promoted by the cooperative activity of Wnt

proteins and R-spondins (RSPO1-RSPO4). Lgr5 functions (with Lgr4 and Lgr6) as a cognate

receptor for R-spondins, and R-spondins are expressed in mesenchymal stromal cells of the ISC

niche [12, 13], including Pdgfra^{lo}Grem1+ trophocytes as an essential source of R-spondins (RSPO1-

RSPO3, especially RSPO3) [14]. Foxl1+ mesenchymal cells (telocytes) and other non-epithelial

stromal cells express Wnt ligands in the ISC niche [13, 15, 16]. Both the number of Lgr5$^+$ ISCs and

the regenerative response to intestinal radiation injury are reduced by cotreatment with R-

spondin 2- and R-spondin 3-neutralizing antibodies [17], and the mucosal damage induced by

intestinal I/R injury can be rescued by treatment with R-spondin 3 [18]. Most importantly, intestinal

RSPO3 [12, 14, 19] and Wnt2 [14] are highly produced by LECs.

The CXC chemokine CXCL12, also called stromal cell-derived factor 1 (SDF-1), is a

homeostatic chemokine expressed in many cell types such as stromal cells, ECs, and fibroblasts

in various tissues [20]. CXCL12 is an essential factor for angiogenesis that involves EC

proliferation and migration to form neo-vessel networks [21]. CXCL12 can be induced by hypoxic

stress [22, 23] and regulates angiogenesis in an autocrine/paracrine manner by interacting with the

CXCR4 and CXCR7 receptors [24]. In the small intestine, CXCR4 is expressed in the crypt

epithelial cells [25, 26]. CXCR4 deficiency in the intestinal epithelium impairs re-epithelialization

after acute dextran sodium sulfate (DSS)-induced injury in mice [26], whereas CXCL12-

overexpressing mesenchymal stem cells (MSCs), distributed in the villous stroma of the small

intestine when administered to mice, activate CXCR4 in the intestinal crypts by activating β-

catenin and improve survival and intestinal epithelial recovery after radiation injury [25]. Thus,

CXCL12-CXCR4 signaling and the canonical Wnt/β-catenin pathway appear to interact during

intestinal regeneration.

FOXC1 and FOXC2 are closely related members of the FOX transcription factor family and

have numerous essential roles in cardiovascular development, health, and disease [27]. Mutations

or changes in the copy number of human *FOXC1* are associated with autosomal-dominant

Axenfeld-Rieger syndrome, which is characterized by abnormalities in the eye and extraocular

defects [28], while inactivating mutations of human *FOXC2* are responsible for the autosomal

dominant syndrome Lymphedema-distichiasis, which is characterized by obstructed lymph

drainage in the limbs and the growth of extra eyelashes [29]. There is also some evidence that

Foxc2 haploinsufficiency in mice increases their susceptibility to DDS-induced colitis [30];

however, the precise function of FOXC1 and FOXC2 in vascular repair and intestinal

regeneration after ischemic injury have yet to be determined.

In this study, we report that FOXC1 and FOXC2 in intestinal BECs and LECs contributes to

vascular repair and intestinal regeneration after I/R injury by regulating expression of paracrine

signaling factors. Inducible, EC- and LEC-specific, single and compound mutant mice for *Foxc1*

and *Foxc2* showed that the *Foxc* mutations impair regeneration of the small intestine after I/R

injury, accompanied by (1) defective repair of intestinal BECs and LECs, (2) reduced expression

of CXCL12 and R-spondin 3 in intestinal BECs and LECs, respectively, and (3) decreased

activation of the Wnt/β-catenin pathway in ISCs. Importantly, treatment with either CXCL12 or

R-spondin 3 partially rescues the defects in intestinal repair and regeneration associated with EC-

and LEC-*Foxc1/c2* deficiency. Together, our data show a new role for FOXC1 and FOXC2 as

key mediators of paracrine signaling in the intestinal blood/lymphatic vasculature during post-

ischemic intestinal repair/regeneration, and our findings may have important implications for

other ischemic conditions that are associated with impairments in tissue regeneration.

Results

Foxc1 and Foxc2 expression in the murine intestine

Foxc1 and *Foxc2* expression in intestinal BECs and LECs was evaluated via quantitative PCR (qPCR) of $CD45^-/CD31^+$ ECs, which include both BECs and LECs isolated from the small intestine in adult mice (Figure 1A). Intestinal ECs are particularly vulnerable to I/R injury, and they undergo apoptosis in response to oxidative stress [2]. Thus, we next investigated whether I/R injury alters *Foxc1/c2* expression in the small intestine of adult mice. Intestinal I/R injury [31] was induced by clamping the superior mesenteric artery for 30 minutes to induce occlusion; then, the clip was removed, and the small intestine was allowed to re-perfuse. *Foxc1/Foxc2* expression was significantly greater in the intestinal $CD45^-/CD31^+$ ECs of mice 4 hours after I/R injury than in the intestinal ECs of sham-operated animals (Figure 1, B and C). These results are consistent with previous studies showing that FOXC2 expression in renal tubular cells of the cortex and outer medulla also increases 24 hours after kidney I/R injury [32]. Comparison of the relative expression levels of *Foxc1* and *Foxc2* in the intestinal ECs 4 hours after I/R injury (Figure 1D) and with those under sham (Figure 1A) revealed that both *Foxc1* and *Foxc2* were increased to a similar extent after I/R injury. Notably, *Foxc1* was more highly expressed than *Foxc2* under the basal and I/R-induced conditions.

To further investigate FOXC1 expression in the adult mouse intestinal ECs, we performed whole-mount immunostaining labeled with CD31 (EC marker), LYVE1 (LEC marker) and FOXC1. In villus, FOXC1 was hardly detectable in both BECs and LECs, while in submucosa, FOXC1 was weakly expressed in the nuclei of BECs, but was still hardly detectable in submucosal LECs (Figure 1E, no surgery). Four hours after I/R injury, blood and lymphatic

vessels were damaged severely, and both FOXC1 and FOXC2 protein levels were upregulated in

both intestinal BECs and LECs (Figure 1E and Supplementary Figure 1A). To investigate *Foxc2*

expression cells in the intestinal vasculature, we crossed tamoxifen-inducible *Foxc2-Cre[ERT2]*

knock-in mice [33] with dual Rosa26mTmG reporter mice [34]. Adult *mTmG/+;Foxc2-Cre[ERT2]* mice

were treated with tamoxifen (150 mg/kg) by oral gavage for 5 consecutive days. Consistent with

immunohistochemical analysis of FOXC2, *Foxc2*-expressing, recombined EGFP+ cells were

also detected in lacteals and villous capillaries of small intestines in adult *mTmG/+;Foxc2-*

Cre[ERT2] mice after intestinal I/R injury (Supplementary Figure 1, B and C).

Expression patterns of FOXC1 and FOXC2 in the intestine were also examined in a mouse NEC

model [35], which includes initial orogastric inoculation of neonatal mice with a standardized adult

mouse commensal bacteria preparation and lipopolysaccharide (LPS) to perturb the normal

intestinal colonization process, gavage with formula every 3 hours, and exposure to brief

episodes of hypoxia for 1 min followed immediately by cold stress (10 min at 4°C) twice daily.

With this protocol, about 50-70% of mice typically develop between 36-72 hours intestinal

injuries ranging from epithelial injury to transmural necrosis [35]. At 24 hours after the neonates

were subjected to the NEC protocol, immunohistochemical analyses of BEC (CD31 and

endomucin [EMCN]) and LEC (PROX1) markers revealed that levels of FOXC1 and FOXC2

were increased in intestinal BECs and LECs (Supplementary Figures 2 and 3), compared to dam-

fed littermate controls.

Generation of tamoxifen-inducible, EC-specific, *Foxc1/c2*-mutant mice in the adult

Murine *Foxc1* and *Foxc2* are both required for vascular development [36, 37, 38], but attempts to

determine how the two genes function during pathological (lymph)angiogenesis have been

generally unsuccessful, because global single and compound *Foxc1/c2* mutant mice die

perinatally with severe cardiovascular abnormalities [37]. Therefore, we crossed conditional-null

Foxc1^{fl} and *Foxc2^{fl}* mutant mice [39] with *Cdh5-Cre^{ERT2}* mice [40] to generate tamoxifen-inducible,

EC-specific, compound *Foxc1;Foxc2*-mutant (*Cdh5-Cre^{ERT2};Foxc1^{fl/fl};Foxc2^{fl/fl}*) mice, which

(after the mutation is induced) are referred to as EC-*Foxc*-DKO mice [41]. To induce the mutations,

adult mice were treated with tamoxifen (150 mg/kg) by oral gavage for 5 consecutive days, and

12 days after tamoxifen treatment was completed (Figure 1F), qPCR and immunohistochemical

analyses confirmed that *Foxc1*- and *Foxc2*-expression was significantly reduced in intestinal ECs

of EC-*Foxc*-DKO mice than in the corresponding cells of control littermates (Figure 1, G-H).

Importantly, the small intestines of EC-*Foxc*-DKO mice appeared morphologically normal 12

days after tamoxifen treatment (Supplementary Figure 1D), suggesting that EC expression of

Foxc1 and *Foxc2* is not required for maintaining intestinal epithelium homeostasis. We also

crossed EC-*Foxc*-DKO mice with Rosa26mTmG mice [34], then treated their adult offspring with

tamoxifen as described above, and identified recombined EGFP^+ cells in the small intestines to

confirm the efficiency of Cre-mediated recombination in intestinal BECs and LECs

(Supplementary Figure 4A).

EC-specific deletion of *Foxc1* and *Foxc2* decreases intestinal mucosal recovery after I/R

When intestinal I/R injury was induced 12 days after tamoxifen treatment in adult mice (Figure

2A), tissue from the distal jejunum was histologically graded and quantified 24 hours after I/R

injury according to the Chiu scoring system [42]. Compared to the control mice, the intestinal

mucosa remained severely injured in EC-*Foxc*-DKO mice (Figure 2, B and C). To further

characterize how the loss of EC-specific *Foxc1/c2* expression affects repair of the intestinal

mucosa during recovery from I/R injury, we also examined mice carrying tamoxifen-inducible,

EC-specific mutations of each individual gene (i.e., EC-*Foxc1*-KO and EC-*Foxc2*-KO mice) and

in their control littermates. Recovery from intestinal damage was impaired in both EC-*Foxc1*-

KO and EC-*Foxc2*-KO mice 24 hours after I/R injury (Figure 2, D and E; Supplementary Figure

5, A and B). As the extent of intestinal injury in the EC-specific *Foxc* single mutant mice was

less than that in the double *Foxc1/c2* mutant mice, EC-*Foxc1* and -*Foxc2* expression is required

for intestinal epithelial regeneration. Since I/R-induced local inflammatory response is critically

associated with intestinal damage [18, 43], mRNA levels of proinflammatory regulators (*TNFα*, *IL-6*

and *Cox2*) were measured 3 hours after I/R injury via qPCR. Expression levels of *TNFα* and

Cox2 were significantly higher in EC-*Foxc*-DKO mice than in their control mice (Figure 2F),

whereas increased *IL-6* expression in EC-*Foxc*-DKO mice exhibited a trend toward significance

(P=0.2014). Furthermore, EC-specific loss of *Foxc* genes significantly reduced the proliferative

response of intestinal epithelial cells to I/R injury as assessed via immunostaining of EpCAM

(intestinal epithelial cell marker) and BrdU (proliferative marker) (Figure 2G), as well as the

quantification of BrdU positive epithelial cell number per crypt (Figure 2H).

Wnt signaling in the small intestine is impaired in EC-*Foxc*-DKO mice after I/R injury

β-catenin regulates the maintenance and regeneration of intestinal epithelial cells [10, 11] by

translocating from the cytosol to the nucleus of ISCs and altering gene expression in response to

activation of the canonical Wnt signaling pathway. This mechanism is consistent with our

observations in control mice, because although β-catenin was located at the adherens junctions of

epithelial cells in the villus, nuclear β-catenin was detected in ISCs co-immunostained with the

ISC marker OLFM4 [44] at the crypt base after I/R injury [13]. However, nuclear localization of β-

catenin was impaired (Figure 3, A and B) and the number of cells expressing the Wnt target

cyclin D1 (CCND1) [13] was significantly reduced in the crypt of EC-*Foxc*-DKO mice both 24 and

48 hours after I/R injury (Figure 3, C and D). Together, the loss of EC-specific *Foxc1/c2*

expression impedes I/R-induced Wnt signaling in ISCs.

Defective mucosal recovery in LEC-specific *Foxc* mutant mice after intestinal I/R injury

As *Cdh5* is expressed in both the blood and lymphatic vasculatures, we generated mice carrying

a tamoxifen-inducible, LEC-specific, compound homozygous $Foxc1^{-/-};Foxc2^{-/-}$ mutation

(referred to herein as LEC-*Foxc*-DKO mice) by breeding conditional-null $Foxc1^{fl/fl}$ and $Foxc2^{fl/fl}$

mice [39] with LEC-specific $Vegfr3\text{-}Cre^{ERT2}$ mice [45] to investigate intestinal LEC-specific functions

of *Foxc1* and *Foxc2* during I/R. We then confirmed *Vegfr3-Cre*-mediated recombination limited

to intestinal LECs of adult LEC-*Foxc*-DKO mice crossed with the Rosa26mTmG reporter mice

(Supplementary Figure 4B). The LEC-*Foxc*-DKO mice and their littermate control mice were

subjected to the sham or I/R injury procedures, and LEC-specific deletion of *Foxc1* and *Foxc2*

significantly increased the severity of intestinal mucosa injury 24 hours after I/R (Figure 4A and

Supplementary Figure 5C). Furthermore, increased intestinal damage was also noted in single

LEC-*Foxc1*-KO and LEC-*Foxc2*-KO mice after I/R (Figure 4, B and C; Supplementary Figure 5,

D and E), suggesting that both FOXC1 and FOXC2 are required in intestinal LECs for intestinal

repair in response to I/R injury. In addition, proliferating (BrdU+) intestinal epithelial cells were

quantified in the crypts of the LEC-*Foxc*-DKO mice 24 hours after I/R injury, and the LEC-

specific loss of *Foxc1/c2* reduced the proliferative response of intestinal epithelial cells to I/R

injury (Figure 4D). Consistent with the increased severity of intestinal mucosa defects in the

LEC-*Foxc*-DKO mice (Figure 4A), the activation of Wnt signaling (i.e., nuclear localization of

β-catenin, Figure 4, E and F) in ISCs and the number of Cyclin D1+ (CCND1+) cells per crypt

after intestinal I/R injury were both reduced in the LEC-*Foxc*-DKO mice compared to the control

mice (Figure 4G, and Supplementary Figure 6).

Defective vascular recovery in EC- and LEC-specific deletions of *Foxc1/c2* genes after

intestinal I/R injury

The vascular recovery of villus BECs and LECs after I/R injury proceeds via a stepwise process

in which blood capillaries (BECs) regrow earlier than lacteals (LECs) in the villous stroma [46].

We found that the EC-*Foxc*-DKO mutation was associated with defective vascular repair of

intestinal BECs and LECs after I/R injury (Figure 5). Vascular endothelial growth factor (VEGF)

receptor (R) 2 (VEGFR2) and VEGFR3 were highly expressed in the growing tips of villous

BECs and LECs, respectively, from control mice after intestinal I/R injury but were severely

diminished in EC-*Foxc*-DKO mice (arrows in Figure 5). Notably, proliferation of intestinal

BECs and LECs was also significantly reduced (Figure 6, A and B), while the number of

apoptotic BECs and LECs was significantly increased (Figure 6, C and D), in EC-*Foxc*-DKO

mice than in their control mice after I/R injury. Following whole-mount immunostaining for

CD31 and LYVE1 (Supplementary Figure 7, A and B; Supplementary videos), the length of

blood capillaries and lacteals (Figure 6E) as well as the percentage of lacteal length to blood

capillary length (Figure 6F) were measured. The regrowth of both blood capillaries and lacteals

in the villi were significantly decreased in EC-*Foxc*-DKO mice after intestinal I/R injury,

whereas EC-*Foxc*-double mutant lacteals were shorter than controls in sham treatments.

Similarly, shorter lacteals were found in proximal jejunums in EC-*Foxc*-DKO neonatal mice at

P7 after Tm treatment from P1 to P5 (Supplementary Figure 7, C-E), suggesting that *Foxc1/c2*

are required for the maintenance of lacteal length. Furthermore, whole-mount immunostaining

for VEGFR2 (Figure 5A) and subsequent quantification reveal a reduction in branches (Figure

6G) and branching points (Figure 6H) of blood capillaries in EC-*Foxc*-DKO mice after intestinal

I/R injury. Together, these results suggest that EC-*Foxc1/c2* expression contributes to the repair

of ischemic intestinal mucosa by promoting the recovery of BECs and LECs.

***Reduced R-spondin 3* and *Cxcl12* expression in intestinal LECs and BECs of EC-*Foxc*-DKO**

mice, respectively, after I/R injury

To investigate molecular mechanisms associated with impaired intestinal regeneration in EC-

Foxc-DKO mutants following I/R, we performed single-cell RNA sequencing (scRNA-seq)

analyses of distal jejuna from control and EC-*Foxc*-DKO mice 18.5 hours after I/R injury.

Dimensionality reduction and clustering analysis identified 22 transcriptionally distinct cell

clusters (Figure 7A and Supplementary Figure 8A) based on known gene markers for each

specific cell type (Supplementary Table 1), including BECs, LECs, stromal cells, epithelial cells

and other cell types. When interpreted according to the recent classification of stromal cell

populations in the ISC niche [12], the results from our scRNA-seq experiments indicates that *R-*

spondin 3 (*Rspo3*) is mainly expressed in two clusters (LECs and Telocytes/Trophocytes) after

intestinal I/R injury (Figure 7B, cluster 12 and 20, respectively). Low expression of *Rspo3* was

found in Myocytes/Pericytes cluster (Figure 7B, cluster 17). Further sub-clustering performed on

Telocytes/Trophocytes cluster based on their known markers [14] identified 3 cell clusters,

Trophocytes, Pdgfra^{lo} Cd81- stromal cells, and Telocytes (Supplementary Figure 8B). A dot plot

used for visualizing differential gene expression (mean expression level) as well as gene

expression frequency in different cell clusters (Figure 7C) showed that *Rspo3* was decreased in

both LECs and Trophocyte clusters in EC-*Foxc*-DKO intestines compared with controls after I/R

injury. In Trophocyte cluster, the moderately decreased trend of *Rspo3* was not significant

(Supplementary Figure 8C). Similar *Rspo3* levels were found in the other 3 cell clusters between

the two groups. Since the percentage of the LECs captured by scRNA-seq was very low (0.373%,

Supplementary Figure 8A), the number of LECs obtained and used for analysis was very limited,

and the decrease of *Rspo3* expression in LECs was not significant (Supplementary Figure 8D).

However, by validating via IHC, RSPO3 protein was lower in intestinal LECs from EC-*Foxc*-

DKO mice than control mice 24 hours after I/R injury (Figure 7D).

The results from our scRNA-seq experiments also identified numerous genes that were

differentially expressed in intestinal BECs from EC-*Foxc*-DKO and control mice after I/R injury

(Figure 7E), including the anti-angiogenic factor *Adamts1* [47, 48] and the CXC chemokine *Cxcl12*,

which were up- and down-regulated, respectively, in EC-*Foxc*-DKO BECs. Notably, although

Cxcl12 was also expressed in both Telocyte/Trophocyte and Fibroblast clusters (Figure 7, F and

G), it was significantly downregulated only in BECs (Figure 7, E and G), which have a larger

cell population than the other two cell clusters (Figure 7F and Supplementary Figure 8A).

Furthermore, qPCR analysis validated that *Cxcl12* expression was significantly reduced in

isolated intestinal CD45$^-$/CD31$^+$ ECs of both sham- and I/R-treated EC-*Foxc*-DKO mice

compared to their littermate controls (Figure 7H).

R-spondin 3 treatment rescues the defective repair of small intestines in EC- and LEC-
301

302 *Foxc*-DKO mice after I/R injury

303 RSPO3 is a key regulator of Wnt signaling during intestinal regeneration [17, 18], and RSPO3

304 prevents I/R-induced intestinal tissue damage and vascular leakage [18]. Accordingly, we

305 investigated whether RSPO3 treatment can rescue the defective repair of small intestines in EC-

306 *Foxc*-DKO mice following I/R injury. Adult EC-*Foxc*-DKO mice were randomly assigned to

307 two experimental groups and were treated with PBS or RSPO3 (5 μg in 100 μL PBS per mouse)

308 by retro-orbital injection 30 min before intestinal ischemia. Quantification of Chiu scores was

309 then performed 24 hours after I/R injury. RSPO3 treatment partially rescued the defective repair

310 of small intestines in EC-*Foxc*-DKO mice (Figure 8, A and B). Given that RSPO3 is expressed

311 in LECs of the small intestine, but not in BECs (Figure 7B), equivalent experiments were

312 subsequently performed with adult mice carrying LEC-specific mutations of both *Foxc* genes.

313 Similar rescue effects of the RSPO3 treatment were observed in LEC-*Foxc*-DKO intestines 24

314 hours after I/R injury with improvement of the Chiu score to the level similar to I/R-exposed

315 control mice (i.e., grade 1) (Figure 8, C and D). Thus, treatment with RSPO3 is sufficient to fully

316 recover intestinal I/R injury in LEC-specific *Foxc1/c2* deficient mice.

317

318 Immunoglobulin M (IgM) and complement also contribute to intestinal I/R injury [18], but RSPO3

319 treatment suppresses the deposition of IgM and complement in damaged intestinal tissue [18],

320 likely by enhancing the activation of Wnt signaling in ISCs of the small intestine. Therefore,

321 IgM deposition was evaluated in damaged intestines of control and EC-*Foxc*-DKO mice 24

322 hours after I/R injury via immunostaining (Supplementary Figure 9). Compared to control mice,

323 IgM deposition was increased in EC-*Foxc*-DKO mice, whereas administration of RSPO3

reduced the level of IgM in EC-*Fox*-DKO mice. The levels of IgM deposition were further

quantified via western blotting [18, 31]. I/R-enhanced IgM levels in the intestinal tissues were

greater in EC-*Foxc*-DKO mice compared to controls (Figure 8, E and F). More importantly,

RSPO3 treatment significantly reduced the levels of IgM in the EC-*Foxc*-DKO mice after I/R

injury (Figure 8, G and H). Collectively, these observations demonstrate that *Foxc1* and *Foxc2*

expression in the intestinal vasculature contributes to intestinal mucosal recovery and

regeneration after I/R injury.

CXCL12 treatment partially rescue the defective repair of small intestines in EC-*Foxc*-

DKO mice after I/R injury

Cxcl12 was only downregulated in intestinal BECs, but not in the Telocyte/Trophocyte and

Fibroblast clusters of EC-*Foxc*-DKO mutant intestine after I/R injury (Figure 7, E and G). Given

evidence that hypoxia upregulates CXCL12 [22, 23] as a key regulator of angiogenesis [24], we

examined whether pre-treatment with CXCL12 can rescue the defects in the vascular regrowth

and intestinal repair associated with EC-*Foxc1/c2* deficiencies. Adult EC-*Foxc*-DKO mice were

treated with PBS or CXCL12 via retro-orbital injection as described previously [49], and I/R

induction was performed 30 minutes later. The degree of intestinal mucosal damage 24 hours

after I/R injury was then quantified via the Chiu scoring system. CXCL12 treatment partially

rescued the defective repair of small intestines in the EC-*Foxc*-DKO mice (Figure 9, A and B).

Moreover, nuclear localization of β-catenin was observed in crypt ISCs of CXCL12-pretreated

EC-*Foxc*-DKO mice after I/R injury compared to the PBS-treatment (Figure 9, C and D), and the

number of cells expressing the Wnt target cyclin D1 (CCND1) [13] was significantly increased in

the crypt ISCs of EC-*Foxc*-DKO mice treated with CXCL12 after I/R injury (Figure 9E).

Administration of CXCL12 also improved the defective vascular recovery (CD31+ vessel

density) of EC-*Foxc*-DKO mice 24 hours after I/R injury (Figure 9, F and G). Therefore,

CXCL12 treatment rescues the defects in vascular regrowth and intestinal repair associated with

EC-*Foxc1/c2* deficiencies by enhancing vascular recovery and stimulating Wnt signaling in the

crypts of the intestinal epithelium.

Discussion

Little is known about the mechanisms that regulate expression of BEC/LEC-derived paracrine factors in tissue regeneration. In the present study, we demonstrate that EC-*Foxc1/Foxc2* expression is crucial for repair of the intestinal mucosa, BECs, and LECs after I/R injury, and that the EC- and LEC-*Foxc*-DKO mutations in mice impair canonical Wnt/β-catenin signaling in ISCs at the crypt base. Furthermore, our scRNA-seq data indicate that RSPO3 expression is attenuated in LECs and stromal cells of the EC-*Foxc*-DKO mice after intestinal I/R injury, which is at least partially attributable to impairments in intestinal regeneration because ISC activity appears to be crucially dependent on Wnt/β-catenin signaling in the subepithelial cellular microenvironment in the ISC niche [10, 11]. We also show that CXCL12 is reduced in intestinal BECs of the EC-*Foxc*-DKO mice after I/R injury. Most significantly, treatment with RSPO3 and CXCL12 rescues the defective repair of small intestines in the LEC-*Foxc*-DKO and EC-*Foxc*-DKO mice, respectively. Consequently, this study elucidates the novel mechanisms that mediate the role of EC-specific FOXC1/C2 in repair of intestinal mucosa and vasculature, regulation of paracrine signaling factors, and activation of ISCs after intestinal I/R injury.

Adult single and compound EC- (or LEC)-specific *Foxc1/c2* mutant mice provide us with the *first* opportunity to comprehensively characterize how *Foxc* genes regulate the post-ischemic repair of intestinal blood/lymphatic vessels after I/R injury and intestinal epithelial regeneration by modulating Wnt/RSPO and CXCL12 signaling. Because the EC-*Foxc*-DKO and LEC-*Foxc*-DKO mutations are induced by different drivers (i.e., *Cdh5-Cre^{ERT2}* and *Vegfr3-Cre^{ERT2}*), the extent of *Foxc* downregulation in the two mutant mouse lines may not be equal. However, the degree of impairments in intestinal repair is consistently greater in the EC-*Foxc*-DKO mutant

line than in the LEC-*Foxc*-DKO mutant line, suggesting that *Foxc1* and *Foxc2* are required in

both BECs and LECs for intestinal tissue repair. Equivalent experiments were also performed

with adult mice carrying EC- (and LEC-) specific KO mutations of each individual *Foxc* genes

to determine the similarities and differences between the phenotypes associated with each

deletion. While our qPCR and scRNA-seq analyses show that expression levels of *Foxc1* in

intestinal ECs are higher than those of *Foxc2*, the phenotypic differences between EC- (or LEC-)

Foxc1-KO and EC- (or LEC-) *Foxc2*-KO mice are not particularly distinct. Although the

reason(s) for the phenotypic similarities remains unclear, recent evidence indicates that *Foxc2* is

essential for the maintenance of intestinal LECs, and that treatment with antibiotics to deplete

gut microbiota rescues the phenotype of LEC-specific *Foxc2* mutants, including lymphatic

dilation in the small intestine [50]. Bacterial translocation occurs when the intestinal barrier is

disrupted by I/R injury, and I/R-induced intestinal damage can be attenuated by pretreatment of

an antibiotic cocktail to deplete gut commensal bacteria [31]. It remains to be elucidated whether

commensal bacteria contribute to aberrant intestinal recovery and regeneration in our EC/LEC-

specific *Foxc* mutant mice.

The EC-specific mutations of both *Foxc* genes in mice impair the regulation of RSPO3 in LECs

(Figure 7), accompanied by reduced expression of ISC markers and Wnt target genes in crypt

cells (Supplementary Figure 8E), and that RSPO3 treatment rescues the defective repair of small

intestines in both EC-*Foxc*-DKO and LEC-*Foxc*-DKO mice after I/R injury (Figure 8). These

observations are consistent with reports that RSPO3 prevents I/R-induced intestinal tissue

damage [18], and that RSPO3 is a key regulator of Wnt signaling during intestinal regeneration [17,

18]. While intestinal RSPO3 is known to be produced by LECs [12, 19], we demonstrate that

pretreatment with RSPO3 almost completely rescues the impaired intestinal repair in the LEC-

Foxc-DKO mice, implicating a paracrine effect of LEC-mediated RSPO3 signaling on the

activation of ISCs after I/R injury. We analyzed data deposited in the ENCODE database [51] with

HOMER software [52], as recently described in our analysis of the *PRICKLE1* locus [41], and

identified putative FoxC binding sites (RYMAAYA or RYACACA) [53, 54, 55] in the vicinity of the

human *RSPO3* locus (Supplementary Figure 10A and Methods). These sites are conserved

between human and mouse, and contain histone methylated and acetylated regions (e.g.,

H3K4Me1 and H3K27Ac ChIP peaks), DNaseI hypersensitive regions, and transcriptionally

active regions (identified by Transcription Factor ChIP-seq data). Thus, FOXC1 and FOXC2 are

likely to regulate *RSPO3* expression directly in LECs. During our manuscript preparation, others

have submitted bioRxiv preprints [56, 57, 58] showing that RSPO3 produced by intestinal LECs is

required for maintaining ISCs in homeostasis and regeneration, reinforcing our finding that

FOXC transcription factors mediate LEC-derived RSPO3 expression as a critical paracrine

regulator in intestinal regeneration after I/R injury [59].

The chemokine CXCL12 is an angiocrine factor that is secreted from BECs and regulates organ-

specific tissue repair, regeneration, and homeostasis, including the liver, bone marrow, and lung [1,

60, 61]. However, the function of BEC-derived CXCL12 paracrine/autocrine signaling in damaged

organs after injury remains largely unknown. The results from our study demonstrate that

CXCL12 expression is downregulated in the intestinal BEC cluster and intestinal CD45-/CD31+

ECs of the EC-*Foxc*-DKO mice after I/R injury (Figure 7), and that CXCL12 treatment partially

rescues the defective repair of small intestines in EC-*Foxc*-DKO mice after I/R injury (Figure 9).

Our results are in accord with recent evidence that FOXC1 controls CXCL12 expression in

CXCL12-abundant reticular (CAR) progenitors and Schwann cells [62, 63], and we also identified

putative FOXC binding sites [53, 54, 55] in evolutionarily conserved regions of the human *CXCL12*

locus (Supplementary Figure 10B and Methods). EC-derived CXCL12 promotes angiogenesis

via an *autocrine* mechanism [24], and CXCL12-CXCR4 signaling cooperates with VEGF to

enhance angiogenic processes such as the morphogenesis and sprouting of vascular tubes [64].

Since VEGFR2 levels at the angiogenic front of growing blood capillaries in the villus are lower

in EC-*Foxc*-DKO mice than in control mice after intestinal I/R injury (Figure 5A), it is likely

that intestinal EC-derived CXCL12 autocrine signaling is defective in EC-*Foxc*-DKO mice.

Consistently, impaired vascular recovery of the EC-*Foxc*-DKO mice after intestinal I/R injury is

rescued by CXCL12 treatment (Figure 9). Since the vascular regrowth of villus BECs and LECs

after I/R injury proceeds via a stepwise, interactive process, and regeneration of villus blood

vessels precedes reconstruction of lacteals [46] (Figure 6 and Supplementary Videos), the severity

of mucosal damage is greater in the EC-*Foxc*-DKO mice than in the LEC-*Foxc*-DKO mice.

When the CXCR4 receptor in intestinal epithelial cells is activated by treatment with CXCL12-

overexpressing MSCs, canonical Wnt/β-catenin is stimulated in the ISCs [25]. Since CXCL12

treatment activates Wnt signaling in the ISC of the EC-*Foxc*-DKO mice (Figure 9), our results

demonstrate that FOXC1 and FOXC2 promote endothelial-derived CXCL12 signaling in

intestinal ischemia, thereby controlling vascular regrowth and intestinal regeneration.

In summary, our study demonstrates that BEC/LEC-FOXC1/FOXC2 expression regulates the

repair of the intestinal vasculature and mucosal damage during intestinal regeneration after I/R

injury by controlling the RSPO3 and CXCL12 signaling pathways. Thus, FOXC1 and FOXC2

regulate blood/lymphatic vascular function in the small intestine, as well as vascular mediated

signaling to ISCs and the ISC niche during intestinal repair. Collectively, this study provides new

insights into fundamental processes that are critically involved in recovery from ischemic disease

and injury and may have important implications for the treatment of other ischemic conditions

that are associated with impairments in tissue regeneration and stem-cell activity, including

cardiovascular disease.

Methods

Animal husbandry

$Foxc1^{fl/fl}$, $Foxc2^{fl/fl}$, $Foxc1^{fl/fl};Foxc2^{fl/fl}$ [39], $Cdh5$-Cre^{ERT2} [40], $Vegfr3$-Cre^{ERT2} [45] and $Foxc2$-Cre^{ERT2}

[33] mice were used. EC-specific or LEC-specific $Foxc1$, $Foxc2$, and compound $Foxc1;Foxc2$

mutant mice were generated by crossing $Foxc$-floxed females ($Foxc1^{fl/fl}$, $Foxc2^{fl/fl}$, and

$Foxc1^{fl/fl};Foxc2^{fl/fl}$) with $Cdh5$-$Cre^{ERT2};Foxc1^{fl/fl}$ (EC-$Foxc1$-KO), $Cdh5$-$Cre^{ERT2};Foxc2^{fl/fl}$ (EC-

$Foxc2$-KO), $Cdh5$-$Cre^{ERT2};Foxc1^{fl/fl};Foxc2^{fl/fl}$ (EC-$Foxc$-DKO), $Vegfr3$-$Cre^{ERT2};Foxc1^{fl/fl}$ (LEC-

$Foxc1$-KO), $Vegfr3$-$Cre^{ERT2};Foxc2^{fl/fl}$ (LEC-$Foxc2$-KO), $Vegfr3$-$Cre^{ERT2};Foxc1^{fl/fl};Foxc2^{fl/fl}$

(LEC-$Foxc$-DKO) males, respectively, as described previously [41]. For Cre recombination

efficiency detection, $mTmG/+;Cdh5$-$Cre^{ERT2};Foxc1^{fl/fl};Foxc2^{fl/fl}$ (mTmG/EC-$Foxc$-DKO),

$mTmG/+;Vegfr3$-$Cre^{ERT2};Foxc1^{fl/fl};Foxc2^{fl/fl}$ (mTmG/LEC-$Foxc$-DKO) and $mTmG/+;Foxc2$-

Cre^{ERT2} mice were generated by crossing mTmG females ($mTmG/mTmG;Foxc1^{fl/fl};Foxc2^{fl/fl}$ and

$mTmG/mTmG$) with EC-$Foxc$-DKO, LEC-$Foxc$-DKO and $Foxc2$-Cre^{ERT2} males, respectively.

Genotyping of mice was performed by Transnetyx Inc.

Tamoxifen treatment

For adult mice, Tamoxifen (Tm, Cayman Chemical #13258) was dissolved in corn oil (Sigma

#C8267) at 40 mg/mL by shaking at 37°C for 3-4 hours. 7–8-week-old male adult mice were

treated with 150mg/kg Tm by oral gavage once daily for 5 consecutive days. For neonatal mice,

each individual was treated with Tm (20 mg/mL, 75 μg) by oral gavage once daily from

postnatal day 1 (P1) to day 5 (P5) [41].

Cre recombination efficiency detection

mTmG/EC-*Foxc*-DKO, mTmG/LEC-*Foxc*-DKO and *mTmG/+;Foxc2-Cre^{ERT2}* mice were used.

The Cre negative mTmG mice were used as control. Twelve days after Tm treatment, the distal

jejunum was harvested and fixed in 4% paraformaldehyde (PFA), followed by dehydration in 30%

sucrose, and OCT (Sakura Finetek, USA) embedding. 10 or 15 μm cryosections were cut and

immunostained with CD31 or LYVE1 antibody (Supplementary Table 2), counterstained

with/without GFP antibody and a nuclear-specific dye DAPI. EGFP fluorescent signal was

detected by imaging to evaluate the Cre recombination efficiency.

Mouse small intestinal ischemia and reperfusion (I/R) surgery

Twelve days after Tm treatment, mice were subjected to small intestinal ischemia and

reperfusion (I/R) surgery as previously described [31]. Briefly, mice were anaesthetized with

inhalation of Isoflurane. A midline laparotomy was performed, and the superior mesenteric

artery (SMA) was then identified, isolated, and clamped by a small nontraumatic vascular clip.

Heparinized-saline (500 μL, 10 U/ml) was added into the peritoneal cavity via syringe to avoid

coagulation of blood. After this ischemic phase for 30 min, the clip was removed, and the

intestine was allowed to reperfuse. 500 μL of sterile saline is administered to the peritoneal

cavity to compensate for the fluid loss during surgery. A Chromic Gut Suture (4–0) was then

used to close the muscle layer, followed by the skin closure with wound clips. Sham-operated

mice were subjected to the exact same surgical procedure, aside from clip placement.

BrdU treatment

Mice were treated with one dose of BrdU (Sigma #B5002, 10 mg/mL in PBS, 5 mg/kg) by

intraperitoneal (i.p.) injection 2 h or 18.5h (for evaluation of proliferative intestinal epithelium or

proliferative BECs and LECs, respectively) before tissue dissection (at I/R-24h and I/R-18.5h

respectively).

Tissue collection

Distal jejunum was selected for this study because this segment has the most severe mucosal

damage after intestinal I/R surgery compared to other segments according to pilot experiments.

Different time points (3 h, 4 h for an early injury stage, 18.5 h, 24 h and 48 h for a late repair

stage) were chosen for different analysis purposes. For histological analysis, transcardial

perfusion was performed on the adult mice with cold PBS followed by 4% PFA after anesthesia.

Distal jejunum was harvested and cut longitudinally to expose the lumen. After several washes

with PBS, the intestine was post-fixed in 4% PFA at 4°C for 4 h (for frozen or whole-mount

samples) or for O/N (for paraffin embedded samples). For qPCR and Western blot on whole

tissue lysates of the small intestine, blood was removed by transcardial perfusion with cold PBS.

Distal jejunum was harvested, opened longitudinally, and washed with cold PBS, then snap-

frozen in liquid nitrogen for the RNA isolation and protein extraction. For the tissue collection

for neonatal intestinal whole-mount staining, the neonates were euthanized at P7 after Tm

treatment from P1 to P5. The proximal jejuna were collected, washed with cold PBS, and fixed

in 4% PFA at 4°C for 4 h, then subjected to the whole-mount staining protocol. Proximal

jejunum was collected from neonatal mouse due to the ease of operation and similar lacteal

length/blood capillary network length ratio between proximal and distal jejuna [65].

Histopathological evaluation of intestinal mucosal damage

Paraffin sections of distal jejuna from the mice 24 h after intestinal I/R were stained with

Hematoxylin and Eosin (H&E). Based on the H&E staining, Chiu Score [42] was used for

evaluating the intestinal mucosal damage after I/R: grade 0, normal mucosa; grade 1,

development of subepithelial Gruenhagen's space at the apex of the villus; grade 2, extension of

the space with moderate epithelial lifting from the lamina propria; grade 3, massive epithelial

lifting with a few denuded villi; grade 4, denuded villi with exposed lamina propria and dilated

capillaries; and grade 5, digestion and disintegration of the lamina propria, hemorrhage and

ulceration. Higher scores represent more severe damage.

Whole-mount (WM) staining

Whole-mount staining of small intestine was performed as previously described [66]. Briefly,

Distal jejunum was dissected, and fixed in fixative (0.5% PFA, 15% picric acid and 100 mM

phosphate buffer, pH 7.0) at 4°C for O/N. Samples were washed with PBS, and subsequently

dehydrated by 10% sucrose for 3 h, and 20% sucrose+10% glycerol in PBS for O/N at 4°C. After

PBS-wash, samples were incubated with blocking buffer (5% donkey serum, 0.5% BSA, 0.3%

Triton X-100, 0.1% NaN_3 in PBS) for 2 h at 4°C, and then incubated with the indicated primary

antibodies (Supplementary Table 2) diluted in the blocking buffer for O/N at 4°C. Samples were

washed with PBST (0.3% Triton X-100 in PBS) for several times, followed by incubation with

indicated fluorochrome-conjugated secondary antibodies (Supplementary Table 2) diluted in the

blocking buffer for O/N at 4°C. The samples were washed again with PBST, post-fixed with 4%

PFA, cut into thin strips (one or two villi wide), cleared with FocusClear (CelExplorer Labs

#FC-101) and mounted on slides in mounting medium.

Immunohistochemistry (IHC) staining

For IHC-P, 4 or 15 μm paraffin sections were deparaffinized, rehydrated, subjected to antigen

retrieval, permeabilized with PBST, blocked with blocking buffer containing 5% Donkey serum

in PBST for 30 min at room temperature (RT) and incubated with indicated antibodies

(Supplementary Table 2) in blocking solution for O/N at 4°C. The sections were washed with

PBS and incubated with indicated fluorochrome-conjugated secondary antibodies

(Supplementary Table 2) in PBS for 1 h at RT. After wash with PBS, the sections were

counterstained with DAPI and mounted with mounting medium. For IHC-F, 10 or 15 μm frozen

sections were washed with PBS, and then subjected to the same blocking and antibody

incubation protocols as IHC-P but without the antigen retrieval step. TUNEL staining was

performed using *In Situ* Cell Death Detection Kit (Roche #11684795910) according to the

manufacturer's instructions.

Imaging

H&E staining images were acquired using an Olympus Vanox AHBT3 Research Microscope

(original magnification 100x, Tokyo, Japan). Fluorescent images were acquired using a Zeiss

AxioVision fluorescence microscope or a Nikon A1 Confocal Laser Microscope with the

software of Zeiss AxioVision SE64 Rel. 4.9.1 or NIS-Elements Viewer 4.20, respectively.

Images were processed and analyzed with Adobe Photoshop, Imaris and Fiji (ImageJ) software.

Imaris imaging software was used to create videos (Supplementary Videos, 1 and 2) for the 3-D

structure of the blood and lymphatic vasculatures in villi after intestinal I/R.

Rescue experiments

In rescue experiments, mice were treated with 5 µg RSPO3 (R&D systems #4120RS025CF) in

100 µL PBS [18], or CXCL12α (PeproTech #250-20A) in PBS at a dose of 50µg/kg of body weight

(BW) [49] by retro-orbital injection 30 min before intestinal ischemia. Distal jejunum was

harvested after I/R at 18.5 h (WB) or 24 h (H&E, IHC) for further analysis.

Endothelial cell isolation from small intestine

ECs were isolated from the distal jejunum for further qPCR analysis. Briefly, the tissue was

processed upon dissociating into single cell suspension in a digestion buffer (2 mg/mL

Collagenase D, 0.2mg/mL DNAase I, 2 mg/mL Dispase II, 100 units/mL penicillin and 100

µg/mL streptomycin in HBSS) for 35 min at 37°C, followed by filtration through a 70µm and a

40µm cell strainer. After several washes, the cell suspension was then incubated with magnetic

Dynabeads (Invitrogen #11035) labeled with CD45 antibody (Biolegend #103102) to deplete the

CD45+ cell population. The CD45- cell suspension was then incubated with Dynabeads labeled

with CD31 antibody (BD #553369) for the isolation of CD45-CD31+ cells. Finally, the CD45-

CD31+ ECs were collected for RNA isolation and qPCR analysis.

RNA isolation and qPCR analysis

A RNeasy Mini Kit (Qiagen #74104) was used for RNA extraction from isolated ECs. TRIzol™

Reagent (Invitrogen #15596026) was used to isolate RNA from whole distal jejunum. The

concentration of RNA was determined using NanoDrop™ 2000 Spectrophotometers (Thermo

Scientific). cDNA was synthesized using an iScript reverse transcriptase kit (Bio-Rad #170-

8891). qPCR was performed on triplicates of cDNA samples by using QuantStudio® 3 Real-

Time PCR System (Applied Biosystems), Fast SYBR reaction mix (Applied Biosystems), and

gene specific primer sets. 18S was used as an internal standard for mRNA expression. Primer

sequences are provided in Supplementary Table 3.

Western Blot

The frozen intestinal tissue was grinned using mortar and pestle chilled with liquid nitrogen,

followed by lysis in RIPA buffer (150mM NaCl, 1% Nonidet P-40, 0.5% sodium deoxycholate,

0.1% SDS and 50mM Tris, pH 7.4) containing protease inhibitors (Roche #4693116001). After

centrifugation, the supernatant of the tissue lysates was collected and mixed with 5x Protein

Loading Buffer. Equal amount of total protein for each sample was loaded and run on an SDS-

PAGE gel. Samples were transferred to 0.45 µm nitrocellulose (Invitrogen) and Western blotted

with the antibodies listed in Supplementary Table 2, followed by the reaction with ECL

substrates. The chemiluminescent signal was then detected by imaging the blot with Azure c600

imaging system. Bands were quantified using ImageJ software.

Preparation of single cell suspension from mouse small intestine

For single-cell RNA sequencing, mouse distal jejunums were collected 18.5h after intestinal I/R

surgery. Two mice were used for each group: Control I/R-18.5h and EC-*Foxc*-DKO I/R-18.5h.

Briefly, mice were anesthetized by isoflurane. Blood was removed by cardiac perfusion with

cold PBS. The distal jejunum was then dissected, washed with cold PBS and cut into small

pieces. The tissue was processed for scRNA-seq upon dissociating into single cell suspension in

a digestion buffer as mentioned above for 35 min at 37°C, followed by filtration through a 70µm

and a 40µm cell strainer. Cells were washed with washing buffer (0.5% BSA, 2mM EDTA in

PBS, pH 7.4) for three times, and resuspended in PBS with 0.04% BSA at a concentration of

1200 cells/µL (according to 10x Genomics Document #CG00053 Rev B) before being passed

through a 30 µm MACS SmartStrainer. The cell viability was tested by using the Cellometer

Auto 2000 Cell Viability Counter (Nexcelom Bioscience, USA). The cell sample was processed

for scRNA-seq only when the cell viability was more than 70%. Average cell viability for

samples was determined to be 80.96%.

Single cell 3' gene expression library construction and sequencing

Single cell 3' gene expression libraries were constructed by using the Chromium Next GEM

Single Cell 3' Reagent Kits v3.1 (10x Genomics, Pleasanton, CA, USA) according to the

manufacturer's manual CG000204 Rev D. The single cell libraries were assessed for quality

(TapeStation 4200, Agilent, Santa Clara, CA, USA) and then run by using paired-end 50 bp

sequencing on the Illumina HiSeq 4000 platform (Illumina, San Diego, CA, USA). 10,000 cells

were targeted for each sample with a sequencing depth of 20,000 read pairs per cell.

Pre-processing of single-cell RNA data

Following library generation and sequencing, raw sequencing data were de-multiplexed and

mapped to the mouse reference genome (mm10) using the CellRanger toolkit (10X Genomics,

version 2.1.0). Gene expression matrices were then generated from both Control and EC-*Foxc*-

DKO mice using CellRanger. The matrix files were then utilized for data processing and

downstream analysis using the BIOMEX browser-based software platform and its incorporated

packages developed in R [67]. Quality control and data pretreatment was performed in BIOMEX

with the following manually set parameters: i) genes with a row average of <.001 were excluded

for downstream analysis and ii) cells in which over 10% of unique molecular identifiers (UMIs)

were derived from the mitochondrial genome were considered as dead cells and removed from

downstream analysis. The data were then normalized in BIOMEX using similar methodology to

the *NormalizeData* function as implemented in the *Seurat* package [68].

Variable gene identification, dimensionality reduction, clustering analysis, and differential

gene expression analysis

Following data pretreatment, BIOMEX was utilized for downstream dimensionality reduction of

data and clustering analysis using the incorporated R packages. First, highly variable genes were

identified utilizing the following feature selections: mean lower threshold = 0.01, mean higher

threshold = 8, dispersion threshold = 0.5. Data (using highly variable genes only) was then auto-

scaled and summarized by principal component analysis (PCA), followed by visualization using

Uniform Manifold Approximation and Projection (UMAP; top 15 principal components (PCs))

to reduce the data into a two-dimensional space. Graph-based clustering was then performed in

BIOMEX to cluster cells according to their respective gene expression profile using

methodology similar to the *FindClusters* function in *Seurat* (clustering resolution = 0.5, k-

nearest neighbors = 15). Marker set analysis was then performed in BIOMEX on highly variable

genes to identify the top 10 gene markers expressed in each initial cluster using similar

methodology described previously [69]. Marker genes were then compared with single cell RNA-

seq data from the small intestine of adult mice available from the Mouse Cell Atlas (MCA,

http://bis.zju.edu.cn/MCA) and previously reported data [70] to identify transcriptionally unique

cell populations. Clusters with highly similar expression patterns indicative of the same cell

phenotype were merged into the same cluster. Differential gene expression analysis between

Control and EC-*Foxc*-DKO mice for individual cell clusters was performed in BIOMEX using

the Model-based Analysis of Single-cell Transcriptomics (MAST) package [71]. *P*-values <.05

were considered statistically significant for differentially expressed genes.

Data visualization

BIOMEX implementation of the *Plotly* software was used for UMAP and volcano plot

visualization.

Forkhead box C transcription factor binding prediction analysis in *RSPO3* and *CXCL12*

loci

Putative FOX-binding sites in the *RSPO3* and *CXCL12* loci were first identified using the

Hypergeometric Optimization of Motif EnRichment (HOMER) [52] suite of tools to scan the entire

Genome Reference Consortium Human Build 37 (GRCh37 or hg19) genome corresponding to

the conserved RYMAAYA FOX transcription factor binding motif or the reported RYACACA

FOXC transcription factor binding motif [72]. The output files were then uploaded to the UCSC

genome browser [51] to identify putative binding sites corresponding to transcriptionally active

areas denoted by histone modification, DNAse sensitivity, and additional transcription factor

chromatin immunoprecipitation data as per work reported and summarized on the Encyclopedia

of DNA Elements (ENCODE; https://genome.ucsc.edu/ENCODE/)(2012). Putative sites in the

human genome were then searched against the Genome Reference Consortium Mouse Build 38

(mm10)   genome   using   the   Evolutionary   Conserved   Region   (ECR)   Browser

(https://ecrbrowser.dcode.org) and rVista 2.0 tools to identify conserved and aligned putative

binding sites between mouse and human sequences. Conserved and aligned putative FOX

binding site sequences are underlined and bolded within the human *RSPO3* and *CXCL12* ECRs

as shown in the Supplementary Methods.

Mouse NEC model

24-hour-old mice were submitted to a well characterized NEC model [35] conducted in a 33-35°C

infant incubator or left with the dam. The NEC protocol includes: 1) initial orogastric inoculation

with a standardized adult mouse commensal bacteria preparation (10^8 colony-forming units) and

LPS (5 mg/kg) to perturb the normal intestinal colonization process; 2) gavage with formula

every 3 h (Esbilac, 200 mL/kg/day); and 3) exposure to brief episodes of hypoxia (60 s in 100%

N_2) followed immediately by cold stress (10 min at 4°C) twice daily. This protocol induces

intestinal injuries ranging from epithelial injury to transmural necrosis resembling human NEC

which typically develop after 36 hours and has been widely used to study NEC pathogenesis [73].

Pups were euthanized by decapitation at 24 hours into NEC and whole intestinal tissues were

collected and fixed in formalin for 24 hours before tissue processing and paraffin block

preparation.

Quantification

For quantification of BrdU+ or CCND1+ cells in crypts, images from different fields of section

under a 20x objective were acquired and about 15~50 crypts were analyzed for each sample. The

number of BrdU+ or CCND1+ cells was counted for each crypt. For quantification of β-catenin

staining, about 5~16 ISCs (OLFM4+) per sample were analyzed for the fluorescent intensity of

β-catenin within the ISC by ImageJ. For analysis of BEC and LEC proliferation and apoptosis,

confocal Z-stacks were acquired using a 20x objective from about 4~ 8 different fields of 15 µm

paraffin sections for each sample. Area of blood vessels (CD31+LYVE1-) and lymphatic vessels

(CD31+LYVE1+) were measured using ImageJ software. Then the number of BrdU+ or

TUNEL+ cells per $0.1mm^2$ vessel areas was calculated and compared between groups.

Measurements for the length of blood capillaries and lacteals was performed as previously

described [66]. The numbers of branches and branching points for the villous blood vasculature

was calculated based on the whole-mount staining of VEGFR2 as previously described [66].

Around 30~50 villi were analyzed for each sample. For CD31+ vessel density quantification,

5~10 images were acquired using a 20x objective and the CD31+ vessel area and the intestinal

tissue area in the images were measured using ImageJ software. CD31+ vessel density

percentage was determined as the total CD31+ vessel area/total intestinal tissue area x 100%.

Statistics

For quantification, statistical analysis was performed using GraphPad Prism 8.0 (GraphPad

Software). *P* values were obtained by performing Mann-Whitney *U* test or Kruskal-Wallis One-

way ANOVA test. Data are presented as box-and-whisker plots of representative experiments

from at least three biological replicates. *P* values <0.05 were considered statistically significant.

For scRNA-seq data, differential gene expression analysis between groups for individual cell

clusters was performed in BIOMEX using the Model-based Analysis of Single-cell

Transcriptomics (MAST) package [71]. Adjusted *P* values <0.05 were considered statistically

significant for differentially expressed genes.

Study approval

All experimental protocols and procedures used in this study were approved by the Institutional

Animal Care and Use Committee (IACUC) at Northwestern University.

727
728

729 **References**

730 1.   Rafii S, Butler JM, Ding BS. Angiocrine functions of organ-specific endothelial cells.
731   *Nature* **529**, 316-325 (2016).
732

733 2.   Bertoni S, Ballabeni V, Barocelli E, Tognolini M. Mesenteric ischemia-reperfusion: an
734   overview of preclinical drug strategies. *Drug Discov Today* **23**, 1416-1425 (2018).
735

736 3.   Gonzalez LM, Moeser AJ, Blikslager AT. Animal models of ischemia-reperfusion-
737   induced intestinal injury: progress and promise for translational research. *Am J Physiol*
738   *Gastrointest Liver Physiol* **308**, G63-75 (2015).
739

740 4.   Bowker RM, Yan X, De Plaen IG. Intestinal microcirculation and necrotizing
741   enterocolitis: The vascular endothelial growth factor system. *Semin Fetal Neonatal Med*
742   **23**, 411-415 (2018).
743

744 5.   Barker N. Adult intestinal stem cells: critical drivers of epithelial homeostasis and
745   regeneration. *Nat Rev Mol Cell Biol* **15**, 19-33 (2014).
746

747 6.   Barker N*, et al.* Identification of stem cells in small intestine and colon by marker gene
748   Lgr5. *Nature* **449**, 1003-1007 (2007).
749

750 7.   Gonzalez LM*, et al.* Preservation of reserve intestinal epithelial stem cells following
751   severe ischemic injury. *Am J Physiol Gastrointest Liver Physiol* **316**, G482-G494 (2019).
752

753 8.   Richmond CA*, et al.* JAK/STAT-1 Signaling Is Required for Reserve Intestinal Stem
754   Cell Activation during Intestinal Regeneration Following Acute Inflammation. *Stem Cell*
755   *Reports* **10**, 17-26 (2018).
756

757 9.   Murata K*, et al.* Ascl2-Dependent Cell Dedifferentiation Drives Regeneration of Ablated
758   Intestinal Stem Cells. *Cell stem cell* **26**, 377-390 e376 (2020).
759

760 10.   Hageman JH, Heinz MC, Kretzschmar K, van der Vaart J, Clevers H, Snippert HJG.
761   Intestinal Regeneration: Regulation by the Microenvironment. *Developmental cell* **54**,
762   435-446 (2020).
763

764 11.   Santos AJM, Lo YH, Mah AT, Kuo CJ. The Intestinal Stem Cell Niche: Homeostasis and
765   Adaptations. *Trends Cell Biol* **28**, 1062-1078 (2018).
766

767 12.   McCarthy N, Kraiczy J, Shivdasani RA. Cellular and molecular architecture of the
768   intestinal stem cell niche. *Nature cell biology* **22**, 1033-1041 (2020).
769

770 13.   Shoshkes-Carmel M*, et al.* Subepithelial telocytes are an important source of Wnts that
771   supports intestinal crypts. *Nature* **557**, 242-246 (2018).
772

773 14. McCarthy N, *et al.* Distinct Mesenchymal Cell Populations Generate the Essential
774 Intestinal BMP Signaling Gradient. *Cell stem cell* **26**, 391-402 e395 (2020).
775
776 15. Farin HF, Van Es JH, Clevers H. Redundant sources of Wnt regulate intestinal stem cells
777 and promote formation of Paneth cells. *Gastroenterology* **143**, 1518-1529 e1517 (2012).
778
779 16. Valenta T, *et al.* Wnt Ligands Secreted by Subepithelial Mesenchymal Cells Are
780 Essential for the Survival of Intestinal Stem Cells and Gut Homeostasis. *Cell reports* **15**,
781 911-918 (2016).
782
783 17. Storm EE, *et al.* Targeting PTPRK-RSPO3 colon tumours promotes differentiation and
784 loss of stem-cell function. *Nature* **529**, 97-100 (2016).
785
786 18. Kannan L, *et al.* R-spondin3 prevents mesenteric ischemia/reperfusion-induced tissue
787 damage by tightening endothelium and preventing vascular leakage. *Proceedings of the*
788 *National Academy of Sciences of the United States of America* **110**, 14348-14353 (2013).
789
790 19. Ogasawara R, *et al.* Intestinal Lymphatic Endothelial Cells Produce R-Spondin3.
791 *Scientific reports* **8**, 10719 (2018).
792
793 20. García-Cuesta EM, Santiago CA, Vallejo-Díaz J, Juarranz Y, Rodríguez-Frade JM,
794 Mellado M. The Role of the CXCL12/CXCR4/ACKR3 Axis in Autoimmune Diseases.
795 *Front Endocrinol (Lausanne)* **10**, 585 (2019).
796
797 21. Herbert SP, Stainier DY. Molecular control of endothelial cell behaviour during blood
798 vessel morphogenesis. *Nat Rev Mol Cell Biol* **12**, 551-564 (2011).
799
800 22. Hitchon C, Wong K, Ma G, Reed J, Lyttle D, El-Gabalawy H. Hypoxia-induced
801 production of stromal cell-derived factor 1 (CXCL12) and vascular endothelial growth
802 factor by synovial fibroblasts. *Arthritis Rheum* **46**, 2587-2597 (2002).
803
804 23. Santiago B, *et al.* CXCL12 gene expression is upregulated by hypoxia and growth arrest
805 but not by inflammatory cytokines in rheumatoid synovial fibroblasts. *Cytokine* **53**, 184-
806 190 (2011).
807
808 24. Santagata S, *et al.* CXCR4 and CXCR7 Signaling Pathways: A Focus on the Cross-Talk
809 Between Cancer Cells and Tumor Microenvironment. *Front Oncol* **11**, 591386 (2021).
810
811 25. Chang P, *et al.* Mesenchymal stem cells over-expressing cxcl12 enhance the
812 radioresistance of the small intestine. *Cell death & disease* **9**, 154 (2018).
813
814 26. Zimmerman NP, Vongsa RA, Faherty SL, Salzman NH, Dwinell MB. Targeted intestinal
815 epithelial deletion of the chemokine receptor CXCR4 reveals important roles for
816 extracellular-regulated kinase-1/2 in restitution. *Lab Invest* **91**, 1040-1055 (2011).
817

818 27. Golson ML, Kaestner KH. Fox transcription factors: from development to disease.
819 *Development* **143**, 4558-4570 (2016).
820

821 28. Seifi M, Walter MA. Axenfeld-Rieger syndrome. *Clinical genetics* **93**, 1123-1130 (2018).
822

823 29. Fang J*, et al.* Mutations in FOXC2 (MFH-1), a forkhead family transcription factor, are
824 responsible for the hereditary lymphedema-distichiasis syndrome. *American journal of*
825 *human genetics* **67**, 1382-1388 (2000).
826

827 30. Becker F*, et al.* Downregulation of FoxC2 Increased Susceptibility to Experimental
828 Colitis: Influence of Lymphatic Drainage Function? *Inflammatory bowel diseases* **21**,
829 1282-1296 (2015).
830

831 31. Yoshiya K*, et al.* Depletion of gut commensal bacteria attenuates intestinal
832 ischemia/reperfusion injury. *Am J Physiol Gastrointest Liver Physiol* **301**, G1020-1030
833 (2011).
834

835 32. Hader C, Marlier A, Cantley L. Mesenchymal-epithelial transition in epithelial response
836 to injury: the role of Foxc2. *Oncogene* **29**, 1031-1040 (2010).
837

838 33. Amin MB*, et al.* Foxc2CreERT2 knock-in mice mark stage-specific Foxc2-expressing
839 cells during mouse organogenesis. *Congenital anomalies* **57**, 24-31 (2017).
840

841 34. Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. A global double-fluorescent Cre
842 reporter mouse. *Genesis* **45**, 593-605 (2007).
843

844 35. Tian R*, et al.* Characterization of a necrotizing enterocolitis model in newborn mice. *Int J*
845 *Clin Exp Med* **3**, 293-302 (2010).
846

847 36. Kume T, Deng KY, Winfrey V, Gould DB, Walter MA, Hogan BL. The forkhead/winged
848 helix gene Mf1 is disrupted in the pleiotropic mouse mutation congenital hydrocephalus.
849 *Cell* **93**, 985-996 (1998).
850

851 37. Kume T, Jiang H, Topczewska JM, Hogan BL. The murine winged helix transcription
852 factors, Foxc1 and Foxc2, are both required for cardiovascular development and
853 somitogenesis. *Genes & development* **15**, 2470-2482 (2001).
854

855 38. Seo S, Fujita H, Nakano A, Kang M, Duarte A, Kume T. The forkhead transcription
856 factors, Foxc1 and Foxc2, are required for arterial specification and lymphatic sprouting
857 during vascular development. *Developmental biology* **294**, 458-470 (2006).
858

859 39. Sasman A*, et al.* Generation of conditional alleles for Foxc1 and Foxc2 in mice. *Genesis*
860 **50**, 766-774 (2012).
861

40. Sorensen I, Adams RH, Gossler A. DLL1-mediated Notch activation regulates
endothelial identity in mouse fetal arteries. *Blood* **113**, 5680-5688 (2009).

41. Norden PR, *et al.* Shear stimulation of FOXC1 and FOXC2 differentially regulates
cytoskeletal activity during lymphatic valve maturation. *eLife* **9**, (2020).

42. Chiu CJ, McArdle AH, Brown R, Scott HJ, Gurd FN. Intestinal mucosal lesion in low-
flow states. I. A morphological, hemodynamic, and metabolic reappraisal. *Arch Surg* **101**,
478-483 (1970).

43. Cerqueira NF, Hussni CA, Yoshida WB. Pathophysiology of mesenteric
ischemia/reperfusion: a review. *Acta Cir Bras* **20**, 336-343 (2005).

44. van der Flier LG, Haegebarth A, Stange DE, van de Wetering M, Clevers H. OLFM4 is a
robust marker for stem cells in human intestine and marks a subset of colorectal cancer
cells. *Gastroenterology* **137**, 15-17 (2009).

45. Martinez-Corral I, *et al.* Vegfr3-CreER (T2) mouse, a new genetic tool for targeting the
lymphatic system. *Angiogenesis* **19**, 433-445 (2016).

46. Meng FW, Shimoda H, Kajiwara T, Matsuda M, Kato S. Reconstruction of central
lacteals in the murine jejunum following ischemia-reperfusion injury. *Archives of*
histology and cytology **70**, 135-146 (2007).

47. Lee NV, *et al.* ADAMTS1 mediates the release of antiangiogenic polypeptides from
TSP1 and 2. *The EMBO journal* **25**, 5270-5283 (2006).

48. Obika M, *et al.* Tumor growth inhibitory effect of ADAMTS1 is accompanied by the
inhibition of tumor angiogenesis. *Cancer science* **103**, 1889-1897 (2012).

49. Garnica MR, de Moraes LV, Rizzo LV, de Andrade HF, Jr. Supplementation of CXCL12
(CXCL12) induces homing of CD11c+ dendritic cells to the spleen and enhances control
of Plasmodium berghei malaria in BALB/c mice. *Immunology* **115**, 399-406 (2005).

50. Gonzalez-Loyola A, *et al.* FOXC2 controls adult lymphatic endothelial specialization,
function, and gut lymphatic barrier preventing multiorgan failure. *Sci Adv* **7**, (2021).

51. Kent WJ, *et al.* The human genome browser at UCSC. *Genome research* **12**, 996-1006
(2002).

52. Heinz S, *et al.* Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-
589 (2010).

53. Kaufmann E, Knochel W. Five years on the wings of fork head. *Mechanisms of*
development **57**, 3-20. (1996).

54. Overdier DG, Porcella A, Costa RH. The DNA-binding specificity of the hepatocyte
nuclear factor 3/forkhead domain is influenced by amino-acid residues adjacent to the
recognition helix. *Molecular and cellular biology* **14**, 2755-2766. (1994).

55. Pierrou S, Hellqvist M, Samuelsson L, Enerback S, Carlsson P. Cloning and
characterization of seven human forkhead proteins: binding site specificity and DNA
bending. *The EMBO journal* **13**, 5002-5012. (1994).

56. Goto N, Imada I, Deshpande V, Yilmaz ÖH. Lymphatics constitute a novel component of
the intestinal stem cell niche. *bioRxiv* **https://doi.org/10.1101/2022.01.28.478205** (2022).

57. Niec RE, *et al.* A lymphatic-stem cell interactome regulates intestinal stem cell activity.
bioRxiv **https://doi.org/10.1101/2022.01.29.478341**, (2022).

58. Palikuqi B, Rispal J, Vaka D, Boffelli D, Klein O. Lymphangiocrine signals are required
for proper intestinal repair after cytotoxic injury. *bioRxiv*
https://doi.org/10.1101/2022.03.08.479995, (2022).

59. Tan C, *et al.* Endothelial FOXC1 and FOXC2 promote intestinal regeneration after
ischemia-reperfusion injury. *bioRxiv* **https://doi.org/10.1101/2022.03.03.482713** (2022).

60. Ribatti D, Tamma R, Annese T. The role of vascular niche and endothelial cells in
organogenesis and regeneration. *Experimental cell research* **398**, 112398 (2021).

61. Yi D, *et al.* Endothelial Autocrine Signaling through CXCL12/CXCR4/FoxM1 Axis
Contributes to Severe Pulmonary Arterial Hypertension. *Int J Mol Sci* **22**, (2021).

62. Omatsu Y, Seike M, Sugiyama T, Kume T, Nagasawa T. Foxc1 is a critical regulator of
haematopoietic stem/progenitor cell niche formation. *Nature* **508**, 536-540 (2014).

63. Xia W, *et al.* Overexpression of Foxc1 regenerates crushed rat facial nerves by promoting
Schwann cells migration via the Wnt/beta-catenin signaling pathway. *Journal of cellular*
physiology **235**, 9609-9622 (2020).

64. Stratman AN, Davis MJ, Davis GE. VEGF and FGF prime vascular tube morphogenesis
and sprouting directed by hematopoietic stem cell cytokines. *Blood* **117**, 3709-3719
(2011).

65. Bernier-Latmani J, *et al.* DLL4 promotes continuous adult intestinal lacteal regeneration
and dietary fat transport. *The Journal of clinical investigation* **125**, 4572-4586 (2015).

66. Bernier-Latmani J, Petrova TV. High-resolution 3D analysis of mouse small-intestinal
stroma. *Nature protocols* **11**, 1617-1629 (2016).

67. Taverna F, *et al.* BIOMEX: an interactive workflow for (single cell) omics data
interpretation and visualization. *Nucleic acids research* **48**, W385-W394 (2020).

68.   Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).

69.   Kalucka J*, et al.* Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**,
764-779.e720 (2020).

70.   Haber AL*, et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333-
339 (2017).

71.   Finak G*, et al.* MAST: a flexible statistical framework for assessing transcriptional
changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome*
biology **16**, 278 (2015).

72.   Chen X*, et al.* Structural basis for DNA recognition by FOXC2. *Nucleic acids research*
47, 3752-3764 (2019).

73.   Yan X*, et al.* Lack of VEGFR2 signaling causes maldevelopment of the intestinal
microvasculature and facilitates necrotizing enterocolitis in neonatal mice. *Am J Physiol*
Gastrointest Liver Physiol **310**, G716-725 (2016).

975
976

Acknowledgments

We thank William Muller (Northwestern University) and Bona Jabri (University of Chicago) for

helpful advice. *Cdh5-Cre^ERT2* mice were kindly provided by Dr. Ralf Adams at the Max-Planck-

Institute for Molecular Biomedicine, Germany. Single cell RNA-seq experiments were

performed at the NUSeq Core Facility at Northwestern University. **Funding:** This work was

supported by the NIH (R01HL126920 and R01HL144129 to TK). Imaging work was performed

at the Northwestern University Center for Advanced Microscopy generously supported by NCI

CCSG P30 CA060553 awarded to the Robert H Lurie Comprehensive Cancer Center. **Author**

Contributions: CT, PRN, NU, XY, IGD, and TK designed and analyzed the experiments. CT,

XY, TL, NU, and TL conducted the experiments. PRN analyzed scRNA-seq data. KA and SO

provided Cre mouse lines. CT, PRN, and TK wrote the manuscript. **Competing Interests: C.**

Tan, None; **P.R. Norden**, None; **T. Liu**, None; **N. Ujiie,** None; **X. Yan**, None; **K Aoto**, None; **S.**

Ortega, None; **I.G. De Plaen,** None; **T. Kume,** None. **Data and materials availability:** All

data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Materials. Additional data related to this paper may be requested from the

authors.

Figure 1

Figure 1. Expression levels of *Foxc1* and *Foxc2* in the mouse small intestine. Relative

expression level(s) of **(A)** *Foxc1* and *Foxc2* in sham intestines, **(B)** *Foxc1* and **(C)** *Foxc2* in sham

and I/R-4h intestines, **(D)** both *Foxc1* and *Foxc2* at I/R-4h, in the isolated CD45-CD31+ ECs

from *Foxc1^{fl/fl};Foxc2^{fl/fl}* mouse distal jejuna. Data are box-and-whisker plots, Mann-Whitney *U*

test, each symbol represents one mouse, N = 4~7, *P<0.05, **P<0.01, ***P<0.001. **(E)**

Representative confocal images of the whole-mount distal jejunum stained with

FOXC1/CD31/LYVE1 in *Foxc1^{fl/fl};Foxc2^{fl/fl}* mice show the up-regulation of FOXC1 in BECs (B)

and LECs (L) at the level of villus and submucosa 4 h after I/R. Scale bars = 20 μm. **(F)**

Schematic showing the time of Tamoxifen (Tm) injection and tissue dissection for Figure 1, G-I.

Relative expression of *Foxc1* **(G)** and *Foxc2* **(H)** in isolated CD45-CD31+ ECs from distal

jejuna 12d after Tm treatment. Data are box-and-whisker plots, Mann-Whitney *U* test, each

symbol represents one mouse, N = 4~5, *$P<0.05$. **(I)** Representative confocal images of

submucosal blood vessels (B)/lymphatics (L) in whole-mount intestines stained with

FOXC1/CD31/LYVE1. In control, FOXC1 is expressed in the nuclei of the BECs, but hardly

detectable in LECs. FOXC1 is down-regulated in BECs after Tm treatment in EC-*Foxc*-DKO

mice compared with control. Scale bars = 50 μm. Note that the red tiny spots outside the

vasculatures are non-specific staining.

1031

 Figure 2

Figure 2. Characterization of defects in the intestinal mucosa in mice with EC-specific deletion of *Foxc1/2* after I/R. (**A**) Schematic showing the time of Tamoxifen (Tm) injection, I/R surgery and tissue collection (TC). (**B**) Representative H&E staining images of the distal jejuna in EC-*Foxc*-DKO and control mice 24h after I/R. The intestinal ischemic injury grading in the Chiu scoring system is indicated by red numbers (0~5). Scale bars = 100 μm. Quantification of Chiu Score for (**C**) Control and EC-*Foxc*-DKO, (**D**) Control and EC-*Foxc1*-KO, (**E**) Control and EC-*Foxc2*-KO groups at I/R-24h. Figure 2D and 2E are based on Supplementary Figure 5, A and B. Data are box-and-whisker plots, Mann-Whitney U test, each symbol represents one mouse, N = 3 in C sham groups, N = 7~12 in C, D, E I/R-24h groups, *P<0.05, ****P<0.0001. (**F**) Relative mRNA expression of proinflammatory mediators *Cox-2*, *TNF-α* and *IL-6* from

intestinal tissue lysates at I/R-3h. Data are box-and-whisker plots, Mann-Whitney U test, each

symbol represents one mouse, N = 6~11, *P<0.05, *n.s.* = not significant. **(G)** Representative

immunostaining images of intestinal crypts labeled with BrdU (proliferative marker, injection

performed 2h before tissue collection) and EpCAM (epithelial marker) show the proliferation of

epithelial cells in crypts. Paraffin sections (4 µm), scale bars = 20 μm. **(H)** Quantification of the

number of BrdU+ epithelial cells per crypt based on Figure 2G. Data are box-and-whisker plots,

Kruskal-Wallis One-way ANOVA test, each symbol represents one mouse, N = 3~6, *P<0.05,

n.s. = not significant.

1063

Figure 3

Figure 3. Wnt signaling in the small intestine is impaired in EC-*Foxc*-DKO mice after I/R injury. **(A)** Representative images of intestinal crypts immunostained with β-catenin and the intestinal epithelial stem cell (ISC) marker OLFM4. At I/R-48h the nuclear translocation of β-catenin in ISCs (dotted circles) was found in control, whereas it's seldom found in EC-*Foxc*-

DKO. Paraffin sections (4 µm), scale bars = 20 µm. **(B)** Quantification of relative fluorescent

intensity of β-catenin immunostaining within ISC based on Figure 3A. Data are box-and-whisker

plots, Mann-Whitney U test, each symbol represents one mouse, N = 4, *$P<0.05$. **(C)**

Representative images of intestinal mucosa labeled with Cyclin D1 (CCND1) in EC-*Foxc*-DKO

mice compared with Control group in sham, 24h and 48h after I/R. Paraffin sections (4 µm),

scale bars = 100 µm. **(D)** Quantification of the number of CCND1+ epithelial cells per crypt

based on Figure 3C. Data are box-and-whisker plots, Kruskal-Wallis One-way ANOVA test,

each symbol represents one mouse, N = 4~6. **$P<0.01$, ****$P<0.0001$, *n.s.*=not significant.

1095

 Figure 4

Figure 4. Characterization of defects in the intestinal mucosa in mice with LEC-specific
deletion of *Foxc1/2* after I/R. Chiu Score analysis from H&E stained distal jejunum 24h after
I/R for: (**A**) Control (*Foxc1^{f/f};Foxc2^{f/f}*) and LEC-*Foxc*-DKO (*Vegfr3-Cre^{ERT2};Foxc1^{f/f};Foxc2^{f/f}*),
(**B**) Control (*Foxc1^{f/f}*) and LEC-*Foxc1*-KO (*Vegfr3-Cre^{ERT2};Foxc1^{f/f}*), (**C**) Control (*Foxc2^{f/f}*) and
LEC-*Foxc2*-KO (*Vegfr3-Cre^{ERT2};Foxc2^{f/f}*) based on Supplementary Figure 5 C-E. Data are box-

and-whisker plots, Mann-Whitney U test, each symbol represents one mouse, N=9~13, *P<0.05,

P<0.01. **(D) Quantification of the number of BrdU+ epithelial cells per crypt in control and

LEC-*Foxc*-DKO mice at 24h after I/R. Data are box-and-whisker plots, Mann-Whitney U test,

each symbol represents one mouse, N = 5~6. **P<0.01. **(E)** Representative images of crypts

immunostained with OLFM4 and β-catenin in control and LEC-*Foxc*-DKO mice 24h after I/R.

The accumulation of β-catenin in the nuclei of ISCs (dotted circles) was found in Control but

inhibited in LEC-Foxc-DKO mice. Paraffin sections (4 µm), scale bars = 20 μm. **(F)**

Quantification of relative fluorescent intensity of β-catenin immunostaining within ISC based on

Figure 4E. Data are box-and-whisker plots, Mann-Whitney U test, each symbol represents one

mouse, N = 9~10, *P<0.05. **(G)** Quantification of the number of CCND1+ epithelial cells per

crypt at I/R-24h from the intestinal immunostaining of CCND1 based on Supplementary Figure

6. Data are box-and-whisker plots, Mann-Whitney U test, each symbol represents one mouse, N

= 7~8. **P<0.01.

1132

Figure 5

Figure 5. Defects in vascular regeneration after intestinal I/R injury in EC-*Foxc*-DKO mouse.

(**A**) Representative images of intestinal whole-mount VEGFR2/CD31 (green/red) immunostaining show increased VEGFR2 (green) expression at the angiogenic front of villous blood capillaries in control mice at I/R-18.5h. In EC-*Foxc*-DKO, the increase of VEGFR2 is inhibited in villous blood vessels and the repair of blood vasculatures is also impaired (arrow).

1154   Scale bars = 200 μm. **(B)** Representative images of intestinal whole-mount VEGFR3/LYVE1

1155   (red/green) immunostaining. 18.5h after I/R, VEGFR3 (red) is increased in lacteals especially at

1156   the lacteal tips in the control but is inhibited in the EC-*Foxc*-DKO lacteals (arrow). Scale bars =

1157   200 μm.

1158

Figure 6

Figure 6. Impaired vascular regeneration after intestinal I/R injury in EC-*Foxc*-DKO mouse.

(**A**) Representative BrdU/CD31/LYVE1/DAPI immunostaining images of intestinal paraffin sections (15 μm) from mice injected with BrdU 18.5h before euthanasia for the analysis of proliferative BECs (B in 1, 3, 5) and LECs (L in 2, 4, 6) in intestines. Arrow heads show BrdU+ BECs or LECs. White/yellow bars = 100 or 10 μm, respectively. (**B**) The numbers of BrdU+

BECs and LECs per 0.1 mm^2 blood vessel (CD31+LYVE1-) and lymphatic vessel

(CD31+LYVE1+) area in intestinal mucosa were quantified respectively based on Figure 6A.

Data are box-and-whisker plots, Kruskal-Wallis One-way ANOVA test, each symbol represents

one mouse, N = 3~8. **(C)** Representative TUNEL/CD31/LYVE1/DAPI immunostaining images

of distal jejunums for the analysis of apoptotic BECs/LECs in intestinal paraffin sections (15

µm). High magnification images are from the dotted line boxes (1~4). Arrow heads show the

apoptotic BECs (B in 1, 2) and LECs (L in 3, 4) at villus (1, 3) or submucosa (2, 4).

White/yellow scale bars = 100 or 20 µm, respectively. **(D)** The numbers of apoptotic BECs and

LECs per 0.1 mm^2 blood vessel (CD31+LYVE1-) and lymphatic vessel (CD31+LYVE1+) area

in intestinal mucosa were quantified respectively based on Figure 6C. Data are box-and-whisker

plots, Mann Whitney *U* test, each symbol represents one mouse, N = 3~6. The length of blood

capillary vasculature and lacteals were measured **(E)** based on Supplementary Figure 7, A and B.

The percentage (%) of the lacteal length/blood capillary network length was then calculated **(F)**.

The numbers of branches **(G)** and the branching points **(H)** of the villous blood vasculatures

were counted based on the whole-mount staining of VEGFR2 (Figure 5A) as previously

described [66] using ImageJ software. Data are box-and-whisker plots, Mann-Whitney U test, each

symbol represents one mouse, N = 3~5. *$P<0.05$, **$P<0.01$. *n.s.* = not significant.

1197

 Figure 7

Figure 7. Single-cell RNA sequencing characterization of the distal jejunum from Control and EC-*Foxc*-DKO mice after I/R at 18.5h. (A) Visualization of unsupervised clustering of 22 distinct clusters by UMAP from the distal jejunum of both control and EC-*Foxc*-DKO mice after I/R at 18.5h. **(B)** UMAP visualization of *Rspo3* expression in LECs (12), Myocytes/Pericytes (17), and Telocytes/Trophocytes (20) cell clusters identified in Figure 7A. **(C)** Dot plot showing

relative expression of *Rspo3* in 5 cell clusters identified by scRNA-seq. Fill colors represent

normalized mean expression levels and circle sizes represent the within-cluster frequency of

positive gene detection. **(D)** Representative images of lymphatic vessels (surrounded by the

white dotted lines) in intestinal mucosa labeled with RSPO3/LYVE1. Scale bars = 10 µm. **(E)**

Volcano plots of differential expression analysis of BECs in Control and EC-*Foxc*-DKO mice at

I/R-18.5h using the MAST model. Blue dots denote genes with significant differential expression,

P < 0.05. **(F)** UMAP visualization of *Cxcl12* expression in BECs (1), fibroblasts (9) and

Telocytes/Trophocytes (20) cell clusters identified in Figure 7A **(G)** Dot plot showing relative

expression of *Cxcl12* in 5 cell clusters identified by scRNA-seq. **(H)** Relative mRNA expression

of *Cxcl12* in Dynabeads-isolated ECs (CD45-CD31+) from distal jejunum. Data are box-and-

whisker plots, Mann-Whitney U test, each symbol represents one mouse, N = 6~10, *$P<0.05$.

1232

 Figure 8

Figure 8. RSPO3 partially rescues impaired regeneration of intestinal mucosa in EC-*Foxc*-DKO and LEC-*Foxc*-DKO mice after I/R. In RSPO3 rescue experiment, each mouse was treated with 5 μg RSPO3 in 100μL PBS by retro-orbital injection 30 min before ischemia. PBS treated mice were used as control. Representative images of H&E staining show the rescue

effects of RSPO3 in intestinal mucosa in EC-*Foxc*-DKO **(A)** and LEC-*Foxc*-DKO **(C)** mouse

strains 24h after I/R. Red numbers indicate the Chiu scores. Scale bars = 100 μm. Quantification

of Chiu Score for RSPO3 rescued **(B)** EC-*Foxc*-DKO and **(D)** LEC-*Foxc*-DKO intestines 24h

after I/R. Data are box-and-whisker plots, Mann-Whitney U test, each symbol represents one

mouse, N = 7~13, ***P*<0.01. Representative Western blots (**E**, **G**) and densitometry

measurements (**F**, **H**) show IgM (heavy chain) in intestinal tissue lysates from control and EC-

Foxc-DKO mice in sham and after I/R at 18.5h (**E**, **F**) as well as IgM (heavy chain) in PBS- and

RSPO3- treated EC-*Foxc*-DKO mice after I/R at 18.5h (**G**, **H**). Data are box-and-whisker plots,

Mann-Whitney U test, each symbol represents one mouse, N = 4~7, **P*<0.05, ***P*<0.01.

n.s.=not significant.

1266

1267 **Figure 9**

Figure 9. CXCL12 partially rescues impaired regeneration of intestinal mucosa in EC-

***Foxc*-DKO mice after I/R.** In CXCL12 rescue experiments, mice were treated with 50μg/kg

CXCL12 in PBS by retro-orbital injection 30 min before ischemia. Mice treated with PBS were

used as control. **(A)** Representative images of H&E staining show the rescue effects of CXCL12

in intestinal mucosa in EC-*Foxc*-DKO mice 24h after I/R. Red numbers indicate Chiu scores.

Scale bars = 100 μm. **(B)** Quantification of Chiu Scores for CXCL12-rescued EC-*Foxc*-DKO

mice 24h after I/R based on Figure 9A. Data are box-and-whisker plots, Mann-Whitney U test,

each symbol represents one mouse, N = 10~11, *$P<0.05$. **(C)** Representative images of crypts

immunostained with OLFM4 and β-catenin in PBS- and CXCL12-treated EC-*Foxc*-DKO mice

24h after I/R. The total protein signal of β-catenin is up-regulated in CXCL12-treated crypts

compared with PBS-treated group. The accumulation of β-catenin in the nuclei of ISCs (dotted

circles) was found in CXCL12-treated mice but inhibited in PBS-treated mice. Paraffin sections

(4 µm), scale bars = 20 μm. **(D)** Quantification of relative fluorescent intensity of β-catenin

immunostaining within ISC based on Figure 9C. Data are box-and-whisker plots, Mann-Whitney

U test, each symbol represents one mouse, N = 5~7, *$P<0.05$. **(E)** Quantification of CCND1+

epithelial cells per crypt after I/R at 24h. Data are box-and-whisker plots, Mann-Whitney U test,

each symbol represents one mouse, N = 7~8. ***$P<0.001$. **(F)** Representative confocal images of

CD31 immunostaining of distal jejunums in PBS- and CXCL12- treated EC-*Foxc*-DKO mice

after I/R at 24h. Paraffin sections (15 µm), scale bars = 100 μm. **(G)** Quantification of CD31+

vessel density (% = total CD31+ vessel area/total intestinal tissue area x 100%) based on Figure

9F. Data are box-and-whisker plots, Mann-Whitney U test, each symbol represents one mouse, N

= 5~6. **$P<0.01$.

1307