It Takes Two: Bimodal interactions between the SARS-CoV-1 fusion peptide and Ca^{2+} ions promote membrane insertion and viral entry

Juliana Debrito Carten [1±], Miya K. Bidon [1±], George Khelashvili [4,5±], Marco R. Straus [2], Tiffany Tang [1], Javier A. Jaimes [2], Harel A. Weinstein [4,5*], Gary R. Whittaker [2,3*], and Susan Daniel [1*]

[1] Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University, Ithaca, NY, USA

[2] Departments of Microbiology & Immunology and [3] Public & Ecosystem Health, Cornell University, Ithaca, NY, USA

[4] Department of Physiology & Biophysics, Weill Cornell Medicine, New York, NY, USA

[5] Institute for Computational Biomedicine, Weill Cornell Medicine, New York, NY 10065, USA

*Corresponding authors:

Susan Daniel: sd386@cornell.edu

Gary Whittaker: grw7@cornell.edu

Harel Weinstein: haw2002@med.cornell.edu

[±] These authors contributed equally.

Abstract

In coronaviruses, the fusion peptide (FP) is situated within the membrane fusion domain

of the spike protein, becoming exposed following proteolytic cleavages at the S1/S2 and S2'

sites. After receptor binding-induced conformational changes, the FP penetrates the host cell

membrane and mediates membrane fusion. Previous work has revealed the importance of

calcium for SARS-CoV-1 FP structural stability and host membrane insertion. In this follow-up

study, we systematically introduced charge-neutralizing alanine mutations in the negatively

charged amino acids within the SARS-CoV-1 fusion peptide (E801A, D802A, D812A, E821A,

D825A, D830A) to identify residues that likely bind to calcium. We assayed fusion competency

by performing a syncytia-formation assay in VeroE6 cells and infectivity using pseudoparticles.

The loss of single negatively charged residues D812 or D830 greatly reduced syncytia formation

and produced noninfectious pseudoparticles. Furthermore, we observed a calcium-dependent

decrease in the infectivity of the D825A and D830A pseudoparticles, as well as fewer syncytia in

the cells expressing E821A/D825A double mutant FP. To clarify which residue pairs in the FP are

most likely to bind calcium and promote host membrane insertion, we carried out molecular

dynamics (MD) simulations of the various FP constructs. From our modeling, residue E801 is

predicted to pair with either D830 or D802 to coordinate one calcium; a second calcium ion

likely pairs residue E821 with either D812 or D825. We propose a model of bimodal calcium

binding in the FP1 and FP2 domains, which anchors the SARS-CoV-1 FP in the host cell

membrane to promote membrane insertion and fusion.

Introduction

Coronaviruses (CoVs) are a diverse family of enveloped single-stranded RNA viruses responsible for causing respiratory and enteric diseases across a wide range of species (1, 2). Three human coronaviruses have posed significant public health threats during the early 21[st] century: the severe acute respiratory syndrome coronavirus (SARS-CoV-1) in 2002, (3), Middle East respiratory syndrome coronavirus (MERS-CoV) in 2012 (4), and currently SARS-CoV-2 (5, 6). SARS-CoV-2 vaccines have proven to be highly effective at preventing morbidity and death during the pandemic; however, current experience demonstrates that there is an ever-present need for antivirals as a parallel line of defense. Development of such therapeutics requires a better understanding of the fundamental biology of coronaviruses and the intrinsic and extrinsic determinants of viral entry.

In coronaviruses, viral entry is mediated by the highly conserved and complex spike (S) glycoproteins, which form homotrimers in the virion membrane and interact with host cell receptors, such as angiotensin-converting enzyme 2 (ACE2) and DPP4 (7). Upon spike-mediated interaction with the host cell surface receptor, CoVs are proposed to utilize two different entry pathways: an "early" route involving proteolytic priming at the cell surface and a "late" route that follows the canonical endosomal/lysosomal pathways (8). The entry pathway utilized is largely dependent on the local protease environment (9, 10). Previous work has shown that SARS-CoV-1 can enter host cells via the early pathway when the transmembrane serine protease 2 (TMPRSS2) and the virus receptor ACE2 are both present in the plasma membrane, or when activated by exogenous trypsin after binding to the cell surface (11). SARS-CoV-1 can also enter cells via the endosomal route where low pH-dependent proteases, such as cathepsin

L, cleave and prime the FP for membrane insertion and fusion. This latter pathway is likely the

predominant one used by SARS-CoV-1 in most laboratory cell lines (*e.g.* Vero cells)(9, 10, 12).

The entry flexibility seen across coronaviruses arises from the differential requirements

needed for protease priming and activation of the S protein. At the boundary between the S1

and S2 domains is the S1/S2 cleavage site, which is cleaved at position R667 by trypsin-like

proteases in SARS-CoV-1 (13), with an additional downstream cathepsin site present at T678

(14). This structurally exposed cleavage loop, common across the sarbecoviruses, separates the

S1 receptor-binding domain from the S2 fusion machinery of the spike protein (15). In SARS-

CoV-1, as with other CoVs, an additional S2' prime site (R797) is located N-terminally adjacent

to the fusion peptide (FP), which becomes liberated to facilitate membrane fusion following

cleavage (16). S2' cleavage allows the S2 domain to undergo a large conformational change

that favorably positions the FP for insertion into the host cell membrane (12). Given the

importance of the FP in initiating membrane fusion, it is not surprising to find significant

conservation of the amino acid sequence in this region of the S protein across many CoVs (17,

18). A second fusion active domain was also identified in SARS-CoV-1 downstream of S2 (16),

with the two domains subsequently considered to act together as part of an extended fusion

platform, FP1/FP2 (17).

Our previous work has focused on elucidating how calcium ions impact the structure

and function of the FP (FP1/FP2) during coronavirus entry. We have previously shown that for

SARS-CoV-1 and also for MERS-CoV, intra-and extracellular depletion of calcium leads to a

significant drop in pseudoparticle infectivity, with SARS-CoV-1 pseudoparticles showing a

greater reduction in infectivity than for MERS-CoV (17, 19). In addition, we have shown using

electron spin resonance (ESR) that the FPs of these viruses induce membrane ordering in a

calcium-dependent manner. We have further predicted the stoichiometry between the FP and

calcium ions to be 1:2 for SARS-CoV-1 and 1:1 for MERS-CoV using isothermal calorimetry (ITC)

(19), and observed that the SARS-CoV-1 FP adopts a conformation with a higher degree of α-

helicity in the presence of calcium and synthetic membranes (17, 19). Taken together, these

results indicate that direct interaction between calcium and the SARS-CoV-1 FP facilitates host

membrane insertion and fusion required for host cell entry.

The CoV FP sequence contains several highly conserved negatively charged amino acids

that flank the hydrophobic residues LLF that are known to interact with the host membrane

upon FP insertion (17). We and others have posited that like the SARS-CoV-2 FP (20), the SARS-

CoV-1 FP binds calcium at the conserved negatively-charged residues, which stabilize the local

structure of the FP and position it for optimal membrane penetration(17, 19); however, a

detailed mechanistic understanding of how these residues act together to facilitate membrane

insertion has remained unclear. Here, we build off of earlier mutagenic (18) and biophysical

characterizations (21) of the SARS-CoV FP which show that the negatively charged residues of

the SARS-CoV-FP (E801, D802, D812, E821A, D825, and D830) interact with calcium and are

important for infectivity to varying degrees. We present specific mechanistic insight regarding

the preferences and conformational consequences of the binding gathered from a combination

of mutagenesis studies and computational simulations. We first systematically examined the

effects of mutating charged residues in the SARS-CoV-1 FP on syncytia formation and viral

infectivity in the presence or absence of calcium to identify residues that likely interact with

calcium. We then used corresponding molecular dynamics simulations of the system to

determine which pairs of residues likely work together to mediate calcium binding. From these

simulations we have identified the FP residue pairs predicted to bind to calcium and position

the FP favorably for host membrane insertion.

Results

Generation and characterization of SARS-CoV-1 Spike fusion peptide mutants

113 Various structural conformations of the SARS-CoV-1 spike monomer have been solved using

114 Cryo-EM (PDB: 5XLR, 5WRG, 5X58) (22). From these pre-fusion structures, it is apparent that

115 the fusion peptide segment containing the core hydrophobic residues LLF adopts an α-helical

116 conformation, followed by a proximal disordered loop known to participate in cysteine-

117 mediated intramolecular disulfide bonding **(Fig 1).**

Figure 1. Structure of the SARS-CoV-1 Spike protein. The monomeric SARS-CoV-1 spike glycoprotein in the prefusion conformation (Gui et al. 2017; PDB: 5XLR). The β strand rich S1 subunit of the spike protein contains the C-terminal domain 1 (CTD1), which mediates ACE2 binding, and the N-terminal domain (NTD). The S2 subdomain contains mostly of α-helices and is comprised of two heptad repeat regions (HR1 and HR2) and the fusion peptide (FP). The fusion peptide is highlighted in light blue and expanded to show its predicted structure and the residues of interest in this study. The hydrophobic LLF residues appear in green, negatively charged residues in red, and the disulfide bond in yellow. Beneath the structure is the 40-amino acid long sequence of the WT FP (GenBank accession no. AAT74874.1). The single charge-to-alanine mutations generated in the aspartic (D) and glutamic (E) acids are indicated in red. The two cysteine residues that form a disulfide bond within the FP2 region are highlighted in yellow.

121   To determine the roles of the negatively charged residues found in the SARS-CoV-1 FP and their

122   involvement in calcium binding, we performed alanine substitutions of the six negatively

123   residues that are highly conserved across SARS-CoV-1, MERS-CoV, and SARS-CoV-2 **(Figs 1 and**

124   **2)**.

Figure 2. Generation of the single charge-to-alanine mutations in the SARS-CoV fusion peptide.
(A) Alignment of the amino acid sequences of SARS-CoV, MERS-CoV, and SARS-CoV-2 FPs showing the highly conserved LLF motif (green), the negatively charged residues (red), and the intramolecular disulfide bond. **(B)** The sequences of the wild-type (WT) SARS CoV fusion peptide (GenBank accession no. AAT74874.1) and various mutants, containing the single charge-to-alanine mutations generated in the aspartic (D) and glutamic (E) acids. Charged residues appear in red and substitutions appear in green. Cysteine residues that form a disulfide bond appear in yellow.

We first made single amino acid (aa) substitutions (E801A, D802A, D812A, E821A, D825A,

D830A) in the wild-type SARS-CoV-1 FP. The wildtype and mutant forms of the SARS-CoV-1 S

gene were then cloned into pcDNA3.1 expression vectors to facilitate their transient expression

and characterization. To verify the expression of folded and active protein we assessed the

ability of the S to be appropriately cleaved for fusion activation. To achieve this, TPCK-trypsin

(1mg/mL) was added to cells 24 hours after transfection and the S proteins were isolated by

cell-surface biotinylation. Biotinylated proteins were retrieved following cell lysis using

streptavidin beads, resolved by gel electrophoresis, and immunoblotted for using a SARS-CoV-1

S polyclonal antibody (**Fig 3**).

Figure 3. Cell surface biotinylation assay of SARS-CoV wild-type and single charge-to-alanine spike mutants. Spike immunoblot analysis of biotinylated WT and mutant S proteins transiently expressed in HEK293T cells, then left untreated (-) or treated (+) with 1 mg/mL Trypsin for 10 minutes. S_0 indicates the full-length, uncleaved spike protein. S_1 and S_2 note the S1 and S2 cleaved subunits of spike. Spike immunoblot showing the cell surface accumulation of WT S and the single charge-to-alanine mutants. Dotted lines indicate where lanes were re-ordered on the same immunoblot; n = 3 biological replicates

The resulting spike immunoblots displayed the full-length uncleaved S species (S_0) migrating at

~185kDa in all untreated samples (**Fig 3, even lanes**). SARS-CoV-1 S mutants containing the

single mutations D802A, E821A, D825A, and D830A had comparable cell surface levels of S

protein to the wild-type protein, indicating that these mutants are not impaired in their

synthesis or trafficking to the plasma membrane. Significantly lower cell surface levels of the

D812A mutant were retrieved in comparison to WT spike (**Fig 3, lanes 5 and 6**); however,

comparable levels of the cleaved S1 and S2 subunits were observed occasionally following

TPCK-Trypsin treatment, indicating variability in the production or retrieval of this mutant in

this assay (**SI Fig 1B, lane 6**). Additionally, proteolytic cleavage of the D802A, D812A, E821A,

D825A, and D830A mutants occurred following treatment with TPCK-trypsin, indicating that

these mutants each had an accessible trypsin cleavage site and were able to be primed for

downstream fusion evens. The two lower molecular weight bands that appear at ~100kDa and

~80kDa in the all TPCK-trypsin treated lanes are likely the S_1 and S_2 subunits, respectively (**Fig**

3A and B). These subunits have been detected previously following cleavage of a recombinant S

protein monomer by Trypsin at the S1/S2 cleavage site.

It should be noted that we were unable to detect surface levels of the E801A mutant on

the spike immunoblots (**SI Fig S2**). To further probe the nature of this residue's importance for S

protein stability, we substituted in the larger, nonpolar methionine (E801M), a polar and

uncharged glutamine (E801Q), a positively charged lysine (E801K) or even the close mimetic

negatively charged aspartic acid (E801D). None of these mutations, including the charge

mimetic E801D, restored the cell surface levels of the S protein to that of wild-type, indicating

the specific requirement of glutamic acid in this region of the FP (**Fig S2**).

Analysis of the SARS-CoV-1 FP single point mutants' ability to promote syncytia formation

Functional tests of the ability of S mutants to induce cell-cell fusion were performed

using a syncytia assay. WT or mutated S constructs were transiently expressed in VeroE6 cells, a

kidney epithelial cell line that expresses the ACE2 receptor (23). 24 hours following

transfection, cell-cell fusion was induced by treating cells with TPCK-Trypsin (2mg/mL) to cleave

the FP at the S1/S2 site. Syncytia were visualized by immunofluorescence using a SARS-CoV-1 S

antibody and cells were co-stained with DAPI to identify multinucleated cells. Syncytia were

quantified by counting every group of fused cells with a minimum of 4 nuclei and expressed as a

percentage of the total number of generic fluorescent particles counted in a field of view

(yes/no binary binning method). VeroE6 cells expressing the WT S readily formed syncytia with

an average of ~47.7% of fluorescent regions of interest (ROIs) being classified as syncytia (**Fig**

4A and B).

Figure 4. Syncytia formation in SARS-CoV wild-type and single charge-to-alanine mutant S-expressing VeroE6 cells. (**A**) Representative immunofluorescence images of VeroE6 cells expressing WT or single charge-to-alanine mutant S constructs. 24 h following transfection, cells were treated with 2mg/mL TPCK-trypsin for 5 minutes to cleave S proteins and induce syncytia formation. Cells were fixed and syncytia were visualized using a SARS-CoV S antibody (*green*). Nuclei were stained with DAPI and appear in blue. Images were taken at 10X magnification. Scale bars = 90 µm (**B**) Quantification of syncytia across each condition. **** denotes significance of $p < 0.0001$. n = a minimum of 3 biological replicates.

Similarly, VeroE6 cells expressing the D802A, E821A, or D825A mutants all exhibited

comparable syncytia formation to cells expressing WT S, indicating that loss of a negative

charge in these individual residues does not affect FP function in this assay (**Fig 4A and B**).

Conversely, cells expressing the D812A and D830A S mutants largely failed to form syncytia,

indicating the importance of these residues in FP mediated cell-cell fusion (**Fig 4A and B**).

Analysis of the SARS-CoV-1 FP single mutants' effect on viral infectivity

We next tested the ability of the SARS-CoV-1 single mutants to infect cells using SARS-CoV-1

pseudoparticles as a surrogate for infectious virus. SARS-CoV-1 is designated a Risk Group 3

Select Agent that requires a specialized biosafety level 3 facilities for experimentation.

Pseudoparticles offer a safe alternative to live SARS-CoV-1 virus and enables experiments to be

carried out in BSL-2 level conditions (24). To generate SARS-CoV-1 pseudoparticles in HEK293T

cells, we used a three-plasmid co-transfection system with plasmids encoding: (1) the full

length CoV-1 surface S protein, (2) the murine leukemia virus (MLV) core proteins *gag* and *pol*,

or (3) a firefly luciferase reporter gene containing the MLV-MJ RNA packaging signal and flanked

by long terminal repeat (LTR) sequences (24). Upon successful particle fusion with the host cell

membrane, the luciferase reporter RNA transcript is reverse transcribed and expressed,

enabling a measurable readout of pseudoparticle entry. Our previous studies have shown that

this approach works well for assessing the infectivity of MERS-CoV (25), SARS-CoV-1 (17, 24),

and SARS-CoV-2 (26) pseudoparticles.

Incorporation of the WT and mutant S proteins into the SARS-CoV-1 pseudoparticles was

assessed using S immunoblotting (**Fig. 5A**). Pseudoparticles with WT or the mutant S proteins

D802A, D812A, E821A, D825A, and D830A show comparable levels of S protein incorporation

(**Fig 5A**). We believe that the observed higher molecular weight species on the S immunoblots

represents unresolved, detergent resistant S trimers present in the pseudoparticles (**Fig 5A**).

The slightly lower molecular weight band at ~185 kDa (S_0) represents full-length monomeric S

protein. The smallest band appearing at ~80kDa represents the S2 subunit (S_2). As expected,

we did not detect the E801A mutant in the SARS-CoV-1 pseudoparticles, likely due to its low

cellular expression or synthesis. After confirming the incorporation of the spike proteins into

the SARS-CoV-1 pseudoparticles, we tested their infectivity. Exposure of VeroE6 cells to WT

SARS-CoV-1 pseudoparticles results in a robust luminescent signal 72 hours post infection,

indicating successful viral entry and infection (**Fig 5B**).

Figure 5. Infectivity of wild-type and single mutant SARS-CoV spike-containing pseudoparticles. (A) Spike immunoblot of pseudoparticles containing wild-type or single charge-to-alanine spike mutants. S_0 indicates the full-length, uncleaved spike protein. S_2 indicates the cleaved S2 subunit of spike. The prominent band above 185 kDa is likely higher order species of spike multimers (gray arrow). (**B & C**) WT and mutant SARS-CoV pseudoparticle infectivity assays in VeroE6 cells left untreated (B) or treated with 50 µM BAPTA-AM (C). *,*** and **** denote significance of $p < 0.05$, 0.001, or 0.0001, respectively. n = 3 biological replicates.

After confirming the incorporation of the spike proteins into the SARS-CoV-1

pseudoparticles, we tested their infectivity. Exposure of VeroE6 cells to WT SARS-CoV-1

pseudoparticles results in a robust luminescent signal 72 hours post infection, indicating

successful viral entry and infection (**Fig 5B**). E821A-containing pseudoparticles did not exhibit a

significant change in infectivity compared to wild-type particles, suggesting this residue does

not have a major effect on FP function (**Fig 5B**). In contrast, the D802A, D812A, and D830A

mutant pseudoparticles all showed a pronounced decrease in infectivity, with the D812A and

D830A mutations resulting in essentially noninfectious particles. D825A-containing

pseudoparticles showed a slight but significant (20%) decrease in infectivity. These results

indicate that mutations in three highly conserved negatively charged residues in the S protein's

FP (D802, D812, and D830) significantly decrease SARS-CoV-1 pseudoparticle infectivity.

We then examined the effect of depleting the extra- and intracellular calcium on

pseudoparticle infectivity by using the cell-permeable calcium chelator 1,2-bis(2-

aminophenoxy)ethane- *N,N,N'N'*- tetraacetic acid tetrakis (BAPTA-AM). We have previously

optimized the concentrations of BAPTA-AM so as not to significantly impact cell viability (17,

25). VeroE6 cells were treated with 50 µM BAPTA-AM briefly prior to and during pseudoparticle

infection, harvested, and assayed for luminescence (**Fig 5C**). Depletion of extra- and

intracellular calcium rendered the WT pseudoparticles noninfectious and similar results were

observed across all the single mutant pseudoparticles tested; none of the pseudoparticles were

infectious in the absence calcium. These data indicate that depletion of extra- and intracellular

calcium exerts a severe effect on pseudoparticle infectivity and is consistent with our previous

findings. Due to the differential effect observed on infectivity with individual mutations and the

loss of available calcium, we deduced that the FP uses multiple negatively charged residues to

bind to calcium ions. We hypothesized that mutating certain residues (E821A, D825A)

individually does not lead to a complete loss in pseudoparticle infectivity since other charged

residues in the FP can compensate by serving as alternate or primary calcium binders.

Multiple negatively charged residues mediate SARS-CoV-1 fusion peptide's calcium binding

The FP contains two defined regions: the N-terminal region starting just upstream from

the S2' cleavage site (aa798-819), termed FP1, and the downstream region that includes a

critical disulfide bond (aa820-837), termed FP2 (16, 18). Lai *et al.* reported a stoichiometry of

two calcium ions per single SARS-CoV-1 FP, therefore we reasoned that FP1 and FP2 regions

each contain residues that can coordinate calcium binding (17). In FP1, the alanine substitutions

of residues D802 and D812 rendered their pseudoparticles non-infectious, as did the D830A

substitution in FP2. To test the hypothesis that the SARS-CoV-1 FP utilizes multiple residues to

mediate calcium binding, we created double residue substitutions to generate the

E821A/D825A mutant. Neither residue when individually mutated affected syncytia formation

(**Fig 4B**); however, the D825A mutant exhibited a mild reduction in infectivity (**Fig 5B**). We

hypothesized that these residues might serve as alternate calcium binding residues. Thus, we

predicted the E821A/D825A double mutant to have a greater effect on syncytia formation and

pseudoparticle infectivity. We also generated the E821A/D830A mutant as a negative control

because the D830A mutation is phenotypically dominant, blocking syncytia formation and

reducing pseudoparticle infectivity (**Fig 6A**).

As with the single charge-to-alanine S mutants, we transiently expressed the double

mutant constructs in HEK-293T cells and performed a biotinylation assay to determine if they

were synthesized, trafficked to the plasma membrane, and able to be cleaved by Trypsin (**Fig**

6B). The double mutants have comparable cell surface levels of full-length protein compared to

WT S, indicating that they are similarly expressed and trafficked to the plasma (**Fig 6B, odd**

lanes). In addition, the nearly complete disappearance of the full-length mutant S species

following trypsin treatment indicates that all mutants are cleaved by trypsin similar to WT S (**Fig**

6B, even lanes).

Figure 6. Generation and validation of double charge-to-alanine SARS-CoV spike mutants. (A) The sequence of the wild-type SARS fusion peptide, beneath which appear the double charge-to-alanine mutants, with mutated aspartic (D) and glutamic (E) acids. WT charged residues appear in red and substitutions appear in green. Yellow indicates the cysteine residues that form an intramolecular disulfide bond. (B) Cell surface biotinylation assay of SARS-CoV wild-type and mutant spike proteins generated transiently in HEK293T cells. Spike western blot analysis of biotinylated WT and mutant S proteins left untreated (-) or treated (+) with 1 mg/mL TPCK-trypsin for 10 minutes. S_0 indicates the full-length, uncleaved spike protein. S_1 and S_2 indicate the S1 and S2 cleaved subunits of spike; n = 3 biological replicates.

We tested the functionality of these S mutants using the previously described syncytia

assay. VeroE6 cells transiently expressing the double mutant E821A/D830A displayed a

significant defect in syncytia formation (< 10% syncytia) in comparison to WT S-expressing cells

(Fig 7A and B). Because the individual D830A mutant exhibited a severe defect in syncytia

formation, this phenotypic effect dominates the functional readout and we see no additional

effect on syncytia formation. The E821A/D825A double mutant exhibited a reduction in

syncytia formation (29.7% syncytia) in comparison to WT, indicating a potential additive effect

when the two individual mutations are combined.

We then generated pseudoparticles containing the E821A/D825A double mutant and

assayed the incorporation of the mutant spike into the pseudoparticles via spike

immunoblotting.

Figure 7. Assessment of syncytia formation ability and infectivity of SARS-CoV double mutant E821A/D825A. (A) Representative immunofluorescence images of VeroE6 cells expressing WT, E821A/D825A, or E821A/D825A constructs. The E821A/D825A construct served as a negative control. 24 h following transfection, cells were treated with 2mg/mL TPCK-trypsin for 5 minutes to cleave S proteins and induce syncytia formation. Syncytia were visualized using a SARS-CoV S antibody (*green*) following fixation. Nuclei were stained with DAPI and appear in blue. Images were taken at 10X magnification. Scale bars = 90μm (B) Quantification of syncytia across each condition.(C) Spike immunoblots of pseudoparticles containing VSVG (no spike control), WT, or mutant E821A/D825A spike. S_0 indicates the full-length, uncleaved spike protein. (D) WT and E821A/D825A SARS-CoV pseudoparticle infectivity in VeroE6 cells left untreated (-) or treated with 50 μM BAPTA-277   AM (+).** and **** denote significance of $p < 0.05$ and 0.0001, respectively. n = 3 biological replicates

The E821A/D825A mutant pseudoparticles exhibited similar S protein levels to the WT

pseudoparticles (**Fig 7C**). Interestingly, VeroE6 cells infected with the mutant E821A/D825A

pseudoparticles exhibited a 28% increase in infectivity compared to WT pseudoparticle infected

cells (**Fig 7D**). As was previously demonstrated, depletion of extra- and intracellular calcium via

BAPTA-AM treatment causes a severe loss of infectivity for both WT and mutant E821A/D825A

pseudoparticles **(Fig 7D).**

Molecular Dynamics simulations identify the modes of the SARS-CoV-1 FP's calcium binding

The effects on cell-cell fusion and infectivity exhibited by the D802A, D812A and D830A

mutants, as well as the calcium-dependent loss of infectivity in the E821 and D825 mutants,

implied that the FP contains multiple sites of calcium binding required for viral entry. Our ability

to determine which residue pairs in the FP coordinate calcium binding was limited using our *in*

vivo mutagenesis approaches; therefore, we carried out atomistic molecular dynamics (MD)

simulations of the SARS-CoV-1 FP to predict the preferred modes of calcium binding and

membrane insertion. In particular, this MD approach allowed us to examine residue 801, which

was refractory to mutagenesis.

To monitor the spontaneous binding of calcium to the FP, we collected 18 independent MD

trajectories of 640ns in length for each construct. Following the analysis protocols described

recently for SARS-CoV-2 FP simulations (20, 27), the various modes of interactions between the

SARS-CoV-1 FP and calcium ions were assessed in the trajectories by monitoring: i) the

distances between the calcium ions in solution and the side chains of all acidic residues in the

peptide; and ii) the pairwise distances between the side chains of all acidic residues.

Summarized in **Fig S3** are the observed events of simultaneous association of two calcium ions

with various pairs of FP residues in the individual trajectories of the WT and the mutant systems

(see *red* and *blue* rectangles). The combined statistics for each construct (*i.e.,* the total number

of binding events for different pairs of residues) are summarized in **Fig 8**.

These results show the modes of calcium coordination in the WT protein, identifying the

most frequent modes involving residue pairs E821/D825 (5 out of 18); E801/D802 and

E801/D830 (4/18 each); and D812/E821 (2/18). Notably, a similar pattern of coordination

preference was observed in our recent computational studies of calcium association with SARS-

CoV-2 FP (20). Moreover, simultaneous binding of calcium ions to the SARS-CoV-2 FP residue

pairs equivalent to E801/D802 and D812/E821 produced the peptide conformations suitable for

membrane penetration. In contrast, conformations that stabilized calcium binding to residues

equivalent to the E821/D825 pair did not enable sustained bilayer insertion.

Figure 8. Predicted modes of Ca²⁺ and SARS-CoV fusion peptide binding. Number of Ca²⁺ binding events to different pairs of anionic residues in SARS-CoV FP in the simulations of the WT and the mutant constructs (labeled in each panel). The experimentally measured phenotypes for each construct are shown (active or inactive). The mode of Ca²⁺ ion association predicted to be inhibitory for membrane insertion (E821-D825) is shown in *red*, and the modes of Ca²⁺association predicted to facilitate membrane insertion are depicted in *green*.

From these MD simulations of the SARS-CoV-1 FP constructs, we were able to predict the

phenotypes for each construct (active or inactive). The mode of calcium ion association

predicted to be inhibitory for membrane insertion (E821-D825) is shown in *red*, and the modes

of calcium association predicted to facilitate membrane insertion are depicted in *green* boxes

(**Fig 8**). In those constructs with single mutations that were predicted to not have fusion activity

(E801A, D802A, D812A, and D830A) we showed that only the calcium-coordination mode

involving the E821/D825 pair persists, but not the D812/E821 pair of residues (**Fig 8**).

Conversely, in the SARS-CoV-1 FP single mutants that maintained WT-like fusion activity (E821A

and D825A), we did not observe the E821/D825 residues predicted to participate in calcium

binding. In these function-preserving mutants we identified additional modes of calcium

binding that are enhanced in comparison to the WT system: E801/D802 in mutant E821A, or

E801/D830 in mutant D825A. For the SARS-CoV-1 FP constructs with multiple mutations, our

MD trajectories revealed an overall reduced calcium binding ability (**Fig 8** and **Fig S4**).

SARS-CoV-1 FP propensity for membrane insertion is regulated by modes of calcium binding

Overall, the above computational results reveal that the calcium binding patterns of SARS-

CoV-1 FP are very similar to those of SARS-CoV-2 FP (20). On this basis, membrane insertion of

the SARS-CoV-1 FP could be expected to be enhanced by the modes of calcium binding

involving the E801/D802 and D812/E821 pairs of residues, and to be reduced by the ones

involving the E821/D825 pair. To test this premise, we carried out MD simulations of two

models of the WT SARS-CoV-1 FP spontaneously associating with the lipid membrane (see

Methods). In one, the peptide was interacting with 2 calcium ions at the E801/D802 and

D812/E821 sites (Model 1), and in the second, 2 calcium ions were bound to the E801/D802

and E821/D825 pairs (Model 2). Each structure was simulated in 36 independent replicates,

each run for ~0.9-1.0 μs (see **Methods**).

Analysis of these trajectories revealed that, indeed, Model 1 extensively penetrated the

membrane, while the membrane insertion of the Model 2 was negligible. This can be seen from

the plots presented in **Fig 9** comparing frequencies of membrane insertion for each SARS-CoV-1

FP residue in the simulations of Model 1 and Model 2 structures.

Figure 9. Membrane insertion for FP1 and FP2 domains of SARS-CoV fusion peptide. (A-B) Frequency of membrane insertion for each residue of SARS-CoV FP in the MD simulations of Model 1 (**A**) and Model 2 (**B**) constructs, which differ in the mode of Ca^{2+} coordination (see text). The ribbon diagrams in the panels show the corresponding structures. Ca^{2+} coordinating anionic residues are indicated in black text; Ca^{2+} ions are shown as blue spheres. The FP1 and FP2 parts of the fusion peptide are colored in silver and green, respectively. (**C-D**) The same frequency calculations but done separately for two representative trajectories in Model 1 set in which the observed membrane insertions involved the N-terminal LLF motif (**C**) and the more centrally located (F815-M816) hydrophobic segment (**D**). The corresponding ribbon diagrams illustrate the structural features of these two distinct insertion modes highlighting (in black) positions of the anionic residues, the hydrophobic residues penetrating the membrane, and the R829 residue interacting with D825. The same color code applies here as in panels A-B.

The membrane insertion was quantified by monitoring the z-coordinate of the C_α atom of each

residue in the peptide. A residue was considered inserted into the membrane if the z-distance

between its C_α atom and the second carbon atom in the tail of a POPC lipid (atom C22 in

CHARMM36 notation) was <5Å. In Model 1 (**Fig 9A**), the N-terminal FP1 segment of the fusion

peptide shows strong propensity for bilayer insertion, while in Model 2 the insertion is minimal

(**Fig 9B**).

The detailed analysis of the individual trajectories in the Model 1 set revealed two distinct

modes of bilayer penetration for SARS-CoV-1 FP, similar to our findings for the SARS-CoV-1-2

FP. Thus, the Model 1 construct penetrates the bilayer either with its N-terminal LLF motif (**Fig

9C**) or with the more centrally located hydrophobic F815-M816 segment (**Fig 9D**). Interestingly,

the two insertion modes appear to alternate which calcium ion is neighboring the inserted

portion. Indeed, when the LLF is inserted, the calcium ion associated with the neighboring

E801/D802 residues is bound to the membrane while the other calcium binding site

(D812/E821) is situated away from the membrane surface (**Fig 9C**). In the case of the F815-

M816 insertion, the position of the calcium binding loci with respect to the membrane is

reversed – the one associated with the D812/E821 pair is membrane-bound, while the

E801/D802 pair is located farther from the bilayer (**Fig 9D**). We also note that in both cases, the

remaining anionic residues in the peptide (*i.e.* the ones not engaged with the calcium ions) are

either solvent exposed (D830) or engaged with electro-neutralizing interactions with

neighboring basic residues (D825/R829) (**Fig 9C and D**). These results support a mechanistic

model in which membrane penetration of the SARS-CoV-1 FP is significant only for specific

modes of calcium binding to the peptide, i.e., to the E801/D802 and D812/E821 pairs of

conserved acidic residues. Moreover, calcium binding at the E821/D825 pair is predicted to

inhibit membrane insertion.

Discussion

Previous work done on the SARS-CoV-1 FP has implicated all of the negatively charged

residues as calcium binders, but to varying degrees. Early structure-function work on the SARS-

CoV-1 FP showed the importance of general FP regions first, with the residues flanked by

cysteines (C822-C833) identified as important for membrane fusion activity (18). FP residues

that when mutated caused pronounced effects on syncytia formation and pseudoparticle

infectivity were then identified (D812 and D830), but before the importance of calcium during

viral entry was known (16). Recent biophysical characterizations of the FP found that mutation

of residue D812 in the FP1 region has the greatest effect on membrane ordering, calcium-

specific FP binding, and calcium-induced FP structural changes. Residue D830 in the FP2 region

(labeled 831 in their manuscript) was also found to be important, causing of loss of membrane

ordering and calcium sensitivity when mutated. In this study, we have taken a two-pronged

approach to identify how the negatively charged residues in the SARS-CoV-1 FP likely bind to

calcium during viral entry: biological studies and molecular dynamics simulations. We first

carried out syncytia formation and infectivity assays with single charge-to alanine FP mutants

E801A, D802A, D812A, E821A, D825A, and D830A (**Fig 2**), confirming that all mutants were

synthesized, accumulated at the cell surface comparable to WT S, and were able to be cleaved

by Trypsin (**Fig 3**).

We next tested the fusion competency of the single FP mutants using a syncytia

formation assay. VeroE6 cells expressing the D802A, E821A, and D825A mutants all formed

syncytia similar to WT S expressing cells (**Fig 4**). Conversely, cells expressing the D812A or

D830A mutants exhibited a pronounced cell-cell fusion defect (**Fig 4**). Infectivity assays

performed with pseudoparticles largely produced the same results: infections with E821A and

D825A S-containing pseudoparticles resulted in luminescence levels comparable or minorly

reduced to WT-containing pseudoparticles. However, pseudoparticles containing the D802A,

D812A, or D830A mutations were unable to infect VeroE6 cells as well. These results indicate

that when calcium levels remain unchanged, loss of an individual negatively charged residue in

FP position D802, D812, or D830 is sufficient to cause a substantial decrease in infectivity. As

previously mentioned, mutagenic studies of the SARS-CoV-1 FP found that mutations in the

SARS-CoV-1 residues D812 (D812A) and D830 (D830L) led to a decrease in syncytia formation

and pseudotyped virion infectivity (16, 18). Taken together, these data suggest that these

residues are essential in a non-redundant manner to the function of the SARS-CoV-1 fusion

peptide.

We then tested the requirement of intracellular calcium on pseudoparticle infectivity by

treating cells with a membrane permeable calcium chelator, BAPTA-AM, prior to pseudoparticle

infection. Following cell treatment with BAPTA-AM, WT and all mutant FP-containing

pseudoparticles are no longer infectious (**Fig 5C**). It is apparent from these data that this

approach could not assay specific FP residues' sensitivity to intracellular calcium. Disruption of

physiological concentrations of calcium is known to impact the fusion and infectivity of SARS-

CoV-1 and SARS-CoV-2 (28). We propose that calcium chelation via BAPTA-AM treatment

disrupted endosomal calcium concentrations causing a major reduction in endosomal fusion of

WT pseudoparticles, as SARS-CoV-1 enters VeroE6 cells primarily via the endosomal/lysosomal

pathway. BAPTA-AM treatment likely disrupted endosomal calcium levels which left

pseudoparticles unable to fuse with the endosomal/lysosomal membranes. We acknowledge

the potential impacts, both direct and indirect, that BAPTA-mediated depletion of calcium may

have on early pathway entry mechanisms. Calcium levels are proposed to affect host cell

receptor interactions at the plasma membrane; the SARS-CoV-2 spike protein's receptor

binding domain contains an RGD (Arginine-Glycine-Aspartate tripeptide) motif that is predicted

to bind to integrins in a calcium-dependent manner (29). ACE2 receptor turnover has also been

proposed to be a calcium-dependent process based on numerous lines of evidence

demonstrating the involvement of the calcium-binding protein, calmodulin (30-32). Because

calcium plays many important roles in cells and during viral infection, we believe that the loss in

infectivity we observed in our pseudoparticle assay reflects the cumulative effect of calcium

disruption on multiple pseudoparticle entry steps.

Previous mutagenic studies of the FP using biophysical readouts of function suggested

that FP residues display varying sensitivities to calcium, with the FP2 residues E821 and D825

showing modest calcium sensitivities and effects on function when mutated (21). We therefore

generated a double mutant (E821A/ E825A) to determine if these residues, when mutated

together, caused a more pronounced biological phenotype. We also generated the

E821A/D830A double mutant to serve as a negative control. Consistent with the single D830A

mutant, we observed a fusion defect in VeroE6 cells transiently expressing the E821A/D830A

mutant (**Fig 7A and B**). Furthermore, the E821A/E825A double mutant showed a significant

reduction in the percentage of syncytia formed, suggesting that the genetic effect is additive at

these loci (**Fig 7B**). When then assayed the infectivity of this mutant using pseudoparticles,

showing first that the double mutant E821A/D825A was successfully incorporated into

pseudoparticles (**Fig 7C**). In comparison to WT FP-containing pseudoparticles, the E821A/D825A

double mutant showed an increase in infectivity, while treatment with BAPTA-AM rendered

both types of pseudoparticles essentially noninfectious (**Fig 7D**). Although we cannot fully

reconcile why we observe a decrease in syncytia formation and an increase in infectivity for the

E821A/D825A-containing pseudoparticles, we hypothesize that these mutations may be

detrimental to spike-mediated fusion but advantageous at a downstream (post-fusion) event

which contributes favorably to viral replication.

From our *in vivo* findings, which confirmed the functional requirements of the

negatively charged residues in the FP, we then focused on identifying the residues that form a

calcium-binding platform. Because mutation of either FP residue D812 or D830 resulted in a

complete loss in S fusion competency and pseudoparticle infectivity, we predicted that they are

essential residues for calcium binding. In addition, residues E821 and D825 appeared to serve

as calcium binding alternates but did not appear to be supporting FP function in a major way.

These inferences were probed using molecular dynamics (MD) simulations of the calcium-

loaded SARS-CoV-1 peptide with the membrane. Exploring the probability of each FP residue

interacting with the membrane in the identified calcium-binding modes reveals the preferred

mode of peptide insertion to be with the N-terminal end interacting with a calcium ion

coordinated by residues E801 and D802 (**Figs 8 and 10**). The depth of insertion and the

determinant role of the calcium ion in facilitating membrane insertion are identical to the

findings for the SARS-CoV-2 peptide and are in agreement with the experimental

measurements of SARS-CoV-1 FP membrane insertion by Lai *et al* (17, 25, 33). That calcium

interacts with the same highly conserved, charged residues in the FPs of SARS-CoV-1 and SARS-

CoV-2 is not surprising.

Results from our computational modeling of SARS-CoV-1 FP-membrane interactions in the

presence of calcium show that the propensity of the SARS-CoV-1 FP for membrane insertion is

dictated by specific modes of calcium binding. The calcium binding mode that enables the most

sustained membrane penetration involves the initial association of the peptide with the bilayer

through a calcium binding site located near the N-terminus of the peptide. This site

(E801/D802) is made accessible following enzymatic cleavage and is inserted into the

membrane via the insertion of the juxtaposed hydrophobic segment (the LLF motif). In this

process, all anionic residues of the peptide are either engaged with calcium ions, with

neighboring basic residues, or remain solvent exposed away from the membrane. Based on the

preferred modes of calcium-loaded FP interaction with the membrane, we propose that the

binding of calcium following S protein cleavage at the S2' site creates an energetically feasible

membrane insertion process composed of the following steps: (1) steering of the peptide

towards the bilayer surface; (2) exposure of the hydrophobic layer of the membrane to the

conserved N-terminal LLF motif of the FP anchored by the calcium interactions with the

phospholipid headgroups; and (3) shielding of the anionic residues of the peptide by the

preferred inserting conformation of the FP to prevent their negative charge from interfering

with membrane insertion (**Figure 9**). These three components became evident from the

analysis of the MD simulation trajectories presented here for the SARS-CoV-1 FP and bear a

very high similarity to results from computational studies of the SARS-CoV-2 FP (20).

Remarkably, in both the SARS-CoV-2 FP simulations (34) and here, we observed a calcium

binding mode that inhibits long-lasting membrane insertion. In this mode calcium association

with the E821/D825 pair (corresponding to the D839/D843 in SARS-CoV-2 FP) positions the LLF

motif near the negatively charged D812 residue at the membrane surface (**Fig 9B**). This

eliminates the favorable effect of the negative charge screening by the calcium, so that the

peptide bilayer encounters are transient and do not lead to penetration by the LLF motif. This

mechanistic model, which highlights the role of the positioning of the D812 residue in specific

calcium binding modes, is supported by our findings for the SARS-CoV-1 S mutants, as it was

previously found to be an impactful mutation in the SARS-CoV-1 FP (16). A recent study from

Koppisetti *et al* asserts that a calcium ion may be binding to D843/D849 in SARS-CoV-2 FP

(corresponding to D825/D830 in SARS-CoV-1 FP); however, as shown from the simulations in

this study and Khelashvili *et al*, there is a high likelihood that the second calcium ion will bind to

D812/D821 (corresponding to the D830/D839 in SARS-CoV-2 FP) for the maximal membrane

penetration (20, 35). Recent NMR studies regarding the membrane interaction with the fusion

peptide still have yet to be fully resolved, as current models use bicelles (35) and micelles (36),

which do not completely capture the composition of the bilayer as was done for the

computational modeling shown here; though the model of membrane insertion proposed by

Koppisetti *et al (35)* most aligns with our proposed CoV FP interaction with bilayer (**Fig 9**).

Thus, as shown in **Figs 8** and **9** and summarized in **Fig 10**, mutations that create a high

propensity for the calcium binding mode involving the E821/D825 pair, stabilize peptide

conformations that are non-productive for membrane penetration and are found here to

inactivate the FP (**Fig 9**).

Figure 10. Summary of SARS-CoV-1 FP mutagenesis and MD simulations work. (A) The FP residues found to be necessary to promote cell-cell fusion and infectivity (biological studies) are the same residues predicted to pair with calcium to promote membrane insertion (MD simulations). The mutant E801A was not detected in our biological studies and appears in gray. **(B)** The FP residue pairings with calcium ions predicted to promote (green dashed lines) or inhibit (red dashed lines) membrane insertion. **(C)** A proposed model of the key charged residues in SARS-CoV-1 FP and their interactions with calcium ions and the membrane (outer leaflet). Depicted is the insertion of the FP1 subdomain, with the E801/D802-Ca^{2+} pairing predicted to promote membrane insertion of the LLF residues. D812 is predicted to bind Ca^{2+} with E821 to promote membrane interactions. The FP structure is based on the SARS-CoV-1 pre-fusion conformation (PDB: 5XLR). **(D)** A model of the SARS-CoV-1 trimer interacting with the calcium ions (in yellow, only 5 in view) and the host membrane. Upon interaction with calcium, the FP1 region adopts a more helical structure and can penetrate the host cell membrane to facilitate membrane fusion. BioRender was used to generate part of this figure.

Current studies on SARS-CoV-2 S protein classified the disordered portion of the FP in the

FP2 domain as the FPPR (fusion peptide proximal region), which includes the charged residues

D839/D843/D848 or E821/D825/D830 equivalent in SARS-CoV-1 FP (20, 37). This FPPR region

was determined to be important, as it binds to the RBD through the CTD1, and maintains the

closed pre-fusion S trimer(37). This is due to the tight packing around the disulfide bond

reinforced by a bond between K835 and D848 (K817 and D830 in SARS-CoV-1 FP). The

equivalent residue in SARS-CoV-1 FP, based on biological studies, is a critical residue (37).

Mutating D830 removes the binding partner of K817, thereby loosening the 'knot' of the

structured FPPR in maintaining the closed form of the trimeric S protein (37). Without the

reinforcements, we may lose the optimal structural conformation of the S protein upon

receptor binding to expose the S2' cleavage site to promote successful proteolytic cleavage.

More broadly, the E801 and D830 residues are conserved within the CoV family and the

D812 residue is conserved within betacoronaviruses (16, 17). The implication is that calcium

interactions are a conserved mechanism that serves to better position the FP for membrane

insertion. Different coronaviruses exhibit different requirements for calcium; MERS-CoV binds

to one calcium ion in its FP1 domain (25), thus, it is important to investigate the role of calcium

and FP interactions across the CoV family. Recent work using a FRET-assay has shown that

calcium interactions with the fusion peptide may drive the pre- to post-fusion conformational

changes in SARS CoV-2 spike (27). The authors of this study showed that alpha and beta

coronaviruses display a high sensitivity to calcium; the fusion activity of spike drops

precipitously as calcium levels rise or decrease past predicted late endosomal concentrations.

The conservation of calcium-binding residues in the FP of many coronaviruses suggests that the

CoV fusion mechanisms are essential for function and can serve as potential targets for broad-

spectrum antiviral drugs (38-40). Repurposing FDA-approved calcium channel blocking (CCB)

drugs to inhibit CoV entry, particularly for SARS-CoV-2, is one option worth exploring. Recent

studies have shown that the CCB felodipine is a potential candidate to inhibit SARS-CoV-2 entry

(41). CCBs can target conserved viral functions, providing a rapid solution to address new and

future SARS-CoV-2 variants. It will be important to identify the mechanisms of CCBs CoV

inhibition, as they may directly inhibit a viral target or indirectly inhibit viral entry by affecting

host cell processes. Interestingly, SARS-CoV-2 variants have arisen as part of Clade 20A that

contain a D839G or D839Y mutation (E821 equivalent in SARS-CoV-1), with this mutation

predicted to affect FP-Calcium interactions (42-44). To date, it is not known if there is any

selective advantage to the virus conferred by this mutation or whether the emergence of these

variants simply represents a founder effect. Future studies will continue to illustrate the role of

the spike FP in controlling both emerging and existing coronaviruses.

Materials and Methods

Cells, plasmids, and reagents. Human embryonic kidney 293 (HEK293T) and African green

monkey kidney epithelial (VeroE6) cells were obtained from the American Type Culture

Collection (ATCC, Manassas, VA). Both cell lines were grown in complete DMEM, composed of

Dulbecco's Modified Eagle medium (DMEM, CellGro), supplemented with 10% HyClone

FetalClone II (GE) and 10 mM HEPES (CellGro), and grown at 37°C with 5% CO_2. The plasmids

used for generating pseudoparticles were the pCMV-MLV gag-pol murine leukemia virus (MLV)

packaging construct, the pTG-Luc transfer vector encoding the luciferase reporter gene, and the

pCAGGS-VSVG plasmids. They were generously provided by Jean Dubuisson (Lille Pasteur

Institute, Lille, France) and co-transfected as previously described (24). The plasmid encoding

the C9-tagged SARS-CoV-1 spike protein (pcDNA3.1-C9-SARS-CoV-1 S) was provided by Dr.

Michael Farzan from the New England Primate Research Center. Recombinant L-1-tosylamide-

2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin was obtained from Sigma. The

calcium chelators EGTA and BAPTA-AM (acetoxymethyl ester) were obtained from VWR and

Tocris, respectively. Both compounds were diluted in dimethyl sulfoxide (ThermoFisher) prior to

use.

Site-directed mutagenesis. Site-directed mutagenesis was performed on the SARS-CoV-1 spike encoding plasmid, pcDNA3.1-SARS-CoV-1-S, via the QuikChange Lightning site-directed mutagenesis kit (Aligent). PCRs and transformations were performed based on the manufacturer's recommendations. The primers used to generate the SARS-CoV-1 S mutants can be found in the supplementary information (**SI, Table S1**). Mutations were confirmed via Sanger Sequencing at the Cornell University Life Sciences Core Laboratories Center.

Cell Surface Biotinylation Assay. HEK293T cells were seeded in poly-D-lysine coated 6-well dishes. 24 hours later, cells were transfected with 1.5 ug of WT or mutant SARS-CoV-1 S plasmid using polyethylenimine (PEI) (Thermo Fisher Scientific). Cells were transfected for 24 hours, washed once with 1X DPBS, and then left untreated or treated with 1 mg/mL of TPCK-trypsin in DPBS for 5 minutes at 37°C. Cell surface proteins were then biotinylated by incubating rinsed cells in a biotin-containing buffer (250 ug/mL in PBS) for 20 minutes. The biotin buffer was then replaced with a quenching solution (50mM glycine in DPBS) and incubated for 30 minutes. Cells were then lysed in ice for 15 minutes using a lysis buffer (0.1% Triton in 1x TBS) with protease inhibitors (cOmplete Protease Inhibitor Cocktail). Lysates were collected and spun at 13,000 rpm for 10 minutes to pellet insoluble material. The lysates were then incubated overnight at 4° C with equilibrated steptavidin beads to retrieve the biotinylated spike proteins. The captured spike proteins were eluted by boiling the beads in 1x LDS sample buffer in the presence of DTT (NuPAGE) for 10 minutes. Proteins were resolved on 4-12% gradient Bis-Tris gels (NuPAGE) and transferred to PVDF membranes. The SARS-CoV-1 S protein was detected using the SARS-CoV-1 S rabbit polyclonal primary antibody (NR-4569, BEI resources) and Alexa Fluor 488-labeled goat

anti-rabbit secondary antibody (Invitrogen). The resulting protein bands were visualized using a

Chemidoc system with Image Lab image capture software (BioRad).

Cell-cell fusion assay. VeroE6 cells were seeded onto glass slides in a 24-well chamber slides

(Millipore). After 24 hours, cells were transfected using Lipofectamine according to the

manufacturer instructions. Following 18 hours of transfection, cells were washed with 1X DPBS

and trysinized with 2mg/mL TPCK-trypsin for 5 minutes at 37°C to. Cells were then fixed with

4% paraformaldehyde (PFA) (ThermoFisher) for 15 minutes and washed three times with DPBS.

To permeabilize the cells, a 0.1% Triton X-100 in DPBS solution was added to each well and cells

were incubated for 5 minutes. After three washes with DPBS, the cells were blocked with 5%

normal goat serum for 30 minutes. Cells were again washed and then labeled with SARS-CoV-1

S rabbit polyclonal antibody (NR-4569, BEI resources), followed by labeling with Alexa Fluor

488-labeled goat anti-rabbit secondary antibody (Invitrogen). The cell nuclei were labeled with

a DAPI stain present in the mounting media (DAPI Fluoromount-G, Southern Biotech). Images

were acquired using an upright microscope (Echo Revolve) with a 10x objective. To quantify the

number of nuclei per syncytia, three randomly selected fields were acquired and spike-

expressing cells and syncytia were manually counted. The total number of syncytia we counted

and expressed as a percentage of the total fluorescent "particles" that were counted in the

image. The data was graphed and statistics were carried out using Microsoft Excel and

GraphPad Prism 7.

SARS-CoV-1 pseudoparticle infectivity assay. SARS-CoV-1 pseudoparticles were produced as

previously described (24, 25). Briefly, to generate pseudoparticles 3.5 x10^5 HEK293T cells were

seeded in 6-well plates. Pseudoparticles were prepared by transfecting HEK293T cells with 600

µg of their respective SARS WT or mutant S plasmids, 800 µg of pTG-Luc plasmid, and 600 µg of

pCMV-MLV gag-pol plasmid using polyethylenimine (PEI) as the transfection reagent. The cell

supernatant was harvested 48 hours post-transfection, centrifuged at 1200 rpm for 7 minutes

to separate the pseudoparticles from residual cellular debris, and filtered through a 0.45 µm

syringe filter. The pseudoparticles were stored at -80°C for one freeze-thaw cycle. To perform

the pseudoparticle infectivity assay, $5x10^5$ VeroE6 cells were seeded into 24-well plates and

infected with 200uL of the pseudoparticle suspension for 72 hrs at 37°C. Cells were lysed for 10

minutes at room temperature using the Luciferase cell lysis reagent (Promega). Luminescence

readings were performed using a Glomax 20/20 Luminometer system (Promega). Each

experiment contained three technical replicates and was repeated at least three times. Data

analysis was performed using GraphPad Prism 7.

For the calcium depletion studies, calcium was chelated from the extracellular and

intracellular environment using EGTA or BAPTA-AM, respectively. To deplete extracellular

calcium levels, VeroE6 cells were pre-treated with DMEM lacking calcium (DMEM- lacks L-

glutamine, sodium pyruvate, HEPES, and calcium; it contains 2% HyClone and 10 mM HEPES) for

1 hour at 37°C. After the DMEM- pretreatment, the VeroE6 cells were infected with 200 µL of

pseudoparticle suspension in the presence of 50 µM EGTA for 2 hours at 37°C. Following

infection, cells were supplemented with 300 µL of calcium-containing cDMEM.

To chelate the remaining intracellular calcium, VeroE6 cells were pretreated with 50 µM

BAPTA-AM in DMEM+ (DMEM without L-glutamine, sodium pyruvate, HEPES; 2% HyClone; 10

mM HEPES; Gibco) in the same conditions as described above. Cells were then infected using

200 µL of the pseudoparticle suspension in the presence of 50 µM BAPTA-AM for 2 hours and

supplemented with cDMEM post-infection. Because BAPTA-AM was dissolved in DMSO, VeroE6

cells were pretreated with the equivalent percentage of DMSO used in the BAPTA-AM treated

cells. This served as an important control for the effects of DMSO alone on our infectivity assay.

Analysis of pseudoparticle infectivity under calcium depletion conditions was performed as

described above. To confirm the presence of the S protein in the pseudoparticles, 1.5 mL of the

harvested pseudoparticles were spun down using an ultracentrifuge in a TLA-55 rotor

(Beckman-Coulter) at 42,000 × g for 2 hours at 4°C. The supernatant was aspirated and the S

protein-containing pellet was resuspended and boiled in 1x LDS sample buffer in the presence

of DTT for 10 minutes. The samples were resolved and immunoblotted for the spike protein as

described above.

Modeling of the SARS-CoV-1 fusion peptide. The structural model of the SARS-CoV-1 S protein

fusion peptide is based on the SARS-CoV-1 S prefusion structure from the Protein Data Bank,

PDB 5XLR. The sequence of the SARS-CoV-1 S protein (GenBank accession no. AAT74874.1) was

aligned to the PDB 5XLR SARS-CoV-1 structure sequence using Geneious software (**v.2020.1.1**).

Structural models of the SARS-CoV-1 S monomer were generated using the Modeller

comparative modeling tool (**v.9.23**) within the Chimera software (v.1.13; University of

California). Images were created using Adobe Illustrator CC (**v.24.03**).

Molecular dynamics (MD) simulations of the SARS-CoV-1 FP in water. For all the atomistic MD

simulations, the SARS-CoV-1 FP segment was capped with neutral N- and C-termini (ACE and

CT3, respectively, in the CHARMM force-field nomenclature). Protonation states of all the

titratable residues were predicted at pH 7 using Propka 3.1 software (45).

For the simulations in water, one copy of the peptide (wild-type or a mutant) was

embedded in a rectangular solution box and ionized using VMD tools ("Add Solvation Box" and

"Add Ions", respectively) (46). The box of dimensions ~90 Å x 80 Å x 82 Å included a Na^+Cl^- ionic

solution as well as 2 Calcium ions, and ~18000 water molecules. The total number of atoms in

the system was ~54,540.

The system was equilibrated with NAMD version 2.13 (47) following a multi-step protocol

during which the backbone atoms of the SARS-CoV-1 FP as well as Calcium ions in the solution

were first harmonically constrained and subsequently gradually released in four steps (totaling

~3ns), changing the restrain force constants k_F from 1, to 0.5, to 0.1 kcal/ (mol $Å^2$), and 0 kcal/

(mol $Å^2$). These simulations implemented all option for rigidbonds, 1fs (for k_F 1, 0.5, and 0.1

kcal/ (mol $Å^2$)) or 2fs (for k_F of 0) integration time-step, PME for electrostatics interactions (48),

and were carried out in NPT ensemble under isotropic pressure coupling conditions, at a

temperature of 310 K. The Nose-Hoover Langevin piston algorithm (49) was used to control the

target P = 1 atm pressure with the "LangevinPistonPeriod" set to 200 fs and

"LangevinPistonDecay" set to 50 fs. The van der Waals interactions were calculated applying a

cutoff distance of 12 Å and switching the potential from 10 Å.

After this initial equilibration phase, the velocities of all atoms in the system were reset and

ensemble MD runs were initiated with OpenMM version 7.4 (50) during which the system was

simulated in 18 independent replicates, each for 640ns (i.e., cumulative time of ~11.5 µs for

each FP construct). These runs implemented PME for electrostatic interactions and were

performed at 310K temperature under NVT ensemble. In addition, 4fs time-step was used, with

hydrogen mass repartitioning and with "friction" parameter set to 1.0/picosecond. Additional

parameters for these runs included: "EwaldErrorTolerance" 0.0005, "rigidwater" True, and

"ConstraintTolerance" 0.000001. The van der Waals interactions were calculated applying a

cutoff distance of 12 Å and switching the potential from 10 Å.

MD simulations of SARS-CoV-1 FP interactions with lipid membranes. Interactions of selected

two models of the Calcium-bound WT SARS-CoV-1 FP with lipid membranes were investigated

with atomistic MD simulations. These runs were initiated by placing each of the models in the

proximity of a bilayer composed of 3:1:1 POPC/POPG/Cholesterol that had been pre-

equilibrated for 25ns as described previously (20).

After the FP-membrane complexes were embedded in a solution box (containing 150 mM

Na^+Cl^- salt concentration), each system was equilibrated with NAMD version 2.13 following the

same multi-step protocol described above during which the backbone atoms of the FP as well

as the Calcium ions were first harmonically constrained and subsequently gradually released in

four steps. After this phase, the velocities of all atoms of the system were reset, and ensemble

MD runs were initiated with OpenMM version 7.4. Each system was simulated in 18

independent replicates, each ran for ~ 1 μs (i.e., cumulative time of ~18 μs for each FP-

membrane complex). These runs implemented PME for electrostatic interactions and were

performed at 298K temperature under NPT ensemble using semi-isotropic pressure coupling,

with 4fs time-steps, using hydrogen mass repartitioning and with "friction" parameter set to

1.0/picosecond. Additional parameters for these runs included: "EwaldErrorTolerance" 0.0005,

"rigidwater" True, and "ConstraintTolerance" 0.000001. The van der Waals interactions were

calculated applying a cutoff distance of 12 Å and switching the potential from 10 Å. For all

simulations we used the latest CHARMM36 force-field for proteins and lipids (51), as well as the

recently revised CHARMM36 force-field for ions which includes non-bonded fix (NBFIX)

parameters for Na$^+$ and Calcium ions (52).

Acknowledgements

This work was supported by The National Institute of Health research grant R01AI35270,

National Science Foundation RAPID grant 2027070 and Fast Grant from the Mercatus Center at

George Mason University. TT is supported by the National Science Foundation Graduate

Research Fellowship Program under Grant No. DGE-1650441 and the Samuel C. Fleming Family

Graduate Fellowship. H.W. and G.K. gratefully acknowledge support from the 1923 Fund, and

the access to computational resources awarded through the COVID-19 High Performance

Computing Consortium at the Center for Computational Innovations (CCI) at the Rensselaer

Polytechnic Institute and at the Oak Ridge Leadership Computing Facility, which is a DOE Office

of Science User Facility supported under Contract DE-AC05-00OR22725. We would like to thank

members of the Daniel, Whittaker, and Abbott groups as well as the Weinstein group at Weill

Cornell for helpful discussions.

Author Contributions

710 Conceived and designed the experiments: MKB GK MRS HW GRW SD. Performed the

711 experiments: MKB JDC GK MRS TT. Analyzed the data: MKB JDC GK MRS TT HW GRW SD.

712 Wrote and edited the paper: JDC MKB GK MRS TT HW GRW SD. Acquired funding: GRW SD.

References

1.   Lai MM, Cavanagh D. The molecular biology of coronaviruses. Adv Virus Res. 1997;48:1-
100.
2.   Masters PS. The molecular biology of coronaviruses. Adv Virus Res. 2006;66:193-292.
3.   Cherry JD, Krogstad P. SARS: the first pandemic of the 21st century. Pediatr Res.
2004;56(1):1-5.
4.   de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, et al. Middle East
respiratory syndrome coronavirus (MERS-CoV): announcement of the Coronavirus Study Group.
J Virol. 2013;87(14):7790-2.
5.   Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated
with human respiratory disease in China. Nature. 2020;579(7798):265-9.
6.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak
associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3.
7.   de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights
into emerging coronaviruses. Nat Rev Microbiol. 2016;14(8):523-34.
8.   Shulla A, Heald-Sargent T, Subramanya G, Zhao J, Perlman S, Gallagher T. A
transmembrane serine protease is linked to the severe acute respiratory syndrome coronavirus
receptor and activates virus entry. J Virol. 2011;85(2):873-82.
9.   Millet JK, Whittaker GR. Host cell proteases: Critical determinants of coronavirus
tropism and pathogenesis. Virus Res. 2015;202:120-34.
10.   Stevens CS, Oguntuyo KY, Lee B. Proteases and variants: context matters for SARS-CoV-2
entry assays. Curr Opin Virol. 2021;50:49-58.
11.   Matsuyama S, Ujike M, Morikawa S, Tashiro M, Taguchi F. Protease-mediated
enhancement of severe acute respiratory syndrome coronavirus infection. Proc Natl Acad Sci U
S A. 2005;102(35):12543-7.
12.   Tang T, Bidon M, Jaimes JA, Whittaker GR, Daniel S. Coronavirus membrane fusion
mechanism offers a potential target for antiviral development. Antiviral Res. 2020;178:104792.
13.   Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike protein via
sequential proteolytic cleavage at two distinct sites. Proc Natl Acad Sci U S A.
2009;106(14):5871-6.
14.   Bosch BJ, Bartelink W, Rottier PJ. Cathepsin L functionally cleaves the severe acute
respiratory syndrome coronavirus class I fusion protein upstream of rather than adjacent to the
fusion peptide. J Virol. 2008;82(17):8887-90.
15.   Jaimes JA, Andre NM, Chappie JS, Millet JK, Whittaker GR. Phylogenetic Analysis and
Structural Modeling of SARS-CoV-2 Spike Protein Reveals an Evolutionary Distinct and
Proteolytically Sensitive Activation Loop. J Mol Biol. 2020;432(10):3309-25.

16.   Madu IG, Roth SL, Belouzard S, Whittaker GR. Characterization of a highly conserved
domain within the severe acute respiratory syndrome coronavirus spike protein S2 domain with
characteristics of a viral fusion peptide. J Virol. 2009;83(15):7411-21.

17.   Lai AL, Millet JK, Daniel S, Freed JH, Whittaker GR. The SARS-CoV Fusion Peptide Forms
an Extended Bipartite Fusion Platform that Perturbs Membrane Order in a Calcium-Dependent
Manner. J Mol Biol. 2017;429(24):3875-92.

18.   Madu IG, Belouzard S, Whittaker GR. SARS-coronavirus spike S2 domain flanked by
cysteine residues C822 and C833 is important for activation of membrane fusion. Virology.
2009;393(2):265-71.

19.   Lai AL, Freed JH. SARS-CoV-2 Fusion Peptide has a Greater Membrane Perturbating
Effect than SARS-CoV with Highly Specific Dependence on Ca(2). J Mol Biol.
2021;433(10):166946.

20.   Khelashvili G, Plante A, Doktorova M, Weinstein H. Ca(2+)-dependent mechanism of
membrane insertion and destabilization by the SARS-CoV-2 fusion peptide. Biophys J.
2021;120(6):1105-19.

21.   Lai AL, Freed JH. Negatively charged residues in the membrane ordering activity of SARS-
CoV-1 and -2 fusion peptides. Biophys J. 2022;121(2):207-27.

22.   Gui M, Song W, Zhou H, Xu J, Chen S, Xiang Y, et al. Cryo-electron microscopy structures
of the SARS-CoV spike glycoprotein reveal a prerequisite conformational state for receptor
binding. Cell Res. 2017;27(1):119-29.

23.   Wang H, Yang P, Liu K, Guo F, Zhang Y, Zhang G, et al. SARS coronavirus entry into host
cells through a novel clathrin- and caveolae-independent endocytic pathway. Cell Res.
2008;18(2):290-301.

24.   Millet JK, Tang T, Nathan L, Jaimes JA, Hsu HL, Daniel S, et al. Production of Pseudotyped
Particles to Study Highly Pathogenic Coronaviruses in a Biosafety Level 2 Setting. J Vis Exp.
2019(145).

25.   Straus MR, Tang T, Lai AL, Flegel A, Bidon M, Freed JH, et al. Ca(2+) Ions Promote Fusion
of Middle East Respiratory Syndrome Coronavirus with Host Cells and Increase Infectivity. J
Virol. 2020;94(13).

26.   Tang T, Jaimes JA, Bidon MK, Straus MR, Daniel S, Whittaker GR. Proteolytic Activation
of SARS-CoV-2 Spike at the S1/S2 Boundary: Potential Role of Proteases beyond Furin. ACS
Infect Dis. 2021;7(2):264-72.

27.   Singh P, Mukherji S, Basak S, Hoffmann M, Das DK. Dynamic Ca(2+) sensitivity stimulates
the evolved SARS-CoV-2 spike strain-mediated membrane fusion for enhanced entry. Cell Rep.
2022;39(3):110694.

28.   Sultan F, Ahuja K, Motiani RK. Potential of targeting host cell calcium dynamics to curtail
SARS-CoV-2 infection and COVID-19 pathogenesis. Cell Calcium. 2022;106:102637.

29.   Dakal TC. SARS-CoV-2 attachment to host cells is possibly mediated via RGD-integrin
interaction in a calcium-dependent manner and suggests pulmonary EDTA chelation therapy as
a novel treatment for COVID 19. Immunobiology. 2021;226(1):152021.

30.   Lambert DW, Clarke NE, Hooper NM, Turner AJ. Calmodulin interacts with angiotensin-
converting enzyme-2 (ACE2) and inhibits shedding of its ectodomain. FEBS Lett.
2008;582(2):385-90.

31.   Lai ZW, Lew RA, Yarski MA, Mu FT, Andrews RK, Smith AI. The identification of a
calmodulin-binding domain within the cytoplasmic tail of angiotensin-converting enzyme-2.
Endocrinology. 2009;150(5):2376-81.
32.   Garcia-Escobar A, Vera-Vera S, Jurado-Roman A, Jimenez-Valero S, Galeote G, Moreno
R. Calcium Signaling Pathway Is Involved in the Shedding of ACE2 Catalytic Ectodomain: New
Insights for Clinical and Therapeutic Applications of ACE2 for COVID-19. Biomolecules.
2022;12(1).
33.   Lai AL, Freed JH. SARS-CoV-2 Fusion Peptide has a Greater Membrane Perturbating
Effect than SARS-CoV with Highly Specific Dependence on Ca2+. Journal of Molecular Biology.
2021;433(10):166946.
34.   Khelashvili G, Plante A, Doktorova M, Weinstein H. Ca^{2+}-dependent mechanism of
membrane insertion and destabilization by the SARS-CoV-2 fusion peptide. Biophysical Journal.
2021;120(6):1105-19.
35.   Koppisetti RK, Fulcher YG, Van Doren SR. Fusion Peptide of SARS-CoV-2 Spike
Rearranges into a Wedge Inserted in Bilayered Micelles. J Am Chem Soc. 2021;143(33):13205-
11.
36.   Schaefer SL, Jung H, Hummer G. Binding of SARS-CoV-2 Fusion Peptide to Host
Endosome and Plasma Membrane. J Phys Chem B. 2021;125(28):7732-41.
37.   Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, Jr., et al. Distinct conformational
states of SARS-CoV-2 spike protein. Science. 2020;369(6511):1586-92.
38.   Straus MR, Bidon MK, Tang T, Jaimes JA, Whittaker GR, Daniel S. Inhibitors of L-Type
Calcium Channels Show Therapeutic Potential for Treating SARS-CoV-2 Infections by Preventing
Virus Entry and Spread. ACS Infect Dis. 2021;7(10):2807-15.
39.   Crespi B, Alcock J. Conflicts over calcium and the treatment of COVID-19. Evol Med
Public Health. 2021;9(1):149-56.
40.   Danta CC. Calcium Channel Blockers: A Possible Potential Therapeutic Strategy for the
Treatment of Alzheimer's Dementia Patients with SARS-CoV-2 Infection. ACS Chem Neurosci.
2020;11(15):2145-8.
41.   Straus MR, Bidon M, Tang T, Whittaker GR, Daniel S. FDA approved calcium channel
blockers inhibit SARS-CoV-2 infectivity in epithelial lung cells. bioRxiv. 2020:2020.07.21.214577.
42.   Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Tracking
Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus.
Cell. 2020;182(4):812-27 e19.
43.   Li Q, Wu J, Nie J, Zhang L, Hao H, Liu S, et al. The Impact of Mutations in SARS-CoV-2
Spike on Viral Infectivity and Antigenicity. Cell. 2020;182(5):1284-94 e9.
44.   Olsson MH, Sondergaard CR, Rostkowski M, Jensen JH. PROPKA3: Consistent Treatment
of Internal and Surface Residues in Empirical pKa Predictions. J Chem Theory Comput.
2011;7(2):525-37.
45.   Olsson MHM, Søndergaard CR, Rostkowski M, Jensen JH. PROPKA3: Consistent
Treatment of Internal and Surface Residues in Empirical pKa Predictions. Journal of Chemical
Theory and Computation. 2011;7(2):525-37.
46.   Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. J Mol Graph.
1996;14(1):33-8, 27-8.

47.   Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, et al. Scalable molecular
dynamics with NAMD. J Comput Chem. 2005;26(16):1781-802.
48.   Essmann U, Perera L, Berkowitz ML, Darden T, Lee H, Pedersen LG. A smooth particle
mesh Ewald method. The Journal of Chemical Physics. 1995;103(19):8577-93.
49.   Evans DJ, Holian BL. The Nose−Hoover thermostat. The Journal of Chemical Physics.
1985;83(8):4069-74.
50.   Eastman P, Swails J, Chodera JD, McGibbon RT, Zhao Y, Beauchamp KA, et al. OpenMM
7: Rapid development of high performance algorithms for molecular dynamics. PLoS Comput
Biol. 2017;13(7):e1005659.
51.   Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, et al. CHARMM36m: an
improved force field for folded and intrinsically disordered proteins. Nature Methods.
2017;14(1):71-3.
52.   Venable RM, Luo Y, Gawrisch K, Roux B, Pastor RW. Simulations of anionic lipid
membranes: development of interaction-specific ion parameters and validation using NMR
data. J Phys Chem B. 2013;117(35):10183-92.
854