The interaction of calcium ions with specific residues in the SARS-CoV

fusion peptide and the regulation of viral infectivity

Miya K. Bidon[1±], George Khelashvili[3±], Marco R. Straus[2], Tiffany Tang[1], Juliana Debrito Carten[1],

Harel Weinstein[3*], Gary R. Whittaker[2*], and Susan Daniel[1*]

[1]Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University,

Ithaca, NY, USA

[2]Department of Microbiology and Immunology, Cornell University, Ithaca, NY, USA

[3]Department of Physiology & Biophysics, Weill Cornell Medicine, New York, NY, USA

*Corresponding authors:

Susan Daniel: sd386@cornell.edu

Gary Whittaker: grw7@cornell.edu

Harel Weinstein: haw2002@med.cornell.edu

[±]These authors contributed equally.

Abstract

Viral envelope fusion with the host cell membrane is dependent on a specific viral fusion peptide (FP) or loop, which becomes exposed during virus entry to drive the process of membrane fusion. In coronaviruses, the FP is a highly conserved domain that sits in the center of spike protein and in SARS-CoV, is adjacent to the S2' proteolytic cleavage site. This peptide contains a hydrophobic LLF motif, as well as several conserved negatively charged amino acids that interact with Ca2+ ions to promote membrane fusion. In this work we perform a systematic mutagenesis study of the negatively charged amino acids within the SARS-CoV fusion peptide (FP1/FP2) and combine this with molecular dynamics simulations to define the membrane interactions that regulate virus infectivity. We show that the E801/D802 amino acid pair in the SARS-CoV FP is predicted to bind to one Ca^{2+} ion to promote FP-membrane interaction, with a second Ca^{2+} ion likely pairing residue D812 with either E821 or D825. The D812/D821 residue pair promotes membrane interaction, whereas the D821/D825 is inhibitory to membrane insertion. Taken together, our results demonstrate the dynamic nature of the coronavirus FP region that likely facilitates its interactions with and insertion into the host cell membrane.

Author Summary

Coronaviruses have reemerged as a highly pathogenic virus family through the rise of SARS-CoV, MERS-CoV, and more recently, SARS-CoV-2. As more transmissible variants of SARS-CoV-2 arise, it is imperative that we understand the mechanisms of CoV viral entry to enable the development of effective therapeutics. Recent reviews have suggested the repurposing of FDA-approved calcium channel blockers to treat infection by coronaviruses; however, calcium's method of

action on viral-host cell fusion events is unknown. We have found that increased calcium

availability leads to increased viral infection across the CoV family, suggesting that calcium is

involved in mediating the interaction between the viral fusion peptide and the host cell

membrane. As such, we hypothesize that the highly conserved fusion peptide interacts directly

with calcium and this interaction is required for viral entry and infection. Through mutagenesis

studies of specific negatively charged residues in the fusion peptide, we have identified residues

that impact viral infectivity. We have also compared the infectivity of wild-type and mutant CoV

pseudoparticles in calcium-rich or -depleted environments using chelating drugs. Our data

mirrors the residue coordination observed SARS-CoV-2, as both between SARS-CoV and SARS-

CoV-2 FPs bind to two calcium ions. These results demonstrate the importance of Ca^{2+} for CoV FP

function during viral entry and opens the possibility of utilizing FDA-approved calcium-blocking

drugs as a treatment for COVID-19.

Introduction

Coronaviruses (CoVs) are a diverse family of enveloped, positive-sense, single-stranded RNA

viruses responsible for causing respiratory and enteric diseases across a wide range of species [1,

2]. Three pathogenic human coronaviruses have emerged during the early 21st century: in 2002,

severe acute respiratory syndrome (SARS-CoV) [3]; in 2012, Middle East respiratory syndrome

(MERS-CoV) [4]; and presently, SARS-CoV-2 [5, 6]. The World Health Organization has listed

coronaviruses as one of the top five emerging pathogens likely to cause major epidemics due to

the minimal countermeasures that exist [7-9]. While vaccines are one strategy that has proven

successful during this SARS-CoV-2 pandemic, there is an outstanding need for antiviral drugs as

a necessary parallel line of defense. To develop new antivirals, we need to understand the

fundamental biology of the virus and its host interactions. From such knowledge we can

therapeutically target well-conserved parts of the viral life cycle in order to confer the broadest

protection. Viral entry is indeed a conserved function [10, 11], with the coronavirus spike (S)

protein controlling virus entry through its interaction with the host cell plasma membrane.

Examining the entry functions of the S protein can point to vulnerable targets that can be

exploited for drug intervention.

The CoV S protein is composed of the S1 and S2 domains, which are responsible for

facilitating receptor binding and membrane fusion, respectively [10, 12, 13]. CoVs utilize two

different pathways to enter host cells: an "early" pathway, often referred to as the plasma

membrane route, and a "late" pathway that follows the canonical endosomal pathway [10]. The

pathway used by the virus is dependent on the host cell type, and more specifically, on the local

protease environment [14]. The current thinking is that SARS-CoV, MERS-CoV and SARS-CoV-2

enter host cells via the early pathway when transmembrane-bound proteases (TTSPs), such as

the TMPRSS2, are present. In cell lines lacking TTSPs, the virus enters cells via the endosomal

route where it interacts with lysosomal proteases (*e.g.* cathepsin L) [10, 15, 16]. This pathway

flexibility arises from the requirement for protease cleavage to prime the S protein for

subsequent conformational changes needed to carry out membrane fusion. Two possible

cleavage sites reside within the S protein: the S1/S2 site and the S2' site [17]. The first site is

located at the boundary of the S1 and S2 domains and is typically cleaved by furin during virus

assembly. It is this cleavage by furin that separates the binding domain from the fusion

machinery. The S2' prime site is located next to the N-terminus of the conserved fusion peptide

(FP) segment that is liberated upon cleavage [18]. S2' cleavage allows the S2 domain to undergo

a large conformational change that positions the FP for insertion into the host cell membrane

[10]. FP membrane insertion is a key step in commencing the process of merging the host cell

membrane with the viral envelope, resulting in the transfer of the viral genome into the host cell.

Given the importance of the FP in initiating membrane fusion, it is not surprising to find

significant conservation of the amino acid sequence in this region of the S protein across many

CoVs [19]. Due to its high degree of conservation, the FP stands out as a potential antiviral target.

Thus, work carried out by our team and others has focused on understanding the structure and

function of the FP. This work has identified conserved residues that interact with calcium ions

(Ca^{2+}) and mediate the interactions between the host cell membrane and the FP that lead to viral

infection [19-23].

In our previous studies, we showed that for both SARS-CoV and MERS-CoV, calcium depletion

in cell culture leads to a significant drop in viral infection [19, 20]. We connected this decrease in

infectivity to a defect in membrane fusion, observing that syncytia formation during cell-to-cell

fusion between S- and ACE2 receptor-expressing VeroE6 cells also drops in calcium-depleted

media. We then sought to connect this observed impact on membrane fusion with the molecular

scale features of S that may interact directly with calcium ions. Within the S2 domain of S, the

FP sequence contains highly conserved charged amino acids that flank the hydrophobic residues

LLF, which interact with the host membrane upon FP insertion [19]. We posited that these

charged residues function as sites for calcium ion binding that may stabilize the local structure of

the FP and position it for optimal membrane insertion. Through a series of biophysical and

biochemical techniques, our group confirmed that calcium interacts directly with the FP and that

this interaction can be linked to infectivity. Using isothermal calorimetry, we determined the

stoichiometry of the interaction between the FP and calcium ions to be two calcium ions for SARS-

CoV [19, 22] and one calcium ion for MERS-CoV [20]. From circular dichroism (CD) measurements,

we observed that in the presence of calcium and membranes, the FP adopts a conformation with

a higher degree of alpha helicity. From electron spin resonance spectroscopy (ESR) results we

found lipid ordering increased upon FP insertion, which can create favorable conditions for

membrane fusion; we also observed greater lipid ordering in the presence of Ca^{2+} [19, 24]. Recent

findings show that calcium is necessary to promote viral entry across multiple coronaviruses,

including SARS-CoV, MERS-CoV, and SARS-CoV-2 [22, 23]. Taken together, these results revealed

that the FP requires calcium to adopt the proper structure needed for membrane insertion.

Despite this initial model for the interaction of calcium with the negatively charged residues of

the SARS-CoV FP [19], the field still lacks a detailed structural understanding of how Ca^{2+} ions

stabilize the FP for membrane insertion.

Here, we present specific mechanistic insight gathered from mutagenesis studies and

computational simulations of the SARS-CoV FP and calcium. This work predicts the preferred

modes of calcium binding to the acidic residues of the SARS-CoV FP and how those modes likely

regulate membrane interactions and subsequent infectivity. We first examined the effects of

mutating individual and multiple charged residues in the SARS-CoV FP on cell fusion and viral

infectivity in the presence or absence of calcium. This was done to identify FP charged residues

that likely interact with calcium during viral entry. We then used corresponding molecular

dynamics simulations of the system to interpret our mutagenesis results structurally. From these

simulations we identified the structures and probabilities of the FP's modes of calcium binding in

1:1 and 2:1 stoichiometries, and predicted the structure of the SARS-CoV FP that is best suited

for membrane penetration.

Results

Generation and characterization of the SARS-CoV wild-type and mutated fusion peptides

Various structural conformations of the SARS-CoV spike monomer have been solved using Cryo-

EM [26]. From these structures, it is apparent that the fusion peptide contains an N-terminal

alpha helix, with the hydrophobic LLF residues, and a C-terminal disordered loop known to

participate in cysteine-mediated intramolecular disulfide bonding **(Fig 1A).** To begin to identify

the SARS-CoV FP's modes of calcium binding, we performed charged-to-alanine substitutions of

six highly conserved residues within the FP sequence (**Fig 1B**). We first made single amino acid

(aa) substitutions (E801A, D802A, D812A, E821A, D825A, D830A) in the wild-type SARS-CoV FP,

which was used as a control. The majority of the negatively charged residues that we mutated

are highly conserved across SARS-CoV, MERS-CoV, and SARS-CoV-2 (**Fig 1C**).

The wild-type and mutant forms of the SARS-CoV S protein were then cloned into

standard pcDNA3.1 expression vectors to facilitate their transient expression in HEK293T cells.

Following confirmation of high transfection efficiency in our cells, S protein synthesis, trafficking

to the cells' surface, and proteolytic cleavage were all assessed. To assay proteolytic cleavage of

the WT and mutant S proteins, TPCK-trypsin was added to the cells' surface 24 hours after

transfection and the S proteins were then isolated by cell-surface biotinylation. Biotinylated

proteins were retrieved following cell lysis using streptavidin beads, resolved by gel

electrophoresis and immunoblotted using a SARS-CoV S polyclonal antibody. The resulting

immunoblots displayed full length, uncleaved (S_0), and cleaved S2 subunit (S_2) SARS-CoV S species

migrating at 180kDa and 80kDa, respectively (**Fig 2A**). SARS-CoV S mutants containing the single

mutations D802A, D812A, E821A, D825A, and D830A have comparable steady-state levels of full-

length protein in comparison to the wild-type protein, indicating that these mutations in the FP

did not impair the synthesis or trafficking of S (**Fig 2A**). Additionally, proteolytic cleavage of the

D802A, D812A, E821A, D825A, and D830A mutants occurred following treatment with TPCK-

trypsin, indicating that these mutants were able to be primed for downstream fusion evens. We

observed a higher molecular weight species running above the full-length S protein (**Fig 2**, (Higher

order S)). We have determined that heating samples at 95°C with 5mM DTT for 10 minutes

resolves this higher molecular weight band (data not shown), indicating that this species likely

results from spike-based protein-protein interactions.

We were unable to detect steady-state levels of the E801A mutant on the spike

immunoblots. To further probe the nature of this residue's importance, we substituted in the

larger, nonpolar methionine (E801M), the polar and uncharged glutamine (E801Q), positively

charged lysine (E801K) or negatively charged aspartic acid (E801D). None of these mutations,

including the charge mimetic E801D, restored the steady-state levels of the S protein to that of

wild-type, indicating the specific requirement of glutamic acid in this region of the FP (**Fig 2B**).

Analysis of the SARS-CoV FP single point mutations' effect on syncytia formation

To functionally test the ability of S protein mutants to induce cell-cell fusion, we

performed a syncytia assay. To do this we transiently expressed the WT and mutated S proteins

in VeroE6 cells, which are kidney epithelial cells that express the ACE2 receptor [26]. 24 hours

following transfection, we induced cell-cell fusion by treating the cells with trypsin to cleave the

FP at the S2' site (R797). Syncytia formation was visualized by immunofluorescence using the

fluorescently labeled SARS-CoV S antibody and DAPI stain to identify multinucleated S-expressing

cells that had fused. Syncytia were quantified by counting every group of fused cells that had a

minimum of 4 nuclei. As expected, VeroE6 cells expressing the WT S protein readily formed

syncytia (**Fig 3A and B**). Conversely, cells expressing the D802A, D812A, E821A, D825A, and

D830A mutants all exhibited very few syncytia, indicating a defect in cell-cell fusion (**Fig 3A and**

B; Fig S1).

Analysis of the SARS-CoV FP mutants' effect on viral infectivity

SARS-CoV is designated a Risk Group 3 Select Agent that requires a specialized biosafety level

3 (BSL-3) setting for experiments. Pseudoparticles can be used as safe surrogates of live SARS-

CoV to enable experimentation in BSL-2 level conditions [27]. To generate SARS-CoV

pseudoparticles in HEK293T cells, we used a three-plasmid co-transfection system, with plasmids

encoding: (1) the full length CoV surface S protein, (2) the murine leukemia virus (MLV) core

proteins *gag* and *pol*, or (3) a firefly luciferase reporter gene containing the MLV-MJ RNA

packaging signal and flanked by long terminal repeat (LTR) sequences [27]. Upon successful

particle fusion with a host cell membrane, the luciferase reporter RNA transcript is reverse

transcribed, integrated into the cell genome and expressed, enabling a measurable readout of

pseudoparticle entry. Our previous studies have shown that this approach works well for

assessing the infectivity of MERS-CoV S [20], SARS-CoV S [19, 27], and SARS-CoV-2 S [28].

Incorporation of the WT and mutant S proteins into the SARS-CoV pseudoparticles was

assessed using immunoblot analysis. Pseudoparticles with WT or the mutant S proteins D802A,

D812A, E821A, D825A, and D830A show comparable levels of S protein incorporation (**Fig 4A**).

As expected, we did not detect the E801A mutant in the SARS-CoV pseudoparticles, likely due to

its low cellular expression or synthesis.

After confirming the incorporation of the wild-type and mutant spike proteins into the

SARS-CoV pseudoparticles, we tested the infectivity of the pseudoparticles. As we have

demonstrated previously, exposure of VeroE6 cells to WT SARS CoV pseudoparticles results in a

robust luminescent signal 72 hours post infection, indicating successful viral entry **(Fig 4B)**.

E821A- and D825A-containing pseudoparticles did not exhibit a significant change in infectivity

compared to wild-type particles, suggesting these individual residues do not have a major effect

on FP function when calcium is present (**Fig 4B**). In contrast, the D802A, D812A, and D830A

mutant pseudoparticles all showed a pronounced decrease in infectivity, with the D812A and

D830A mutations resulting in essentially noninfectious particles. These results indicate that

mutations in these three highly conserved negatively charged residues in the S protein's FP

significantly decrease SARS-CoV pseudoparticle infectivity in calcium-containing conditions.

We next tested the effect of depleting extracellular calcium on the pseudoparticles' ability to

infect VeroE6 cells using the chelator ethylene glycol-bis (ß-aminoethyl ether)-*N,N,N'N'*-

tetraacetic acid (EGTA). We have previously studied the effects of calcium on viral infectivity and

adopted similar calcium depletion methods in this study [19, 20]. Briefly, we treated VeroE6 cells

with 50 µM EGTA prior to and during infection. Following incubation with the chelating agent,

VeroE6 cells were inoculated with WT or mutant S protein-containing pseudoparticles and

incubated for 72 hours. Cells were then lysed and luminescence was quantified as a measure of

pseudoparticle infectivity (**Fig 4B**). In support of our previous work [19], depletion of extracellular

calcium had a marked effect on WT pseudoparticle infectivity, decreasing it to approximately half

that of cells infected with WT pseudoparticles in the absence of EGTA (**Fig 4B**). Cells exposed to

pseudoparticles containing either D802A or the D825A mutations also exhibited a further drop

in infectivity, with the D825A mutation having the greatest additional reduction in infectivity (**Fig**

4B). The depletion of extracellular calcium had an additive effect on the decrease in infectivity in

the context of specific FP mutations (D802A, D825A). These results suggest that these negatively

charged residues in the FP are affected either directly or indirectly by loss of extracellular calcium.

We then examined the effect of intracellular calcium depletion on pseudoparticle infectivity

by using the cell-permeable calcium chelator 1,2-bis(2-aminophenoxy)ethane- *N,N,N'N'-*

tetraacetic acid tetrakis (BAPTA-AM). We have previously optimized the concentrations of

BAPTA-AM so as not to significantly impact cell viability [19, 20]. VeroE6 cells were treated with

50 μM BAPTA-AM briefly prior to and during pseudoparticle infection, then harvested and

assayed for luminescence (**Fig 4C**). Interestingly, depletion of intracellular calcium rendered the

WT pseudoparticles noninfectious; similar results were observed across the WT and all mutant

pseudoparticles tested; none of the pseudoparticles were infectious. In general, these data

suggest that depletion of intracellular calcium exerts a more severe effect on pseudoparticle

infectivity and point to the endosomal pathway as the primary route of pseudoparticle entry.

From our infectivity data we deduced that the FP likely uses multiple negatively charged

residues to bind to multiple calcium ions. This is supported by the further decrease in infectivity

of Spike E802A- and D825A-containing pseudoparticles when extracellular calcium is depleted.

We hypothesize that mutating those residues individually does not lead to a complete loss in

pseudoparticle infectivity, since other charged residues in the FP can compensate by binding

calcium; however, removal of calcium from the medium mimics the loss of those additional

residues, leading to the FP's inability to bind calcium and a loss in infectivity.

Multiple negatively charged residues mediate SARS-CoV fusion peptide's calcium binding

To test the hypothesis that the FP utilizes alternative residues to mediate calcium binding,

we created pairwise and triple residue substitutions within the S protein's FP. The FP contains

two defined regions: the N-terminal region starting just upstream from the S2' cleavage site

(aa798-819), termed FP1, and the downstream region that includes a critical disulfide bond

(aa820-837), termed FP2 [18, 29]. Lai *et al.* has reported a stoichiometry of two calcium ions per

single SARS-CoV FP, therefore we predict that FP1 and FP2 regions each contain residues that can

coordinate calcium binding [19]. In FP1, the alanine substitutions of residues D802, and D812

rendered their pseudoparticles non-infectious; in FP2, the D825A and D830A substitution also

led to non-infectious pseudoparticles. To investigate if individual substitutions of these three

residues changed the mode of calcium binding, we created a complete set of their double

mutants   (E821A/D825A,   E821A/D830A,   D825A/D830A),   and   a   triple   mutant

(E821A/D825A/D830A) (**Fig 5A**).

We first confirmed that these mutants were synthesized, trafficked to the plasma

membrane, and were able to be cleaved by trypsin, as we had done with the single mutants (**Fig**

5B). We also tested the functionality of these S mutants using the previously described syncytia

assay. VeroE6 cells transiently expressing either a double mutant (E821A/D825A, E821A/D830A,

D825A/D830A) or the triple mutant (E821A/D825A/D830A) all displayed fewer syncytia

compared to WT S-expressing cells (**Fig 5C and D)**.

We then generated pseudoparticles containing the double and triple mutants and

assayed the incorporation of the various mutants into the particles by immunoblotting for the

spike protein. The double mutant E821/D830A- and D825A/D830A-containing pseudoparticles

incorporated roughly equal amounts the S protein, while the triple mutant pseudoparticles

(E821A/D825A/D830A) showed slightly reduced protein levels (**Fig 5E**). The E821A/D825A mutant

particles exhibited significantly decreased protein levels indicating that although this mutant

could be expressed and trafficked to the plasma membrane, its incorporation into

pseudoparticles was diminished. We acknowledge the general low protein levels of the double

and triple S mutants in our pseudoparticles as being a potential confounding variable.

Nevertheless, we infected VeroE6 cells with the WT and mutant S protein pseudoparticles to

investigate if multiple mutations in negatively charged residues of the FP caused a more

pronounced decrease in infectivity. As was previously demonstrated, WT pseudoparticles are

able to infect VeroE6 cells, with subsequent depletion of extra- or intra-cellular calcium causing

a partial and then complete loss of infectivity, respectively **(Fig 5F and G).** Infectivity was

dramatically reduced for all mutants in comparison to WT pseudoparticles, irrespective of extra-

and intracellular calcium depletion. (**Fig 5F and G**). These data support the hypothesis that the

SARS-CoV FP contains multiple sites of calcium binding, which when mutated result in a

nonfunctional spike protein irrespective of calcium levels.

Molecular Dynamics simulations identify the modes of the SARS-CoV FP's Ca^{2+} binding

The effects that the various S mutants have on SARS CoV pseudoparticle infectivity suggests

that the FP contains multiple sites of Ca^{2+} binding that are required for viral entry. This confirmed

requirement of $Ca^{2+}$for FP function prompted an analysis of the predicted modes of Ca^{2+} binding

in the wild-type and mutant SARS-CoV FPs. To this end, we carried out extensive atomistic

molecular dynamics (MD) simulations of all the SARS-CoV FP constructs studied here (see

Methods section). To monitor the spontaneous binding of Ca^{2+} to the FP, we collected 18

independent MD trajectories of 640ns in length for each construct. Following the analysis

protocols described recently for SARS-CoV-2 FP simulations [23], the various modes of

interactions between the SARS-CoV FP and Ca^{2+} ions were assessed in the trajectories by

monitoring: i) the distances between the Ca^{2+} ions in solution and the side chains of all acidic

residues in the peptide; and ii) the pairwise distances between the side chains of all acidic

residues. Summarized in **Fig S2** are the observed events of simultaneous association of two Ca^{2+}

ions with various pairs of FP residues in the individual trajectories of the WT and the mutant

systems (see *red* and *blue* rectangles). The combined statistics for each construct (*i.e.,* the total

number of binding events for different pairs of residues) are summarized in **Fig 6**.

These results show the modes of Ca^{2+} coordination in the WT protein, identifying the most

frequent modes involving residue pairs E821/D825 (5 out of 18); E801/D802 and E801/D830

(4/18 each); and D812/E821 (2/18). Notably, a similar pattern of coordination preference was

observed in our recent computational studies of Ca^{2+} association with SARS-CoV-2 FP [23].

Moreover, simultaneous binding of Ca^{2+} ions to the SARS-CoV-2 FP residue pairs equivalent to

E801/D802 and D812/E821 produced the peptide conformations prone to membrane

penetration. In contrast, conformations that stabilized Ca^{2+} binding to residues equivalent to the

E821/D825 pair did not enable sustained bilayer insertion.

From these MD simulations of the SARS-CoV FP constructs, we were able to predict the

phenotypes for each construct (active or inactive). The mode of Ca^{2+} ion association predicted to

be inhibitory for membrane insertion (E821-D825) is shown in *red*, and the modes of Ca^{2+}

association predicted to facilitate membrane insertion are depicted in *green* boxes (**Fig 6**). In

those constructs with single mutations that were predicted to not have fusion activity (i.e.,

E801A, D802A, D812A, and D830A) we showed that only the Ca^{2+}-coordination mode involving

the E821/D825 pair persists, but not the D812/E821 pair of residues (**Fig 6**). Conversely, in the

SARS-CoV FP single mutants that maintained WT-like fusion activity (E821A and D825A), we did

not observe the E821/D825 residues predicted to participate in calcium binding. In these

function-preserving mutants we identified additional modes of Ca^{2+} binding that are enhanced in

comparison to the WT system: E801/D802 in mutant E821A, or E801/D830 in mutant D825A. For

the SARS-CoV FP constructs with multiple mutations, our MD trajectories revealed an overall

reduced Ca^{2+} binding ability (**Fig 6** and **Fig S3**), consistent with our experimental findings that (i)

these constructs are severely defective in the cell-cell fusion (**Fig 5C and D**) and infectivity assays,

and (ii) their function does not depend on the levels of either extracellular or intracellular Ca^{2+}

(**Fig 5F and G**).

SARS-CoV FP propensity for membrane insertion is regulated by modes of Ca^{2+} binding.

Overall, the above computational results reveal that the Ca^{2+} binding patterns of SARS-

CoV FP are very similar to those of SARS-CoV-2 FP. On this basis, membrane insertion of the SARS-

CoV FP could be expected to be enhanced by the modes of Ca^{2+} binding involving the E801/D802

and D812/E821 pairs of residues, and to be reduced by the ones involving the E821/D825 pair.

To test this premise, we carried out MD simulations of two models of the WT SARS-CoV FP

spontaneously associating with the lipid membrane (see **Methods**). In one, the peptide was

interacting with 2 Ca^{2+} ions at the E801/D802 and D812/E821 sites (Model 1), and in the second,

2 Ca^{2+} ions were bound to the E801/D802 and E821/D825 pairs (Model 2). Each structure was

simulated in 36 independent replicates, each run for ~0.9-1.0μs (see **Methods**).

Analysis of these trajectories revealed that, indeed, Model 1 extensively penetrated the

membrane, while the membrane insertion of the Model 2 was negligible. This can be seen from

the plots presented in **Fig 7** comparing frequencies of membrane insertion for each SARS-CoV FP

residue in the simulations of Model 1 and Model 2 structures. The membrane insertion was

quantified by monitoring the z-coordinate of the C_α atom of each residue in the peptide. A residue

was inserted into the membrane if the z-distance between its C_α atom and the second carbon

atom in the tail of a POPC lipid (atom C22 in CHARMM36 notation) was <5Å. As shown in **Fig 7A**,

in Model 1, the N-terminal FP1 segment of the fusion peptide shows strong propensity for bilayer

insertion, while in Model 2 the insertion is minimal (**Fig 7B**).

The detailed analysis of the individual trajectories in the Model 1 set revealed two distinct

modes of bilayer penetration for SARS-CoV FP, similar to our findings for the SARS-CoV-2 FP. Thus,

the Model 1 construct penetrates the bilayer either with its N-terminal LLF motif (**Fig 7C**) or with

the more centrally located hydrophobic F815-M816 segment (**Fig 7D**). Interestingly, the two

insertion modes appear to alternate which Ca^{2+} ion is neighboring the inserted portion. Thus,

when the LLF is inserted, the Ca^{2+} ion associated with the neighboring E801/D802 residues is

bound to the membrane while the other Ca^{2+} binding site (D812/E821) is situated away from the

membrane surface (see snapshot in **Fig 7C**). In case of the F815-M816 insertion, the position of

the Ca^{2+} binding loci with respect to the membrane is reversed – the one associated with the

D812/E821 pair is membrane-bound, while the E801/D802 pair is located farther from the bilayer

(snapshot in **Fig 7D**). We also note that in both cases, the remaining anionic residues in the

peptide (*i.e.* the ones not engaged with the Ca^{2+} ions) are either solvent exposed (D830) or

engaged with electro-neutralizing interactions with neighboring basic residues (D825/R829, **Fig**

7C and D). These results support a mechanistic model in which membrane penetration of the

SARS-CoV FP is significant only for specific modes of Ca^{2+} binding to the peptide, i.e., to the

E801/D802 and D812/E821 pairs of conserved acidic residues. Moreover, Ca^{2+} binding at the

E821/D825 pair is predicted to inhibit membrane insertion.

Discussion

The resurgence of pathogenic human coronaviruses brings about an immediate need for

improving countermeasures to combat these global health threats. Increasing our knowledge of

the mechanisms of viral entry is essential to develop broad-spectrum vaccines and antivirals. Viral

entry is a critical step in the viral life cycle that is mediated by the S viral protein [8, 30, 31]. The

S protein's FP is responsible for mediating viral and host cell membrane fusion. This region is

highly conserved across the *Coronaviridae* viral family and understanding it how it functions may

lead to the development of more broad-acting therapeutics. Previous work has highlighted the

role that cations play in viral fusion, specifically Ca^{2+}. The electrostatic interactions between

cations and negatively charged residues in the fusion peptide promote membrane fusion, leading

to higher infectivity [19-22, 32]. Lai et al. has proposed that two Ca^{2+} ions were needed to interact

with SARS-CoV FP; in each of the subdomains (FP1 and FP2), one Ca^{2+} coordinates two negatively

charged residues to allow the FP to adopt a conformation that promotes membrane insertion

[19].

To interrogate the requirement of the negatively charged residues in the SARS CoV fusion

peptide for its function, we first made single charge-to alanine substitutions in those residues

(**Fig 1**). Following transient expression of these mutants in Vero6 cells, we confirmed that

mutants D802A, D812A, E821A, D825A, and D830A were synthesized, accumulated to levels

comparable to the WT protein, and were trafficked to the plasma membrane (**Fig 2**). We further

confirmed the cell surface localization of these same FP mutants by treating VeroE6 cells

expressing these mutants with trypsin to assess their cleavage, after which we retrieved the FPs

using a cell surface biotinylation assay (**Fig 2**). The majority of the FP mutants assayed exhibited

cleavage following treatment with trypsin, indicating the accessibility of the S1/S2 cleavage site.

The E801A mutant was not detected on our spike immunoblots and further attempts to

understand this occurrence through additional substitutions did not result in the detection of this

mutant. We hypothesize that the absence of glutamate at position 801 in the fusion peptide

causes a loss in protein stability, though further work is needed to determine the significance of

this residue in the fusion peptide.

We next tested the fusion competency of the single FP mutants we had created using a

syncytia assay, which utilizes fusion events in cells transiently expressing the FP as a readout of

FP fusion activity. All of the single FP charged-to-alanine mutants the were detected on the spike

immunoblot (E801A, D802A, D812A, E821A, D825A, and D830A) exhibited a pronounced fusion

defect, as evidenced by the low number of fused VeroE6 cells, or syncytia, that were observed

(**Fig 3**). Taken together, these data suggest that the highly conserved, negatively-charged

residues within the FP individually contribute in a non-redundant manner to the function of the

fusion peptide. However, due to the limitations and variability of the syncytia assay, we chose to

use SARS-CoV2 pseudoparticles to mimic a more in vivo-like system to examine the functionality

of the various FP mutants.   Successful pseudoparticle entry into host cells results in the

integration of the luciferase reporter gene into the cellular genome. Luminescence can therefore

be used as a readout of pseudoparticle infectivity. We first confirmed the incorporation of the

WT and single-charged-to-alanine mutant FPs into the pseudoparticles (**Fig 4A).** Nearly all FP

mutants generated (E801A, D802A, D812A, E821A, D825A, and D830A) were incorporated in the

SARS-CoV2 pseudoparticles; the E801A mutant was not detected.

We then infected VeroE6 cells using the SARS-CoV2 pseudoparticles we had generated

and measured luminescence as a readout of infectivity and a proxy for viral entry. Introduction

of WT pseudoparticles into VeroE6 cells results in a robust luminescence signal, indicating

successful viral entry and fusion competency of the viral particles when calcium levels are

unperturbed (**Fig 4B and C**). Infections with E821A and D825A-containing pseudoparticles at

physiological levels of calcium also resulted in luminescence levels comparable to WT-containing

pseudoparticles, suggesting that these residues are not required for FP function. To the contrary,

pseudoparticles containing the D802A, D812A, or D830A mutations were unable to infect VeroE6

cells, resulting in aa significant drop in luminescence. These data indicate that when intra- and

extra-cellular calcium levels remain unchanged, loss of an individual negative charges at positions

D802, D812, or D830 is sufficient to cause a substantial decrease in infectivity, likely due to a

defect in FP-mediated viral fusion and subsequent viral entry.

We then proceeded to test the requirement of extracellular calcium on the infectivity of

our FP-containing pseudoparticles by treating the cells with EGTA, a calcium-preferring chelator,

prior to infection. Removal of extracellular calcium resulted in a significant drop in the infectivity

of WT FP-containing particles, which is consistent with the known requirement of calcium for

SARS CoV2 viral entry (**Fig 4B**). Interestingly, pseudoparticles containing either the D802A or

D825A mutant showed a further reduction in infectivity when the extracellular calcium was

depleted. This suggests that multiple negatively charged residues in the FP are involved in calcium

binding and while loss of a single negative charge may not be sufficient to completely disrupt

infectivity, multiple "hits" are. Thus, removal of extracellular calcium mimics the loss of additional

electrostatic interactions needed for FP function, resulting in a further decrease in infectivity.

Pseudoparticles containing either the D812A or D830A FP mutant were essentially non-infectious

in the presence or absence of extracellular calcium. These results implicate residues D812A and

D830A in FP function; however, their specific roles as they relate to calcium cannot be teased

apart in this system.

We also depleted intracellular calcium levels using the cell permeable calcium chelator

BAPTA-AM, in order to test the contribution of intracellular (endosomal) during pseudoparticle

infectivity.

Following treatment with BAPTA-AM, WT FP-containing pseudoparticles are no longer

infectious (**Fig 4C**), indicating that intracellular calcium, specifically in the endosome, contributes

to pseudoparticle entry (**Fig 4B and C**).   We propose that chelation with EGTA likely removed

the majority of extracellular calcium causing a 50% reduction in infectivity of WT FP-containing

pseudoparticles; however, it is known that SARS-CoV enters cells via two pathways: the plasma

membrane and endosomal pathways. Hence, the partial reduction in viral infectivity in the

presence of EGTA and then the complete loss of infectivity in the presence of BAPTA-AM may

reflect the SARS-CoV pseudoparticles' usage of these two pathways as well. It is important for

us to acknowledge that we cannot rule out the potential impact that depletion of intracellular

calcium may have on the integration, expression, or synthesis of the reporter transgene in the

pseudoparticle. However, we performed cell viability assays to optimize the concentrations of

both chelators used in this study and are confident that treatments with EGTA and BAPTA-AM

did not induce toxicity in the VeroE6 cells.

Given that the results from our infectivity assay suggested the involvement of multiple

calcium-binding residues in the fusion peptide, we then generated double mutants (E821A,

E825A; E821A, D830A; E825A, D830A) and a triple mutant (E821A:E825A;D830A) of the FP (**Fig**

6A). We validated the expression, synthesis, and cell surface localization of these mutants, as

well as their ability to be cleaved by trypsin (**Fig 6B**). As with the single mutants, we observed a

fusion defect in VeroE6 cells transiently expressing these mutant constructs, evidenced by the

few number of syncytia formed in comparison to cells expressing the WT FP (**Fig 6D**). When then

assayed the fusion activity of these mutants using the previously described pseudoparticle

infectivity assay. The double mutants E821A; D830A and E825A; D830A and a triple mutant

E821A; E825A; D830A were all incorporated into the pseudoparticles; the E821A; E825A mutant

was not (**Fig 6C**). If the FP binds calcium using more than one negatively charged residue, then

pseudoparticles containing the double or triple charge-to alanine mutants should exhibit a

further decrease in infectivity. In comparison to WT FP-containing pseudoparticles, the FP double

mutants E821A; D830A and E825A; D830A and the triple mutant E821A;E825A;D830A we assayed

showed a complete loss in infectivity. Because the FP double mutant E821A, E825A was not

incorporated into the pseudoparticles, we could not fully assess the impact of these mutations

on FP function; however, the low number of syncytia observed in cells expressing this mutant

suggests these residues are important for FP function. In summary, these results support the

hypothesis that the SARS-CoV2 fusion peptide requires multiple negatively charged residues to

bind to calcium during viral entry. Specifically, our data implicate FP residues D802 and D825 in

coordinating the FP's interaction with calcium.

To better understand how the various charged residues in the SARS-CoV FP coordinate

calcium binding, we undertook molecular dynamic (MD) simulations with the FP and calcium

ions. The MD simulations of the Ca^{2+}-loaded peptide with the membrane illuminated the way

Ca^{2+} binding to FP binding may affect its function. Exploring the probability of each FP residue

interacting with the membrane in the identified Ca^{2+}-binding modes reveals the preferred mode

of peptide insertion to be with the N-terminal end interacting with a calcium ion coordinated by

residues E801 and D802 (**Fig 7**). The depth of insertion and the determinant role of the Ca^{2+} ion

in facilitating membrane insertion are identical to the findings for the SARS-CoV-2 peptide and

are in agreement with the experimental measurements of SARS-CoV FP membrane insertion by

Lai *et al* [19, 20, 22]. That Ca^{2+} interacts with the same highly conserved, charged residues in the

FPs of SARS-CoV and SARS-CoV-2 is not surprising.

Results from our computational modeling of SARS-CoV FP-membrane interactions in the

presence of Ca^{2+} show that the propensity of the SARS-CoV FP for membrane insertion is dictated

by specific modes of Ca^{2+} binding. The Ca^{2+} binding mode that enables the most sustained

membrane penetration involves the initial association of the peptide with the bilayer through a

Ca^{2+} binding site located near the N-terminus of the peptide. This site (E801/D802) is made

accessible following enzymatic cleavage and is inserted into the membrane via the insertion of

the juxtaposed hydrophobic segment (the LLF motif). In this process, all anionic residues of the

peptide are either engaged with Ca^{2+} ions, with neighboring basic residues, or remain solvent

exposed away from the membrane. Based on the preferred modes of Ca^{2+}-loaded FP interaction

with the membrane, we propose that the binding of Ca^{2+} following S protein cleavage at the S2'

site creates an energetically feasible membrane insertion process (**Fig 7A**) composed of the

following steps: (1) steering of the peptide towards the bilayer surface; (2) exposure of the

hydrophobic layer of the membrane to the conserved N-terminal LLF motif of the FP anchored

by the Ca^{2+} interactions with the phospholipid headgroups; and (3) shielding of the anionic

residues of the peptide by the preferred inserting conformation of the FP to prevent their

negative charge from interfering with membrane insertion. These three components became

evident from the analysis of the MD simulation trajectories presented here for the SARS-CoV FP

and bear a very high similarity to results from computational studies of the SARS-CoV-2 FP

(PMID: 33631204).

Remarkably, in both the SARS-CoV-2 FP simulations [23] and here, we observed a Ca^{2+} binding

mode that inhibits long-lasting membrane insertion. In this mode Ca^{2+} association with the

E821/D825 pair (corresponding to the D839/D843 in SARS-CoV-2 FP) positions the LLF motif near

the negatively charged D812 residue at the membrane surface (**Fig 7B**). This eliminates the

favorable effect of the negative charge screening by the Ca^{2+}, so that the peptide bilayer

encounters are transient and do not lead to penetration by the LLF motif. This mechanistic model,

which highlights the role of the positioning of D812 residue in specific Ca^{2+} binding modes, is

supported by our findings for the SARS-CoV S mutants, as it was previously found to be an

impactful mutation in the SARS-CoV FP [18]. A recent study from Koppisetti *et al* asserts that a

Ca^{2+} may be binding to D843/D849 in SARS-CoV-2 FP (corresponding to D825/D830 in SARS-CoV

FP); however, as shown from the simulations in this study and Khelashvili *et al*, there is a high

likelihood that second Ca^{2+} ion will bind to D812/D821 (corresponding to the D830/D839 in SARS-

CoV-2 FP) for the maximal membrane penetration [23, 25]. Recent NMR studies regarding the

membrane interaction with the fusion peptide still have yet to be fully resolved as current models

use bicelles [25] and micelles [33], which do not completely capture the composition of the

bilayer as was done for the computational modeling shown here; though the model of membrane

insertion proposed by Koppisetti *et al* [25] most aligns with our proposed CoV FP interaction with

bilayer (**Fig 8**). Thus, as shown in **Figs 6 and 7**, mutations that creates a high propensity for the

Ca^{2+} binding mode involving the E821/D825 pair, stabilize peptide conformations that are non-

productive for membrane penetration and are found here to inactivate the FP (**Fig 6**).

Interestingly, though our computational studies did not confirm the role of D830 with Ca^{2+},

our biological studies still reaffirmed its importance in the FP. Current studies on SARS-CoV-2 S

protein classified the disordered portion of the FP in the FP2 domain as the FPPR (fusion peptide

proximal region), which includes the charged residues D839/D843/D848 or E821/D825/D830

equivalent in SARS-CoV FP [23, 34]. This FPPR region was determined to be important, as it binds

to the RBD through the CTD1, and maintains the closed pre-fusion S trimer [34]. This is due to

the tight packing around the disulfide bond reinforced by a bond between K835 and D848 (K817

and D830 in SARS-CoV FP). The equivalent residue in SARS-CoV FP, based on biological studies, is

a critical residue [34]. Mutating D830 removes the binding partner of K817, thereby loosening

the 'knot' of the structured FPPR in maintaining the closed form of the trimeric S protein [34].

Without the reinforcements, we may lose the optimal structural conformation of the S protein

upon receptor binding to expose the S2' cleavage site to promote successful proteolytic cleavage

[30].

More broadly, the E801 and D830 residues are conserved within the CoV family and the D812

residue is conserved within betacoronaviruses [18, 19]. The implication is that calcium

interactions are a conserved mechanism that serves to better position the FP for membrane

insertion. Different coronaviruses exhibit different requirements for calcium; MERS-CoV binds

to one Ca^{2+} ion in its FP1 domain [20], thus, it is important to investigate the role of Ca^{2+} and FP

interactions across the CoV family. The conservation of calcium-binding residues in the FP of

many coronaviruses suggests that the CoV fusion mechanisms can be potential targets for broad-

spectrum antiviral drugs [35-37]. Repurposing FDA-approved calcium channel blocking (CCB)

drugs to inhibit CoV entry, particularly for SARS-CoV-2, is one option worth exploring. Recent

studies have shown that the CCB felodipine is a potential candidate to inhibit SARS-CoV-2 entry

[35]. CCBs can target conserved viral functions, providing a rapid solution to address new and

future SARS-CoV-2 variants. It will be important to identify the mechanisms of CCBs CoV

inhibition, as they may directly inhibit a viral target or indirectly inhibit viral entry by affecting

host cell processes

In this study, we elucidate the relationship between highly conserved residues in the SARS

CoV FP and the critical role they have in coordinating calcium binding to facilitate viral entry.

Interestingly, SARS-CoV-2 variants have arisen as part of Clade 20A that contain a D839G or

D839Y mutation (E821 equivalent in SARS-CoV), with this mutation predicted to affect FP-Ca^{2+}

interactions [38-40]. To date, it is not known if there is any selective advantage to the virus

conferred by this mutation or whether the emergence of these variants simply represents a

founder effect.

Regarding to the role calcium plays during SARS-CoV entry, our data points to the necessity

of intra- and extra-cellular calcium during infection. Loss of extracellular calcium results in a 50%

reduction in PP infectivity, while the loss of intracellular calcium rendered pseudoparticles

noninfectious. Together with the MD simulations, we propose a model of SARS-CoV viral entry

mediated, in part, by calcium. In this model, upon S2' cleavage the spike protein's FP is exposed

and stabilized by binding to 2 calcium ions through electrostatic interactions. We believe that the

residues that likely mediate these essential interactions with calcium are E801/D802 and

D812/E821. Stabilization of the FP is critical for host membrane insertion, given the unstuctured

and flexible FP2 loop in this peptide. We hypothesize that calcium interaction helps stabilize the

FP structure prior to membrane insertion, thus in the absence of a single negatively charged

residue, the FP can compensate through alternative residues to bind to calcium. Overall, more

studies are needed to fully understand the role calcium plays in CoV FP function and its effects

on viral pathogenesis.

562
563   **Materials and Methods**
564

Cells, plasmids, and reagents. Human embryonic kidney 293 (HEK293T) and African green

monkey kidney epithelial (VeroE6) cells were obtained from the American Type Culture

Collection (ATCC, Manassas, VA). Both cell lines were grown in complete DMEM, composed of

Dulbecco's modified Eagle medium (DMEM, CellGro), supplemented with 10% HyClone

FetalClone II (GE) and 10 mM HEPES (CellGro) at 37°C and 5% CO_2. The plasmids used for

generating pseudoparticles were the pCMV-MLV gag-pol murine leukemia virus (MLV) packaging

construct, the pTG-Luc transfer vector encoding the luciferase reporter gene, and the pCAGGS-

VSVG plasmids were provided by Jean Dubuisson (Lille Pasteur Institute, Lille, France) and co-

transfected as previously described [27]. The plasmid encoding the C9-tagged SARS-CoV spike

protein (pcDNA3.1-C9-SARS-CoV S) was provided by Dr. Michael Farzan from the New England

Primate Research Center. Recombinant L-1-tosylamide-2-phenylethyl chloromethyl ketone

(TPCK)-treated trypsin was obtained from Sigma. The calcium chelators EGTA and BAPTA-AM

(acetoxymethyl ester) were obtained from VWR and Tocris, respectively. Both compounds were

diluted in dimethyl sulfoxide (ThermoFisher) prior to use.

Site-directed mutagenesis. Site-directed mutagenesis was performed on the SARS-CoV spike

protein vector, pcDNA3.1-SARS-CoV-S, via the QuikChange Lightning site-directed mutagenesis

kit (Aligent). PCRs and transformations were performed based on the manufacturer's

recommendations. Primers obtained from IDT Technologies were designed using the primer

design tool from Aligent. The primers used to generate the SARS-CoV S mutants can be found in

the supplementary information (**SI, Table S1**). Mutations were confirmed via Sanger Sequencing

at the Cornell University Life Sciences Core Laboratories Center.

Cell Surface Biotinylation Assay. HEK293T cells were seeded at a density of 8×10^5 cells per well

in poly-D-lysine coated 6-well plates. 24 hours later, cells were transfected with 2ug of WT or

mutant SARS-CoV S protein expressing plasmids using polyethylenimine (PEI) (Fisher). To

transfect each well, 2 μg of plasmid DNA and 6 μL of a 1ug/uL solution of PEI were incubated with

50 μL of serum free OptiMEM (Gibco) for 20 minutes. 2 mL of complete DMEM were then added

to this transfection mixture, which was used to replace the cell media. Cells were transfected for

24 hours, washed once with 1X DPBS, and then left untreated or treated with 1mL of 0.8 nM

TPCK-trypsin in DPBS for 10 minutes at 37°C. Cell surface proteins were then biotinylated on ice

by incubating rinsed cells in a biotin-containing buffer (250 ug/mL in PBS) for 20 minutes. The

biotin buffer was then replaced with a quenching solution (50mM glycine in DPBS) and incubated

for 30 minutes on ice. Cells were then lysed for 15 minutes using a lysis buffer containing 0.1%

Triton in 1x TBS with protease inhibitors (cOmplete Protease Inhibitor Cocktail). Lysates were

collected and spun at 13000 rpm for 10 minutes to pellet the insoluble material. The lysates were

then incubated overnight at 4° C with equilibrated steptavidin beads to retrieve the biotinylated

spike proteins. The captured spike proteins were eluted by boiling the beads in 1x LDS sample

buffer in the presence of DTT (NuPAGE) for 10 minutes. Proteins were resolved on 4-12% gradient

Bis-Tris gels (NuPAGE) and transferred to PVDF membranes. The SARS-CoV S protein was

detected using the SARS-CoV S rabbit polyclonal primary antibody (NR-4569, BEI resources) and

Alexa Fluor 488-labeled goat anti-rabbit secondary antibody (Invitrogen). The resulting protein

bands were visualized using a Chemidoc system with Image Lab image capture software (BioRad).

Bands have only been adjusted in contrast and brightness using the Image Lab software.

Cell-cell fusion assay. 3.5×10^5 VeroE6 cells were seeded in 8-well chamber slides (Millipore).

After 24 hours, cells were transfected with a mixture containing 0.75 µL of Turbofect

(Thermofisher), 0.5 µg of S-expressing plasmid, and 11.75 µL OptiMEM for each well. Following

18 hours of transfection, cells were washed with 1X DPBS and trysinized with 0.8 nM TPCK-trypsin

for 5 minutes at 37°C to. Cells were then fixed with 4% paraformaldehyde (PFA) (ThermoFisher)

for 15 minutes and washed three times with DPBS. To permeabilize the cells, 0.1% Triton X-100

was added to each well and cells were incubated for 5 minutes on ice. After three washes with

DPBS, the cells were blocked with 5% normal goat serum for 30 minutes. Cells were again washed

and then labeled with SARS-CoV S rabbit polyclonal antibody (NR-4569, BEI resources), followed

by labeling with Alexa Fluor 488-labeled goat anti-rabbit secondary antibody (Invitrogen). The

cell nuclei were labeled with a DAPI stain present in the mounting media (DAPI Fluoromount-G,

Southern Biotech). Microscopy images were acquired using an upright microscope (Echo Revolve)

with a 10x objective. To quantify the number of nuclei per syncytia, three randomly selected

fields were acquired and spike-expressing cells and syncytia were manually counted. The average

number of nuclei per syncytium and standard deviation were calculated and visualized using

Microsoft Excel and GraphPad Prism 7.

SARS CoV pseudoparticle infectivity assays. SARS-CoV pseudoparticles were produced as

previously described [20, 27]. To generate pseudoparticles, 3.5×10^5 HEK293T cells were seeded

in 6-well plates. Pseudoparticles were prepared by transfecting HEK293T cells with 600 µg of their

respective SARS WT or mutant S plasmids, 800 µg of pTG-Luc, and 600 µg of pCMV-MLV gagpol

using polyethylenimine (PEI) as the transfection reagent. The cell supernatant was harvested 48

hours post-transfection, centrifuged at 1200 rpm for 7 minutes to separate the pseudoparticles

from residual cellular debris, and filtered through a 0.45 µm syringe filter. The pseudoparticles

were stored at -80°C for one freeze-thaw cycle. To determine pseudoparticle infectivity, 5×10^5

VeroE6 cells were seeded into 24-well plates and infected 200uL of pseudoparticles for 72 hrs at

37°C. Cells were lysed for 10 minutes at room temperature using the Luciferase cell lysis reagent

(Promega). Luminescence readings were performed using a luminometer, Glomax 20/20 system

(Promega). Each experiment contained three technical replicates and was repeated at least three

times. Data analysis was performed using graphical software, GraphPad Prism 7.

For the calcium depletion studies, calcium was chelated from the extracellular or intracellular

environment using EGTA or BAPTA-AM, respectively. To deplete extracellular calcium levels,

VeroE6 cells were pre-treated with DMEM lacking calcium (DMEM- lacks L-glutamine, sodium

pyruvate, HEPES, and calcium; it contains 2% HyClone and 10 mM HEPES) for 1 hour at 37°C. After

the DMEM- pretreatment, the VeroE6 cells were infected with 200 µL of pseudoparticles in the

presence of 50 µM EGTA for 2 hours at 37°C. Following infection, cells were supplemented with

300 µL of calcium-containing cDMEM.

To chelate intracellular calcium, VeroE6 cells were pretreated with 50 µM BAPTA-AM

(dissolved in DMSO) in DMEM+ (DMEM without L-glutamine, sodium pyruvate, HEPES; 2%

HyClone; 10 mM HEPES; Gibco) in the same conditions as described above. Cells were then

infected with 200 µL of pseudoparticles in the presence of 50 µM BAPTA-AM for 2 hours and

supplemented with cDMEM post-infection. Because BAPTA-AM was dissolved in DMSO, VeroE6

cells were pretreated with the equivalent percentage of DMSO used in the BAPTA-AM treated

cells. This served as an important control for the effects of DMSO alone on our infectivity assay.

Pseudoparticles infections carried out in calcium-containing DMEM+ served as a positive control

for infection in absence of the extracellular chelator. Analysis of pseudoparticle infectivity under

these calcium depletion conditions was performed as described above.

To confirm the presence of the S protein in the pseudoparticles, 1.5 mL of the harvested

pseudoparticles were spun down using an ultracentrifuge in a TLA-55 rotor (Beckman-Coulter) at

$42,000 \times g$ for 2 hours at 4°C. The supernatant was aspirated and the S protein-containing pellet

was resuspended and boiled in 1x LDS sample buffer in the presence of DTT for 10 minutes. The

samples were resolved and immunoblotted for the spike protein as described above.

Modeling of the SARS-CoV fusion peptide. The structural cartoon model of the SARS-CoV S

protein fusion peptide is based on the SARS-CoV S prefusion structure from the Protein Data

Bank, PDB 5XLR. The sequence of the SARS-CoV S protein (GenBank accession no. AAT74874.1)

was aligned to the PDB 5XLR SARS-CoV structure sequence using Geneious software (**v.2020.1.1**).

Structural models of the SARS-CoV S monomer were generated using the Modeller comparative

modeling tool (**v.9.23**) within the Chimera software (v.1.13; University of California). Images were

created using Adobe Illustrator CC (**v.24.03**).

Molecular dynamics (MD) simulations of the SARS-CoV FP in water. For all the atomistic MD

simulations, the SARS-CoV FP segment was capped with neutral N- and C-termini (ACE and CT3,

respectively, in the CHARMM force-field nomenclature). Protonation states of all the titratable

residues were predicted at pH 7 using Propka 3.1 software [42].

For the simulations in water, one copy of the peptide (wild-type or a mutant) was embedded

in a rectangular solution box and ionized using VMD tools ("Add Solvation Box" and "Add Ions",

respectively) [43]. The box of dimensions ~90 Å x 80 Å x 82 Å included a Na^+Cl^- ionic solution as

well as 2 Ca^{2+} ions, and ~18000 water molecules. The total number of atoms in the system was

~54,540.

The system was equilibrated with NAMD version 2.13 [44] following a multi-step protocol

during which the backbone atoms of the SARS-CoV FP as well as Ca^{2+} ions in the solution were

first harmonically constrained and subsequently gradually released in four steps (totaling ~3ns),

changing the restrain force constants k_F from 1, to 0.5, to 0.1 kcal/ (mol Å^2), and 0 kcal/ (mol Å^2).

These simulations implemented all option for rigidbonds, 1fs (for k_F 1, 0.5, and 0.1 kcal/ (mol Å^2))

or 2fs (for k_F of 0) integration time-step, PME for electrostatics interactions [45], and were carried

out in NPT ensemble under isotropic pressure coupling conditions, at a temperature of 310 K.

The Nose-Hoover Langevin piston algorithm [46] was used to control the target P = 1 atm

pressure with the "LangevinPistonPeriod" set to 200 fs and "LangevinPistonDecay" set to 50 fs.

The van der Waals interactions were calculated applying a cutoff distance of 12 Å and switching

the potential from 10 Å.

After this initial equilibration phase, the velocities of all atoms in the system were reset and

ensemble MD runs were initiated with OpenMM version 7.4 [47] during which the system was

simulated in 18 independent replicates, each for 640ns (i.e., cumulative time of ~11.5 μs for each

FP construct). These runs implemented PME for electrostatic interactions and were performed

at 310K temperature under NVT ensemble. In addition, 4fs time-step was used, with hydrogen

mass repartitioning and with "friction" parameter set to 1.0/picosecond. Additional parameters

for these runs included: "EwaldErrorTolerance" 0.0005, "rigidwater" True, and

"ConstraintTolerance" 0.000001. The van der Waals interactions were calculated applying a

cutoff distance of 12 Å and switching the potential from 10 Å.

MD simulations of SARS-CoV FP interactions with lipid membranes. Interactions of selected

two models of the Ca^{2+}-bound WT SARS-CoV FP with lipid membranes were investigated with

atomistic MD simulations. These runs were initiated by placing each of the models in the

proximity of a bilayer composed of 3:1:1 POPC/POPG/Cholesterol that had been pre-equilibrated

for 25ns as described previously [23].

After the FP-membrane complexes were embedded in a solution box (containing 150 mM

Na^+Cl^- salt concentration), each system was equilibrated with NAMD version 2.13 following the

same multi-step protocol described above during which the backbone atoms of the FP as well as

the Ca^{2+} ions were first harmonically constrained and subsequently gradually released in four

steps. After this phase, the velocities of all atoms of the system were reset, and ensemble MD

runs were initiated with OpenMM version 7.4. Each system was simulated in 18 independent

replicates, each ran for ~ 1 μs (i.e., cumulative time of ~18 μs for each FP-membrane complex).

These runs implemented PME for electrostatic interactions and were performed at 298K

temperature under NPT ensemble using semi-isotropic pressure coupling, with 4fs time-steps,

using hydrogen mass repartitioning and with "friction" parameter set to 1.0/picosecond.

Additional parameters for these runs included: "EwaldErrorTolerance" 0.0005, "rigidwater" True,

and "ConstraintTolerance" 0.000001. The van der Waals interactions were calculated applying a

cutoff distance of 12 Å and switching the potential from 10 Å.

For all simulations we used the latest CHARMM36 force-field for proteins and lipids [48], as well

as the recently revised CHARMM36 force-field for ions which includes non-bonded fix (NBFIX)

parameters for Na^+ and Ca^{2+} ions [49].

Acknowledgements

This work was supported by The National Institute of Health research grant R01AI35270, National

Science Foundation RAPID grant 2027070 and Fast Grant from the Mercatus Center at George

Mason University. TT is supported by the National Science Foundation Graduate Research

Fellowship Program under Grant No. DGE-1650441 and the Samuel C. Fleming Family Graduate

Fellowship. We would like to thank members of the Daniel, Whittaker, and Abbott groups as well

as the Weinstein group at Weill Cornell for helpful discussions.

Author Contributions

Conceived and designed the experiments: MKB GK MRS HW GRW SD. Performed the

experiments: MKB GK MRS TT JDC. Analyzed the data: MKB GK MRS TT HW GRW SD JDC. Wrote

and edited the paper: MKB GK MRS TT HW GRW SD JDC. Acquired funding: GRW SD.

Figure 1. Generating the charge-to-alanine mutations in the SARS-CoV fusion peptide. (A) The predicted structure of the SARS-CoV fusion peptide (FP) modeled on the SARS PDB: 5XLR. (B) The sequence of the wild-type (WT) SARS FP showing all 40 amino acids (GenBank accession no. AAT74874.1) Listed below are the single charge-to-alanine mutations generated in the aspartic (D) and glutamic (E) acids. The red charged WT residues and the green substitution are shown in each position for the single mutants. Gold highlights a critical disulfide bond within the FP that is necessary for its function. (C) Alignment of the amino acid sequences of SARS-CoV, MERS-CoV, and SARS-CoV-2 FPs showing the high conservation of the LLF motif (green), the charges residues (red) and the disulfide bond. Note that the green LLF residues were not mutated in this study.

Figure 2. Cleavage comparison of SARS-CoV wild-type and mutant spike. Western blot analysis of WT and mutant S proteins with (*+*) and without (-) trypsin. S_0 indicates the full-length, uncleaved spike protein. S_2 indicates the cleaved S2 subunit of the spike protein. (A) Spike immunoblots showing the accumulation of WT S and single charge-to-alanine mutants. (B) Spike immunoblots showing the additional mutants generated at the E801 position. N = 4 (biological replicates).

Figure 3. Fusion activity of the SARS-CoV wild-type and mutant S-expressing cells. (A) Representative immunofluorescence images of VeroE6 cells expressing WT or mutant D802A S constructs. Following transfection, cells were treated with trypsin to cleave SARS-CoV S proteins and induce syncytium formation. Syncytia were visualized using a SARS-CoV S antibody (*green*) and nuclei appear in blue (DAPI). Images were taken at 10X magnification. Scale bars =170μM (B) Quantification of nuclei per syncytium. *** denotes significance of $p < 0.001$; **** denotes significance of $p < 0.0001$.

Figure 4. Infectivity of wild-type and mutant SARS-CoV pseudoparticles. (A) Spike immunoblots of pseudoparticles containing wild-type and mutant spike. S_0 indicates the full-length, uncleaved spike protein. S_2 indicates the cleaved S2 subunit of the spike protein. (B) WT and mutant SARS-CoV pseudoparticle infectivity in VeroE6 cells treated with 50 μM EGTA (+) or left untreated (-). (C) WT and mutant SARS-CoV pseudoparticle infectivity in VeroE6 cells treated with 50 μM BAPTA-AM (+) or left untreated (-). * denotes significance of $p < 0.05$; ** denotes significance of $p < 0.01$; *** denotes significance of $p < 0.001$; **** denotes significance of $p < 0.0001$.

Figure 5. Assessing the potential redundancy in fusion peptide and calcium interactions (A) Double and triple mutants were generated in the FP to determine if residues E821 and D825 are functionally redundant. Multiple charged residues (red) in the SARS FP were mutated to alanines (green) to generate double and triple mutants. Gold highlights a critical disulfide bond within the FP that is necessary for its function. (B) Western blot analysis of WT and double/ triple mutant S proteins with (*+*) and without (-) trypsin. S_0 indicates the full-length, uncleaved spike protein. S_2 indicates the cleaved S2 subunit of the spike protein.

 (C) Representative immunofluorescence images of VeroE6 cells expressing WT or the triple mutant (E821A/D825A/D830A) S constructs. Following transfection, cells were treated with

trypsin to cleave SARS-CoV S proteins and induce syncytium formation. Syncytia were visualized
using a SARS-CoV S antibody (*green*) and nuclei appear in blue (DAPI). Images were taken at 10X
magnification. Scale bars =170μM (D) Quantification of nuclei per syncytium in the WT and
double/triple mutant expressing VeroE6 cells. (E) Spike immunoblots of pseudoparticles
containing wild-type and double/triple mutant spike. S_0 indicates the full-length, uncleaved spike
protein. S_2 indicates the cleaved S2 subunit of the spike protein. (F) WT and double/triple mutant
SARS-CoV pseudoparticle infectivity in VeroE6 cells treated with 50 μM EGTA (+) or left untreated
(-). (G) WT and double/triple mutant SARS-CoV pseudoparticle infectivity in VeroE6 cells treated
with 50 μM BAPTA-AM (+) or left untreated (-). * denotes significance of $p < 0.05$; *** denotes
significance of $p < 0.001$; **** denotes significance of $p < 0.0001$.

Figure 6. Modes of Ca^{2+} binding to SARS-CoV fusion peptide. Number of Ca^{2+} binding events to
different pairs of anionic residues in SARS-CoV FP in the simulations of the WT and the mutant
constructs (labeled in each panel). The experimentally measured phenotypes for each construct
are shown (active or inactive). The mode of Ca^{2+} ion association predicted to be inhibitory for
membrane insertion (E821-D825) is shown in *red*, and the modes of Ca^{2+} association predicted
to facilitate membrane insertion are depicted in *green*.

Figure 7. Membrane insertion for FP1 and FP2 domains of SARS-CoV fusion peptide. (A-B)
Frequency of membrane insertion for each residue of SARS-CoV FP in the MD simulations of
Model 1 (A) and Model 2 (B) constructs (differing in the mode of Ca^{2+} coordination, see text). The
snapshot insertions in the panels show the corresponding structures. Ca^{2+} coordinating anionic
residues are highlighted by licorice rendering; Ca^{2+} ions are shown as blue spheres. The FP1 and
FP2 parts of the fusion peptide are colored in silver and lime, respectively. (C-D) The same
frequency calculations but done separately for two representative trajectories in Model 1 set in
which the observed membrane insertions involved the N-terminal LLF motif (C) and the more
centrally located (F815-M816) hydrophobic segment (D). The corresponding snapshots illustrate
structural features of these two distinct insertion modes highlighting (in licorice) positions of the
anionic residues, the hydrophobic residues penetrating the membrane, and R829 residue
interacting with D825. Same color code as in panels A-B.

Figure 8. Predicted model of CoV FP interaction with a lipid bilayer. This model is based on the
pre-fusion FP domain (from PDB: 5XLR), and summarizes the data obtained from our work,
highlighting the role of key charged residues within the fusion peptide. We depict the insertion
of the FP1 domain with E801/D802-Ca^{2+} (green) needed for membrane association and with D812
able to coordinate Ca^{2+} and pair with either E821 to promote membrane interaction (green) or
with D825 to inhibit membrane interaction (red).

References

1. Lai MMC, Cavanagh D. The Molecular Biology of Coronaviruses. Elsevier; 1997. p. 1-100.
2. Masters PS. The Molecular Biology of Coronaviruses. Advances in Virus Research2006. p. 193-292.
3. Cherry JD, Krogstad P. SARS: the first pandemic of the 21st century. Pediatr Res. 2004;56(1):1-5. Epub 2004/05/21. doi: 10.1203/01.PDR.0000129184.87042.FC. PubMed PMID: 15152053; PubMed Central PMCID: PMCPMC7086556.
4. de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, et al. Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the Coronavirus Study Group. J Virol. 2013;87(14):7790-2. Epub 2013/05/17. doi: 10.1128/JVI.01244-13. PubMed PMID: 23678167; PubMed Central PMCID: PMCPMC3700179.
5. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus associated with human respiratory disease in China. Nature. 2020;579(7798):265-9. Epub 2020/02/06. doi: 10.1038/s41586-020-2008-3. PubMed PMID: 32015508; PubMed Central PMCID: PMCPMC7094943.
6. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3. Epub 2020/02/06. doi: 10.1038/s41586-020-2012-7. PubMed PMID: 32015507; PubMed Central PMCID: PMCPMC7095418.
7. M SM, P M, F A-S, C R, MP K, B M. World Health Organization methodology to prioritize emerging infectious diseases in need of research and development. Emerg Infect Dis. 2018. doi: 10.3201/eid2409.171427.
8. da Costa VG, Moreli ML, Saivish MV. The emergence of SARS, MERS and novel SARS-2 coronaviruses in the 21st century. Arch Virol. 2020;165(7):1517-26. Epub 2020/04/22. doi: 10.1007/s00705-020-04628-0. PubMed PMID: 32322993.
9. Velavan TP, Meyer CG. The COVID-19 epidemic. Trop Med Int Health. 2020;25(3):278-80. Epub 2020/02/16. doi: 10.1111/tmi.13383. PubMed PMID: 32052514.
10. Tang T, Bidon M, Jaimes JA, Whittaker GR, Daniel S. Coronavirus membrane fusion mechanism offers a potential target for antiviral development. Antiviral Res. 2020;178:104792. Epub 2020/04/10. doi: 10.1016/j.antiviral.2020.104792. PubMed PMID: 32272173; PubMed Central PMCID: PMCPMC7194977.
11. de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. Nat Rev Microbiol. 2016;14(8):523-34. Epub 2016/06/28. doi: 10.1038/nrmicro.2016.81. PubMed PMID: 27344959.
12. Belouzard S, Millet JK, Licitra BN, Whittaker GR. Mechanisms of coronavirus cell entry mediated by the viral spike protein. Viruses. 2012;4(6):1011-33. Epub 2012/07/21. doi: 10.3390/v4061011. PubMed PMID: 22816037; PubMed Central PMCID: PMCPMC3397359.
13. Millet JK, Whittaker GR. Physiological and molecular triggers for SARS-CoV membrane fusion and entry into host cells. Virology. 2018;517:3-8. Epub 2017/12/26. doi: 10.1016/j.virol.2017.12.015. PubMed PMID: 29275820.
14. Lee C-Y, Lowen AC. Animal models for SARS-CoV-2. Current Opinion in Virology. 2021;48:73-81. doi: 10.1016/j.coviro.2021.03.009.

15. Millet JK, Whittaker GR. Host cell proteases: Critical determinants of coronavirus tropism and
pathogenesis. Virus Res. 2015;202:120-34. Epub 2014/12/03. doi:
10.1016/j.virusres.2014.11.021. PubMed PMID: 25445340; PubMed Central PMCID:
PMCPMC4465284.

16. Stevens CS, Oguntuyo KY, Benhur L. Proteases and variants: context matters for SARS-CoV-2
entry assays. Current Opinion in Virology. 2021. doi: 10.1016/j.coviro.2021.07.004.

17. White JM, Whittaker GR. Fusion of Enveloped Viruses in Endosomes. Traffic. 2016;17(6):593-
614. Epub 2016/03/05. doi: 10.1111/tra.12389. PubMed PMID: 26935856; PubMed Central
PMCID: PMCPMC4866878.

18. Madu IG, Roth SL, Belouzard S, Whittaker GR. Characterization of a highly conserved domain
within the severe acute respiratory syndrome coronavirus spike protein S2 domain with
characteristics of a viral fusion peptide. J Virol. 2009;83(15):7411-21. Epub 2009/05/15. doi:
10.1128/JVI.00079-09. PubMed PMID: 19439480; PubMed Central PMCID:
PMCPMC2708636.

19. Lai AL, Millet JK, Daniel S, Freed JH, Whittaker GR. The SARS-CoV Fusion Peptide Forms an
Extended Bipartite Fusion Platform that Perturbs Membrane Order in a Calcium-Dependent
Manner. J Mol Biol. 2017;429(24):3875-92. Epub 2017/10/24. doi:
10.1016/j.jmb.2017.10.017. PubMed PMID: 29056462; PubMed Central PMCID:
PMCPMC5705393.

20. Straus MR, Tang T, Lai AL, Flegel A, Bidon M, Freed JH, et al. Ca(2+) Ions Promote Fusion of
Middle East Respiratory Syndrome Coronavirus with Host Cells and Increase Infectivity. J
Virol. 2020;94(13). Epub 2020/04/17. doi: 10.1128/JVI.00426-20. PubMed PMID: 32295925;
PubMed Central PMCID: PMCPMC7307142.

21. Nathan L, Lai AL, Millet JK, Straus MR, Freed JH, Whittaker GR, et al. Calcium Ions Directly
Interact with the Ebola Virus Fusion Peptide To Promote Structure-Function Changes That
Enhance Infection. ACS Infect Dis. 2020;6(2):250-60. Epub 2019/11/21. doi:
10.1021/acsinfecdis.9b00296. PubMed PMID: 31746195; PubMed Central PMCID:
PMCPMC7040957.

22. Lai AL, Freed JH. SARS-CoV-2 Fusion Peptide has a Greater Membrane Perturbating Effect
than SARS-CoV with Highly Specific Dependence on Ca2+. Journal of Molecular Biology.
2021;433(10):166946.

23. Khelashvili G, Plante A, Doktorova M, Weinstein H. Ca^{2+}-dependent mechanism
of membrane insertion and destabilization by the SARS-CoV-2 fusion peptide. Biophysical
Journal. 2021;120(6):1105-19. doi: 10.1016/j.bpj.2021.02.023.

24. Basso LG, Vicente EF, Crusca E, Jr., Cilli EM, Costa-Filho AJ. SARS-CoV fusion peptides induce
membrane surface ordering and curvature. Sci Rep. 2016;6:37131. Epub 2016/11/29. doi:
10.1038/srep37131. PubMed PMID: 27892522; PubMed Central PMCID: PMCPMC5125003.

25. Koppisetti RK, Fulcher YG, Van Doren SR. Fusion Peptide of SARS-CoV-2 Spike Rearranges into
a Wedge Inserted in Bilayered Micelles. Journal of the American Chemical Society. 2021. doi:
10.1021/jacs.1c05435.

26. Wang H, Yang P, Liu K, Guo F, Zhang Y, Zhang G, et al. SARS coronavirus entry into host cells
through a novel clathrin- and caveolae-independent endocytic pathway. Cell Res.
2008;18(2):290-301. Epub 2008/01/30. doi: 10.1038/cr.2008.15. PubMed PMID: 18227861;
PubMed Central PMCID: PMCPMC7091891.

27. Millet JK, Tang T, Nathan L, Jaimes JA, Hsu HL, Daniel S, et al. Production of Pseudotyped
Particles to Study Highly Pathogenic Coronaviruses in a Biosafety Level 2 Setting. J Vis Exp.
2019;(145). Epub 2019/03/19. doi: 10.3791/59010. PubMed PMID: 30882796.
28. Tang T, Jaimes JA, Bidon MK, Straus MR, Daniel S, Whittaker GR. Proteolytic Activation of
SARS-CoV-2 Spike at the S1/S2 Boundary: Potential Role of Proteases beyond Furin. ACS Infect
Dis. 2021;7(2):264-72. Epub 2021/01/13. doi: 10.1021/acsinfecdis.0c00701. PubMed PMID:
33432808; PubMed Central PMCID: PMCPMC7839419.
29. Madu IG, Belouzard S, Whittaker GR. SARS-coronavirus spike S2 domain flanked by cysteine
residues C822 and C833 is important for activation of membrane fusion. Virology.
2009;393(2):265-71. Epub 2009/09/01. doi: 10.1016/j.virol.2009.07.038. PubMed PMID:
19717178; PubMed Central PMCID: PMCPMC3594805.
30. Du L, He Y, Zhou Y, Liu S, Zheng BJ, Jiang S. The spike protein of SARS-CoV--a target for vaccine
and therapeutic development. Nat Rev Microbiol. 2009;7(3):226-36. Epub 2009/02/10. doi:
10.1038/nrmicro2090.   PubMed   PMID: 19198616;   PubMed   Central   PMCID:
PMCPMC2750777.
31. Lu G, Wang Q, Gao GF. Bat-to-human: spike features determining 'host jump' of
coronaviruses SARS-CoV, MERS-CoV, and beyond. Trends Microbiol. 2015;23(8):468-78. Epub
2015/07/25. doi: 10.1016/j.tim.2015.06.003. PubMed PMID: 26206723.
32. Dubé M, Rey FA, Kielian M. Rubella Virus: First Calcium-Requiring Viral Fusion Protein. PLOS
Pathogens. 2014;10(12):e1004530. doi: 10.1371/journal.ppat.1004530.
33. Schaefer SL, Jung H, Hummer G. Binding of SARS-CoV-2 Fusion Peptide to Host tmpr and
Plasma Membrane. The Journal of Physical Chemistry B. 2021;125(28):7732-41. doi:
10.1021/acs.jpcb.1c04176.
34. Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, et al. Distinct conformational states of
SARS-CoV-2 spike protein. Science. 2020;369(6511):1586. doi: 10.1126/science.abd4251.
35. Straus MR, Bidon M, Tang T, Whittaker GR, Daniel S. FDA approved calcium channel blockers
inhibit SARS-CoV-2 infectivity in epithelial lung cells. bioRxiv. 2020:2020.07.21.214577. doi:
10.1101/2020.07.21.214577.
36. Crespi B, Alcock J. Conflicts over calcium and the treatment of COVID-19. Evolution, Medicine,
and Public Health. 2021;9(1):149-56. doi: 10.1093/emph/eoaa046.
37. Danta CC. Calcium Channel Blockers: A Possible Potential Therapeutic Strategy for the
Treatment of Alzheimer's Dementia Patients with SARS-CoV-2 Infection. ACS Chem Neurosci.
2020;11(15):2145-8. Epub 2020/07/15. doi: 10.1021/acschemneuro.0c00391. PubMed
PMID: 32662982.
38. Borges V, Isidro J, Cortes-Martins H, Duarte S, Vieira L, Leite R, et al. Massive dissemination
of a SARS-CoV-2 Spike Y839 variant in Portugal. Emerging Microbes & Infections.
2020;9(1):2488-96. doi: 10.1080/22221751.2020.1844552.
39. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Tracking Changes
in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell.
2020;182(4):812-27.e19.
40. Li Q, Wu J, Nie J, Zhang L, Hao H, Liu S, et al. The Impact of Mutations in SARS-CoV-2 Spike on
Viral Infectivity and Antigenicity. Cell. 2020;182(5):1284-94.e9.

41. Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike protein via
sequential proteolytic cleavage at two distinct sites. Proceedings of the National Academy of
Sciences. 2009;106(14):5871. doi: 10.1073/pnas.0809524106.
42. Olsson MHM, Søndergaard CR, Rostkowski M, Jensen JH. PROPKA3: Consistent Treatment of
Internal and Surface Residues in Empirical pKa Predictions. Journal of Chemical Theory and
Computation. 2011;7(2):525-37. doi: 10.1021/ct100578z.
43. Humphrey W, Dalke A, Schulten K. VMD: Visual molecular dynamics. Journal of Molecular
Graphics. 1996;14(1):33-8.
44. Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, et al. Scalable molecular
dynamics with NAMD. Journal of Computational Chemistry. 2005;26(16):1781-802.
45. Essmann U, Perera L, Berkowitz ML, Darden T, Lee H, Pedersen LG. A smooth particle mesh
Ewald method. The Journal of Chemical Physics. 1995;103(19):8577-93. doi:
10.1063/1.470117.
46. Evans DJ, Holian BL. The Nose–Hoover thermostat. The Journal of Chemical Physics.
1985;83(8):4069-74. doi: 10.1063/1.449071.
47. Eastman P, Swails J, Chodera JD, McGibbon RT, Zhao Y, Beauchamp KA, et al. OpenMM 7:
Rapid development of high performance algorithms for molecular dynamics. PLOS
Computational Biology. 2017;13(7):e1005659. doi: 10.1371/journal.pcbi.1005659.
48. Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, et al. CHARMM36m: an
improved force field for folded and intrinsically disordered proteins. Nature Methods.
2017;14(1):71-3. doi: 10.1038/nmeth.4067.
49. Venable RM, Luo Y, Gawrisch K, Roux B, Pastor RW. Simulations of Anionic Lipid Membranes:
Development of Interaction-Specific Ion Parameters and Validation Using NMR Data. The
Journal of Physical Chemistry B. 2013;117(35):10183-92. doi: 10.1021/jp401512z.
973

Figures

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Supplementary Information

SI Fig S1. Fusion activity of the SARS-CoV wild-type and mutant S-expressing cells.
Immunofluorescence images of VeroE6 cells expressing WT or mutant (single/double/triple) S
constructs. Following transfection, cells were treated with trypsin to cleave SARS-CoV S proteins
and induce syncytium formation. Syncytia were visualized using a SARS-CoV S antibody (*green*)
and nuclei appear in blue (DAPI). Images were taken at 10X magnification.
1007

1008

1009

SI Fig S2. Models of Ca^{2+} binding to the WT and single mutant SARS-CoV FPs. The tables show,
for each construct, acidic residues implicated in Ca^{2+} binding in 18 independent atomistic MD
simulations (each 640ns in length). In a particular trajectory, residue pairs simultaneously
engaging with the bound Ca^{2+} are denoted by *red* or *blue* rectangles, whereas instances of Ca^{2+}
ion associating with a single acidic residue is depicted with *grey-striped* rectangle. The
trajectories in which simultaneous binding of two Ca^{2+} ions to different pairs of residues were
observed are highlighted in *green*. The simulations in which a single Ca^{2+} ion was bound to a pair
of acidic residues are shown in *yellow*.
1018

1019

SI Fig S3. Models of Ca^{2+} binding to the WT and multiple mutant SARS-CoV FPs. The tables show, for each construct, acidic residues implicated in Ca^{2+} binding in 18 independent atomistic MD simulations (each 640ns in length). In a particular trajectory, residue pairs simultaneously engaging with the bound Ca^{2+} are denoted by *red* or *blue* rectangles, whereas instances of Ca^{2+} ion associating with a single acidic residue is depicted with *grey-striped* rectangle. The trajectories in which simultaneous binding of two Ca^{2+} ions to different pairs of residues were observed are highlighted in *green*. The simulations in which a single Ca^{2+} ion was bound to a pair of acidic residues are shown in *yellow*.

Mutation	Forward Primer	Reverse Primer
E801A	GCGCAGCTTCATCGCGGACCTGCTCTTCA	TGAAGAGCAGGTCCGCGATGAAGCTGCGC
E801M	TGTTGAAGAGCAGGTCCATGATGAAGCTGCGCTTGG	CCAAGCGCAGCTTCATCATGGACCTGCTCTTCAACA
E801Q	GAAGAGCAGGTCCTGGATGAAGCTGCGCT	AGCGCAGCTTCATCCAGGACCTGCTCTTC
E801K	TTGAAGAGCAGGTCCTTGATGAAGCTGCGCTTG	CAAGCGCAGCTTCATCAAGGACCTGCTCTTCAA
E801D	GTTGAAGAGCAGGTCGTCGATGAAGCTGCGC	GCGCAGCTTCATCGACGACCTGCTCTTCAAC
D802A	CTTGTTGAAGAGCAGGGCCTCGATGAAGCTGCG	CGCAGCTTCATCGAGGCCCTGCTCTTCAACAAG
D812A	TGAAGCCGGCGGCGGCCAGCGTC	GACGCTGGCCGCCGCCGGCTTCA
E821A	CGCCCAGGCACGCGCCGTACTGC	GCAGTACGGCGCGTGCCTGGGCG
D825A	CGGGCGTTGATGGCGCCCAGGCACT	AGTGCCTGGGCGCCATCAACGCCCG
D830A	GCGCAGATCAGGGCGCGGGCGTTGA	TCAACGCCCGCGCCCTGATCTGCGC
E821A/D825A	CGGGCGTTGATGGCGCCCAGGCACG	CGTGCCTGGGCGCCATCAACGCCCG
E821A/D830A	GCGCAGATCAGGGCGCGGGCGTTGA	TCAACGCCCGCGCCCTGATCTGCGC
D825A/D830A	GCGCAGATCAGGGCGCGGGCGTTGA	TCAACGCCCGCGCCCTGATCTGCGC
E821A/D825A/D830A	GCGCAGATCAGGGCGCGGGCGTTGA	TCAACGCCCGCGCCCTGATCTGCGC

1029

1030   **SI Table S1. List of primers used to generate SARS-CoV S mutants.**

1031