1

A tissue boundary orchestrates the segregation of inner ear sensory organs

Ziqi Chen, Shuting Xu, Magdalena Żak and Nicolas Daudet*

UCL Ear Institute, University College London, London WC1X 8EE, UK

* Corresponding author: n.daudet@ucl.ac.uk,

Abstract

The inner ear contains distinct sensory organs, produced sequentially by segregation from a large sensory-competent domain in the developing otic vesicle. To understand the mechanistic basis of this process, we investigated the changes in prosensory cell patterning, proliferation and character during the segregation of some of the vestibular organs in the mouse and chicken otic vesicle. We discovered a specialized boundary domain, located at the interface of segregating organs. It is composed of prosensory cells that gradually enlarge, elongate and are ultimately diverted from a prosensory fate. Strikingly, the boundary cells align their apical borders and constrict basally at the interface of cells expressing or not the Lmx1a transcription factor, an orthologue of drosophila Apterous. The boundary domain is absent in *Lmx1a*-deficient mice, which exhibit defects in sensory organ segregation, and is disrupted by the inhibition of ROCK-dependent actomyosin contractility. Altogether, our results suggest that actomyosin-dependent tissue boundaries ensure the proper separation of inner ear sensory organs and uncover striking homologies between this process and the compartmentalization of the drosophila wing disc by lineage-restricted boundaries.

INTRODUCTION

The inner ear of vertebrates is composed of a series of liquid-filled chambers containing distinct sensory organs. Each organ contains a sensory epithelium populated with mechanosensory 'hair cells', separated from one another by supporting cells. In amniotes, the ventral part of the inner ear forms the cochlea, which hosts an auditory epithelium called the basilar papilla in birds and crocodiles, or the organ of Corti in mammals. The dorsal part forms the vestibular system, with three cristae (anterior, posterior and lateral) and their associated semi-circular canals responding to the angular rotations of the head, and the maculae of the utricle and saccule, sensitive to gravity and linear acceleration. These sensory organs are separated from one another by non-sensory territories with essential roles in inner ear function and homeostasis (Ekdale, 2016).

The development of the inner ear is a remarkable example of 3D tissue morphogenesis that starts with the formation of the otic placode, an epithelial thickening located on both sides of the hindbrain. The placode rapidly invaginates into the underlying mesenchyme to form the otic cup. Subsequently, the cup pinches off from the surface ectoderm and closes into the otic vesicle (or otocyst), which then undergoes a drastic remodelling to give rise to the inner ear (reviewed in Alsina and Whitfield, 2017; Basch et al., 2016a).

How are the different cell types and territories of the inner ear specified and positioned during its embryonic development? The prevalent view is that diffusible signals (eg Wnt, Hedgehog, FGF, Retinoic Acid) secreted by tissues surrounding the otocyst direct the expression of key factors (transcription factors and signalling molecules) to specify various otic cell identities (reviewed in Fekete and Wu, 2002; Ohta and Schoenwolf, 2018). It was also proposed that once specified, adjacent but distinct populations of otic progenitors might be prevented from intermixing by lineage-restricted tissue boundaries akin to those discovered at the interface of "embryonic compartments" in the Drosophila wing disc (Brigande et al., 2000b; Fekete, 1996). This 'compartment boundary' model has received some support from the analysis of mouse mutants exhibiting defects in restricted domains of the inner ear and some fate map studies, but it remains hypothetical with respect to the existence and precise location of tissue boundaries.

One aspect of inner ear development that results in the spatial separation of adjacent groups of cells is the formation of its sensory organs. In the chicken or mouse otic vesicle, the prospective sensory domains, or prosensory domains, are recognizable as thickened epithelial patches in the ventro-medial and posterior regions (Knowlton, 1967). The prosensory patches express the Notch ligand Jag1, which promotes the maintenance of the prosensory fate by Notch-mediated lateral induction (reviewed in Daudet and Żak, 2020; Kiernan, 2013). They also express the High-Mobility Group transcription factor Sox2, which is essential for the formation of all inner ear sensory organs (Dabdoub et al., 2008; Kiernan et al., 2005) as well as otic neurogenesis (Steevens et al., 2019, 2017). At early developmental stages, Sox2 marks a wide antero-posterior "pan-sensory" domain

occupying the ventro-medial wall of the otocyst (Steevens et al., 2019), due at least in part to the inhibition of Sox2 expression by Wnt activity in the dorsal aspect of the otocyst (Żak and Daudet, 2021). Subsequently, the pan-sensory domain splits into the posterior crista, and several sensory organs in the anterior part of the otocyst: the anterior crista, the lateral crista, and the utricle (Mann et al., 2017; Sánchez-Guardado et al., 2013). This segregation process was first described in the inner ear of fish and amphibians more than a century ago (Norris, 1892) and could have played an essential role in the evolution of the vertebrate inner ear, by allowing new sensory organs to acquire specific functions (reviewed in Fritzsch et al., 2002). Genetic studies in the mouse have shown that mutations or absence of specific transcription factors and signalling molecules can lead to defects in sensory organ segregation (reviewed in Alsina and Whitfield, 2017). One such factor is Lmx1a (cLmx1b in the chicken), a LIM-homeodomain transcription factor whose orthologue Apterous acts as a "selector gene" for the dorsal compartment of the Drosophila wing disc [ref]. In the mouse inner ear, Lmx1a is expressed in the non-sensory territories separating the sensory organs, and its absence causes severe morphogenesis defects and partial fusions between adjacent sensory territories (Koo et al., 2009; Nichols et al., 2008; Steffes et al., 2012). Our previous work has shown that Lmx1a antagonizes Jag1/Notch signalling within some early sensory-competent cells, thereby leading those cells to adopt a non-sensory fate between adjacent sensory organs (reviewed in Daudet and Żak, 2020; Mann et al., 2017). Despite these insights into the genetic regulation of sensory organ segregation, the cellular basis of this process is still unclear. Is it solely dependent on changes in cell character regulated by Notch and Lmx1a, or does it involve other processes and if so, of which type?

To tackle this question, we examined the changes in cell morphology during the segregation of the lateral (LC) and anterior (AC) cristae from the pan-sensory domain in the embryonic chicken and mouse otocyst. We identified a population of prosensory cells with enlarged surfaces, which appear at the interface of segregating organs and gradually elongate and re-align along the border of the prospective cristae before down-regulating Sox2 expression. As segregation proceeds, these large cells constrict basally at the interface of Lmx1a-positive (on the cristae side) and negative (on the pan-sensory domain side) cells, strongly suggesting the formation of a lineage-restricted tissue boundary. In Lmx1a-null mouse otocysts, the boundary domain is disrupted; the AC and the LC initiate their segregation from the utricle but they remain fused to one another, indicating that Lmx1a is required for the differentiation of the non-sensory cells separating the two cristae. Finally, we show that the perturbation of ROCK-dependent actomyosin contractility, a common player in the establishment of lineage-restricted boundaries, leads to defects in the organisation of the boundary domain and abnormal sensory organ segregation.

Altogether, our results strongly suggest that a lineage-restricted tissue boundary forms at the interface of segregating sensory organs, providing support to the compartment-boundary model of inner ear development. They also uncover striking similarities in the genetic circuits and cellular mechanisms of tissue segregation in the inner ear and the Drosophila wing disc.

RESULTS

A specialized boundary domain with a basal constriction forms between segregating vestibular organs in the chicken otocyst

We focused in this study on the formation of the anterior (AC) and lateral (LC) cristae, using whole-mount preparations of the otic vesicle immunostained for the prosensory marker Sox2. In the chicken inner ear, the AC is the first to segregate from the large anterior pan-sensory domain between stage HH23-25 (E4-E5), followed by the LC between HH25-27 (E4.5-E5.5) (**Figure 1a-c**; see also Mann et al., 2017). When we examined at high magnification phalloidin-stained samples, we noticed that the cells located at the interface of the cristae and the pan-sensory domain had a larger apical surface than those of surrounding territories.

Focusing on the LC, which could be identified from the earliest stages of its formation, we distinguished 3 stages of segregation as a function of Sox2 expression and the extent of separation from the pan-sensory domain. At stage 1 (HH25-26), the prospective LC formed a budding that was not yet separated from the pan-sensory domain (**Figure 1a and d-d'''**). Seven to nine rows of Sox2-positive cells located at the interface of the future LC and the pan-sensory domain appeared slightly larger than those located on either side. At stage 2 (HH26-27; **Figure 1b and e-e'''**), Sox2 started to be downregulated within the interface domain separating the LC from the pan-sensory domain. Scattered Sox2-positive nuclei were also present at the edge of a second Sox2-positive lobe of the LC, which remained connected to the pan-sensory domain. Of note, the second lobe of the LC does not give rise to sensory cells at later developmental stages and as discussed later, it is not present in the mouse inner ear. In some cases, the large cells appeared to align their apical borders along the edge of the crista, forming a multi-cellular actin cable-like structure suggesting a coordinated increase in cell bond tension (**Figure 1e-e'''**). At stage 3 (HH27-28; **Figure 1c and f-f'''**), the two lobes of the LC were clearly separated from the pan-sensory domain by an 80-180 μm wide territory containing cells with no (or very reduced) Sox2 expression. At the border of the LC, 7-10 rows of large cells remained visible. Comparable changes were observed at the border of the AC (**Figure 1-Supplementary Figure 1**), although we were not able to analyze the early stages of segregation for this organ.

Concomitant to these changes in cell surface morphology, we noticed a progressive enrichment in F-actin staining at the base of the epithelium along the presumptive border of the LC (**Figure 2**). At stages 2-3, it was associated to a clear basal constriction of the cells and an upward "pull" of the basal lamina along the presumptive border of the LC. This suggests that the basal constriction could at least partly explain the enlarged apical surface of the cells composing the specialized boundary domain.

Figure 1. Segregation of the anterior and lateral cristae in the chicken inner ear. a-c) Low magnification surface views and transverse line optical reconstructions (x,y) of whole-mount preparations of the anterior part of the chicken otocyst at different stages of development (HH=Hamburger and Hamilton stages), immunostained for Sox2 expression. The anterior (AC) and lateral (LC) cristae segregate sequentially from the pan-sensory domain (PD), which then becomes the utricle (UT). The LC is connected to a second Sox2-expressing domain (stars) that does not give rise to a sensory organ. **d-i")** Stages 1 (**d-e"**), 2 (**f-g"**) and 3 (**h-i"**) of the segregation of the LC, characterized by a progressive down-regulation of Sox2 expression in the domain separating the LC from the PD domain. High magnification views (**e-e"**, **g-g"** and **i-i"** correspond to the boxed areas in respectively **d-f-h)** reveal striking changes in cellular morphology and organization during LC segregation. A specialized boundary domain composed of large cells that progressively align their membranes along the border of the LC (arrowheads in **e'-i"**) becomes clearly visible from stage 2 onwards.

Figure 1 – Supplementary Figure 1. Segregation of the anterior crista in the chicken inner ear.

Low and high magnification views of the anterior crista (AC) and pan-sensory domain (PD) at

different stages of development (HH=Hamburger and Hamilton stages), immunostained for Sox2

expression and labelled with fluorescent phalloidin. At stage 1 of segregation (**a-b”**), the two patches

are in contact and few cells have enlarged surfaces at their interface (arrowheads in **b’-b”**). At stage

2 (**c-d”**), several rows of larger cells are seen at the interface of the PD and prospective AC

(arrowheads in **d’-d”**). At stage 3 (**e-f”**), the two patches are separated by a Sox2-negative domain

and several rows of large cells (arrowheads in **f’-f”**).

Figure 2. A multicellular basal constriction forms at the interface of the lateral crista and pan-sensory domain. a-c) Samples stained for F-actin and Sox2 expression at stages 1-3 of LC segregation in the chicken inner ear. Optical sections collected from the surface and basal planes, and orthogonal reconstructions (z). In all images, the pan-sensory domain is on the left and the LC on the right. **a)** F-actin is enriched at the base of the interface of the LC from stage 1 (arrowheads). **b-c)** At stage 2-3, the F-actin enrichment becomes very clear along the prospective border of the LC (arrowheads). A basal constriction of the cells (arrows in z) and an upward 'pull' of the basal lamina (stars) are also noticeable.

We next used the Epitool (Heller et al., 2016) and Icy softwares (see Methods) to analyse in greater
detail the changes in surface cell morphology during the segregation of the LC. Cell skeletons were
generated from phalloidin-stained preparations (n=3 samples per stage), allowing the quantification
of cell surfaces, elongation (the ratio between their long and short axis) and orientation (**Figure 3**).
This analysis showed that the large cells at the interface of the LC and pan-sensory domain
expanded their apical surface and increased in number from stage 1 to stage 3 (**Figure 3b-c**); they
became more elongated than neighbouring cells from stage 2 (**Figure 3d**) and their long axis
became gradually aligned along the same planar polarity vector, parallel to the edge of the crista, at
stage 2-3 (**Figure 3e**). The increase in cell surface area was maximum in the interface domain but
observed throughout the tissue, explaining the overall reduction in cell density observed between
stage 1 and stage 3. The elongation and re-orientation of the cells was also noticeable throughout
the epithelium, suggesting that these are primarily driven by global anisotropic changes in tissue
tension. Altogether these results indicate that very dynamic changes in cell morphology and tissue
patterning occur at the interface of the prospective LC and the pan-sensory domain before (stage 1)
and during (stage 2-3) their segregation.

A multicellular basal constriction forms at the interface of Lmx1a-positive and Lmx1a-negative cells in the mouse otocyst

To find out if a comparable interface domain forms in the mammalian inner ear, we analysed the
otocysts of Lmx1a^{GFP} knock-in mouse embryos. In a previous study, we showed that the inner ear of
heterozygous Lmx1a$^{GFP/+}$ mice develops normally and GFP expression becomes gradually
upregulated in the non-sensory territories surrounding the vestibular prosensory patches (Mann et
al., 2017). However, we did not investigate in detail the earliest stages of segregation of the two
cristae from the pan-sensory domain.

In E11.5 otocysts (n=4), two small Sox2-positive patches corresponding to the prospective AC and
LC were visible at the anterior edge of the pan-sensory domain (**Figure 4a**). However, in contrast to
what was observed in the chicken otocyst, the LC did not have a second Sox2-positive lobe and the
segregation of the AC and LC was synchronous. In fact, the AC was still connected to the pan-
sensory domain at the time the LC budding emerged from it. In some way, the size and location of
the AC and LC in E11.5 mouse otocysts resembled that of the two Sox2-positive lobes of the LC in
the chicken otic vesicle at stage 2 (compare with **Figure 1b**). At this stage, Lmx1a/GFP expression
pattern overlapped with that of Sox2 in between the presumptive cristae and at their interface with
the large pan-sensory domain (**Figure 4b-b'**). Seen from the surface, the border of the Lmx1a-GFP
positive territory was relatively jagged, although GFP-positive and negative cells remain clearly
separated. The interface domain contained cells with a slightly enlarged surface in 1 out of 4 samples
only, and there was no basal constriction (**Figure 4c-c'**). We conclude that E11.5 corresponds to an
early stage of segregation of the cristae, equivalent to 'stage1' in the chicken inner ear.

Figure 3. Analysis of the morphological changes during segregation of the lateral crista in the chicken otocyst. a) Cell skeletons generated from phalloidin stained preparations at three stages of LC segregation. In all images, the interface domain is in the middle, the pan-sensory domain on the left, and the LC on the right. **b-c**) Colour scale and pooled scatter plots of the surface cell area show that the largest cells are located at the interface of the LC and pan-sensory domain from stage 1. **d**) Color-coded representation of the cell elongation ratio. **e**) Visualization of the long axis of the cells showing a progressive elongation between stages 1-3 and a clear alignment of the cells along the border of the LC at stage 3.

Figure 4. Segregation of the anterior and lateral cristae in the heterozygous *Lmx1a* $^{GFP/+}$ knock-in mouse inner ear. Whole-mount surface views and orthogonal reconstructions (x) of the anterior part of the otocysts from *Lmx1a* $^{GFP/+}$ mouse at different stage of development, immunostained for Sox2 expression and stained for F-actin. **a)** At E11.5, the prospective anterior (AC) and lateral (LC) cristae are visible at the anterior edge of the pan-sensory domain (PD). Lmx1a/GFP expression pattern overlaps with that of Sox2 in between the presumptive cristae and at their interface with the PD **(b-b')**. The border of the Lmx1a-GFP positive territory is jagged, but GFP-positive and negative cells are clearly separated **(arrowheads in c')**. Cell surfaces appear slightly enlarged at the limit of the Lmx1a/GFP **(star in c). d)** At E13.5, the AC, LC and UT are separated. Lmx1a/GFP expression is present in Sox2-positive cells at the border of the AC and LC **(arrowheads in e-e')** and in prospective non-sensory cells in between the two cristae, which exhibited a clear reduction in Sox2 expression. Cells with enlarged surface areas are clearly recognizable in between the cristae and UT **(brackets in f-f')**. Lmx1a/GFP is detected in the half of the large cell domain facing the cristae. A basal constriction is present at the precise interface of the Lmx1a-expressing and non-expressing cells **(arrows in f-f')**. At P0, Lmx1a/GFP expression persisted in a wide non-sensory territory separating the two cristae from the UT **(g)**. There is no overlap between Sox2 and GFP expression at the border of the sensory organs **(stars & bracket in h-h')**. No basal constriction is observed **(bracket in i)**.

At E13.5 (n=5), Sox2 expression marked 3 discrete patches corresponding to the AC, the LC and

the prospective utricle (**Figure 4d**). Lmx1a/GFP expression was still present in Sox2-positive cells

at the border of the AC and LC and in prospective non-sensory cells in between the two cristae which

exhibited a clear reduction in Sox2 expression (**Figure 4e-e'**). Furthermore, an interface domain with

enlarged cells was clearly recognizable in between the cristae and prospective utricle (**Figure 4f-f'**).

Remarkably, Lmx1a-GFP was detected in the half of the large cell domain facing the cristae; the

other half, facing the prospective utricle, had no or very low levels of GFP fluorescence and exhibited

reduced Sox2 expression. The apical membranes of the cells at the interface of GFP-positive and

negative domains were in some samples markedly aligned (**Figure 4 - Supplementary Figure 1**),

suggesting an increase in cell bond tension along the border of Lmx1a expression.

Figure 4 - Supplementary Figure 1. Alignment of cell borders at the interface of Lmx1a-

positive and negative cells in the mouse inner ear. (a-a') Whole-mount preparations of the

interface between the pan-sensory domain (PD) and lateral crista (LC) in an E13.5 *Lmx1a* $^{GFP/+}$

mouse otocyst. F-actin labelling reveals a marked alignment of apical cell borders along the

boundary of Lmx1a-GFP expression (arrowheads).

Remarkably, a basal constriction similar to that observed in the chicken otocyst (at stage 2-3) was

present at the precise interface of the Lmx1a-expressing and non-expressing cells (transverse view

in **Figure 4f-f'**).

In neonate mice (n=3; **Figure 4g**), Lmx1a/GFP expression persisted in a wide non-sensory territory

separating the two cristae from the utricle. In contrast to earlier stages, there was no overlap between

Sox2 and GFP expression at the border of the sensory organs (**Figure 4h-h'**) and a basal

constriction was either absent or impossible to precisely locate in the samples examined (**Figure 4i-**

i').

Altogether, these data indicate that the cellular correlates of cristae segregation are comparable in

the mouse and chicken inner ear, despite some differences in its dynamics and outcomes. In both

species, a specialized population of boundary cells with enlarged apical surfaces appears at the

interface of the two cristae and the utricle at early stages of their segregation. Furthermore, a basal

constriction and alignment of apical cell borders occur in the middle of the large cell domain, at the

precise interface of Lmx1a-expressing and non-expressing cells. This strongly suggests that a

lineage-restricted tissue boundary separates these two cell populations.

The boundary domain is disrupted in *Lmx1a*-null mice

The complete absence of Lmx1a function leads to severe defects in inner ear morphogenesis,

including absence of semi-circular canals or tissue constrictions and an abnormal growth and

segregation of sensory organs (Koo et al., 2009; Mann et al., 2017; Nichols et al., 2008; Steffes et

al., 2012). To find out if Lmx1a is required for the formation of the boundary domain, we compared

cell morphology and organisation in E14 *Lmx1a*$^{GFP/+}$ heterozygous (n=8) versus *Lmx1a*$^{GFP/GFP}$

homozygous (*Lmx1a*-null; n=5) knock-in mice (**Figure 5**). Otocysts from homozygous *Lmx1a*$^{GFP/GFP}$

animals had very severe defects in the morphology of the AC and LC, which in all samples exhibited

various degrees of expansion and fusion with one another (compare **Figure 5a** and **5e**). Consistent

with our previous observations (Mann et al., 2017), Sox2 expression was maintained in the GFP-

positive cells within the fused cristae, suggesting that the fusion of the two cristae is to a large extent

due to a specification defect: in the absence of Lmx1a, sensory-competent cells fail to differentiate

into non-sensory cells at the interface of the AC and LC, producing instead a large and continuous

sensory territory. However, a relatively straight border of GFP-expression was present at the edge

of the fused cristae (**Figure 5b** and **5f**) and a domain with reduced Sox2 expression was present

between the cristae and utricle of the *Lmx1a*$^{GFP/GFP}$ samples, although not as clearly defined than in

the *Lmx1a*$^{GFP/+}$ samples (stars in **Figure 5c** and **5g**). This suggests that the initial segregation of the

crista and utricle pools of sensory progenitors is to some extent independent from Lmx1a function.

Nevertheless, there were some differences in GFP pattern, cell morphology and organisation at the

interface of the fused cristae and utricle. Firstly, GFP pattern was very mosaic across *Lmx1a*$^{GFP/GFP}$

cells, including at the border of the GFP-positive domain. This may be due to the abnormal

differentiation of cells of the Lmx1a-lineage as well as increased mixing between cells belonging or

not to the Lmx1a-lineage at the edge of the cristae. Secondly, there was no basal constriction at the

interface of the GFP-positive and negative cells in the *Lmx1a*$^{GFP/GFP}$ samples (compare **Figure 5d**

and **5h**). Finally, the cellular organisation of the boundary domain was markedly different in the

Lmx1a$^{GFP/GFP}$ samples. In fact, a quantitative analysis of cell surface morphology showed that cells

on both sides of the boundary domain in *Lmx1a*$^{GFP/GFP}$ samples (n=3) tended to be larger, more

aligned with one another and more elongated than in *Lmx1a*$^{GFP/+}$ ones (n=2) (**Figure 5 – Figure**

Supplementary 1). Interestingly, these changes affected both GFP-positive and negative domains,

suggesting non-cell autonomous consequences to the absence of *Lmx1a*.

Figure 5 – Comparison of the boundary domain in *Lmx1a* ^{GFP/+} and *Lmx1a* ^{GFP/GFP} mouse inner ear. Whole-mount preparations of the utricle (UT), lateral (LC) and anterior (AC) cristae region, after immunostaining for Sox2 (magenta) and F-actin labelling with phalloidin (white). **(a)** In the E15 *Lmx1a* ^{GFP/+} inner ear (**a-d**), the three sensory patches are segregated and Lmx1a-GFP is present around the utricle and in between the two cristae (star in **a**). High magnification surface **(b-c)** and transverse reconstruction (**d**) of the interface domain (brackets). The boundary of Lmx1a-GFP expression is relatively straight (arrowheads) and Sox2 expression is reduced in the cells immediately adjacent to it and facing the utricle (stars in **c**). A basal constriction is visible at the border of the Lmx1a-GFP expression domain (arrow in **d**). In the E15 *Lmx1a* ^{GFP/GFP} inner ear (**e-h**) GFP expression is reduced at the interface of the UT and the fused AC and LC (stars in **e**). At higher magnification (**f-h**), note the presence of GFP-negative cells in between GFP-positive cells at the interface domain (arrows in **f-g**) but a domain of reduced Sox2 expression (stars in **g**) is maintained between the fused cristae and the UT. (**h**) In transverse reconstructions, note the absence of a basal constriction at the border of the cristae domain.

Altogether, these results show that Lmx1a is required for the differentiation of the non-sensory

territories between the AC and LC and the maintenance of the boundary domain between the cristae

and the utricle. This is consistent with the notion that Lmx1a acts as a "selector" for the non-sensory

fate in the domain separating the two cristae. Lmx1a function is also required for the formation of the

large cell domain and its associated basal constriction at the limit of the Lmx1a-expressing domain,

but neither appear necessary for the initial separation of the utricle from the cristae.

Figure 5 – Figure Supplementary 1. Quantitative analysis of the boundary domain in *Lmx1a*GFP/+ (left column) and *Lmx1a*GFP/GFP (right column) E15 mouse inner ear. a) Representative cell skeletons generated from phalloidin stained preparations. In all images, the interface domain is in the middle, the utricle on the left, and the LC/AC domain on the right. b-c) Colour scale representation (from skeletons shown in **a**) and pooled scatter plots (from 2 *Lmx1a*GFP/+ and 3 *Lmx1a*GFP/GFP samples) of the surface cell areas. Note the more uniform distribution of the large cells on both sides of the interface domain in the Lmx1a-null inner ear. d-e) Visualization of the

1   long axis (from skeletons shown in **a**) and pooled elongation ratio (from 2 *Lmx1a*$^{GFP/+}$ and 3

2   *Lmx1a*$^{GFP/GFP}$ samples) of the cells. Note that cells are more aligned with one another and more

3   elongated in the homozygous than in the heterozygous samples.

Pharmacological blockade of ROCK-dependent actomyosin contractility disrupts sensory organ segregation

In other contexts, tissue boundary formation or maintenance depends on increased actomyosin

contractility at the interface of lineage-restricted compartments (reviewed in Dahmann et al., 2011;

Monier et al., 2011). The resulting increase in cell bond tension is thought to create a physical 'fence'

that restricts cell intermixing. To find out if this might also be the case in the inner ear, we set up free-

floating organotypic cultures of HH25 chicken otocysts, an early stage of segregation of the LC, and

treated these with Y-27632, a potent inhibitor of ROCK-dependent actomyosin contractility (**Figure**

6).

After short-term (3h) culture in control DMEM medium (n=3; **Figure 6a-c**), HH26 otocysts at early

stages of LC segregation had a well-recognizable large cell domain, but only one out of three

samples examined had a clear basal constriction. Multicellular rosettes were frequently observed at

the surface of the epithelium (**Figure 6b'**). However, the interface between the Sox2-positive nuclei

of the lateral crista and utricle was not as well defined as *in vivo*, suggesting a partial disruption of

tissue organisation *in vitro*. In samples treated with 20μM Y-27632 (**n=3; Figure 6d-f**), the large cell

domain was poorly defined due to the increase in the size of the cells bordering it and there was no

basal constriction. Multicellular rosettes were completely absent, suggesting a rapid inhibition of

cortical actomyosin contractility by Y-27632. We also noted in all samples some folding and

invaginations of the epithelium, likely resulting from a global increase in apical cell surface caused

by the inhibition of actomyosin contractility. Nevertheless, the interface between the LC and the

utricle was still visible, indicating that a short blockade of actomyosin contractility does not result in

extensive cell mixing at the interface of the two organs.

Next, we examined the consequences of longer Y-27632 treatments. We first performed some

experiments to determine whether the initial stages of LC segregation can occur *in vitro*. We

compared LC size and segregation in 15 pairs of otocysts, with each pair from the same embryo

(HH26 – stage 1 or 2 of LC segregation) composed of an otocyst immediately fixed after dissection

and the other one maintained in culture for 24h before fixation (**Figure 6g-l**). All of the cultured LC

showed some progression in LC formation compared to their contralateral control, with a clear

boundary between weakly and strongly Sox2- expressing cells in 12/15 samples and a basal

constriction present in 14/15 samples. The size of the LC was greater in cultured samples,

suggesting growth and survival of the prosensory cells. However, the distance between the LC and

the pan-sensory domain ranged from 15.82-49.82 μm only, which was reduced compared to what

would normally be observed *in ovo* in a stage HH27 otocyst (n=9; distance = 46.3-179.62 μm; see

1 also Mann et al., 2017). These results show that the initial stages of specification and segregation of

2 the LC do occur *in vitro*, albeit at a slower pace than *in vivo*.

3 We next compared pairs of contralateral HH26 otocysts cultured for 24h in either control medium or

4 20μm Y-27631 (**Figure 6m-r**). Only three (out of Y-27631-treated samples could be analysed, due

5 to the extreme fragility of the tissue which made impossible the dissection and mounting of the

6 samples. Compared to their contralateral controls, the Y-27631-treated samples exhibited an

7 extensive fusion and expansion of both the AC and LC domains, which were nevertheless present

8 and distinguishable from the pan-sensory domain. Signs of severe epithelial disorganisation included

9 the frequent absence of F-actin staining at the apical surface, suggesting a disassembly of the

10 adherens junction complexes, and Sox2-positive nuclei located very close to the surface. The

11 interface domain with large cells and a reduced Sox2 expression was absent, indicating that ROCK

12 activity might be dispensable for the specification and growth of the LC, but it is absolutely required

13 for the maintenance of its epithelial character and its segregation from the pan-sensory domain.

Figure 6 – Effects of the ROCK inhibitor Y-27632 on the segregation of the lateral crista in organotypic cultures of embryonic chicken otocysts. a-f) Surface and orthogonal views (x, c, f) of organotypic cultures of stage HH26 otocysts maintained for 3 hours in control medium (a-c) or medium supplemented with 20μm Y-27632 (d-f). Panels b-c and e-f are high magnification views of the boxed area in respectively a and d; the position of the large cell domain is indicated by a bracket and an example of multicellular rosette is highlighted (arrowhead in b'). Treatment with Y-27632 induces a loss of multicellular rosettes and abnormal folding of the epithelium (stars in e'-f). g-l) Surface and orthogonal views (x, i, l) of a pair of otocysts from the same HH26 embryo; one otocyst was fixed immediately (g-i) and the other maintained for 24 hours in culture (j-l). Panels h-i and k-l are high magnification views of the boxed area in respectively g and j; the position of the large cell domain is indicated by a bracket. After 24 hours, the lateral crista has progressed in its segregation from the pan-sensory domain. There is a clear reduction in Sox2 expression levels at the edge of the prospective LC and a basal constriction is noticeable (arrows in k-k'). m-r) Surface and orthogonal views (x, o, r) of a pair of otocysts from the same HH26 embryo; one otocyst was cultured 24 hours in control medium (m-o) and the other maintained for 24 hours in 20μm Y-27632 (p-r). Panels n-o and q-r are high magnification views of the boxed area in respectively m and p. Treatment with Y-27632 induces a loss of the typical epithelial cell morphology and large cell domain (compare n' and q') and a disruption of sensory organ segregation (stars in p). Abbreviations: LC=lateral crista; AC=anterior crista; PD=pan-sensory domain.

1 **Perturbation of ROCK activity in vivo disrupts the organization of the boundary**
2 **domain**

3 In light of the very severe defects in epithelial morphology caused by long-term Y-27631 treatment
4 and the limited extent of cristae segregation *in vitro*, we turned to a genetic approach to interfere
5 with ROCK-dependent actomyosin contractility *in vivo* in the developing otocyst. We used *in ovo*
6 electroporation (EP) of the otic cup to overexpress in the otocyst a truncated dominant-negative form
7 of ROCK fused to EGFP (RIIC1-EGFP), able to bind to its upstream regulator Shroom3, but lacking
8 the kinase domain necessary for Myosin II regulation (Nishimura and Takeichi, 2008). Pilot
9 experiments on HH23 samples transfected with a transient expression vector confirmed that RIIC1-
10 EGFP localizes at the cell membrane (Figure **7a**). The analysis of surface cell morphologies with
11 Epitool (Figure **7b-d**) showed that RIIC1-EGFP causes a mild (approximately 23%) but statistically
12 significant ($p < 0.01$, Mann Whitney test $U = 10622.5$) increase in the average apical surface area of
13 transfected otic cells ($18.22 \mu m^2$; SD=11.04) compared to untransfected cells ($14.84 \ \mu m^2$; SD=8.38),
14 consistent with an inhibition of cortical actomyosin contractility.

15 Next, we used the RCAS retroviral system (Hughes, 2004) to investigate the effects of RIIC1-EGFP
16 on cristae segregation, since EP of a replication-competent retrovirus proviral DNA leads to a
17 sustained and widespread transgene expression (Freeman et al., 2012). The right otic cups of E2
18 (HH13-14) embryos were electroporated with either the RCAS-EGFP (control) or the RCAS-RIIC1-
19 EGFP proviral DNA and the embryos were incubated for 3 to 5 days post-EP. In controls (n=8; **Figure**
20 **7e-f"**), the LC was present in 6 out of 8 samples and separated from the utricle by distances ranging
21 from 55 to 155 μm. We hypothesize that the absence of LC in two of the controls may be due to
22 abnormal or delayed development of the inner ear after EP. The pattern of EGFP expression was
23 very variable among samples, but in general it encompassed sensory and non-sensory regions.
24 However, in two samples, we noted a very marked difference in EGFP fluorescence levels on each
25 side of the boundary of the LC (**Figure 7f'**). Although several factors could potentially explain this
26 observation (e.g. different rate of EGFP production or degradation in different cell populations), it fits
27 with the idea that there is limited cell intermixing at this particular location.

28 In the RCAS-RIIC1-EGFP electroporated samples (n=6; **Figure 7g-j"**), the LC was detected in all
29 cases but it was either connected to the utricle (n=2/6), or migrated at a smaller distance than in
30 controls (n=4/6; range = 24-64μm), suggesting an impaired segregation. The examination of the
31 putative LC boundary domain at high magnification showed that as previously observed with the
32 short-term overexpression experiments, RIIC1-EGFP fluorescence was enriched at the cell surface
33 and baso-lateral membranes (**Figure 7h'** and **j'**).

Figure 7 – Overexpression of a dominant-negative form of ROCK in the developing chicken inner ear disrupts sensory organ segregation. a) surface view of otic epithelial cells 24h after in ovo electroporation with the pRCII-GFP expression plasmid. **b-d)** analysis of cell skeletons of transfected (in green) versus untransfected (UT) cells indicate that overexpression of the RCII-GFP fusion protein induces an increase in apical cell surface area. **e-f")** surface views of an E6 control otocyst electroporated at E2 with an RCAS-GFP proviral DNA. In this sample, the limit of Sox2 expression (arrowheads in f) matches that of strong GFP fluorescence (arrowheads in f') and the presumptive tissue boundary in the middle of the large cell domain (bracket in f") separating the

lateral crista (LC) from the utricle (UT). **g-j")** surface views of two E6 samples electroporated at E2

with RCAS-RCII-GFP and showing abnormal segregation of the LC. In the first example, the

interface domain contains poorly aligned Sox2-expressing cells (arrowheads in h). A basal

constriction is still visible (arrows in **h'** and **h"**, transverse reconstructions) but the large cells of the

interface domain are not properly aligned (arrowheads in **h"**). In the second example (**i-j"**), a mosaic

pattern of RCII-GFP fluorescence is observed in the interface domain. Cells with low levels of RCII-

GFP fluorescence (star in **j'**) exhibit a more regular alignment of their apical borders (arrowheads in

j") than transfected cells.

Despite some variability in severity, clear defects in the alignment of Sox2-positive cells were

observed at the border of the prospective LC in regions with high levels of RCII-EGFP expression,

suggesting potential intermixing between cells of the LC and those of prospective non-sensory

domains. Although it was not always possible to visualize it clearly, a basal constriction appeared

to be present, including in domains with high levels of RCII-EGFP expression (**Figure 7h'-h"**).

Strikingly, in one sample with a mosaic pattern of RCII-EGFP expression in the boundary domain,

the untransfected cells were the only ones exhibiting a clear alignment of their apical borders

reminiscent of what was observed in controls (**Figure j'-j"**).

We conclude that ROCK-dependent actomyosin contractility is required for the coordination of

cortical cell tension at the boundary between the LC and pan-sensory domain and for normal

sensory organ segregation. However, additional mechanisms are likely to limit the intermixing of

prosensory cells at the boundary since the separation of Sox2-high and Sox2-low cells was still

visible in these experiments.

1 The boundary domain is highly proliferative

Figure 8. Analysis of cell proliferation during the segregation of the lateral crista in the chicken otocyst. whole-mount preparations of samples collected one hour after an EdU pulse in ovo at stage 1 (**a-b**) or stage 2-3 (**c-d**) of segregation of the LC, immunostained for EdU, Sox2 an labelled with DAPI. Numerous EdU-positive cells (**arrows in a-c**) and mitotic figures (**arrows in b-d**) are present at the interface between the LC and PD (**dotted line in b-d**) at both stages. **e)** Spatial distribution of the mitotic figures within a 36x72μm region centred on the middle of the boundary domain, for early and late stages of segregation (n=7 embryos for each stage).

One important function of actomyosin-dependent tissue boundaries is to limit the disruptive impact of cell proliferation on the spatial organisation of cells during tissue growth and morphogenesis (reviewed in Monier et al., 2010). This prompted us to examine the spatial pattern of cell proliferation during LC segregation. To detect S-phase cells, we administered *in ovo* a short (2 hour) pulse of the thymidine analogue Edu to stage HH23-25 embryos before collecting inner ear tissue and process

it for Click-iT EdU reaction and Sox2 immunostaining. Numerous Edu-positive cells were present
throughout the boundary domain at both early (stage 1, **Figure 8a-b**; n=7) and advanced (stage 2-
3, **Figure 8c-d**; n=7) segregation stages, and these always included some of the cells located at the
interface of the prospective LC and pan-sensory domain. This was confirmed by the seemingly
random distribution of mitotic figures (visualized by DAPI staining) within the boundary domain of
fixed samples (n=7 for each stage; **Figure 8b**, **d** and **e**). We conclude that the boundary at the
interface of the LC and pan-sensory domains contains mitotic cells throughout the segregation
stages.

DISCUSSION

We discovered a specialized actomyosin-dependent boundary domain at the interface of segregating
sensory organs in the developing inner ear. We propose that this boundary restricts intermixing
between Lmx1a-positive and negative cells during the segregation of adjacent sensory organs and
act as a signalling centre guiding the differentiation of non-sensory territories. Our findings provide
strong support to the 'compartment boundary' model for inner ear morphogenesis (Brigande et al.,
2000b; Fekete, 1996; Fekete and Wu, 2002) and suggest that an evolutionary conserved module for
tissue boundary formation has been co-opted in the inner ear for separating its distinct sensory
organs.

A tissue boundary forms during the segregation of inner ear sensory organs

The morphogenesis of complex tissues depends on the harmonious coordination of cell proliferation,
cell specification, and tissue patterning events. Tissue boundaries play an important role in this
process by preventing, at specific locations, the intermingling of cells destined to form distinct
anatomical structures. Besides this 'physical' role, tissue boundaries can act as signalling centres or
'organizers', regulating the proliferation and differentiation of adjacent cells through various
intercellular signalling pathways. Tissue boundaries were discovered (Garcia-Bellido et al., 1973)
and most studied in the imaginal wing disc of Drosophila, where they establish lineage-restricted
compartments along the dorso-ventral and antero-posterior axis (Wang and Dahmann, 2020). In
vertebrates, they are present in the nervous system (Addison and Wilkinson, 2016; Langenberg and
Brand, 2005), in the ectoderm of the limb bud (Altabef et al., 1997), or in the gut (Smith and Tabin,
2000). Their formation typically involves three consecutive steps (Addison and Wilkinson, 2016;
Dahmann et al., 2011; Wang and Dahmann, 2020). Firstly, 'selector' genes (encoding transcription
factors) assign a specific identity to different populations of cells. Secondly, these cells interact
through signalling molecules and sort themselves into adjacent domains according to their genetic
identity through adhesive or repulsive interactions. Finally, actomyosin-dependent processes
increase cell tension at the interface of adjacent compartments to create and maintain a stable 'fence'
preventing cell mixing (Major and Irvine, 2006; Monier et al., 2011, 2010).

Figure 9. Schematic representation of sensory organ segregation. a) The up-regulation of Lmx1a expression within the pan-sensory domain coincides with the formation of a specialised boundary domain composed of cells with enlarged cell surfaces. A basal constriction and an alignment of cell borders occur at the precise interface of Lmx1a-positive and Lmx1a-negative cells. We propose that it is a transient lineage-restricted boundary, dependent on actomyosin contractility, which separates adjacent pools of sensory progenitors. As the spatial segregation proceeds, the Lmx1a-expressing cells give rise to non-sensory cells separating sensory organs. **b)** Hypothetical regulatory gene network regulating sensory organ segregation. Notch-dependent lateral induction maintains Sox2 expression and promotes the prosensory fate. Lmx1a antagonizes Notch activity and promotes adoption of a non-sensory fate; it is also required for the proper formation of the boundary domain between the cristae and utricle.

The morphogenesis of the inner ear has long been proposed to depend on the formation of lineage-restricted embryonic compartments analogous to those present in the fly wing disc. In fact, some genes have sharply defined domains of expression in the otic vesicle before any sign of morphological differentiation and their absence prevents the formation of specific structures of the adult inner ear, suggesting that these may act as selector genes for a particular otic fate (Brigande et al., 2000b; Fekete, 1996; Fekete and Wu, 2002). Fate map and lineage studies have also suggested the existence of lineage-restricted boundaries in the dorsal part of the chicken otic vesicle, which gives rise to the endolymphatic duct and sac, although their exact location and cellular features remain unknown (Brigande et al., 2000a; Kil and Collazo, 2002; Sánchez-Guardado et al., 2014). In this study, we provide strong evidence that the segregation of the anterior and lateral cristae from the prospective utricle is associated to the formation of a tissue boundary. It is composed of a subset of prosensory cells which progressively enlarge, elongate, align their apical cell borders and constrict

basally at the precise junction between Lmx1a-positive and negative cells. In the absence of lineage-tracing or live-imaging experiments, we do not have definitive proof that these cells generate a strict lineage-restricted boundary. However, their position (at the interface of two distinct "genetic compartments") and cellular features (basal constriction and alignment of their apical cell borders) are consistent with this idea. Furthermore, a lineage-restricted boundary could explain why we observed, in some of our RCAS-infection experiments, an uneven distribution of GFP-expressing cells on either side of the basal constriction separating the lateral crista from the utricle. In a previous study, we argued that the changing patterns of Lmx1a and Sox2 expression in the embryonic inner ear suggested some form of dynamic competition for the adoption of the sensory versus non-sensory fates at the lateral border of sensory organs, as opposed to the existence of a strict lineage boundary (Mann et al., 2017). Our new results, which focused on earlier stages of sensory organ formation, call for a revision of this conclusion: the early prosensory cells are indeed labile, but their initial separation into distinct pools of sensory progenitors (for the cristae and the utricle at least) appears to be associated to the formation of a transient lineage-restricted tissue boundary.

Which mechanisms establish and maintain the boundary domain between segregating sensory organs? In a previous study, we showed that Lmx1a is essential for the specification of the non-sensory territories separating sensory organs and its overexpression in the chicken inner ear antagonizes prosensory specification, suggesting that it acts as a "selector" gene for the non-sensory fate (Mann et al., 2017). The present study confirms this hypothesis: in its absence (in $Lmx1a^{GFP/GFP}$ embryos), cells of the Lmx1a-lineage commit to a prosensory fate, leading to a fusion between the lateral and anterior cristae. On the other hand, the fused cristae and the utricle are separated by a Sox2-negative domain, suggesting that the initial segregation of these two sensory domains proceeds independently from Lmx1a function. Nevertheless, the typical features (large cell domain and basal constriction) of the boundary domain separating the cristae from the utricle were absent in all but one of the $Lmx1a$-null samples examined. Altogether, these results suggest that Lmx1a plays a role in the maintenance or continued differentiation of the boundary domain, but is not required for its initial formation. Finally, the patterning abnormalities in $Lmx1a^{GFP/GFP}$ samples occurred in both GFP-positive and negative territories, which points at some type of interaction between Lmx1a-expressing and non-expressing cells, and the possibility that the boundary domain is also a signalling centre influencing the differentiation of adjacent territories.

Given its conserved role in tissue boundaries, we hypothesized that actomyosin contractility could contribute to the formation or maintenance of the boundary domain. The *in vitro* results showed that short-term inhibition of ROCK-dependent actomyosin contractility with Y-27632 disrupts the morphology and patterning of boundary cells. Interestingly, the large cells were still present, but more elongated and more polarised than in control cultures. This suggests that global anisotropic forces within the developing otocyst contribute to a large extent to the elongated cellular shapes and long axis alignments observed at the interface of segregating organs. Nevertheless, the spatial separation of cells expressing either high or low levels of Sox2 was maintained. Longer-term treatments led to

an apparent fusion of adjacent sensory territories, but this may have been caused by the complete loss of epithelial integrity observed in these experiments. On the other hand, the overexpression in the chicken inner ear of a dominant-negative form of ROCK, unable to interact with its upstream regulator Shroom3 (Nishimura and Takeichi, 2008), produced more informative results. Firstly, the LC failed to segregate normally from the pan-sensory domain, indicating a requirement for ROCK activity in this process. Secondly, the border of the LC was very irregular, with some intermingling between cells expressing high and low levels of Sox2. This would be expected if an actomyosin-dependent lineage boundary had been disrupted. Altogether, these results show that ROCK-dependent coordination of actomyosin contractility is crucial for the segregation of the sensory organs and the maintenance of a straight boundary between these, yet other mechanisms are likely to contribute to the initial sorting of the Lmx1a-expressing and non-expressing cells. One strong candidate is the basal constriction at the edge of the cristae, which resembles the one described at the midbrain-hindbrain boundary (Gutzman et al., 2018, 2008) and could contribute to the physical separation of sensory progenitors. It has been proposed that a zone of non-proliferating cells could impede cell movements between adjacent compartments at the DV wing disc boundary (Blair, 1993; see however O'Brochta and Bryant, 1985 for a different view) or between rhombomeres in the hindbrain (Guthrie et al., 1991). However, the results of our Edu-incorporation assays show extensive proliferation of the cells composing the boundary domain at early and late stages of segregation, arguing against this hypothesis. Other mechanisms involved in boundary formation in the hindbrain and insect compartments include differential cell adhesion or repulsive interactions mediated by eph-ephrin signalling (Dahmann et al., 2011; Wilkinson, 2018). Further studies will be needed to determine which of these mechanisms, if any, contributes to sensory organ segregation in the inner ear.

Deep homologies between sensory organ segregation and drosophila wing disc compartmentalization: co-option of a tissue boundary module?

An increase in cortical cell tension or a multicellular actin-cable have been described along many tissue boundaries (Fagotto, 2014; Wang and Dahmann, 2020): the dorso-ventral (Major and Irvine, 2006, 2005) and antero-posterior (Landsberg et al., 2009) boundaries of the wing disc, the parasegments (Monier et al., 2010) and salivary gland placode (Röper, 2012) boundaries in drosophila and at the interface of rhombomeres of the hindbrain (Calzolari et al., 2014) in vertebrates. Hence, the implication of actomyosin-dependent processes to prevent cell intermixing is a conserved feature of tissue boundaries, which is also present in the inner ear. Perhaps more surprisingly, the core elements of the gene regulatory network regulating sensory organ segregation in the inner ear and the dorso-ventral (DV) compartmentalization of the drosophila wing disc are remarkably similar (Daudet and Żak, 2020). In fact, Notch signalling acts by lateral induction in both tissues: to maintain prosensory character in the inner ear, or to enhance Notch activity at the DV boundary in the wing disc. We have argued that *Lmx1a* and *Apterous*, which are orthologous genes,

play a selector-like function in both tissues, specifying the identity of dorsal cells in the wing disc and

that of the non-sensory cells separating the vestibular organs in the inner ear. In both tissues, the

interactions between Apterous/Lmx1a and Notch signalling have a comparable outcome: a reduction

in the levels of Notch activity within Apterous/Lmx1a expressing cells. In the wing disc, Apterous

drives the expression of Fringe, a glycosyl-transferase that makes the Notch receptor less sensitive

to the Serrate ligand in dorsal cells, potentiating signalling through Delta, expressed by ventral cells

(de Celis et al., 1996; de Celis and Bray, 1997; Doherty et al., 1996; Irvine and Wieschaus, 1994,

1994; Panin et al., 1997; Rauskolb et al., 1999). We do not know yet if any of the vertebrate Fringe

orthologues contribute to the regulation of Notch activity at the border of inner ear vestibular organs.

However, it has been shown that Lfng and Mfng are expressed at the medial border of the prosensory

domain in the embryonic mouse cochlea, and *Lnfg;Mfng* double mutants exhibit patterning defects

consistent with an abnormal positioning of the medial border of the organ of Corti (Basch et al.,

2016b). Further studies should investigate the roles of Fringe proteins during vestibular organ

segregation. By extension, it would be worth investigating the potential role of Wnt signalling in this

process, since one of the important consequences of Notch activity at the dorso-ventral wing disc

boundary is to induce the expression of Wingless, which then regulates the growth and patterning

of the wing (de Celis et al., 1996; Diaz-Benjumea and Cohen, 1995; Rulifson and Blair, 1995). It is

worth noting that a similar genetic circuit including Notch signalling (Fringe, Jagged) and Lmx1 is

deployed in vertebrate limbs to position the apical ectodermal ridge, the domain that act as a growth

zone and lineage boundary between the dorsal and the ventral ectoderm of the limb (Altabef et al.,

1997). This 'deep homology', defined as "the sharing of the genetic regulatory apparatus that is

used to build morphologically or phylogenetically disparate animal features" (Shubin et al., 2009),

suggests that Notch, Lmx1 and actomyosin contractility are part of an ancestral module for tissue

boundary formation that has been co-opted and adapted for the patterning of a great variety of

tissues.

Within the inner ear itself, it is possible that modifications of this core module are responsible for

some of the interspecies differences in sensory organ numbers and morphologies. It is thought that

the ancestral inner ear resembled that of jawless vertebrates (the lamprey and the hagfish), with only

two cristae (anterior and posterior) and a single ventral macula (Hammond and Whitfield, 2006;

Higuchi et al., 2019; Jørgensen et al., 1998; Lowenstein et al., 1968). Jawed vertebrates harbour

three semi-circular canals and cristae as well as additional sensory organs in the vestibular and

ventral part of the inner ear (Retzius, 1884, 1881), enabling in particular the detection of air-borne

sound in land vertebrates. Differences in inner ear sensory organ numbers, size and morphologies

are common place among vertebrates and it is possible that some of these originate from changes

in the genetic networks controlling sensory organ segregation. In the present study, we found at least

a clear indication of interspecies variations in the segregation of the cristae: in the chicken inner ear,

the anterior crista segregates before the lateral one, whilst in the mouse the segregation of the two

cristae occurs simultaneously. Furthermore, the chicken lateral crista is formed by two Sox2-positive

lobes, one of which turns into a non-sensory territory. The fact that the specification of prosensory cells and their segregation is regulated by signalling pathways with well-known spatial patterning activities (FGF, Wnt and Notch signalling) may have offered ample opportunities for "sensory organ experimentation", facilitating in the course of vertebrate evolution the emergence of additional organs with new functionalities, or on the contrary the suppression of redundant ones.

MATERIAL AND METHODS

Animals

Fertilised White Leghorn chicken (*Gallus gallus*) eggs were obtained from Henry Stewart UK and incubated at 37.8°C for the designated times. Embryonic stages refer to embryonic days (E), with E1 corresponding to 24 hours of incubation or to Hamburger and Hamilton stages (Hamburger and Hamilton, 1951). Embryos older than E5 were sacrificed by decapitation.

Lmx1a^{GFP} knock-in mice (Griesel et al., 2011) were maintained on a C57BL/6 mixed background at the UCL Ear Institute animal facility and bred to generate Lmx1a$^{GFP/GFP}$ (homozygous/Lmx1a null) and Lmx1a$^{GFP/+}$ (heterozygous) mice.

All animal procedures were carried out in accord with United Kingdom legislation outlined in the Animals (Scientific Procedures) Act 1986 and approved by University College London local Ethics Committee.

In ovo electroporation

Electroporation of the otic placode/cup of E2 chick embryos (stage HH 10–14) was performed using a BTX ECM 830 Electro Square Porator as previously described (Freeman et al., 2012). The RIIC2-EGFP plasmid was obtained from Nishimura, T. & Takeichi (Nishimura and Takeichi, 2008) and the insert subcloned into RCAS using conventional methods. All plasmids used for *in ovo* electroporation were purified using the 16 Qiagen Plasmid Plus Midi Kit (Qiagen). The total concentration of DNA ranged for each set of experiments between 0.5–1µg/µl. The electroporated inner ears were collected from the embryos between E5-7 and processed for whole-mount immunohistochemistry.

Immunohistochemistry

Entire chicken embryos (E3-E4) or their heads (>E5) were collected, fixed for 1.5-2 hours in 0.1 M phosphate buffered saline (PBS) with 4% paraformaldehyde and processed for whole-mount immunostaining as described previously (Mann et al., 2017). In brief, the inner ear epithelium was

dissected out from surrounding mesenchyme and incubated for 30 minutes in a blocking solution
consisting of PBS with 0.3% Triton and 10% goat serum. The primary and secondary antibodies
(diluted in PBS with 0.1% Triton, or PBT) were incubated overnight at 4C or for 2 hours at room
temperature. Several rinses in PBT were done after antibody incubation. The following reagents
were used: rabbit anti-Sox2 (Abcam, UK; 97959, 1:500); mouse IgG1 monoclonal anti-Sox2 (BD
Biosciences, San Jose, CA; 561469, 1:500); secondary goat anti-mouse IgG antibodies and
phalloidin conjugated to Alexa dyes (1:1000, Thermo Fischer Scientific UK). At the end of the
immunostaining protocol, the samples were further dissected under a Leica MZ16F
stereomicroscope equipped with a fluorescence light source to visualize precisely the position of the
Sox2-expressing sensory organs and facilitate their mounting in Vectashield (Vector labs, UK). A
fine layer of vacuum grease was applied between the slide and coverslip to avoid excessive
flattening of the tissue. Confocal stacks were acquired using a Zeiss LSM880 inverted confocal
microscope or a Perkin Elmer spinning-disc confocal and further processed with ImageJ.

Analysis of cell morphologies

Phalloidin labelling was used to visualize and analyse apical cell surfaces. Confocal z-stacks of the
surface of the samples were processed with Image J and exported into Epitool for automated cell
segmentation (Heller et al., 2016). The cell skeletons were manually corrected using CellEdit plugin
in the Icy software (de Chaumont et al., 2012). Colour scale of cell surface area and elongation ratio
were generated based on the corrected cell skeleton using CellOverlay plugin in Icy. Quantitative
data on cell morphologies were exported into Excel spreadsheet and scatter plots, radial histogram,
box plots and statistics of the data were generated using OriginPro 2017.

Organotypic cultures

The anterior domain of the otocysts and its underlying mesenchyme were dissected from E5 (HH26)
chicken embryos in room-temperature L-15 medium (Leibovitz). The dissected tissues were either
fixed at t=0 or cultured in 150-300µL DMEM medium (Invitrogen) containing 1% HEPES with or
without 20 µm Y-27632 (Sigma-Aldrich, cat # Y0503, lot # 103M4711V) in a tissue culture incubator
(5% CO_2, 37 degree) for 3h or 24hr. For the 3h-long experiments, the tissues were placed in 96-well
plates and free floating in DMEM medium. For the 24h-long experiments, the tissues were placed in
35mm Mattek dishes and coated with a small (approximately 20µl) drop of 0.4%-0.8% agarose (low
gelling temperature, Sigma-Aldrich) in 0.1M PBS before adding DMEM medium. The cultured tissues
were then fixed and processed for immunostaining as previously described.

Cell proliferation assays

Fertilized eggs were windowed at E4-5 (HH23-25) and a 200µl drop of a PBS solution containing
200µM of EdU (5-ethynyl-2´-deoxyuridine) was applied on top of the embryo. The eggs were sealed
with tape and returned to incubation for 2 hours, then the embryos heads were collected and fixed

for 30 min at room temperature in PBS with 4% Formaldehyde. Following several rinses in PBS and PBT, the inner ear tissue was dissected and processed for Click-iT EdU reaction (Thermofisher UK) according to the manufacturer's protocol. At the end of the reaction, the tissue was further processed for Sox2 immunostaining as described above and labelled with DAPI before mounting. For the mapping of mitotic figures (identified by DAPI staining), a 72x36μm "region-of-interest" centred on the boundary between the lateral crista and pan-sensory domain was selected from a confocal stack acquired with a 40x objective. The position of each mitotic figure was recorded in x,y and the results plotted for different samples collected at stage 1 (n=7) or 2-3 (n=7).

Acknowledgments

We thank Thea Støle and Caitlin Broadbent (UCL Ear Institute) for excellent technical support. This work was supported by an Action on Hearing Loss International Research Grant (G76; to ND and MZ) and the UK Research and Innovation (Medical Research Council MR/S003029/1 and Biotechnology and Biological Sciences Research Council BB/L003163/1 to ND).

Competing interests

The authors declare no competing interests

REFERENCES

Addison, M., Wilkinson, D.G., 2016. Chapter Thirty-Four - Segment Identity and Cell Segregation in the Vertebrate Hindbrain, in: Wassarman, P.M. (Ed.), Current Topics in Developmental Biology, Essays on Developmental Biology, Part B. Academic Press, pp. 581–596. https://doi.org/10.1016/bs.ctdb.2015.10.019

Alsina, B., Whitfield, T.T., 2017. Sculpting the labyrinth: Morphogenesis of the developing inner ear. Seminars in Cell & Developmental Biology. https://doi.org/10.1016/j.semcdb.2016.09.015

Altabef, M., Clarke, J.D., Tickle, C., 1997. Dorso-ventral ectodermal compartments and origin of apical ectodermal ridge in developing chick limb. Development 124, 4547–4556.

Basch, M.L., Brown, R.M., Jen, H.-I., Groves, A.K., 2016a. Where hearing starts: the development of the mammalian cochlea. J. Anat. 228, 233–254. https://doi.org/10.1111/joa.12314

Basch, M.L., Ii, R.M.B., Jen, H.-I., Semerci, F., Depreux, F., Edlund, R.K., Zhang, H., Norton, C.R., Gridley, T., Cole, S.E., Doetzlhofer, A., Maletic-Savatic, M., Segil, N., Groves, A.K., 2016b. Fine-tuning of Notch signaling sets the boundary of the organ of Corti and establishes sensory cell fates. eLife 5, e19921. https://doi.org/10.7554/eLife.19921

Blair, S.S., 1993. Mechanisms of compartment formation: evidence that non-proliferating cells do

not play a critical role in defining the D/V lineage restriction in the developing wing of Drosophila. Development 119, 339–351.

Brigande, J.V., Iten, L.E., Fekete, D.M., 2000a. A Fate Map of Chick Otic Cup Closure Reveals Lineage Boundaries in the Dorsal Otocyst. Developmental Biology 227, 256–270. https://doi.org/10.1006/dbio.2000.9914

Brigande, J.V., Kiernan, A.E., Gao, X., Iten, L.E., Fekete, D.M., 2000b. Molecular genetics of pattern formation in the inner ear: Do compartment boundaries play a role? Proceedings of the National Academy of Sciences 97, 11700–11706.

Calzolari, S., Terriente, J., Pujades, C., 2014. Cell segregation in the vertebrate hindbrain relies on actomyosin cables located at the interhombomeric boundaries. The EMBO Journal 33, 686–701. https://doi.org/10.1002/embj.201386003

Dabdoub, A., Puligilla, C., Jones, J.M., Fritzsch, B., Cheah, K.S., Pevny, L.H., Kelley, M.W., 2008. Sox2 signaling in prosensory domain specification and subsequent hair cell differentiation in the developing cochlea. Proceedings of the National Academy of Sciences 105, 18396–18401.

Dahmann, C., Oates, A.C., Brand, M., 2011. Boundary formation and maintenance in tissue development. Nature Reviews Genetics 12, 43–55. https://doi.org/10.1038/nrg2902

Daudet, N., Żak, M., 2020. Notch Signalling: The Multitask Manager of Inner Ear Development and Regeneration. Adv. Exp. Med. Biol. 1218, 129–157. https://doi.org/10.1007/978-3-030-34436-8_8

de Celis, J.F., Bray, S., 1997. Feed-back mechanisms affecting Notch activation at the dorsoventral boundary in the Drosophila wing. Development 124, 3241–3251.

de Celis, J.F., Garcia-Bellido, A., Bray, S.J., 1996. Activation and function of Notch at the dorsal-ventral boundary of the wing imaginal disc. Development 122, 359–369. https://doi.org/10.1242/dev.122.1.359

de Chaumont, F., Dallongeville, S., Chenouard, N., Hervé, N., Pop, S., Provoost, T., Meas-Yedid, V., Pankajakshan, P., Lecomte, T., Le Montagner, Y., Lagache, T., Dufour, A., Olivo-Marin, J.-C., 2012. Icy: an open bioimage informatics platform for extended reproducible research. Nat Methods 9, 690–696. https://doi.org/10.1038/nmeth.2075

Diaz-Benjumea, F.J., Cohen, S.M., 1995. Serrate signals through Notch to establish a Wingless-dependent organizer at the dorsal/ventral compartment boundary of the Drosophila wing. Development 121, 4215–4225. https://doi.org/10.1242/dev.121.12.4215

Doherty, D., Feger, G., Younger-Shepherd, S., Jan, L.Y., Jan, Y.N., 1996. Delta is a ventral to dorsal signal complementary to Serrate, another Notch ligand, in Drosophila wing formation. Genes Dev 10, 421–434. https://doi.org/10.1101/gad.10.4.421

Ekdale, E.G., 2016. Form and function of the mammalian inner ear. Journal of Anatomy 228, 324–337. https://doi.org/10.1111/joa.12308

Fagotto, F., 2014. The cellular basis of tissue separation. Development 141, 3303–3318.

https://doi.org/10.1242/dev.090332

Fekete, D.M., 1996. Cell fate specification in the inner ear. Current opinion in neurobiology 6, 533–541.

Fekete, D.M., Wu, D.K., 2002. Revisiting cell fate specification in the inner ear. Current Opinion in Neurobiology 12, 35–42. https://doi.org/10.1016/S0959-4388(02)00287-8

Freeman, S., Chrysostomou, E., Kawakami, K., Takahashi, Y., Daudet, N., 2012. Tol2-Mediated Gene Transfer and In Ovo Electroporation of the Otic Placode: A Powerful and Versatile Approach for Investigating Embryonic Development and Regeneration of the Chicken Inner Ear, in: Mace, K.A., Braun, K.M. (Eds.), Progenitor Cells, Methods in Molecular Biology. Humana Press, pp. 127–139. https://doi.org/10.1007/978-1-61779-980-8_10

Fritzsch, B., Beisel, K.W., Jones, K., Fariñas, I., Maklad, A., Lee, J., Reichardt, L.F., 2002. Development and evolution of inner ear sensory epithelia and their innervation. J. Neurobiol. 53, 143–156. https://doi.org/10.1002/neu.10098

Garcia-Bellido, A., Ripoll, P., Morata, G., 1973. Developmental Compartmentalisation of the Wing Disk of Drosophila. Nature 245, 251–253. https://doi.org/10.1038/10.1038/newbio245251a0

Guthrie, S., Butcher, M., Lumsden, A., 1991. Patterns of cell division and interkinetic nuclear migration in the chick embryo hindbrain. J Neurobiol 22, 742–754. https://doi.org/10.1002/neu.480220709

Gutzman, J.H., Graeden, E., Brachmann, I., Yamazoe, S., Chen, J.K., Sive, H., 2018. Basal constriction during midbrain-hindbrain boundary morphogenesis is mediated by Wnt5b and focal adhesion kinase. Biol Open 7, bio034520. https://doi.org/10.1242/bio.034520

Gutzman, J.H., Graeden, E.G., Lowery, L.A., Holley, H.S., Sive, H., 2008. Formation of the zebrafish midbrain–hindbrain boundary constriction requires laminin-dependent basal constriction. Mechanisms of Development 125, 974–983. https://doi.org/10.1016/j.mod.2008.07.004

Hamburger, V., Hamilton, H.L., 1951. A series of normal stages in the development of the chick embryo. Journal of morphology 88, 49–92.

Hammond, K.L., Whitfield, T.T., 2006. The developing lamprey ear closely resembles the zebrafish otic vesicle: otx1 expression can account for all major patterning differences. Development 133, 1347–1357. https://doi.org/10.1242/dev.02306

Heller, D., Hoppe, A., Restrepo, S., Gatti, L., Tournier, A.L., Tapon, N., Basler, K., Mao, Y., 2016. EpiTools: An Open-Source Image Analysis Toolkit for Quantifying Epithelial Growth Dynamics. Developmental Cell 36, 103–116. https://doi.org/10.1016/j.devcel.2015.12.012

Higuchi, S., Sugahara, F., Pascual-Anaya, J., Takagi, W., Oisi, Y., Kuratani, S., 2019. Inner ear development in cyclostomes and evolution of the vertebrate semicircular canals. Nature 565, 347–350. https://doi.org/10.1038/s41586-018-0782-y

Hughes, S.H., 2004. The RCAS vector system. FOLIA BIOLOGICA-PRAHA- 50, 107–119.

Irvine, K.D., Wieschaus, E., 1994. fringe, a boundary-specific signaling molecule, mediates

interactions between dorsal and ventral cells during Drosophila wing development. Cell 79, 595–606. https://doi.org/10.1016/0092-8674(94)90545-2

Jørgensen, J.M., Shichiri, M., Geneser, F.A., 1998. Morphology of the Hagfish Inner Ear. Acta Zoologica 79, 251–256. https://doi.org/10.1111/j.1463-6395.1998.tb01163.x

Kiernan, A.E., 2013. Notch signaling during cell fate determination in the inner ear. Seminars in Cell & Developmental Biology 24, 470–479. https://doi.org/10.1016/j.semcdb.2013.04.002

Kiernan, A.E., Pelling, A.L., Leung, K.K.H., Tang, A.S.P., Bell, D.M., Tease, C., Lovell-Badge, R., Steel, K.P., Cheah, K.S.E., 2005. Sox2 is required for sensory organ development in the mammalian inner ear. Nature 434, 1031–1035. https://doi.org/10.1038/nature03487

Kil, S.-H., Collazo, A., 2002. A review of inner ear fate maps and cell lineage studies. Journal of Neurobiology 53, 129–142. https://doi.org/10.1002/neu.10127

Knowlton, V.Y., 1967. Correlation of the development of membranous and bony labyrinths, acoustic ganglia, nerves, and brain centers of the chick embryo. J. Morphol. 121, 179–207. https://doi.org/10.1002/jmor.1051210302

Koo, S.K., Hill, J.K., Hwang, C.H., Lin, Z.S., Millen, K.J., Wu, D.K., 2009. Lmx1a maintains proper neurogenic, sensory, and non-sensory domains in the mammalian inner ear. Developmental Biology 333, 14–25. https://doi.org/10.1016/j.ydbio.2009.06.016

Landsberg, K.P., Farhadifar, R., Ranft, J., Umetsu, D., Widmann, T.J., Bittig, T., Said, A., Jülicher, F., Dahmann, C., 2009. Increased Cell Bond Tension Governs Cell Sorting at the Drosophila Anteroposterior Compartment Boundary. Current Biology 19, 1950–1955. https://doi.org/10.1016/j.cub.2009.10.021

Langenberg, T., Brand, M., 2005. Lineage restriction maintains a stable organizer cell population at the zebrafish midbrain-hindbrain boundary. Development 132, 3209–3216. https://doi.org/10.1242/dev.01862

Lowenstein, O., Osborne, M.P., Thornhill, R.A., 1968. The Anatomy and Ultrastructure of the Labyrinth of the Lamprey (Lampetra fluviatilis L.). Proceedings of the Royal Society of London B: Biological Sciences 170, 113–134. https://doi.org/10.1098/rspb.1968.0029

Major, R.J., Irvine, K.D., 2006. Localization and requirement for Myosin II at the dorsal-ventral compartment boundary of the Drosophila wing. Developmental Dynamics 235, 3051–3058. https://doi.org/10.1002/dvdy.20966

Major, R.J., Irvine, K.D., 2005. Influence of Notch on dorsoventral compartmentalization and actin organization in the Drosophila wing. Development 132, 3823–3833. https://doi.org/10.1242/dev.01957

Mann, Z.F., Gálvez, H., Pedreno, D., Chen, Z., Chrysostomou, E., Żak, M., Kang, M., Canden, E., Daudet, N., 2017. Shaping of inner ear sensory organs through antagonistic interactions between Notch signalling and Lmx1a. Elife 6. https://doi.org/10.7554/eLife.33323

Monier, B., Pélissier-Monier, A., Brand, A.H., Sanson, B., 2010. An actomyosin-based barrier inhibits cell mixing at compartmental boundaries in Drosophila embryos. Nature Cell

Biology 12, 60–65. https://doi.org/10.1038/ncb2005

Monier, B., Pélissier-Monier, A., Sanson, B., 2011. Establishment and maintenance of
compartmental boundaries: role of contractile actomyosin barriers. Cell. Mol. Life Sci. 68,
1897–1910. https://doi.org/10.1007/s00018-011-0668-8

Nichols, D.H., Pauley, S., Jahan, I., Beisel, K.W., Millen, K.J., Fritzsch, B., 2008. Lmx1a is required
for segregation of sensory epithelia and normal ear histogenesis and morphogenesis. Cell
and Tissue Research 334, 339–358. https://doi.org/10.1007/s00441-008-0709-2

Nishimura, T., Takeichi, M., 2008. Shroom3-mediated recruitment of Rho kinases to the apical cell
junctions regulates epithelial and neuroepithelial planar remodeling. Development 135,
1493–1502. https://doi.org/10.1242/dev.019646

Norris, H.W., 1892. Studies on the development of the ear of amblystoma. Part I.: Development of
the auditory vesicle. J. Morphol. 7, 23–34. https://doi.org/10.1002/jmor.1050070103

O'Brochta, D.A., Bryant, P.J., 1985. A zone of non-proliferating cells at a lineage restriction
boundary in Drosophila. Nature 313, 138–141. https://doi.org/10.1038/313138a0

Ohta, S., Schoenwolf, G.C., 2018. Hearing crosstalk: the molecular conversation orchestrating
inner ear dorsoventral patterning. WIREs Developmental Biology 7, e302.
https://doi.org/10.1002/wdev.302

Panin, V.M., Papayannopoulos, V., Wilson, R., Irvine, K.D., 1997. Fringe modulates Notch-ligand
interactions. Nature 387, 908–912.

Rauskolb, C., Correia, T., Irvine, K.D., 1999. Fringe-dependent separation of dorsal and ventral
cells in the Drosophila wing. Nature 401, 476. https://doi.org/10.1038/46786

Retzius, G., 1884. Das Gehororgan der Reptilien, der Vogel und der Saugetiere. In: Das
Gehorogan der Wirbeltiere, vol 2. Samson and Wallin, Stockholm.

Retzius, G., 1881. Das Gehörorgan der Fische und Amphibien. In: Das Gehörorgan der
Wirbelthiere, vol 1. Samson and Wallin, Stockholm.

Röper, K., 2012. Anisotropy of Crumbs and aPKC Drives Myosin Cable Assembly during Tube
Formation. Developmental Cell 23, 939–953. https://doi.org/10.1016/j.devcel.2012.09.013

Rulifson, E.J., Blair, S.S., 1995. Notch regulates wingless expression and is not required for
reception of the paracrine wingless signal during wing margin neurogenesis in Drosophila.
Development 121, 2813–2824. https://doi.org/10.1242/dev.121.9.2813

Sánchez-Guardado, L.Ó., Puelles, L., Hidalgo-Sánchez, M., 2014. Fate map of the chicken otic
placode. Development 141, 2302–2312. https://doi.org/10.1242/dev.101667

Sánchez-Guardado, L.Ó., Puelles, L., Hidalgo-Sánchez, M., 2013. Fgf10 expression patterns in the
developing chick inner ear. Journal of Comparative Neurology 521, 1136–1164.
https://doi.org/10.1002/cne.23224

Shubin, N., Tabin, C., Carroll, S., 2009. Deep homology and the origins of evolutionary novelty.
Nature 457, 818–823. https://doi.org/10.1038/nature07891

Smith, D.M., Tabin, C.J., 2000. Clonally Related Cells Are Restricted to Organ Boundaries Early in

the Development of the Chicken Gut to Form Compartment Boundaries. Developmental Biology 227, 422–431. https://doi.org/10.1006/dbio.2000.9874

Steevens, A.R., Glatzer, J.C., Kellogg, C.C., Low, W.C., Santi, P.A., Kiernan, A.E., 2019. SOX2 is required for inner ear growth and cochlear nonsensory formation before sensory development. Development 146, dev170522. https://doi.org/10.1242/dev.170522

Steevens, A.R., Sookiasian, D.L., Glatzer, J.C., Kiernan, A.E., 2017. SOX2 is required for inner ear neurogenesis. Scientific Reports 7, 4086. https://doi.org/10.1038/s41598-017-04315-2

Steffes, G., Lorente-Cánovas, B., Pearson, S., Brooker, R.H., Spiden, S., Kiernan, A.E., Guénet, J.-L., Steel, K.P., 2012. Mutanlallemand (mtl) and Belly Spot and Deafness (bsd) Are Two New Mutations of Lmx1a Causing Severe Cochlear and Vestibular Defects. PLoS ONE 7, e51065. https://doi.org/10.1371/journal.pone.0051065

Wang, J., Dahmann, C., 2020. Establishing compartment boundaries in Drosophila wing imaginal discs: An interplay between selector genes, signaling pathways and cell mechanics. Seminars in Cell & Developmental Biology. https://doi.org/10.1016/j.semcdb.2020.07.008

Wilkinson, D.G., 2018. Establishing sharp and homogeneous segments in the hindbrain. F1000Research 7, 1268. https://doi.org/10.12688/f1000research.15391.1

Żak, M., Daudet, N., 2021. A gradient of Wnt activity positions the neurosensory domains of the inner ear. eLife 10, e59540. https://doi.org/10.7554/eLife.59540