Title: Unifying adaptive stress and adaptive habitat selection hypotheses through movement ecology

Authors: Newediuk, Levi[1], Mastromonaco, Gabriela F.[2], & Vander Wal, Eric[1]

[1] Department of Biology, Memorial University, St. John's Newfoundland, Canada A1B 3X9

[2] Reproductive Physiology, Toronto Zoo, Toronto Ontario, Canada M1B 5K7

Corresponding Author: Levi Newediuk

email: ljnewediuk@mun.ca; ORCiD: https://orcid.org/0000-0003-0719-9596

Abstract

Habitat selection is adaptive, but habitats where animals exhibit elevated glucocorticoid production are often associated with lower fitness. The perception of fitness costs persists in part because glucocorticoid samples rarely represent the internal state of individuals at the time of sampling, making it difficult to demonstrate a causal link between glucocorticoid production and habitat selection at fine temporal and spatial scales. We show how to test this fine-scale causal link by integrating movement ecology with stress physiology. We first synthesize physiological evidence that effect of glucocorticoid production on movement aligns with habitat selection theory. We distinguish whether habitat selection exposes individuals to stressors that elevate their glucocorticoid levels, or whether elevated glucocorticoids stimulate selection for better habitat. We then test the directionality of these two pathways using movement-based habitat selection analysis at the individual level. We capitalize on a combination of individual movement data and the predictable time required for glucocorticoids to peak in samples like hair, feathers, and feces from individuals. Our proof-of-concept shows how to distinguish when glucocorticoids stimulate adaptive habitat selection. Finally, we suggest a potential evolutionary role for glucocorticoids in habitat specialization, offering a mechanism by which populations might adapt to their environments over generations.

Keywords: Glucocorticoids, stress physiology, optimal foraging theory, integrated step selection analysis, environmental challenges, non-invasive sampling, individual differences

1 Introduction

Animals are adapted to their environments, both physiologically through glucocorticoid

production and behaviourally through habitat selection. Habitat selection is the process by which

animals use some habitats disproportionately to their availability. For example, animals might

relocate to sparser foraging patches that maximize energetic intake [1] while mitigating the stress

of competition for forage [2]. Movement between foraging patches is driven in part by

glucocorticoid hormone production. Glucocorticoids stimulate appetite when energy reserves are

low [3] and make energy available for movement to find foraging resources [4]. Their role in

energy regulation makes glucocorticoids a key indicator for internal state, which is integral to

movement and habitat selection [5]. But glucocorticoid production is also associated with

exposure to stressors that can compromise fitness [6]. Glucocorticoids have pleiotropic effects

under stress, for example reducing body mass when they stimulate catabolism of energy reserves

[7], and lowering reproductive success when they stimulate behaviours that interfere with

reproduction [8]. While these costs usually only arise when animals face chronic and inescapable

stressors [9], studies considering possible adaptive benefits of glucocorticoids are the minority.

The benefits of glucocorticoid production are least considered when their production coincides

with selection of stressful habitats.

Glucocorticoids decrease fitness directly when their long-term production becomes

physiologically harmful or otherwise compromises reproduction or survival. Reproductive

output, for example, declines after long-term experimental exposure to stressors [10]. Oxidative

stress can accumulate even after short-term glucocorticoid exposure [11]. Though the

physiological and behavioural consequences of glucocorticoid production can compromise

fitness, glucocorticoid production is not inherently pathological. In fact, its production is often an

adaptive response to stressors faced in the wild [12]. For example, depletion of body reserves is

associated with elevated glucocorticoids [13,14], but breakdown of protein and lipid reserves are

is not always harmful over longer time scales. For example, when food availability is low or

energetic demand is high, mobilizing lipid and protein reserves prevents starvation [7].

Sacrificing body condition might eventually compromise survival, but lifetime reproductive

success may also be higher for animals that select habitat favouring their own survival when

conditions are not favourable for offspring survival [15]. Quantifying survival and reproductive

success associated with glucocorticoid production can help distinguish when glucocorticoid-

driven habitat selection is adaptive. However, the distinction requires knowing whether

glucocorticoids caused habitat selection, or habitat selection caused glucocorticoid production.

One way to establish the causal direction between glucocorticoids and habitat selection is

to identify the timing of habitat selection relative to associated glucocorticoid production. When

glucocorticoids are already elevated before moving to a new habitat, stressors in the selected

habitat could not have caused their production. Instead, glucocorticoid production might have

driven selection of the habitat, motivated by stressors experienced elsewhere. For example,

migratory animals elevate glucocorticoids in spring as stressors intensify on winter ranges [16],

energetically preparing them for migration to summer ranges [17]. Thus, during migration the

glucocorticoids causing habitat selection are adaptive. In contrast, when glucocorticoids become

elevated after selecting habitat, their production might represent reactions to stressors. For

example, sandgrouse (*Pterocles alchata*) inhabiting a recreational area had elevated

glucocorticoids only after weekends with high tourist traffic, suggesting human disturbance

triggered their stress responses [18].

The sequence from glucocorticoid production to habitat selection is clearer when

glucocorticoids are elevated in response to measurable stressors like human disturbance.

However, many natural stressors facing wild animals are not readily observable. For example,

when a wild animal selects a risky habitat where it briefly encounters a predator, the

hypothalamic-pituitary-adrenal axis might stimulate an acute stress response. Eventually,

glucocorticoids produced during the stress response can be measured in non-invasive samples

like hair and feces. However, non-invasive samples are difficult to age and could represent

stressors that occurred hours, days, or even weeks in the past [19]. The temporal disconnect

between stressor and sample makes the exact timing of the stressor, and therefore the relative

order of habitat selection and glucocorticoid production, ambiguous.

The spatial scale at which non-invasive samples are collected also complicates the causal

relationship between glucocorticoids and habitat selection. Glucocorticoids are primarily

responsible for fine-scale habitat selection decisions by individuals at the so-called second and

third orders of selection (*sensu* Johnson [20]). However, the difficult task of linking individuals

to non-invasive samples in the field [21] limits inferences to the effects of population-average

glucocorticoids on coarser-scale habitat selection (i.e., first order selection [20]). These coarse-

scale population-level comparisons create uncertainty around which stressors cause habitat

selection. For example, predator abundance can be compared between habitats, but assumed

stress responses to predation risk may instead be caused by other associated stressors like food

shortages [22]. Furthermore, individual differences in diet and glucocorticoid metabolism

between populations, rather than stressors per se, may result in different glucocorticoid levels

between them [23]. The consequence is that glucocorticoids and habitat selection may be either

correlated or share a causal relationship. Still, the most common interpretation of any

relationship between them is that selection of a stressful habitat both caused glucocorticoid

production and reduced fitness (Table 1).

We show how the movement ecology lens (*sensu* [5]) can help distinguish adaptive

production of glucocorticoids from their purported harmful effects. To demonstrate, we first

synthesize evidence that glucocorticoids are mechanistically responsible for adaptive habitat

selection behaviour. Our examples show how habitat selection changes when animals in different

life-history stages manipulate glucocorticoid levels to support either reproduction or survival.

Next, we discuss the time lag between stressor and peak glucocorticoid concentrations in tissues

like blood, feces, and hair. Normally considered a challenge for researchers aiming to connect

glucocorticoids with movement and habitat selection, we demonstrate how the consistency of

this time lag presents an opportunity to pinpoint the stressor responsible for habitat selection.

Once the timing of the stressor and individual identity of the sample are known, movement-

based analyses can test causal links between glucocorticoids and adaptive habitat selection

responses to the stressor. We demonstrate this analysis with a proof-of-concept. Our proof-of-

concept acts as a template for promising research questions to which the same approach could be

applied in the future.

Physiology of glucocorticoids as a mechanism for movement

Patterns of glucocorticoid production correlate with movement activity over daily and seasonal

time periods. Baseline glucocorticoid levels, regulated by the autonomic nervous system [24],

coincide with energy needs for locomotion at predictable times. For example, circadian activity

patterns in mice coincide with the rise and fall of circulating glucocorticoids [25], and in turn,

glucocorticoid rhythms adapt to regular feeding periods when energy is available for movement

[26]. The interplay among energy availability, movement, and these baseline glucocorticoid

levels is a mechanistic explanation for diel habitat selection patterns that help prey species access

forage while avoiding their predators [27]. Seasonal activity patterns like migration track longer-

term baseline glucocorticoid rhythms [28]. Because baseline glucocorticoid levels are seasonally

elevated during migration [17], birds are primed to adjust movement quickly in response to cues

like wind assistance [29] and stressors upon arrival at migration stopover sites [30]. Non-

migratory birds also elevate baseline glucocorticoids during reproduction when offspring require

the most active provisioning [6]. Thus, via extension of their effects on activity and locomotion,

glucocorticoids form the physiological connection between environmental pressures and

adaptive patterns of behaviour to overcome them.

Environmental pressures, like predation risk, induce stress responses that make prey

species avoid risky foraging habitat. Predator cues stimulate a cascade of hormone release via the

hypothalamic-pituitary-adrenal (HPA) axis, the main products of which include glucocorticoids

and other faster-acting hormones such as catecholamines. While catecholamines are mainly

involved in increasing blood pressure, heart rate, and driving other more immediate responses to

stress [31], glucocorticoids have a primary role in helping to regulate blood glucose levels.

Glucose is mobilized when glucocorticoids prevent its uptake by adipose and muscle tissue [32]

and stimulate gluconeogenesis [4], thereby providing needed energy to forgo foraging while

escaping predators. Continued effects of elevated predation risk on glucocorticoid production can

have long-term effects on habitat selection patterns. For example, decadal shifts in predation

pressure by grey seals in Atlantic Canada caused Atlantic cod (*Gadus morhua*) and white hake

(*Urophycis tenuis*) to redistribute to safer areas with less food [33]. In such cases, the potential

cost of predation outweighs the value of energy gain.

The continued need to replenish calories under long-term predation risk, however, can

also encourage selection for risky habitats that offer more productive foraging. As risk attenuates

after a stress response, glucocorticoid levels return to baseline levels through a combination of

negative feedback on the HPA axis and reduced glucocorticoid production. When reduction in

HPA axis activity follows a stress response, still-elevated baseline levels stimulate feeding

behaviour [4] and a preference for high-calorie foods [3]. If lower feeding rates during the stress

response eventually compromise energy storage, peripheral satiety signals also decline and

animals seek to replace energy stores [4]. For wildlife, replacing energy stores means movement

to foraging habitats. Consequent hunger encourages animals exposed to chronic risk to spend

even more time foraging [34]. Preferences for richer foods are reinforced over time because their

consumption gradually blunts mounting of stress responses [35], keeping baseline glucocorticoid

levels in a slightly elevated state. Similarly, predation risk elevates respiration rate, changing

C:N ratios of animal tissue and stimulating selection of forage with greater carbon content [36].

Over time, habitat selection patterns emerge that prioritize energy gain over avoidance of

predation risk. For example, rather than moving to unproductive areas with less poaching,

elephant groups in Kenya responded to intense ivory poaching by moving to ranges with higher

primary productivity [37]. Thus, glucocorticoid-stimulated movement mediates trade-offs

between energy-motivated and risk-avoidant habitat selection, which is ultimately adaptive.

Integrating glucocorticoids into individual habitat selection

Testing when glucocorticoids stimulate adaptive habitat selection is difficult in the wild because

glucocorticoid samples are imprecise measures of stressors faced by individuals. Precision comes

from placing movement appropriately along the timeline from individual glucocorticoid

production to sample collection. Glucocorticoids are incorporated into growing hair cells from

the plasma, and depending on the rate of hair growth, represent circulating levels over periods

from weeks to months [38]. Feathers also require weeks or months to incorporate

glucocorticoids, and only do so during growth periods prior to moult [19]. Fecal samples

represent circulating glucocorticoids processed by the liver and deposited as metabolites within

several hours to days depending on species-specific gut transport times (reviewed in [39]). In

contrast, the temporal resolution of biotelemetry data is often hourly or finer, making

glucocorticoid samples a relatively coarse representation of stressors faced during habitat

selection. Moreover, samples are often collected without knowledge of individual identity. The

mismatch between habitat selection and glucocorticoid production makes it necessary to pool

population movement data over weeks, months, or even years. The more data are pooled,

however, the more challenging it becomes to assess whether glucocorticoid production is a cause

or consequence of habitat selection behaviour. Addressing the problem of causation requires

validating the time required for glucocorticoids to peak in samples after stressors, then

determining whether individual behaviour occurred before or after the hormone peak.

Determining whether behaviours caused or were caused by elevated glucocorticoid levels

is especially important in the wild where stress responses are often inferred instead of observed.

The general expectation is that perceived stressor severity determines the quantity of

glucocorticoids produced in response [40]. However, the bioactive component of

glucocorticoids, and not the overall amount produced in response to stressors, triggers

behavioural responses. Habitat selection behaviours driven by baseline glucocorticoid levels, like

foraging and migration, arise when glucocorticoids bind to low-affinity mineralocorticoid (MR)

receptors [28]. For immediate behavioural effects during a stress response, glucocorticoids bind

high-affinity glucocorticoid (GR) receptors [41]. Receptor binding is controlled in part by

corticosterone binding globulins (CBGs) which bind glucocorticoids in the plasma and prevent

them binding receptors [42]. As a result, the effects of glucocorticoids on behaviour saturate

when either high CBG concentrations, downregulated receptor transcription [43], or negative

feedback on the stress response slow receptor binding rates downstream [44]. Estimating the

amount of circulating glucocorticoids available to bind to receptors, however, requires either

blood samples (e.g., [42]) or experimental hormone challenges (e.g., [45]). The controlled

conditions required for these protocols make it difficult to link CBGs, receptors, and negative

feedback to habitat selection under natural conditions. Alternatively, modelling individual

glucocorticoid-habitat selection relationships could reveal when these downstream mediators

adjust how the hormones affect behaviour. This is because when downstream mediators interact

with glucocorticoids at high concentrations, their relationship with behaviour becomes nonlinear.

For example, Adélie penguin (*Pygoscelis adeliae*) locomotory behaviour had a U-shaped

relationship with glucocorticoid supplementation, where glucocorticoid effects on behaviour at

high circulating concentrations were buffered by high CBG concentrations [46].

Adjusting behavioural responses to glucocorticoids downstream is beneficial for some

individuals and during some seasons. For example, as day lengths shorten in winter, the effect of

cortisol supplementation on white-crowned sparrow (*Zonotrichia leucophrys gambelii*) activity

saturates [47]. Saturating effects of glucocorticoids on behaviour are particularly common during

reproduction [43], when habitat selection must adjust to meet the needs of breeding individuals

and those provisioning offspring [48]. Individuals having more experience with a stressor

sometimes develop stronger negative feedback to prevent pathology from chronic stress [49].

The result is that the same stressor might produce different peak glucocorticoid levels in two

individuals or differences in how glucocorticoid effects are mediated downstream [50]. These

differences might manifest in different adaptive habitat selection patterns by individual or by

season in response to the environment.

Differences in how individuals respond to the environment pose a problem for

disentangling cause and effect because they introduce variation into the glucocorticoid-habitat

selection relationship. To control for this variation, individual movement must be linked to

glucocorticoid samples collected from individual animals at known times. While repeated

sampling of individual animals is often infeasible in the wild, biotelemetry data can help link

individuals with opportunistically collected field samples. Feathers, hair, and feces contain

genetic material in addition to glucocorticoids, allowing anonymous samples from populations to

be matched to individuals with genetic data using genetic capture-mark-recapture techniques

[39]. Conspicuous species like ungulates can be observed defecating or shedding hair [51]. When

individuals also carry biotelemetry tags, areas like trails and locations of kill or rest sites can be

targeted for sampling specific individuals (e.g., [52]), especially when fine-scale timestamped

location data link individual movements to their circulating glucocorticoids during the same

period [21].

When the timing of sample collection, estimated age of the sample, and the time required

for glucocorticoids to peak in hair, feces, or feathers are known, the approximate timing of

stressors can also be pinpointed. Movement-based models can then be used to interpret whether

the stressor caused or was a product of habitat selection. One possible interpretation of elevated

glucocorticoids is that the habitat selected by an animal exposed it to a stressor that affected its

circulating glucocorticoid levels. This interpretation might be tested with double hierarchical

models that include a coefficient for movement characteristics describing behaviours related to

foraging (e.g., tortuous movements) over the "steps" preceding the glucocorticoid peak (outlined

in [53]). Another interpretation is that glucocorticoid levels became elevated by the stressor,

which then stimulated habitat selection. Habitat selection analysis might be used to test this

interpretation. Integrated step selection analysis [54], a specialized class of habitat selection

analysis, can be used to test whether glucocorticoids produced before the beginning of a step

affect the relative probability of habitat selection at its end (Box 1). In the following section, we

provide a proof-of-concept demonstrating this approach.

Proof-of-concept: Glucocorticoid-mediated fine-scale habitat selection by elk

Background

For species that provide parental care, the fitness consequences of habitat selection manifest

around parturition as an energetically demanding life stage. Female ungulates must both

maintain sufficient body condition prior to calving and compensate for the demanding energetic

costs of spring lactation via summer nutrition [55]. In part, these nutritional needs are met by

shifting habitat use toward better forage. Increased used of forage rich habitat also coincides with

elevated baseline glucocorticoid levels typical of ungulate calving periods [56]. However, calves

are vulnerable to predation and in some species of ungulate are typically hidden for a few weeks

following birth. To ensure reproductive success, mothers must compromise some of their

nutritional needs for safety needs of the calf [57]. Safety typically requires use of cover habitat

like forests and shrubland despite better foraging opportunities elsewhere [58].

Methods

We used integrated step selection analysis (Box 1; [54]) to test whether selection for cover

habitat by 13 female elk (*Cervus canadensis*) shifted in response to fecal glucocorticoid

metabolite levels, and whether the response changed during the approximate 16-d window when

their calves are most vulnerable (see Supplementary 1 for detailed methods). We collected fecal

glucocorticoid samples, which for elk correspond to circulating levels approximately 20 hours

prior to defecation [59,60]. Our global positioning system-collared study population in

southeastern Manitoba, Canada (49.134, -96.557) uses forest as cover habitat [61] to safely avoid

humans in the largely agricultural landscape that also acts as a source of calving season forage

[48]. Because animals forage to recover from stressors [4], we predicted that prior to calving and

during the approximately 20 hours following glucocorticoid peaks, elk would select locations

farther from cover. Foraging to recover energy after stress responses risks calf safety by

exposing them to potential encounters with predators. However, lactating elk also require more

energy. Thus, during the 16-d window of calf vulnerability we expected elk to use locations

either farther from or closer to cover habitat depending on their priority for energy versus safety.

If elk prioritized safety, those with young calves should remain closer to cover habitat even after

glucocorticoid production increased. Alternatively, if they prioritized energy, they might have an

even stronger preference for areas further from cover when glucocorticoid production increased.

Importantly, a change in behaviour after calving might indicate either CBG concentration,

receptor number, negative feedback, or all three are adjusted downstream to support calf

survival.

Results & Discussion

Elk habitat selection differed with glucocorticoid levels and presence of young calves. Before

calving and when calves were able to escape predators, their mothers were 50% more likely to

select locations farther from cover habitat for each unit increase in glucocorticoids (Fig. 2, "pre"

period). Elevated glucocorticoids increased the likelihood that elk selected locations further from

cover even more strongly within the 16-d window when calves were most vulnerable (Fig. 2,

"post" period). These results suggest the non-linear effects of glucocorticoids on habitat selection

may be mediated post-partum either by CBG concentration, receptor availability, or negative

feedback on glucocorticoid production. Thus, changes in glucocorticoid-mediated habitat

selection by our elk population characterize life-history transitions, making glucocorticoid

production and its downstream mediation an adaptive process that supports lifetime reproductive

success. See Supplementary 2 for comprehensive results.

Conclusions and future directions

One of the most pervasive ideas about glucocorticoid overproduction is that it is an indicator of

exposure to stressors that decrease fitness [6]. We interpreted the fitness implications of

glucocorticoid production through its relationship to adaptive behaviours like habitat selection.

In our proof-of-concept, elk with elevated glucocorticoid levels selected locations further from

cover habitat, suggesting a preference for forage-rich agriculture. Selection for forage aligns with

physiological evidence that glucocorticoids help animals recover energetically after a stress

response [4]. When they had young calves, elk with elevated glucocorticoids exhibited even

stronger selection for locations further from cover, suggesting they prioritized energy over

remaining close to cover habitat for their calves (Fig. 1b). We suggest the change in behaviour

associated with care for young calves might have been brought about by physiological changes

that altered the downstream effects of glucocorticoids. We demonstrated that glucocorticoid

production and downstream mediation are adaptive with a modelling approach that could

demonstrate a causal relationship between glucocorticoid production and habitat selection. Key

to our inference was knowing the time required for glucocorticoids to peak in fecal pellet

samples and which samples belonged to which individuals. These two pieces of information,

already available or collected in many studies, hold promise for testing existing hypotheses in

habitat selection and stress physiology and for exploring new links between individuals and their

environments (Box 2).

One promising extension of our approach is to test whether adjusting habitat selection

through glucocorticoid production helps individuals and populations respond to stressors over

multiple generations. Individuals respond to stressors by adjusting glucocorticoid production

over their lifetimes to support either survival or reproduction. For example, as they age and

fewer reproductive opportunities remain, individual terns (*Sterna hirundo*) produce fewer

glucocorticoids to support reproductive behaviours [62]. These age-dependent changes in

reproduction are adaptive at the population level because they align with age-dependent

reproductive investment as predicted by life-history theory. Individuals also differ in their

glucocorticoid responses to the same stressors [63]. These stress-coping mechanisms are

transmissible to offspring [64], meaning selection might act on habitat-specific glucocorticoid

production as individuals are exposed to different stressors in the habitats they occupy [65].

Disruptive selection on both stress-coping mechanisms and habitat selection could reinforce

differences among individuals, resulting in both the physiological and spatial separation of

populations over time [66]. For example, frogs adapted to habitats with long-term exposure to

anthropogenic noise both produced fewer glucocorticoids in response to noise and had developed

physiological coping mechanisms to equalize fitness with their noise-free habitat counterparts

[67]. The evolutionary effects of glucocorticoid-habitat selection feedback might be preceded by

habitat specialization at smaller scales, likely testable using our movement-based approach (Box

2).

Our synthesis highlighted a limitation in the way we currently interpret associations

between habitat selection and glucocorticoids: namely that we rarely consider glucocorticoids an

agent of adaptive habitat selection. The negative association between habitat selection and

glucocorticoid production persists because it is more challenging to test when glucocorticoids

drive adaptive habitat selection. We showed the tools to test an adaptive relationship between the

two already exist in study systems where ecologists collect biotelemetry data and non-invasive

glucocorticoid samples. The challenges going forward will be to integrate glucocorticoids into

movement ecology and to explore alternative interpretations about glucocorticoid production in

the context of habitat selection.

Conflicts of interest: The authors have no conflicts of interest to report.

Author contributions: LN and EVW conceived of the ideas. LN designed the proof-of-concept

methodology, analyzed the data, and wrote the original draft of the manuscript. GFM oversaw

hormone extractions. All authors contributed to reviewing & editing and approved the final

manuscript for publication.

Funding: This work was supported by a funding agreement from the Manitoba Fish and Wildlife

Enhancement Fund (#5951) awarded to EVW and LN and a Mitacs Accelerate Fellowship (with

partner organization Nature Conservancy of Canada) awarded to LN, and a NSERC Discovery

Grant awarded to EVW.

Data availability: Data and code to reproduce the analysis are available at

https://doi.org/10.5281/zenodo.6325510

References

1. Charnov EL. 1976 Optimal foraging, the marginal value theorem. *Theoretical Population Biology* **9**, 129–135.

2. Fretwell SD, Lucas HL Jr. 1969 On territorial behavior and other factors influencing habitat distribution in birds: I. Theoretical development. *Acta Biotheoretica* **19**, 16–36.

3. Pecoraro N, Reyes F, Gomez F, Bhargava A, Dallman MF. 2004 Chronic stress promotes palatable feeding, which reduces signs of stress: feedforward and feedback effects of chronic stress. *Endocrinology* **145**, 3754–3762. (doi:10.1210/en.2004-0305)

4. Dallman MF, Strack AM, Akana SF, Bradbury MJ, Hanson ES, Scribner KA, Smith M. 1993 Feast and famine: Critical role of glucocorticoids with insulin in daily energy flow. *Frontiers in Neuroendocrinology* **14**, 303–347. (doi:10.1006/frne.1993.1010)

5. Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE. 2008 A movement ecology paradigm for unifying organismal movement research. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 19052–19059. (doi:10.1073/pnas.0800375105)

6. Bonier F, Martin PR, Moore IT, Wingfield JC. 2009 Do baseline glucocorticoids predict fitness? *Trends in Ecology & Evolution* **24**, 634–642. (doi:10.1016/j.tree.2009.04.013)

7. Landys MM, Ramenofsky M, Guglielmo CG, Wingfield JC. 2004 The low-affinity glucocorticoid receptor regulates feeding and lipid breakdown in the migratory Gambel's white-crowned sparrow *Zonotrichia leucophrys gambelii*. *Journal of Experimental Biology* **207**, 143–154. (doi:10.1242/jeb.00734)

8.   Vitousek MN, Jenkins BR, Safran RJ. 2014 Stress and success: individual differences in the

glucocorticoid stress response predict behavior and reproductive success under high

predation risk. *Hormones and Behavior* **66**, 812–819. (doi:10.1016/j.yhbeh.2014.11.004)

9.   Boonstra R, Hik D, Singleton GR, Tinnikov A. 1998 The impact of predator-induced stress

on the snowshoe hare cycle. *Ecological Monographs* **68**, 371–394.

10. Vitousek MN, Taff CC, Ardia DR, Stedman JM, Zimmer C, Salzman TC, Winkler DW.

2018 The lingering impact of stress: brief acute glucocorticoid exposure has sustained, dose-

dependent effects on reproduction. *Proc. R. Soc. B.* **285**, 20180722.

(doi:10.1098/rspb.2018.0722)

11. Majer AD *et al.* 2019 Is there an oxidative cost of acute stress? Characterization, implication

of glucocorticoids and modulation by prior stress experience. *Proc. R. Soc. B.* **286**,

20191698. (doi:10.1098/rspb.2019.1698)

12. Boonstra R. 2013 Reality as the leading cause of stress: rethinking the impact of chronic

stress in nature. *Funct Ecol* **27**, 11–23. (doi:10.1111/1365-2435.12008)

13. Jeanniard du Dot T, Rosen DAS, Richmond JP, Kitaysky AS, Zinn SA, Trites AW. 2009

Changes in glucocorticoids, IGF-I and thyroid hormones as indicators of nutritional stress

and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*). *Comparative*

Biochemistry and Physiology Part A: Molecular & Integrative Physiology **152**, 524–534.

(doi:10.1016/j.cbpa.2008.12.010)

14. Krause JS, Pérez JH, Meddle SL, Wingfield JC. 2017 Effects of short-term fasting on stress

physiology, body condition, and locomotor activity in wintering male white-crowned

sparrows. *Physiology & Behavior* **177**, 282–290. (doi:10.1016/j.physbeh.2017.04.026)

15. Jaatinen K, Seltmann MW, Öst M. 2014 Context☐dependent stress responses and their

connections to fitness in a landscape of fear. *J Zool* **294**, 147–153. (doi:10.1111/jzo.12169)

16. Anderwald P, Campell Andri S, Palme R. 2021 Reflections of ecological differences? Stress

responses of sympatric Alpine chamois and red deer to weather, forage quality, and human

disturbance. *Ecology and Evolution* **11**, 15740–15753. (doi:10.1002/ece3.8235)

17. Landys☐Ciannelli MM, Ramenofsky M, Piersma T, Jukema J, Group CR, Wingfield JC.

2002 Baseline and stress☐induced plasma corticosterone during long☐distance migration in

the bar☐tailed godwit, *Limosa lapponica*. *Physiological and Biochemical Zoology* **75**, 101–

110. (doi:10.1086/338285)

18. Casas F, Benítez-López A, Tarjuelo R, Barja I, Viñuela J, García JT, Morales MB, Mougeot

F. 2016 Changes in behaviour and faecal glucocorticoid levels in response to increased

human activities during weekends in the pin-tailed sandgrouse. *Sci Nat* **103**, 91.

(doi:10.1007/s00114-016-1416-6)

19. Gormally BMG, Romero LM. 2020 What are you actually measuring? A review of

techniques that integrate the stress response on distinct time☐scales. *Funct Ecol* **34**, 2030–

2044. (doi:10.1111/1365-2435.13648)

20. Johnson DH. 1980 The comparison of usage and availability measurements for evaluating

resource preference. *Ecology* **61**, 65–71. (doi:10.2307/1937156)

21. Newediuk L, Vander Wal E. 2021 Predicting the individual identity of non-invasive faecal

and hair samples using biotelemetry clusters. *Mamm Biol* (doi:10.1007/s42991-021-00173-8)

22. Petrullo L, Delaney D, Boutin S, McAdam AG, Lane JE, Boonstra R, Palme R, Dantzer B.

2022 The glucocorticoid response to environmental change is not specific to agents of

natural selection in wild red squirrels. *Hormones and Behavior* **146**, 105262.

(doi:10.1016/j.yhbeh.2022.105262)

23. Goymann W. 2012 On the use of non-invasive hormone research in uncontrolled, natural

environments: the problem with sex, diet, metabolic rate and the individual: *Non-invasive*

hormone research. *Methods in Ecology and Evolution* **3**, 757–765. (doi:10.1111/j.2041-

210X.2012.00203.x)

24. Herman JP, Figueiredo H, Mueller NK, Ulrich-Lai Y, Ostrander MM, Choi DC, Cullinan

WE. 2003 Central mechanisms of stress integration: hierarchical circuitry controlling

hypothalamo–pituitary–adrenocortical responsiveness. *Frontiers in Neuroendocrinology* **24**,

151–180. (doi:10.1016/j.yfrne.2003.07.001)

25. Malisch JL, Breuner CW, Gomes FR, Chappell MA, Garland T. 2008 Circadian pattern of

total and free corticosterone concentrations, corticosteroid-binding globulin, and physical

activity in mice selectively bred for high voluntary wheel-running behavior. *General and*

Comparative Endocrinology **156**, 210–217. (doi:10.1016/j.ygcen.2008.01.020)

26. Rovirosa MJ, Levine S, Gordon MK, Caba M. 2005 Circadian rhythm of corticosterone

secretion in the neonatal rabbit. *Developmental Brain Research* **158**, 92–96.

(doi:10.1016/j.devbrainres.2005.06.007)

27. Smith JA, Donadio E, Pauli JN, Sheriff MJ, Middleton AD. 2019 Integrating temporal

refugia into landscapes of fear: prey exploit predator downtimes to forage in risky places.

Oecologia **189**, 883–890. (doi:10.1007/s00442-019-04381-5)

28. Landys MM, Ramenofsky M, Wingfield JC. 2006 Actions of glucocorticoids at a seasonal

baseline as compared to stress-related levels in the regulation of periodic life processes.

General and Comparative Endocrinology **148**, 132–149. (doi:10.1016/j.ygcen.2006.02.013)

29. Eikenaar C, Müller F, Leutgeb C, Hessler S, Lebus K, Taylor PD, Schmaljohann H. 2017

Corticosterone and timing of migratory departure in a songbird. *Proc. R. Soc. B.* **284**,

20162300. (doi:10.1098/rspb.2016.2300)

30. Falsone K, Jenni-Eiermann S, Jenni L. 2009 Corticosterone in migrating songbirds during

endurance flight. *Hormones and Behavior* **56**, 548–556. (doi:10.1016/j.yhbeh.2009.09.009)

31. Sapolsky RM, Romero LM, Munck AU. 2000 How do glucocorticoids influence stress

responses? integrating permissive, suppressive, stimulatory, and preparative actions. **21**, 35.

32. Carroll TB, Aron DC, Findling JW, Tyrrell JB. 2018 Chapter 9: Glucocorticoids and adrenal

androgens. In *Greenspan's Basic & Clinical Endocrinology* (eds DG Gardner, D Shoback),

p. 65. McGraw-Hill Education.

33. Swain DP, Benoît HP, Hammill MO. 2015 Spatial distribution of fishes in a Northwest

Atlantic ecosystem in relation to risk of predation by a marine mammal. *J Anim Ecol* **84**,

1286–1298. (doi:10.1111/1365-2656.12391)

34. McNamara JM, Houston AI. 1990 State-dependent ideal free distributions. *Evol Ecol* **4**, 298–

311. (doi:10.1007/BF02270929)

35. Foster MT, Warne JP, Ginsberg AB, Horneman HF, Pecoraro NC, Akana SF, Dallman MF.

2009 Palatable foods, stress, and energy stores sculpt corticotropin-releasing factor,

adrenocorticotropin, and corticosterone concentrations after restraint. *Endocrinology* **150**,

2325–2333. (doi:10.1210/en.2008-1426)

36. Hawlena D, Schmitz OJ. 2010 Physiological stress as a fundamental mechanism linking

predation to ecosystem functioning. *The American Naturalist* **176**, 537–556.

(doi:10.1086/656495)

37. Goldenberg SZ, Douglas-Hamilton I, Wittemyer G. 2018 Inter-generational change in

African elephant range use is associated with poaching risk, primary productivity and adult

mortality. *Proc. R. Soc. B.* **285**, 20180286. (doi:10.1098/rspb.2018.0286)

38. Burnard C, Ralph C, Hynd P, Hocking Edwards J, Tilbrook A. 2017 Hair cortisol and its

potential value as a physiological measure of stress response in human and non-human

animals. *Anim. Prod. Sci.* **57**, 401. (doi:10.1071/AN15622)

39. Palme R. 2019 Non-invasive measurement of glucocorticoids: advances and problems.

Physiology & Behavior **199**, 229–243. (doi:10.1016/j.physbeh.2018.11.021)

40. Luttbeg B, Grindstaff JL. 2022 Optimal hormonal regulation when stressor cues are

imperfect. *Hormones and Behavior* **144**, 105227. (doi:10.1016/j.yhbeh.2022.105227)

41. Dallman MF. 2005 Fast glucocorticoid actions on brain: Back to the future. *Frontiers in*

Neuroendocrinology **26**, 103–108. (doi:10.1016/j.yfrne.2005.08.001)

42. Breuner CW, Beyl HE, Malisch JL. 2020 Corticosteroid-binding globulins: lessons from

biomedical research. *Molecular and Cellular Endocrinology* **514**, 110857.

(doi:10.1016/j.mce.2020.110857)

43. Lattin CR, Romero LM. 2013 Seasonal variation in corticosterone receptor binding in brain,

hippocampus, and gonads in House Sparrows (*Passer domesticus*). *The Auk* **130**, 591–598.

(doi:10.1525/auk.2013.13043)

44. Lattin CR, Breuner CW, Michael Romero L. 2016 Does corticosterone regulate the onset of

breeding in free-living birds?: The CORT-Flexibility Hypothesis and six potential

mechanisms for priming corticosteroid function. *Hormones and Behavior* **78**, 107–120.

(doi:10.1016/j.yhbeh.2015.10.020)

45. Mastromonaco GF, Gunn K, McCurdy-Adams H, Edwards DB, Schulte-Hostedde AI. 2014

Validation and use of hair cortisol as a measure of chronic stress in eastern chipmunks

(*Tamias striatus*). *Conservation Physiology* **2**, cou055–cou055.

(doi:10.1093/conphys/cou055)

46. Spée M, Marchal L, Thierry A-M, Chastel O, Enstipp M, Maho YL, Beaulieu M, Raclot T.

2011 Exogenous corticosterone mimics a late fasting stage in captive Adélie penguins

(*Pygoscelis adeliae*). *American Journal of Physiology-Regulatory, Integrative and*

Comparative Physiology **300**, R1241–R1249. (doi:10.1152/ajpregu.00762.2010)

47. Breuner CW, Wingfield JC. 2000 Rapid behavioral response to corticosterone varies with

photoperiod and dose. *Hormones and Behavior* **37**, 23–30. (doi:10.1006/hbeh.1999.1554)

48. Amor JM, Newman R, Jensen WF, Rundquist BC, Walter WD, Boulanger JR. 2019 Seasonal

home ranges and habitat selection of three elk (*Cervus elaphus*) herds in North Dakota. *PLoS*

ONE **14**, e0211650. (doi:10.1371/journal.pone.0211650)

49. Cyr NE, Romero LM. 2009 Identifying hormonal habituation in field studies of stress.

General and Comparative Endocrinology **161**, 295–303. (doi:10.1016/j.ygcen.2009.02.001)

50. Baldan D, Negash M, Ouyang JQ. 2021 Are individuals consistent? Endocrine reaction

norms under different ecological challenges. *Journal of Experimental Biology* **224**,

jeb240499. (doi:10.1242/jeb.240499)

51. Dulude☐de Broin F, Hamel S, Mastromonaco GF, Côté SD. 2020 Predation risk and

mountain goat reproduction: evidence for stress☐induced breeding suppression in a wild

ungulate. *Funct Ecol* **34**, 1003–1014. (doi:10.1111/1365-2435.13514)

52. Giroux M-A, Dussault C, Lecomte N, Tremblay J-P, Côté SD. 2012 A new way of assessing

foraging behaviour at the individual level using faeces marking and satellite telemetry. *PLoS*

ONE **7**, e49719. (doi:10.1371/journal.pone.0049719)

53. Hertel AG, Niemelä PT, Dingemanse NJ, Mueller T. 2020 A guide for studying among-

individual behavioral variation from movement data in the wild. *Mov Ecol* **8**, 30.

(doi:10.1186/s40462-020-00216-8)

54. Avgar T, Potts JR, Lewis MA, Boyce MS. 2016 Integrated step selection analysis: bridging

the gap between resource selection and animal movement. *Methods Ecol Evol* **7**, 619–630.

(doi:10.1111/2041-210X.12528)

55. Cook RC *et al.* 2013 Regional and seasonal patterns of nutritional condition and

reproduction in elk. *Wild. Mon.* **184**, 1–45. (doi:10.1002/wmon.1008)

56. Lehman CP, Rota CT, Millspaugh JJ. 2019 Nutritional condition and physiological stress

levels of elk in the Black Hills, South Dakota. *PeerJ* **7**, e7185. (doi:10.7717/peerj.7185)

57. Duchamp JE, DeVivo MT, DeBerti JM, Heffernan LM, Kougher JD, Larkin JL. 2019

Selection of calf bedding sites by female elk in the Central Appalachian Mountains. *Journal*

of Fish and Wildlife Management **10**, 91–101. (doi:10.3996/102015-JFWM-097)

58. Alves J, Alves da Silva A, Soares AMVM, Fonseca C. 2013 Sexual segregation in red deer:

is social behaviour more important than habitat preferences? *Animal Behaviour* **85**, 501–509.

(doi:10.1016/j.anbehav.2012.12.018)

59. Ashley NT, Barboza PS, Macbeth BJ, Janz DM, Cattet MRL, Booth RK, Wasser SK. 2011

Glucocorticosteroid concentrations in feces and hair of captive caribou and reindeer

following adrenocorticotropic hormone challenge. *General and Comparative Endocrinology*

172, 382–391. (doi:10.1016/j.ygcen.2011.03.029)

60. Huber S, Palme R, Zenker W, Mostl E. 2003 Non-invasive monitoring of the adrenocortical

response in red deer. *The Journal of Wildlife Management* **67**, 258. (doi:10.2307/3802767)

61. Hinton JW, Freeman AE, St☐Louis V, Cornicelli L, D'Angelo GJ. 2020 Habitat selection by

female elk during Minnesota's agricultural season. *Jour. Wild. Mgmt.* **84**, 957–967.

(doi:10.1002/jwmg.21850)

62. Heidinger BJ, Nisbet ICT, Ketterson ED. 2006 Older parents are less responsive to a stressor

in a long-lived seabird: a mechanism for increased reproductive performance with age? *Proc.*

R. Soc. B. **273**, 2227–2231. (doi:10.1098/rspb.2006.3557)

63. Guindre-Parker S, Mcadam AG, van Kesteren F, Palme R, Boonstra R, Boutin S, Lane JE,

Dantzer B. 2019 Individual variation in phenotypic plasticity of the stress axis. *Biol. Lett.* **15**,

20190260. (doi:10.1098/rsbl.2019.0260)

64. Jenkins BR, Vitousek MN, Hubbard JK, Safran RJ. 2014 An experimental analysis of the

heritability of variation in glucocorticoid concentrations in a wild avian population. *Proc. R.*

Soc. B. **281**, 20141302. (doi:10.1098/rspb.2014.1302)

65. Patterson SH, Hahn TP, Cornelius JM, Breuner CW. 2014 Natural selection and

glucocorticoid physiology. *J. Evol. Biol.* **27**, 259–274. (doi:10.1111/jeb.12286)

66. Lema SC. 2020 Hormones, developmental plasticity, and adaptive evolution: endocrine

flexibility as a catalyst for 'plasticity-first' phenotypic divergence. *Molecular and Cellular*

Endocrinology **502**, 110678. (doi:10.1016/j.mce.2019.110678)

67. Tennessen JB, Parks SE, Swierk L, Reinert LK, Holden WM, Rollins-Smith LA, Walsh KA,

Langkilde T. 2018 Frogs adapt to physiologically costly anthropogenic noise. *Proc. R. Soc.*

B. **285**, 20182194. (doi:10.1098/rspb.2018.2194)

68. Northrup JM, Vander Wal E, Bonar M, Fieberg J, Laforge MP, Leclerc M, Prokopenko CM,

Gerber BD. 2022 Conceptual and methodological advances in habitat□selection modeling:

guidelines for ecology and evolution. *Ecological Applications* **32**. (doi:10.1002/eap.2470)

69. Boyce MS, McDonald LL. 1999 Relating populations to habitats using resource selection

functions. *Trends in Ecology & Evolution* **14**, 268–272. (doi:10.1016/S0169-5347(99)01593-

1)

70. Fieberg J, Signer J, Smith B, Avgar T. 2021 A 'How to' guide for interpreting parameters in

habitat□selection analyses. *J Anim Ecol* **90**, 1027–1043. (doi:10.1111/1365-2656.13441)

71. Fortin D, Morales JM, Boyce MS. 2005 Elk winter foraging at fine scale in Yellowstone

National Park. *Oecologia* **145**, 335–343.

72. MacLeod KJ, Sheriff MJ, Ensminger DC, Owen DAS, Langkilde T. 2018 Survival and

reproductive costs of repeated acute glucocorticoid elevations in a captive, wild animal.

General and Comparative Endocrinology **268**, 1–6. (doi:10.1016/j.ygcen.2018.07.006)

73. Laundré JW, Hernández L, Altendorf KB. 2001 Wolves, elk, and bison: reestablishing the

'landscape of fear' in Yellowstone National Park, U.S.A. *Can. J. Zool.* **79**, 1401–1409.

(doi:10.1139/z01-094)

74. Peers MJL *et al.* 2018 Quantifying fear effects on prey demography in nature. *Ecology* **99**,

1716–1723. (doi:10.1002/ecy.2381)

75. Stuber EF, Carlson BS, Jesmer BR. 2022 Spatial personalities: a meta-analysis of consistent

individual differences in spatial behavior. *Behavioral Ecology* **33**, 477–486.

(doi:10.1093/beheco/arab147)

76. Brass KE, Herndon N, Gardner SA, Grindstaff JL, Campbell P. 2020 Intergenerational

effects of paternal predator cue exposure on behavior, stress reactivity, and neural gene

expression. *Hormones and Behavior* **124**, 104806. (doi:10.1016/j.yhbeh.2020.104806)

77. Boudreau MR, Seguin JL, Boonstra R, Palme R, Boutin S, Krebs CJ, Murray DL. 2019

Experimental increase in predation risk causes a cascading stress response in free-ranging

snowshoe hares. *Oecologia* **191**, 311–323. (doi:10.1007/s00442-019-04500-2)

78. Leroux SJ *et al.* 2017 Stoichiometric distribution models: ecological stoichiometry at the

landscape extent. *Ecol Lett* **20**, 1495–1506. (doi:10.1111/ele.12859)

79. Lamb CT, Mowat G, McLellan BN, Nielsen SE, Boutin S. 2017 Forbidden fruit: human

settlement and abundant fruit create an ecological trap for an apex omnivore. *J Anim Ecol*

86, 55–65. (doi:10.1111/1365-2656.12589)

80. Gangoso L *et al.* 2021 Avian scavengers living in anthropized landscapes have shorter

telomeres and higher levels of glucocorticoid hormones. *Science of the Total Environment*

782, 146920. (doi:10.1016/j.scitotenv.2021.146920)

81. Lea JMD, Walker SL, Kerley GIH, Jackson J, Matevich SC, Shultz S. 2018 Non-invasive

physiological markers demonstrate link between habitat quality, adult sex ratio and poor

population growth rate in a vulnerable species, the Cape mountain zebra. *Functional Ecology*

32, 300–312. (doi:10.1111/1365-2435.13000)

82. Narayan EJ, Jessop TS, Hero JM. 2015 Invasive cane toad triggers chronic physiological

stress and decreased reproductive success in an island endemic. *Functional Ecology* **29**,

1435–1444. (doi:10.1111/1365-2435.12446)

Box 1 – Primer on habitat selection analysis and glucocorticoids

Habitat selection analysis (HSA) quantifies how stressors influence which habitats — or points in space characterized by a unique set of resources, risks and conditions [68] — are used preferentially or avoided by animals:

$$w_i = \frac{u_i}{a_i}$$

Where w_i is the "selection ratio" for habitat i, u_i is the proportion of habitat i used by the animal, and a_i is the proportional representation of habitat i within a landscape [69]. Note that use is often quantified using locations from biotelemetry devices. Unused locations are usually unknown, rendering proportion of use indeterminable. In these cases, selection for habitats can be estimated using logistic regression:

$$w_x = exp(\beta_0 + \beta_1 k_i + \cdots + \beta_n k_j)$$

Where the used locations are compared to a larger sample of "available" locations. Instead of probability of selection as in the selection ratio, coefficients β_1 to β_n represent selection for habitat variables, k, *relative* to j others in the model [70].

Later adjustments to the HSF model stemmed from technological advances in biotelemetry data collection. Modern biotelemetry devices collect used location data at regular time intervals. Intervals are often frequent enough that not all available locations are necessarily accessible, making biological constraints on movement a likely constraint on habitat availability. For example, locations visited within a day may not be accessible between hourly locations because the distance an animal can travel within an hour is constrained. Step selection functions

(SSF) address this constraint by drawing available locations from a set of trajectories to which

the animal could move from each location point (Fig. 1; [71]). Turn angle and step length

between location points become additional variables in the model, drawn from empirical

distributions of the animal's movements. Movement, however, is also influenced by stressors

habitat-specific stressors encountered along a movement path. Integrated step selection functions

(iSSF) extend the SSF approach by incorporating this interdependency between movement and

habitat selection within the modelling process [54].

Aside from its statistical justification, constraining availability and integrating movement

into the habitat selection process makes iSSF models convenient for testing hypotheses about

glucocorticoids and movement. Just as movement depends on habitats encountered along a

movement path, it also depends on the animal's glucocorticoid production. Glucocorticoid

samples from known time points along the movement path can be integrated into the iSSF model

as interactions between habitat and movement covariates (Fig. 1). Interaction coefficients test

hypotheses about the effects of glucocorticoids on both capacity and motivation for movement.

For example:

Step length and turn angle interaction with glucocorticoids at the start of a step: When

animals produce more glucocorticoids, are their movements faster and more directed (Fig. 1 a)?

Interaction between glucocorticoids at the start of a step and habitat at the end of the step:

Do animals that produce more glucocorticoids select safer habitat or habitats with resources to

replenish energy reserves (Fig. 1 b)?

Box 2 – Future directions

Integrating glucocorticoid hormones and habitat selection can both test the assumptions of some foundational hypotheses in stress physiology and space use ecology and generate new testable hypotheses that integrate both.

Glucocorticoid production can cause oxidative stress and reduce reproductive success (e.g., [72]), but elevated baseline levels often increase lifetime reproductive success if they facilitate immediate investment in current offspring (*cort-adaptation hypothesis*; [6]). Some investment behaviours, such as foraging trips by nesting birds, are based in habitat selection. By integrating glucocorticoids and habitat selection with iSSF models, we can detect proximate effects of glucocorticoids on investment behaviours. Are these investment behaviours repeatable within individuals? If so, how much does investment strategy contribute to individual differences in lifetime reproductive success?

The *landscape of fear* is a popular space use ecology concept linking habitat selection to predation risk: animals facing predation are expected to avoid unsafe habitats or use antipredator strategies like vigilance or safety in numbers to mitigate risk [73]. However, predator cues rather than presence are often used in practice as a proxy for risk, making fear itself difficult to quantify [74]. Can we use glucocorticoids to quantify the landscape of fear, and iSSF models to measure its responsibility for antipredator habitat selection strategies?

Just as glucocorticoids drive adaptive habitat selection behaviour, the habitats individual animals select should also affect future changes in their hormone levels. This two-way relationship between habitat and hormones may drive a positive feedback loop that reinforces habitat

selection behaviour. Evidence is emerging that habitat selection behaviour is indeed repeatable

[75] and that adults continuously exposed to stressors can epigenetically alter offspring hormonal

responses to the same stressors [76]. Do intergenerational glucocorticoid-habitat selection

feedback loops underlie specialization of populations on habitats?

Looking beyond populations, an integrated approach to studying glucocorticoids and habitat

selection may also provide insights into ecosystem functioning and the ability of species to

respond to environmental change.

The *carbon stress hypothesis* [36] purports animals facing predation stress have higher cellular

respiration rates, making them carbon limited. To maintain C:N ratios, stressed animals should

seek forage with more of digestible carbon. Animals also select foraging habitat when they

produce more glucocorticoids under stress physiology theory [77]. Can we use iSSF and spatial

data of elemental landscape composition [78] to reconcile expectations of stress physiology with

the carbon-stress hypothesis? Preferences for carbon-rich forage and resulting spatial variation in

carbon-rich additions to detrital pools could fundamentally alter ecosystem structure [36].

Even if glucocorticoids drive adaptive habitat selection, animals unable to access habitats that

counteract stressors may instead behave in a manner that compromises fitness. Fitness costs may

be particularly high in human-modified landscapes. For example, anthropogenic structures

present physical barriers to movement and limit assessment of habitat selection costs (i.e.,

ecological traps; e.g., [79]). Can we use the null expectation that both habitat selection and

glucocorticoid production are adaptive to identify cases where animals are not free to move?

Glucocorticoids in these cases may measure need for management intervention.

Table 1. Examples of recent field studies reporting negative relationships among habitat-

associated stressors, glucocorticoid production, and fitness, without testing for causal

relationships among the three. We searched for articles on Web of Science in June 2021 using

the search query *habitat* AND *glucocorticoid** OR *corticosterone* OR *cortisol* AND *fitness* AND

chronic stress.

Topic	Findings	Interpretation	Citation
Effect of human footprint on feather glucocorticoids and survival rate of Eurasian griffon vultures (*Gyps fulvus*)	Glucocorticoids elevated among birds living in areas with higher human footprint. Previous studies found lower survival rates in areas with greater footprint.	Chronic stress is associated with living in areas with higher human footprint	[80]
Effect of habitat quality on fecal glucocorticoids and population growth rate in Cape mountain zebra (*Equus zebra zebra*)	Glucocorticoids elevated in lower-quality habitat. Glucocorticoids also highest among populations with lower female fecundity and growth rates.	Populations in lower-quality habitat suffering from chronic stress due to poorer nutrition	[81]
Effect of invasive species on urinary glucocorticoids on reproduction and survival in Fijian ground frog (*Platymantis vitiana*)	Glucocorticoids elevated in enclosures with invader. Body condition and reproduction, but not survival, also lower with invader.	Competition with invader causes chronic stress that suppresses reproduction.	[82]

Figure 1. Schematic representation of used locations (circles) along a movement path from

$t = 0$ through $t = 2$. Solid arrows represent the movement path between location points, and

dashed arrows represent five available trajectories an animal could have taken at $t = 1$, drawn

from assumed distributions of the animal's step lengths (SL) and turn angles (TA) between

movement paths with parameters based on its movement characteristics. Effects of

glucocorticoids (CORT) at $t = 1$ on movement between $t = 1$ and $t = 2$ can be tested using its

interaction with TA (between $t = 0$ and $t = 1$) and SL (between $t = 1$ and $t = 2$) as variables

in an iSSF model (a). Its interaction with habitat at $t = 2$ tests how glucocorticoids at $t = 1$

affect habitat selection.

Figure 2. Log relative selection strength (RSS) for a location within cover (dashed lines) versus

450 m from cover (solid lines) with increasing fecal glucocorticoid metabolite (FGM) levels.

Blue ribbons show the change in RSS between these locations in the 16-d window ("pre"

period) when young calves are most vulnerable, and orange ribbons outside of this period

("post" period). Distance between the solid and dashed lines indicate relative selection strength

is stronger for locations closer to cover as FGM levels increase. Higher elevation of the orange

ribbon indicates a stronger effect during the 16-d window of calf vulnerability, suggesting

lower corticosterone binding globulin concentration, fewer receptors, or reduced negative

feedback may be driving selection for foraging habitat when energy needs are greatest.

a) CORT: TA + CORT: SL

b) CORT: habitat