A domesticated fungal cultivar recycles its cytoplasmic contents as nutritional rewards for its leafcutter ant farmers

Caio Ambrosio Leal-Dutra[1*], Lok Man Yuen[2], Pedro Elias Marques,[3] Bruno Augusto Maciel Guedes[4], Marta Contreras-Serrano[1], Jonathan Zvi Shik[1,5]

[1] Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark

[2] Department of Life Sciences, Imperial College London, London SW7 2AZ, UK

[3] Laboratory of Molecular Immunology, Department of Microbiology, Immunology and Transplantation, Rega Institute, KU Leuven, Leuven, Belgium

[4] Departamento de Ciências Básicas da Vida - Universidade Federal de Juiz de Fora, Campus Governador Valadares, MG, 35020-360, Brasil

[5] Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Republic of Panama

* caio@bio.ku.dk

Authors email in order:

lok.yuen19@imperial.ac.uk

pedro.marques@kuleuven.be

bruno.guedes@ufjf.edu.br

contrerasserranomarta@gmail.com

jonathan.shik@bio.ku.dk

Competing Interests

The authors declare no competing interest.

Keywords: autophagy, *Leucoagaricus gongylophorus*, leafcutter ant, gongylidia, fungus, attini, attines

ABSTRACT

Leafcutter ants farm a fungal cultivar (*Leucoagaricus gongylophorus*) that converts inedible

vegetation into food that sustains colonies with up to millions of workers. Analogous to edible

fruits of crops domesticated by humans, *L. gongylophorus* has evolved specialized nutritional

rewards—swollen hyphal cells called gongylidia that package metabolites ingested by ant

farmers. Yet, little is known about how gongylidia form, and thus how fungal physiology and ant

provisioning interact to farming performance. We explored the mechanisms governing

gongylidium formation using microscopy imaging of ant-cultivated fungus and controlled *in vitro*

experiments with the cultivar grown in isolation from ant farmers. First, *L. gongylophorus* is

polykaryotic (up to 17 haploid nuclei/cell) and our results suggest intracellular nucleus

distributions govern gongylidium morphology with their absence in expanding edges arresting

apical growth and their presence mediating complex branching patterns. Second, nanoscale

imaging (SEM, TEM) shows that the cultivar recycles its own cellular material (*e.g.* cytosol,

mitochondria) through a process called 'autophagy' and stores the resulting metabolites in

gongylidia. This autophagic pathway is further supported by gongylidium suppression when

isolated fungal cultures are grown on media with autophagy inhibitors, and differential transcript

expression (RNA-seq) analyses showing upregulation of multiple autophagy genes in

gongylidia. We hypothesize that autophagic nutritional reward production is *the* ultimate cultivar

service and reflects a higher-level organismality adaptation enabled by strict symmetric lifetime

commitment between ant farmers and their fungal crop.

Introduction

The advent of domesticated agriculture some 10 000 years ago was a turning point for humans and for the domesticated crops whose derived traits would likely have been maladaptive in their free-living ancestors [1-3]. Key crop adaptations include whole genome duplication events (resulting in polyploidy) that can increase functional heterozygosity [4] and selection for specific regulatory genes that can reduce seed shattering or enhance fruit size, color, and sweetness [5-8]. Fascinatingly, humans are not the only farmers. Several insect lineages independently evolved obligate farming systems of fungal cultivars that produce specialized edible reward structures [9]. However, while human farmers modify growth environments in well-known ways to maximize crop yield (e.g. adding fertilizers, controlling watering conditions, etc.), the analogous mechanisms by which insect farmers promote expression of edible reward structures in fungal cultivars remain poorly understood.

The largest-scale insect farmers are the *Atta* leafcutter ants, the crown group of the fungus-farming 'attine' ant lineage [10, 11]. Despite clear analogies with farming systems of humans [9], farming by leafcutter ants is fundamentally different because it is 'organismal' in the sense that it represents a strictly symmetrical obligate mutualistic dependence [12]. Such an arrangement usually does not allow alternative crops, but does sustain selection for co-evolutionary integration and higher-level adaptation that cannot evolve when farming practices are asymmetrically promiscuous [13, 14]. These differences make it interesting to explore how leafcutter farmers regulate crop productivity, since they can help to: 1) understand the broad eco-evolutionary success of these naturally selected farming systems and 2) provide nutritional insights into whether the leafcutter ectosymbiosis has achieved an organismal level of conflict-free trait evolution typically only seen in lifetime committed endosymbioses (*e.g.* the mitochondria-nucleus partnership in eukaryotic cells [15, 16]).

A mature rainforest colony of the leafcutter ant *Atta colombica* can have millions of specialized ants that divide the work of foraging fresh plant fragments and caring for the fungal cultivar [17]. In this way, colonies convert foraged fragments from hundreds of plant species [18] into a mulch used to provision their fungal cultivar *Leucoagaricus gongylophorus*. In return, the cultivar converts inedible plant biomass into edible reward structures called gongylidia, that are swollen hyphal cells *ca.* 30 μm in diameter and grow in bundles called staphylae [19-27].

Gongylidia are a defining trait of irreversible crop domestication and are unique to the fungal

lineage farmed by leafcutter ants and other higher-neoattine genera including *Trachymyrmex*,

Sericomyrmex, *Mycetomoellerius*, and *Paratrachymyrmex* [20, 28, 29].

Gongylidia mediate functional integration with their ant symbionts in two main ways.

First, they contain enzymes (*e.g.*, laccases, pectinases, proteases) that ants ingest and then

vector to patches of newly deposited vegetation to catalyze fungus-mediated digestion and

detoxification [30-34]. Second, they contain nutrients (*e.g.*, amino acids, lipids and glycogen)

that are the ants' primary food source [35, 36]. The ability to regulate the quantity and quality of

gongylidia would thus provide clear benefits for the ant farmers. However, the mechanisms

linking substrate provisioning by farming ants and the production of the cultivar's edible yield

have remained poorly known. To better understand these mechanisms, we: 1) visualized the

morphology of gongylidia and staphylae using scanning electron microscopy (SEM), 2)

described the cellular reorganizations that mediate gongylidium formation by combining light,

fluorescence, confocal and transmission electron microscopy (TEM), and 3) used a

transcriptomics experiment to compare gene expression in hyphae and gongylidia to help

resolve the metabolic pathways underlying gongylidia production.

We next examined the cellular origins of the edible resources contained in the large

vacuole that fills each gongylidium cell. Previous evidence suggests that *L. gongylophorus*

directly metabolizes provisioned plant fragments to produce these edible resources. First, the

cultivar can metabolize lipids rich in alpha-linoleic acid (18:3) from foraged plant fragments into

linolenic acid (18:2) that is enriched in gongylidia [37]. This synthesized metabolite is thought to

mediate interkingdom communication by eliciting attractive behaviors in ant workers, in contrast

to the precursor 18:3 lipid that elicits antagonistic behaviors [37]. Second, isotopic enrichment

studies have shown that the cultivar quickly (within two days) shunts C and N from provisioned

substrates (glucose and ammonium nitrate, respectively) into edible gongylidia [38]. Third,

different substrate types are associated with increased expression of genes regulating targeted

pathways for nutritional metabolism [21, 36, 39, 40]. However, it is also reasonable to predict

that the diversity of compounds found within gongylidia have a diversity of biochemical origins.

Autophagy is a plausible alternative and/or complementary pathway underlying

gongylidia formation that involves the recycling of the cultivar's own metabolic source material

and potentially the fine-tuning of its composition. The metabolic pathways for autophagy are

conserved across eukaryotes and are known to mediate development, cellular differentiation

[41-44], and housekeeping [45, 46] in fungal cells. During autophagy, the cultivar's own

cytoplasmic components (*i.e.*, glycogen, proteins, organelles) are incorporated into a vacuole

for enzymatic degradation and the resulting catabolites are then recycled as nutrients to sustain

other cellular processes and produce new cellular components [47, 48]. Initial evidence for

autophagy in *L. gongylophorus* was first obtained in 1979 by Angeli-Papa and Eymé [49] who

used TEM imaging to observe endoplasmic reticulum membranes engulfing mitochondria during

gongylidium formation. However, to our knowledge, this preliminary evidence for autophagic

recycling of the cultivar's own intracellular content during gongylidium formation has not been

subsequently explored.

We propose that confirmation of an autophagic pathway(s) would have important

implications for understanding the leafcutter symbiosis since it implies that natural selection has

targeted the farming symbiosis in ways that made provisioning more robust and less dependent

on the variable quality and quantity of foraged vegetation. Specifically, we predict that

autophagic nutrient recycling of cellular contents would: 1) reduce variability in the quality of the

cultivar's nutritional rewards and provide opportunities to optimize the composition of metabolic

source material, 2) constrain the ability of ants to directly regulate cultivar productivity through

their provisioning decisions, and 3) function in a complementary manner to provisioned plant

substrates by providing a stable supply of metabolic precursor substrates during periods of

environmental vegetation shortage.

We tested for autophagic gongylidium formation in three ways. First, autophagy

encompasses two main types of cellular recycling mechanisms: 1) macroautophagy in which

cytoplasmic content (i.e. cytosolic metabolites and organelles) are sequestered into double-

membraned vesicles that fuse with vacuoles, and 2) microautophagy in which the vacuolar

membrane invaginates and directly engulfs cytoplasmic cargo [47, 48]. We determined whether

and how these autophagic processes influence gongylidium formation by examining organelle

rearrangements in TEM images and tracking experimentally supplied fluorescent-labeled

nutrients using confocal microscopy. Second, we tested whether autophagy is necessary for

gongylidium formation by performing an *in vitro* experiment where the density of staphyla was

measured in cultivars grown with known inhibitors and promoters of autophagy in fungal cells.

Third, we tested whether autophagic pathways are differentially expressed in developing

gongylidium cells by performing transcriptomic analyses of the mycelia and differentiated

staphylae of the cultivar when grown under controlled *in vitro* conditions on a standardized

medium without ant farmers.

Results

Nutritional reward structures

Each gongylidium cell consists of two sections that we term the bulb (swollen section)

and the filament (elongated section) (Fig. 1 A). Gongylidia are often connected by intercalary

bulbs (between filaments) and intercalary filaments (between bulbs) (Fig. 1 B-C) with

multifaceted branching patterns and individual gongylidium cells bearing two or more terminal

bulbs (Fig. 1 D). Gongylidia bulbs have variable diameters ranging from 12 μm to 50 μm and

variable filament lengths ranging from 40 μm to > 250 μm. We hypothesize that variable bulb

sizes reflect indeterminate growth trajectories of expanding gongylidium cells. All gongylidium

cells also contained at least eight nuclei usually concentrated at the intersection of the bulb and

the filament (Fig. 1 E) and usually one single vacuole (Fig. 1 F) that comprised up to half of

each bulb's total volume.

Individual staphylae range widely in size, with tens to hundreds of individual

gongylidium cells (Fig. 1 G, Fig. S1), and always formed on the surface of fungus garden

mycelial matrix where ants can easily detach them from surrounding hyphae. Staphylae also

form in the absence of ants under *in vitro* (Petri dish) growth conditions, but they have the

following key morphological differences compared to those growing in ant-tended fungus

gardens, being: 1) less detachable because they are usually covered by filamentous hyphae, 2)

larger in area and with more individual gongylidium cells, and 3) comprised of several 'burst'

gongylidium cells. We thus propose that under farming conditions, ants harvest staphylae

earlier in their development before vacuoles can produce turgor pressure exceeding the

retaining capacity of their exceptionally thin (ca. 120 to 220 nm) cell walls (Fig. 1 H).

An autophagic mechanism of gongylidia formation

Microscopy images (TEM, light, fluorescence and confocal) of gongylidium cells revealed structures that are diagnostic of macroautophagic processes. First, gongylidia were enriched with long stretches of endoplasmic reticulum that produce double-membraned vesicles called autophagosomes (Fig. 2), within which recycling of cellular materials is initiated. We confirm that autophagosomes contained cytosol (Fig. 2 A), glycogen (Fig. 2 B) and mitochondria (Fig. 2 C), and predict that other metabolites (*e.g.*, lipids, amino acids, enzymes [28, 30, 37, 40, 64]) are likely abundant, but are too small to be detected with TEM. Second, large numbers of damaged mitochondria were present in gongylidia and were often associated with endoplasmic reticulum membranes (Fig. 2 C) where they were likely destined to be sequestered into autophagosomes, digested, and recycled into edible metabolites. Third, vacuoles within gongylidium bulbs often contained single-membraned autophagic bodies (Fig. 2 D-F). These vesicles indicate the delivery of metabolites into vacuoles, since they are autophagosomes that lost their outer membrane after vacuolar fusion. This was confirmed by confocal images showing that autophagic bodies within gongylidium vacuoles contained fluorescently labeled sugars from the cultivar's cytosol (Fig. 2 E-F, Video S1). Given these hallmarks of macroautophagy (and the lack of evidence for microautophagy), we henceforth use 'autophagy' to refer to macroautophagy.

An autophagic mechanism for gongylidium formation was further supported by significant *in vitro* treatment effects of chemical autophagy inhibitors on staphyla density (Kruskal-Wallis: $H_3 = 34.7$, $p < 0.001$, Fig. 3). Pairwise post-hoc comparisons indicated that both autophagy inhibitors (3-MA and CQ) had significantly reduced staphyla density compared to both the control (PDA) ($p_{adj} < 0.001$ and $p_{adj} = 0.001$, respectively) and the autophagy-promoter (RAP) treatment ($p_{adj} < 0.001$ and $p_{adj} = 0.026$, respectively). There was also a significant treatment effect on mycelial growth area ($H_3 = 16.3$, $p = 0.001$). However, this was due to differences between 3-MA and all other treatments (PDA:3-MA, $p_{adj} = 0.004$; RAP:3-MA, $p_{adj} = 0.015$; CQ:3-MA, $p_{adj} = 0.001$), with no other significant pairwise comparisons (Fig. S2). Thus, while both autophagy inhibition treatments resulted in staphyla reduction, it is possible that 3-MA negatively influenced staphyla density through unknown indirect effects on cultivar

performance. The autophagy promotor (RAP) did not significantly increase staphyla density

relative to control (PDA) or either of the autophagy inhibition treatments (Fig. 3).

Transcriptome assembly and autophagy pathway analysis

Using a *de-novo* assembly and a clustering analysis by similarity, we recovered 78,820

transcripts from the *L. gongylophorus* transcriptome of which, 4,755 were differentially

expressed (log2 fold-change > 1.0, Benjamini–Hochberg adjusted $P < 0.001$) in staphylae (n =

3,011 transcripts) or in non-differentiated mycelia (n = 1,744 transcripts). Of these differentially

expressed transcripts (DETs), 31 were assigned a KEGG orthology term associated with the

yeast autophagy pathway (n = 22 in staphylae, n = 9 in mycelia, Table 1). Several key genes

were upregulated in staphylae that are typically over-transcribed during autophagy [48]. These

include genes (ATG7, ATG8, ATG10) linked to the recruitment of cargo into incipient unclosed

membranes (*i.e*., phagophores) and mature autophagosomes, as well as genes related to

starvation signaling (Sch9, Tap42, ATG13 and RIM15), vacuole fusion machinery (Ypt7, Mon1

and Vps3), and degradation of autophagic bodies (PRB1 and ATG15) (Table 1, Fig. S3). In

contrast, the few autophagy-specific genes upregulated in undifferentiated mycelia appear

related to the starvation signaling step of autophagy induction, rather than a fully functioning

autophagy pathway.

Discussion

Our results provide novel insights into the mechanisms of higher-level homeostatic

integration in a uniquely 'organismal' insect-microbe ectosymbiosis. Several lines of evidence

indicate that the fungal cultivar of leafcutter ants uses autophagic recycling to convert its own

cellular material into edible metabolites within specialized nutritional reward structures. First,

nanoscale imaging shows the cellular hallmarks of autophagy (*e.g.* autophagosomes,

autophagic bodies, abundant endoplasmic reticula) and indicates rapid delivery of labeled

cytosol nutrients into gongylidium vacuoles (ca. 30 minutes). Second, experimental suppression

of autophagy suppresses gongylidium density. Third, we find an upregulated autophagy

pathway associated with differentiated gongylidium cells. We hypothesize that this autophagic

recycling pathway represents a final domestication step where the cultivar came to
unambiguously prioritize nutritional services to its hosts even at the expense (up to a point) of
its own mycelial health. In this sense, the autophagic recycling pathway expresses an obligately
symmetric commitment between symbionts achieved after the domesticated fungal cultivar lost
the capacity for a free-living existence and became fully integrated into the host colony's germ
line.

We further propose that autophagic recycling facilitates homeostasis at higher levels of
organization by stabilizing the quantity and nutritional quality of the cultivar's nutritional rewards
in fluctuating environments. First, the seasonal and spatial availability of preferred plant
fragments may vary in suboptimal ways [65, 66]. Second, foraged plant fragments can contain
key nutrients—but these nutrients can occur in suboptimal ratios and concentrations relative to
the cultivar's intrinsic needs and tolerances [19]. Third, plant fragments contain a wealth of
recalcitrant compounds (e.g. cellulose and lignin) and toxic metabolites that can reduce cultivar
performance [67-70]. During such periods of plant-fragment shortage [71-73], the cultivar may
use autophagic recycling of its own organelles to yield reliably available and chemically
predictable metabolic precursor compounds. Analogous adaptations aimed solely regulating
homeostasis at higher levels of organization are absent from all other ectosymbioses that are
promiscuous by comparison, where the interests of symbionts are not completely aligned, and
partners must screen, sanction, and police to dissuade cheating [74-76].

The discovery of autophagic recycling also provides a new lens to interpret well-known
gardening behaviors in leafcutter ants. For instance, gardening ants constantly prune the
cultivar's fungal mycelia and this has been hypothesized to cause mechanical disruptions that
stimulate gongylidium formation [77]. We propose that such pruning behaviors sever hyphal
connections and block the flow of nutrients to newly isolated fungal cells, which in turn induces
gongylidium formation by causing an autophagic response to starvation—which is a common
driver of autophagy in cells [50]. Additionally, previous *in vitro* studies have observed highest
staphylae densities at the lowest nutrient concentrations suggesting a link between staphyla
formation and nutrient depletion [19, 40, 78-80]. Thus, while much research assumes that
cultivar production hinges on a maximized flow of provisioned plant fragments [19, 53, 81], the

behaviors linked to targeted nutritional suppression may also be important for the production of nutritional rewards.

While gongylidia-linked autophagic recycling appears common, it likely complements other nutritional reward production mechanisms. First, leafcutter ants frequently ingest gongylidia contents (crushing the gongylidia to ingest their cellular content) and vector the cultivar's enzymes [31, 32, 34, 35] and nutrients [35, 36] as fecal droplets to catalyze degradation and detoxification of newly deposited plant fragments [30]. Metabolites within these fecal droplets are assimilated by the cultivar and some subset likely enters biosynthetic pathways linked to gongylidium formation. Second, key nutrients also appear to derive from bacterial symbionts (rather than plant fragments or autophagic recycling) [82-84]. For instance, attine ants have lost the ability to synthesize arginine [80]—and depend on the cultivar's metabolism to produce this nitrogen-rich amino acid [28]. In turn, the ants have evolved tight mutualistic associations with specialized bacterial symbionts that convert excess arginine in the ants' guts into ammonia (the Mollicutes EntAcro1, [84]) that is an N-rich fertilizer vectored by the ants back to their fungal symbiont. As further evidence that the autophagic-recycling is one of several mechanisms by which gongylidia fill with metabolites, we observed that staphyla production was still possible (even though significantly reduced) when autophagy was inhibited in the *in vitro* experiment. Resolving whether and how these nutritional pathways fluctuate relative to the specific resource needs of the colony thus represents an important next step.

Reconstructing the cellular reorganizations enabling gongylidium formation

The combined evidence we present throughout this study enables us to propose a complete pathway for gongylidia development (Fig. 4). Tissue differentiation initiates when a hyphal cell growing at surface interstices within the fungus garden matrix widens (Fig. 4 A) and continues as a vacuole begins expanding in response to autophagic recycling. This process accelerates autophagosomes fuse with the vacuole and delivers cytoplasmic cargo (Fig. 4 B). The location of this vacuole shapes the morphology of developing gongylidia because it mediates the spatial distributions of the up to 17 [86] haploid nuclei within individual gongylidium cells by excluding them from the apical tip (Fig. 4 C).

We hypothesize that the obstruction of nuclear migration causes the bulb's balloon-like

expansion by blocking communication between nuclei and the Spitzenkörper—the centralized

machinery for hyphal growth located in the hyphae tip. At the transcript level, a structurally

modified and upregulated transcript carrying a domain associated with microtubule related

proteins [40] may mediate such nuclear migration in association with motor protein complexes

[87-89]. This mechanism would resemble growth dynamics in most filamentous fungi where

nuclei are distributed evenly throughout the hyphal compartment and promote tip elongation by

migrating apically [88, 90]. Further supporting this hypothesis, when nuclei are occasionally

aggregated in different regions of the filament and bulb, they are associated with ramified

branching gongylidia and intercalary bulb formation. Finally, we propose that staphylae arise

from this patchy ramification of tangled gongylidia (Fig. 4 D).

These results can also inform our understand of the functional consequences of

polyploidy in the domesticated *L. gongylophorus* cultivar which contains up to 7 distinct

haplotypes per cell [86]. Specifically, key next steps involve moving beyond distributions of

nuclei in gongylidium cells, to testing whether factors like nucleus-specific expression and

nuclear dominance are linked to gongylidium formation. Such a mechanism has recently been

observed in the production of edible reward structures produced by the heterokaryon human-

domesticated champignon fungus (*Agaricus bisporus*), where two distinct nuclear types exhibit

differential expression in distinct tissues during mushroom formation [91]. The convergent

existence of such molecular mechanisms in gongylidium formation would provide a further

means of testing the hypothesis that these nutritional rewards are derived from cystidia [20],

which are modified hypha found in the hymenium of several groups of basidiomycetes [92, 93].

Autophagic induction of cystidia may have provided crucial pre-adaptations harnessed by

natural selection to generate the unique gongylidium reward structures.

Methods

Sample acquisition and fungal symbiont isolation

Two colonies of *Atta colombica* (Ac2012-1 and Ac2019-1) were used in this study that

were collected in Soberanía National Park, Panama and thereafter maintained at the University

of Copenhagen in a climate-controlled room (25°C, 70% RH, minimal daylight). For microcopy

and imaging, we used staphylae collected directly from the colonies' fungus gardens, and for

the autophagy inhibition experiments and transcriptome sequencing, we used axenic cultures

isolated from the fungal cultivar (*L. gongylophorus*) grown in 90 mm Petri dishes filled with 20 ml

potato dextrose agar (PDA) that were kept in the dark at 25°C.

Imaging morphology of staphyla and gongylidia

Using samples collected directly from the fungus gardens of live leafcutter colonies, we

first used scanning electron microscopy (SEM) to visualize the external morphology of

gongylidia and staphylae. The samples were fixed in PBS with 0.1% Tween 20 and fixatives

(4% glutaraldehyde, 4% formaldehyde) and then dehydrated in an ethanol series (35%, 55%,

75%, 85%, 95%, and 2x in 100%) for 30 minutes per concentration, critical-point dried, coated

with platinum, and imaged on a JSM-840 scanning electron microscope (JEOL, Tokyo, Japan)

at 7.0 kV at the Zoological Museum of the University of Copenhagen. A slight wrinkled

appearance of the surface of gongylidium cells in the resulting SEM images was due to

unavoidable plasmolysis caused by the preparation process.

We then used light, fluorescence and confocal microscopy to view the cultivar's internal

morphology (*e.g.* septa, vacuoles, nuclei, etc.) and examine cellular reorganizations associated

with gongylidium induction. For all imaging, staphylae were first placed in a drop of mounting

solution (dH$_2$O, PBS, 3% KOH) on a glass slide. Gongylidia were then separated under a stereo

microscope (16x or 25x magnification) with 0.16-mm diameter acupuncture needles and

stained. For visualization with white light, we placed samples in either 0.1% Congo-red (in 150

mM NaCl) for one minute followed by a wash with 150 mM NaCl, or 1.5% phloxine followed by a

wash with 3% KOH. For nucleus visualization under UV light, we stained samples for 10

minutes using "Vectashield with DAPI" (Vector Laboratories, Burlingame, CA, USA). For

confocal imaging, we stained the staphylae with dextran conjugated with Alexa Fluor 647

(Invitrogen, MA, USA) in the concentration of 100 μg/ml in PBS for 30 minutes and washed

twice in PBS. We then acquired images at magnifications of up to 400x by performing bright-

field, dark-field, phase-contrast, and fluorescence microscopy under an Olympus BX63

microscope (Olympus, Tokyo, Japan). The microstructures were measured and photographed

using a mounted QImaging Retiga 6000 monochrome camera and cellSens Dimension v1.18

(Olympus) image-processing software. Confocal images were acquired with a Dragonfly

spinning-disk confocal system (Andor Technology, Belfast, Northern Ireland) equipped with a

25x water-immersion objective. Samples were excited with a 637 nm laser line and

fluorescence was collected with a 698/77 emission filter. Images were processed using FIJI

software.

We next used transmission electron microscopy (TEM) to visualize fungal cells with

greater magnification and resolution (i.e. the 500 nm scale). Staphylae were collected from

fragments of intact fungus gardens, fixed in 2% glutaraldehyde in 0.05 M PBS (pH 7.2) and then

post-fixed in 1% w/v OsO_4 with 0.05M $K_3Fe(Cn)_6$ in 0.12 M sodium phosphate buffer (pH 7.2) for

2 hr at room temperature. Fixed samples were washed three times in ddH_2O for 10 minutes and

dehydrated in a series of increasing ethanol concentrations series (70%, 96% and 99.9%). Each

dehydration lasted 15 min and was performed twice per concentration. Samples were then

repeatedly infiltrated for 20 to 40 min with increasing Resin Epon:Propylene oxide ratios (1:3,

1:1, 3:1) and subsequently embedded in 100% Epon and polymerized overnight at 60°C.

Sections of 60 nm thickness were then cut with an Ultracut 7 ultramicrotome (Leica, Vienna,

Austria), collected on copper grids with Formvar supporting membranes, and stained with both

uranyl acetate and lead citrate. These samples were TEM imaged on a CM100 BioTWIN

(Philips, Eindhoven, The Netherlands) at an accelerating voltage of 80 kV. Digital images were

recorded with a side-mounted OSIS Veleta digital slow scan 2 × 2 k CCD camera and the ITEM

software package (Olympus Soft Imaging Corp, Münster, Germany). This sample preparation

and imaging was performed at the Core Facility for Integrated Microscopy at the University of

Copenhagen.

Autophagy inhibition assay

We tested the role of autophagy in gongylidium production using an *in vitro* growth

assay with four treatment groups. Autophagy is often initiated in cells when the target of

rapamycin kinase (TOR) is inhibited. Autophagy can thus be induced *in vitro* by adding

rapamycin (RAP) [50] an allosteric TOR inhibitor [48]. Autophagy is often inhibited *in vitro* using

Chloroquine (CQ) or 3-methyladenine (3-MA), as these compounds respectively block

autophagosome-vacuole fusion [51] and suppress an enzyme (class III PtdIns3K) required to

initiate autophagosome formation [52]. We compared cultivars grown in the dark for 46 days at

25°C on a baseline Potato Dextrose Agar (PDA) medium containing nutrients known to

maximize cultivar performance [19, 53] (n = 40) with cultivars grown on plates with RAP (n =

27), CQ (n = 28), or 3-MA (n = 30). Briefly, 5-mm diameter fungus plugs from previously isolated

and reinoculated PDA culture were placed in 60-mm Petri dishes containing 10 ml of PDA

(control), PDA + 300 ng/ml rapamycin (Medchem Express, Monmouth Junction, NJ, USA), PDA

+ 1.5 mM chloroquine diphosphate (Sigma-Aldrich, St. Louis, Missouri, USA), or PDA + 10 mM

3-MA (Medchem Express). We then photographed plates to measure growth area (mm^2) using

ImageJ [54] and directly counted the staphylae on these Petri dishes under a stereo microscope

(40x magnification) to quantify staphyla density (number of staphylae/growth area). The

measurement of growth area also enabled assessment of other unintended inhibitory effects of

the added chemicals on cultivar performance. We tested for treatment effects (PDA-Control,

RAP, CQ, 3-MA) on mycelial growth and staphyla density in R version 4.0.2 [55] using a

Kruskal-Wallis test in *rstatix* version 0.7.0 [56] with pairwise post-hoc tests performed using

Dunn's Test in *rstatix* with adjusted p-values calculated using the false-discovery rate method.

392   *Transcriptome sequencing, assembly and differential expression analyses*

To detect whether upregulated transcripts in staphylae were enriched with autophagy

genes, we first collected staphylae and undifferentiated mycelia from axenic fungal cultures

isolated from an *A. colombica* colony (Ac2012-1) and grown on PDA medium for 30 days as

specified above. From individual Petri dishes, we then used a RNeasy plant mini kit (Qiagen,

Germany) to extract total RNA from differentiated staphyla (pooled 200 staphylae individually

collected with sterile acupuncture needles, n = 5 Petri dishes samples) and undifferentiated

mycelia (adjacent mycelia lacking staphylae, n = 5 Petri dishes samples). Samples were

immediately placed in Qiagen RLC buffer containing 10uM DTT and RNA was then extracted

using the manufacturer-specified protocol. These samples were sent to BGI Europe

(Copenhagen, Denmark) where mRNA enrichment with oligo dT and strand-specific libraries

were constructed using dUTP in the cDNA synthesis. For each sample, between 24 and 30

million clean 100bp paired-end reads were generated by a DNBseq-G400 sequencer (MGI
Tech, Shenzhen, China).

We used pooled clean reads from all samples (staphylae and mycelia) to assemble a *de*
novo transcriptome using Trinity-v2.12.0 [57] and default settings with the addition of Jaccard-
clip and strand-specific (SS) options, to reduce the generation of chimeric transcripts and
account for the SS library construction, respectively. All downstream tools accounted for SS
sequences (analyzing only positive strands or sequences). Using CD-HIT [58] to cluster highly
similar sequences (98% similarity), we reduced the assembly from 93,470 to 78,820 transcripts
ranging from 187 to 26,203 bp. We used Trinity built-in pipelines to first estimate transcript
abundance with RSEM v1.3.1 [59] and then build a transcript expression matrix. We then
performed a differential expression analysis with a trinity built-in pipeline using DESeq2 [60]
setting the analysis to filter for the differentially expressed transcripts (DETs) with $\log_2$ fold
change > 1.0 and *P* < 0.001. To annotate the DETs, we first converted transcripts to amino acid
sequences using Transdecoder (https://github.com/TransDecoder/TransDecoder) to identify the
longest open read frames (orfs) per transcript and translate them into amino acid sequences,
and then annotated the DETs with KEGG database [61] using the online tools BlastKOALA,
GhostKOALA [62] and KofamKOALA [63].

Data availability

423 All the generated RNA-seq datasets are available at NCBI under the BioProject ID
424 PRJNAXXXXXX.

Acknowledgements

This study was funded by a European Research Council Starting Grant (ELEVATE:
ERC-2017-STG-757810) to JZS. Benjamin Conlon assisted with statistical analyses. Useful
comments were provided by Jacobus Boomsma, Gareth W. Griffith and Pedro Elias
Marques. The images recreating stages of gongylidium formation in Figure 4 were
produced by Damond Kyllo. Assistance with transmission and scanning electron
microscopies was provided by the Core Facility for Integrated Microscopy, Faculty of

Health and Medical Sciences, and the Zoological Museum, both at University of

Copenhagen. Data processing and analysis were performed using Computerome, the National

Life Science Supercomputer at DTU (www.computerome.dk). Sylvia Mathiasen and Rasmus

Larsen provided general laboratory assistance.

References

1.   Solberg MF, Robertsen G, Sundt-Hansen LE, Hindar K, Glover KA. Domestication leads to increased predation susceptibility. Scientific Reports. 2020;10(1):1929.

2.   Milla R, Osborne CP, Turcotte MM, Violle C. Plant domestication through an ecological lens. Trends Ecol Evol. 2015;30(8):463-9.

3.   Gering E, Incorvaia D, Henriksen R, Conner J, Getty T, Wright D. Getting Back to Nature: Feralization in Animals and Plants. Trends Ecol Evol. 2019;34(12):1137-51.

4.   Comai L. The advantages and disadvantages of being polyploid. Nature Reviews Genetics. 2005;6(11):836-46.

5.   Stetter MG, Gates DJ, Mei W, Ross-Ibarra J. How to make a domesticate. Curr Biol. 2017;27(17):R896-R900.

6.   Edger PP, Poorten TJ, VanBuren R, Hardigan MA, Colle M, McKain MR, et al. Origin and evolution of the octoploid strawberry genome. Nat Genet. 2019;51(3):541-7.

7.   Piperno DR. Patterns, Process, and New Developments

The Origins of Plant Cultivation and Domestication in the New World Tropics. CurrAnthr. 2011;52(S4):S453-S70.

8.   Renner SS, Wu S, Pérez-Escobar OA, Silber MV, Fei Z, Chomicki G. A chromosome-level genome of a Kordofan melon illuminates the origin of domesticated watermelons. Proc Natl Acad Sci. 2021;118(23):e2101486118.

9.   Mueller UG, Gerardo NM, Aanen DK, Six DL, Schultz TR. The evolution of agriculture in insects. Annual Review of Ecology, Evolution, and Systematics. 2005:563-95.

10.   Schultz TR, Brady SG. Major evolutionary transitions in ant agriculture. Proc Natl Acad Sci. 2008;105(14):5435-40.

11.   Barrera CA, Sosa-Calvo J, Schultz TR, Rabeling C, Bacci Jr M. Phylogenomic reconstruction reveals new insights into the evolution and biogeography of Atta leaf-cutting ants (Hymenoptera: Formicidae). Syst Entomol. 2021;n/a(n/a).

12.   Boomsma JJ. Lifetime Commitment between Social Insect Families and Their Fungal Cultivars Complicates Comparisons with Human Farming. In: Schultz TR, Gawne R, Peregrine PN, editors. The Convergent Evolution of Agriculture in Humans and Insects. Vienna Series in Theoretical Biology. Cambridge, Massachusetts: MIT Press; 2022. p. 73-86.

13.   Frank SA. Host-symbiont conflict over the mixing of symbiotic lineages. Proceedings of the Royal Society of London Series B: Biological Sciences. 1996;263(1368):339-44.

14.   Axelrod R, Hamilton WD. The Evolution of Cooperation. Science. 1981;211(4489):1390-6.

15.   Boomsma JJ, Gawne R. Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. Biological Reviews. 2018;93(1):28-54.

16.   Leigh Jr EG. The evolution of mutualism. J Evol Biol. 2010;23(12):2507-28.

17.   Hölldobler B, Wilson EO. The leafcutter ants: civilization by instinct. New York: W. W Norton & Co.; 2011.

18.   Wirth R, Herz H, Ryel RJ, Beyschlag W, Hölldobler B. Herbivory of Leaf-Cutting
Ants. Germany: Springer-Verlag Berlin Heidelberg; 2003. 233 p.

19.   Crumiere AJJ, James A, Lannes P, Mallett S, Michelsen A, Rinnan R, et al. The
multidimensional nutritional niche of fungus-cultivar provisioning in free-ranging
colonies of a neotropical leafcutter ant. Ecol Lett. 2021;24(11):2439-51.

20.   Mueller UG, Schultz TR, Currie CR, Adams RM, Malloch D. The origin of the
attine ant-fungus mutualism. Q Rev Biol. 2001:169-97.

21.   Quinlan RJ, Cherrett JM. The role of fungus in the diet of the leaf-cutting ant
Atta cephalotes (L.). Ecol Entomol. 1979;4(2):151-60.

22.   Weber NA. Dry season adaptations of fungus-growing ants and their fungi. The
Anatomical Record. 1957;128(3):1.

23.   Wheeler WM. The fungus-growing ants of North America. Bulletin of the
American Museum of Natural History. 1907;23:139.

24.   Möller A. Die Pilzgärten einiger südamerikanischer Ameisen. A.F.W. S, editor.
Jena: Gustav Fischer Verlag; 1893.

25.   Powell RJ. The influence of substrate quality on fungus cultivation by some
Attine ants: University of Exeter; 1984.

26.   Weber NA. Fungus-growing ants. Science. 1966;153(3736):587-604.

27.   Swingle WT. Fungus Gardens in the Nest of an Ant (Atta tardigrada Buckl.)
near Washington. Proc Am Assoc Adv Sci. 1896;44th Meet.:2.

28.   Nygaard S, Hu H, Li C, Schiøtt M, Chen Z, Yang Z, et al. Reciprocal genomic
evolution in the ant–fungus agricultural symbiosis. Nat Commun. 2016;7:12233.

29.   Solomon SE, Rabeling C, Sosa-Calvo J, Lopes CT, Rodrigues A, Vasconcelos
HL, et al. The molecular phylogenetics of Trachymyrmex Forel ants and their fungal
cultivars provide insights into the origin and coevolutionary history of 'higher-attine' ant
agriculture. Syst Entomol. 2019;44(4):939-56.

30.   De Fine Licht HH, Schiøtt M, Rogowska-Wrzesinska A, Nygaard S, Roepstorff
P, Boomsma JJ. Laccase detoxification mediates the nutritional alliance between leaf-
cutting ants and fungus-garden symbionts. Proc Natl Acad Sci. 2013;110(2):583-7.

31.   Kooij PW, Rogowska-Wrzesinska A, Hoffmann D, Roepstorff P, Boomsma JJ,
Schiott M. *Leucoagaricus gongylophorus* uses leaf-cutting ants to vector proteolytic
enzymes towards new plant substrate. ISME J. 2014;8(5):1032-40.

32.   Schiøtt M, Rogowska-Wrzesinska A, Roepstorff P, Boomsma JJ. Leaf-cutting
ant fungi produce cell wall degrading pectinase complexes reminiscent of
phytopathogenic fungi. BMC Biol. 2010;8(1):156.

33.   Aylward FO, Khadempour L, Tremmel DM, McDonald BR, Nicora CD, Wu S, et
al. Enrichment and Broad Representation of Plant Biomass-Degrading Enzymes in the
Specialized Hyphal Swellings of Leucoagaricus gongylophorus, the Fungal Symbiont of
Leaf-Cutter Ants. PLOS ONE. 2015;10(8):e0134752.

34.   Schiøtt M, Boomsma JJ. Proteomics reveals synergy between biomass
degrading enzymes and inorganic Fenton chemistry in leaf-cutting ant colonies. eLife.
2021;10:e61816.

35.   Martin MM. The Biochemical Basis of the Fungus-Attine Ant Symbiosis.
Science. 1970;169(3940):16.

36.   Martin MM, Martin JS. The biochemical basis for the symbiosis between the
ant, Atta colombica tonsipes, and its food fungus. J Insect Physiol. 1970;16(1):109-19.

37. Khadempour L, Kyle JE, Webb-Robertson B-JM, Nicora CD, Smith FB, Smith
RD, et al. From Plants to Ants: Fungal Modification of Leaf Lipids for Nutrition and
Communication in the Leaf-Cutter Ant Fungal Garden Ecosystem. mSystems.
2021;6(2):e01307-20.

38. Shik JZ, Rytter W, Arnan X, Michelsen A. Disentangling nutritional pathways
linking leafcutter ants and their co-evolved fungal symbionts using stable isotopes.
Ecology. 2018;99(9):1999-2009.

39. Martin MM, Carman RM, MacConnell JG. Nutrients Derived from the Fungus
Cultured by the Fungus-Growing Ant Atta colombica tonsipes1. Ann Entomol Soc Am.
1969;62(1):11-3.

40. De Fine Licht HH, Boomsma JJ, Tunlid A. Symbiotic adaptations in the fungal
cultivar of leaf-cutting ants. Nat Commun. 2014;5(1):5675.

41. Kikuma T, Ohneda M, Arioka M, Kitamoto K. Functional Analysis of the *ATG8*
Homologue Ao*atg8* and Role of Autophagy in Differentiation and Germination in
Aspergillus oryzae. Eukaryot Cell. 2006;5(8):1328-36.

42. Pinan-Lucarré B, Paoletti M, Dementhon K, Coulary-Salin B, Clavé C.
Autophagy is induced during cell death by incompatibility and is essential for
differentiation in the filamentous fungus Podospora anserina. Mol Microbiol.
2003;47(2):321-33.

43. Liu X-H, Gao H-M, Xu F, Lu J-P, Devenish RJ, Lin F-C. Autophagy vitalizes the
pathogenicity of pathogenic fungi. Autophagy. 2012;8(10):1415-25.

44. Kües U, Navarro-González M. How do Agaricomycetes shape their fruiting
bodies? 1. Morphological aspects of development. Fungal Biology Reviews.
2015;29(2):63-97.

45. Elander PH, Minina EA, Bozhkov PV. Autophagy in turnover of lipid stores:
trans-kingdom comparison. J Exp Bot. 2017;69(6):1301-11.

46. Bartoszewska M, Kiel JA. The role of macroautophagy in development of
filamentous fungi. Antioxid Redox Signal. 2011;14(11):2271-87.

47. Levine B, Klionsky DJ. Development by Self-Digestion: Molecular Mechanisms
and Biological Functions of Autophagy. Dev Cell. 2004;6(4):463-77.

48. Klionsky DJ, Abdel-Aziz AK, Abdelfatah S, Abdellatif M, Abdoli A, Abel S, et al.
Guidelines for the use and interpretation of assays for monitoring autophagy (4th
edition)1. Autophagy. 2021;17(1):1-382.

49. Angeli-Papa J, Eymé J. Le champignon cultivé par la" fourmi-manioc",
Acromyrmex octospinosus Reich en Guadeloupe; résultats préliminaires sur le
mycélium en culturue pure et sur l'infrastructure des hyphes. C r hebd Séanc Acad Sci,
Paris D   1979;281:21-4.

50. Noda T, Ohsumi Y. Tor, a Phosphatidylinositol Kinase Homologue, Controls
Autophagy in Yeast*. J Biol Chem. 1998;273(7):3963-6.

51. Mauthe M, Orhon I, Rocchi C, Zhou X, Luhr M, Hijlkema K-J, et al. Chloroquine
inhibits autophagic flux by decreasing autophagosome-lysosome fusion. Autophagy.
2018;14(8):1435-55.

52. Wu Y-T, Tan H-L, Shui G, Bauvy C, Huang Q, Wenk MR, et al. Dual Role of 3-
Methyladenine in Modulation of Autophagy via Different Temporal Patterns of Inhibition
on Class I and III Phosphoinositide 3-Kinase*. J Biol Chem. 2010;285(14):10850-61.

53.   Shik JZ, Kooij PW, Donoso DA, Santos JC, Gomez EB, Franco M, et al.
Nutritional niches reveal fundamental domestication trade-offs in fungus-farming ants.
Nat Ecol Evol. 2021;5(1):122-34.

54.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of
image analysis. Nat Methods. 2012;9(7):671-5.

55.   R Core Team. R: a language and environment for statistical computing.
Foundation for Statistical Computing. Vienna, Austria 2020.

56.   Kassambara A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R
package version 0.7.0. ed2021.

57.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-
length transcriptome assembly from RNA-Seq data without a reference genome. Nat
Biotechnol. 2011;29(7):644-52.

58.   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-
generation sequencing data. Bioinformatics. 2012;28(23):3150-2.

59.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. BMC Bioinformatics. 2011;12(1):323.

60.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550.

61.   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes.
Nucleic Acids Res. 2000;28(1):27-30.

62.   Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG
Tools for Functional Characterization of Genome and Metagenome Sequences. J Mol
Biol. 2016;428(4):726-31.

63.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.
KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score
threshold. Bioinformatics. 2019;36(7):2251-2.

64.   Grell MN, Linde T, Nygaard S, Nielsen KL, Boomsma JJ, Lange L. The fungal
symbiont of *Acromyrmex* leaf-cutting ants expresses the full spectrum of genes to
degrade cellulose and other plant cell wall polysaccharides. BMC Genomics. 2013;14.

65.   Mueller UG, Mikheyev AS, Hong E, Sen R, Warren DL, Solomon SE, et al.
Evolution of cold-tolerant fungal symbionts permits winter fungiculture by leafcutter ants
at the northern frontier of a tropical ant–fungus symbiosis. Proc Natl Acad Sci.
2011;108(10):4053-6.

66.   Nichols-Orians CM. Environmentally Induced Differences in Plant Traits:
Consequences for Susceptibility to a Leaf-Cutter Ant. Ecology. 1991;72(5):1609-23.

67.   Howard JJ. Leafcutting Ant Diet Selection: The Role of Nutrients, Water, and
Secondary Chemistry. Ecology. 1987;68(3):503-15.

68.   Howard JJ. Leafcutting and Diet Selection: Relative Influence of Leaf Chemistry
and Physical Features. Ecology. 1988;69(1):250-60.

69.   Nichols-Orians CM, Schultz JC. Interactions among leaf toughness, chemistry,
and harvesting by attine ants. Ecol Entomol. 1990;15(3):311-20.

70.   Crumiere AJJ, Mallett S, Michelsen A, Rinnan R, Shik JZ. Nutritional challenges
of feeding a mutualist: Testing for a nutrient-toxin tradeoff in fungus-farming leafcutter
ants. Ecology. 2022;103(6):e3684.

71.   Mundim FM, Costa AN, Vasconcelos HL. Leaf nutrient content and host plant
selection by leaf-cutter ants, Atta laevigata, in a Neotropical savanna. Entomol Exp
Appl. 2009;130(1):47-54.

72.   Berish CW. Leaf-Cutting Ants (Atta cephalotes) Select Nitrogen-Rich Forage.
The American Midland Naturalist. 1986;115(2):268-76.

73.   Howard JJ, Cazin J, Wiemer DF. Toxicity of terpenoid deterrents to the
leafcutting antAtta cephalotes and its mutualistic fungus. J Chem Ecol. 1988;14(1):59-
69.

74.   Jandér KC, Herre EA. Host sanctions and pollinator cheating in the fig tree-fig
wasp mutualism. Proc R Soc B. 2010;277(1687):1481-8.

75.   Heil M, McKey D. Protective Ant-Plant Interactions as Model Systems in
Ecological and Evolutionary Research. Annual Review of Ecology, Evolution, and
Systematics. 2003;34(1):425-553.

76.   Kiers ET, Rousseau RA, West SA, Denison RF. Host sanctions and the
legume–rhizobium mutualism. Nature. 2003;425(6953):78-81.

77.   Bass M, Cherrett JM. Leaf-Cutting Ants (Formicidae, Attini) Prune Their Fungus
to Increase and Direct Its Productivity. Funct Ecol. 1996;10(1):55-61.

78.   Schiøtt M, De Fine Licht HH, Lange L, Boomsma JJ. Towards a molecular
understanding of symbiont function: Identification of a fungal gene for the degradation
of xylan in the fungus gardens of leaf-cutting ants. BMC Microbiol. 2008;8(1):40.

79.   Moller IE, De Fine Licht HH, Harholt J, Willats WGT, Boomsma JJ. The
Dynamics of Plant Cell-Wall Polysaccharide Decomposition in Leaf-Cutting Ant Fungus
Gardens. PLOS ONE. 2011;6(3):e17506.

80.   Nygaard S, Zhang G, Schiøtt M, Li C, Wurm Y, Hu H, et al. The genome of the
leaf-cutting ant Acromyrmex echinatior suggests key adaptations to advanced social
life and fungus farming. Genome Res. 2011;21(8):1339-48.

81.   Shik JZ, Gomez EB, Kooij PW, Santos JC, Wcislo WT, Boomsma JJ. Nutrition
mediates the expression of cultivar–farmer conflict in a fungus-growing ant. Proc Natl
Acad Sci. 2016;113(36):10121-6.

82.   Pinto-Tomás AA, Anderson MA, Suen G, Stevenson DM, Chu FST, Cleland
WW, et al. Symbiotic Nitrogen Fixation in the Fungus Gardens of Leaf-Cutter Ants.
Science. 2009;326(5956):1120-3.

83.   Khadempour L, Fan H, Keefover-Ring K, Carlos-Shanley C, Nagamoto NS,
Dam MA, et al. Metagenomics Reveals Diet-Specific Specialization of Bacterial
Communities in Fungus Gardens of Grass- and Dicot-Cutter Ants. Frontiers in
Microbiology. 2020;11.

84.   Sapountzis P, Zhukova M, Shik JZ, Schiott M, Boomsma JJ. Reconstructing the
functions of endosymbiotic Mollicutes in fungus-growing ants. eLife. 2018;7:e39209.

85.   Zhukova M, Sapountzis P, Schiøtt M, Boomsma J. Phylogenomic analysis and
metabolic role reconstruction of mutualistic Rhizobiales hindgut symbionts of
Acromyrmex leaf-cutting ants. FEMS Microbiol Ecol. In press.

86.   Kooij PW, Aanen DK, Schiøtt M, Boomsma JJ. Evolutionarily advanced ant
farmers rear polyploid fungal crops. J Evol Biol. 2015;28(11):1911-24.

87.   Xiang X, Beckwith SM, Morris NR. Cytoplasmic dynein is involved in nuclear
migration in Aspergillus nidulans. Proc Natl Acad Sci. 1994;91(6):2100-4.

88.   Xiang X. Nuclear movement in fungi. Semin Cell Dev Biol. 2018;82:3-16.

89.   Yamamoto A, Hiraoka Y. Cytoplasmic dynein in fungi: insights from nuclear
migration. J Cell Sci. 2003;116(22):4501-12.

90.   Plamann M, Minke PF, Tinsley JH, Bruno KS. Cytoplasmic dynein and actin-
related protein Arp1 are required for normal nuclear distribution in filamentous fungi. J
Cell Biol. 1994;127(1):139-49.

91.   Gehrmann T, Pelkmans JF, Ohm RA, Vos AM, Sonnenberg ASM, Baars JJP, et
al. Nucleus-specific expression in the multinuclear mushroom-forming fungus *Agaricus
bisporus* reveals different nuclear regulatory programs. Proc Natl Acad Sci U S A.
2018;115(17):4429-34.

92.   Stalpers JA. Identification of wood-inhabiting Aphyllophorales in pure culture.
Studies in Mycology. 1978;16:248.

93.   Nobles MK. Identification of cultures of wood-inhabiting Hymenomycetes.
Canadian Journal of Botany. 1965;43(9):1097-139.

Figures and table

Fig. 1: The *Leucoagaricus gongylophorus* fungal cultivar produces gongylidia as specialized nutritional reward structures for leafcutter ants. A-C) Gongylidium cells are typically depicted as a bulb at the end of a filament in the apical hyphal compartment separated by a septum (arrowheads). **D)** Gongylidia frequently exhibit more complex branching, with bulbs between filaments or in lateral branches of single hyphal cells delimited by septa. **E)** Individual gongylidium cells are polykaryotic [86], meaning that they have many haploid nuclei (white dots). Here, we show that in mature non-branching gongylidium cells, these nuclei occur at the base of the bulb (below a single large vacuole) and in the filament. Nuclei were visualized using DAPI staining. **F)** Each gongylidium cell contains a large vacuole (arrowheads). **G)** Staphyla grow in discrete patches at the surface of the fungus garden matrix in the middle garden stratum (arrowheads). **H)** Gongylidium cells have thin cell walls ranging from 120 to 220 nm (cw). Images produced by light microscopy (panels A, D, F), fluorescence microscopy stained with DAPI (panel E), SEM (panels B, C, G) and TEM (panel H). Scale bars: A-F = 20 μm, G = 100 μm, H = 200 nm.

Fig. 2: Autophagic recycling of cellular material. A) The autophagosomes that are diagnostic of autophagy are vesicles with double-layered membranes (arrows) and are produced by stretches of endoplasmic reticulum membranes (arrowheads). **B)** Autophagosomes sequester cytoplasmic components including glycogen (arrowhead) and **C)** mitochondria which are then delivered to vacuoles in gongylidia bulbs for further degradation. **D-E)** Vacuolar expansion is mediated by autophagosomes that lose their outer membrane after fusing with the vacuole. These single-membraned autophagic bodies are vesicles that can be seen inside the vacuole prior to their degradation (arrowhead). Images acquired by TEM (A-C), phase contrast microscopy (D) and confocal microscopy stained with dextran-Alexa fluor 647 (E). Scale bars A-C = 500 nm, D-E = 20 μm.

Fig. 3: Experimental evidence that autophagic recycling of the fungal cultivar's own cellular material mediates gongylidia formation. Gongylidium density was significantly inhibited when *L. gongylophorus* was grown on potato dextrose agar supplemented with one autophagy inhibitor chloroquine (CQ, n = 28) or 3-methyladenine (3-MA, n = 30)) relative to control (PDA, n = 40) and an autophagy promoter rapamycin (RAP, n = 27)). Representative Petri-dishes with control (PDA; top) and inhibition (CQ; bottom) are displayed at the right, black outlines indicate the radial growth area of cultivars and white fungal clusters in the control are the staphylae. Different letters above the boxes indicate significant differences determined by a post-hoc Dunn's pairwise test ($p < 0.05$) and horizontal bars indicate the distribution means.

Fig. 4: The hypothesized stages of autophagy-mediated gongylidium development. A) An unknown mechanism (potentially starvation mediated by ant pruning [77]) triggers the widening of ordinary hyphae. As the hypha elongates, the nuclei (orange circles) naturally migrate towards the hyphal tip. **B)** Mediated primarily by an autophagic process, a large vacuole expands with the fusion of newly formed double membrane vesicles called autophagosomes (blue membraned vesicles) that sequester material present in the cytosol like glycogen (black and gray aggregates) and damaged mitochondria (brown indented ovals). This process is indicated by the proliferation of endoplasmic reticulum membranes (blue membranes around nuclei) that produce autophagosomes. **C)** The fusion of autophagosomes into vacuoles mediates their expansion and either forces the apical bulb swelling while also halting further apical growth by excluding nuclei from the hyphal tip or produce an intercalary bulb when the vacuole is located among nuclei. **D)** This process repeats in up to hundreds of adjacent hyphae that become tangled to form the staphyla.

732 **Table 1:** Upregulated transcripts in autophagy pathway annotated with KEGG database (see Fig. S3).

Tissue	KEGG	Gene names	No. of 733 transcripts
Staphylae	K06902	ATG22; UMF1; MFS transporter, UMF1 family	1 734
	K07897	Ypt7; RAB7A; Ras-related protein Rab-7A	1
	K08331	ATG13; autophagy-related protein 13	1
	K08337	ATG7; ubiquitin-like modifier-activating enzyme ATG7	3
	K12767	RIM15; serine/threonine-protein kinase RIM15	2
	K17606	IGBP1, TAP42; immunoglobulin-binding protein 1	1
	K17888	ATG10; ubiquitin-like-conjugating enzyme ATG10	1
	K17900	ATG15, AUT5; lipase ATG15	1
	K01336	PRB1; cerevisin	5
	K08341	ATG8, GABARAP, LC3; GABA(A) receptor-associated protein	1
	K19800	SCH9; serine/threonine protein kinase SCH9	1
	K20177	VPS3, TGFBRAP1; vacuolar protein sorting-associated protein 3	2
	K20195	MON1; vacuolar fusion protein MON1	2
Mycelium	K01336	PRB1; cerevisin	1
	K04464	MAPK7; mitogen-activated protein kinase 7	2
	K08337	ATG7; ubiquitin-like modifier-activating enzyme ATG7	2
	K12761	SNF1; carbon catabolite-derepressing protein kinase	1
	K17906	ATG2; autophagy-related protein 2	1
	K21157	SAK1; SNF1-activating kinase 1	1
	K06655	PHO85; negative regulator of the PHO system	1

735   **Supplementary figures**

738   **Fig. S1: Representative counts of gongylidia per staphylae in *L. gongylophorus* from**

739   **different leafcutter ants' species.** A) Staphylae from in vitro culture without ant manipulation.

740   B-E) staphylae from colonies fungus garden. Yellow marks indicate individual gongylidia count

741   on ImageJ. Scale bars sizes are indicated in each image.

Fig. S2: Mycelial growth distribution (area per treatment). The mycelial growth showed significant differences between 3-MA and all other treatments (PDA:3-MA, p_{adj} = 0.004; RAP:3-MA, p_{adj} = 0.015; CQ:3-MA, p_{adj} = 0.001), but no other significant pairwise comparisons. Thus, while both autophagy inhibition treatments resulted in staphyla reduction, it is possible that 3-MA negatively influenced staphyla density through unknown indirect effects on cultivar performance. Horizontal bars indicate the distribution means.

Fig. S3: Autophagy metabolic pathway map displaying gene expression in *L. gongylophorus.* Highlighted genes show upregulated transcripts in staphylae (red), mycelium (cyan) or in both tissues (magenta). The steps of the pathway are illustrated on the right showing where the products of these genes will act during autophagy. Adapted from KEGG website (www.kegg.jp).

Video S1: Time lapse of staphyla stained with dextran-Alexa Fluor 647 under confocal microscope over 20 minutes. Red arrows point gongylidia in which is possible to observe autophagic bodies trapped within the vacuole. Some of these vesicles are filled with dextran and appear more fluorescent than the vacuole lumen, others are filled with non-fluorescent material and appear darker than the vacuole.