Title

Cardio-audio synchronization elicits neural and cardiac surprise responses in human

wakefulness and sleep

Short Title

Cardio-audio based prediction of sounds in wakefulness and sleep

Authors and Affiliations

Andria Pelentritou[1*], Christian Pfeiffer[2], Sophie Schwartz[3,4] & Marzia De Lucia[1*]

*Corresponding authors: Antria.Pelentritou@chuv.ch ; Marzia.De-Lucia@chuv.ch

[1]Laboratoire de Recherche en Neuroimagerie (LREN), University Hospital (CHUV) and

University of Lausanne (UNIL), Lausanne, Switzerland

[2]Robotics and Perception Group, University of Zurich, Zurich, Switzerland

[3]Department of Neuroscience, Faculty of Medicine, University of Geneva, Geneva, Switzerland

[4]Swiss Center for Affective Sciences, University of Geneva, Geneva, Switzerland

Abstract

The human brain can infer temporal regularities in auditory sequences with fixed sound-to-sound intervals and in pseudo-regular sequences where sound onsets are locked to cardiac inputs. Here, we investigated auditory and cardio-audio regularity encoding during sleep, when reduced vigilance may result in altered bodily and environmental stimulus processing. Using electroencephalography and electrocardiography in healthy volunteers (N=26) during wakefulness and sleep, we measured the response to unexpected sound omissions within three regularity conditions: isochronous, with fixed sound-to-sound intervals, synchronous, where sound and heartbeat are temporally coupled, and a control condition without specific regularity. The isochronous and synchronous sequences induced a modulation of the omission-evoked neural response in wakefulness and N2 sleep, the latter accompanied by a background oscillatory activity reorganization. Cardio-audio regularity encoding was accompanied by a heartbeat deceleration upon omissions in all vigilance states. The violation of auditory and cardio-audio regularity elicits neural and cardiac surprise responses across vigilance stages.

Introduction

The processing of auditory regularity is a basic brain mechanism that enables the rapid detection of unexpected stimuli which can persist even in altered states of consciousness such as sleep and coma [1]. The mechanism underlying auditory regularity encoding has mostly been studied by investigating the neural responses to deviant sounds interrupting a sequence of repeated standard stimuli as reported in healthy human wakefulness [2,3], during sleep (*e.g.* [4–6]), and in disorders of consciousness patients [7–9]. This rudimental component of auditory discrimination, known as mismatch negativity (MMN), has often been interpreted in the framework of the predictive coding theory [2,3,10,11]. According to this theory, the MMN may arise from the contribution of multiple, non-exclusive mechanisms including repetition suppression in response to frequent stimuli and generation of a 'prediction error' following an unexpected mismatch between the predicted and presented stimuli. The predictive nature of the neural responses to violations within regular auditory sequences has received experimental support from studies on unexpected omissions [12–16] wherein the top-down prediction is not confounded by the neural response to deviant sound stimuli. In these paradigms, the neural response to rare deviant stimuli arises from a violation of 'what' is expected in the sequence. The mechanisms underlying the expectation of 'when' a sound is expected in the context of regular auditory stimuli remain far less studied and largely unresolved [17]. In this context, recent studies demonstrated that auditory prediction is not limited to fixed sound-to-sound regularities but it extends to pseudo-regular sequences in which temporal prediction is induced by locking the sound onsets to interoceptive inputs, i.e. internally-generated signals such as the heartbeat [18–20]. To date, it remains unknown whether interoceptive signals similarly affect auditory processing during sleep, when consciousness and attentional levels are altered, and perception of the external environment is reduced compared to wakefulness.

Here, first, we aimed at investigating whether auditory regularity encoding is preserved across sleep stages as revealed by the neural response to sound omissions and second, we aimed

at testing whether detecting regularity in trains of auditory stimuli may benefit from their temporal alignment (or misalignment) with bodily signals such as the heartbeats. We hypothesized that – as the brain gradually disconnects from the environment during sleep – bodily signals may play an important role in informing auditory regularity encoding and the detection of unexpected violation of such regularities. Healthy volunteers underwent two separate sessions of simultaneous electroencephalography (EEG) and electrocardiography (ECG) recordings during wakefulness and full night sleep. Participants passively listened to two possible varieties of auditory regularities (Fig. 1). In the first condition, sounds were presented at a fixed short delay relative to the ongoing heartbeat (synchronous or synch condition), in the second, sounds were presented at a fixed sound-to-sound interval (isochronous or isoch condition) and a third condition, where sounds were presented without any specific regularity (asynchronous or asynch condition) served as a control condition. To assess the strength of regularity encoding in all conditions, we measured the EEG and cardiac responses - measured by ECG - to unexpected omissions interspersed within the auditory sequences. Upon sound omissions in wakefulness, we expected prediction error generation in the synch and isoch conditions driven by auditory regularity and cardio-audio regularity.

Fig. 1. Experimental paradigm overview. The data acquisition was organized in four block types, administered in pseudo-random order. No auditory stimulation was performed during the Baseline condition. In the Synchronous condition, the R peaks (circles) were detected online in the ECG signal and the sounds were administered at a fixed 52 ms R-to-sound. In the Asynchronous condition, sound-to-sound intervals from a preceding Synchronous blocks were administered resulting in variable R-to-sound and sound-to-sound intervals. In the Isochronous condition, the median RR interval from a

80   preceding Synchronous block was computed and served as the fixed sound-to-sound interval. Sound

81   omissions occurred pseudo-randomly for 20% of trials within each block.

Results

1. Participant demographics and sleep characteristics

1.1. Participants

Twenty-six healthy volunteers participated in the study (14 female; 1 left-handed; mean age: 27 years, range: 20-35 years) and each took part in a wakefulness and sleep recording session. All 26 participants were included for wakefulness while the sleep dataset included 25 participants (13 female; 1 left-handed; mean age: 26 years, range: 20-35 years) due to malfunctioning equipment during the sleep session for 1 participant.

We performed the analyses of the simultaneously acquired EEG and ECG signals. For the EEG analysis, the data of 5 participants were excluded for the wakefulness session due to excessive artifactual periods during acquisition (which yielded a loss of more than 50% of the trials), resulting in the inclusion of 21 out of 26 participants. For the sleep session, from 25 participants, we included 12 for N1, 23 for N2, 14 for N3, and 13 for REM sleep. For the ECG analysis, we included all 26 participants for the wakefulness session and 15 for N1 sleep, 24 for N2 sleep, 18 for N3 sleep, and 15 for REM sleep (S1 Table & S2 Table).

1.2. Sleep characteristics

The sleep characteristics recorded during the experimental night across the eligible population (N=25) are summarized in Table 1.

Table 1. Group averaged sleep characteristics. Total Bed Time calculated based on Lights Off and Lights On time points. Total Sleep Time computed as the period between the Sleep Onset and final awakening, uncovered by the sleep staging, excluding wakefulness periods but including possible micro-arousal periods during the night. Percentages of sleep stage periods (N1, N2, N3, REM) during the session were computed in relation to the Total Sleep Time. Sleep Efficiency (%) = Total Sleep Time/Total Bed Time for each participant×100. SD = Standard Deviation.

N=25	Mean	SD
Total Bed Time (min)	420.3	93.8
Total Sleep Time (min)	302.4	107.9
Sleep Onset (min)	24.6	19.1
Sleep Efficiency (%)	69.8	12.9
N1 (%)	12.9	4.5
N2 (%)	57.4	6.9
N3 (%)	15.5	8.8
REM (%)	14.1	5.5

2. Quality control analysis

We conducted a series of control analyses on the heart rate and the relation between heartbeat

and sound onsets across auditory conditions in order to identify possible factors influencing

the neural and cardiac response to sound omissions during the wakefulness (AWAKE: N=21)

and sleep (N1: N=12; N2: N=23; N3: N=14; REM: N=13) sessions. In the following, we will

refer to R as the latency of the R peak of the ECG waveform (Fig. 1), S represents the sound

onset latency, RR the R peak-to-R peak latency, SS the sound-to-sound latency, RS the R

peak-to-sound latency, SR the sound to the next R peak latency, and variability is calculated

as the standard error of the mean (SEM). One way repeated measures Friedman tests with

factor Condition (synch, asynch, isoch and baseline - a control condition without auditory

stimulation - where relevant) were performed on the group RR interval, RR variability, SS

interval, SS variability, RS interval and RS variability for sound trials, as well as SR interval

and SR variability for omission trials (Fig. S1-4).

In the synch condition, we observed an average RS interval of 51.8 ms (SEM = 1.1 ms) for

sound trials in wakefulness and sleep, and -2.2 ms (SEM = 5.2 ms) for the other auditory

conditions (figure A in S1 Fig. for neural quality control analyses; see Materials and Methods,

Auditory conditions for the full cohort results). By construction, we expected lower RS variability

in the synch relative to the two other conditions (isoch and asynch), as was confirmed by one

way repeated measures Friedman tests with factor Auditory Condition (synch, asynch, isoch)

(figure B in S1 Fig.; p<0.0005; AWAKE: $\chi^2_{(1,3)}$=31.5; N1: $\chi^2_{(1,3)}$=18.2; N2: $\chi^2_{(1,3)}$=34.5; N3:

$\chi^2_{(1,3)}$=21.1; REM: $\chi^2_{(1,3)}$=19.5; all post-hoc paired Wilcoxon signed-rank tests showed lower

variability in the synch compared to the asynch or isoch condition). In addition, as expected,

SR within omission was more variable in the asynch and isoch conditions compared to the

synch condition (figure B in S2 Fig.; p<0.0005; AWAKE: $\chi^2_{(1,3)}$=31.7; N1: $\chi^2_{(1,3)}$=18.2; N2:

$\chi^2_{(1,3)}$=35.0; N3: $\chi^2_{(1,3)}$=21.6; REM: $\chi^2_{(1,3)}$=19.5; post-hoc Wilcoxon signed-ranked tests

confirmed significant differences). This first series of control analyses demonstrated that the

experimental manipulation, inducing an online fixed temporal alignment between R peak and

sound in the synch and a variable one in the other conditions, was successful.

The analysis of auditory regularity based on the SS interval confirmed higher variability during

the synch and asynch compared to the isoch condition (figure B in S3 Fig.; p<0.0005; AWAKE:

$\chi^2_{(1,3)}$=28.6; N1: $\chi^2_{(1,3)}$=18.2; N2: $\chi^2_{(1,3)}$=36.3; N3: $\chi^2_{(1,3)}$=19.0; REM: $\chi^2_{(1,3)}$=16.8; corroborated by

post-hoc Wilcoxon signed-rank tests) with an average variability value of 1.5 ms in wakefulness

and 3.4 ms across sleep stages for the isoch condition. This second analysis confirmed that

the isoch condition was characterized by highly regular sound-to-sound intervals in comparison

to the other conditions.

During the sleep session, the heartbeat was not globally affected during N2, N3 or REM sleep

by the different cardio-audio and audio-audio stimulus presentation conditions as shown by

1x4 repeated measures Friedman tests on average RR intervals with factor Condition (synch,

asynch, isoch, baseline) (figure A in S4 Fig.; p>0.05). Conversely, the same analysis

performed in wakefulness and N1 sleep revealed significant differences in RR intervals (figure

A in S4 Fig.; AWAKE: $\chi^2_{(1,3)}$=14.4, p<0.005; N1: $\chi^2_{(1,3)}$=12.3, p<0.05). Post-hoc Wilcoxon

signed-rank tests uncovered that significantly reduced RR intervals were specific to the

baseline condition (AWAKE: p<0.005 for synch/asynch/isoch *vs* baseline; N1: p<0.005 for

asynch/isoch *vs* baseline) while no differences were observed across auditory conditions (p>0.05).

Finally, we carried out a control analysis on the ECG waveforms to exclude the potential confound of a different degree of contamination of the ECG artifacts in the EEG between conditions of interest upon interpretating the differential EEG signals locked to the R peak. To this aim, we performed a time-wise ECG waveform comparison of the grand averaged ECG trials time-locked to R peaks during sound omissions. Statistical analysis based on non-parametric cluster permutation statistics (p<0.05, two tailed) contrasting paired experimental conditions (synch, asynch, isoch, baseline) for each vigilance state (S5 Fig.) revealed no significant differences (p>0.05) in any of the sleep stages and for any of the comparisons. In wakefulness, significant differences (p<0.05) were observed in the synch *vs* baseline comparison between 365 ms and 446 ms and in the asynch *vs* baseline comparison between 368 ms and 443 ms.

Overall, these last series of analyses suggest that the heartbeat and ECG characteristics were well matched across auditory conditions and as a consequence, they do not represent a main confounding factor for explaining the cardiac and neural differential responses across regularity types.

3. Neural omission response

3.1. Auditory response comparisons

Auditory evoked potentials (AEPs) were computed as the average of the peri-stimulus EEG epochs between 100 ms pre-stimulus and 500 ms post-stimulus onset in the synch, asynch and isoch conditions for the wakefulness session (AWAKE: N=21) and all sleep stages (N1: N=12; N2: N=23; N3: N=14; REM: N=13). The AEPs revealed the expected event-related potential components including the N100 at approximately 100 ms post-stimulus onset and similar AEP morphologies across sleep stages provided qualitative evidence that sound stimuli were processed by the participants' brain (Fig. 2). Cluster-based permutation statistical

analysis confirmed no significant differences between the isoch and asynch AEPs in

wakefulness or any of the sleep stages (p>0.05); suggesting auditory processing of the single

sounds within the sequence was similar across conditions. We refrained from performing these

comparisons in the synch condition as the AEPs in that case were contaminated with the

response to the heartbeat.

Fig. 2. AEPs in wakefulness and sleep. Grand averaged AEPs in the isoch (blue lines) and asynch

(black lines) across 19 EEG sensors (10-20 system; for visualization purposes) in the wakefulness

session (AWAKE) and all stages of sleep. AEPs in wakefulness display the expected

electrophysiological signatures such as the N100, as the first and highest in amplitude component. AEPs

are also observed in all sleep stages, yet with differences in signal morphology, signal amplitude, and

response latency. Note that the synch condition AEPs are not depicted as they would be superimposed

with the heartbeat-evoked potential.

3.2. Omission response in auditory temporal regularities

3.2.1. Omission response comparisons

In order to test our hypothesis that auditory regularities induced an expectation of incoming

stimuli, we derived omission responses during sound omissions in the isoch and asynch

conditions in wakefulness (AWAKE: N=21) and all sleep stages (N1: N=12; N2: N=23; N3:

N=14; REM: N=13). Average sound-based omission evoked potentials (OEPs) were calculated

by extracting epochs from the continuous EEG recordings that were time-locked to the average

sound interstimulus interval. A random selection of epochs were extracted from continuous

baseline recordings, such that the latencies between epoch onset and closest heartbeat (i.e.

R peak) were matched to the trial onsets in the sound-based isoch and asynch conditions at

the single-trial level.

Based on previous reports of neural responses to omissions within regular auditory sequences

during wakefulness (*e.g.* [12,15,21]), here, we expected central negativity in the isoch condition

(where temporal regularity existed in sound stimuli) between 100 ms and 250 ms. In the

wakefulness session, the cluster-based permutation test comparing the isoch *vs* asynch

condition yielded a significant negative cluster ($p < 0.05$, Cohen's d = 0.67) at 226 ms to 288

ms following expected sound onset in central scalp electrodes (Fig. 3A). The existence of an

omission response in the isoch condition is further confirmed by the isoch *vs* baseline

comparison (figure A in S6 Fig.), revealing a significant negative cluster ($p < 0.05$, Cohen's d =

0.91) at 103 ms to 170 ms following expected sound onset on anterior-central scalp electrodes.

A second significant negative cluster ($p < 0.05$, Cohen's d = 1.23) was observed for the same

comparison at 251 ms to 295 ms on posterior electrodes and a significant positive cluster

($p < 0.05$, Cohen's d = 1.02) at roughly the same latency between 246 ms and 303 ms on

anterior-central electrodes. Despite the absence of a regularity rule in the asynch condition,

the asynch *vs* baseline comparison in wakefulness also revealed a significant positive cluster

($p < 0.05$, Cohen's d = 0.82) on central scalp electrodes at 197 ms to 288 ms following expected

sound onset (figure B in S6 Fig.). This points to some degree of inference of the temporal

relationship across auditory stimuli being established during asynch sequence administration

in wakefulness, potentially driven by the pseudo-regularity of auditory inputs.

Sleep OEPs differences across auditory conditions were only significant in N2 sleep

comparisons. In addition, the isoch *vs* baseline and the asynch *vs* baseline comparisons during

N2 sleep did not reveal significant differences. Isoch *vs* asynch OEP differences were largely

similar between wakefulness and N2 sleep, at least in terms of latency relative to expected

sound onset. In more detail, statistical evaluation of the isoch and asynch condition differences

(Fig. 3B) identified a significant negative cluster ($p<0.05$, Cohen's d = 0.66) at 85 ms to 223

ms following expected sound onset localized to posterior-central scalp electrodes. Our results

in wakefulness and N2 sleep suggest that in the isoch condition, the fixed sound-to-sound

interval induced an expectation of upcoming sounds and resulted in a neural surprise response

upon violation of the regularity rule during omissions.

3.2.2. Sound *vs* omission response comparisons

We contrasted the isoch AEPs and OEPs in wakefulness and N2 sleep, in order to ensure that

the observed neural omission responses were indeed related to the unfulfilled sound

expectation and not due to differences of neural responses to stimuli across blocks. In

wakefulness (figure A in S7 Fig.), cluster permutation statistical analysis contrasting the isoch

AEPs and isoch OEPs revealed two significant positive clusters, a first ($p<0.0005$) between 72

ms and 362 ms and a second ($p<0.05$) between 394 ms and 500 ms, both localized to posterior

electrodes. Two negative clusters were additionally identified, a first ($p<0.0005$) between 66

ms and 190 ms and a second between 259 ms and 360 ms, both on central scalp electrodes.

In N2 sleep (figure B in S7 Fig.), the isoch AEP and OEP contrast yielded one positive cluster

($p<0.005$) between 136 ms and 407 ms on central electrodes and one negative cluster

($p<0.0005$) between 138 ms and 500 ms on outer electrodes. These latencies overlap with

those obtained when comparing OEPs between isoch *vs* baseline (and isoch *vs* asynch) and

provide support to our interpretation that these neural omission responses were due to

regularity violation upon omission of expected sounds.

3.3. Omission response in cardio-audio regularities

3.3.1. Omission response comparisons

To investigate the effect of cardio-audio synchronicity on omission responses, we derived

heartbeat-evoked potentials during sound omissions (OHEPs) in the synch and asynch

conditions for the wakefulness session (AWAKE: N=21) and all sleep stages (N1: N=12; N2:

N=23; N3: N=14; REM: N=13). Average OHEPs were calculated by extracting epochs from

the continuous EEG recordings that were time-locked to the first R peak of the ECG signal

during omissions. As an additional control condition for this analysis, we extracted a random

selection of R peaks in the ECG signal to derive average HEPs from the continuous EEG

recordings in the baseline condition.

In the wakefulness session, we expected to observe differences in the HEPs when comparing

the synch to asynch and the synch to baseline conditions, as a consequence of the

predictability of sound onset in the synch condition, based on the fixed delay between R peaks

and sounds [19]. We expected no differences when comparing asynch *vs* baseline. For the

synch *vs* asynch comparison (Fig. 3C), the cluster-based permutation test ($p<0.05$, two-tailed)

revealed a significant positive cluster ($p<0.05$, Cohen's d = 1.36) at 223 ms to 274 ms following

R peak on posterior-central scalp electrodes. A significant negative cluster ($p<0.05$, Cohen's

d = 0.91) at 162 ms to 267 ms following R peak was additionally observed in this comparison

within two distinct intervals, a first at 162 ms to 213 ms on anterior-central electrodes and a

second at 217 ms to 267 ms on posterior-lateral rightmost electrodes. The synch *vs* baseline

comparison (figure C in S6 Fig.) yielded one significant negative cluster ($p<0.0005$, Cohen's d

= 1.21) on posterior-central scalp electrodes at 36 ms to 275 ms. As anticipated, the asynch

vs baseline comparison showed no significant differences (figure D in S6 Fig.).

Fig. 3. Neural omission response in wakefulness and N2 sleep. OEPs (time-locked to mean SS interval) and OHEPs (time-locked to the R peak) comparisons during sound omissions for the isoch (blue lines) and synch (red lines) conditions compared to the asynch condition (black lines) during wakefulness (AWAKE) and N2 sleep (N2). Top panels show grand average EEG waveforms averaged

over all electrodes in the significant negative clusters. Shaded regions indicate ±SEM across participants. Middle panels display the number of electrodes within each significant cluster derived from cluster permutation statistical analysis (p<0.05, two-tailed). Bottom panels demonstrate topography differences at cluster peaks with significant electrodes highlighted in black. Significant differences between the isoch and asynch condition OEPs in wakefulness (A) and N2 sleep (B) demonstrate that the isoch regularity produces an expectation of incoming sounds, in wakefulness and N2 sleep. Significant differences are observed between the synch and asynch OHEPs in wakefulness (C) and N2 sleep (D), suggesting that the synch cardio-audio regularity induces a modulation of OHEPs both in wakefulness and N2 sleep.

In N2 sleep, we found differences between the OHEPs of the synch and asynch conditions (Fig. 3D) at -99 ms to 117 ms (p<0.05, Cohen's d = 0.84) and at 322 ms to 500 ms (p<0.05, Cohen's d = 0.62) following R peak onset; both clusters were localized to central electrodes. As sound omissions are unexpected, the early significant difference between the synch and asynch conditions observed only during N2 sleep could be explained by changes in the background oscillatory activity during this vigilance state (addressed in 4. Slow oscillation analysis). Similar to wakefulness, no significant clusters were identified for the asynch *vs* baseline comparison in N2 sleep. However, contrary to wakefulness results, comparing the synch and baseline OHEPs in N2 sleep revealed no significant differences between the two conditions. The same statistical tests carried out in the remaining sleep stages provided no significant results.

3.3.2. Sound *vs* omission response comparisons

We additionally contrasted the synch AEPs (time-locked to the R peak) and OHEPs in wakefulness and N2 sleep in order to identify the latencies at which the neural correlates of the unfulfilled auditory prediction arise within the synch condition. In wakefulness (figure C in S7 Fig.), cluster permutation statistics revealed two significant positive clusters, a first (p<0.0005) between 125 ms and 313 ms and a second (p<0.05) between 325 ms and 443 ms,

both on posterior electrodes. Two negative clusters were additionally identified, a first (p<0.0005) between 113 ms and 246 ms and a second (p<0.05) between 314 ms and 442 ms, both on central electrodes. In N2 sleep (figure D in S7 Fig.), the synch AEPs *vs* OHEPs contrast yielded one positive cluster (p<0.0005) between 208 ms and 500 ms on central electrodes and one negative cluster (p<0.0005) between 203 ms and 500 ms on outer electrodes. Put together, the latencies of differential responses between synch AEPs and OHEPs included those observed when comparing the OHEPs between the synch and asynch conditions and between the synch and baseline conditions and are therefore coherent with our interpretation that the neural omission responses arise from the neural response to violation of cardio-audio regularity.

3.3.3. Heartbeat response comparisons

The OHEPs comparison between the synch and asynch conditions and between the synch and baseline conditions contains both the neural response to heartbeat signals and that to the sound omission. In an attempt to disentangle the contribution of these two neural responses, we built an estimation of the neural response to heartbeat in the synch condition free from the neural response to sounds. This was obtained by subtracting from the AEPs of the synch condition (which by construction includes both the neural response to cardiac and auditory signals), the AEPs of the asynch condition which is not contaminated in a systematic manner by the neural response to heartbeat. We then performed two comparisons for this clean 'synch HEPs' estimation, one to the asynch HEPs and the other to the baseline HEPs during sound presentations using cluster permutation statistical tests (p<0.05, two-tailed). In N2 sleep the 'synch HEPs' did not differ from the asynch HEPs or baseline HEPs (p>0.05). In wakefulness, we found no significant differences (p>0.05) between the synch and asynch condition HEPs. Significant differences (p<0.05) were only observed between the synch and baseline HEPs at 330 ms to 493 ms and 293 to 443 ms, as a positive and negative cluster respectively. The fact that these latencies occur later than those at which we observed differences between the OHEPs in the synch *vs* baseline condition corroborate our interpretation that the neural

omission responses (Fig. 3CD) are a result of the unfulfilled prediction upon sound omission

in the synch condition (and not a result of a modulation of the response to cardiac input).

4. Slow oscillations analysis

As outlined above (Fig. 3D), the analysis of OHEPs during N2 sleep (N=23) revealed that

cardio-audio synchronization (compared to the asynch) gave rise to a central early onset

positivity (at -99 ms to 117 ms) and late onset negativity (322 ms to 500 ms). This effect is

reminiscent of the up and down states of slow oscillations (SOs) during N2 sleep (0.5-1.2 Hz

oscillations [22,23]). This interpretation was confirmed by the analysis of the band-passed EEG

data at 0.5 Hz to 4 Hz (delta band; reflecting the range of SO frequencies) which produced a

highly similar profile for the synch *vs* asynch OHEPs during N2 sleep with an early onset central

positivity difference (p<0.05) between -99 ms and 125 ms and a late onset central negativity

difference (p<0.05) between 259 ms to 500 ms following heartbeat onset.

In light of this observation and since sound presentations are known to alter the background

oscillatory activity in sleep, notably the SOs during NREM sleep [24–26], we investigated

whether the three auditory conditions had differential effects on the ongoing oscillations during

N2 sleep. We selected the SO positive half-wave peak latency at electrode Cz (Fig. 4AB) as

the representative SO latency and computed the median sound-to-SO latency for all auditory

conditions and median R peak-to-SO latency for all auditory conditions and the baseline.

Statistical assessment using an 1x3 repeated measures Friedman test on the mean sound-to-

SO latencies yielded significant differences across auditory conditions (Fig. 4C; $\chi^2_{(1,3)}$=7.9,

p<0.05). Post-hoc Wilcoxon signed-rank tests confirmed lower latencies in synch compared to

asynch (p<0.005) and in isoch compared to asynch (p<0.05). These results suggested a

possible readjustment of the SOs with respect to sound onset depending on the regularity

condition (Fig. 4C) since when regularity was present, either in the synch or isoch condition,

SOs tended to align to the sound onset. Because of the fixed RS delay in the synch condition,

the alignment between sound onset and SOs was also reflected in a lower heartbeat to SO

peak in the synch compared to the asynch and isoch conditions (Fig. 4D). In order to rule out that this SO readjustment was due to a specific relation between R peak and SO irrespective of sound presentation, we carried out a 1x4 repeated measures Friedman test on the mean R peak-to-SO latencies which revealed significant differences across conditions (Fig. 4D; $\chi^2_{(1,4)}$=7.4, p>0.05), suggesting that potential readjustment of SOs was specific to auditory regularities and not to cardiac input.

Fig. 4. Modulation of SO activity by auditory stimulation in N2 sleep. Grand averaged (N=23) SO waveforms at electrode Cz centred around sound onset (A) and R peak (B) in the synch (red lines), asynch (black lines), isoch (blue lines), and baseline (green lines) conditions during N2 sleep. Grand average and subject-wise sound-to-SO median latency (C) and R peak-to-SO median latency (D) distributions with statistically significant differences (p<0.05; black vertical lines) across conditions. SO

peaks are more likely to occur closer to the sound onset in the synch and isoch conditions than in the asynch conditions (C). These results are not trivially driven by a temporal proximity between R peaks and SO peaks in the synch condition (D). n.s. = not significant.

5. Cardiac omission response

We further investigated whether regularity violation upon omission of expected sounds could also elicit a cardiac response across vigilance states. We analyzed heartbeat changes based on the RR intervals extracted from the ECG in response to sound omissions as a function of auditory conditions (synch, asynch, isoch).

5.1. Auditory condition contrast

We compared average RR intervals before, during and after omissions using two-way repeated measures ANOVAs with factors Auditory Condition (synch, asynch, isoch) and Trial Order (one trial before, trial during, first trial after, second trial after sound omissions) in the wakefulness and sleep sessions. To take full advantage of the available data in each vigilance state (S1 Table & S2 Table), separate repeated measures ANOVAs were performed for each vigilance state (AWAKE: N=26; N1: N=15; N2: N=24; N3: N=18; REM: N=15). These analyses were conducted on normalized RR intervals (Fig. 5; by subject-wise division of each of the investigated average RR intervals by the average RR interval prior to sound omission), so that the results would not be attributable to inter-subject variability in RR intervals. Of note, analyses on non-normalized RR intervals displayed similar results (data in S2 Text).

Overall, in wakefulness and sleep, we observed that cardio-audio synchronization led to an increase in the average RR interval during sound omissions (RR_{om}) compared to the trial before omission (RR_{-1}) that persisted over the following trial (RR_{+1}) and decreased over the second trial following omission (RR_{+2}). This deceleration in the heart rate upon sound omission was absent or inconsistent in the asynch and isoch conditions.

Fig. 5. Cardiac omission response in wakefulness and sleep. (A) Grand averaged normalized RR intervals prior (RR_{-1}), during (RR_{om}), and following (RR_{+1} and RR_{+2}) sound omissions across auditory conditions (synch: red line; asynch: black line; isoch: blue line) in the wakefulness session (AWAKE) and all sleep stages (N1, N2, N3, REM). (B) Grand averaged normalized RR intervals for participants with sufficient data across all vigilance states, prior (RR_{-1}), during (RR_{om}), and following (RR_{+1} and RR_{+2}) sound omissions in the wakefulness session (AWAKE, yellow) and all sleep stages (N1, light blue; N2,

darker blue; N3, darkest blue; REM, green) for each auditory condition. Error bars indicate the SEM (upper limit). Statistical analyses highlighted a heart rate deceleration upon sound omission in the synch condition across all vigilance states.

Statistical analysis confirmed this difference across auditory conditions as follows. 3x4 repeated measures ANOVAs revealed a significant interaction of Auditory Condition x Trial Order for all vigilance states (Fig. 5A; $p < 0.0005$; AWAKE: $F_{(3,4)} = 14.1$; N1: $F_{(3,4)} = 6.6$; N2: $F_{(3,4)} = 11.6$; N3: $F_{(3,4)} = 6.2$; REM: $F_{(3,4)} = 6.2$) with a main effect of Auditory Condition (AWAKE: $F_{(3,1)} = 27.2$, $p < 0.0005$; N1: $F_{(3,1)} = 11.4$, $p < 0.0005$; N2: $F_{(3,1)} = 14.4$, $p < 0.0005$; N3: $F_{(3,1)} = 9.0$, $p < 0.005$; REM: $F_{(3,1)} = 17.6$, $p < 0.0005$) and a main effect of Trial Order (AWAKE: $F_{(1,4)} = 20.1$, $p < 0.0005$; N1: $F_{(1,4)} = 5.3$, $p < 0.005$; N2: $F_{(1,4)} = 18.4$, $p < 0.0005$; N3: $F_{(1,4)} = 3.5$, $p < 0.05$; REM: $F_{(1,4)} = 7.5$, $p < 0.0005$). Permutation testing followed by Wilcoxon signed-rank tests evaluated the distribution of post-permutation F values against the original F values and confirmed the significance of the main effects and interactions ($p < 0.0005$). Post-hoc paired Wilcoxon signed-rank tests with Bonferroni correction for multiple comparisons across conditions corroborated that in the synch condition (Fig. 5A, red line), omissions elicited a long-lasting heart rate deceleration, with higher RR interval during and immediately after the omission than before the omission across all vigilance states (AWAKE: $p < 0.0005$; N1: $p < 0.0005$; N2: $p < 0.0005$; N3: $p < 0.005$; REM: $p < 0.005$).

5.2. Vigilance State Contrast

To better characterize how sound omissions modulated RR intervals as a function of auditory conditions and vigilance states (Fig. 5B), we computed a three-way repeated measures ANOVA with factors Vigilance State (AWAKE, N1, N2, N3, REM), Auditory Condition (synch, asynch, isoch), and Trial Order (one trial before, trial during, first trial after, second trial after sound omission), including only participants who had sufficient data in all vigilance states (N=6; S2 Table).

The ANOVA revealed a significant main effect of Auditory Condition ($F_{(1,3,1)}=27.5$, $p<0.0005$),

due to overall higher RR interval values (*i.e.* enhanced deceleration) during the synch

condition, and a main effect of Trial Order ($F_{(1,1,4)}=3.9$, $p<0.05$) but no main effect of Vigilance

State ($p>0.05$). Critically, replicating the analyses on separate vigilance states (Fig. 5A), the

interaction effect of Auditory Condition and Trial Order was also significant ($F_{(1,3,4)}=4.2$,

$p<0.0005$). Permutation testing followed by Wilcoxon signed-rank tests verified the significance

of the main effects and interaction ($p<0.0005$). The interactions Vigilance State x Auditory

Regularity, Vigilance State x Trial Order, and the triple interaction across all factors (Vigilance

State x Auditory Regularity x Trial Order) were not significant ($p>0.05$). This pattern of results

further confirmed that the observed heart rate deceleration upon sound omission in the synch

condition and lack thereof in the asynch and isoch conditions was consistent across vigilance

states.

Discussion

We investigated the neural and cardiac correlates of cardio-audio regularity processing by

administering sounds at fixed temporal pace (isoch), in synchrony to the ongoing heartbeat

(synch) and in a control condition without specific temporal regularity (asynch) while

maintaining matched average sound-to-sound intervals across conditions in wakefulness and

sleep. We tested whether auditory regularity encoding would result in a violation detection

response as measured by the cardiac and neural signals upon unexpected omitted sounds.

Neural responses to sound omissions revealed that sound sequences induced prediction of

upcoming sounds both when sounds occurred at a fixed pace and when temporally

synchronized to the ongoing heartbeat during wakefulness and N2 sleep. Enhanced prediction

related to cardio-audio synchronization was also evident in all vigilance states observed as a

strong cardiac deceleration after sound omission. Analysis of the SOs during N2 sleep

revealed a reorganization of the ongoing background brain activity both when sounds occurred

in synchrony with the ongoing heartbeat and at fixed temporal pace.

Auditory regularity and omission response in wakefulness

The neural response to sound omissions differed between the isoch and asynch conditions at 226 ms to 288 ms (Fig. 3A), a later onset compared to classic omission responses in the literature, observed at 100 ms to 250 ms post-stimulus onset ([12,15,16,21] and others). Two main factors can possibly explain this divergence. First, in our study, the sound-to-sound intervals were individually adjusted, in order to match the average heartbeat, which resulted in a variable SS interval across participants. Second, the neural omission response was derived from the difference between unexpected omissions in the isoch condition, inducing both auditory and temporal prediction ('what' and 'when'), and in the asynch condition, which can only generate an auditory prediction ('what' without 'when') [17]. In previous reports, the omission response was instead based on the comparison of expected *vs* unexpected sound omissions, revealing a violation detection response typically in the range of 100 ms and up to 250 ms post-stimulus onset [15,27]. It is likely that in our analysis, the earliest of these effects was canceled as a result of having similar 'what' prediction about incoming sounds in both isoch and asynch. That some level of auditory prediction is also elicited in the asynch sequence is also confirmed by the differential response between the OEPs in the asynch *vs* baseline (figure B in S6 Fig.).

Evidence of a violation detection upon omission of expected sounds was further confirmed by the difference between isoch and baseline at 103 ms to 170 ms and 251 ms to 295 ms post-stimulus onset (figure A in S6 Fig.). The early difference (~100ms) of central negative polarity in the isoch *vs* baseline comparison is consistent with previous reports on a sensory template formation at the predicted sound onset occurring in the same period as the auditory N100 response and traced back to the auditory cortex [15,16,21,28,29].

Cardio-audio regularity and omisssion response in wakefulness

During wakefulness, the temporal coupling between cardiac and auditory inputs in the synch condition resulted in different OHEPs (time-locked to the R peak) compared to the asynch

condition (Fig. 3C) or the baseline condition (figure C in S6 Fig.). The observed centrally-

located positive peak in the synch *vs* asynch comparison around 240 ms (Fig. 3C) closely

matches previous results using a similar paradigm, where a difference of the same polarity and

localization was observed at 260 ms [19]. When considering the average offset of 52 ms (*i.e.*

average heartbeat to sound onset latency in the present study), the latencies of observed

differences in the present study (peaking at synch *vs* asynch: 240 ms; synch *vs* baseline: 200

ms) overlap well with previously reported classic omission responses between 150 ms and

200 ms ([15,16,21] and others). Moreover, during the synch condition, the cardio-audio

synchronicity was essential for inducing the OHEP response, as demonstrated by the lack of

a differential OHEP response between the asynch and baseline conditions (time-locked to the

R peak). Overall, these results suggest that the brain utilizes the temporal cue provided by the

ongoing heartbeat to predict the onset of upcoming sounds.

Omission response in auditory and cardio-audio regularity in N2 sleep

During N2 sleep, we showed a neural response to violation of both auditory and cardio-audio

regularity upon omission of expected sounds. First, OEPs during the isoch compared to the

asynch conditions revealed a negative polarity difference at approximately 100 ms to 200 ms

(Fig. 3B). Since, to the best of our knowledge, no other studies in sleep have utilized omissions

to study violation detection, we can only refer to the deviant sound literature in interpreting

these results. Accordingly, the observed isoch *vs* asynch negative polarity is in line with the

finding of the classic MMN response in NREM sleep (*e.g.* [30–32]) and suggests an implicit

tracking of the temporal relationship between sounds despite altered vigilance in N2 sleep. In

the comparison of the OHEPs during the synch *vs* asynch condition, we found a positive and

negative cluster between -99 ms and 117 ms and 322 ms and 500 ms, respectively (Fig. 3D).

In order to corroborate the interpretation of the early OHEPs difference, we compared AEPs

and OHEPs locked to the R peak during the synch condition (figure D in S7 Fig.), which

confirmed that the only sound omission related effect was the one occurring at latencies

starting at 322 ms.

Sound administration modulates SO activity in N2 sleep

The modulation of the OHEP in N2 sleep prompted us to look into the possible influence of auditory stimulation on the background oscillatory activity. Indeed, the latencies and topography unveiled in the synch *vs* asynch N2 comparison (*i.e.* fronto-central positivity at -99 ms to 117 ms and fronto-central negativity at 322 ms to 500 ms) could reflect the up and down states of SOs during N2 sleep; keeping in mind that SO peak to trough latencies typically range between 400 ms to 1000 ms (reflecting 0.5-1.2 Hz oscillations; [22–24]). Here, this influence was demonstrated by a significantly reduced median latency between the sound stimulus onset and the peak of SOs for the synch and isoch conditions compared to the asynch condition (Fig. 4C). This indicates that sound presentations induced a modulation of slow oscillatory activity in N2 sleep in our subjects in such a way that, when auditory prediction could be generated (*i.e.* synch and isoch), sound onset was more likely to occur close to the SO peaks. This evidence is reminiscent of the closed-loop auditory stimulation literature wherein temporary synchronization of sound stimuli to ongoing SOs induced an enhancement of the SO rhythm during NREM sleep [24–26,33]. In the present paradigm as well as in closed-loop auditory stimulation studies, the temporal proximity between sound onset and the SO positive peak suggests the existence of a preferential time window of stimulus processing which may coincide with the positive phase of the SO cycle when neuronal firing is maximal [23,34]. On this basis, auditory regularity encoding would induce a reorganization of the ongoing SO activity, in order to facilitate the neural processing of expected sounds in a sequence when sound onset can be predicted (see also [35] for consistent findings in associative learning bound to the SO peaks).

Of note, not only sounds have been shown to have an impact on the latency of SOs in NREM sleep but also R peaks tend to occur close to the positive peak of the SO compared to other latencies [36,37]. With this in mind, we additionally investigated the potential impact of heartbeat signals on SOs (Fig. 4BD). In this study, we revealed no significant differences between R peak onset and SO peak across conditions. This finding suggests that SO latency modulation was specific to auditory regularities and not driven by a systematic temporal

readjustmenet of the slow oscillations by the heartbeat, which would otherwise compromise

our results in the synch condition where heartbeat and sound are temporally aligned.

Cardio-audio regularities result in heart rate deceleration upon sound omissions

In addition to studying whether and how the brain might track regularities across cardiac and

auditory inputs, we investigated the ECG responses to sound omissions as another potential

marker of auditory prediction. We found a deceleration of the heart rate during omissions

across wakefulness and all sleep stages that was most evident and consistently observed in

the synch condition (Fig. 5). One possible interpretation of the cardiac deceleration upon

omission across vigilance states is in terms of attention reorientation following an unexpected

and potentially dangerous event, a parasympathetically-driven effect often reported in

conditioning paradigms [38–40]. This response can also be associated to a startle reaction and

a physiological freezing response, linked to heart rate deceleration, pupil dilation and skin

conductance alterations [41–45]. The heart rate deceleration upon threat detection may result

from cholinergic system engagement, which modulates arousal, with similar mechanisms

observed in rats when awakened from NREM sleep by activation of basal forebrain cholinergic

neurons using chemical and optogenetic techniques [46,47].

Yet, another plausible explanation for the heart rate deceleration following unexpected

omission in the synch condition relates to top-down adjustment of cardiac rhythm in order to

account for the unexpected silence. As a sound is predicted following a heartbeat (in the

synch), the omission may prolong the generation of the next heartbeat within physiologically

plausible boundaries so as to 'wait' for a delayed auditory stimulus within the sequence,

followed by a rapid readjustment to the original rhythm upon subsequent sound presentations.

By construction, unlike the synch condition where the heartbeat deceleration was consistently

observed, such deceleration was either reduced or absent in conditions where sound

presentations bear no temporal relation to ongoing heartbeats (*i.e.* asynch and isoch). In a

previous study investigating the cardiac response to sound omission as a function of heartbeat-

to-sound onset delay and of interoceptive *vs* exteroceptive attention, a similar cardiac

deceleration was reported only in the condition of external attention [20]. While this result seems at odds with our finding of preserved cardiac deceleration across vigilance states (and potentially attentional resources), a straightforward comparison is prevented by our lack of control of the focus of attentional resources during wake, also shown to be preserved in sleep [48,49].

Cardiac but not neural omission responses observed in N1, N3 and REM sleep

Despite previous reports of preserved deviance detection in auditory oddball paradigms during N1 [4,6,31], N3 [30] and REM [4–6] sleep, in the present study, we did not identify neural omission responses to violations of the auditory and cardio-audio regularity in these sleep stages. This is possibly explained by the low number of participants providing sufficient EEG based artifact-free trials in N1, N3 and REM sleep (S1 Table & S2 Table), not the case in N2 sleep which occupies a much larger portion of the healthy sleep cycle. It therefore remains unresolved whether the lack of neural omission responses during these sleep stages resulted from a low statistical power or was due to truly impaired auditory regularity processing and deviance detection.

The analysis of the cardiac response to unexpected omissions suggests that auditory regularity processing, especially when induced by the cardio-audio coupling, is preserved across all sleep stages and would possibly be detectable at the neural level provided sufficient available data. What also emerges from this observation is a possible strength of the cardiac omission response. Detecting the heart rate deceleration with a measure as easily obtained and artifact-free as the ECG signal could offer a simple and reproducible biomarker of the preservation of sensory regularity processing in health and disease. In line with this proposal, recent work posits that the analysis of heart rate fluctuations may be informative of the degree of preservation of conscious sensory processing in healthy populations and disorders of consciousness patients [50–53]. A further investigation into the origins of the cardiac omission response through the measure of pupil dilation, respiration or changes in skin conductance will

help identify to what extent this response is a consequence of autonomic nervous system

activity.

Limitations

First, as the OHEPs are contaminated by the heartbeat related artifacts, differential neural

responses to sound omission across conditions could trivially be driven by differences between

these artifacts. Nonetheless, in our study, we could reasonably exclude this confounding factor

when interpreting the differences in the OHEPs because of the highly similar ECG signal

waveforms during the omission period across auditory conditions and both in wakefulness and

sleep (S5 Fig.). The only relevant difference was observed when comparing the baseline

average ECG waveforms to that of the synch and the asynch conditions which nevertheless

occurred at different latencies to those emerging from the comparison of the OHEPs. This

difference could be explained by the fact that in the wakefulness session, the baseline was

administered once at the start of the experiment, causing elevated heart rate and reduced

heart rate variability (S4 Fig.), until participants settled with the experimental setting.

A final possible confound to mention relates to how sensory processing is influenced by the

stimulus onset relative to the cardiac cycle [54–59], thought to rely on whether stimulus

presentation occurs within the systole or diastole period [57,60]. Although in our case, the

latencies of sound onset and periods of investigation are solely within the systole range, the

slower heart rate in sleep in comparison to wakefulness may have resulted in sounds being

presented at different timings within the cardiac cycle across vigilance states. Nonetheless,

within the range of the systole period, we expect that sensory processing would be similar [60]

both in wakefulness and sleep. In support of this, we already verified that changes of the order

of 10 ms between R peak and sound onset produce similar results in the comparison between

synch and asynch in wakefulness, upon consideration of the different average RS interval in

the synch condition for this study, M = 52 ms, and an earlier study with M = 41 ms [19].

Key contributions

The experimental paradigm implemented herein has several advantages over previous

investigations into the role of heartbeat signals in auditory regularity processing. First, the use

of omissions as opposed to deviant sounds commonly employed in sleep MMN investigations

(*e.g.* [4–6]) and interoceptive-auditory stimulus interactions [18], allows for a direct

investigation of sensory regularity encoding, free from bottom-up auditory stimulus

contributions. Second, this experimental paradigm enabled a counterbalanced ECG artifact

between the synch, asynch and baseline conditions when contrasting the OHEPs time-locked

to the R peaks. Third, by investigating both wakefulness and sleep in the same healthy

volunteers, we now demonstrate that the human brain infers on the temporal relationship

across cardiac and auditory inputs in making predictions about upcoming auditory events

across vigilance states. Finally, in the current investigation, we did not only focus on potential

neural responses to omissions but additionally identified a cardiac deceleration as a result of

violation detection. Importantly, the cardio-audio synchronicity created *ad hoc* in the

experimental environment might reflect real life readjustment of the heartbeat rhythm in order

to optimize the temporal relationship between bodily signals and exteroceptive inputs for

optimal sensory encoding.

Conclusion

Collectively, the present results suggest that the human brain can keep track of temporal

regularities between exteroceptive inputs, and across interoceptive and exteroceptive inputs

during both wakefulness and sleep. Inferring on this relationship may allow humans to form

predictions about upcoming sensory events and may aid violation detection when such

predictions are unfulfilled. Our findings replicate earlier findings in wakefulness [19] and

support theories of an interoceptive predictive coding mechanism [61–63]. To our knowledge,

this is the first study to investigate auditory regularity processing using omissions and to offer

evidence for a potential role of interoceptive inputs under the predictive coding framework in

sleep. The conscious and unconscious brain may implicitly process relationships across

interoceptive and exteroceptive inputs in order to optimize the signalling and prediction of potential upcoming dangers.

Materials and Methods

Data and code availability

All the data used in the main analyses are available for review and will be made publically available upon publication. Costum-made code used to run the quality control analyses and EEG and ECG data analyses are available on https://github.com/DNC-EEG-platform/CardioAudio_Sleep/. Any additional information required to reanalyze the data reported in this paper is available from the corresponding authors upon request.

Ethics statement

Approval for the study was obtained by the local ethics committee (La Commission Cantonale d'Ethique de la Recherche sur l'Etre Humain), in accordance with the Helsinki declaration.

Human participants

Twenty-six self-reported good sleeper volunteers took part in both the wakefulness and sleep arms of this study. Participants were considered eligible if they had no history of psychiatric, neurological, respiratory or cardiovascular conditions, no sleep apnoea, and a regular sleep schedule, evaluated during a phone interview. Hearing conditions were an additional exclusion criterion. All participants gave written informed consent and received approximately 150 Swiss Francs as monetary compensation.

Experimental design

A two-way crossover experimental design was implemented in this study. Participants attended one wakefulness and one sleep session on two occasions separated by a minimum of one day and a maximum of ten days (wakefulness session first for 12 out of 26 participants). In both sessions, participants were instructed to passively listen to the administered sound sequences. They were naïve to the experimental manipulation, as suggested by informal

verbal inquiry regarding the experimental design after the experiment. At the end of the second

session, and if desired, participants were debriefed about the purpose and design of the

experiment.

The sleep session recordings took place in a sound-attenuated hospital room equipped with a

comfortable hospital bed to allow for overnight sleep recordings. During the sleep session,

participants arrived at the laboratory at approximately 9 pm and following set-up preparation,

they were instructed to lie down and inform the experimenters when they were ready to sleep.

Lights were then switched off, the auditory stimulus administration commenced and the

volunteers were left alone to naturally fall asleep. Although participants had the liberty to leave

at any time, we explained that their inclusion in the study required a minimum of four hours of

continuous data acquisition after sleep onset. They were free to choose to spend the night at

the sleep laboratory and to be woken up at a desired time or by approximately 7 am the next

morning, at which time lights were switched on.

In both sessions, participants were equipped with electrodes for heartbeat (ECG), eye

movement (EOG), and EEG recordings (see below). For the sleep session, additional

electrodes for submental electromyography (EMG) were attached, in accordance with the 2007

AASM guidelines for sleep scoring [64]. In-ear phones (Shure SE 215, Niles, IL) were utilized

during both sessions instead of external headphones, in order to increase sound attenuation,

subject comfort during sleep and to prevent physical contact with and thus displacement of

EEG cap and electrodes. The online EEG, ECG, EOG and EMG were continuously monitored

by the experimenters to ensure effective stimulus administration, data acquisition, heartbeat

detection and sleep quality throughout both sessions.

Assessment of sleep onset was qualitatively inspected by monitoring the electrophysiological

signals in real-time to identify well-known sleep markers in accordance with Iber *et al.* [64].

Specifically, indicative of sleep onset in the EEG signals, were the dissipation of low-amplitude

alpha (8-13 Hz) posterior activity, the appearance of high-amplitude slow wave activity (1-4

Hz) and observable K complexes or sleep spindles. In addition, the reduction in eye

movements and muscle tone demonstrated that volunteers were asleep, as observed in the

EOG and submental EMG peripheral electrophysiological signals, respectively.

Stimuli

Sound stimuli were 1000 Hz sinusoidal tones of 100 ms duration (including 7 ms rise and fall

times) and 0 μs inter-aural time difference. A 10 ms linear amplitude envelope was applied at

stimulus onset and offset to avoid clicks. Stimuli were 16-bit stereo sounds sampled at

44.1 kHz and were presented binaurally with individually adjusted intensity to a comfortable

level for wakefulness. A considerably lower than wakefulness intensity of approximately 45 dB

was chosen for sleep, in order to facilitate a non-fragmented sleep session without multiple

awakenings.

Experimental procedure

During the wakefulness session, volunteers sat comfortably on a chair in a sound-attenuated

experimental room and were instructed to keep their eyes open, avoid excessive eye blinking,

body and jaw movements, and to breathe regularly; the aforementioned measures served in

ensuring high signal quality. Each participant was presented with four types of stimulation

conditions administered in separate experimental blocks in a pseudo-random order and was

asked to passively listen to the sounds while keeping the eyes fixed on a cross centrally located

in the visual field. The conditions were a baseline without auditory stimulation and three

auditory conditions, namely synch, asynch and isoch. During wakefulness, the baseline lasted

ten minutes and was acquired prior to auditory stimulation. During sleep, numerous two-minute

baseline blocks were acquired in alternation to the sound sequences in an attempt to ensure

that baseline background activity was comparable to the preceding sound stimulation during

all stages of sleep. The three auditory conditions lasted five minutes each and corresponded

to separate experimental blocks, which were repeated six times during wakefulness in a semi-

randomized order. During the sleep session, sounds were administered for the entire length of

the sleep recording in sequences of three auditory blocks always followed by a baseline (*e.g.*

isoch-synch-asynch-baseline or synch-asynch-isoch-baseline).

Auditory conditions

All auditory conditions consisted of the sequential presentation of 250 stimuli (80% sounds and

20% omissions) administered in a pseudo-random order wherein at least one sound stimulus

intervened between two subsequent omissions. Details for each auditory condition are given

below and a thorough post-hoc evaluation of the experimental manipulation is provided in the

Results and Supporting Information (Results 2. Quality control analysis & Fig. S1-4).

In the synch condition, the temporal onset of each sound stimulus was triggered by the online

detection of R peaks from raw ECG recordings. To enable effective online R peak detection,

raw ECG recordings were analyzed in real-time using a custom MATLAB Simulink script

(R2019b, The MathWorks, Natick, MA). The variance over the preceding 50 ms time window

was computed and an R peak was detected when the online ECG value exceeded an

individually adjusted 10–15 mV^2 variance threshold, which in turn triggered the presentation of

a sound stimulus or an omission. This procedure resulted in a fixed R peak-to-sound average

delay (*i.e.* RS interval) of 52 ms (SD = 5 ms) for wakefulness and sleep across participants.

In the asynch condition, the onset of sound presentation was based on the RR intervals

extracted from a previously acquired synch block. Specifically, the ECG recorded during the

preceding synch block was analyzed offline to extract RR intervals by automatic detection of

R peaks and computation of RR intervals. 250 RR intervals were selected if they were above

the 25[th] and below the 75[th] percentile of RR interval distribution in the synch block, in order to

take into account possible missed R peaks in the online detection during the synch block. Next,

RR interval order was shuffled giving rise to a predefined pseudo-random sequence closely

resembling the participant's heartbeat rhythm. By construction, differences between the synch

and asynch conditions in terms of average and variance of the RR intervals were minimized,

contrary to the RS interval being fixed in the synch condition and variable in the asynch

condition. The variable sound to heartbeat relationship resulted in an average RS interval of -3 ms (SD = 25 ms, range = −206 to 96 ms), as well as an average SR interval of 848 ms (SD = 136 ms, range = 504-1259 ms) for wakefulness and sleep.

In the isoch condition, the onset of sound presentations was based on the median RR interval calculated during a previously acquired synch block. This procedure produced similar sound-to-sound intervals across the synch, asynch and isoch conditions. However, unlike the synch and asynch conditions, sound-to-sound intervals in the isoch condition were of a low variability (*i.e.* SS variability: SEM of SS interval) of 1 ms for wakefulness and 3 ms for sleep across participants. Similar to asynch, in the isoch condition, the variable sound to heartbeat relationship resulted in an average RS interval of -1 ms (SD = 32 ms, range = −116 to 165 ms) in addition to an average SR interval of 853 ms (SD = 141 ms, range = 516-1263 ms) for wakefulness and sleep.

Data acquisition

Continuous EEG (g.HIamp, g.tec medical engineering, Graz, Austria) was acquired at 1200 Hz from 63 active ring electrodes (g.LADYbird, g.tec medical engineering) arranged according to the international 10–10 system and referenced to the right ear lobe. Biophysical data were acquired using single-use Ag/AgCl electrodes. Three-lead ECG was recorded by attaching two electrodes (a third was a reference) to the participant's chest on the infraclavicular fossae and above the heart. A vertical EOG electrode was attached below the right eye and a horizontal EOG electrode was attached to the outer right canthus. Since changes in muscle tone are associated with alterations in consciousness during sleep and are an essential marker for effective sleep staging [64], EMG was additionally acquired sub-mentally during the sleep session alone. Impedances of all active electrodes were kept below 50 kΩ. All electrophysiological data were acquired with an online band-pass filter between 0.1 and 140 Hz and a band-stop filter between 48 and 52 Hz to reduce electrical line noise.

Sleep scoring

An experienced sleep scoring specialist (Somnox SRL, Belgium), blind to the experimental

manipulation in this study, performed the scoring of the continuous sleep electrophysiological

data in order to pinpoint the periods of wakefulness and micro-arousal as well as periods of

N1, N2, N3 and REM sleep. Sleep scoring was performed via visual inspection of contiguous

30-second segments of the EEG, EOG and EMG time-series, as outlined in the 2007 AASM

guidelines for sleep scoring [64]. Segments scored as periods of wakefulness or micro-

arousals in the sleep recordings were excluded from further analysis.

Data analysis

Electrophysiological data analyses were performed in MATLAB (R2019b, The MathWorks,

Natick, MA) using open-source toolboxes EEGLAB (version 13.4.4b, [65]), Fieldtrip (version

20201205, [66]), as well as using custom-made scripts. Raincloud plots were generated using

the Raincloud plot toolbox [67].

R peak detection

The R peaks in the continuous raw ECG signal were selected offline using a semi-automated

approach as in Pfeiffer & De Lucia [19]. The custom-made MATLAB script, *rpeakdetect.m*

(https://ch.mathworks.com/matlabcentral/fileexchange/72-peakdetect-

m/content/peakdetect.m), was utilized to automatically identify the sharp R peaks in the raw

ECG signal. Visual inspection of the online and offline detected R peaks ensured that the

selected peaks fitted within the expected structure of the QRS complex in the continuous raw

ECG signal. Frequent flawed online identification of the R peaks or faulty auditory stimulus

presentation in a given block resulted in the exclusion of the block from a given participant's

dataset. For blocks that were included, unrealistic RR, RS and SR interval values, observed

as a result of infrequent flawed offline marking of R peaks, were identified and excluded using

the *rmoutliers* MATLAB function (R2019b, The MathWorks, Natick, MA) and visual inspection

of the detected R peaks and selected outliers.

Quality control analysis

A series of control analyses were performed to investigate whether the experimental

manipulation was producing the expected RS, RR and SS mean and variances, and the

presence of possible confounding factors. RR and SS intervals (for sound trials preceded by

sound trials) were extracted for the synch, asynch, isoch and baseline conditions where

relevant. In addition, RS intervals for sound trials and SR intervals for omission trials were

computed to quantify the degree of cardio-audio synchronization and heartbeat onset

variability during sound omission, respectively. Variability in the same interval measures,

computed as the SEM was additionally investigated. Of note, unlike RR, SR and RS variability,

SS variability was first computed within a given experimental block and then across

experimental blocks, to account for ongoing changes in RR intervals (and hence SS intervals)

in sleep. Non-parametric one way repeated measures Friedman tests ($p<0.05$) were performed

on the average RR, SS, RS and SR intervals and variabilities of each participant for

wakefulness and all sleep stages with within-subject factor Condition (3 levels for SS, RS, SR

intervals and variability: synch, asynch, isoch; 4 levels for RR intervals and variability: synch,

asynch, isoch, baseline). Post-hoc paired Wilcoxon signed-rank tests ($p<0.05$) identified any

significant pairwise comparisons (no multiple comparisons correction was applied since

pairwise differences were of interest).

To ensure no significant differences in the ECG signal across experimental conditions, ECG

waveforms during omissions were extracted from ECG recordings between -100 ms and 500

ms relative to R peak onset, matching the EEG trial-based analysis (see below, EEG Data

Analysis). The non-parametric cluster-based permutation statistical analysis approach [68]

was employed to investigate ECG waveform differences between the various experimental

conditions outlined herein. In order to reject the null hypothesis that no significant differences

existed in the given set of experimental conditions being contrasted, maximum cluster-level

statistics were determined by shuffling condition labels (5000 permutations), allowing for a

chance-based distribution of maximal cluster-level statistics to be estimated. Since maxima

are utilized by this method, it enables the correction of multiple comparisons over time. A two-

tailed Monte-Carlo p-value allowed for the definition of a threshold of significance from the

distribution (p<0.05, two-tailed).

EEG data analysis

Continuous raw EEG data were band-pass filtered using second-order Butterworth filters

between 0.1 and 40 Hz for the wakefulness session and 0.5 and 30 Hz for the sleep session.

A larger high pass cut-off was chosen for sleep electrical signals in order to exclude periodic

breathing and sweating artifacts which may occur in that frequency range in sleep and a lower

low-pass cut-off was employed in agreement with the sleep literature (*e.g.* [24]). To look

specifically at slow oscillatory activity, raw electrophysiological data were band-pass filtered

between 0.5 and 4 Hz, a well-established frequency range for investigations of slow-wave

activity in the delta band range [69].

Ocular and muscular activity in the continuous EEG was identified for wakefulness data by

performing Independent Component Analysis (ICA) as implemented in Fieldtrip [65,66] and

any selected eye movement or muscle activity related components were removed. We

refrained from applying the ICA algorithm to the continuous sleep EEG data since changes in

periodic eye movements and muscle activity form an integral part of the human sleep cycle.

Therefore, contrary to wakefulness, excluding ocular and muscular contributions to the

acquired EEG signal may have resulted in loss of relevant activity in sleep electrophysiology.

Wakefulness and sleep continuous EEG data were epoched between -100 ms and 500 ms

relative to the selected event onset. This range was chosen in reflection of previously reported

classic omission response latencies ([15,16,21] and others) and importantly, to ensure no

overlap between consecutive auditory stimulation trials (as in [19], assuming a range of 60-

100 beats per minute across volunteers). Various event onsets were selected, namely,

heartbeat onset, sound onset and omission onset, depending on the comparison performed

(Results 3. Neural omission response). Of note, while the isoch condition had fixed SS

intervals, this was not the case in the asynch condition. To test whether sound prediction was

based on computing the time of maximal sound occurrence probability, average sound-based

omission responses were calculated by extracting epochs from the continuous EEG asynch

and isoch recordings, now time-locked to the latency occurring during the omission at the

average SS interval.

The resting-state baseline recordings acquired as part of the experimental procedure (without

auditory stimulation) were utilized to extract EEG evoked potentials with matched event onsets

to auditory stimulation trials and epoched between -100 ms and 500 ms. First, we extracted

HEPs based on heartbeat onset in the resting state for upcoming statistical comparisons to the

synch and asynch OHEP. Second, in the absence of sound stimulation, the baseline allowed

for comparing the OEPs of the isoch and asynch conditions to a control condition. In this case,

a random selection of epochs was extracted from continuous baseline recordings, such that

the latencies between epoch onset and closest heartbeat (*i.e.* R peak) were matched on the

single-trial level to the trial onsets in the sound-based isoch and asynch conditions. Upcoming

pre-processing steps were replicated for auditory stimulation and baseline trials, for each

participant, and for wakefulness and all sleep stages as follows.

Artifact electrodes and trials were identified using a semi-automated approach for artifact

rejection as implemented in Fieldtrip [66]. Noisy EEG electrodes were excluded based on a

signal variance criterion (3 z-score Hurst exponent) and substituted with data interpolated from

nearby channels using spherical splines [70]. Across eligible participants, an average of 7.3

(SD = 2.2) electrodes (M = 11.7%, SD = 3.6%) were interpolated in wakefulness and an

average of 8.9 (SD = 1.1) electrodes (M = 14.4% SD = 1.7%) were interpolated in sleep.

Extended recording lengths, participant sweating, and head and body movement during the

night could account for the greater number of artifactual electrodes observed in sleep

recordings. In wakefulness data, trials containing physiological artifacts (*e.g.* eye movement,

excessive muscle activity), not accounted for by ICA, were identified by visual inspection and

by using a 70 µV absolute value cut-off applied to the EEG signal amplitude and were excluded

from further analysis. A higher absolute value threshold of 300 µV was employed for the

selection of artifactual epochs in sleep, in order to prevent the exclusion of high-amplitude slow

wave activity [71]. Extreme outliers in signal kurtosis and variance were additional criteria for

the rejection of artifactual trials from sleep recordings. Finally, common average re-referencing

was applied.

The 30-second sleep stage labeled epochs were used to label artifact-free trials for all event

onset types and for all three auditory conditions and the baseline. Hence, five different sets of

trials formed the final processed dataset per participant for each of the five vigilance states:

AWAKE, N1, N2, N3 and REM sleep where available (S1 Table & S2 Table). We note that we

chose not to employ pre-stimulus baseline correction in wakefulness or sleep trials. In addition,

since one way ANOVAs on artifact-free trial numbers for each evoked response of interest

confirmed that no significant differences ($p > 0.05$) existed across auditory conditions alone, all

trials were kept in auditory condition comparisons. Conversely, since acquired baseline data

were significantly less ($p < 0.05$) than the three auditory conditions, baseline trial numbers were

quantitatively matched for each participant and for each vigilance state between auditory

conditions and baselines. The number of available trials for each vigilance state, averaged

across experimental conditions and event onsets of interest were AWAKE: M = 245, SD = 26

trials; N1: M = 97, SD = 22 trials; N2: M = 428, SD = 184 trials; N3: M = 159, SD = 72 trials;

REM: M = 163, SD = 65 trials. Finally, grand average evoked responses to sound omissions

were derived.

EEG statistical analysis

EEG statistical analyses were performed in MATLAB (R2019b, The MathWorks, Natick, MA)

using Fieldtrip (version 20201205, [66]), as well as custom made scripts. For the analysis of

the electrophysiological signals during the wakefulness and sleep sessions, we imposed a

minimum of 60 artifact-free trials for the EEG data analysis, chosen based on the signal-to-

noise ratio required to meaningfully interpret EEG statistical analysis results (Luck, 2005).

The non-parametric cluster-based permutation statistical analysis approach [72] was

employed to investigate sensor-level EEG-based differences between the various

experimental conditions outlined herein. Under this statistical framework, statistically

significant individual data samples were grouped based on the degree of their shared spatial

and temporal characteristics. The resulting clusters were statistically evaluated by summating

the t-values for all samples forming up a given cluster. In order to reject the null hypothesis

that no significant differences existed in the given set of experimental conditions being

contrasted, maximum cluster-level statistics were determined by shuffling condition labels

(5000 permutations), allowing for a chance-based distribution of maximal cluster-level statistics

to be estimated. A two-tailed Monte-Carlo p-value allowed for the definition of a threshold of

significance from the distribution ($p < 0.05$, two-tailed). In wakefulness, this procedure was

performed from 0 ms to 350 ms relative to the event onset of interest (heartbeat onset for

OHEP comparisons and expected sound onset based on average SS interval for OEP

comparisons). The aforementioned time window was chosen in reflection of a previous

publication from our laboratory [19], suggesting that changes of interest occurred within the

selected period. Conversely, in sleep and in the absence of prior knowledge specific to this

paradigm, we chose to apply cluster-based permutation statistics over the entire epoch length

from -100 ms to 500 ms relative to event onset. Finally, in order to evaluate the size of the

observed effects, the Cohen's d statistic was calculated at the latenies of the largest significant

clusters [73].

SO data analysis

To examine whether N2 sleep EEG statistical analysis results could have been influenced by

the relationship between sounds and SO activity [24,26,33] and/or between heartbeats and

SO activity [36] in NREM sleep, we identified SOs in N2 sleep artifact-free EEG data. We then

computed the latency at which the SOs occurred compared first, to sound presentations in the

auditory conditions and second, to heartbeats in the auditory conditions and baseline.

SOs in the continuous EEG time-series were detected across experimental conditions over frontocentral electrode Cz where a high probability for SO detection is to be expected [22]. We marked SOs based on the method described in Ngo *et al.* [24] and in Besedovsky *et al.* [33] In brief, the 0.5 to 30 Hz band-pass filtered data were downsampled from 1200 Hz to 100 Hz and a low pass finite impulse response filter of 3.5 Hz was used in order to improve the detection of SO components. Next, N2 sleep labeled segments were extracted for the identification of SOs. Consecutive positive-to-negative zero crossings were picked out and were selected only if their temporal distance was between 0.833 s and 2 s, yielding a frequency between 0.5 Hz and 1.2 Hz for the designated oscillations. Negative and positive peak potentials in the oscillations were defined as the minima and maxima present within eligible consecutive positive-to-negative zero crossings. The mean negative and mean positive-to-negative amplitude differences were calculated across selected oscillations at electrode Cz. For each identified oscillation that satisfied the frequency criterion, if the negative amplitude was 1.25 times lower than the mean negative amplitude and the positive to negative amplitude difference was 1.25 times higher than the mean positive to negative amplitude difference, the oscillation was marked as a SO.

The positive half-wave peak time point was chosen as the representative latency for each SO in light of relevant literature [23,34] and following visual inspection of the SO and sound presentation time-series. Sound to SO latency for the synch, asynch and isoch conditions and R peak to SO latency for the synch, asynch, isoch and baseline conditions were computed for all artifact-free sound trials. Omission trials were excluded from this analysis since in this case, we were interested in how sounds and the sound to heartbeat relationship may modulate SOs in N2 sleep. Latencies were considered as valid only if they were between -800 and 800ms, a range chosen to minimize potential contamination of the sound or R peak to SO relationship by upcoming sounds or heartbeats. Median latencies were calculated for each subject and condition separately.

SO statistical analysis

In N2 sleep, a non-parametric 1x3 repeated measures Friedman test was calculated on the

median sound to SO latencies at electrode Cz with within-subject factor Auditory Condition

(synch, asynch, isoch) and a 1x4 repeated measures Friedman test were computed on the

median heartbeat to SO latencies with within-subject factor Condition (synch, asynch, isoch,

baseline). Pairwise comparisons were calculated using post-hoc paired Wilcoxon signed-rank

tests between all investigated within-subject variables (no multiple comparisons correction was

applied since pairwise differences were of interest). Latency trial numbers were matched for

the median heartbeat to SO latency comparison but not for the sound to SO latency

comparison, since in the former, baseline recordings lengths were significantly lower to

auditory condition recordings (as confirmed by 1x4 and 1x3 repeated measures ANOVAs on

latency trial numbers, which significantly differed only across heartbeat to SO trials and not

sound to SO trials).

ECG data analysis

The identified R peaks were used to derive omission trial related RR intervals. The

representative R peak for an omission trial was selected as the first R peak following an

omission in the continuous ECG signal. The omission RR Interval (RR_{om}) was therefore

calculated as the latency between the selected omission R peak and the R peak immediately

preceding it. In order to investigate potential heartbeat alterations associated with the omission

trial, additional RR intervals were identified using contiguous R peaks to reflect the RR interval

for the trial prior to omission (RR_{-1}) and up to two trials following omission (RR_{+1} and RR_{+2}).

Omission trials were considered only if they were followed by at least two sound stimuli to

ensure no overlap between investigated RR intervals.

The 30-second sleep stage labeled epochs were used to label omission trials for all auditory

conditions. For the analysis of the ECG signals during the wakefulness and sleep sessions,

we imposed a minimum of 30 artifact-free trials for the ECG data analysis and for each

condition of interest. Hence, different sets of RR intervals formed the final dataset per

participant and consisted of five vigilance states: AWAKE, N1, N2, N3 and REM sleep where

available (S1 Table & S2 Table). One way ANOVAs on RR interval quantities showed no

significant differences ($p>0.05$) in trial numbers across auditory conditions within each

vigilance state, therefore all RR intervals were included in the statistical analysis. RR intervals

for each omission trial order type were averaged for each auditory condition, participant, and

each vigilance state.

ECG statistical analysis

First, 3x4 repeated measures ANOVAs were performed on the average RR intervals with

within-subject factors Auditory Condition (synch, asynch, isoch) and Trial Order (one trial

before, trial during, first trial after, second trial after sound omissions) for the wakefulness

session (AWAKE) and each of the sleep stages (N1, N2, N3, REM) separately. Second, for a

sub-set of participants (N=6) who had sufficient trials across wakefulness and all sleep stages,

a 5x3x4 repeated measures ANOVA was performed on the average RR intervals with within-

subject factors Vigilance State (AWAKE, N1, N2, N3, REM), Auditory Condition (synch,

asynch, isoch) and Trial Order (one trial before, trial during, first trial after, second trial after

sound omission). Of note, unlike comparisons within vigilance states, trial numbers were

matched across vigilance states by randomly selecting a subset of RR intervals based on the

minimum number ($\geqslant30$ trials) of available RR intervals across participants to ensure accurate

comparison across vigilance states. In order to reject the null hypothesis that no significant

differences existed in the given set of ANOVA factors being compared, each of the factors'

condition labels were shuffled (2000 permutations), allowing for an estimation of the chance-

based distribution of maximal cluster-level statistics to be estimated. Wilcoxon signed-ranked

tests ($p<0.05$) were used to evaluate whether significant differences existed between the

distribution of post-permutation F values and the true main effect and interaction effect F

values. Post-hoc Wilcoxon signed-rank tests with Bonferroni correction for multiple

comparisons ($p<0.018$ for auditory condition and $p<0.013$ for omission trial order comparisons)

were utilized for pairwise comparisons between all investigated within-subject variables.

 Table 2. Hypotheses and tests performed for the main analyses of the study.

Analysis	Hypothesis	Description	Test
EEG neural heartbeat-based omission response	**H1**	**Omission response time-locked to heartbeats during synch in wakefulness and N2 sleep.**	Non-parametric cluster-based permutation statistics comparing synch *vs* asynch, synch *vs* baseline and asynch *vs* baseline for trials time-locked to heartbeat onset for wakefulness (AWAKE) and all stages of sleep (N1, N2, N3, REM) separately.
	H0	No omission responses time-locked to heartbeats in all vigilance states.	
EEG neural sound-based omission response	**H1**	**Omission response time-locked to expected sound onset during isoch in wakefulness and N2 sleep.**	Non-parametric cluster-based permutation statistics comparing isoch *vs* asynch, isoch *vs* baseline and asynch *vs* baseline for trials time-locked to omission onset for wakefulness (AWAKE) and all stages of sleep (N1, N2, N3, REM) separately.
	H0	No omission response time-locked to expected sound onset in all vigilance states.	
EEG SOs	**H1**	**Sound to SO and heartbeat to SO latency differences in synch and Sound to SO latency differences in isoch compared no regularity conditions (asynch, baseline) in N2 sleep.**	Non-parametric one way repeated measures Friedman tests on the median sound to SO and median heartbeat to SO latency with factor Condition (synch, asynch, isoch where baseline where relevant).
	H0	No sound to SO and/or heartbeat to SO latency	

		differences across auditory conditions in N2 sleep.	
ECG cardiac omission response	**H1**	**RR intervals in relation to omission onset significantly different across experimental conditions in wakefulness and all stages of sleep.**	3x4 repeated measures ANOVAs on the average RR intervals with factors Auditory Condition (synch, asynch, isoch) and Trial Order (one trial before, trial during, first trial after, second trial after sound omission) for each vigilance state separately (AWAKE, N1, N2, N3, REM).
	H0	No significant differences across experimental conditions independent of vigilance state.	5x3x4 repeated measures ANOVA on the average RR intervals with factors Vigilance State (AWAKE, N1, N2, N3, REM), Auditory Condition (synch, asynch, isoch) and Trial Order (one trial before, trial during, first trial after, second trial after sound omission).

Acknowledgments

We express our gratitude to Nathalie Nguepnjo Nguissi, Elsa Dosi and Diana Ortolani for assistance during data acquisition. We extend our thanks to Dr Aurore Guyon Postalci from SOMNOX for performing the stage scoring of the sleep data. We thank Rupert Ortner and the g.tec team for technical support.

Funding Sources

1007 This work was supported by a Spark SNSF Grant (no. 196194) awarded to M.D.L.

Declaration of interests

1009 The authors declare no competing interests.

Author contributions

1011 A.P., C.P., S.S. and M.D.L. designed the experiment and conceived the analysis. A.P.

1012 conducted the experiment and acquired the data. A.P. and M.D.L. analyzed the data and wrote

1013 the manuscript. C.P. and S.S. edited the manuscript.

References

1.   Tivadar RI, Knight RT, Tzovara A. Automatic Sensory Predictions: A Review of Predictive Mechanisms in
the Brain and Their Link to Conscious Processing. Frontiers in Human Neuroscience. Frontiers Media
S.A.; 2021. doi:10.3389/fnhum.2021.702520

2.   Garrido MI, Kilner JM, Stephan KE, Friston KJ. The mismatch negativity: a review of underlying
mechanisms. Clin Neurophysiol. 2009;120: 453–463. Available:
http://www.zora.uzh.chhttp//www.zora.uzh.chhttp://www.zora.uzh.ch

3.   Näätänen R, Paavilainen P, Rinne T, Alho K. The mismatch negativity (MMN) in basic research of central
auditory processing: A review. Clinical Neurophysiology. 2007. pp. 2544–2590.
doi:10.1016/j.clinph.2007.04.026

4.   Ruby P, Caclin A, Boulet S, Delpuech C, Morlet D. Odd Sound Processing in the Sleeping Brain. J Cogn
Neurosci. 2008;20: 296–311. Available: http://direct.mit.edu/jocn/article-
pdf/20/2/296/1936782/jocn.2008.20023.pdf

5.   Sculthorpe LD, Ouellet DR, Campbell KB. MMN elicitation during natural sleep to violations of an auditory
pattern. Brain Res. 2009;1290: 52–62. doi:10.1016/j.brainres.2009.06.013

6.   Strauss M, Sitt JD, King JR, Elbaz M, Azizi L, Buiatti M, et al. Disruption of hierarchical predictive coding
during sleep. Proc Natl Acad Sci U S A. 2015;112: E1353–E1362. doi:10.1073/pnas.1501026112

7.   Tzovara A, Simonin A, Oddo M, Rossetti AO, De Lucia M. Neural detection of complex sound sequences
in the absence of consciousness. Brain. 2015;138: 1160–1166. doi:10.1093/brain/awv041

8.   Perez P, Valente M, Hermann B, Sitt J, Faugeras F, Demeret S, et al. Auditory Event-Related "Global

Effect" Predicts Recovery of Overt Consciousness. Front Neurol. 2021;11: 1–10.

doi:10.3389/fneur.2020.588233

9.   Tzovara A, Rossetti AO, Spierer L, Grivel J, Murray MM, Oddo M, et al. Progression of auditory

discrimination based on neural decoding predicts awakening from coma. Brain. 2013;136: 81–9.

doi:10.1093/brain/aws264

10.   Heilbron M, Chait M. Great Expectations: Is there Evidence for Predictive Coding in Auditory Cortex?

Neuroscience. Elsevier Ltd; 2018. pp. 54–73. doi:10.1016/j.neuroscience.2017.07.061

11.   Barascud N, Pearce MT, Griffiths TD, Friston KJ, Chait M. Brain responses in humans reveal ideal

observer-like sensitivity to complex acoustic patterns. Proc Natl Acad Sci U S A. 2016;113: E616-25.

doi:10.1073/pnas.1508523113

12.   Chennu S, Noreika V, Gueorguiev D, Shtyrov Y, Bekinschtein TA, Henson R. Silent expectations:

Dynamic causal modeling of cortical prediction and attention to sounds that weren't. J Neurosci. 2016;36:

8305–8316. doi:10.1523/JNEUROSCI.1125-16.2016

13.   Demarchi G, Sanchez G, Weisz N. Automatic and feature-specific prediction-related neural activity in the

human auditory system. Nat Commun. 2019;10: 3440. doi:10.1038/s41467-019-11440-1

14.   Recasens M, Uhlhaas PJ. Test–retest reliability of the magnetic mismatch negativity response to sound

duration and omission deviants. Neuroimage. 2017;157: 184–195. doi:10.1016/j.neuroimage.2017.05.064

15.   SanMiguel I, Widmann A, Bendixen A, Trujillo-Barreto N, Schröger E. Hearing silences: Human auditory

processing relies on preactivation of sound-specific brain activity patterns. J Neurosci. 2013;33: 8633–

8639. doi:10.1523/JNEUROSCI.5821-12.2013

16.   Wacongne C, Labyt E, Van Wassenhove V, Bekinschtein T, Naccache L, Dehaene S. Evidence for a

hierarchy of predictions and prediction errors in human cortex. Proc Natl Acad Sci U S A. 2011;108:

20754–20759. doi:10.1073/pnas.1117807108

17.   SanMiguel I, Saupe K, Schröger E. I know what is missing here: Electrophysiological prediction error

signals elicited by omissions of predicted "what" but not "when." Front Hum Neurosci. 2013;7: 407.

doi:10.3389/fnhum.2013.00407

18.   Van Elk M, Lenggenhager B, Heydrich L, Blanke O. Suppression of the auditory N1-component for

heartbeat-related sounds reflects interoceptive predictive coding. Biol Psychol. 2014;99: 172–182.

doi:10.1016/j.biopsycho.2014.03.004

19.   Pfeiffer C, De Lucia M. Cardio-audio synchronization drives neural surprise response. Sci Rep. 2017;7:

14842. doi:10.1038/s41598-017-13861-8

20.   Banellis L, Cruse D. Skipping a Beat: Heartbeat-Evoked Potentials Reflect Predictions during

Interoceptive-Exteroceptive Integration. Cereb Cortex Commun. 2020;1: 1–14.

doi:10.1093/texcom/tgaa060

21.   van Laarhoven T, Stekelenburg JJ, Vroomen J. Temporal and identity prediction in visual-auditory events:

Electrophysiological evidence from stimulus omissions. Brain Res. 2017;1661: 79–87.

doi:10.1016/j.brainres.2017.02.014

22.   Massimini M, Huber R, Ferrarelli F, Hill S, Tononi G. The sleep slow oscillation as a traveling wave. J

Neurosci. 2004;24: 6862–6870. doi:10.1523/JNEUROSCI.1318-04.2004

23.   Nir Y, Staba RJ, Andrillon T, Vyazovskiy V V., Cirelli C, Fried I, et al. Regional Slow Waves and Spindles

in Human Sleep. Neuron. 2011;70: 153–169. doi:10.1016/j.neuron.2011.02.043

24.   Ngo HV V., Martinetz T, Born J, Mölle M. Auditory closed-loop stimulation of the sleep slow oscillation

enhances memory. Neuron. 2013;78: 545–553. doi:10.1016/j.neuron.2013.03.006

25.   Ong JL, Patanaik A, Chee NIYN, Lee XK, Poh JH, Chee MWL. Auditory stimulation of sleep slow

oscillations modulates subsequent memory encoding through altered hippocampal function. Sleep.

2018;41: 5. doi:10.1093/sleep/zsy031

26.   Schneider J, Lewis PA, Koester D, Born J, Ngo HV V. Susceptibility to auditory closed-loop stimulation of

sleep slow oscillations changes with age. Sleep. 2020;43: 12. doi:10.1093/sleep/zsaa111

27.   Wacongne C, Changeux JP, Dehaene S. A neuronal model of predictive coding accounting for the

mismatch negativity. J Neurosci. 2012;32: 3665–3678. doi:10.1523/JNEUROSCI.5003-11.2012

28.   van Laarhoven T, Stekelenburg JJ, Eussen MLJM, Vroomen J. Atypical visual-auditory predictive coding

in autism spectrum disorder: Electrophysiological evidence from stimulus omissions. Autism. 2020;24:

1849–1859. doi:10.1177/1362361320926061

29.   Dercksen TT, Widmann A, Schröger E, Wetzel N. Omission related brain responses reflect specific and

unspecific action-effect couplings. Neuroimage. 2020;215: 11684. doi:10.1016/j.neuroimage.2020.116840

30.   Sabri M, Labelle S, Gosselin A, Campbell KB. The effects of digital filtering on mismatch negativity in

wakefulness and slow-wave sleep. J Sleep Res. 2002;11: 123–127.

31.   Sabri M, Labelle S, Gosselin A, Campbell KB. Effects of sleep onset on the mismatch negativity (MMN) to

frequency deviants using a rapid rate of presentation. Cogn Brain Res. 2003;17: 164–176.

doi:10.1016/S0926-6410(03)00090-9

32.   Winter O, Kok A, Kenemans L, Elton M. Auditory event-related potentials to deviant stimuli during

drowsiness and stage 2 sleep. Electroencephalogr Clin Neurophysiol. 1995;96: 398–412.

33.   Besedovsky L, Ngo HV V., Dimitrov S, Gassenmaier C, Lehmann R, Born J. Auditory closed-loop

stimulation of EEG slow oscillations strengthens sleep and signs of its immune-supportive function. Nat

Commun. 2017;8. doi:10.1038/s41467-017-02170-3

34.   Navarrete M, Schneider J, Ngo HV V., Valderrama M, Casson AJ, Lewis PA. Examining the optimal

timing for closed-loop auditory stimulation of slow-wave sleep in young and older adults. Sleep. 2020;43:

6: 1–14. doi:10.1093/sleep/zsz315

35.   Züst MA, Ruch S, Wiest R, Henke K. Implicit Vocabulary Learning during Sleep Is Bound to Slow-Wave

Peaks. Curr Biol. 2019;29: 541-553.e7. doi:10.1016/j.cub.2018.12.038

36.   Lechinger J, Heib DPJ, Gruber W, Schabus M, Klimesch W. Heartbeat-related EEG amplitude and phase

modulations from wakefulness to deep sleep: Interactions with sleep spindles and slow oscillations.

Psychophysiology. 2015;52: 1441–1450. doi:10.1111/psyp.12508

37.   Mensen A, Zhang Z, Qi M, Khatami R. The occurrence of individual slow waves in sleep is predicted by

heart rate. Sci Rep. 2016;6: 6–13. doi:10.1038/srep29671

38.   Pappens M, Schroijen M, Sütterlin S, Smets E, Van Den Bergh O, Thayer JF, et al. Resting heart rate

variability predicts safety learning and fear extinction in an interoceptive fear conditioning paradigm. PLoS

One. 2014;9. doi:10.1371/journal.pone.0105054

39.   Seligowski A V., Reffi AN, Phillips KA, Orcutt HK, Auerbach RP, Pizzagalli DA, et al. Neurophysiological

responses to safety signals and the role of cardiac vagal control. Behav Brain Res. 2021;396: 112914.

doi:10.1016/j.bbr.2020.112914

40.   Szeska C, Richter J, Wendt J, Weymar M, Hamm AO. Attentive immobility in the face of inevitable distal

threat—Startle potentiation and fear bradycardia as an index of emotion and attention. Psychophysiology.

2021;58: 1–17. doi:10.1111/psyp.13812

41.   Blanchard DC, Griebel G, Pobbe R, Blanchard RJ. Risk assessment as an evolved threat detection and

analysis process. Neurosci Biobehav Rev. 2011;35: 991–998. doi:10.1016/j.neubiorev.2010.10.016

42.   Gladwin TE, Hashemi MM, van Ast V, Roelofs K. Ready and waiting: Freezing as active action

preparation under threat. Neurosci Lett. 2016;619: 182–188. doi:10.1016/j.neulet.2016.03.027

43.   Jennings JR, Van Der Molen MW. Preparation for speeded action as a psychophysiological concept.

1123   Psychol Bull. 2005;131: 434–459. doi:10.1037/0033-2909.131.3.434

1124   44.   Lojowska M, Gladwin TE, Roelofs K. Supplemental Material for Freezing Promotes Perception of Coarse

1125   Visual Features. J Exp Psychol Gen. 2015;144: 1080–1088. doi:10.1037/xge0000117.supp

1126   45.   Roelofs K. Freeze for action: Neurobiological mechanisms in animal and human freezing. Philos Trans R

1127   Soc B Biol Sci. 2017;372: 20160. doi:10.1098/rstb.2016.0206

1128   46.   McKenna JT, Thankachan S, Uygun DS, Shukla C, McNally JM, Schiffino FL, et al. Basal Forebrain

1129   Parvalbumin Neurons Mediate Arousals from Sleep Induced by Hypercarbia or Auditory Stimuli. Curr Biol.

1130   2020;30: 2379-2385.e4. doi:10.1016/j.cub.2020.04.029

1131   47.   Zant JC, Kim T, Prokai L, Szarka S, McNally J, McKenna JT, et al. Cholinergic neurons in the basal

1132   forebrain promote wakefulness by actions on neighboring non-cholinergic neurons: An opto-dialysis

1133   study. J Neurosci. 2016;36: 2057–2067. doi:10.1523/JNEUROSCI.3318-15.2016

1134   48.   Kouider S, Andrillon T, Barbosa LS, Goupil L, Bekinschtein TA. Inducing task-relevant responses to

1135   speech in the sleeping brain. Curr Biol. 2014;24: 2208–2214. doi:10.1016/j.cub.2014.08.016

1136   49.   Legendre G, Andrillon T, Koroma M, Kouider S. Sleepers track informative speech in a multitalker

1137   environment. Nat Hum Behav. 2019;3: 274–283. doi:10.1038/s41562-018-0502-5

1138   50.   Endoh H, Kamimura N, Honda H, Nitta M. Early prognostication of neurological outcome by heart rate

1139   variability in adult patients with out-of-hospital sudden cardiac arrest. Crit Care. 2019;23: 323.

1140   doi:10.1186/s13054-019-2603-6

1141   51.   Pérez P, Madsen J, Banellis L, Türker B, Raimondo F, Perlbarg V, et al. Conscious processing of

1142   narrative stimuli synchronizes heart rate between individuals. Cell Rep. 2021;36: 109692.

1143   doi:10.1016/j.celrep.2021.109692

1144   52.   Raimondo F, Rohaut B, Demertzi A, Valente M, Engemann DA, Salti M, et al. Brain–heart interactions

1145   reveal consciousness in noncommunicating patients. Ann Neurol. 2017;82: 578–591.

1146   doi:10.1002/ana.25045

1147   53.   Riganello F, Zubler F, Haenggi M, De Lucia M. Heart rate complexity: An early prognostic marker of

1148   patient outcome after cardiac arrest. Clin Neurophysiol. 2022;134: 27–33.

1149   doi:10.1016/j.clinph.2021.10.019

1150   54.   Pfeifer G, Garfinkel SN, Gould van Praag CD, Sahota K, Betka S, Critchley HD. Feedback from the heart:

1151   Emotional learning and memory is controlled by cardiac cycle, interoceptive accuracy and personality.

1152   Biol Psychol. 2017;126: 19–29. doi:10.1016/j.biopsycho.2017.04.001

1153   55.   Waselius T, Wikgren J, Halkola H, Penttonen M, Nokia MS. Learning by heart: cardiac cycle reveals an

1154   effective time window for learning. J Neurophysiol. 2018;120: 830–838. doi:10.1152/jn.00128.2018

1155   56.   Kimura K. Cardiac cycle modulates reward feedback processing: An ERP study. Neurosci Lett. 2019;711:

1156   13447. doi:10.1016/j.neulet.2019.134473

1157   57.   Adelhöfer N, Schreiter ML, Beste C. Cardiac cycle gated cognitive-emotional control in superior frontal

1158   cortices. Neuroimage. 2020;222: 11727. doi:10.1016/j.neuroimage.2020.117275

1159   58.   Petzschner FH, Weber LA, Wellstein K V., Paolini G, Do CT, Stephan KE. Focus of attention modulates

1160   the heartbeat evoked potential. Neuroimage. 2019;186: 595–606. doi:10.1016/j.neuroimage.2018.11.037

1161   59.   Kunzendorf S, Klotzsche F, Akbal M, Villringer A, Ohl S, Gaebler M. Active information sampling varies

1162   across the cardiac cycle. Psychophysiology. 2019;56: e13322. doi:10.1111/psyp.13322

1163   60.   Al E, Iliopoulos F, Forschack N, Nierhaus T, Grund M, Motyka P, et al. Heart-brain interactions shape

1164   somatosensory perception and evoked potentials. Proc Natl Acad Sci U S A. 2020;117: 10575–10584.

1165   doi:10.1073/pnas.1915629117

1166   61.   Seth AK, Suzuki K, Critchley HD. An interoceptive predictive coding model of conscious presence. Front

1167   Psychol. 2012;2: 395. doi:10.3389/fpsyg.2011.00395

1168   62.   Park HD, Tallon-Baudry C. The neural subjective frame: From bodily signals to perceptual consciousness.

1169   Philos Trans R Soc B Biol Sci. 2014;369: 20130. doi:10.1098/rstb.2013.0208

1170   63.   Park HD, Blanke O. Coupling Inner and Outer Body for Self-Consciousness. Trends Cogn Sci. 2019;23:

1171   377–388. doi:10.1016/j.tics.2019.02.002

1172   64.   Iber C, Ancoli-Israel S, Chesson Jr. AL, Quan SF. The AASM Manual for the Scoring of Sleep and

1173   Associated Events. 2007.

1174   65.   Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics

1175   including independent component analysis. J Neurosci Methods. 2004;134: 9–21.

1176   doi:10.1016/j.jneumeth.2003.10.009

1177   66.   Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software for advanced analysis of

1178   MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci. 2011;2011: 1568.

1179   doi:10.1155/2011/156869

67.   Allen M, Poggiali D, Whitaker K, Marshall TR, Kievit RA. Raincloud plots: A multi-platform tool for robust

data visualization. Wellcome Open Res. 2019;4: 1–51. doi:10.12688/wellcomeopenres.15191.1

68.   Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods.

2007;164: 177–190. doi:10.1016/j.jneumeth.2007.03.024

69.   Bouchard M, Lina JM, Gaudreault PO, Lafreniret A, Duba J, Gosselin N, et al. Sleeping at the switch.

Elife. 2021;10: 1–17. doi:10.7554/eLife.64337

70.   Nolan H, Whelan R, Reilly RB. FASTER: Fully Automated Statistical Thresholding for EEG artifact

Rejection. J Neurosci Methods. 2010;192: 152–162. doi:10.1016/j.jneumeth.2010.07.015

71.   Andrillon T, Pressnitzer D, Léger D, Kouider S. Formation and suppression of acoustic memories during

human sleep. Nat Commun. 2017;8: 179. doi:10.1038/s41467-017-00071-z

72.   Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods.

2007;164: 177–190. doi:10.1016/j.jneumeth.2007.03.024

73.   Sawilowsky SS. New Effect Size Rules of Thumb. J Mod Appl Stat Methods. 2009;8: 597–599.

doi:10.22237/jmasm/1257035100

1194