Parallel Mechanosensory Systems are Required for Negative Gravitaxis in *C. elegans*

Authors

Caroline Ackley[1], Neda Ziaei Kajbaf[1], Lindsey Washiashi[1], Ruchira Krishnamurthy[1], Pradeep Joshi[1], Joel H. Rothman[1]

[1]Department of MCD Biology and the Neuroscience Research Institute, University of California, Santa Barbara, CA, USA

Correspondence

carolineackley@ucsb.edu

Abstract

The ability to sense Earth's gravitational pull is essential for orientation, navigation, and proprioception in most organisms. Here, we demonstrate that *C. elegans* adults and dauers exhibit negative gravitaxis and that this behavior is modulated by other environmental inputs such as light and electromagnetic fields. We found that components of the DEG/ENaC pathway involved in gentle touch sensation – specifically the neuronally-enriched microtubule subunits MEC-7 and MEC-12, but not the related MEC-4 and MEC-10 mechanosensory subunits – are required for negative gravitaxis. In addition, we found that TRPA-1, a receptor protein associated with cold and mechanosensation in both mammals and invertebrates, is critical for this behavior. Finally, we show that PVD neurons, which utilize TRPA-1 to detect harsh cold, are also essential for gravity sensation. Conversely, TRNs are not required for this behavior. These findings implicate a complex, interconnected mechanism for gravity sensation involving an ion channel that is also present in the mammalian vestibular system.

Introduction

Gravitropism is a common behavior among Eukaryotes. Members of plants, animals, fungi, and even

protists depend on gravity sensation for survival. Small changes in position or orientation result in a

mechanical force that is transduced by graviperceptive organs. This force is often conveyed using dense

organelles or mineral-rich structures whose displacement triggers a signaling pathway that ultimately

results in a behavior[1]. Another common characteristic of gravity transduction pathways is the use of

ciliated neurons, in which the deflection of hair-like "stereocilia" opens mechanically gated ion

channels[2], [3]. Although this general mechanism has been well characterized, less is known about how

mechanoreceptors and mechanosensory cells transduce such minute forces – calculated to be as small as

0.57-1.13pN in *Euglena*, for example –into a robust signal[4].

Given the availability of detailed connectome of the entire nervous system and powerful genetic and

optogenetic tools, *C. elegans* has been a powerful model organism for elucidation of the neural circuitry

and molecular mechanism governing sensory perception, learning, memory, and behavior[5]–[12].

However, because they are traditionally studied in a two-dimensional environment (on petri dishes

containing agar), few studies have examined complex behavior exhibited by *C. elegans* in three-

dimensions. In the wild, *C. elegans* nematodes are typically found in compost in wet areas (such as under

shrubs or along riverbanks) that can change drastically from season to season[13], [14]. *C. elegans* has

evolved an adaptive alternative dauer phase that prioritizes dispersal over reproduction during adverse

environmental conditions. Dauers exhibit characteristic nictation behavior in which they "stand" on their

tails and wave their heads. This is believed to facilitate dispersal, possibly by allowing worms to "hitch a

ride" on nearby isopods[15]. Therefore, we believe that gravity may be a critical input that *C. elegans* use

in combination with other cues to navigate to the surface before traveling longer distances (Fig. 1A).

Surprisingly, previous studies have shown that *C. elegans* dauers do not demonstrate a gravitactic

preference, unlike *C. japonica*, which negatively gravitax[16] and that adult worms orient downwards

when swimming in liquid[17]. However, as in *Drosophila* and *Ciona*, gravitaxis in worms is likely context-

dependent[18], [19]. *C. elegans* has a strong aversive response to light[6] and responds to electromagnetic

fields[20]–[22]. Therefore, integration with other sensory modalities may influence or mask gravitactic

behavior. No gravitactic experiments have been performed in the absence of either stimulus. In this study

we show that in the absence of light, *C. elegans* dauers exhibit preferential negative gravitaxis which is

enhanced when shielded from electromagnetic fields.

While there is less known about behavioral responses to gravity, the effects of hyper- and microgravity

on *C. elegans* physiology is well documented[23]–[27]. In response to hypergravitational forces, signaling

by the mechanosensory receptor sodium channel proteins MEC-4/MEC-10 DEG/ENaC lead to nuclear

localization of DAF-16 FoxO transcription factor[28]. These 'mec' genes were found to be expressed in the

six touch receptive neurons (TRNs) in *C. elegans* that specifically mediate gentle touch sensation[29].

Additionally, MEC-10 is expressed in the proprioceptive/multimodal neurons PVDL/R and FLPL/R, while

MEC-4 is found in the FLP pair only (Fig. 1B). Mechanosensation is also mediated by Trp cation channels

which are common in vertebrates and invertebrates[30]–[33]. The TRPA-1 receptor is a polymodal sensor,

capable of conveying either high or low temperatures as well as light and the presence of noxious

stimuli[34]. In worms, TRPA-1 confers sensitivity to noxious cold in PVDs as well as mechanosensation in

OLQ and IL1 neurons. The *trpa-1* homologs *pyx* and *pain* are necessary for gravitaxis in

Drosophila[35].Therefore, it seems likely that *C. elegans* would have a system for detecting normal

gravitational force that may utilize components of known mechanosensory pathways and that some of

these neurons might be involved in graviperception.

We discovered that *C. elegans* worms prefer to migrate vertically against the force of gravity – negative

gravitaxis – and that this behavior is most pronounced in dauer larvae. Further, gravitaxis is influenced by

environmental cues such as light and background electromagnetic fields. We also show that the TRPA-1

channel and MEC-7/12 microtubules, but not the associated channel proteins DEG/ENaC MEC-4/10, are

required for this behavior. Finally, we identified the polymodal PVD neurons, which spans the length of

the worm, as essential for this directional bias. Based on these findings, we propose a model of gravity

transduction involving parallel mechanosensory pathways and PVD neurons.

Results

***C. elegans* exhibits negative gravitactic behavior**

Under stressful conditions, including overcrowding, lack of resources, and extreme temperatures, *C.*

elegans larvae develop into dauers. Through drastic physical, metabolic, and behavioral changes, dauers

become efficient vectors for dispersal that can survive for months without food[36]. Under normal

laboratory growth conditions where *C. elegans* are cultivated on flat agar surface in petri dishes, *C. elegans*

dauers do not nictate unless contaminated by fungi. Nictation behavior is also observed when grown on

three dimensional habitable scaffolds[37]; however, 3D scaffolds are not very convenient to study

migratory behavior.

To address this issue, we adapted a setup used to study neuromuscular integrity in *C. elegans*[38]

to investigate if *C. elegans* dauers exhibit directional bias in response to gravity. We injected dauers into

gravitaxis chambers comprised of two serological pipettes that, when stacked end to end, allowed for ~25

cm of movement in either direction away from the injection site (Fig.1C). Pipette chambers loaded with

worms were oriented either vertically or horizontally and were left overnight and scored the following

day. In the absence of any cues, worms would not exhibit any migratory bias in either direction when the

chambers are oriented horizontally. For all gravitaxis assays, we included a horizontal comparison to

control for any directional preference attributable to the construction and design of the chamber itself. If

worms perceive gravity they would exhibit a directional bias in the vertically oriented chamber – a

migratory bias to the bottom if exhibiting positive gravitaxis and a migratory bias to the top in case of

negative gravitaxis.

Contrary to prior assays conducted on petri plates that showed *C. elegans* dauers do not favor

any direction in response to gravity[16], N2 dauers exhibited a small but significant directional bias in

migration to the top of the vertical chamber in stark contrast to lack of any directional preference in

horizontal controls ((n = 3,816 worms over 14 trials; $p < 0.001$) (Fig. 2A, B)). A Gravitaxis Index (GI)

analogous to the chemotaxis Index (CI)[39], that represents the fraction of worms that migrate to the top

or bottom of the gravitaxis chamber did not represent the migratory behavior of the worms (GI vertical =

0.43, SEM = 0.17; GI horizontal = 0.20, SEM = 0.16; $p > 0.05$, two-tailed t test). Instead, we analyzed the

distribution of worms across each assay and compared vertical and horizontal distributions using Kruskal-

Wallis. We performed Dunn's test for multiple comparisons with Bonferroni's correction when comparing

across multiple conditions.

Gravity perception competes with light and electromagnetic signals in governing migration

C. elegans exhibits strong negative phototactic behavior. It is possible that the vertical negative gravitactic

migration might be adversely influenced by the strong response to light (Fig. 1A). We hypothesized that

N2 dauer larvae might exhibit a stronger negative gravitaxis effect in the dark. We repeated the assay in

the dark using a light-blocking blackout cloth. We found that N2 dauers demonstrated a strong preference

for migration towards the top of the chambers under these conditions in stark contrast to lack of any

directional preference in horizontal controls (n = 37,869 worms over 108 trials; $p < 0.001$) (Fig. 2A, B).

Significantly, this preference increased relative to worms tested under normal laboratory lighting

conditions ($p < 0.01$) (Fig. 2C).

In addition to photosensitivity, *C. elegans* adults have been shown to respond to electromagnetic

fields and their direction of migration is influenced by their satiety state – well fed worms migrate up

while starved worms migrated down a vertical column[20]. To test if electromagnetic fields (EMF)

influence the gravitaxis behavior of dauers, we decided to shield our assay from background EM fields

present in the lab using a simple Faraday cage made of aluminum foil. To our surprise, removing

background EMF also had a drastic effect on gravitactic behavior. Dauers in the -light -EMF condition

negatively gravitaxed in comparison with horizontal controls (p < 0.001; n = 37,869 worms over 108 trials)

(Fig. 2A, B) and were significantly more negatively gravitactic than either the -light +EMF or +light +EMF

conditions (p < 0.001 in both cases) (Fig. 2C). Over the course of several months, we noticed that gravitaxis

in our Faraday cage gradually decreased. After reinforcing the cage once more, we saw a recovery of

strong negative gravitatic behavior, adding to our observation that worms rely on both EMF and gravity

inputs in determining vertical movement (Fig. S1). Thus, *C. elegans* exhibits negative gravitaxis behavior

which becomes pronounced when sensory stimulus by light is blocked and is further enhanced when

shielded from electromagnetic fields suggesting that *C. elegans* gravitaxis behavior is complex and a

consequence of integration of multiple sensory inputs. We continued to use a Faraday cage for the

remainder of the experiments described in this study.

In addition to external cues, we were curious if development has any effect on gravitaxis behavior.

According to our hypothesis, after worms find suitable conditions and reach adulthood, there is less

motivational drive to migrate long distances in any direction. Instead, adult worms are cannibalized by

their offspring when resources are scarce allowing the offspring to enter the alternate

developmental/dispersal dauer stage. We hypothesized that adult worms would either lose motivation to

migrate against the force of gravity or would lose the ability to sense gravity altogether. In both cases, this

would result in diminished gravitaxis in our assay.

As expected, N2 adults were less gravitactic than dauers in the absence of light and EMF (p <

0.001) (Fig. 2C). However, adults negatively gravitaxed significantly more than horizontal controls (p <

0.001; n = 1,297 worms over 11 trials) (Fig. 2A, B). Moreover, reduced gravitaxis in adults may be an

artifact of our experimental setup that was designed for dauers. Because of their length, adults can cover

a greater distance in a short amount of time with less exertion. It is possible that between the start of the

experiment and when the worms were scored, adults were able to reach the end of the chamber and turn

back. Scaling up the assay or monitoring the adults as they move would provide more insight into this

question.

Our results that *C. elegans* dauers and adults exhibit negative gravitaxis on solid surface in

contradiction to prior study are likely attributable to assay design. Previous experiments have mostly been

performed using Petri dishes, which limit the distance worms can travel. We were able to find robust

differences between horizontal and vertically oriented chambers in part owing to the long distances they

traversed. In addition, the environment of a worm crawling in a thin space between the agar and tube

wall may be somewhat more analogous to crawling through a column of soil than the environment worms

experience on the side of an open plate. Likewise, earlier experiments on gravitaxis in *C. elegans* make no

mention of the lighting conditions used, leading us to believe that signal integration may have complicated

these initial findings. A correlation was found between the number of worms in the chamber and the

range of distances travelled, which points to a social aspect of this behavior that may also have been

unaccounted for (Fig. S2).

Neuron-specific microtubules are required for gravitaxis

Gravitational forces perceived by invertebrates can be very small and consequently invertebrates have

evolved extremely sensitive mechanisms for perceiving these tiny forces. In some invertebrates, stretch

receptors are imbedded in the cuticle which results in an ion influx upon deformation in a manner

analogous to stretch and pressure sensors in mammals[40]. Gentle touch in *C. elegans* is perceived by the

mechanosensory DEG/ENaC channels MEC-4 and MEC-10[29] which form a pressure-sensitive channel in

Touch Receptor Neurons (TRNs) that respond to cuticle deflections through extracellular matrix proteins

MEC-1, MEC-5, and MEC-9[41] (Fig. 3). MEC-12 and MEC-7, which encode alpha and beta tubulin

respectively, form unique 15 protofilament microtubules only in the six TRNs. These microtubule bundles

are thicker than the typical 11 protofilament microtubules found in most cells, including other

neurons[42]. Although the exact role of these unique microtubule structures in touch sensation is not well

understood, the 15-protofilament structures function in part by anchoring the MEC-4/10 receptor

complex within the cell[43]

Dauers defective in some components of the gentle touch perception pathway – *mec-2(e75)*, *mec-*

18(u228), *mec-4(e1339))*., *mec-10(tm1552)*, *mec-12(e1605),* and *mec-7(ok2152)* – exhibited impaired

negative gravitaxis when compared with N2 dauers (Fig. 4A-C). While the mechanoreceptor subunit

mutants *mec-4* and *mec-10* showed a modest, non-significant decrease in their ability to gravitax

compared with wildtype controls (Fig. 4C), vertically oriented worms exhibited a significant directional

bias from their horizontal controls (n = 1,750 worms over 12 trials; p < 0.001 and n = 4,422 worms over

25 trials; p < 0.001 for *mec-4* and *mec-10*, respectively) (Fig. 4A, B) suggesting that the ability to perceive

gravity is not seriously impaired in mutants lacking MEC-4/MEC-10 mechanosensory channel subunits. As

a caveat, the *mec-4* allele used here (*e1339*) is a missense mutation that exhibits weaker phenotypes

generally than the null, *tu253*. When tested in our assay, *mec-4(tu253)* worms do not gravitax (Fig. S3);

however, these worms are also sluggish, which has led to *e1339* being used in place of *tu253* in at least

one other study[44]. In contrast to our *mec-4*/*mec-10* results, both microtubule subunit mutants (*mec-7*

and *mec-12*) lose directional preference entirely and behave like the horizontal controls (n = 1,441 over 8

trials; p > 0.5 and n = 10,094 worms over 21 trials; p > 0.5, respectively) (Fig. 4A-C).

We examined the known *mec-7/12* phenotypes to understand how these genes alone could alter

gravitaxis behavior. Because MEC-7 and MEC-12 are tubulin proteins, mutations in either gene alter

axonal outgrowth to varying degrees[45], [46]. However, multiple studies have shown that the single

missense mutation H192Y in MEC-12, which we used in our experiments, eliminates gentle touch

sensation even though TRN development is not significantly affected[43], [47], [48]. MEC-12 is unique in

part because it is the only *C. elegans* microtubule subunit that is acetylated at K40[45], [49] and mutations

that modify K40 or MEC-17, a transacetylase necessary for microtubule acetylation and stability in TRNs,

also prohibit mechanosensation[50]–[52].　We tested whether acetylation of MEC-12 by MEC-17 is

necessary for gravitaxis in *mec-17(u265)* worms that lack functional MEC-17. This mutation did not affect

gravitaxis, as vertically oriented *mec-17* worms differed significantly from horizontal controls (n = 1,101

worms over 7 trials; $p < 0.001$) and were not statistically different from N2 dauers ($p > 0.05$) (Fig. 4A-C).

In addition to anchoring MEC-4/MEC-10 ion channels in TRNs, MEC-7/MEC-12 microtubules are

involved in regulating gene expression through MAPK signaling, ultimately targeting the transcription

factor CEBP-1. MEC-7/MEC-12 could be regulating gravitaxis through CEBP-1. However, the deletion

mutant *cebp-1(tm2807)* showed no defect in our gravitaxis assay; *cebp-1* dauers migrated significantly

higher in vertically oriented chambers than to the equivalent distance from the origin horizontally and

were statistically indistinguishable from N2 dauer controls (n = 1,140 worms over 9 trials; $p < 0.001$ and p

> 0.5, respectively) (Fig. 4A-C). Thus, CEBP-1 activation by MEC-7/MEC-12 is not essential for transducing

gravitational force.

Other candidate genes within the DEG/ENaC pathway were tested to determine the amount of

crossover between the canonical system and a potentially new gravitactic mechanism. *mec-2* encodes an

accessory protein that is part of the MEC-4/MEC-10 channel complex that acts as a linker between

receptors and 15 protofilament microtubules. Like the *mec-4* and *mec-10* mutants, *mec-2(e75)* dauers

were not significantly impaired in their ability to gravitax (n = 4,147 worms over 20 trials; $p < 0.001$) (Fig.

4A, B). *mec-18(u228)* mutants also exhibited similar results (n = 5,820 worms over 9 trials; $p < 0.001$),

although the exact function of MEC-18 and its role in gentle touch sensation is not well characterized[53],

[54]. Finally, an allele resulting in defective MEC-3 – a transcription factor required for differentiation of

several sensory neurons – did not significantly affect gravitaxis (n = 2,882 worms over 12 trials; $p < 0.001$).

Taken together, these results support the conclusion that MEC-7/MEC-12 microtubules might be involved

in a distinct role in gravity perception aside from their structural role in axon and dendrite length,

anchoring the DEG/ENaC channel subunits, or regulation of CEBP-1 in transducing mechanical force. The

lack of a strong effect in *mec-4* and *mec-10* mutants also indicates that other types of mechanosensory

channels may be involved in gravity perception.

TRPA-1 is critical for gravity sensation

Ion channel protein TRPA-1 is one member of a superfamily of TRP proteins involved in many sensory

modalities, including sensitivity to touch, hot and cold temperatures, noxious chemicals, and light[31],

[34]. Orthologs have been found across the animal kingdom, including in mammals, fish, and other

invertebrates such as *D. melanogaster*. Mouse TRPA-1 is also present in the vestibular system, the primary

organ where gravity sensation occurs in mammals[55]. In *C. elegans*, *trpa-1* is expressed in the

proprioceptive/nociceptive pair of neurons PVDL/R, among others. Although TRPA-1 is known to play a

role in noxious cold sensation in these neurons, *Drosophila* homologs of TRPA channels, *pain* and *pyx,*

are required for gravity sensation[35]. Additionally, *trpa-1* knockout mice are shown to have impairments

in mechanosensation as well as perceiving noxious cold[33]. Likewise, *C. elegans* TRPA-1 is likely to have

mechanosensory functions in addition to its known role in cold sensation[33].

The *trpa-1(ok999)* mutation is a large deletion that removes about 1.4 kbp from the N-terminus of the

TRPA-1 protein[33]. When *trpa-1* mutants were tested in our assay, they showed no directional

preference when compared with horizontal controls (n = 2,692 worms over 17 trials; p > 0.05) (Fig. 5A, B).

Predictably, vertically oriented *trpa-1* mutants differed significantly from vertical N2 controls (p < 0.001)

(Fig 5A, B). *trpa-1(ok999)* mutant adult worms exhibit several movement defects including decreased

forward locomotion as well as several variations in the sinusoidal movement typical to wildtype

worms[56] and as such could also affect vertical migration leading to misinterpretation of the role of

TRPA-1 in graviperception. However, in our assay, *trpa-1(ok999)* dauers were distributed similarly across

the chambers in both the horizontal and vertical conditions, demonstrating that they can travel long

distances regardless of orientation.

PVD neurons, and not TRNs, are required for gravitaxis

The role of *trpa-1, mec-7,* and *mec-12* and not *mec-4/10* in affecting this behavior presented an interesting

challenge. Although MEC-7 and MEC-12 are present in many neurons, their function is not well

understood outside of TRNs. There is no reported connection between microtubules containing MEC-7/12

and TRPA-1 function. Of the six TRNs, only one expresses *trpa-1*[57]. However, we noticed that PVDs,

which detect muscle movement in addition to a host of other stimuli, express all three candidate genes.

We hypothesized, therefore, that PVD neurons could be integration centers for two proprioceptive inputs:

muscle movement and relative position to Earth's gravity.

To test this hypothesis, we genetically ablated the TRNs and PVDs by expression of constitutively active

MEC-4 channels and DEG-3, respectively. The *mec-4(d)* gain of function (GOF) mutation results in

constitutive MEC-4/10 activation that degrades TRNs specifically, which we confirmed through the

absence of *mec-3p::*RFP marked TRNs (Fig. 6A-F). PVD neurons were ablated (PVD-) by the expression of

a constitutively active version of the nicotinic acetylcholine receptor (nAChR) channel subunit, *deg-*

*3(u662)/*DEG-3, under the *ser2prom3* promoter[58]. By adulthood, the highly branched PVD neurons have

developed 4° processes (Fig. 6A). 1°-3° processes can be observed in dauers (Fig. 6E). This ablation was

confirmed using *ser2prom3::*GFP (Fig. 6D).

As expected, ablation of the TRN neurons did not alter the ability of the worms to perceive gravity.

TRN- worms did not perform differently from N2 in our gravitaxis assay ($p > 0.05$) (Fig. 7B) and showed a

directional bias to the top compared with horizontal controls (n = 2,680 worms over 19 trials; $p < 0.001$)

(Fig. 7A, B). Meanwhile, PVD- worms did not gravitax, although their distribution differed significantly

from the horizontal condition (n = 4,855 worms over 15 trials; $p < 0.001$) (Fig. 7A, B). This difference is

attributable to a higher percentage of worms clustered at the center of the vertical assay, whereas worms

tested in the horizontal condition showed a slight (but likely random) preference for one side of the

chamber. As before, we suspect that this difference is due to choice and not ability, as PVD- worms appear

to have normal movement and are capable of reaching the ends of the chamber in any orientation.

Discussion

In this study, we found that *C. elegans* demonstrate a strong negative gravitactic preference especially in

the dispersal dauer phase, and less prominently in adults. In response to stressors such as overcrowding,

lack of food, and poor environmental conditions, *C. elegans* enter an altered developmental dauer

pathway that facilitates survival and enables them to seek a better habitat[14], [59]. Therefore, if a habitat

underground becomes poor, it is crucial that dauers find their way to the surface and nictate in the

absence of other guiding cues. Recent studies have also found that worms are sensitive to the earth's

magnetic fields[20]–[22]in addition to photosensitivity[6]. We found that the normal lighting conditions

and EM field in our laboratory setting was enough to alter gravitactic behavior. This supports our

understanding of signal integration in the nervous system, in which several sensory inputs are constantly

compared when making decisions[60].

The extent to which light, EM fields, and other inputs override gravity sensation, as well as the circuits

required to calculate their respective influence on behavior, are important questions for future studies.

Previous assays for gravitaxis by *C. elegans* dauers conducted on plates did not detect any directional

preference while *C. japonica* dauers under similar assay conditions were strongly negatively

gravitactic[16]. Our altered assay has shown that C. elegans dauers as well as adults also exhibit negative

gravitaxis. The reduced ability of N2 dauers to exhibit negative gravitaxis might be an outcome of myriad

adaptations to laboratory cultivation as has been observed for several other traits including social feeding,

egg laying behavior, oxygen tolerance, as well as nictation[11], [14], [15], [61]–[63]. Interestingly, our

results suggest that negative gravitaxis is maintained (albeit weakened) in the LSJ1 sister strain, which has

been raised in liquid media over several generations (Fig. S4). It would be interesting to see how these

two strains compare gravitactically when swimming rather than crawling on solid media.

Finding that *trpa-1* and components of the DEG/ENaC system are both involved in gravitaxis reveals a

complex mechanism for transducing gravitational force. Unlike the DEG/ENaC channels, TRPA-1 receptors

do not require internal or external machinery to function[33]. Additionally, *trpa-1* is only expressed in one

of six TRNs, which are the only neurons containing the 15 protofilament structures formed by MEC-

7/MEC-12 tubulins. As described previously, MEC-7/MEC-12 microtubules may have a function in

mechanosensation beyond their structure; whether this function can take place when they are not

assembled into 15 protofilament structures has not been explored. These findings point to at least two

redundant pathways involved in gravity sensation that need to be explored in more detail as well as their

possible role in nictation.

Finally, we found evidence pointing to a pair of neurons, PVDL/R, as primary gravity-sensors in worms.

Unlike TRN- worms, which behave normally in our gravitaxis assay when compared with controls, PVD-

worms have a distinct lack of preference in response to gravity. These results, in conjunction with our

trpa-1 data, provides a sketch of how gravity may be transduced at a molecular and cellular level. We

propose that PVD neurons utilize TRPA-1 to provide information about the direction of gravity. However,

we are still missing key details about this mechanism – we cannot conclude, for example, that TRPA-1 is

the primary gravity receptor in worms. As with all TRPs, TRPA-1 is known to have primary and secondary

roles in a variety of sensory pathways[30]. Likewise, we do not know how MEC-7 and MEC-12 microtubule

subunits may be acting in gravity transduction. They may be performing a vital function within PVD

neurons that is required to transmit a signal; it is also possible that they play a role in other important

supportive or downstream neurons. As stated previously, ciliated neurons are frequently used in gravity

sensation[2], [3]. C. elegans have 60 ciliated neurons[58] – of these, IL2 neurons may also be involved in

detecting gravity because of their known role in nictation[16]. Other ciliated neurons express MEC-12,

TRPA-1, and the likely gravity sensitive mechanoreceptor in mammals, TMC-1[60]–[66]. Thus, PVDs may

be secondary neurons that act within a circuit of cells that make decisions about movements with respect

to gravity.

Understanding how *C. elegans* perceives gravity has implications for both worm and mammalian

biology. For years, *C. elegans* has been used as a model for understanding the effects of microgravity on

human physiology in space[23], [24], [26], [67]–[71]. Despite this, only a few studies have investigated

whether worms are capable of sensing gravity in the nervous system. Additionally, gravity perception is

linked evolutionarily and developmentally with hearing[72], [73]. The vestibular and auditory systems in

mammals share similar sensory structures, transduction machinery, and gene expression[74].

Mammalian TRPA1 has already been identified for its putative role in sensory transduction within inner

ear hair cells in mice[75]. The similarities between these systems makes the study of gravity sensation

important not only for understanding mammalian vestibular sensation, but auditory sensation as well. In

addition to its potential function in the inner ear, *trpa-1* homologs may also be involved in pain and

inflammation, as well as childhood asthma in humans[76], [77]. Remarkably a recent study shows that C.

elegans is able to sense sound through mechanical perturbations of the skin and involves the multimodal

PVD neurons and the FLP mechanosensory neurons[78]. Uncovering the role of TRPA-1 as a

mechanoreceptor opens the door to using *C. elegans* as a model for studying audition and auditory

disorders and has further implications for translational research.

Materials and Methods

Worm rearing

C. elegans worms were grown and maintained on OP50 *E.coli* at room temperature (23°C) using standard methods[85]. Dauer formation was induced by overcrowding, which occurred 8-12 days after chunking onto seeded NGM plates. Adults used for gravitaxis assays and imaging were taken from synchronized populations 2 days after hatching. Crosses were performed as described previously[86], using fluorescence microscopy and/or PCR to confirm the outcome of each cross. All strains used in this study are included in Table S1.

Dauer isolation

Dauer larvae were isolated from 1-5 starved NGM plates using 1% SDS and a 30% sucrose gradient as described previously[59], [87]. Briefly, worms were rinsed and collected in M9, then pelleted and resuspended in 7mL 1% SDS in a 15 mL falcon tube. Worms were left rotating for 30 minutes in SDS to kill all non-dauer lifestages. After 3-5 rinses with M9, worms were resuspended in 10 mL 30% sucrose solution and centrifuged for 5 minutes. Surviving dauers were collected at the interface of the water-sucrose gradient with a large bore glass Pasteur pipette and rinsed 3-5X in M9. Worm were used immediately or after rotating in solution overnight.

Synchronization

Mixed stage worms were rinsed from confluent plates with M9 and collected in 1.5 mL tubes. Worms were then rinsed and pelleted. A 1 mL bleach solution containing 150 μL bleach and 50 μL 5N KOH was added and worms were lysed in this solution for 3 minutes, or until most corpses were dissolved. The embryo pellet was then rinsed 3-5X in M9 and plated onto OP50 NGM plates until adulthood (~2 days at 23°C).

Assay preparation

Gravitaxis assay chambers were created using 5 mL serological pipettes and 4% NGM agar. To assemble each chamber, a heated blade was used to cut the tapered end of one pipette and the cotton plug was removed from the end of a second pipette. The two pipettes were then joined end-to-end by melting the two ends slightly over a Bunsen burner and fusing them with moderate pressure, ensuring that no gaps remained. The NGM media was then autoclaved and then taken up by each double pipette using a standard serological pipettor while the agar was still molten. Best results were achieved by orienting each pipette parallel with the benchtop as much as possible during the procedure and as the agar solidified. Pipettes were used within 24 hours of construction or stored at 4°C to be used within 48 hrs.

Once solidified, a heated 3mm Allen key was used to punch a hole through one wall of the pipette about 5mm below the fusion point (for reference, the cotton-plug end is the "top"). Each end of the pipette was removed using a heated blade and sealed with parafilm.

Gravitaxis assay

Worms were pelleted in M9 and were either extracted directly from the pellet or pipetted onto a square of parafilm. After cutting the tip to increase the bore, a pipette was used to aspirate 0.5-2 uL of the concentrated worm pellet, letting in a small volume of air at the end to facilitate injection. Worms were

injected into the agar at the opening created by the Allen key. This opening was then sealed with parafilm, and the chamber was immediately oriented vertically or horizontally. Worms were allowed to gravitax overnight and were scored the following day (typically 12-24 hrs after injection).

Pipettes were examined under a dissecting microscope and a marker was used to mark the location of each worm. Pipettes were scored immediately after removal from the assay location. Worms were only scored if they appeared alive, healthy, and were not swimming (chambers were only scored if a majority of worms were alive and crawling). Worms were not scored within 2.5 cm to either side of the injection point.

To tally the location of each worm, pipettes were divided into 7 sections on each side of the 5 cm injection site. Each section was 3.5 cm long (the same length as a 1 mL volume indicated on the pipettes). The number of marker points in each section was then recorded.

Imaging

Fluorescent micrographs were taken using a Leica SP8 Resonant Scanning Confocal microscope at 10-40x. Brightness and contrast were adjusted and overlays added using ImageJ.

Quantification and Statistical Analysis

Data were collected in Excel and analyzed using RStudio. Statistical tests, p values, sample sizes, and replicates are detailed in the results and figure legends.

References

[1] M. D. Ross, K. G. Pote, A. Miller, D. C. Phillips, and R. J. P. Williams, "Some properties of otoconia," *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, vol. 304, no. 1121, pp. 445–452, Feb. 1984, doi: 10.1098/rstb.1984.0038.

[2] L. A. Bezares-Calderón, J. Berger, and G. Jékely, "Diversity of cilia-based mechanosensory systems and their functions in marine animal behaviour," *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, vol. 375, no. 1792, p. 20190376, Feb. 2020, doi: 10.1098/rstb.2019.0376.

[3] F. Lacquaniti *et al.*, "Multisensory Integration and Internal Models for Sensing Gravity Effects in Primates," *BioMed Res. Int.*, vol. 2014, pp. 1–10, 2014, doi: 10.1155/2014/615854.

[4] D.-P. Häder and R. Hemmersbach, "Gravitaxis in Euglena," in *Euglena: Biochemistry, Cell and Molecular Biology*, S. D. Schwartzbach and S. Shigeoka, Eds. Cham: Springer International Publishing, 2017, pp. 237–266. doi: 10.1007/978-3-319-54910-1_12.

[5] D. L. Riddle, T. Blumenthal, B. J. Meyer, and J. R. Priess, *C. elegans Responds to a Variety of Chemicals*. Cold Spring Harbor Laboratory Press, 1997. Accessed: Nov. 27, 2018. [Online]. Available: https://www.ncbi.nlm.nih.gov/books/NBK19972/

[6] A. Ward, J. Liu, Z. Feng, and X. Z. S. Xu, "Light-sensitive neurons and channels mediate phototaxis in C. elegans," *Nat. Neurosci.*, vol. 11, no. 8, pp. 916–922, Aug. 2008, doi: 10.1038/nn.2155.

[7] C. I. Bargmann, E. Hartwieg, and H. R. Horvitz, "Odorant-selective genes and neurons mediate olfaction in C. elegans," *Cell*, vol. 74, no. 3, pp. 515–527, Aug. 1993, doi: 10.1016/0092-8674(93)80053-H.

[8] S. Ward, "Chemotaxis by the Nematode Caenorhabditis elegans: Identification of Attractants and Analysis of the Response by Use of Mutants," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 70, no. 3, pp. 817–821, Mar. 1973.

[9] D. Ramot, B. L. MacInnis, H.-C. Lee, and M. B. Goodman, "Thermotaxis is a Robust Mechanism for Thermoregulation in C. elegans Nematodes," *J. Neurosci. Off. J. Soc. Neurosci.*, vol. 28, no. 47, pp. 12546–12557, Nov. 2008, doi: 10.1523/JNEUROSCI.2857-08.2008.

[10] J. Russell, A. G. Vidal-Gadea, A. Makay, C. Lanam, and J. T. Pierce-Shimomura, "Humidity sensation requires both mechanosensory and thermosensory pathways in Caenorhabditis elegans," *Proc. Natl. Acad. Sci.*, vol. 111, no. 22, pp. 8269–8274, Jun. 2014, doi: 10.1073/pnas.1322512111.

[11] J. M. Gray *et al.*, "Oxygen sensation and social feeding mediated by a C. elegans guanylate cyclase homologue," *Nature*, vol. 430, no. 6997, pp. 317–322, Jul. 2004, doi: 10.1038/nature02714.

[12] A. J. Bretscher, K. E. Busch, and M. de Bono, "A carbon dioxide avoidance behavior is integrated with responses to ambient oxygen and food in Caenorhabditis elegans," *Proc. Natl. Acad. Sci.*, vol. 105, no. 23, pp. 8044–8049, Jun. 2008, doi: 10.1073/pnas.0707607105.

[13] E. Kodama-Namba, L. A. Fenk, A. J. Bretscher, E. Gross, K. E. Busch, and M. de Bono, "Cross-Modulation of Homeostatic Responses to Temperature, Oxygen and Carbon Dioxide in C. elegans," *PLoS Genet.*, vol. 9, no. 12, Dec. 2013, doi: 10.1371/journal.pgen.1004011.

[14] L. Frézal and M.-A. Félix, "C. elegans outside the Petri dish," *eLife*, vol. 4, Mar. 2015, doi: 10.7554/eLife.05849.

[15] H. Lee *et al.*, "Nictation, a dispersal behavior of the nematode Caenorhabditis elegans, is regulated by IL2 neurons," *Nat. Neurosci.*, vol. 15, no. 1, pp. 107–112, Jan. 2012, doi: 10.1038/nn.2975.

[16] E. Okumura, R. Tanaka, and T. Yoshiga, "Negative gravitactic behavior of Caenorhabditis japonica dauer larvae," *J. Exp. Biol.*, vol. 216, no. 8, pp. 1470–1474, Apr. 2013, doi: 10.1242/jeb.075739.

[17] W.-L. Chen, H. Ko, H.-S. Chuang, D. M. Raizen, and H. H. Bau, "Caenorhabditis elegans exhibits positive gravitaxis," *BMC Biol.*, vol. 19, no. 1, p. 186, Sep. 2021, doi: 10.1186/s12915-021-01119-9.

[18] M. Bostwick *et al.*, "Antagonistic Inhibitory Circuits Integrate Visual and Gravitactic Behaviors," *Curr. Biol.*, vol. 30, no. 4, pp. 600-609.e2, Feb. 2020, doi: 10.1016/j.cub.2019.12.017.

[19] J.-E. Bae *et al.*, "Positive geotactic behaviors induced by geomagnetic field in Drosophila," *Mol. Brain*, vol. 9, no. 1, p. 55, Dec. 2016, doi: 10.1186/s13041-016-0235-1.

[20] A. Vidal-Gadea *et al.*, "Magnetosensitive neurons mediate geomagnetic orientation in Caenorhabditis elegans," *eLife*, vol. 4, 2015, doi: 10.7554/eLife.07493.

[21] L. Landler, S. Nimpf, T. Hochstoeger, G. C. Nordmann, A. Papadaki-Anastasopoulou, and D. A. Keays, "Comment on 'Magnetosensitive neurons...,'" *eLife*, vol. 7, 2018, doi: 10.7554/elife.30187.001.

[22] "Response to 'Comment on "Magnetosensitive neurons..."'", doi: 10.7554/elife.31414.001.

[23] Y. Gao, D. Xu, L. Zhao, and Y. Sun, "The DNA damage response of C. elegans affected by gravity sensing and radiosensitivity during the Shenzhou-8 spaceflight," *Mutat. Res. Mol. Mech. Mutagen.*, vol. 795, pp. 15–26, Jan. 2017, doi: 10.1016/j.mrfmmm.2017.01.001.

[24] Y. Gao, D. Xu, L. Zhao, M. Zhang, and Y. Sun, "Effects of microgravity on DNA damage response in Caenorhabditis elegans during Shenzhou-8 spaceflight," *Int. J. Radiat. Biol.*, vol. 91, no. 7, pp. 531–539, Jul. 2015, doi: 10.3109/09553002.2015.1043754.

[25] J. N. Saldanha, S. Pandey, and J. A. Powell-Coffman, "The effects of short-term hypergravity on Caenorhabditis elegans," *Life Sci. Space Res.*, vol. 10, pp. 38–46, Aug. 2016, doi: 10.1016/j.lssr.2016.06.003.

[26] D. Xu, Y. Gao, L. Huang, and Y. Sun, "Changes in miRNA expression profile of space-flown Caenorhabditis elegans during Shenzhou-8 mission," *Life Sci. Space Res.*, vol. 1, pp. 44–52, Apr. 2014, doi: 10.1016/j.lssr.2013.12.001.

[27] S. S. Kalichamy, T. Y. Lee, K. Yoon, and J. I. Lee, "Hypergravity hinders axonal development of motor neurons in Caenorhabditis elegans," *PeerJ*, vol. 4, p. e2666, Nov. 2016, doi: 10.7717/peerj.2666.

[28] N. Kim *et al.*, "Gravity Force Transduced by the MEC-4/MEC-10 DEG/ENaC Channel Modulates DAF-16/FoxO Activity in Caenorhabditis elegans," *Genetics*, vol. 177, no. 2, pp. 835–845, Oct. 2007, doi: 10.1534/genetics.107.076901.

[29] M. Chalfie and J. Sulston, "Developmental genetics of the mechanosensory neurons of Caenorhabditis elegans," *Dev. Biol.*, vol. 82, no. 2, pp. 358–370, Mar. 1981, doi: 10.1016/0012-1606(81)90459-0.

[30] C. Montell, "The venerable inveterate invertebrate TRP channels," *Cell Calcium*, vol. 33, no. 5–6, pp. 409–417, Jun. 2003, doi: 10.1016/s0143-4160(03)00053-8.

[31] M. Chatzigeorgiou *et al.*, "Specific roles for DEG/ENaC and TRP channels in touch and thermosensation in C. elegans nociceptors," *Nat. Neurosci.*, vol. 13, no. 7, pp. 861–868, Jul. 2010, doi: 10.1038/nn.2581.

[32] L. Han *et al.*, "Two novel DEG/ENaC channel subunits expressed in glia are needed for nose-touch sensitivity in Caenorhabditis elegans," *J. Neurosci. Off. J. Soc. Neurosci.*, vol. 33, no. 3, pp. 936–949, Jan. 2013, doi: 10.1523/JNEUROSCI.2749-12.2013.

[33] K. S. Kindt *et al.*, "Caenorhabditis elegans TRPA-1 functions in mechanosensation," *Nat. Neurosci.*, vol. 10, no. 5, pp. 568–577, May 2007, doi: 10.1038/nn1886.

[34] K. Venkatachalam, J. Luo, and C. Montell, "Evolutionarily conserved, multitasking TRP channels: lessons from worms and flies," *Handb. Exp. Pharmacol.*, vol. 223, pp. 937–962, 2014, doi: 10.1007/978-3-319-05161-1_9.

[35] Y. Sun, L. Liu, Y. Ben-Shahar, J. S. Jacobs, D. F. Eberl, and M. J. Welsh, "TRPA channels distinguish gravity sensing from hearing in Johnston's organ," *Proc. Natl. Acad. Sci.*, vol. 106, no. 32, pp. 13606–13611, Aug. 2009, doi: 10.1073/pnas.0906377106.

[36] Y. Wang, A. N. Ezemaduka, Y. Tang, and Z. Chang, "Understanding the mechanism of the dormant dauer formation of C. elegans: From genetics to biochemistry," *IUBMB Life*, vol. 61, no. 6, pp. 607–612, 2009, doi: 10.1002/iub.211.

[37] A. Guisnet, M. Maitra, S. Pradhan, and M. Hendricks, "A three-dimensional habitat for C. elegans environmental enrichment," *PLOS ONE*, vol. 16, no. 1, p. e0245139, Jan. 2021, doi: 10.1371/journal.pone.0245139.

[38] C. Bainbridge, A. Schuler, and A. G. Vidal-Gadea, "Method for the assessment of neuromuscular integrity and burrowing choice in vermiform animals," *J. Neurosci. Methods*, vol. 264, pp. 40–46, May 2016, doi: 10.1016/j.jneumeth.2016.02.023.

[39] C. I. Bargmann, "Chemosensation in C. elegans," 2006.

[40] J. A. Bender and M. A. Frye, "Invertebrate solutions for sensing gravity," *Curr. Biol.*, vol. 19, no. 5, pp. R186–R190, Mar. 2009, doi: 10.1016/j.cub.2008.12.024.

[41] J. G. Cueva, A. Mulholland, and M. B. Goodman, "Nanoscale Organization of the MEC-4 DEG/ENaC Sensory Mechanotransduction Channel in Caenorhabditis elegans Touch Receptor Neurons," *J. Neurosci.*, vol. 27, no. 51, pp. 14089–14098, Dec. 2007, doi: 10.1523/JNEUROSCI.4179-07.2007.

[42] M. Chalfie and J. N. Thomson, "Structural and functional diversity in the neuronal microtubules of Caenorhabditis elegans.," *J. Cell Biol.*, vol. 93, no. 1, pp. 15–23, Apr. 1982, doi: 10.1083/jcb.93.1.15.

[43] A. Bounoutas, R. O'Hagan, and M. Chalfie, "The multipurpose 15-protofilament microtubules in C. elegans have specific roles in mechanosensation," *Curr. Biol. CB*, vol. 19, no. 16, pp. 1362–1367, Aug. 2009, doi: 10.1016/j.cub.2009.06.036.

[44] S. M. Maguire, C. M. Clark, J. Nunnari, J. K. Pirri, and M. J. Alkema, "The C. elegans Touch Response Facilitates Escape from Predacious Fungi," *Curr. Biol.*, vol. 21, no. 15, pp. 1326–1330, Aug. 2011, doi: 10.1016/j.cub.2011.06.063.

[45] D. Lockhead *et al.*, "The tubulin repertoire of Caenorhabditis elegans sensory neurons and its context-dependent role in process outgrowth," *Mol. Biol. Cell*, vol. 27, no. 23, pp. 3717–3728, Nov. 2016, doi: 10.1091/mbc.E16-06-0473.

[46] C.-H. Chen, A. Lee, C.-P. Liao, Y.-W. Liu, and C.-L. Pan, "RHGF-1/PDZ-RhoGEF and retrograde DLK-1 signaling drive neuronal remodeling on microtubule disassembly," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 111, no. 46, pp. 16568–16573, Nov. 2014, doi: 10.1073/pnas.1410263111.

[47] M. Chalfie and M. Au, "Genetic control of differentiation of the Caenorhabditis elegans touch receptor neurons," *Science*, vol. 243, no. 4894, pp. 1027–1033, Feb. 1989, doi: 10.1126/science.2646709.

[48] A. Bounoutas, J. Kratz, L. Emtage, C. Ma, K. C. Nguyen, and M. Chalfie, "Microtubule depolymerization in Caenorhabditis elegans touch receptor neurons reduces gene expression through a p38 MAPK pathway," *Proc. Natl. Acad. Sci.*, vol. 108, no. 10, pp. 3982–3987, Mar. 2011, doi: 10.1073/pnas.1101360108.

[49] J. S. Akella *et al.*, "MEC-17 is an α-tubulin acetyltransferase," *Nature*, vol. 467, no. 7312, pp. 218–222, Sep. 2010, doi: 10.1038/nature09324.

[50] T. Shida, J. G. Cueva, Z. Xu, M. B. Goodman, and M. V. Nachury, "The major α-tubulin K40 acetyltransferase αTAT1 promotes rapid ciliogenesis and efficient mechanosensation," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 107, no. 50, pp. 21517–21522, Dec. 2010, doi: 10.1073/pnas.1013728107.

[51] I. Topalidou *et al.*, "Enzymatic and non-enzymatic activities of the tubulin acetyltransferase MEC-17 are required for microtubule organization and mechanosensation in C. elegans," *Curr. Biol.*, vol. 22, no. 12, pp. 1057–1065, Jun. 2012, doi: 10.1016/j.cub.2012.03.066.

[52] B. Neumann and M. A. Hilliard, "Loss of MEC-17 leads to microtubule instability and axonal degeneration," *Cell Rep.*, vol. 6, no. 1, pp. 93–103, Jan. 2014, doi: 10.1016/j.celrep.2013.12.004.

[53] G. Gu, G. A. Caldwell, and M. Chalfie, "Genetic interactions affecting touch sensitivity in Caenorhabditis elegans.," *Proc. Natl. Acad. Sci.*, vol. 93, no. 13, pp. 6577–6582, Jun. 1996, doi: 10.1073/pnas.93.13.6577.

[54] Y. Zhang and M. Chalfie, "MTD-1, a touch-cell-specific membrane protein with a subtle effect on touch sensitivity," *Mech. Dev.*, vol. 119, no. 1, pp. 3–7, Nov. 2002, doi: 10.1016/S0925-4773(02)00293-9.

[55] T. Kamakura *et al.*, "Functional expression of TRPV1 and TRPA1 in rat vestibular ganglia," *Neurosci. Lett.*, vol. 552, pp. 92–97, Sep. 2013, doi: 10.1016/j.neulet.2013.07.019.

[56] E. Yemini, T. Jucikas, L. J. Grundy, A. E. X. Brown, and W. R. Schafer, "A database of Caenorhabditis elegans behavioral phenotypes," *Nat. Methods*, vol. 10, no. 9, Art. no. 9, Sep. 2013, doi: 10.1038/nmeth.2560.

[57] S. R. Taylor *et al.*, "Molecular topography of an entire nervous system," Neuroscience, preprint, Dec. 2020. doi: 10.1101/2020.12.15.422897.

[58] A. Albeg *et al.*, "C. elegans multi-dendritic sensory neurons: Morphology and function," *Mol. Cell. Neurosci.*, vol. 46, no. 1, pp. 308–317, Jan. 2011, doi: 10.1016/j.mcn.2010.10.001.

[59] M. C. Ow and S. E. Hall, "A Method for Obtaining Large Populations of Synchronized Caenorhabditis elegans Dauer Larvae," in *Methods in Molecular Biology (Clifton, N.J.)*, vol. 1327, 2015, pp. 209–219.

[60] A. Metaxakis, D. Petratou, and N. Tavernarakis, "Multimodal sensory processing in Caenorhabditis elegans," *Open Biol.*, vol. 8, no. 6, p. 180049, doi: 10.1098/rsob.180049.

[61] M.-A. Félix and C. Braendle, "The natural history of Caenorhabditis elegans," *Curr. Biol.*, vol. 20, no. 22, pp. R965–R969, Nov. 2010, doi: 10.1016/j.cub.2010.09.050.

[62] M. de Bono and C. I. Bargmann, "Natural Variation in a Neuropeptide Y Receptor Homolog Modifies Social Behavior and Food Response in C. elegans," *Cell*, vol. 94, no. 5, pp. 679–689, Sep. 1998, doi: 10.1016/S0092-8674(00)81609-8.

[63] E. E. Large, R. Padmanabhan, K. L. Watkins, R. F. Campbell, W. Xu, and P. T. McGrath, "Modeling of a negative feedback mechanism explains antagonistic pleiotropy in reproduction in domesticated Caenorhabditis elegans strains," *PLoS Genet.*, vol. 13, no. 5, p. e1006769, May 2017, doi: 10.1371/journal.pgen.1006769.

[64] D. P. Corey and J. R. Holt, "Are TMCs the Mechanotransduction Channels of Vertebrate Hair Cells?," *J. Neurosci. Off. J. Soc. Neurosci.*, vol. 36, no. 43, pp. 10921–10926, Oct. 2016, doi: 10.1523/JNEUROSCI.1148-16.2016.

[65] C. A. Nist-Lund *et al.*, "Improved TMC1 gene therapy restores hearing and balance in mice with genetic inner ear disorders," *Nat. Commun.*, vol. 10, no. 1, p. 236, Jan. 2019, doi: 10.1038/s41467-018-08264-w.

[66] M. Chatzigeorgiou, S. Bang, S. W. Hwang, and W. R. Schafer, "tmc-1 encodes a sodium-sensitive channel required for salt chemosensation in C. elegans," *Nature*, vol. 494, no. 7435, pp. 95–99, Feb. 2013, doi: 10.1038/nature11845.

[67] Y. Honda *et al.*, "Genes down-regulated in spaceflight are involved in the control of longevity in *Caenorhabditis elegans*," *Sci. Rep.*, vol. 2, p. 487, Jul. 2012, doi: 10.1038/srep00487.

[68] F. Selch *et al.*, "Genomic response of the nematode Caenorhabditis elegans to spaceflight," *Adv. Space Res. Off. J. Comm. Space Res. COSPAR*, vol. 41, no. 5, pp. 807–815, 2008, doi: 10.1016/j.asr.2007.11.015.

[69] L. Zhao, Q. Rui, and D. Wang, "Molecular basis for oxidative stress induced by simulated microgravity in nematode Caenorhabditis elegans," *Sci. Total Environ.*, vol. 607–608, pp. 1381–1390, Dec. 2017, doi: 10.1016/j.scitotenv.2017.07.088.

[70] L. Qiao, S. Luo, Y. Liu, X. Li, G. Wang, and Z. Huang, "Reproductive and Locomotory Capacities of Caenorhabditis elegans Were Not Affected by Simulated Variable Gravities and Spaceflight During the Shenzhou-8 Mission," *Astrobiology*, vol. 13, no. 7, pp. 617–625, Jul. 2013, doi: 10.1089/ast.2012.0962.

[71] Y. Honda, S. Honda, M. Narici, and N. J. Szewczyk, "Spaceflight and Ageing: Reflecting on Caenorhabditis elegans in Space," *Gerontology*, vol. 60, no. 2, pp. 138–142, 2014, doi: 10.1159/000354772.

[72] B. Fritzsch, B. J. Kopecky, and J. S. Duncan, "Chapter 12 - Development of the Mammalian 'Vestibular' System: Evolution of Form to Detect Angular and Gravity Acceleration," in *Development of Auditory and Vestibular Systems*, R. Romand and I. Varela-Nieto, Eds. San Diego: Academic Press, 2014, pp. 339–367. doi: 10.1016/B978-0-12-408088-1.00012-9.

[73] M. Lipovsek and R. J. Wingate, "Conserved and divergent development of brainstem vestibular and auditory nuclei," *eLife*, vol. 7, p. e40232, Dec. 2018, doi: 10.7554/eLife.40232.

[74] D. I. Scheffer, J. Shen, D. P. Corey, and Z.-Y. Chen, "Gene Expression by Mouse Inner Ear Hair Cells during Development," *J. Neurosci.*, vol. 35, no. 16, pp. 6366–6380, Apr. 2015, doi: 10.1523/JNEUROSCI.5126-14.2015.

[75] D. P. Corey *et al.*, "TRPA1 is a candidate for the mechanosensitive transduction channel of vertebrate hair cells," *Nature*, vol. 432, no. 7018, Art. no. 7018, Dec. 2004, doi: 10.1038/nature03066.

[76] V. Gallo *et al.*, "TRPA1 gene polymorphisms and childhood asthma," *Pediatr. Allergy Immunol.*, vol. 28, no. 2, pp. 191–198, 2017, doi: https://doi.org/10.1111/pai.12673.

[77] A. Koivisto *et al.*, "TRPA1: A Transducer and Amplifier of Pain and Inflammation," *Basic Clin. Pharmacol. Toxicol.*, vol. 114, no. 1, pp. 50–55, 2014, doi: https://doi.org/10.1111/bcpt.12138.

[78] A. J. Iliff *et al.*, "The nematode C. elegans senses airborne sound," *Neuron*, vol. 109, no. 22, pp. 3633-3646.e7, Nov. 2021, doi: 10.1016/j.neuron.2021.08.035.

[79] Z. Altun, "Individual Neuron List," *WormAtlas*. https://www.wormatlas.org/neurons/Individual%20Neurons/Neuronframeset.html (accessed Mar. 03, 2022).

[80] L. Emtage, G. Gu, E. Hartwieg, and M. Chalfie, "Extracellular Proteins Organize the Mechanosensory Channel Complex in C. elegans Touch Receptor Neurons," *Neuron*, vol. 44, no. 5, pp. 795–807, Dec. 2004, doi: 10.1016/j.neuron.2004.11.010.

[81] M. B. Goodman and E. M. Schwarz, "Transducing touch in Caenorhabditis elegans," *Annu. Rev. Physiol.*, vol. 65, pp. 429–452, 2003, doi: 10.1146/annurev.physiol.65.092101.142659.

[82] X. Chen, M. D. Cuadros, and M. Chalfie, "Identification of Nonviable Genes Affecting Touch Sensitivity in Caenorhabditis elegans Using Neuronally Enhanced Feeding RNA Interference," *G3 GenesGenomesGenetics*, vol. 5, no. 3, pp. 467–475, Jan. 2015, doi: 10.1534/g3.114.015776.

[83] A. L. Brown, Z. Liao, and M. B. Goodman, "MEC-2 and MEC-6 in the Caenorhabditis elegans Sensory Mechanotransduction Complex: Auxiliary Subunits that Enable Channel Activity," *J. Gen. Physiol.*, vol. 131, no. 6, pp. 605–616, Jun. 2008, doi: 10.1085/jgp.200709910.

[84] N. Tavernarakis and and M. Driscoll, "Molecular Modeling of Mechanotransduction in the Nematode Caenorhabditis Elegans," *Annu. Rev. Physiol.*, vol. 59, no. 1, pp. 659–689, 1997, doi: 10.1146/annurev.physiol.59.1.659.

[85] T. Stiernagle, "Maintenance of C. elegans," *WormBook*, 2006, doi: 10.1895/wormbook.1.101.1.

[86] D. Fay, "Genetic mapping and manipulation: Chapter 1-Introduction and basics," *WormBook*, 2006, doi: 10.1895/wormbook.1.90.1.

[87] X. Karp, "Working with dauer larvae," *WormBook*, pp. 1–19, Aug. 2018, doi: 10.1895/wormbook.1.180.1.

Figure 1. Behavioral model, wiring diagram, and gravitaxis assay design

A

LIGHT

GRAVITY

Environmental Stimuli:
- Light
- Gravity
- Temperature
- Moisture/Humidity
- Geomagnetic Field
- Odorants
- Oxygen
- Pressure
- Sound
- Texture

C

50 cm

+7
+6
+5
+4
+3
+2
+1
-1
-2
-3
-4
-5
-6
-7

12-24 hrs

B

ALM(L)

PVD(L)

FLP(L)

AVM

PVM

PLM(L)

TRNs
- gentle touch responsive
- *mec-4, mec-10* +

FLPs/PVDs
- proprioceptive, multimodal
- *mec-10* +
- *mec-4* + (FLPs only)
- *trpa-1* + (PVDs only)

Figure 2. *C. elegans* negatively gravitax under different environmental conditions and developmental stages

Figure 3. DEG/ENaC components

Figure 4. MEC-7/12 microtubule subunits are necessary for gravitaxis in contrast to other DEG/ENaC components

A

B

Figure 6. PVD and TRN neurons in adult and dauer worms

mec-3p::RFP + Podr1::RFP (TRNs + AWC) ser-2prom-3::myrGFP (PVD + OLL)

DIC

Figure 7. Genetic ablation of PVD but not TRNs affects gravitaxis

Figure Legends

Figure 1. Behavioral model, wiring diagram, and gravitaxis assay design

(A) Graphic depicting environmental stimuli that inform *C. elegans* behavior in a natural setting. **(B)** Diagram of key neurons mentioned in this study. TRN neurons are labeled in red, while FLP and PVD neurons are green (for neurons with left and right pairs, only the left neuron is displayed). Adapted from individual neuron illustrations available on WormAtlas[79]. **(C)** Gravitaxis chamber and experimental design used in this study. Figures **(B)** and **(C)** created with BioRender.com.

Figure 2. *C. elegans* negatively gravitax under different environmental conditions and developmental stages

(A) Histograms depicting cumulative distribution of *C. elegans* worms over multiple trials in vertical (blue) and horizontal (orange) assays. Presence or absence of normal overhead light and background EMF (no Faraday cage) are denoted with +/-. **(B-C)** Boxplots depicting data shown in A. Vertical conditions only compared in **C**. Dauers tested in a dark Faraday cage (yellow) are used in future comparisons. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; n.s. is not significant using Kruskal-Wallis followed by Dunn's test with Bonferroni correction. Notches on boxplots represent 95% confidence intervals; mean values are indicated with a diamond.

Figure 3. DEG/ENaC components

MEC-4 and MEC-10 subunits form a sodium channel that is required for gentle touch sensation in TRNs. MEC-1, -5, and -9 components of the extracellular matrix (ECM) provide connection points between touch receptors and the cuticle[80]. MEC-7/12 microtubules form 15 protofilament structures in the TRNs only; however, these β and α subunits (respectively) are found in other neuronal 11 protofilament structures[43]. MEC-2 proteins associate with both the Deg/ENaC channels and TRN microtubules and enhance channel activity[81]–[83]. (Adapted from images by *Tavernarakis et. al 1997* [84] and others). Created with BioRender.com.

Figure 4. MEC-7/12 microtubule subunits are necessary for gravitaxis in contrast to other DEG/ENaC components

(A-C) Dauers with mutations in DEG/ENaC genes tested compared with N2 dauers. **(A)** Histograms comparing horizontal (orange) and vertical (blue) distributions of each strain over multiple trials. **(B-C)** Boxplots summarizing data shown in **A**; vertical comparisons with N2 dauers shown in **C** (yellow boxplot)**.** * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; n.s. is not significant using Kruskall-Wallis followed by Dunn's test with Bonferroni correction. Notches on boxplots represent 95% confidence intervals; mean values are indicated with a diamond.

Figure 5. TRPA-1 channels are also required for gravitaxis

(A-B) *trpa-1* mutants in vertical (blue) and horizontal (assays) compared with N2 controls. * p < 0.05, ** p < 0.01, *** p <0.001 , **** p < 0.0001; n.s. is not significant using Kruskall-Wallis followed by Dunn's test with Bonferroni correction. Notches on boxplots represent 95% confidence intervals; mean values are indicated with a diamond.

Figure 6. PVD and TRN neurons in adult and dauer worms

Confocal fluorescent **(A, C, and E)** and DIC **(B, D, and F)** micrographs of *mec-3p::RFP* and *ser-2prom-3::GFP* expression labelling the TRN and AWC neurons and PVD and OLL neurons, respectively. **(A-B)** Wildtype adult hermaphrodites; 4° process indicated with white arrowhead in inset. **(C-D)** *mec-4(e1611)* gain of function adult hermaphrodite, in which TRNs are ablated. **(E-F)** *mec-4(e1611)* dauer worm showing development of 2° and 3° processes in inset. Scale bar 100µm, 50µm in insets.

Figure 7. Genetic ablation of PVD but not TRNs affects gravitaxis

(A-C) Gravitaxis behavior of dauers lacking TRNs only (*mec-4* gof), dauers lacking PVD neurons only (*ser2prom-3::deg-3*), and N2 dauer controls. * p < 0.05, ** p < 0.01, *** p <0.001 , **** p < 0.0001; n.s. is not significant using Kruskall-Wallis followed by Dunn's test with Bonferroni correction. Notches on boxplots represent 95% confidence intervals; mean values are indicated with a diamond.

Acknowledgements:

We acknowledge the use of the NRI-MCDB Microscopy Facility and the Resonant Scanning Confocal supported by the NSF MRI grant DBI-1625770.
Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).
We are grateful for the generous gifts of transgenic worms provided to us by the Treinin Lab (Hebrew University), Chalfie Lab (Columbia University), and Shen Lab (Stanford University).

Declaration of Interests:

The authors declare no competing interests.