Stabilization of recombinant proteins by fusion with heat-resistant obscure (Hero) proteins

Eri Morimoto[1,2], Kotaro Tsuboyama[2,3], Yukihide Tomari[1,2,†]

[1]Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-0032, Japan.

[2]Laboratory of RNA Function, Institute for Quantitative Biosciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-0032, Japan.

[3]Department of Pharmacology, Northwestern Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA

[†]Corresponding author and lead contact. Email: tomari@iqb.u-tokyo.ac.jp (Y.T.)

Abstract

Although recombinant proteins are widely used in biotechnology and pharmaceutical industries, improving their solubility and stability is often a challenging issue. We recently discovered a class of highly unstructured heat-resistant obscure (Hero) proteins, which function to protect other "client" proteins *in trans* from various stresses *in vitro* and *in vivo*. Here, we show that Hero proteins can enhance the stability of recombinant proteins when fused *in cis*. Fusion with Hero proteins improved the otherwise challenging production of TAR DNA-binding protein of 43 kDa (TDP-43) in *Escherichia coli*. Moreover, fusing with Hero proteins strongly protected the activity of firefly luciferase bearing destabilizing mutations under various stress conditions. These data suggest that Hero proteins can be used as versatile stabilization tags for recombinant protein production.

Introduction

Recombinant proteins have been widely used in biotechnology and pharmaceutical

industries (1). *Escherichia coli* (*E. coli*) is one of the most common hosts to produce

recombinant proteins with high yield and low cost. However, overexpressed proteins in

E. coli often accumulate in inclusion bodies due to improper folding (2,3). To overcome

this limitation, fusion tags such as glutathione-S-transferase (GST) and maltose binding

protein (MBP) are frequently used. While helpful in increasing the solubility, GST forms

a homodimer in solution, which makes it unsuitable for oligomeric proteins (4,5). MBP

can also improve the solubility of tagged proteins, but MBP itself is a protein of ~42.5

kDa and this large size may increase the complexity in protein production and

downstream processes (6,7).

Stability of recombinant proteins after purification is also crucial for their applications.

Proteins are generally prone to denaturation especially under stress conditions such as

heat and freeze-thaw cycles. Maltodextrin-binding protein from *Pyrococcus furiosus* m

(pfMBP) and RNase HI from *Sulfolobus tokodaii* (Sto-RNase HI) are known to increase

not only the solubility but also the thermostability of recombinant proteins (8,9).

However, while "stabilization tags" are in high demand, they remain poorly explored.

We have previously reported that heat-resistant obscure (Hero) proteins, which are

heat-soluble, hydrophilic, highly charged, and widespread in animals, including humans.

Hero proteins can stabilize various "client" proteins *in vitro* and *in vivo* (10). For

example, Hero proteins can protect the enzymatic activity of lactate dehydrogenase

(LDH) from desiccation *in vitro* or that of firefly luciferase (Fluc) from heat shock in

HEK293T cells. Moreover, Hero proteins can prevent amyotrophic lateral sclerosis

(ALS)-associated pathogenic protein aggregations of TAR DNA-binding protein of 43

kDa (TDP-43) in cultured motor neurons and in *Drosophila* models for

neurodegenerative diseases (10). In light of these stabilization effects previously

demonstrated *in trans*, we hypothesized that Hero proteins may help protecting other

proteins of interest *in cis*.

Here we show that, indeed, recombinant proteins can be significantly stabilized by

fusion with Hero proteins. Fusing with Hero proteins improved the otherwise

challenging production of recombinant TDP-43 in *E. coli*. Moreover, fusion with Hero

strongly protected the enzymatic activity of Fluc bearing destabilizing mutations under

stress conditions such as heat, freeze-thaw cycles, and protease treatment. These

data suggest that Hero proteins have the potential as stabilization tags for recombinant

proteins.

Results

Hero tags can improve the protein expression of TDP-43 in *E. coli*.

TDP-43 is intrinsically aggregation-prone, and it is generally difficult to produce TDP-43

as a recombinant protein in *E. coli* (11). Based on our previous observation that Hero

proteins can suppress aggregation of TDP-43 *in trans* in human cells (10), we

wondered if Hero proteins can be used as fusion tags to increase the protein solubility

in cis in the *E. coli* expression system. We constructed a series of expression vectors,

in which TDP-43 was N-terminally tagged with His-FLAG and each of 6 representative

human Hero proteins or GST as a control, or His-FLAG alone (Fig. 1A). After protein

expression, we separated the soluble and insoluble fractions and analyzed the soluble

fraction by capillary-based quantitative Western blotting using anti-FLAG antibody. As

previously reported (11), TDP-43 was mostly found in the insoluble fraction (Fig. 1B),

and many incomplete peptides and/or degradation products were detected in the

soluble fraction (Fig. 1C), highlighting the difficulty of recombinant TDP-43 production in

E. coli. Compared to the His-FLAG alone (no tag), fusion with GST decreased the

protein yield of full-length TDP-43 in the soluble fraction, whereas fusion with Hero9, 11

and 20 did not compromise or slightly increased the yield (Fig. 1D). Importantly,

Hero11-tagging significantly improved the integrity of soluble TDP-43, with much less

degradation products compared to other tags (Fig. 1E). We concluded that fusion with

some Hero proteins has the potential to improve the otherwise challenging protein

expression of TDP-43 in *E. coli*.

Hero tags mitigate the loss of Fluc activity by heat.

We have previously demonstrated that co-expression of Hero proteins in HEK293T

cells mitigate the loss of Fluc activity by heat shock (10). To evaluate the protective

effect *in cis in vitro*, we expressed Hero or GST-tagged Fluc in *E. coli* using the same

expression constructs as our TDP-43 experiment above. Fluc is widely used as a

bioluminescent reporter in various species including *E. coli,* and as expected it was

expressed in the soluble fraction even without Hero or GST tag (Fig. 2A). Because

protein yields with Hero13 and 45 were extremely low, we excluded them from further

experiments. We purified the series of recombinant tagged Fluc proteins and adjusted

their concentrations (Fig. 2B). After confirming that the luminescence activities are

roughly comparable among all the samples (Fig. S1A), we exposed them to heat (Fig.

2C) by incubating them at 33 °C (Fig. 2D) or 37 °C (Fig. 2E) for 20 min. We then

measured the luminescence and calculated the loss of the enzymatic activity by heat

incubation. Except for Hero11, all tags protected the Fluc activity from heat but only

modestly (Fig. 2D and E).

Because wild-type (WT) Fluc is an intrinsically stable protein, fusion tags may have

only little space to improve its stability. It is known that the R188Q single-mutation (SM)

and the R188Q/R261Q double-mutation (DM) can strongly destabilize Fluc at high

temperatures (≥ 25 °C), without severely compromising the enzymatic activity at 20 °C

(12). Therefore, we repeated the heat stress test using Fluc-SM (Fig. 3A–D) and DM

(Fig. 3A and E–G). Fusion with Hero9 or Hero20 significantly protected the activity of

Fluc-SM at both 33 °C and 37 °C (Fig. 3C and D), whereas GST tag did not show any

apparent protection. For Fluc-DM, Hero9 was particularly effective in protecting the

enzymatic activity even at 37 °C (Fig. 3F and G). We concluded that tagging with some

Hero proteins can mitigate the destabilization of proteins by heat, especially for

intrinsically unstable ones.

Hero tags protect Fluc activity from freeze-thaw cycles.

In general, proteins tend to be denatured by freezing and thawing (13). Indeed, 2

cycles of freezing and thawing strongly compromised the Fluc-WT, SM and DM activity

(Fig. 4). Although GST mildly protected the Fluc activity from the freeze-thaw cycles,

Hero9 and Hero20 showed superior protection activity for Fluc-SM and Fluc-DM (Fig.

4B–D). These data suggest that Hero tags can be used to prevent the loss of function

via freeze-thaw cycles (Fig. 4B–D).

Hero tags protect Fluc activity from Proteinase K.

Since proteins are generally prone to degradation by proteases both *in vivo* and *in vitro*,

we tested if Hero tags can protect client proteins from Proteinase K (PK), a

representative serine protease. We incubated the series of Fluc-WT, SM, and DM

proteins with PK for 30 min and measured their luminescence activity (Fig. 5). While

GST showed no shielding effect, Hero9 strongly protected the activity of Fluc-SM and

DM from PK-mediated proteolysis (Fig. 5C and D). Thus, fusion with Hero proteins is a

useful strategy to confer increased protease resistance on client proteins (14).

Hero proteins protect Fluc activity better in *cis* than in *trans*.

Finally, we compared the protective effect of Hero proteins *in cis* and *in trans*. We

purified recombinant GST, Hero9 and Hero11 proteins (which were most effective *in*

cis), added each of them to the "no tag" Fluc protein in equimolar concentrations, and

challenged the mixture with heat, freeze-thaw cycles, or PK. As shown in Fig. S2A–I,

the protective effect was only minimum in all the stress conditions. We concluded that

fusing with Hero proteins *in cis* works better than mixing with them *in trans*, at least to

protect the Fluc activity *in vitro*.

Discussion

In this study, we demonstrated that Hero proteins can be used as useful fusion tags

that can protect recombinant proteins under various stress conditions. We envision that

Hero tags act as a physical shield that prevents collisions of molecules leading to

denaturation. Hero tags may be also helpful in promoting and maintaining the proper

folding of client proteins by improving the molecular environment. In this sense, Hero

may be reminiscent of polyethylene glycol (PEG), a post-production modification

commonly used to increase the solubility and stability for biopharmaceuticals (15,16). It

is known that PEG itself can show immunogenicity albeit rarely, and the long-term

toxicity of PEG-modified products has recently been cautioned (17,18). To overcome

this problem, researchers have developed "PAS," an artificial polypeptide of defined

sequence containing the 3 small amino acids Pro, Ala, and Ser. PAS is biodegradable

and non-immunogenic, yet improves solubility in *E. coli* and protein half-lives *in vivo*

(19). Hero proteins may resemble PAS, except that they are from natural sources. In

our current study, we tested only 6 representative Hero proteins, but the human

genome encodes hundreds of Hero protein candidates, many of which remain to be

characterized (10). Thus, it is possible that there are Hero proteins that act as better

stabilization tags than those tested in this study. Moreover, it will be interesting to

examine in the future if tandem repeats of Hero proteins may increase the stabilization

effect, as was demonstrated for PAS (20).

Toward the application of Hero tags, it is important to note that different Hero

proteins have different preferences for their client proteins. For example, in our

previous research, Hero7 and 11 showed strong resistance to heat shock when co-

expressed with Fluc *in trans* in cultured human cells (10). However, the *in-trans*

protective effect of Hero11 was only minimum *in vitro* (Fig. 2S). When fused *in cis*,

Hero11 rather abolished the Fluc activity *in vitro*, while Hero9 showed the strongest

protection (Fig. 3–4). On the other hand, Hero11 strongly improved the integrity of

soluble TDP-43 in the *E. coli* expression system (Fig. 1C–E). Those data suggest that

even the same Hero protein can behave differently depending on the client protein and

condition. Unfortunately, it is currently difficult to predict the best combination between

Hero proteins and clients, and it will be important to test multiple Hero proteins to

identify one that best protects the protein of interest.

In summary, our current study provides the potential of Hero proteins as versatile

stabilization tags for recombinant proteins and serve as a starting point for further

optimization and engineering.

Figure legends

Figure 1. Hero proteins improve solubility and stability of TDP-43 in *E. coli.*

(A) Schematic representation of Hero-tagged protein expression constructs.

(B) Expression of tagged TDP-43 proteins. TDP-43 fused with GST, each Hero protein

(Hero7, 9, 11, 13, 20, or 45) or no tag was expressed in *E. coli*, and the soluble and

insoluble fractions were analyzed by SDS-PAGE. Expected sizes are 50, 62, 57, 59,

60, 62, 70, and 92 kDa for no tag, GST, Hero7, Hero9, Hero11, Hero13, Hero20,

and Hero 45, respectively. A representative image from 3 independent experiments

is shown. Note that extreme biases in amino acid composition of Hero proteins can

affect the mobility of protein bands on SDS-PAGE (10).

(C) Capillary-based quantitative Western blotting of the soluble fractions in (B). A

representative image from 3 independent experiments is shown. The bands marked

with red arrows represent the full-length proteins.

(D) Relative quantification of the full-length proteins compared to no tag in (C). Mean ±

SD from 3 independent experiments are shown. *P*-values were calculated by the

Steel-Dwass test against no tag. None showed $p < 0.05$.

(E) Quantification of the fraction of the full-length proteins out of the total proteins in (C).

Mean ± SD from 3 independent experiments are shown. Fusion with Hero11

resulted in the highest purity. *P*-values were calculated by Tukey HSD against no

tag. *Hero11: $p = 0.01$.

Figure 2. Hero tags slightly mitigate the loss of Fluc activity by heat.

(A) Expression of tagged Fluc proteins. Fluc fused with GST, each Hero protein (Hero7,

9, 11, or 20) or no tag was expressed in *E. coli*, and the soluble and insoluble

fractions were analyzed by SDS-PAGE.

(B) SDS-PAGE of His-purified Fluc proteins.

(C) Schematic representation of the heat treatment.

(D) Heat treatment of Fluc at 33°C for 20 min. Fractions of the remaining Fluc activities

after the heat incubation were calculated. Mean ± SD from 4 independent

experiments are shown. *P*-values were calculated by Tukey HSD against no tag.

None showed $p < 0.05$.

(E) Heat treatment of Fluc at 37°C for 20 min. Fractions of the remaining Fluc activities

after the heat incubation were calculated. Mean ± SD from 4 independent

experiments are shown. *P*-values were calculated by Tukey HSD against no tag.

*Hero20: $p = 0.0148$.

Figure 3. Hero tags markedly mitigate the loss of mutated Fluc activity by heat.

(A) Schematic representation of the heat treatment.

(B) SDS-PAGE of His-purified Fluc-SM proteins.

(C) Heat treatment of Fluc-SM at 33 °C for 20 min. Fractions of remaining Fluc
activities after the heat incubation were calculated. Mean ± SD from 6 independent
experiments are shown. *P*-values were calculated by Tukey HSD against no tag.
*Hero9: $p = 0.0038$, Hero11: $p = 0.0455$.

(D) Heat treatment of Fluc-SM at 37 °C for 20 min. Fractions of remaining Fluc
activities after the heat incubation were calculated. Mean ± SD from 6 independent
experiments are shown. *P*-values were calculated by Tukey HSD against no tag.
*Hero9: $p = 0.0017$, Hero11: $p = 0.007$, Hero20: $p = 0.001$.

(E) SDS-PAGE of His-purified Fluc-DM proteins.

(F) Heat treatment of Fluc-DM at 33 °C for 20 min. Fractions of remaining Fluc
activities after the heat incubation were calculated. Mean ± SD from 5 independent
experiments are shown. *P*-values were calculated by Tukey HSD against no tag.
*Hero9: $p = 0.0073$.

(G) Heat treatment of Fluc-DM at 37 °C for 10 min. Fractions of remaining Fluc

activities after the heat incubation were calculated. Mean ± SD from 5 independent

experiments are shown. *P*-values were calculated by Tukey HSD against no tag.

*Hero9: $p = 0.001$.

Figure 4. Hero tags protect Fluc activity from freeze-thaw cycles.

(A) Schematic representation of the freeze-thaw cycles.

(B) Freeze-thaw cycles of Fluc-WT. Fractions of remaining Fluc activities after the

second cycle were calculated. Mean ± SD from 5 independent experiments are

shown. *P*-values were calculated by Tukey HSD against no tag. *Hero11: $p = 0.045$,

Hero20: $p = 0.0344$.

(C) Freeze-thaw cycles of Fluc-SM. Fractions of remaining Fluc activities after the

second cycle were calculated. Mean ± SD from 6 independent experiments are

shown. *P*-values were calculated by Tukey HSD against no tag. *GST: $p = 0.014$,

Hero7: $p = 0.0309$, Hero9: $p = 0.001$, Hero20: $p = 0.0018$.

(D) Freeze-thaw cycles of Fluc-DM. Fractions of remaining Fluc activities after the

second cycle were calculated. Mean ± SD from 6 independent experiments are

shown. *P*-values were calculated by Tukey HSD against no tag. *Hero9: $p = 0.0022$,

Hero11: $p = 0.0074$, Hero20: $p = 0.001$.

Figure 5. Hero tags protect Fluc activity from Proteinase K.

(A) Schematic representation of the PK treatment.

(B) PK treatment of Fluc-WT for 30 min. Fractions of remaining Fluc activities after the

PK treatment were calculated. Mean ± SD from 4 independent experiments. *P*-

values were calculated by Tukey HSD against no tag. None showed $p < 0.05$.

(C) PK treatment of Fluc-SM for 30 min. Fractions of remaining Fluc activities after the

PK treatment were calculated. Mean ± SD from 6 independent experiments. *P*-

values were calculated by Tukey HSD against no tag. *Hero9: $p = 0.0051$.

(D) PK treatment of Fluc-WT for 30 min. Fractions of remaining Fluc activities after the

PK treatment were calculated. Mean ± SD from 6 independent experiments. *P*-

values were calculated by Tukey HSD against no tag. *Hero9: $p = 0.001$.

Figure S1. Basal Fluc activities are comparable with and without fusion tags.

(A) Basal Fluc activities before the stress tests in Figure 2. *P*-values were calculated by

the Steel-Dwass test against no tag. None showed $p < 0.05$.

Figure S2. Fluc activities are only minimally protected by Hero proteins *in trans*.

(A) Heat treatment of Fluc-WT mixed with buffer alone, GST, Hero9 or Hero11 at 37 °C

for 20 min.

(B) Heat treatment of Fluc-SM mixed with buffer alone, GST, Hero9 or Hero11 at 37 °C

for 20 min.

(C) Heat treatment of Fluc-DM mixed with buffer alone, GST, Hero9 or Hero11 at 37 °C

for 10 min.

(D) Freeze-thaw cycles of Fluc-WT mixed with buffer alone, GST, Hero9 or Hero11.

(E) Freeze-thaw cycles of Fluc-SM mixed with buffer alone, GST, Hero9 or Hero11.

(F) Freeze-thaw cycles of Fluc-DM mixed with buffer alone, GST, Hero9 or Hero11.

(G) PK treatment of Fluc-WT mixed with buffer alone, GST, Hero9 or Hero11 for 30 min.

(H) PK treatment of Fluc-SM mixed with buffer alone, GST, Hero9 or Hero11 for 30 min.

(I) PK treatment of Fluc-DM mixed with buffer alone, GST, Hero9 or Hero11 for 30 min.

For all the data, p-values were calculated by Tukey HSD against no tag. None showed

$p < 0.05$.

Acknowledgements

We are grateful to Andy Y. Lam for providing recombinant proteins of GST, Hero9 and

Hero11. This work was supported in part by JSPS KAKENHI Grant Number

JP18H05271 and JP21H05278 (to Y.T.).

Competing interests

Y.T. and K.T. have a patent application related to this work. E.M. declares no competing

interests.

Author Contributions

E.M., K.T., and Y.T. designed the project. E.M. performed all the experiments. K.T. and

Y.T. supervised the project. E.M., K.T., and Y.T. wrote the manuscript.

References:

1. Biologics Market _ 2021 - 26 _ Industry Share, Size, Growth - Mordor Intelligence.

2. Chrunyk BA, Evans J, Lillquist J, Young P, Wetzel R. Inclusion body formation and protein stability in sequence variants of interleukin-1β. Journal of Biological Chemistry. 1993;268(24):18053–61.

3. Williams DC, van Frank RM, Muth WL, Burnett JP. Cytoplasmic Inclusion Bodies in Escherichia coli Producing Biosynthetic Human Insulin Proteins. Vol. 215, New Series. 1982.

4. Veal EA, Mark Toone W, Jones N, Morgan BA. Distinct roles for glutathione S-transferases in the oxidative stress response in Schizosaccharomyces pombe. Journal of Biological Chemistry. 2002 Sep 20;277(38):35523–31.

5. Kaplan W, Erhardt J, Sluis-Cremer N, Dirr H, Hüsler P, Klump H. Conformational stability of pGEX-expressed *Schistosoma japonicum* glutathione S-transferase: A detoxification enzyme and fusion-protein affinity tag. Protein Science. 1997 Feb;6(2).

6. Kapust RB, Waugh DS. Escherichia coli maltose-binding protein is uncommonly effective at promoting the solubility of polypeptides to which it is fused (1999).

7.   Dyson MR, Shadbolt SP, Vincent KJ, Perera RL, McCafferty J. Production of

soluble mammalian proteins in Escherichia coli: Identification of protein features

that correlate with successful expression. BMC Biotechnology. 2004 Dec 14;4.

8.   Luke JM, Carnes AE, Sun P, Hodgson CP, Waugh DS, Williams JA.

Thermostable tag (TST) protein expression system: Engineering thermotolerant

recombinant proteins and vaccines. Journal of Biotechnology. 2011 Feb

10;151(3):242–50.

9.   Takano K, Okamoto T, Okada J, Tanaka SI, Angkawidjaja C, Koga Y, et al.

Stabilization by fusion to the C-terminus of hyperthermophile Sulfolobus tokodaii

RNase HI: A possibility of protein stabilization tag. PLoS ONE. 2011;6(1).

10. Tsuboyama K, Kozuka-Hata H, Oyama M, Iwasaki S, Tomari Y. A widespread

family of heat-resistant obscure (Hero) proteins protect against protein instability

and aggregation.

11. Capitini C, Conti S, Perni M, Guidi F, Cascella R, de Poli A, et al. TDP-43

inclusion bodies formed in bacteria are structurally amorphous, non-amyloid and

inherently toxic to neuroblastoma cells. PLoS ONE. 2014 Jan 30;9(1).

12. Gupta R, Kasturi P, Bracher A, Loew C, Zheng M, Villella A, et al. Firefly

luciferase mutants as sensors of proteome stress. Nature Methods. 2011

Oct;8(10):879–84.

13.   Zhang A, Qi W, Singh SK, Fernandez EJ. A new approach to explore the impact

of freeze-thaw cycling on protein structure: Hydrogen/deuterium exchange mass

spectrometry (HX-MS). Pharmaceutical Research. 2011 May;28(5):1179–93.

14.   Markert Y, Köditz J, Mansfeld J, Arnold U, Ulbrich-Hofmann R. Increased

proteolytic resistance of ribonuclease A by protein engineering. Vol. 14, Protein

Engineering. 2001.

15.   Milton Harris J, Chess RB. Effect of pegylation on pharmaceuticals. Vol. 2,

Nature Reviews Drug Discovery. 2003. p. 214–21.

16.   Abuchowski A, van Es T, Palczuk NC, Davis FF. Alteration of immunological

properties of bovine serum albumin by covalent attachment of polyethylene

glycol. Journal of Biological Chemistry. 1977;252(11):3578–81.

17.   Sellaturay P, Nasser S, Islam S, Gurugama P, Ewan PW. Polyethylene glycol

(PEG) is a cause of anaphylaxis to the Pfizer/BioNTech mRNA COVID-19

vaccine. Vol. 51, Clinical and Experimental Allergy. Blackwell Publishing Ltd;

2021. p. 861–3.

18.   Fletcher AM, Tellier P, Douville J, Mansell P, Graziano MJ, Mangipudy RS, et al.

Adverse vacuolation in multiple tissues in cynomolgus monkeys following repeat-

dose administration of a PEGylated protein. Toxicology Letters. 2019 Dec

15;317:120–9.

19.   Schlapschy M, Binder U, Börger C, Theobald I, Wachinger K, Kisling S, et al.

PASylation: A biological alternative to PEGylation for extending the plasma half-

life of pharmaceutically active proteins. Protein Engineering, Design and

Selection. 2013 Aug;26(8):489–501.

20.   Binder U, Skerra A. PASylation®: A versatile technology to extend drug delivery.

Vol. 31, Current Opinion in Colloid and Interface Science. Elsevier Ltd; 2017. p.

10–7.

Experimental Procedures

Plasmid construction

A DNA fragment containing FLAG-tag and GST, Hero7, 9, 11, 13, 20, or 45 (SERF2,

C9orf16, C19orf53, C11orf58, BEX3, or SERBP1 respectively) was inserted into pCold

I (Takara), together with the client protein, TDP-43, Fluc (WT), Fluc (SM; R188Q) or

Fluc (DM; R188Q, R261Q) (12).

Protein purification

Recombinant tagged TDP-43 and Fluc proteins were expressed in *E. coli* BL21 strain.

The cells were cultivated in 6 or 13 mL for TDP-43 and in 250 mL for Fluc to an OD600

of 0.4–0.6 at 37 °C, and then grown at 15 °C overnight with 1 mM isopropyl-β-D-

thiogalactoside (IPTG) following cold-shock on ice for 20 min. For TDP-43, the cells

were resuspended in lysis buffer [200 mM HEPES-KOH pH7.4, 200 mM KOAc, and

200 mM Mg(OAc)$_2$] supplemented with 0.2 mM TCEP, EDTA-free protease inhibitor

cocktail (Roche), and DNase I, sonicated, and centrifuged at 10,000 x g for 10 min. The

pellets were resuspended and sonicated again and the soluble and insoluble fractions

were analyzed by SDS-PAGE and capillary-based Western blotting. For Fluc, the cells

were resuspended in His A buffer [30 mM HEPES-KOH (pH 7.4), 200 mM KOAc, 5%

glycerol] supplemented with EDTA-free protease inhibitor cocktail (Roche), sonicated,

and centrifuged at 10,000 x g for 5 min. The supernatant was added to a slurry of

cOmplete His-Tag Purification Resin (Roche) or Ni Sepharose High Performance

(Merck) and eluted with His B buffer (His A buffer containing 400 mM imidazole). The

eluates were mixed with 20 % Glycerol and 1 mM DTT, snap-frozen by liquid-N_2 and

stored at –80 °C.

Capillary-based Western blotting

Samples were prepared and analyzed by Jess according to the manufacturer's

instruction (Protein Simple). Anti-DDDDK antibody was used as the primary antibody at

1:100 (M185, MBL). Anti-mouse antibody was used as the secondary antibody at 1:100

(Protein Simple).

Stress conditions

High temperature

For the *in-cis* experiments (Fig. 2 and 3), 40 μL of tagged Fluc proteins (~40 nM) were

incubated at 33 °C and 37 °C for 20 min, except that the 37 °C incubation for Fluc-DM

was for 10 min. For the *in-trans* experiments (Fig. S2A–C), Fluc and GST, Hero9,

Hero11, or lysis buffer were mixed (final concentrations ~40 nM) in 40 uL and incubated

at 37 °C for 20 min (WT and SM) or 10 min (DM).

Freeze and thaw cycles

For the *in-cis* experiments, 80 μL of tagged Fluc proteins (~40 nM) were frozen at −

80 °C for 30 min and thawed at room temperature for 10 min. This cycle was repeated

twice. For the *in-trans* experiments, Fluc and GST, Hero9, Hero11, or lysis buffer were

mixed (final concentrations ~40 nM) in 80 uL, frozen, and thawed twice.

Proteinase K treatment

For the *in-cis* experiments, 40 μL of tagged Fluc proteins (~40 nM) were incubated with

10 uL of Proteinase K (0.06 U/mL) for 30 min on ice. For the *in-trans* experiments, each

Fluc and GST, Hero9, Hero11, or lysis buffer were mixed (final concentrations ~40 nM)

in 40 uL and incubated with 10 uL of Proteinase K (0.06 U/mL).

Luciferase assay

The luciferase activities of Fluc were measured before and after the stress treatment,

using sensilite Enhanced Flash Luminescence (Perkin Elmer) and SPARK 10 M plate

reader (TECAN). The fractions of the remaining activity were then calculated.

Figure 1

Figure 2

C

Fluc-WT

33°C or 37°C

Figure 3

Figure 4

Fluc-WT, SM, or DM

Figure 5

Fluc-WT, SM, or DM

Supplemental Figure 1

A

Supplemental Figure 2