ReLo is a simple and quick colocalization assay to identify and characterize direct protein-protein interactions

Harpreet Kaur Salgania[1], Jutta Metz[1], Mandy Jeske[1]*

[1] Heidelberg University Biochemistry Center (BZH), Im Neuenheimer Feld 328, 69120 Heidelberg, Germany

*Correspondence: jeske@bzh.uni-heidelberg.de

ABSTRACT

The characterization of protein-protein interactions (PPIs) is fundamental for understanding biochemical processes. Many methods have been established to identify and study direct PPIs; however, the screening and investigation of PPIs involving large or poorly soluble proteins remain challenging. As a result, we developed ReLo, a simple, rapid, and versatile cell culture-based method for detecting and investigating interactions in a cellular context. Importantly, our data strongly suggest that with ReLo specifically direct binary PPIs are detected. By applying additional bridging experiments ReLo can also be used to determine the binding topology of subunits within multiprotein complexes. Moreover, ReLo has the potential to identify protein domains that mediate complex formation, screen for interfering point mutations, study interactions that depend on conformation or protein arginine methylation, and it is sensitive to drugs that mediate or interfere with an interaction. Taken together, ReLo is a simple and quick alternative for the study of PPIs particularly when established methods fail.

1 INTRODUCTION

The identification and characterization of protein-protein interactions (PPIs) are a

routine laboratory practice and lay the foundation for understanding biological

processes. PPIs can be identified through various established, mass spectrometry-

coupled screening methods, including coimmunoprecipitation (co-IP), tandem affinity

purification, and proximity-dependent labeling approaches, such as BioID or APEX [1–

6]. Thus, a list of interacting protein candidates is obtained, and these candidates are

usually ranked according to their abundance in eluate fractions. Determining which of

these candidates are truly direct binding partners requires subsequent validation

experiments, often involving *in vitro* methods, such as GST pull-down assays, which

depend upon the availability of purified proteins. If the proteins are poorly soluble and

cannot be obtained through recombinant protein expression or if expertise in

recombinant protein expression and purification methods is lacking, PPIs can be

validated using cell-based assays.

Yeast two-hybrid (Y2H) and protein complementation assays (PCA) are well-

established techniques in which an interaction results in the reconstitution and

subsequent detection of a split reporter protein, such as a transcription factor, ubiquitin,

an enzyme, or a fluorescent protein [7–12]. However, standard Y2H and PCA assays are

not particularly suitable for the analysis of potentially unstable proteins, as they are

barely expressed or rapidly degraded in a cell, resulting in unreliable, mostly false-

negative results [7,10]. Therefore, to obtain conclusive findings, negative results require

additional assessment of the protein expression levels, which complicates the process,

particularly when probing many PPIs.

Cell-based PPI methods that are better suited for testing interactions involving

potentially unstable proteins are based on fluorescent protein tagging and

colocalization readouts and thus allow the simultaneous monitoring of both PPIs and

protein expression levels via fluorescence microscopy. The readout of these

colocalization assays is usually the translocation of one protein upon its association

with a second distinctly localized protein (e.g., localization to a membrane, the nucleus,

or granules). 'Cytoskeleton-based assay for protein-protein interaction' (CAPPI),

'membrane recruitment assay' (MeRA), and 'knocksideways in plants' (KSP) are

translocation assays developed for use with plant cells [13–15]. 'Nuclear translocation

assay' (NTA), 'emerging circle of interactive proteins at specific endosomes'

(ECLIPSE), and 'protein interactions from imaging of complexes after translocation'

(PICT) are assays that require the addition of a compound (e.g., rapamycin) to monitor

the translocation upon a PPI [16–18]. Other translocation assays are based on

oligomerization/aggregation readouts [19–21] and may thus not be suitable for studying

interactions with proteins that form granules on their own within a cell. Importantly,

none of the described translocation assays have been assessed based on their ability

to distinguish direct binary interaction from those where the two proteins tested are

eventually bridged by cell-endogenous proteins.

We developed a simple and fast translocation PPI assay named ReLo for use

with an animal cell culture. This assay is based on the relocalization of a protein upon

its interaction with a second membrane-anchored protein. We applied ReLo to many

large proteins, most of which harbor long disordered regions and are known to be

insoluble after recombinant protein expression experiments. Using this set of proteins,

we demonstrate that ReLo can be employed to identify and characterize PPIs.

Importantly, using two structurally well-characterized multidomain protein complexes,

we show that only direct interactions are detected using ReLo, a prerequisite for the

analysis of previously unknown protein complexes through *in vitro* and structural

biology methods. Furthermore, ReLo is responsive to drug treatment, which enables

the study of drug-induced interactions and the screening of small PPI inhibitors. In

summary, ReLo is a simple, rapid and versatile tool that enables a thorough initial

description of direct PPI networks.

RESULTS

ReLo: a simple and robust cell culture-based PPI assay

As a preparation for the ReLo assay, two proteins of interest were fused to a red

fluorescent protein (mCherry) and a green fluorescent protein (EGPF, mEGFP).

Importantly, one of the constructs carried an additional fusion to a membrane-

anchoring protein domain, which led to distinct subcellular membrane localization of

the fusion protein. Upon interaction, the second protein is expected to colocalize with

the anchored protein to the membrane and it would thereby relocalize with respect to

its original location (**Fig. 1a**). Thus, we refer to the assay as the 'relocalization PPI

assay', abbreviated 'ReLo'.

ReLo is based on a simple methodology in which cells are seeded onto 4-well

chambered coverslips and cotransfected with the desired combination of plasmids.

After 48 h, the protein localization is analyzed by live-cell confocal fluorescence

microscopy (**Fig. 1a**). We used S2R+ cells, which were derived from semiadherent

Schneider's-line-2 (S2) cells, which in turn have been established from late *Drosophila*

embryos [22,23]. Compared with S2 cells, S2R+ cells show greater adherence to dishes,

and thus they do not require coated dishes for their adhesion prior to live-cell

microscopy. All ReLo plasmids carry an in-frame blunt end restriction site, allowing the

application of a simple, fast, and straightforward cloning procedure (see Methods

section).

For the anchoring of cytoplasmic proteins of interest to a membrane, we

selected the pleckstrin homology (PH) domain of rat phospholipase Cδ_1 (PLCδ_1), which

specifically recognizes phosphatidylinositol 4,5-bisphosphate [24,25] and thus directs the

fusion construct to the plasma membrane of a cell (**Fig. 1a**). In the ReLo assay, the

membrane localization was independent of whether the PH domain was fused to the

N- or C-terminus of a protein (**Supplementary Fig. 1a**). Unfortunately, nuclear proteins

fused to the PH domain were only inefficiently retained within the cytoplasm and barely

localized to the plasma membrane (**Supplementary Fig. 1c**). Hence, the testing of

PPIs to a nuclear protein may result in false-negative outcomes if the interaction

partner resides in the cytoplasm. Therefore, we tested alternative membrane-

anchoring domains to assess their ability to retain nuclear proteins within the

cytoplasm. Hence, we selected the minimembrane protein subunit 4 of the yeast

oligosaccharyltransferase complex (OST4), which was previously used as N-terminal

fusion protein to localize a nuclear protein to the endoplasmic reticulum (ER) [26,27]

(**Supplementary Fig. 1d**). Because the plasma membrane localization of the PH

domain distinguishes from a ubiquitous cytoplasmic localization better than the ER

localization of OST4, we prefer using the PH domain in ReLo whenever possible.

PPI mapping and mutational analysis

As a proof of concept for interaction studies with ReLo, we tested a previously

characterized protein complex consisting of the extended LOTUS (eLOTUS) domain

of Oskar (Oskar 139-240) and the C-terminal RecA-like domain (CTD) of the ATP-

dependent DEAD-box RNA helicase Vasa (Vasa-CTD, Vasa 463-661) [28] (**Fig. 1b**).

Only the short isoform of Oskar (Short Oskar, aa 139-606) interacts with Vasa [29], and

therefore, only Short Oskar and its domains were used in the following experiments

(**Fig. 1b**). In our setup, the eLOTUS domain of Oskar was fused to PH-mCherry and

localized to the plasma membrane. Vasa-CTD was fused to EGFP and localized

ubiquitously within the cytoplasm and nucleus. When coexpressed with PH-mCherry-

eLOTUS but not with PH-mCherry alone, EGFP-Vasa-CTD relocalized to the plasma

membrane (**Fig. 1c**). The unstructured region of Vasa (Vasa 1-200) and the N-terminal

RecA-like domain (Vasa 200-463) did not interact with the Oskar-eLOTUS domain,

and similarly, neither the unstructured region of Oskar (Oskar 241-387) nor its OSK

domain (Oskar 388-606) interacted with Vasa-CTD (**Fig. 1c**). Surface point mutations

that had been previously shown to interfere with the Vasa-Oskar interaction [28] were

also found to be inhibitory in the ReLo assay (**Fig. 1d**). Together, these data confirmed

the specific interaction between Vasa-CTD and Oskar-eLOTUS and demonstrated that

ReLo can be applied to map PPIs and to identify mutations that interfere with PPIs.

Oskar-eLOTUS and Vasa-CTD form a transient complex characterized by a

dissociation constant (K_D) of ~10 μM, and although this complex has been crystallized,

it is not sufficiently stable to be detected by size-exclusion chromatography [28,30,31].

Nevertheless, the relocalization upon the interaction between Oskar-eLOTUS and

Vasa-CTD was clearly detectable in the ReLo assay: in a total of three independent

replicates, the relocalization of Vasa-CTD toward the plasma membrane was observed

in 91 of 94 (i.e., 97%) cotransfected cells with both red and green fluorescent signals

(**Supplementary Fig. 2**). These data indicate that relocalization is a highly frequent

event and that the assay is well suited for studying low-affinity complexes. Most other

interactions tested revealed relocalization in 100% of the cotransfected cells (see

below; **Supplementary Fig. 3**).

Conformation-dependent interactions

Previous data have suggested that the Oskar-Vasa interaction depends on the

conformation of Vasa [28]. We aimed to test the interaction between Oskar and different

Vasa conformations in a full-length protein context. However, in contrast to the

eLOTUS domain of Oskar, full-length Short Oskar did not localize to the cytoplasm but

exclusively localized to the nucleus in S2R+ cells (**Supplementary Fig. 1b)** [28].

Therefore, we made use of the OST4-mCherry-Oskar construct, which localized to

membranous structures within the cytoplasm (**Supplementary Fig. 1d and Fig. 1e**).

Upon coexpression, wild-type Vasa relocalized and colocalized with OST4-Oskar at

the ER (**Fig. 1e**), confirming the interaction between Vasa and Oskar in the ReLo

assay. Also using the OST4 anchor, the interaction mapped to the Oskar-eLOTUS

domain and the Vasa-CTD **(Supplementary Fig. 1e)** and the known Vasa interface

mutant (F504E) was unable to bind to Oskar (**Fig. 1e**), which is consistent with the

data obtained using the PH domain (**Fig. 1c, 1d**).

The cores of Vasa and other ATP-dependent DEAD-box RNA helicases are

composed of two RecA-like domains, which display different orientations relative to

each other depending on the presence of bound ATP and RNA [32]. In a substrate-

unbound form, the helicase core adopts an open conformation and closes upon

substrate binding (**Fig. 2a**). To assess the conformation-dependent Vasa interaction

with the ReLo assay, we used Vasa variants with well-characterized point mutations in

the ATP-binding pocket that stabilize either the open conformation (K295N; Vasa-

open) or the closed conformation (E400Q; Vasa-closed) [33–35]. When testing the

interaction of OST4-anchored Oskar with the Vasa mutants using the ReLo assay, we

observed an interaction with Vasa-open but not Vasa-closed (**Fig. 2b)**, which revealed

that Oskar prefers to bind to the open conformation of Vasa. These outcomes were

consistent with our previous observations [28] and demonstrated that the ReLo assay

allows the study of PPIs that are dependent on the specific conformation of an

interaction partner.

Protein arginine methylation-dependent interactions

Protein interactions may depend on posttranslational modifications, such as protein

arginine methylation. The symmetric dimethylated arginine (sDMA) modification is

catalyzed by a subset of protein arginine methyltransferases (PRMTs) [36], and sDMA

methylation activity has been previously reported in S2 cells [37]. To test whether the

ReLo assay is suitable for investigating interactions involving sDMA modifications in

S2R+ cells, we tested the previously characterized strictly sDMA-dependent

interaction between the PIWI protein Aubergine (Aub) and Tudor [38,39] (**Fig. 2c**). Using

ReLo, we indeed observed an Aub-Tudor interaction (**Fig. 2d**). Aub carries four sDMAs

within its RG-rich N-terminus (R11, R13, R15, and R17), which are specifically

recognized and bound by extended Tudor (eTud) domains of Tudor [38–41]. The

substitution of these four arginine residues with lysine residues (Aub R->K) rendered

Aub unmodifiable by PRMT5 [41] and abolished the Aub-Tudor interaction (**Fig. 2d**).

Together, these data demonstrated that the S2R+ cells exhibited sufficient sDMA

activity to effectively modify proteins expressed after transient transfection of the cells.

We conclude that ReLo is suitable for the study of PPIs that depend on PRMT5-

catalyzed sDMA modification.

Effect of small molecules on PPIs

We subsequently tested whether the ReLo assay can be used to study PPIs that are

induced by the addition of small molecules to the cell culture medium. To this end, we

tested the previously characterized rapamycin-dependent interaction between human

FK506-binding protein 12 (FKBP12) and the FKBP12-rapamycin-binding domain

(FRB) of human mTOR [42]. In the absence of rapamycin, using the dimethyl sulfoxide

(DMSO)-containing control medium, FKBP12 and FRB did not interact in the ReLo

assay but did interact in the presence of 100 nM rapamycin in the cell culture medium

(**Fig. 2e**).

Furthermore, we tested whether the ReLo assay can be used to inhibit PPIs

through drug treatment. We opted to interfere with the interaction between human p53

and the human ortholog of mouse double minute 2 (MDM2) using the known

peptidomimetic inhibitor nutlin-3 [43]. In the ReLo assay, we used only the N-terminal

domains of p53 and MDM2, which were sufficient to mediate the p53-MDM2 interaction

[44]. Indeed, although an interaction between p53 1-50 and MDM2 1-118 was detected

in the DMSO control experiment, this interaction was not observed when the cells were

incubated with 5 µM nutlin-3 (**Fig. 2f**).

Together, these data suggest that the ReLo assay can be used to study

interactions involving non-*Drosophila* proteins and to screen drugs based on their

ability to either enable or inhibit a specific PPI.

ReLo reveals direct binary interactions

Thus far, we have tested known direct interactions between proteins that are not

endogenously expressed in S2R+ cells. PPIs detected with the ReLo assay may not

necessarily involve direct contact. Instead, PPIs may also result from indirect contacts

caused by incorporation of the coexpressed pair of proteins into cell-endogenous

protein complexes and subsequent indirect bridging of this pair through one or more

common interaction partners. To understand the extent to which direct or indirect

associations underlie a relocalization event observed with ReLo, we assessed

interactions between individual subunits of the CCR4-NOT complex, which is

endogenous to S2R+ cells. The CCR4-NOT complex is an essential eukaryotic

deadenylase comprising six subunits that form the core with an architecture that is

well-characterized at the molecular and structural levels [45,46]. In this complex, NOT1,

which has a size of 281 kDa in *Drosophila,* acts as the scaffolding subunit for the

assembly of all other subunits. Specifically, the CAF1-CCR4 heterodimer and CAF40

bind to the central region of NOT1, and the NOT2-NOT3 subcomplex associates with

the C-terminal region of NOT1 (**Fig. 3a**). Using ReLo, we performed a systematic

pairwise screen in which each subunit was tested against all other subunits of the

Drosophila CCR4-NOT core complex (**Fig. 3b and Supplementary Fig. 4a)**. In S2R+

cells, NOT1 and NOT3 localized exclusively to the cytoplasm, whereas NOT2, CAF1,

CAF40 and CCR4 localized to both the cyto- and the nucleoplasm (**Supplementary**

Fig. 4b). Remarkably, we observed interactions only between proteins that had

previously been shown to exhibit direct associations but not between indirectly linked

combinations: NOT1 bound specifically to CAF1, CAF40, NOT2, and NOT3, while

CAF1 bound to CCR4, and NOT2 bound to NOT3 (**Fig. 3b, c, Supplementary Fig.**

4a, b). We also tested the interactions to CAF1 using the OST4 membrane anchor and

observed results similar to those obtained using the PH anchor (**Supplementary Fig.**

4c). Together, these data suggest that the incorporation of two tagged CCR4-NOT

complex subunits into one endogenous complex is insignificant, and thus direct but not

indirect interactions between the two proteins tested are observed with the ReLo

assay.

Although the subunits of the endogenous CCR4-NOT complex show moderate

to high expression levels in S2R+ cells [47] (**Supplementary Fig. 5a)**, we aimed to

challenge the ReLo assay by testing PPI between subunits of a cytoplasmic complex

that is even more abundant in S2R+ cells. We chose the seven-subunit Arp2/3

complex, which is known to initiate actin polymerization in eukaryotes [48], and of which

several subunits are among the most highly expressed cytoskeletal genes [47,49]

(**Supplementary Fig. 5a)**. In S2R+ cells, the Arp2/3 subunits Arp3, Arpc1, and Arpc4

are expressed about two- to four-fold higher as compared to NOT3, which is the most

abundant subunit of the CCR4-NOT complex (**Supplementary Fig. 5a)**. Compared to

the CCR4-NOT complex, binary PPIs within the Arp2/3 complex are less well

characterized, and we concluded which subunit is directly or indirectly bound to

another subunit based on the crystal structure of the bovine Arp2/3 complex [50]

(**Supplementary Fig. 5b, c**). The crystal structure points to six direct pairwise

contacts. Using ReLo assays, we detected two PPIs, namely the Arpc2-Arpc4 and the

Arpc4-Arpc5 interaction (**Supplementary Fig. 5c, d**). The other four contacts, which

we did not observe with ReLo, were also not detected in an earlier pairwise study of

the human Arp2/3 complex using Y2H assays [51]. Importantly, we did not detect any

false-positive interaction using the ReLo assay. These data suggest that even if a

complex is highly abundant in a cell, the detection of a PPI between two protein

candidates in the ReLo assays is direct rather than indirectly mediated by incorporation

into the same endogenous complex. Nevertheless, we cannot exclude that

occasionally, indirect interactions are detected through a naturally expressed bridging

protein.

Finally, we compared the ReLo assay to the split-ubiquitin yeast two-hybrid

assay [27,52], which we have previously used successfully to map the interaction between

Short Oskar and Vasa [30]. When we assessed the pairwise interactions between the

six CCR4-NOT complex core subunits using Y2H, we observed mostly false negative

and false positive results; the only conclusive interaction that we detected was between

NOT2 and NOT3 (**Supplementary Fig. 6**). This observation is in stark contrast to our

data obtained using ReLo assays. As both Y2H and ReLo worked well to map the

Oskar-Vasa interaction, together the data suggest that the applicability of Y2H for

testing interactions is case/project dependent. ReLo could be expected to be more

reliable than Y2H for testing large proteins from higher eukaryotes, although a larger

scale study would be required to fully test this.

Topological description of multisubunit complexes

As has been described for S2 cells [53], we usually observed a very high cotransfection

efficiency of the S2R+ cells when performing ReLo experiments. Therefore, we tested

whether the bridging of CCR4-NOT subunits that do not directly interact is observable

when one or two common binding partners are added to the mixture. Specifically, S2R+

cells were cotransfected with three or four plasmids: one plasmid expressing the PH-

mCherry fusion construct, one plasmid expressing the mEGFP fusion construct, and

one or two plasmids expressing nonfluorescent bridging factors (**Fig. 3d**). Indeed,

NOT3 failed to interact with CAF1 or CAF40 in the presence of the control plasmid but

interacted when NOT1 was provided in addition (**Fig. 3e, f**). We also tested the bridging

of the NOT3-CCR4 interaction, which requires not only NOT1 but also the CAF1

subunit. The NOT3-CCR4 interaction was not observed if NOT1 alone was

coexpressed but was observed when both NOT1 and CAF1 were coexpressed (**Fig.**

3g). Similar results were obtained from the testing of indirect interactions with CAF40

(**Supplementary Fig. 4d**). Together, these data show that bridging experiments with

ReLo can be used to reconstitute the binding topology of multisubunit complexes such

as the CCR4-NOT complex.

Assessment of PPIs between the CCR4-NOT complex and mRNA repressor

proteins

The CCR4-NOT complex can be recruited specifically to mRNAs through adapter

RNA-binding proteins, which leads to accelerated deadenylation and eventually

degradation and/or translational repression of the targeted mRNA. We then assessed

PPIs between some of these repressor proteins and the six core subunits of the CCR4-

NOT complex using ReLo (**Fig. 4a)**. Co-IP experiments combined with structural

analysis previously revealed the specific subunit(s) of the CCR4-NOT complex through

which adapter proteins recruit the complex to a specific mRNA. For example,

Drosophila Bag-of-marbles (Bam) has been shown to bind specifically to CAF40 [54].

We confirmed this finding with ReLo: Bam bound to CAF40 but not to any other CCR4-

NOT complex subunit (**Fig. 4b and Supplementary Fig. 7a**). A point mutation in Bam

(M24E; Bam MUT), which is known to interfere with CAF40 binding [54], also abolished

CAF40 binding in the ReLo assay (**Fig. 4b**). Likewise, a mutation at the CAF40 surface

(V186E; CAF40 MUT), which was shown to prevent Bam binding [54], abolished the

interaction with Bam in the ReLo assay (**Fig. 4b**).[55]

Drosophila Roquin has been demonstrated to bind to *Drosophila* CAF40. It also

binds to the so-called CAF40 module composed of CAF40 and the CAF40-binding

region of NOT1 and to the NOT1/2/3 module composed of the C-terminal domains of

the NOT1, NOT2, and NOT3 subunits of human CCR4-NOT complex [55]. Of the

subunits tested in the ReLo assay, we detected clear binding of Roquin to *Drosophila*

CAF40, NOT1, and NOT3 but not to NOT2 (**Fig. 4c and Supplementary Fig. 7b**),

essentially confirming the previous data. In addition, we detected a previously unknown

interaction between Roquin and CCR4 (**Fig. 4c**).

Drosophila Nanos was shown to bind to the NOT1/2/3 module of the human

CCR4-NOT complex in GST pull-down experiments, but does not bind to any of these

domains individually [56]. With ReLo, we detected the interaction of Nanos with

Drosophila NOT1 but not with NOT2 or NOT3, which suggests that the NOT1

interaction might be predominant (**Fig. 4d and Supplementary Fig. 7c**). In addition,

we detected Nanos binding to the CAF40 subunit (**Fig. 4d**), an interaction that has not

previously been reported.

Encouraged by these mostly validating data, we tested the interaction of the

CCR4-NOT complex with less well characterized potential adaptor proteins.

Drosophila meiosis regulator and mRNA stability factor 1 (MARF1) is an oocyte-

specific protein that recruits the CCR4-NOT complex to target mRNAs and thereby

controls meiosis [57]. *Drosophila* Smaug recruits the CCR4-NOT complex to *nanos* and

other mRNAs to regulate posterior patterning and nuclear divisions of the early embryo

[58–62]. Whether MARF1 or Smaug directly associate with the CCR4-NOT complex was

unknown. Using ReLo, we found that Smaug interacted with the NOT3 subunit [63], and

that MARF1 bound to the NOT1 subunit (**Fig. 4e and Supplementary Fig. 7d**), which

suggests that Smaug and MARF1 are indeed direct recruiters of the CCR4-NOT

complex.

Drosophila Cup and its orthologous protein 4E-T are translation repressor

proteins that act through their interactions with the eukaryotic translation initiation

factor 4E (eIF4E) and the DEAD-box RNA helicase Me31B/DDX6 [64–68]. Both the

human 4E-T - Me31B and the *Drosophila* Cup-eIF4E complex are structurally

characterized [69,70] and we detected both interactions using ReLo assays (**Fig. 4f, left**

panels). As demonstrated through co-IP experiments, Cup also interacts with the

CAF1, CCR4, NOT1, NOT2, and NOT3 subunits of the CCR4-NOT complex [71]. We

did not detect Cup binding to any subunit of the CCR4-NOT core complex using ReLo

assays (**Supplementary Fig. 7e**), which suggests that if Cup directly recruits the

CCR4-NOT complex, then it probably involves weak and multivalent interactions. Cup

also reportedly binds the repressors Nanos and Bruno [65,72,73], and with the ReLo assay

we confirmed both interactions (**Fig. 4f, right panels**). Consistent with a recent report

[74], we also observed that Bruno binds directly to Nanos (**Fig. 4g**). We then asked if

Bruno and Nanos bind to Cup as a complex. However, when coexpressed with Cup,

Bruno and Nanos did not interact (**Fig. 4g**), which suggests that Bruno and Nanos

compete for binding to the same region of Cup. Bruno has thus far not been shown to

establish direct contact with the CCR4-NOT complex, and we did not detect Bruno

binding to any of the core CCR4-NOT complex subunits (**Supplementary Fig. 7f**).

In summary, the assessment of PPIs using ReLo involving a wide selection of

repressor proteins, confirmed many previously described interactions and revealed

new interactions. Thus, ReLo is a powerful method for identifying novel PPIs in a

candidate approach.

Combining ReLo assays with structural prediction

Lastly, we aimed to obtain some molecular insight into one of the newly identified

interactions. We tested the structural prediction of the Nanos-CAF40 complex using

AlphaFold2-multimer-v3 and obtained a high-confidence model as indicated by the

predicted aligned error (PAE) plot (**Fig. 4h, Supplementary Fig. 8a**) [75–77]. The

predicted structure is composed of full-length CAF40 and a short α-helix spanning

residues 140 to 160 of Nanos, which is bound to the same site of CAF40 that was

shown to accommodate the Bam or Roquin α-helical peptide [54,55] (**Supplementary**

Fig. 8b). The CAF40 point mutation that prevented Bam binding (**Fig. 4b**), also

abolished the interaction with Nanos (**Fig. 4i**). Likewise, deletion of the CAF40-binding

α-helix from Nanos (Nanos Δ140-160) prevented the interaction with CAF40 (**Fig. 4i**).

These data are consistent with the predicted structural model of the Nanos-CAF40

complex. Taken together, our exemplary analysis suggests that the combination of

ReLo assays with structural modeling can lead to fast molecular insight into protein

complexes.

DISCUSSION

The detection and characterization of PPIs are crucial for uncovering regulatory

mechanisms that underlie cellular processes. Here, we described ReLo, a rapid

procedure for identifying and investigating pairwise as well as multisubunit PPIs. We

provided strong evidence showing that pairwise PPIs identified with ReLo are based

on direct contacts. Thus, PPI partners newly identified using ReLo are highly promising

candidates for use in subsequent studies using *in vitro* assays and experimental

structural biology methods. Alternatively, PPI mapping data obtained from ReLo

experiments may be used to guide subsequent modeling experiments using

AlphaFold2-Multimer to obtain structural information on complexes [75,76]. A thus

predicted protein–protein interface may then rapidly be validated through mutational

analysis using ReLo, and the mutations that specifically interfere with a PPI may finally

be tested *in vivo* without the need for purifying a protein or experimentally determining

a structure of a protein complex.

We refrained from quantifying the relocalization events observed with ReLo

through image processing tools to assess or compare PPIs. Although only qualitative,

the results obtained were typically clear. For example, in control experiments, we never

observed relocalization, and in experiments in which proteins interacted, most or all

cells showed relocalization (**Supplementary Figs. 2 and 3**). Furthermore, the

implementation of a relocalization score as a quantified readout of an interaction may

have easily led to the false assumption that the score represents a measure of binding

affinity between proteins. Such false assumptions are also an issue with Y2H dot

assays, which appear to but do not provide quantitative data. The degree of

relocalization identified with the ReLo assay or yeast cell growth in a Y2H assay not

only reflects the binding affinity of the protein complex but also relies on the protein

expression levels, protein stability, subcellular localization, and other factors. With this

thought in mind, we prefer to consider ReLo a qualitative PPI method, the strength of

which lies in its speed and versatility.

ReLo is easy to perform in every laboratory equipped with cell culture

technology and with access to a confocal fluorescence microscope. In contrast to

standard Y2H or PCA assays, the expression levels of the proteins tested in a ReLo

assay are directly monitored during microscopy, which facilitates data interpretation.

The *Drosophila* S2R+ cell line used for our ReLo tests requires very simple handling.

Similar to S2 cells, S2R+ cells grow at ambient temperature, do not need an incubator

with CO_2, and can be passaged without the need for coated dishes nor the scraping or

trypsinization of cells [53]. Using S2R+ cells, we successfully investigated not only

Drosophila PPIs but also human PPIs. However, when a cell line derived from an

alternative organism is needed for a ReLo assay, only the PH or OST4 membrane

anchors need to be inserted into expression vectors compatible with the desired cell

line.

With ReLo assays, we investigated PPIs involving NOT1 and Tudor, two large

proteins comprising 281 kDa (2505 aa) and 285 kDa (2515 aa), respectively. Thus, the

detection of PPIs using the ReLo assay appears to be successful regardless of the

length of the protein of interest, which, in our experience, is a big advantage over Y2H

methods. Similar to any other cell-based PPI assay, attention is needed when working

with toxic proteins. To reduce toxicity, we recommend testing the splitting of the toxic

protein into its domains or, when possible, testing protein variants with mutated active

sites.

PPIs are highly relevant as putative therapeutic targets for the development of

new treatments [78,79]. In ReLo, complex formation is reversible, and we demonstrated

that ReLo is a tool for testing the effect of small molecules on PPIs. Due to its simple

setup, ReLo might be easily adjusted to a high-throughput approach using automated

imaging, which allows for large drug screening experiments. In a setup in which a

single specific PPI is subjected to a drug screening experiment, it might be

advantageous to express the two protein partners from one plasmid to ensure

equimolar protein expression within a cell, which may facilitate data interpretation.

Taken together, our data show that investigations with ReLo are rapid, simple,

and reliable. We recommend using ReLo as an initial tool to screen and characterize

PPIs, particularly in cases in which Y2H or more complicated approaches fail.

Subsequently, ReLo can be complemented with biochemical, structural, or genetic

approaches to further characterize or ultimately validate the biological relevance of a

given PPI.

ACKNOWLEDGMENTS

We thank Elmar Wahle, Julien Bethune, Bernd Bukau, and Ivana Vonkova for DNA

constructs and Aurelio Telemann for the S2R+ cell line. We are grateful to Christian

Bleischwitz, Eva Boberlin, Jana Kubíková, Katharina Müller, Rebecca Reinig, Gabrielė

Ubartaitė, and Xiaohan Zhao for technical assistance provided with some experiments.

We thank the Nikon Imaging Center at the University of Heidelberg for access to

microscopes. We gratefully acknowledge the data storage service SDS@hd supported

by the Ministry of Science, Research and the Arts Baden-Württemberg (MWK) and the

German Research Foundation (DFG) through the grants INST 35/1314-1 FUGG and

INST 35/1503-1 FUGG. We thank Peter Becker, Hüseyin Besir, Julien Béthune, and

Doris Höglinger for their insightful comments on the manuscript. This work was funded

by the Emmy-Noether Program of the German Research Foundation (DFG; JE-827/1-

1).

METHODS

Plasmid backbone construction

pAc5.1-EGFP and pAc5.1-mCherry were described previously [28]. pAc5.1-mEGFP

encodes the monomeric A206K EGFP variant and was generated by site-directed

mutagenesis of pAc5.1-EGFP. The pAc5.1-λN-HA vector [80] was used to create

nonfluorescent constructs. The *Saccharomyces cerevisiae* OST4 sequence was

amplified from a pDHB1 vector [27] and inserted into the KpnI site of pAc5.1-mCherry to

yield pAc5.1-OST4-mCherry. For cloning into the pAc5.1-EGFP, pAc5.1-mEGFP,

pAc5.1-mCherry, pAc5.1-λN-HA, and pAc5.1-OST4-mCherry vectors, sequences of

interest were inserted into the blunt-end EcoRV site. The rat PLCδ_1-PH sequence was

amplified from the pETM11-His6-PH-Sumo3-sfGFP vector [81] and inserted into the KpnI

site of pAc5.1-mCherry and pAc5.1-mEGFP to obtain pAc5.1-PH-mCherry and

pAc5.1-PH-mEGFP, respectively. Alternatively, the PH sequence was inserted into the

EcoRV site of pAc5.1-mCherry to obtain the pAc5.1-mCherry-PH vector. For all PH-

containing vectors, a new unique in-frame FspAI blunt end site was introduced 3' or 5'

with respect to the fluorescent protein sequences and was used to insert the sequence

of interest.

Cloning

Ligation reactions were assembled in a 10 µl reaction containing T4 DNA ligase

(Thermo Scientific), 50 ng of vector DNA, and a 5- to 20-fold molar excess of DNA

inserts and were incubated for 1 to 2 h at room temperature. DNA inserts were

generated by PCR amplification. To prevent re-ligation, the reaction was supplemented

with 0.25 µl of the blunt end restriction enzyme that was also used for linearization of

the respective vector, except when this site was present in the insert sequence.

Positive clones were screened by colony PCR using one primer that binds to the vector

and a second that binds to the insert. All constructs were verified by sequencing.

Detailed information on all DNA constructs used in this study is listed in

Supplementary Table 1.

Cell culture

S2R+ cells were cultured at 25°C in Schneider's *Drosophila* medium + (L)-glutamine

(Thermo Scientific) supplemented with 10% fetal bovine serum (Sigma) and 1 x

Gibco™ Antibiotic-Antimycotic (Thermo Scientific). Cells were seeded in six-well

glass-bottom plates (Cellvis; earlier experiments) or four-well polymer μ-slides (Ibidi;

later experiments) and cotransfected using jetOPTIMUS (Polyplus Transfection)

according to the instruction manual. Specifically, 600 μl of cells with a density of

1×10^6 cells/ml were seeded per well in a four-well slide and 61 μl of transfection

reagent mixture was added. This mixture contained 1 μl of transfection reagent and

600 ng of DNA in total for cotransfecting two (300 ng DNA each), three (200 ng each),

or four (150 ng each) plasmids diluted in jetOPTIMUS buffer. After 48 h of incubation

at 25°C, images of live cells were taken with a 60x or 100x oil objective and a Nikon

TE2000 laser scanning, a Nikon Ti-E spinning disc confocal, or a Nikon Eclipse Ti2

laser scanning fluorescence microscope and processed with Fiji software [82]. Changing

the amount of DNA or transfection reagent during the transfection did not change the

protein expression levels in the cells. However, reducing the total DNA amount to

150 ng and/or the transfection reagent to 0.5 μl significantly reduced the transfection

efficiency.

Drug treatment
484
Rapamycin and nutlin-3 (both from MedChemExpress) were dissolved in DMSO to

obtain stock concentrations of 10.9 mM and 10 mM, respectively, and they were diluted

with SF-4 Baculo express medium (BioConcept) to generate working concentrations

of 100 nM and 5 μM, respectively. S2R+ cells were grown in SF-4 medium and

cotransfected with desired plasmids using FuGENE® HD transfection reagent

(Promega). Specifically, cells were seeded as described above and 26 μl of

transfection reagent mixture was added. This mixture contained 1 μl of transfection

reagent and 400 ng of DNA (i.e., 200 ng DNA each for cotransfecting two plasmids)

diluted in SF-4 Baculo express medium. Twenty-four hours after transfection, the cells

were treated with the drug by replacing the medium with drug-containing medium. Cells

were imaged after 24 h of incubation with the drug or DMSO control medium at 25°C.

AUTHOR CONTRIBUTIONS

H.K.S. and J.M. acquired, analyzed, and interpreted the data. M.J. conceived and

supervised the project, acquired, analyzed, and interpreted the data, and wrote the

manuscript with input from all authors.

COMPETING INTEREST

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

The authors declare that all relevant data supporting the findings of this study are

available within the article and its supplementary information files.

MAIN FIGURES

Fig. 1. ReLo assay and its use in PPI mapping and mutational analysis.
a Plasmids encoding fluorescently tagged proteins 1 and 2 are cotransfected into S2R+ cells. Protein 1 carries an additional fusion to a membrane anchoring (MA) domain, i.e., PH or OST4. After 48 h, protein localization is analyzed by confocal fluorescence microscopy. If protein 2 interacts with protein 1, protein 2 is relocalized to the plasma membrane (PH domain) or ER (OST4). **b** Domain organization of *Drosophila* Oskar and Vasa proteins. Oskar is expressed in two isoforms, and Short Oskar lacks amino acids 1-138. The eLOTUS domain of Oskar was previously shown to interact with the C-terminal domain (CTD) of Vasa (yellow stripe), and the crystal structure of the complex has been resolved [28,30]. **c** Vasa 463-661, but not Vasa 1-200 or 200-463, interacted with the Oskar eLOTUS domain. Vasa 463-661 did not interact with Oskar 241-396 or Oskar 398-606. **d** Oskar A162E/L228E and Vasa F504E point mutations (MUT) interfered with the Oskar-Vasa interaction. **e** OST4-mCherry Oskar localized to the ER (top panel), and Vasa fully relocalized with Oskar to the ER (bottom panel), while Vasa MUT did not. In the "Control" experiments the respective construct was coexpressed with the vector indicated at the top of the panel lacking an insert. The scale bar is 10 µm.

529
530
Fig. 2. PPI studies using ReLo with respect to conformation, posttranslational modification and drug sensitivity.
a Pictorial representation of Vasa N- and C-terminal RecA-like domains (NTD and CTD, respectively) in the open and closed conformations. Vasa-open and Vasa-closed carry the K295N or E400Q mutations, respectively. **b** EGFP-Vasa-open, but not EGFP-Vasa-closed, interacted with OST4-mCherry-Oskar. The white dotted lines indicate cell boundaries. **c** Symmetric dimethyl arginine (sDMA) modifications can be catalyzed by PRMT5 and recognized by eTud domains. **d** Wild-type Aub interacted with Tudor, whereas Aub carrying nonmethylatable 4R->K point mutations (R11K/R13K/R15K/R17K) did not. The profiles indicate the measured fluorescence distribution across the cell along a line indicated by the white arrows (shown for the green channel only). Red and green fluorescence signals are independent and displayed relatively. **e** Rapamycin induced an interaction between human FKBP12 and FRB. The control contained 0.0009% dimethyl sulfoxide (DMSO). **f** The interaction between p53 1-50 and MDM2 1-118 was inhibited by nutlin-3 treatment. The control contained 0.05% DMSO. In the "Control" experiments the respective construct was coexpressed with the vector indicated at the top of the panel lacking an insert. The scale bar is 10 µm.

Fig. 3. ReLo identifies direct PPIs.
a Subunit organization of the *Drosophila* CCR4-NOT core complex. **b** NOT1-mCherry-
PH recruited CAF1, CAF40, NOT2, and NOT3 but not CCR4 to the plasma membrane.
See **Supplementary Fig. 3a** for the PPI analysis of the other CCR4-NOT complex
subunits. **c** Summary of the results from a pairwise screen of CCR4-NOT complex core
subunits using ReLo. The blue color indicates an interaction, and the gray color
indicates no interaction. **d** Schematic showing the ReLo assay involving coexpression
of four different protein constructs. **e** NOT3 interacted with CAF1 in the presence of
NOT1. **f** NOT3 interacted with CAF40 in the presence of NOT1. **g** NOT3 recruited
CCR4 to the plasma membrane only in the presence of both NOT1 and CAF1. In the
"Control" experiments the respective construct was coexpressed with the vector
indicated at the top of the panel lacking an insert. The scale bar is 10 μm.
563

Fig. 4. PPIs between the CCR4-NOT complex and repressor proteins.
Summary of the results (**a**) from a pairwise screen of CCR4-NOT complex core subunits with Bam (**b**), Nanos (**c**), Roquin (**d**), MARF1 (**e**), Bruno and Cup (**Supplementary Fig. 7**), and Smaug [63]. The blue color indicates an interaction, and the gray color indicates no interaction. Bam MUT carries the M24E and CAF40 MUT the V186E point mutation. **f** Cup interacted with eIF4E, Me31B, Bruno, and Nanos. **g** Bruno interacted with Nanos in the absence but not in the presence of Cup. In the "Control" experiments the respective construct was coexpressed with the vector indicated at the top of the panel lacking an insert. The scale bar is 10 µm.

REFERENCES

1. Masters, S. C. Co-immunoprecipitation from transfected cells. *Methods Mol Biol* **261**, 337–350 (2004).
2. Ransone, L. J. Detection of protein-protein interactions by coimmunoprecipitation and dimerization. *Methods Enzymol* **254**, 491–497 (1995).
3. Rigaut, G. *et al.* A generic protein purification method for protein complex characterization and proteome exploration. *Nat Biotechnol* **17**, 1030–1032 (1999).
4. Gingras, A. C., Abe, K. T. & Raught, B. Getting to know the neighborhood: using proximity-dependent biotinylation to characterize protein complexes and map organelles. *Curr Opin Chem Biol* **48**, 44–54 (2019).
5. Bosch, J. A., Chen, C. L. & Perrimon, N. Proximity-dependent labeling methods for proteomic profiling in living cells: An update. *Wiley Interdiscip Rev Dev Biol* **10**, 1–17 (2021).
6. Roux, K. J. Marked by association: techniques for proximity-dependent labeling of proteins in eukaryotic cells. *Cell Mol Life Sci* **70**, 3657–64 (2013).
7. Lalonde, S. *et al.* Molecular and cellular approaches for the detection of protein-protein interactions: Latest techniques and current limitations. *Plant Journal* **53**, 610–635 (2008).
8. Walport, L. J., Low, J. K. K., Matthews, J. M. & Mackay, J. P. The characterization of protein interactions – what, how and how much? *Chem Soc Rev* **50**, 12292–12307 (2021).
9. Remy, I. & Michnick, S. W. Application of protein-fragment complementation assays in cell biology. *Biotechniques* **42**, 137–145 (2007).
10. Cui, Y. *et al.* Techniques for detecting protein-protein interactions in living cells: principles, limitations, and recent progress. *Sci China Life Sci* **62**, 619–632 (2019).
11. Blaszczak, E., Lazarewicz, N., Sudevan, A., Wysocki, R. & Rabut, G. Protein-fragment complementation assays for large-scale analysis of protein-protein interactions. *Biochem Soc Trans* **49**, 1337–1348 (2021).
12. Wang, T. *et al.* Detecting Protein-Protein Interaction Based on Protein Fragment Complementation Assay. *Curr Protein Pept Sci* **21**, 598–610 (2020).
13. Lv, S. *et al.* CAPPI: A Cytoskeleton-Based Localization Assay Reports Protein-Protein Interaction in Living Cells by Fluorescence Microscopy. *Mol Plant* **10**, 1473–1476 (2017).
14. Grefen, C. *et al.* Subcellular localization and in vivo interactions of the Arabidopsis thaliana ethylene receptor family members. *Mol Plant* **1**, 308–320 (2008).
15. Winkler, J. *et al.* Visualizing protein–protein interactions in plants by rapamycin-dependent delocalization. *Plant Cell* **33**, 1101–1117 (2021).
16. Dixon, A. S. & Lim, C. S. The nuclear translocation assay for intracellular protein-protein interactions and its application to the Bcr coiled-coil domain. *Biotechniques* **49**, 519–524 (2010).
17. Lee, K. H., Lee, S., Lee, W. Y., Yang, H. W. & Heo, W. Do. Visualizing dynamic interaction between calmodulin and calmodulin-related kinases via a monitoring method in live mammalian cells. *Proc Natl Acad Sci U S A* **107**, 3412–3417 (2010).
18. Gallego, O. *et al.* Detection and Characterization of Protein Interactions In Vivo by a Simple Live-Cell Imaging Method. *PLoS One* **8**, 1–6 (2013).

627   19.   Taslimi, A. *et al.* An optimized optogenetic clustering tool for probing protein
628   interaction and function. *Nat Commun* **5**, 4925 (2014).
629   20.   Martin, B. R., Deerinck, T. J., Ellisman, M. H., Taylor, S. S. & Tsien, R. Y.
630   Isoform-Specific PKA Dynamics Revealed by Dye-Triggered Aggregation and
631   DAKAP1α-Mediated Localization in Living Cells. *Chem Biol* **14**, 1031–1042
632   (2007).
633   21.   Watanabe, T. *et al.* Genetic visualization of protein interactions harnessing
634   liquid phase transitions. *Sci Rep* **7**, 1–13 (2017).
635   22.   Schneider, I. Cell lines derived from late embryonic stages of Drosophila
636   melanogaster. *J Embryol Exp Morphol* **27**, 353–365 (1972).
637   23.   Yanagawa, S. I., Lee, J. S. & Ishimoto, A. Identification and characterization of
638   a novel line of Drosophila Schneider s2 cells that respond to wingless
639   signaling. *Journal of Biological Chemistry* **273**, 32353–32359 (1999).
640   24.   Garcia, P. *et al.* The Pleckstrin Homology Domain of Phospholipase C-δ1
641   Binds with High Affinity to Phosphatidylinositol 4,5-Bisphosphate in Bilayer
642   Membranes. *Biochemistry* **34**, 16228–16234 (1995).
643   25.   Lemmon, M. A., Ferguson, K. M., O'Brien, R., Sigler, P. B. & Schlessinger, J.
644   Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin
645   homology domain. *Proc Natl Acad Sci U S A* **92**, 10472–10476 (1995).
646   26.   Kim, H., Yan, Q., Von Heijne, G., Caputo, G. A. & Lennarz, W. J. Determination
647   of the membrane topology of Ost4p and its subunit interactions in the
648   oligosaccharyltransferase complex in Saccharomyces cerevisiae. *Proc Natl*
649   *Acad Sci U S A* **100**, 7460–7464 (2003).
650   27.   Möckli, N. *et al.* Yeast split-ubiquitin-based cytosolic screening system to
651   detect interactions between transcriptionally active proteins. *Biotechniques* **42**,
652   725–730 (2007).
653   28.   Jeske, M., Müller, C. W. & Ephrussi, A. The LOTUS domain is a conserved
654   DEAD-box RNA helicase regulator essential for the recruitment of Vasa to the
655   germ plasm and nuage. *Genes Dev* **31**, 939–952 (2017).
656   29.   Breitwieser, W., Markussen, F. H., Horstmann, H. & Ephrussi, A. Oskar protein
657   interaction with vasa represents an essential step in polar granule assembly.
658   *Genes Dev* **10**, 2179–2188 (1996).
659   30.   Jeske, M. *et al.* The crystal structure of the Drosophila germline inducer Oskar
660   identifies two domains with distinct Vasa Helicase- and RNA-binding activities.
661   *Cell Rep* **12**, 587–598 (2015).
662   31.   Yang, N. *et al.* Structure of Drosophila Oskar reveals a novel RNA binding
663   protein. *Proc Natl Acad Sci U S A* **112**, 11541–11546 (2015).
664   32.   Linder, P. & Jankowsky, E. From unwinding to clamping - the DEAD box RNA
665   helicase family. *Nat Rev Mol Cell Biol* **12**, 505–16 (2011).
666   33.   Walker, J. E., Saraste, M., Runswick, M. J. & Gay, N. J. Distantly related
667   sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases
668   and other ATP-requiring enzymes and a common nucleotide binding fold.
669   *EMBO J* **1**, 945–951 (1982).
670   34.   Gorbalenya, A. E., Koonin, E. V., Donchenko, A. P. & Blinov, V. M. A
671   conserved NTP-motif in putative helicases. *Nature* **333**, 22 (1988).
672   35.   Xiol, J. *et al.* RNA clamping by Vasa assembles a piRNA amplifier complex on
673   transposon transcripts. *Cell* **157**, 1698–1711 (2014).
674   36.   Stopa, N., Krebs, J. E. & Shechter, D. The PRMT5 arginine methyltransferase:
675   Many roles in development, cancer and beyond. *Cellular and Molecular Life*
676   *Sciences* **72**, 2041–2059 (2015).

677   37.   Gonsalvez, G. B., Rajendra, T. K., Tian, L. & Matera, A. G. The Sm-Protein
678   Methyltransferase, Dart5, Is Essential for Germ-Cell Specification and
679   Maintenance. *Current Biology* **16**, 1077–1089 (2006).

680   38.   Liu, H. *et al.* Structural basis for methylarginine-dependent recognition of
681   Aubergine by Tudor. *Genes Dev* **24**, 1876–1881 (2010).

682   39.   Kirino, Y. *et al.* Arginine methylation of Aubergine mediates Tudor binding and
683   germ plasm localization. *RNA* **16**, 70–8 (2010).

684   40.   Nishida, K. M. *et al.* Functional involvement of Tudor and dPRMT5 in the
685   piRNA processing pathway in Drosophila germlines. *EMBO Journal* **28**, 3820–
686   3831 (2009).

687   41.   Kirino, Y. *et al.* Arginine methylation of Piwi proteins catalysed by dPRMT5 is
688   required for Ago3 and Aub stability. *Nat Cell Biol* **11**, 652–658 (2009).

689   42.   Choi, J., Chen, J., Schreiber, S. L. & Clardy, J. Structure of the FKBP12-
690   rapamycin complex interacting with the binding domain of human FRAP.
691   *Science* **273**, 239–42 (1996).

692   43.   Vassilev, L. T. *et al.* In Vivo Activation of the p53 Pathway by Small-Molecule
693   Antagonists of MDM2. *Science (1979)* **303**, 844–848 (2004).

694   44.   Kussie, P. H. *et al.* Structure of the MDM2 oncoprotein bound to the p53 tumor
695   suppressor transactivation domain. *Science (1979)* **274**, 948–953 (1996).

696   45.   Collart, M. A. & Panasenko, O. O. The Ccr4-Not Complex: Architecture and
697   Structural Insights. *Subcell Biochem* **83**, 349–379 (2017).

698   46.   Temme, C., Simonelig, M. & Wahle, E. Deadenylation of mRNA by the CCR4-
699   NOT complex in Drosophila: molecular and developmental aspects. *Front
700   Genet* **5**, 143 (2014).

701   47.   Brown, J. B. *et al.* Diversity and dynamics of the Drosophila transcriptome.
702   *Nature 2014 512:7515* **512**, 393–399 (2014).

703   48.   Pizarro-Cerdá, J., Chorev, D. S., Geiger, B. & Cossart, P. The Diverse Family
704   of Arp2/3 Complexes. *Trends Cell Biol* **27**, 93–100 (2017).

705   49.   Hu, Y., Comjean, A., Perrimon, N. & Mohr, S. E. The Drosophila Gene
706   Expression Tool (DGET) for expression analyses. *BMC Bioinformatics* **18**, 98
707   (2017).

708   50.   Robinson, R. C. *et al.* Crystal Structure of Arp2/3 Complex. *Science (1979)*
709   **294**, 1679–1684 (2001).

710   51.   Zhao, X., Yang, Z., Qian, M. & Zhu, X. Interactions among subunits of human
711   Arp2/3 complex: p20-Arc as the hub. *Biochem Biophys Res Commun* **280**,
712   513–517 (2001).

713   52.   Stagljar, I., Korostensky, C., Johnsson, N. & Te Heesen, S. A genetic system
714   based on split-ubiquitin for the analysis of interactions between membrane
715   proteins in vivo. *Proc Natl Acad Sci U S A* **95**, 5187–5192 (1998).

716   53.   Yang, J. & Reth, M. Drosophila S2 Schneider cells: a useful tool for rebuilding
717   and redesigning approaches in synthetic biology. *Methods Mol Biol* **813**, 331–
718   41 (2012).

719   54.   Sgromo, A. *et al.* Drosophila Bag-of-marbles directly interacts with the CAF40
720   subunit of the CCR4-NOT complex to elicit repression of mRNA targets. *RNA*
721   **24**, 381–395 (2018).

722   55.   Sgromo, A. *et al.* A CAF40-binding motif facilitates recruitment of the CCR4-
723   NOT complex to mRNAs targeted by Drosophila Roquin. *Nat Commun* **8**,
724   14307 (2017).

725   56.   Raisch, T. *et al.* Distinct modes of recruitment of the CCR4-NOT complex by
726   Drosophila and vertebrate Nanos. *EMBO J* **35**, 974–90 (2016).

727   57.   Zhu, L., Kandasamy, S. K., Liao, S. E. & Fukunaga, R. LOTUS domain protein
728   MARF1 binds CCR4-NOT deadenylase complex to post-transcriptionally
729   regulate gene expression in oocytes. *Nat Commun* **9**, 4031 (2018).
730   58.   Semotok, J. L. *et al.* Smaug recruits the CCR4/POP2/NOT deadenylase
731   complex to trigger maternal transcript localization in the early Drosophila
732   embryo. *Curr Biol* **15**, 284–94 (2005).
733   59.   Zaessinger, S., Busseau, I. & Simonelig, M. Oskar allows nanos mRNA
734   translation in Drosophila embryos by preventing its deadnylation by
735   Smaug/CCR4. *Development* **133**, 4573–4583 (2006).
736   60.   Smibert, C. A., Wilson, J. E., Kerr, K. & Macdonald, P. M. smaug protein
737   represses translation of unlocalized nanos mRNA in the Drosophila embryo.
738   *Genes Dev* **10**, 2600–9 (1996).
739   61.   Dahanukar, A. & Wharton, R. P. The Nanos gradient in Drosophila embryos is
740   generated by translational regulation. *Genes Dev* **10**, 2610–2620 (1996).
741   62.   Dahanukar, A., Walker, J. A. & Wharton, R. P. Smaug, a novel RNA-binding
742   protein that operates a translational switch in Drosophila. *Mol Cell* **4**, 209–218
743   (1999).
744   63.   Pekovic, F. *et al.* RNA binding proteins Smaug and Cup induce CCR4–NOT-
745   dependent deadenylation of the *nanos* mRNA in a reconstituted system.
746   *Nucleic Acids Res* **51**, 3950–3970 (2023).
747   64.   Nelson, M. R., Leidal, A. M. & Smibert, C. A. Drosophila Cup is an eIF4E-
748   binding protein that functions in Smaug-mediated translational repression.
749   *EMBO J* **23**, 150–9 (2004).
750   65.   Nakamura, A., Sato, K. & Hanyu-Nakamura, K. Drosophila Cup is an eIF4E
751   binding protein that associates with Bruno and regulates oskar mRNA
752   Translation in Oogenesis. *Dev Cell* **6**, 69–78 (2004).
753   66.   Wilhelm, J. E., Hilton, M., Amos, Q. & Henzel, W. J. Cup is an eIF4E binding
754   protein required for both the translational repression of oskar and the
755   recruitment of Barentsz. *Journal of Cell Biology* **163**, 1197–1204 (2003).
756   67.   Zappavigna, V., Piccioni, F., Villaescusa, J. C. & Verrott, A. C. Cup is a
757   nucleocytoplasmic shuttling protein that interacts with the eukaryotic translation
758   initiation factor 4E to modulate Drosophila ovary development. *Proc Natl Acad
759   Sci U S A* **101**, 14800–14805 (2004).
760   68.   Kamenska, A. *et al.* The DDX6-4E-T interaction mediates translational
761   repression and P-body assembly. *Nucleic Acids Res* **44**, 6318–6334 (2016).
762   69.   Ozgur, S. *et al.* Structure of a Human 4E-T/DDX6/CNOT1 Complex Reveals
763   the Different Interplay of DDX6-Binding Proteins with the CCR4-NOT Complex.
764   *Cell Rep* **13**, 703–711 (2015).
765   70.   Kinkelin, K., Veith, K., Grünwald, M. & Bono, F. Crystal structure of a minimal
766   eIF4E-Cup complex reveals a general mechanism of eIF4E regulation in
767   translational repression. *Rna* **18**, 1624–1634 (2012).
768   71.   Igreja, C. & Izaurralde, E. CUP promotes deadenylation and inhibits decapping
769   of mRNA targets. *Genes Dev* **25**, 1955–1967 (2011).
770   72.   Verrotti, A. C. & Wharton, R. P. Nanos interacts with Cup in the female
771   germline of Drosophila. *Development* **127**, 5225–5232 (2000).
772   73.   Kim, G. *et al.* Region-Specific Activation of oskar mRNA Translation by
773   Inhibition of Bruno-Mediated Repression. *PLoS Genet* **11**, 1–22 (2015).
774   74.   Bansal, P., Madlung, J., Schaaf, K., Macek, B. & Bono, F. An Interaction
775   Network of RNA-Binding Proteins Involved in Drosophila Oogenesis. *Molecular
776   and Cellular Proteomics* **19**, 1485–1502 (2020).

777   75.   Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv*
778   2021.10.04.463034 (2021) doi:https://doi.org/10.1101/2021.10.04.463034.
779   76.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold.
780   *Nature* **596**, 583–589 (2021).
781   77.   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature*
782   *Methods 2022 19:6* **19**, 679–682 (2022).
783   78.   Ivanov, A. A., Khuri, F. R. & Fu, H. Targeting protein-protein interactions as an
784   anticancer strategy. *Trends Pharmacol Sci* **34**, 393–400 (2013).
785   79.   Buchwald, P. Small-molecule protein-protein interaction inhibitors: Therapeutic
786   potential in light of molecular size, chemical space, and ligand binding
787   efficiency considerations. *IUBMB Life* **62**, 724–731 (2010).
788   80.   Behm-Ansmant, I. *et al.* mRNA degradation by miRNAs and GW182 requires
789   both CCR4:NOT deadenylase and DCP1:DCP2 decapping complexes. *Genes*
790   *Dev* **20**, 1885–98 (2006).
791   81.   Vonkova, I. *et al.* Lipid Cooperativity as a General Membrane-Recruitment
792   Principle for PH Domains. *Cell Rep* **12**, 1519–30 (2015).
793   82.   Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis.
794   *Nat Methods* **9**, 676–682 (2012).
795