Morph-dependent nematode infection and its effect on host

movement in the land snail *Cepaea nemoralis* (Mollusca,

Gastropoda)

Maxime Dahirel[1,2*], Marine Proux[3], Claudia Gérard[3], Armelle Ansart[3]

[1]Université Côte d'Azur, INRAE, CNRS, ISA, F-06903, Sophia-Antipolis, France

[2]Department of Biology, Ghent University, B-9000, Ghent, Belgium

[3]Univ Rennes, UR1, CNRS, ECOBIO (Ecosystèmes, Biodiversité, évolution), UMR 6553, F-35000,

Rennes, France

[*]corresponding author: maxime.dahirel@yahoo.fr

Abstract

Host behaviour can be influenced by parasitic risk and infection through a variety of direct and

indirect mechanisms. We can expect individuals expressing different phenotypes to also differ in

the ways their behaviour is altered by parasites. We used the land snail *Cepaea nemoralis*, a

species with various shell colour morphs differing in behaviour and physiology, as a model to

study the link between parasite response and individual behaviour variation. We analysed

metazoan parasite abundance and its relation to behaviour (movement and food intake) in

snails from three shell morphs (from light unbanded to darker five-banded) and from two

neighbouring populations from contrasted environments. Snails were parasitized by mites,

trematodes and nematodes, from rarest to most frequent. We confirm that terrestrial

gastropods can defend against infection by trapping parasitic nematodes in their shell. We show

that nematode encapsulated in shells can uncover past infections even when a snail population

is currently nematode-free, possibly due to parasite seasonality. We present new observations

suggesting that not only nematodes but also mites might be encapsulated in shells. Infection

levels varied between morphs, with darker snails harbouring fewer nematodes. Behaviour

(movement and food intake) was dependent on nematode, but not trematode infection.

Individuals with higher nematode load ate less, irrespective of morph and population. The most-

infected morph (unbanded snails) showed reduced activity in the nematode-infected population

compared to the one that was nematode-free at sampling time. As a result, parasites may

cancel previously described behavioural differences between morphs. We discuss the possible

mechanisms explaining morph-dependent responses to parasites, and how parasite risk may be

an unseen force shaping *C. nemoralis* morph variation in space and time. We conclude on the

possible ecological consequences of the link, mediated by shell colour, between thermal and

immune responses.

Keywords: activity; *Brachylaima*; food intake; *Riccardoella*; shell colour polymorphism; snail

host-parasite interactions

Introduction

Parasites and the behaviour of their hosts are interlinked in complex ways: parasites change the environmental context in which potential hosts express their behaviour, and directly alter actual hosts' internal phenotypic state; host behaviour may favour or hinder parasite spread. Focusing on movement-related behaviours (Binning, Shaw & Roche, 2017 for a review), potential hosts may react to the perceived risk of infection by moving to avoid contact with parasites (e.g. Davies & Knowles, 2001). Costs of infection may force infected individuals to limit or delay movements (e.g. Debeffe et al., 2014; O'Dwyer, Kamiya & Poulin, 2014), possibly reshaping the distribution of infection risk in space. All these responses may be shaped by pre-existing behavioural variation, leading to feedbacks between parasite dynamics and the behaviour of uninfected and infected individuals (Ezenwa et al., 2016). To untangle these dynamics, and understand their consequences, we need to map the existing relationships between parasite infection and behavioural variation.

Land snails and slugs host a diverse array of metazoan parasites and parasitoids, including flies, mites, cestodes, trematodes and nematodes (Barker, 2004; Morley & Lewis, 2008; Gracenea & Gállego, 2017; Pieterse, Malan & Ross, 2017; Gérard et al., 2020). Depending on species, parasites use terrestrial gastropods as their only hosts, or as intermediate hosts with definitive hosts often being birds and mammals (Barker, 2004). Some of these parasites can cause diseases (e.g., strongylosis) in domesticated animals and humans (Meerburg, Singleton & Kijlstra, 2009; Butcher, 2016; Giannelli et al., 2016).

Terrestrial gastropods can defend against metazoan parasites in several ways. They may alter their behaviour to avoid parasite-infected microhabitats (Wynne, Morris & Rae, 2016). If parasites manage to enter snails, they can be neutralized through several physiological mechanisms, including cellular and non-cellular immune responses (Furuta & Yamaguchi, 2001; Coaglio et al., 2018). Snails can also trap nematodes directly in the inner layer of their shell (Rae, 2017). As in insects (González-Santoyo & Córdoba-Aguilar, 2012; San-Jose & Roulin, 2018), melanin and the phenoloxidase pathway involved in melanisation play a key role in snail defences against parasites (Coaglio et al., 2018). This association between immunity and

pigment-related pathways could be especially relevant in land snails. Indeed, many species
exhibit shell colour polymorphism (Williams, 2017), which affects behaviour, thermal responses
and vulnerability to predators (e.g. Jones, Leith & Rawlings, 1977; Rosin *et al.*, 2011; Surmacki,
Ożarowska-Nowicka & Rosin, 2013; Köhler *et al.*, 2021).

The grove snail *Cepaea nemoralis* has a long history as an iconic model in evolutionary biology
(Jones *et al.*, 1977; Ożgo, 2009). Individuals can differ in background shell color (yellow, pink or
brown) as well as in the number of dark bands (from zero to five). Decades of research have
aimed to disentangle the roles of thermal selection, predation risk and neutral processes in
driving genetic variation in morph frequencies (Cain & Sheppard, 1954; Lamotte, 1959; Jones *et*
al., 1977; Schilthuizen, 2013; Ożgo *et al.*, 2017). These color differences are associated with
behavioural and physiological differences (e.g. Kavaliers, 1992; Dahirel, Gaudu & Ansart, 2021),
leading to potentially adaptive phenotypic syndromes.

If susceptibility and response to parasite infection are shaped by host behaviour and physiology,
this means that (i) *C. nemoralis* morphs might consistently differ in their infection levels, (ii)
variation in infection between populations may alter host behaviour in a morph-dependent
way, (iii) infection and behaviour may also covary independently of morph. In this context, we
compared natural infections by metazoan parasites across two habitats differing in microclimate
conditions. We examined whether infections varied with morph and potentially morph-
dependent behaviours.

Based on the link between pigments and immune defence outlined above, we expected snails
with more dark bands hosted fewer parasites. Since darker snails are less mobile (Dahirel *et al.*,
2021), this is in line with some predictions from the pace-of-life syndrome hypothesis, namely
that fast active individuals have lower immune responses (Réale *et al.*, 2010; but see Royauté *et*
al., 2018). We also hypothesized that metazoan parasites would have notable negative effects
on snail performance (McElroy & de Buron, 2014), leading to slower behaviours and changed
food intake in more infected snails (both at the morph level and within morphs). We expected
these effects on behaviour to be stronger in morph(s) more vulnerable to parasite infection.

Methods

Snail collection and maintenance

We collected snails in early May 2019 (i.e. at most ≈ 1.5 month after hibernation) from two contrasted sites in and near the village of Arçais, France. From previous studies, we know that snail parasites are present, albeit in a different species (*Cornu aspersum*, Gérard *et al.*, 2020), and that environmental differences between the sites are linked to differences in the frequency of darker *Cepaea nemoralis* morphs (Dahirel *et al.*, 2021).

- The "open habitat" population is in a garden with little to no trees and thus exposed to the sun throughout the year (≈ 46°17′50″N, 0°41′30″ W).

- The "shaded habitat" population (≈ 46°18′01″ N, 0°42′56″ W) is in a small humid woodlot which is not directly exposed to the sun during snail activity months, and occasionally experiences partial winter flooding. The substrate is poorer in calcite than in the open habitat (Dupuis *et al.*, 1983; BRGM, 2005). Five-banded snails are more frequent there, in agreement with hypotheses on shell colour and thermal selection (Dahirel *et al.*, 2021).

We sampled 90 adult individuals per population, divided equally into unbanded, three-banded, and five-banded snails. Adults were recognizable by the presence of a reflected "lip" around the shell opening. We only sampled snails with yellow background shells, leaving aside the rarer brown and pink-shelled morphs. Sampling was random with respect to other shell features (e.g. band fusion).

We kept snails under standard conditions (20 ± 1°C, L:D 16:8) in 30 × 45 × 8 cm^3 plastic crates lined with 1-2 cm of soil kept humid and regularly renewed, with a lid made of a fine plastic mesh. Snails were grouped by morph × population of origin. Food (vegetal flour supplemented with calcium, Hélinove, Saint Paul en Pareds, France) was available *ad libitum* in Petri dishes. Four days before the experiments, we marked snails with a unique identifier on the side of their shell (using Posca pens, uni Mitsubishi Pencil, Tokyo, Japan). At the same time, we measured snail size (greater shell diameter) to the nearest 0.1 mm.

Movement behaviour and food intake

Behavioural experiments began two weeks after snail sampling. We tested each snail twice for each trait, to separate repeatable among-individual variation from within-individual variability. To ensure a manageable workload, we split the 180 snails in three groups of 60, each starting the experimental sequence on a different day, with each group containing snails from all band number × population combinations in equal proportions.

We first tested snails for movement behaviour. We stimulated snail activity by placing them in a Petri dish containing ≈ 1 mm of water until they were out of their shell. We then placed each snail individually on a clean lab bench, and used the time snails took to move more than 10 cm in a straight line from their starting point as a measure of movement (longer times implying slower individuals). Importantly, we only counted time after snails moved more than 2 cm (i.e. about one shell length) from their starting point, to ensure all observed snails were truly active. Some snails stayed inactive and the relevant observations were thus excluded (11 individuals missed one of the two tests, 6 individuals missed both). We stopped observation 20 min after placing snails on the lab bench (not after the start of activity).

Movement tests were immediately followed by food intake tests. We kept all snails individually in $10 \times 7 \times 6$ cm^3 boxes for 18 hours (including 8 hours of darkness) and provided them with 1.5 g of dry food (a non-limiting quantity, large enough to feed one snail for at least three days, based on preliminary tests). Afterwards, we recovered, dried (60°C for 60 hours) and weighed the remaining food (to the nearest 0.1 mg) to estimate food intake.

Snails were then placed back in their original box. The second test for each behaviour was done one week later.

Parasitological research

After freezing at -20°C and thawing, we dissected each snail under a binocular microscope. After removing the body from the shell, we looked for and counted metazoan parasites in various locations: space between shell and soft body, lung, heart, kidney, body cavity and digestive gland, as in Gérard et al. (2020). We found nematodes, larval trematodes and parasitic mites

(Acari). Trematodes were identified to the genus level as in Gérard et al. (2020), and mites using

Fain (2004). That level of taxonomic precision could not be reached for nematodes. However,

nematodes from different families consistently use/infect different organs and reach different

body sizes within snails (Morand, Wilson & Glen, 2004; Wilson, 2012). We assigned nematodes

to size classes (<1 mm, 1-3 mm, 3-5 mm, >5mm) using graph paper, and used nematode

location and size to infer the most likely families infecting our snails. We additionally observed

shells and shell fragments under brighter light to detect encapsulated parasites (Rae, 2017). By

simplicity, we use "live" as a shorthand to designate parasites found free within the body or

between the body and the shell during dissection, and thus presumably still living before snails

were sacrificed, by opposition with "trapped" or "encapsulated" parasites found within shells.

Statistical analyses

We analysed our data in a Bayesian framework, using R (version 4.2.0, R Core Team, 2022) and

the brms R package (Bürkner, 2017) as frontends for the Stan language (Carpenter *et al.*, 2017).

We used the tidybayes, bayesplot, patchwork packages for data preparation, model evaluation

and plotting, as well as the tidyverse suite of packages (Gabry *et al.*, 2019; Kay, 2019; Wickham

et al., 2019; Pedersen, 2020). One snail (out of 180) died during the experiments and was

excluded from statistical analyses, but was used for natural history information (see **Results**).

We ran two models, which both included at least effects of population of origin, number of shell

bands, as well as population × banding interactions as categorical explanatory variables

(hereafter "morph and population effects").

We first used a Gaussian linear model to analyse how snail shell diameter (mean centred and

scaled to unit 1 SD), varied with morph and population effects.

We then ran a multivariate generalized mixed/multilevel model to examine how behaviour and

parasite infections varied between morph and populations, and whether these traits were

correlated at the individual level. Trematodes and nematodes could not be identified to species,

so we used the total numbers of trematodes and nematodes per host snail in our analyses (as in

e.g. Debeffe *et al.*, 2014). Mite infections were too rare to be analysed statistically (**Results**).

This multivariate model included four submodels:

• live trematode abundance was modelled as a function of morph and population effects,

as well as shell size, using a Poisson model. This model included random effects of test

group and individual identity. The latter acts as an observation-level random effect and

accounts for overdispersion (Harrison, 2014); it also allows us to build an individual-level

correlation matrix connecting all four traits.

• live nematode abundance was modeled as trematode abundance, with one change. As

no live nematodes were found in the shaded population, we only fit this submodel to

data from the open population and removed the population fixed effect. Importantly,

keeping all data instead does not change our conclusions (see **Data availability**).

• food consumption (mean-centred and scaled to unit 1SD) was analysed using a Gaussian

model including the same fixed and random effects as trematode abundance. Although

food consumption data are proportions (of total food), and so are well-suited to Beta

models (Douma & Weedon, 2019), the Gaussian model behaved much better regarding

posterior predictive checks (**Data availability**). As individuals were measured twice, the

individual random effect here allows separation between within- and among individual

variation.

• movement behaviour (latency to move more than 10 cm from the starting point) was

inverse-transformed, mean-centred and scaled to unit 1SD, and then analysed using a

Gaussian model including the same fixed and random effects as above. We initially

attempted to analyse untransformed latencies using lognormal or Gamma models, but

the Gaussian model performed better regarding posterior predictive checks (**Data**

availability).

In both models we used weakly informative priors (McElreath, 2020): $\mathrm{Normal}(0,1)$ priors for

fixed effects, $\mathrm{half-Normal}(0,1)$ priors for standard deviations (random effect and residual),

and a $\mathrm{LKJ}(2)$ prior for the correlation matrix in the multivariate model. See **Supporting**

Information S1 for detailed model descriptions. We ran four chains per model for 4000

iterations each, with the first half of each chain used as warmup. We checked convergence and
mixing graphically and using the updated   statistic (Vehtari *et al.*, 2021). All parameters had
satisfactory effective sample sizes (Vehtari *et al.*, 2021). Posteriors are summarised in text and
figures as means [95% Highest Posterior Density Intervals].

We also looked at the proportion of snails with encapsulated nematodes as a function of shell
morph and population. Here we did not run statistical models, and simply present these
proportions as qualitative information about past parasite exposure in both populations. We
explicitly refrained from analysing the prevalence or abundance of encapsulated nematodes at
the individual level within populations, as it is likely linked to cumulative exposure time/age,
and snail age is not known (and cannot be approximated by e.g. size, given *Cepaea nemoralis*
exhibits determinate growth).

Results

Snail size

Figure 1. Snail shell diameter as a function of band number and population of origin. Grey dots are
observed values, displayed along the posterior distributions of the average sizes (black dots and
segments are the means and 95% CI of the posteriors).

Snails from the shaded population were on average smaller than those from the open habitat if

they were three-banded ($\Delta_{[\text{shaded}-\text{open}]}$= -0.70 mm [-1.38; -0.04], **Fig. 1**) or five-banded

($\Delta_{[\text{shaded}-\text{open}]}$= -1.37mm [-2.07; -0.71], **Fig. 1**); unbanded snails were however of similar size in

both habitats ($\Delta_{[\text{shaded}-\text{open}]}$= -0.39 mm [-1.03; 0.33], **Fig. 1**).

Parasitological information

Snails from both populations and all morphs were infected by metazoan parasites (**Figs. 2-3**, see

Supporting Information S2 for detailed descriptive statistics about prevalence and infection

intensity). All trematodes found were live *Brachylaima* sp. metacercariae in snails' kidneys; no

live sporocysts were recorded and no trematodes of any stage were found trapped in shells.

Among the organs we examined, we detected live nematodes only within the lung and in the

space between the body and the shell; most of these nematodes were in the 1-3 mm size class

(297/371; 211/269 excluding the snail that died during the experiments). Based on this, we can

tentatively assume that most nematodes found belonged to the Rhabditidae family (Morand *et*

al., 2004; Wilson, 2012). In addition we frequently (**Fig. 3**) found nematodes encapsulated in

shells. No live nematodes were found in individuals from the shaded population (**Fig.3**), by

contrast to the open population where 65.6% of snails were infected (59/90, including the dead

snail). However, a high number of snails contained encapsulated nematodes in their shells, in

both populations (**Figs 2, 3**), indicating past exposure to nematodes even in the shaded

population. Only one snail (unbanded from the shaded habitat) harboured live mites (N = 93

mites), identified as *Riccardoella* sp. Three other snails (all five-banded from the shaded habitat)

had (unidentified) mites trapped in their shell (N = 1, 2 and 15 mites, **Fig. 2**). As mentioned in

Methods, one snail among 180 died during the experiments; it harboured 102 live nematodes,

more than 4 times the number of the most infected surviving snail (**Fig. 3**).

Figure 2. Examples of encapsulated parasites in *Cepaea nemoralis* shells. (A) mite; (B) nematode. Original
images, including other images of encapsulated parasites, are made available with the rest of the data
(see **Data availability**).

In the open population where live nematodes were found, nematode abundance was morph-
dependent: five-banded snails were on average infected by about a third of the nematodes
infecting other snails (= 0.29 [0.11; 0.52],   = 0.34 [0.12;
0.62]; **Table 1**, **Fig. 3**), mostly due to differences in prevalence between morphs (**Supporting**
Information S2). Five-banded snails also contained fewer trematodes, but only when compared
to three-banded snails and only clearly in the shaded habitat (= 0.19 [0.00;
0.57]; **Fig. 3**, **Supporting Information S3**). Parasite abundance (trematode or nematode) was not
linked to shell size (**Table 1**).

Figure 3. Parasite-related variables as a function of morph and population of origin (N = 179, one dead snail was excluded). (A) Proportion of snails having at least one nematode trapped in their shells; (B) Live nematode abundance; (C) Live *Brachylaima* sp. trematode abundance (metacercariae). For (B) and (C): grey dots are observed values, displayed along the posterior distributions of the average abundances based on the model described **Table 1** (black dots and segments are the means and 95% CI of the posteriors). Posterior predictions are based on fixed effects only, and the effects of shell size are averaged out. See **Supporting Information S3** for pairwise comparisons.

264 **Table 1**. Summary of posterior model coefficients for the parasite and behaviour multivariate model.

265 Individual-level correlations between traits are presented in **Table 2**. See **Methods** for model details, and

266 **Supporting Information S3** for pairwise comparisons among morph × population groups. Fixed effects of

267 population and size that are different from 0, based on 95% intervals are in bold.

	Nematode abundance*	Trematode abundance	Movement	Food intake
Fixed effects				
Intercept$_{[0\ bands]}$	0.84 [0.09; 1.52]	-0.94 [-2.44; 0.48]	-0.26 [-0.66; 0.16]	-0.22 [-0.61; 0.19]
Intercept$_{[3\ bands]}$	0.67 [-0.08; 1.38]	-0.10 [-1.54; 1.33]	0.16 [-0.24; 0.56]	-0.09 [-0.50; 0.31]
Intercept$_{[5\ bands]}$	-0.47 [-1.31; 0.32]	-1.26 [-2.72; 0.23]	-0.09 [-0.50; 0.31]	-0.04 [-0.45; 0.37]
Population = "shaded"$_{[0\ bands]}$	-–	-0.02 [-1.42; 1.39]	**0.71 [0.31; 1.08]**	**0.42 [0.04; 0.83]**
Population = "shaded"$_{[3\ bands]}$	-–	0.06 [-1.31; 1.34]	-0.23 [-0.61; 0.13]	0.22 [-0.18; 0.63]
Population = "shaded"$_{[5\ bands]}$	-–	-0.88 [-2.44; 0.66]	-0.08 [-0.47; 0.32]	0.13 [-0.26; 0.55]
Shell size (standardised)	-0.19 [-0.53; 0.13]	0.06 [-0.57; 0.72]	0.09 [-0.03; 0.21]	**0.23 [0.10; 0.35]**
Random effects SDs				
Test group SD	0.53 [0.00; 1.23]	1.95 [0.94; 3.19]	0.23 [0.00; 0.70]	0.21 [0.00; 0.67]
Individual-level SD	1.14 [0.85; 1.46]	2.57 [1.99; 3.21]	0.32 [0.04; 0.54]	0.56 [0.43; 0.70]
Residual SD	-–	-–	0.92 [0.82; 1.02]	0.8 [0.72; 0.88]

268 * The nematode model is only estimated for the "sun-exposed" population, since no live nematodes

269 were recorded in the "shaded" population

Snail behaviour

Both movement and food intake were repeatable at the individual level (see individual-level SDs Table 1), although adjusted repeatability (sensu Nakagawa & Schielzeth, 2010) was weaker and closer to 0 for movement ($r_{[movement]}$ = 0.11 [0.00; 0.23]; $r_{[food\ intake]}$ = 0.29 [0.15; 0.42]; **Supporting Information S4**).

Movement depended on both population of origin and shell band number (**Table 1**, **Fig. 4**, **Supporting Information S3**). In the shaded population, five-banded snails were on average less mobile than unbanded snails ($\Delta_{[5bands-0bands]}$ = -0.61 SDs [-1.00; -0.21]). In snails from the open habitat however, there was no such difference between morphs ($\Delta_{[5bands-0bands]}$ = 0.17 SDs [-0.19; 0.59]). This was due to unbanded snails being less mobile when coming from the open habitat than when coming from the shaded population ($\Delta_{[shaded-open]}$ = 0.71 SDs [0.31; 1.08]), while there was no clear between-population difference for five-banded snails ($\Delta_{[shaded-open]}$ = -0.08 SDs [-0.47; 0.32])(**Table 1**, **Fig. 4**). In any case, there was no detectable link between movement and shell size (**Table 1**).

Larger snails tended to eat more (**Table 1**). Source population had no effect on food intake if snails were three- ($\Delta_{[shaded-open]}$= 0.03 g [-0.02; 0.08], **Fig. 4**) or five-banded ($\Delta_{[shaded-open]}$= 0.02 g [-0.03; 0.07], **Fig. 4**), but similarly to movement, unbanded snails coming from the shaded population ate more on average than their counterparts from the open habitat ($\Delta_{[shaded-open]}$= 0.05 g [0.00; 0.10], **Fig. 4**).

Figure 4. Behaviours as a function of morph and population of origin. (A) Movement metric (1/latency to move beyond 10 cm, centered and scaled to unit 1SD); (B) Food intake. Grey dots are observed values, displayed along the posterior distributions of the average abundances based on the model described in **Table 1** (black dots and segments are the means and 95% CI of the posteriors). Observed values from the same individual are connected by a line. Posterior predictions are based on fixed effects only, and the effects of shell size are averaged out. See **Supporting Information S3** for pairwise comparisons.

Looking at the individual-level correlation matrix between traits (i.e. once fixed effects of shell

morph, size and population of origin are accounted for), food intake and nematode abundance

are negatively correlated, while all other correlations (including between nematode and

trematode abundance) are not markedly different from zero (**Table 2**). The food-nematode

correlation persists if we refit the nematode model to include data from the shaded population

(**-0.58 [-0.87; -0.29]**).

Table 2. Individual-level correlations between parasite abundances and behavioural traits. Correlations

involving nematode abundance are only based on individuals from the open population (see Methods);

results are not qualitatively different if the model is refitted to include all observations (see **Data**

availability). Correlations that are different from 0 (based on 95% intervals) are in bold.

	Nematode abundance	Trematode abundance	Movement
Trematode abundance	0.01 [-0.35; 0.37]	-–	-–
Movement	-0.24 [-0.73; 0.25]	-0.17 [-0.68; 0.28]	-–
Food intake	**-0.60 [-0.87; -0.30]**	-0.12 [-0.46; 0.19]	0.24 [-0.23; 0.71]

Discussion

To understand how individual variation in behaviour and infection interact, we studied

movement, food intake and natural parasite infection across two populations and three morphs

of the land snail *Cepaea nemoralis*. We found evidence of morph-dependent responses to

parasites, as well as links between parasite infection and behaviour. We additionally present

evidence of land snails encapsulating parasitic arthropods (mites) in their shell.

Snails of the darker morphs were smaller in the shaded population (**Fig. 1**). While the two

habitats differ along various abiotic dimensions (calcium availability in substrate,

microclimate…, see **Methods**), their documented effect on shell size are highly species- and

study-dependent, and there is no consistent evidence for or against their role in driving the

differences we observe (Goodfriend, 1986). Differences in parasite infection between the two
habitats might also influence body size, for instance if high parasite risk drives resource
allocation away from growth and towards immune defence (Rantala & Roff, 2005). In any case,
our models conditioned on size, meaning that any effect of morph/population we discuss below
is "size being equal". In addition, we did not find any effect of size itself on parasites or
movement (**Table 1**), meaning that the effects of morph/population we found are likely not
mediated by size differences.

One snail was found with live *Riccardoella* mites, hemolymph feeders with negative effects on
mollusc life-history and behavior (e.g. Schüpbach & Baur, 2008a). They are transmitted
horizontally during mating, but may find new hosts by leaving their host and following snail
mucus trails (Schüpbach & Baur, 2008b). However, their limited prevalence in our sample
prevented us from investigating their effects. Intriguingly, we found encapsulated mites in three
other snails (**Fig. 2**). Many land snails can defend themselves against nematode infection by
trapping them in their shell (Rae, 2017), including *Cepaea nemoralis* (this study, Williams & Rae,
2016; Rae, 2017). Based on extensive surveys, Rae (2017) wrote that encapsulation in the shell
is a specific response against nematode infection; our observations might contradict this
affirmation. *Riccardoella* infection is highly variable across populations or species, with reported
prevalences of 0% or near 0% being frequently observed (this study; Russo & Madec, 2013; Baur
& Baur, 2005; Segade *et al.,* 2013), which may explain why encapsulated mites were not
previously found. We need further studies to determine with certainty whether shell
encapsulation is a more general response to metazoan parasites or if it is a nematode-specific
response, as posited by Rae (2017), but occasionally and anecdotally catches other foreign
bodies.

A majority of snails in the open population were hosting nematodes, the majority of which were
likely Rhabditidae based on size and organ location (Morand *et al.,* 2004; Wilson, 2012).
Interactions between Rhabditidae and terrestrial molluscs are complex, including
phoresy/endozoochory, facultative parasitism, necromeny (feeding and reproducing on the host
snail's body after its death), and obligate larval parasites that actively infect snails by
e.g. entering the lung cavity (Morand *et al.,* 2004; Pieterse *et al.,* 2017; Sudhaus, 2018).

Delimiting these interaction types is complicated because even when they have no effects on

adult snails, these nematodes can have dramatically negative effects on juveniles (Grannell,

Cutler & Rae, 2021). By contrast to the open population, we found no live nematodes in any

snail from the shaded population (**Fig. 3**; note that our methods were not designed to detect

nematodes embedded in e.g. snail foot tissue). Thanks to nematodes trapped in shells (**Fig. 3**),

which integrate infections over time, we know parasitic nematodes are not truly absent from

the shaded population; they were simply not infecting snails at the time of sampling. The

"shaded" woodlot habitat is sheltered from the sun, more humid and likely colder than the

open one. Infections are often dependent on abiotic parameters, including temperature and

humidity, leading to seasonality (Altizer *et al.*, 2006). Differences in nematode infection

between the habitats may then be explained, at least partly, by differences in microclimate.

First, the different microclimates experienced by the two populations may favour different

nematode species within the soil community, differing in their tendency to parasitism. Second,

even if nematode communities are similar, conditions favourable to nematode activity,

development or infection may be reached only later in the year in the shaded habitat. For

instance, *Deroceras reticulatum* slugs were more negatively impacted by *Phasmarhabditis*

hermaphrodita infection when conditions were artificially warmed during winter (El-Danasoury

& Iglesias-Piñeiro, 2017), indicating that warmer conditions may favour nematode activity. In

any case, these results mean that any between-population differences in snail behaviour in the

present study need to be interpreted not only in relation with microhabitat and microclimate

differences, but also in terms of average infection levels.

In the open population where active nematodes were actually found, morph differences in

infection seem to validate our initial hypothesis of increased immune defence in darker snails

(**Fig. 3**). By contrast, no significant morph difference in immune defence was found in the closely

related *Cepaea hortensis* (Scheil *et al.*, 2013, 2014), but their endpoints were different

(phenoloxidase activity and mortality). In addition, our behavioural results (discussed below) are

also consistent with the hypothesis of morph-dependent immunity. These results are interesting

in the light of recent evidence that melanins may not be the pigments forming the dark bands in

C. nemoralis (Affenzeller *et al.*, 2020). Indeed, our initial hypothesis when designing this

experiment (in 2019) was based on the oft-seen association between immunity and melanin
(San-Jose & Roulin, 2018), including in snails (Coaglio *et al.*, 2018), although other pigments may
also have links to immunity (Tan *et al.*, 2020). While it is still likely that melanin-associated
pathways play a role in immune defence in *Cepaea* (Scheil *et al.*, 2014), it is probably not as
simple as "banded snails constitutively produce more melanin, and thus are better protected".
More studies are needed to understand the mechanisms linking shell morph to nematode
resistance.

Snails from all morphs and both populations were infected by *Brachylaima* trematodes (**Fig. 3**).
We previously documented *Brachylaima* infecting *Cornu aspersum* in the same area (Gérard *et*
al., 2020); finding these parasites a year later and in *C. nemoralis* may indicate that they are
well-implanted in local communities. Contrary to nematodes, we found no evidence that morph
or population had any influence on the abundance of trematodes (**Fig. 3**). This is consistent with
another study analyzing both *C. nemoralis* and *C. hortensis* (Żbikowska *et al.*, 2020). We must
acknowledge however that snail infections by *Brachylaima* vary seasonally, and that spring may
not be the infection peak (Gracenea & Gállego, 2017). Longitudinal studies, possibly combined
with experimental infections or cures, may help confirm whether morphs respond differently to
trematode risk.

Nematode and trematode numbers were uncorrelated (**Table 2**). Terrestrial gastropods are
commonly infected by both nematodes and trematodes at the same time (e.g. Segade *et al.*,
2013; Antzée-Hyllseth *et al.*, 2020). Our results extend these previous studies, and suggest that
despite sharing a host, nematode and trematode niches are sufficiently distinct within snails to
limit spatial and trophic competition. We reiterate however that studies more expansive in
space and time, including seasons and/or populations with higher prevalence and abundance of
trematodes and nematodes, are needed to confirm this. Combined with finer-grained studies
(e.g. at the parasite species level; Morand *et al.*, 2004), this would allow us to understand better
how parasites with similar or different niches within snails interact.

We found that movement behaviour in *C. nemoralis* was morph-dependent, but that this
dependency varied between populations: five-banded morphs were less mobile, but contrary to

Dahirel et al. (2021), this was only the case in the shaded population (**Fig. 4**). In the open

population, snail morphs are mobile at similarly low levels: five-banded snails do not change

behaviour between the two populations, while unbanded snails go down to five-banded snail

levels of mobility (**Fig. 4**). One of the main differences between the present study and Dahirel et

al. (2021) is the collection date. While here snails were collected several weeks after the onset

of spring activity and kept active throughout, in Dahirel et al. (2021) they were collected in late

fall and stored at cold temperatures inducing dormancy for months before being tested. This

difference may be especially relevant: winter or summer dormancy can lead to sharp decreases

in parasite infection in several parasite-snail systems (Solomon, Paperna & Matkovics, 1996;

Haeussler *et al.*, 2012; Segade *et al.*, 2013). We can thus reconcile the behavioral results from

Dahirel et al. (2021) from those of the present study if we assume that in the former

experiment, hibernation reduced parasite load across all morphs and populations, and that in

the present study, lower temperatures in the shaded habitat led to longer hibernation. We can

then interpret population differences in the present study in the light of their different infection

patterns. Five-banded snails' behaviour is almost unchanged between the two populations,

while unbanded snails from the "infected" population are slower than in the nematode-free

population (**Fig. 4**). This is again consistent with the idea that unbanded snails are more

vulnerable to parasites than banded ones. The question then becomes: is this reduction in

activity in infected unbanded snails simply the direct consequence of the costs of infection, or

does it reflect something more? For instance, infective nematodes may be able to follow mucus

trails from active snails (Andrus *et al.*, 2020). Reducing activity and the associated mucus

production may then be protective for snails, by limiting the risk of further infections.

Experimental manipulations of both snail infection and parasite risk in the environment may

help disentangle these two responses. Future studies including species-level identification of

nematodes will also be needed, to determine whether nematodes with more negative impacts

also trigger stronger behavioural responses, or if snails respond indiscriminately to nematode

abundance.

Food intake was influenced by nematode infection: highly parasitized snails ate less (**Table 2**).

Feeding reduction in nematode-infected individuals has been documented in other terrestrial

mollusc species (Antzée-Hyllseth *et al.*, 2020; Saeedizadeh & Niasti, 2020) and is an indication

that parasite infection is indeed costly for snails even if they are non-lethal, at least in adults

(Wilson *et al.*, 2000; Williams & Rae, 2016; Grannell *et al.*, 2021). By contrast, trematode

infection had no effect on behaviour, including feeding behaviour (**Table 2**). We however only

found snails infected by *Brachylaima* metacercariae (in the kidney). At the sporocyst stage,

Brachylaima parasitizes the digestive gland (e.g. Gérard *et al.*, 2020), which would likely have a

stronger effect on snail behaviour. Indeed, sporocysts can cause high damage, direct and

indirect, to that key organ for digestion and for energy storage (Segade *et al.*, 2011; Żbikowska

et al., 2020).

The fact that snail response to parasites seems to be morph-dependent for nematodes, but not

for trematodes, has interesting implications for host-parasite and snail-predator interactions in

a changing world. Trematodes infecting helicid land snails are trophically transmitted to

vertebrate definitive hosts including rodents and birds (Gérard *et al.*, 2020; Sitko & Heneberg,

2020). Visual selection by thrushes (*Turdus philomelos*) has been advanced as one of the key

explanations for the maintenance of *Cepaea* shell polymorphism (Jones *et al.*, 1977), and

predation by rodents may also be morph dependent (Rosin *et al.*, 2011; Rosin *et al.*, 2013). If

confirmed, our results suggest that morph-dependent predation confers no additional

trematode-avoiding advantage. In addition, snail morphs differ in their thermal responses:

lighter-coloured morphs are better adapted to heat, and increase in frequency with global

warming or urbanization (Jones *et al.*, 1977; Silvertown *et al.*, 2011; Ożgo *et al.*, 2017; Kerstes *et*

al., 2019). If lighter morphs are more vulnerable to infection by nematodes, then we need to

understand how snail immunity will interact with snail thermal tolerance to better predict how

snail and nematode dynamics will be altered in response to climate change (El-Danasoury &

Iglesias-Piñeiro, 2017; Aleuy & Kutz, 2020).

Acknowledgements

460   We thank participants to the 4th French Conference on Animal Ecophysiology (CEPA4, Rennes,

461   2019) for interesting discussions and questions. Funding for MD's position during this research

462   was obtained by Elodie Vercken (see Funding).

Data availability

Data and R scripts to reproduce all analyses presented in this manuscript are available on Github (https://github.com/mdahirel/cepaea-parasite-behaviour-2019) and archived in Zenodo (DOI: https://doi.org/10.5281/zenodo.6327265). Representative images of parasites trapped in shells are also archived in Zenodo (DOI: https://doi.org/10.5281/zenodo.6326305).

Funding

MD was a postdoctoral researcher funded by the French Agence Nationale de la Recherche (PushToiDeLa, ANR-18-CE32-0008), during the course of this research.

References

Affenzeller, S., Wolkenstein, K., Frauendorf, H. & Jackson, D.J. (2020). Challenging the concept that eumelanin is the polymorphic brown banded pigment in *Cepaea nemoralis*. *Sci. Rep.* **10**, 2442.

Aleuy, O.A. & Kutz, S. (2020). Adaptations, life-history traits and ecological mechanisms of parasites to survive extremes and environmental unpredictability in the face of climate change. *Int. J. Parasitol. Parasites Wildl.* **12**, 308–317.

Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M. & Rohani, P. (2006). Seasonality and the dynamics of infectious diseases. *Ecol. Lett.* **9**, 467–484.

Andrus, P., Ingle, O., Coleman, T. & Rae, R. (2020). Gastropod parasitic nematodes (*Phasmarhabditis* sp.) are attracted to hyaluronic acid in snail mucus by cGMP signalling. *J. Helminthol.* **94**, e9.

Antzée-Hyllseth, H., Trandem, N., Torp, T. & Haukeland, S. (2020). Prevalence and parasite load of nematodes and trematodes in an invasive slug and its susceptibility to a slug parasitic nematode compared to native gastropods. *J. Invertebr. Pathol.* **173**, 107372.

Barker, G.M. (Ed.). (2004). *Natural enemies of terrestrial molluscs*. Wallingford, UK: CABI.

Baur, A. & Baur, B. (2005). Interpopulation variation in the prevalence and intensity of parasitic
mite infection in the land snail *Arianta arbustorum*. *Invertebr. Biol.* **124**, 194–201.

Binning, S.A., Shaw, A.K. & Roche, D.G. (2017). Parasites and host performance: incorporating
infection into our understanding of animal movement. *Integr. Comp. Biol.* **57**, 267–280.

BRGM. (2005). Geological map at 1:50 000 (BD-CHARM-50), department: Deux-Sèvres (79,
France).

Bürkner, P.-C. (2017). brms: an R package for Bayesian multilevel models using Stan. *J. Stat.*
Softw. **80**, 1–28.

Butcher, A.R. (2016). Children, snails and worms: the *Brachylaima cribbi* story. *Microbiol. Aust.*
37, 30–33.

Cain, A.J. & Sheppard, P.M. (1954). Natural selection in *Cepaea*. *Genetics* **39**, 89–116.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M.,
Guo, J., Li, P. & Riddell, A. (2017). Stan: a probabilistic programming language. *J. Stat. Softw.* **76**,
1–32.

Coaglio, A.L., Ferreira, M.A.N.D., dos Santos Lima, W. & de Jesus Pereira, C.A. (2018).
Identification of a phenoloxidase- and melanin-dependent defence mechanism in *Achatina*
fulica infected with *Angiostrongylus vasorum*. *Parasites Vectors* **11**, 113.

Dahirel, M., Gaudu, V. & Ansart, A. (2021). Boldness and exploration vary between shell morphs
but not environmental contexts in the snail *Cepaea nemoralis*. *Ethology* 13.

Davies, M.S. & Knowles, A.J. (2001). Effects of trematode parasitism on the behaviour and
ecology of a common marine snail (*Littorina littorea* (L.)). *J. Exp. Mar. Biol. Ecol.* **260**, 155–167.

Debeffe, L., Morellet, N., Verheyden-Tixier, H., Hoste, H., Gaillard, J.-M., Cargnelutti, B., Picot,
D., Sevila, J. & Hewison, A.J.M. (2014). Parasite abundance contributes to condition-dependent
dispersal in a wild population of large herbivore. *Oikos* **123**, 1121–1125.

Douma, J.C. & Weedon, J.T. (2019). Analysing continuous proportions in ecology and evolution:
a practical introduction to beta and Dirichlet regression. *Methods Ecol. Evol.* **10**, 1412–1430.

Dupuis, J., Cariou, E., Coirier, B. & Ducloux, J. (1983). *Notice explicative, Carte géologique de la*
France à 1/50 000, feuille Niort (610). Orléans, France: BRGM.

El-Danasoury, H. & Iglesias-Piñeiro, J. (2017). Performance of the slug parasitic nematode
Phasmarhabditis hermaphrodita under predicted conditions of winter warming. *J. Pestic.*
Science **42**, 62–66.

Ezenwa, V.O., Archie, E.A., Craft, M.E., Hawley, D.M., Martin, L.B., Moore, J. & White, L. (2016).
Host behaviour–parasite feedback: an essential link between animal behaviour and disease
ecology. *Proc. Royal Soc. B* **283**, 20153078.

Fain, A. (2004). Mites (Acari) parasitic and predaceous on terrestrial gastropods. In *Natural*
enemies of terrestrial molluscs: 505–524. Barker, G.M. (Ed.). Wallingford, UK: CABI.

Furuta, E. & Yamaguchi, K. (2001). Haemolymph: blood cell morphology and function. In *The*
biology of terrestrial molluscs: 289–306. Barker, G.M. (Ed.). Wallingford, UK: CABI.

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. (2019). Visualization in Bayesian
workflow. *J. Royal Stat. Soc. A* **182**, 389–402.

Gérard, C., Ansart, A., Decanter, N., Martin, M.-C. & Dahirel, M. (2020). *Brachylaima* spp.
(Trematoda) parasitizing *Cornu aspersum* (Gastropoda) in France with potential risk of human
consumption. *Parasite* **27**, 15.

Giannelli, A., Cantacessi, C., Colella, V., Dantas-Torres, F. & Otranto, D. (2016). Gastropod-borne
helminths: a look at the snail–parasite interplay. *Trends Parasitol.* **32**, 255–264.

González-Santoyo, I. & Córdoba-Aguilar, A. (2012). Phenoloxidase: a key component of the
insect immune system. *Entomol. Exp. Appl.* **142**, 1–16.

Goodfriend, G.A. (1986). Variation in land-snail shell form and size and its causes: a review. *Syst.*
Zool. **35**, 204–223.

Gracenea, M. & Gállego, L. (2017). Brachylaimiasis: *Brachylaima* spp. (Digenea: Brachylaimidae)
metacercariae parasitizing the edible snail *Cornu aspersum* (Helicidae) in Spanish public
marketplaces and health-associated risk factors. *J. Parasitol.* **103**, 440–450.

Grannell, A., Cutler, J. & Rae, R. (2021). Size-susceptibility of *Cornu aspersum* exposed to the
malacopathogenic nematodes *Phasmarhabditis hermaphrodita* and *P. californica*. *Biocontrol Sci.*
Technol. **31**, 1149–1160.

Haeussler, E.M., Pizá, J., Schmera, D. & Baur, B. (2012). Intensity of parasitic mite infection
decreases with hibernation duration of the host snail. *Parasitology* **139**, 1038–1044.

Harrison, X.A. (2014). Using observation-level random effects to model overdispersion in count
data in ecology and evolution. *PeerJ* **2**, e616.

Jones, J.S., Leith, B.H. & Rawlings, P. (1977). Polymorphism in *Cepaea*: a problem with too many
solutions? *Annu. Rev. Ecol. Evol. Syst.* **8**, 109–143.

Kavaliers, M. (1992). Opioid systems, behavioral thermoregulation and shell polymorphism in
the land snail, *Cepaea nemoralis*. *J. Comp. Physiol. B* **162**, 172–178.

Kay, M. (2019). *tidybayes: tidy data and geoms for Bayesian models*.
https://doi.org/10.5281/zenodo.1308151

Kerstes, N.A.G., Breeschoten, T., Kalkman, V.J. & Schilthuizen, M. (2019). Snail shell colour
evolution in urban heat islands detected via citizen science. *Commun. Biol.* **2**, 264.

Köhler, H.-R., Capowiez, Y., Mazzia, C., Eckstein, H., Kaczmarek, N., Bilton, M.C., Burmester,
J.K.Y., Capowiez, L., Chueca, L.J., Favilli, L., Gomila, J.F., Manganelli, G., Mazzuca, S.,
Moreno-Rueda, G., Peschke, K., Piro, A., Cardona, J.Q., Sawallich, L., Staikou, A.E., Thomassen,
H.A. & Triebskorn, R. (2021). Experimental simulation of environmental warming selects against
pigmented morphs of land snails. *Ecol. Evol.* **11**, 1111–1130.

Lamotte, M. (1959). Polymorphism of natural populations of *Cepaea nemoralis*. *Cold Spring*
Harb. Symp. Quant. Biol. **24**, 65–86.

McElreath, R. (2020). *Statistical rethinking: a Bayesian course with examples in R and Stan*. 2nd
edition. Boca Raton, USA: Chapman and Hall/CRC.

McElroy, E.J. & de Buron, I. (2014). Host performance as a target of manipulation by parasites: a
meta-analysis. *J. Parasitol.* **100**, 399–410.

Meerburg, B.G., Singleton, G.R. & Kijlstra, A. (2009). Rodent-borne diseases and their risks for
public health. *Crit. Rev. Microbiol.* **35**, 221–270.

Morand, S., Wilson, M.J. & Glen, D.M. (2004). Nematodes (Nematoda) parasitic in terrestrial
molluscs. In *Natural enemies of terrestrial molluscs*: 525–557. Barker, G.M. (Ed.). Wallingford,
UK: CABI.

Morley, N.J. & Lewis, J.W. (2008). The influence of climatic conditions on long-term changes in
the helminth fauna of terrestrial molluscs and the implications for parasite transmission in
southern England. *J. Helminthol.* **82**, 325–335.

Nakagawa, S. & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biol. Rev.* **85**, 935–956.

O'Dwyer, K., Kamiya, T. & Poulin, R. (2014). Altered microhabitat use and movement of littorinid
gastropods: the effects of parasites. *Mar. Biol.* **161**, 437–445.

Ożgo, M. (2009). Current problems in the research of *Cepaea* polymorphism. *Folia Malacol.* **16**,
55–60.

Ożgo, M., Liew, T.-S., Webster, N.B. & Schilthuizen, M. (2017). Inferring microevolution from
museum collections and resampling: lessons learned from *Cepaea*. *PeerJ* **5**, e3938.

Pedersen, T.L. (2020). *patchwork: the composer of plots*. https://CRAN.R-
project.org/package=patchwork

Pieterse, A., Malan, A.P. & Ross, J.L. (2017). Nematodes that associate with terrestrial molluscs
as definitive hosts, including *Phasmarhabditis hermaphrodita* (Rhabditida: Rhabditidae) and its
development as a biological molluscicide. *J. Helminthol.* **91**, 517–527.

R Core Team. (2021). *R: a language and environment for statistical computing*. Vienna, Austria:
R Foundation for Statistical Computing.

Rae, R. (2017). The gastropod shell has been co-opted to kill parasitic nematodes. *Sci. Rep.* **7**,
4745.

Rantala, M.J. & Roff, D.A. (2005). An analysis of trade-offs in immune function, body size and
development time in the Mediterranean Field Cricket, *Gryllus bimaculatus*. *Funct. Ecol.* **19**, 323–
330.

Réale, D., Garant, D., Humphries, M.M., Bergeron, P., Careau, V. & Montiglio, P.-O. (2010).
Personality and the emergence of the pace-of-life syndrome concept at the population level.
Phil. Trans. R. Soc. B **365**, 4051–4063.

Rosin, Z.M., Kobak, J., Lesicki, A. & Tryjanowski, P. (2013). Differential shell strength of *Cepaea*
nemoralis colour morphs—implications for their anti-predator defence. *Naturwissenschaften*
100, 843–851.

Rosin, Z.M., Olborska, P., Surmacki, A. & Tryjanowski, P. (2011). Differences in predatory
pressure on terrestrial snails by birds and mammals. *J. Biosci.* **36**, 691–699.

Royauté, R., Berdal, M.A., Garrison, C.R. & Dochtermann, N.A. (2018). Paceless life? A meta-
analysis of the pace-of-life syndrome hypothesis. *Behav. Ecol. Sociobiol.* **72**, 64.

Russo, J. & Madec, L. (2013). Linking immune patterns and life history shows two distinct
defense strategies in land snails (Gastropoda, Pulmonata). *Physiol. Biochem. Zool.* **86**, 193–204.

Saeedizadeh, A. & Niasti, F. (2020). Response of grey slug to entomopathogenic nematodes.
Bragantia **79**, 447–456.

San-Jose, L.M. & Roulin, A. (2018). Toward understanding the repeated occurrence of
associations between melanin-based coloration and multiple phenotypes. *Am. Nat.* **192**, 111–
130.

Scheil, A.E., Hilsmann, S., Triebskorn, R. & Köhler, H.-R. (2013). Shell colour polymorphism,
injuries and immune defense in three helicid snail species, *Cepaea hortensis*, *Theba pisana* and
Cornu aspersum maximum. *Results Immunol.* **3**, 73–78.

Scheil, A.E., Hilsmann, S., Triebskorn, R. & Köhler, H.-R. (2014). Shell colouration and parasite
tolerance in two helicoid snail species. *J. Invertebr. Pathol.* **117**, 1–8.

Schilthuizen, M. (2013). Rapid, habitat-related evolution of land snail colour morphs on
reclaimed land. *Heredity* **110**, 247–252.

Schüpbach, H.U. & Baur, B. (2008a). Parasitic mites influence fitness components of their host,
the land snail *Arianta arbustorum*. *Invertebr. Biol.* **127**, 350–356.

Schüpbach, H.U. & Baur, B. (2008b). Experimental evidence for a new transmission route in a
parasitic mite and its mucus-dependent orientation towards the host snail. *Parasitology* **135**,
1679–1684.

Segade, P., Crespo, C., García, N., García-Estévez, J.M., Arias, C. & Iglesias, R. (2011).
Brachylaima aspersae n. sp. (Digenea: Brachylaimidae) infecting farmed snails in NW Spain:
Morphology, life cycle, pathology, and implications for heliciculture. *Vet. Parasitol.* **175**, 273–
286.

Segade, P., García-Estévez, J., Arias, C. & Iglesias, R. (2013). Parasitic infections in mixed system-
based heliciculture farms: dynamics and key epidemiological factors. *Parasitology* **140**, 482–497.

Silvertown, J., Cook, L., Cameron, R., Dodd, M., McConway, K., Worthington, J., Skelton, P.,
Anton, C., Bossdorf, O., Baur, B., Schilthuizen, M., Fontaine, B., Sattmann, H., Bertorelle, G.,
Correia, M., Oliveira, C., Pokryszko, B., Ożgo, M., Stalažs, A., Gill, E., Rammul, Ü., Sólymos, P.,
Féher, Z. & Juan, X. (2011). Citizen science reveals unexpected continental-scale evolutionary
change in a model organism. *PLOS ONE* **6**, e18927.

Sitko, J. & Heneberg, P. (2020). Emerging helminthiases of song thrush (*Turdus philomelos*) in
Central Europe. *Parasitol Res.* **119**, 4123–4134.

Solomon, A., Paperna, I. & Matkovics, A. (1996). The influence of aestivation in land snails on
the larval development of *Muellerius* cf. *capillaris* (Metastrongyloidea: Protostrongylidae). *Int. J.*
Parasitol. **26**, 363–367.

Sudhaus, W. (2018). Dispersion of nematodes (Rhabditida) in the guts of slugs and snails. *Soil*
Org. **90**, 101–114.

Surmacki, A., Ożarowska-Nowicka, A. & Rosin, Z.M. (2013). Color polymorphism in a land snail
Cepaea nemoralis (Pulmonata: Helicidae) as viewed by potential avian predators.
Naturwissenschaften **100**, 533–540.

Tan, K., Zhang, H., Lim, L.-S., Ma, H., Li, S. & Zheng, H. (2020). Roles of carotenoids in
invertebrate immunology. *Front. Immunol.* **10**.

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. (2021). Rank-normalization,
folding, and localization: an improved $\hat{R}$ for assessing convergence of MCMC (with discussion).
Bayesian Anal. **16**, 667–718.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J.,
Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K. & Yutani, H.
(2019). Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686.

Williams, A. & Rae, R. (2016). *Cepaea nemoralis* (Linnaeus, 1758) uses its shell as a defence
mechanism to trap and kill parasitic nematodes. *J. Molluscan Stud.* **82**, 349–350.

Williams, S.T. (2017). Molluscan shell colour. *Biol. Rev.* **92**, 1039–1058.

Wilson, M.J. (2012). Pathogens and parasites of terrestrial molluscs. In *Manual of techniques in*
invertebrate pathology: 427–439. Lacey, L.A. (Ed.). Oxford; New York: Academic Press.

Wilson, M.J., Hughes, L.A., Hamacher, G.M. & Glen, D.M. (2000). Effects of *Phasmarhabditis*
hermaphrodita on non-target molluscs. *Pest. Manag. Sci.* **56**, 711–716.

Wynne, R., Morris, A. & Rae, R. (2016). Behavioural avoidance by slugs and snails of the parasitic
nematode *Phasmarhabditis hermaphrodita*. *Biocontrol Sci. Technol.* **26**, 1129–1138.

Żbikowska, E., Marszewska, A., Cichy, A., Templin, J., Smorąg, A. & Strzała, T. (2020). *Cepaea*
spp. as a source of *Brachylaima mesostoma* (Digenea: Brachylaimidae) and *Brachylecithum* sp.
(Digenea: Dicrocoeliidae) larvae in Poland. *Parasitol. Res.* **119**, 145–152.

0.25 mm