Title: Pudendal, but not tibial, nerve stimulation modulates vulvar blood perfusion in anesthetized

rodents

Authors

Elizabeth C. Bottorff, MS[1,2]

Tim M. Bruns, PhD[1,2]

Institution

Full institution details for each author: name of institution, department, city and country.

1. University of Michigan, Biomedical Engineering Department, Ann Arbor, MI, USA

2. University of Michigan, Biointerfaces Institute, Ann Arbor, MI, USA

Corresponding Author Name and Email address:

Tim Bruns

bruns@umich.edu

Conflict of Interest: None

Acknowledgement:

We thank Tim Brown of Moor Instruments for assistance with data collection and analysis, Eric Kennedy

and Ahmad Jiman for their help in experiment preparation, and the University of Michigan Unit for

Laboratory Animal Medicine for animal husbandry. This study was supported in part by NIH Award

T32NS115724.

Abstract

Background: Preclinical studies have shown that neuromodulation can increase vaginal blood perfusion,

but the effect on vulvar blood perfusion is unknown.

Aim: Our goal was to investigate whether pudendal or tibial nerve stimulation can evoke an increase in

vulvar blood perfusion.

Methods: We used female Sprague-Dawley rats for non-survival procedures under urethane anesthesia.

We measured perineal blood perfusion in response to twenty-minute periods of pudendal and tibial nerve

stimulation using laser speckle contrast imaging (LSCI). After a thoracic-level spinalization and a rest

period, we repeated each stimulation trial.

Outcomes: We calculated average blood perfusion before, during, and after stimulation for three perineal

regions (vulva, anus, and inner thigh), for each nerve target and spinal cord condition.

Results: We observed a significant increase in vulvar, anal, and inner thigh blood perfusion during

pudendal nerve stimulation in spinally intact and spinalized rats. Tibial nerve stimulation had no effect on

perineal blood perfusion for both spinally intact and spinalized rats.

Clinical Implications: This study provides further support for neuromodulation as a treatment for women

with sexual arousal disorders.

Strengths and Limitations: This is the first study to examine vulvar hemodynamics with LSCI in response

to nerve stimulation. Studies in unanesthetized animal models with genital arousal disorders are needed

to obtain further insights into the mechanisms of neural control over genital hemodynamics.

Conclusion: This study demonstrates that pudendal nerve stimulation modulates vulvar blood perfusion,

indicating the potential of pudendal neuromodulation to improve genital blood flow as a treatment for

women with sexual dysfunction.

Keywords: blood flow, neuromodulation, pudendal nerve, tibial nerve, vulva, laser speckle imaging

Introduction

Female sexual health is an important determinant in quality of life, contributing to both increased meaning in life and general well-being[1]. Unfortunately, approximately 40% of women suffer from female sexual dysfunction (FSD)[2]. Female sexual dysfunction can present in multiple domains, which can be thought of as physiological (arousal, lubrication, orgasm, pain) and psychological (satisfaction and desire). Due to limited research regarding the etiology of FSD and basic female anatomy, current treatment options for FSD are very limited, especially for women who have deficits in the physiological domains such as arousal[3]. Bremelanotide and flibanserin, both approved to treat hypoactive sexual desire disorder (HSDD), have shown mild efficacy in improving sexual desire and arousal[4,5]. However, bremelanotide has been associated with a high incidence of adverse events[6] and a review of flibanserin, suggests that it has minimal clinical benefit[7]. Sildenafil citrate has been studied as a treatment for female arousal disorders. Studies with sildenafil have demonstrated an increase in clitoral vasocongestion and an association with increased sexual satisfaction[8]. Sildenafil was also used to study vaginal vasocongestion in treatment and placebo groups during erotic visual stimulation[9]. Although this study reported significant increases in vasocongestion for the treatment group, there were no differences in subjective arousal between the sildenafil and placebo groups. The lack of consistent improvements in subjective arousal as well as a high incidence of adverse events led to sildenafil citrate no longer being pursued as a treatment option[10].

Neuromodulation is a potential treatment option for women with FSD. Sacral neuromodulation (SNM) is a standard treatment for overactive bladder and fecal incontinence[11,12]. SNM delivers electrical stimulation to sacral nerves, which contain somatic and sympathetic nerve fibers of the pelvic organs. Clinical studies have found that sexual function can improve in women receiving SNM for bladder function[13–17]. Similar improvements in sexual function have been seen in clinical studies that used tibial nerve stimulation to treat bladder dysfunctions[18,19]. The sexual health benefits of neuromodulation have been studied in women without concomitant pelvic disorders. In a recent study, transcutaneous electrical nerve stimulation (TENS) of the tibial nerve or dorsal genital nerve (DGN), a distal branch of the pudendal nerve, improved survey-reported female sexual function index (FSFI) scores, a standardized metric for evaluating female sexual function [20]. These subjects reported an improvement in their overall FSFI score

and their individual arousal and orgasm sub-scores. Subjects receiving DGN stimulation also saw

improvements in sexual satisfaction. There is a lack of physiological measurements of sexual health and

so the mechanisms by which pudendal and tibial nerve stimulation improve sexual function are not

completely understood. One possible mechanism is that nerve stimulation modulates genital blood flow,

which is necessary to facilitate vasocongestion, lubrication, and sexual receptivity[21]. This is supported by

preclinical studies showing that stimulation of the tibial nerve for 30 minutes can result in a transient

increase in vaginal blood flow 20-35 minutes after stimulation onset[22]. Preclinical studies have also shown

an increase in vaginal blood flow in response to pudendal nerve stimulation, with peak responses

occurring anywhere from 1-2 minutes[23] or up to 30 minutes[24] after stimulation onset depending on

stimulation parameters. Additionally, it is well established that stimulation of the somatic pudendal

nerve[25–27] and tibial nerve[27–29] can modulate spinal control over the bladder and bowel, leading to

improvement in dysfunctional states. We hypothesized that a similar spinal reflex exists for genital

vasocongestion. Both pudendal and tibial nerve stimulation have the potential to improve sexual function

and have specific translational advantages, including the ease of access and existing clinical use for

pelvic dysfunctions[18,19,26,30], however a greater understanding of the underlying mechanisms and neural

circuitry is needed.

Female sexual dysfunction often presents in women with spinal cord injuries (SCI). Women with

SCI retain varying aspects of sexual function, depending on their injury type. For example, research has

shown that sensory impairment to the T11-L2 dermatomes, but not injury level, is associated with

decreased genital arousal[31]. Although paraplegic patients report that regaining sexual function after SCI is

a priority[32,33], women with SCI have the same FSD treatment options as non-neurogenic women. These

treatment options do not consider the impact of neurological dysfunction. Once the neural pathways

controlling genital arousal are better understood, clinicians can use knowledge of the patient's injury to

determine if electrical stimulation is able modulate these pathways to improve genital arousal in women

with neurogenic FSD. However, it is first necessary to identify how different nerve targets of

neuromodulation modulate genital arousal. In healthy women, genital arousal is the physiological

response to receptive sexual stimuli and is characterized by an increase in genital blood flow,

engorgement of the genitals, and the production of lubrication[34]. Most research studying female sexual

responses use vaginal photoplethysmography to measure genital arousal[35]. However, some researchers

are turning towards more non-invasive methods such as laser doppler imaging or laser speckle contrast

imaging (LSCI). These laser-based methodologies are becoming more useful in measuring female sexual

arousal because they show a higher concordance between genital and subjective arousal[36,37]. However,

the importance of genital-subjective arousal synchrony is not well understood[3] and some models for the

sexual response cycle suggest that genital arousal is necessary for satisfactory sexual intercourse,

regardless of subjective arousal[38,39].

In this preclinical study we used LSCI to measure blood perfusion changes in the perineal region

of female rats in response to tibial and pudendal nerve stimulation. Additionally, we looked at how this

blood perfusion response changes after thoracic level spinalization. Although LSCI has been used to

assess genital arousal in clinical studies[36,37,40,41], this is the first time it has been used to measure perineal

blood perfusion in animals. We found that during the stimulation epoch, pudendal, but not tibial, nerve

stimulation causes an increase in perineal blood perfusion. There were no differences in perineal blood

perfusion after stimulation was delivered, for either nerve target, when compared with baseline. We found

that these response patterns were not altered by thoracic spinal transection and are likely due to direct

efferent activation of the pelvic floor musculature.

Methods

Animal Surgery

All experimental procedures were approved by the governing Institutional Animal Care and Use

Committee (IACUC) in accordance with the National Institutes of Health's guidelines for the care and use

of laboratory animals. We designed the primary experimental protocol prior to starting the study. The

study protocol was not pre-registered. We performed non-survival procedures on 17 nulliparous female

Sprague-Dawley rats (Charles River Breeding Labs, Wilmington, MA, USA) weighing 0.25 to 0.30

kilograms. Female Sprague-Dawley rats are a common preclinical model for studying sexual arousal

because their physiological response to sexual stimuli is similar to that found in humans (e.g., increases

in vaginal blood flow, length, and pressure)[42]. We used ten animals for primary experimental testing, as

described below. We selected this sample size based on prior animal studies with similar objectives[24,43,44].

Animals were housed in ventilated cages under controlled temperature, humidity, and photoperiod (12-h

light/dark cycle), and provided laboratory chow (5L0D, LabDiet, St. Louis, MO, USA). Animals were

anesthetized using intraperitoneal urethane (1.5 g/kg), a commonly used anesthetic in rodent surgeries

studying pelvic organ function[23,28,45,46]. Sufficient anesthetic depth was confirmed once the animal no

longer had a toe pinch response. We used a heating pad to maintain body temperature and monitored

vital signs (heart rate, respiration rate, and oxygen saturation levels) every 15 minutes. We used vital

signs as humane endpoints. We performed vaginal cytology prior to surgical access to determine estrous

stage[47], to examine if one stage (e.g. estrus) better facilitates a perineal blood flow response.

With the rat in the prone position, we made a dorsal midline incision through the skin 3-4 cm

rostral to the base of the tail. Then we extended the incision laterally on the right side, and the ischiorectal

fossa was separated. We used retractors to keep the musculature open and isolated the pudendal nerve

using forceps. We placed a bipolar nerve cuff with stranded stainless-steel wire electrodes (400 μm

diameter; Cooner Wire Co, Chatsworth, CA, SA) and silicone elastomer tubing (0.5-mm inner diameter;

Dow Corning, Midland, MI, USA) around both sensory and motor branches of the pudendal nerve within

the ischiorectal fossa, proximal to the division of the motor branch into its dorsal and ventral branches[48].

We then closed the incision and moved the animal to a supine position. We then placed a percutaneous

electromyogram (EMG) wire (stainless steel, 50 μm, MicroProbes, Gaithersburg, MD) subcutaneously,

parallel to the tibial nerve and ipsilateral to the pudendal nerve cuff. We measured perineal blood

perfusion with a laser speckle contrast imaging (LSCI) system (MOORFLPI-2, Moor Instruments,

Wilmington, DE). We aimed the laser at the perineal region, angled downward at 25 degrees, and

positioned such that the path of the laser traveled 15 to 20 cm to the animal (Figure 1A).

Experimental Protocol

We delivered stimulation using an isolated pulse generator (Model 2100, A-M Systems, Sequim,

WA). We determined the motor threshold (MT) for stimulating the pudendal and tibial nerves by slowly

increasing the stimulation amplitude until a motor response (anal or toe twitch respectively) was

observed. Each animal went through a series of 45-minute trials. For each trial, we used LSCI to measure

perineal blood perfusion at the maximum sampling rate of 0.25 Hz, using the temporal processing

settings of the MOORFLPI-2, for 5 minutes prior to stimulation, during 20 minutes of stimulation, and for

20 minutes after stimulation. We delivered stimulation to one of the nerve targets using biphasic,

rectangular pulses (0.2-ms pulse width) at 20 Hz and twice the MT. We repeated the recording protocol

for the other nerve target. After both trials, we placed the rat in the prone position and performed a

laminectomy. We used a rostral/caudal incision to expose T8-10 level vertebrae. After removal of the

dorsal processes, we used a scalpel blade to transect the spinal cord. We closed the musculature with

sutures and closed the skin with skin staples. We then moved the rat to the supine position and allowed it

to rest for at least 15 minutes. Next, we re-assessed the MT and repeated the trials in the same order as

the intact spinal cord trials. After all experimental procedures were complete, we euthanized the animal

with an intraperitoneal injection of sodium pentobarbital (250-300 mg/kg). In the first rat, tibial nerve

stimulation was performed first. We then semi-randomized the order of the stimulated nerve across

following animals to arrive at balanced groups upon study termination. Investigators were not blind to

stimulation order.

We added two sets of supplemental experiments to the protocol after the study began. In some

initial experiments, we saw the blood perfusion signal spontaneously increase over 100 APU for 1 to 3

minutes, at 5-to-6-minute intervals. These increases were too abrupt to be considered a physiological

response in blood perfusion, so we performed another set of experiments to examine LSCI data sampling

(n = 2). The experimental procedure in these experiments followed the same set of 4 trials (pudendal and

tibial stimulation, before and after spinal cord transection), except that the blood perfusion was sampled

at 1 Hz or 25 Hz using the spatial processing settings instead of the maximum sampling rate of 0.25 Hz

with the temporal processing setting. In the second set of supplemental experiments (n = 3), we

transected the pudendal nerve next to the cuff electrode between stimulation trials with an intact spinal

cord. We looped a suture string around the nerve during cuff placement and pulled it tightly to cut the

nerve. We transected the pudendal nerve a different way in each of these experiments: once proximal to

the cuff, once distal to the cuff, and once distal to the cuff with a partial and then complete distal cut.

Data Analysis

We performed all data and statistical analyses using moorFLPI-2 Research Software (Software-

MOORFLPI2-3VX, Moor Instruments, Wilmington, DE) and MATLAB (Mathworks, Natick, MA, USA). We

drew regions of interest (ROIs) around the vulva, the anus, and a region on the inner thigh for each LSCI

trial analysis in moorFLPI-2. We drew a circular ROI centered on the vaginal orifice, with a diameter equal

to the width of the tissue mound surrounding the urethra. We then centered two identical sized ROIs on

the anus and the inner thigh on the contralateral side of where stimulation was delivered, such that the

three ROIs formed an equilateral triangle (Figure 1B). We calculated average perfusion units within each

ROI (vulvar, anal, and inner thigh) and extracted the signal for each 45-minute trial. Each signal is

referred to as vulvar blood perfusion (VBP), anal blood perfusion (ABP), and inner thigh blood perfusion

(ITBP). We observed an artifact when stimulation was turned on and removed it by eliminating the data

from 20 seconds before to 60 seconds after stimulation onset. We then calculated the temporal mean for

each ROI before, during, and after stimulation. We used a linear regression to determine the impact of

estrous stage, stimulation order, experiment number, and weight on the results. Estrous stage and

stimulation order were coded as binary values in this analysis. We made comparisons between average

VBP, ABP, and ITBP before and after spinalization as well as across stimulation epochs (before, during,

and after the stimulation periods) using t-tests (alpha = 0.05). Each animal served as their own control.

Results

We considered animals that completed all 4 nerve stimulation trials (n = 10) as experimental units

and used them for full data analysis. Three animals were used to perform supplementary nerve

transection trials. Two animals had blood perfusion measured using different LSCI parameters to test for

aliasing. The remaining two animals died prematurely during the second stimulation trial and were

excluded from analysis. Estrous stage, stimulation order, experiment number, and weight had no impact

on results. Detailed experimental demographics can be found in Table 1.

Intact Spinal Cord Recordings

During pudendal nerve stimulation trials (Figure 2A), the average VBP at baseline was 184 ± 44

APU, which increased to 331 ± 129 APU during stimulation (+80.9 ± 64.2% change from baseline) and

fell to 185 ± 55 APU after stimulation. Similarly, the average ABP was 172 ± 40 prior to stimulation, 348 ±

164 during stimulation (+97.6 ± 74.4%), and 169 ± 44 APU after stimulation ended. The average ITBP

was 94 ± 33, 154 ± 67 (+67.0 ± 63.7%), and 92 ± 35 before, during, and after pudendal stimulation. The

average blood perfusion during pudendal nerve stimulation for all ROIs was significantly higher than at

baseline and after stimulation ($p < 0.01$).

The average VBP for tibial nerve stimulation trials (Figure 2C) was 183 ± 50 APU prior to

stimulation, 192 ± 48 APU during stimulation ($+5.7 \pm 11.9\%$), and 181 ± 52 APU after stimulation ended

(Figure 3C). For the same trials, ABP was 163 ± 42, 163 ± 42 ($+0.4 \pm 8.0\%$), and 162 ± 47 APU before,

during, and after stimulation. ITBP had mean values for the three periods of 85 ± 26, 93 ± 34 ($+7.8 \pm$

22.5%), and 85 ± 31 APU. Tibial nerve stimulation did not elicit significant changes in any ROIs during

stimulation, and the average blood perfusion was significantly lower in all ROIs compared to pudendal

nerve stimulation ($p < 0.001$). Neither pudendal nor tibial stimulation elicited changes in perineal blood

perfusion that persisted after stimulation was turned off.

Post-Spinalization Recordings

Pudendal nerve stimulation trials after spinalization (Figure 2B) had an average VBP of 193 ± 61,

337 ± 131 ($+74.5 \pm 42.8\%$ change from post-spinalization baseline), and 188 ± 65 APU before, during

and after stimulation respectively. ABP for the same trials was 165 ± 39, 316 ± 161 ($+86.6 \pm 59.1\%$), and

167 ± 37 APU. Average ITBP was 88 ± 30, 153 ± 63 ($+74.4 \pm 60.7\%$), and 86 ± 28 APU (Figure 3B).

There were no significant differences between average blood perfusion before and after spinalization for

each ROI.

Tibial nerve stimulation trials had no significant differences after spinalization (Figure 2D). VBP

was 193 ± 76, 211 ± 87 ($+9.1 \pm 11.8\%$), and 196 ± 65 APU prior to, during, and after stimulation. ABP

was 168 ± 46, 171 ± 41 ($+2.6 \pm 7.8\%$), and 163 ± 32 APU for the same epochs. ITBP was 83 ± 29, $93 \pm$

45 ($+10.0 \pm 21.6\%$), and 79 ± 28 APU (Figure 3D).

Sampling Frequency Experiments

In the sampling frequency experiments, we sampled blood flow using the LSCI system at 1 Hz for

the first 4 minutes (no stimulation) and then at 25 Hz for 2 minutes (1 minute stimulation off, 1 minute on)

to capture any artifact resulting from stimulation being turned on. We then reduced the sampling rate to 1

Hz for 18 minutes during stimulation. Then, we increased the sampling rate to 25 Hz for 3 minutes (2

minutes stimulation on, 1 minute off) to capture the transition during stimulation cessation. Finally, we

used a sampling rate of 1 Hz for the final 19 minutes (stimulation off). In these two experiments, we

observed regular, gradual oscillations in blood perfusion for each ROI that occurred at 1–3-minute

intervals and were 100 to 300 APU in amplitude (e.g., Figure 4B). These oscillations were similar in width

and amplitude as occasional rapid spontaneous increases in blood flow observed in 10 trials across 4

experiments in the main cohort (e.g., Figure 4A). This similarity suggests that the rapid changes during

these trials were an artifact due to the low sampling rate. The stimulation artifact period during these

experiments had the same features as in other experiments, with a brief, high signal period (generally 1

second or less in length), supporting our plan of removing that period during analyses.

Nerve Transection Experiments

We performed three experiments to examine the contributions of afferent and efferent pathways

to the blood perfusion response to pudendal nerve stimulation. For the two pudendal nerve transection

experiments with full cuts (one proximal and one distal to the stimulation location), the increases in VBP,

ABP, and ITBP were abolished and there were no changes in blood perfusion for the duration of the trial.

For the partial pudendal nerve transection experiment, there were still small, but noticeable increases in

VBP, ABP, and ITBP during stimulation (+35.4%, +17.0%, and +43.8% change from baseline) which

returned to baseline after stimulation was ceased. After complete pudendal nerve transection, we

observed no changes in blood perfusion during stimulation for any of the ROIs (+0.9%, -6.5, and +1.0%

change for VBP, ABP, and ITBP respectively).

Discussion

In this study, we examined perineal blood flow changes in response to pudendal and tibial nerve

stimulation. Additionally, we studied the impact of spinalization on these blood flow responses. This is the

first study of its kind to use LSCI to measure perineal blood flow in animals. In intact animals, we saw a

transient increase in perineal blood flow (VBP and ABP) during pudendal nerve stimulation that returned

to baseline immediately after stimulation was turned off (Figure 3). Tibial nerve stimulation in intact

animals did not impact blood flow during or after stimulation was delivered (Figure 2C). After spinal cord

transection, pudendal and tibial nerve stimulation had the same effect as during intact spinal cord trials

(Figure 2C/D).

Perineal blood perfusion increases during pudendal nerve stimulation are likely attributable to the

direct activation of motor efferents, causing the anal and urethral sphincter to contract immediately after

stimulation onset[49]. This muscle activation causes a generalized increase of blood flow to the entire

perineal region. Spinal cord transection (Figure 3B) had no impact on the pudendal stimulation response,

excluding the involvement of a supraspinal pathway. Transecting the pudendal nerve distally abolished

the response immediately, indicating that the neural pathways involved are likely direct efferents, and not

a genital somato-visceral spinal reflex. Furthermore, studies demonstrating a potential genital somato-

visceral reflex that increases vaginal blood perfusion have a response that persists after stimulation is

turned off[23,24], which our VBP responses did not. Transecting the pudendal nerve proximally in one

experiment unexpectedly abolished the perineal blood perfusion response. It is possible that we damaged

the nerve during suture placement or transection. Experimental limitations prevented a repetition of this

trial. Other experimental limitations include our use of anesthesia, which may have dampened the effect

of stimulation on vulvar blood flow, and the duration of stimulation. It is possible that repeated or longer

stimulation sessions are required to evoke a non-direct response, increasing vulvar blood perfusion.

Prior preclinical pudendal nerve stimulation studies have used short duration (20 seconds) and

long duration (30 minutes) stimulation and measured the blood perfusion response using laser doppler

flowmetry (LDF) probes placed on the interior vaginal wall[23,24]. Both studies found an increase in vaginal

blood flow that either began or was transiently maintained after stimulation was turned off. Vaginal blood

flow gradually increased, peaked, and gradually returned to baseline during a 15-second to two-minute

period. Preclinical experiments have also demonstrated the potential for tibial nerve stimulation to

modulate vaginal blood flow[22,50] and that estrogen is necessary to preserve this response[50]. It is not

unexpected that a balanced gonadal hormone milleu is necessary for a genital blood flow response,

however little research has been done to assess the direct impact of peripheral nerve stimulation on

gonadal hormones. One such study found that tibial nerve stimulation did not have an impact on serum

estradiol[50], and we hypothesize that pudendal nerve stimulation may not have a direct impact on gonadal

hormones either. We expected to see increases in perineal blood perfusion in our experiments, however

neither pudendal nor tibial nerve stimulation had an effect on perineal blood perfusion that persisted

beyond the stimulation epoch. The vagina receives its blood supply from the uterine, vaginal, and internal

pudendal arteries[51]. In contrast, the perineal region, which includes the vulva and anus, is only supplied

by the internal pudendal arteries[52]. It is possible that pudendal nerve stimulation elicited changes in blood

perfusion of internal tissues, such as the vagina, that could not be measured using LSCI.

Genital blood flow in women has been most commonly assessed with vaginal

photoplethysmography (VPP)[35]. However, more researchers are using non-invasive techniques such as

Laser Doppler Imaging (LDI)[36,37,41] and LSCI[40] because they report higher levels of agreement between

genital arousal, subjective arousal, and lubrication. Studies using LDI or LSCI to assess genital arousal

have thus far been made in clinical settings in which participants receive visual sexual stimulation. Nerve

stimulation may not be sufficient to evoke a lasting vulvar blood perfusion response in an anesthetized

model of genital arousal. This suggests that vulvar blood perfusion needs further investigation before it is

used in anesthetized animals as a direct measurement of genital arousal.

The absence of blood perfusion changes during tibial nerve stimulation (Figure 2, 3) is not entirely

unexpected, as the tibial nerve innervates the lower leg and does not directly innervate the perineal

region[53]. Spinal cord transection did not change the lack of response. Researchers have previously found

that vaginal blood flow, measured with LDF, can increase in response to tibial nerve stimulation[22,50]. The

absence of blood perfusion changes in these experiments could be attributed to a difference in recording

areas (vagina vs vulva), just as in pudendal nerve stimulation trials. Clinical studies have demonstrated

that tibial nerve stimulation can improve bladder dysfunction in both non-neurogenic and

neurogenic[18,19,54,55] populations in weekly percutaneous stimulation sessions. In those studies, tibial nerve

stimulation improved bladder storage, a function controlled by the sympathetic nervous system. However,

healthy sexual functioning requires coordination of the sympathetic, parasympathetic, and somatic

nervous systems and it is the parasympathetic pathways that cause smooth muscle relaxation and

vaginal vasocongestion. Different stimulation parameters or repeated stimulation sessions (as is common

for clinical treatment) may cause a change in VBP similar to perfusion changes observed in vaginal

tissue.

We did not expect to see changes in ITBP during pudendal nerve stimulation, as the pudendal

nerve does not directly innervate the superficial regions of the inner thigh. Spinalization did not have an

impact on the average ITBP, across all stimulation epochs (Figure 3). Distal pudendal nerve transection

abolished the increase in ITBP observed in intact animals during pudendal nerve stimulation. The

pudendal nerve does not directly innervate any regions of the leg, but one cadaver study discovered that

the sciatic artery can arise from the internal pudendal artery[56]. It is possible that activation of motor

efferents caused blood flow to increase in pudendal arteries that then branch off and supply the inner

thigh. Thus, the increases in ITBP are likely attributable to general blood flow to the region due to the

contraction of the pelvic floor muscles placing an increased demand on blood supply, further supporting

our conclusion that the increases in vulvar blood perfusion are due to somatic activation of the pelvic floor

musculature.

This study sought to assess the impact of two potential treatment modalities for FSD (pudendal

and tibial nerve stimulation) on perineal blood perfusion. We theorize that improvements in FSD

symptoms are related to improvements in genital arousal and that it is possible to modulate genital

arousal via pudendal or tibial nerve stimulation. Both treatment modalities have demonstrated potential in

treating FSD. The dorsal genital nerve, a distal branching of the pudendal nerve, is a promising target

because it can be accessed superficially, and stimulation has shown to modulate sexual function[20]. Tibial

nerve stimulation is ideal for similar reasons as it is both superficial and easy to access, but more work is

necessary to understand how it modulates genital blood flow. However, this preliminary study did not find

any lasting changes in genital arousal as measured by perineal blood perfusion. We suspect that this

inconsistency with prior preclinical studies is due to a difference in blood perfusion location and that the

responses we did observe were due to muscle contractions requiring an increased blood supply to the

region. An improved animal model for evaluating genital blood perfusion would incorporate simultaneous

measurements of vaginal and vulvar blood perfusion. This model is necessary to understand the

complete hemodynamics of different genital structures, in particular the relationship between vaginal and

vulvar blood perfusion during arousal.

Conclusions

This is the first study to measure female rat perineal blood perfusion with LSCI in response to

nerve stimulation. We found that vulvar and anal blood perfusion increased during pudendal but not tibial

nerve stimulation. Spinalization had no impact on the perineal blood perfusion response to either

stimulation target. Further research is needed on the relationships between vaginal blood perfusion,

vulvar blood perfusion, subjective arousal, and vaginal lubrication, and the effects of peripheral nerve

stimulation.

Table 1. Summary of experiment parameters

Animal ID	Weight (kg)	First Nerve Target	Estrous Stage	Intact Spinal Cord Pudendal Nerve MT (µA)	Tibial Nerve MT (µA)	Spinalized Pudendal Nerve MT (µA)	Tibial Nerve MT (µA)
A	0.29	Tibial	Diestrus	300	400	500	1000
B	0.29	Tibial	Inconclusive	3000	3600	2100	1800
C	0.29	Tibial	Proestrus	500	300	500	300
F	0.29	Pudendal+	Metestrus	500	1300	500	3400
G	0.30	Tibial+	Metestrus	400	250	400	900
J	0.28	Pudendal	Proestrus	300	600	500	5000
K	0.26	Pudendal	Proestrus	200	400	1000	400
M	0.25	Tibial	Inconclusive	300	750	600	600
N	0.26	Pudendal	Proestrus	130	400	300	400
O	0.26	Pudendal	Inconclusive	750	1700	750	1500
P	0.26	Tibial	Estrus	140	250	170	700
Q	0.26	Pudendal*	Estrus	170	-	-	-
T	0.25	Tibial	Diestrus	260	110	400	530
U	0.26	Pudendal*	Metestrus	110	-	-	-
V	0.26	Pudendal*	Inconclusive	600	-	-	-

364 MT: motor threshold, *: Nerve Transection Experiment, +: Sampling Frequency Experiment

    Figure Captions

368   **Figure 1**. Experimental set-up. Figure 1A shows a diagram of the LSCI with the laser aimed at the vaginal

369   vestibule of a rat lying in the supine position. Figure 1B shows an example of a laser speckle image live

370   (left) and produced as a heatmap (right) with ROIs drawn around the vaginal vestibule, anus, and a

371   region in the inner thigh in blue, green, and yellow respectively.

Figure 2. Individual (thin, colored lines) and average (solid black line) blood perfusion responses for all

animals that completed the entire nerve stimulation protocol (n=10). Top left panel shows pre-

spinalization pudendal stimulation trials. Blood perfusion returned to baseline after stimulation was turned

off at 25 minutes. The top right panel shows pudendal stimulation trials after spinalization and follows

similar trends as pre-spinalization trials. The bottom panels depict tibial nerve stimulation trials for intact

(left) and spinalized (right) trials.

Figure 3. Boxplots of blood perfusion units across trial epochs (before, during, and after stimulation).

Boxplot central lines give the median, edges indicate the interquartile ranges, whiskers represent range,

and the cross denotes an outlier. Pudendal nerve stimulation trials (A, B) found elevated blood perfusion

across all three ROIs during stimulation. There were no changes during tibial stimulation trials (C, D). *

denotes significant difference (p < 0.05) between before and during stimulation epochs for an ROI.

389 **Figure 4.** Intact spinal cord pudendal nerve stimulation trial periods of an experiment with abrupt

390 increases in blood perfusion (4A) and an experiment with a higher LSCI sampling rate (4B). Top panels

391 depict VBP, ABP, and ITBP waveforms during stimulation (0.25 and 1 Hz sampling frequency for Rat J

392 (4A) and Rat F (4B) respectively). Bottom panels show zoomed-in plots of an individual signal oscillation

393 in each trial of similar width (~ 2 minutes) and amplitude (150-200 APU).

References

1. Kashdan, T. B., Goodman, F. R., Stiksma, M., Milius, C. R. & McKnight, P. E. Sexuality leads to boosts in mood and meaning in life with no evidence for the reverse direction: A daily diary investigation. *Emotion* **18**, 563–576 (2018).

2. Laumann, E. O., Paik, A. & Rosen, R. C. Sexual dysfunction in the United States: Prevalence and predictors. *J. Am. Med. Assoc.* **281**, 537–544 (1999).

3. Meston, C. M. & Stanton, A. M. Understanding sexual arousal and subjective–genital arousal desynchrony in women. *Nat. Rev. Urol.* **16**, 107–120 (2019).

4. Katz, M. *et al.* Efficacy of flibanserin in women with hypoactive sexual desire disorder: Results from the BEGONIA trial. *J. Sex. Med.* **10**, 1807–1815 (2013).

5. Clayton, A. H. *et al.* Bremelanotide for female sexual dysfunctions in premenopausal women: A randomized, placebo-controlled dose-finding trial. *Women's Heal.* **12**, 325–337 (2016).

6. Kingsberg, S. A. *et al.* Bremelanotide for the Treatment of Hypoactive Sexual Desire Disorder: Two Randomized Phase 3 Trials. *Obstet. Gynecol.* **134**, 899–908 (2019).

7. Jaspers, L. *et al.* Efficacy and safety of flibanserin for the treatment of hypoactive sexual desire disorder in women: A systematic review and meta-analysis. *JAMA Intern. Med.* **176**, 453–462 (2016).

8. Cavalcanti, A. L. *et al.* Effect of sildenafil on clitoral blood flow and sexual response in postmenopausal women with orgasmic dysfunction. *Int. J. Gynecol. Obstet.* **102**, 115–119 (2008).

9. Laan, E. *et al.* The enhancement of vaginal vasocongestion by sildenafil in healthy premenopausal women. *J. Women's Heal. Gender-Based Med.* **11**, 357–365 (2002).

10. Basson, R., McInnes, R., Smith, M. D., Hodgson, G. & Koppiker, N. Efficacy and Safety of Sildenafil Citrate in Women with Sexual Dysfunction Associated with Female Sexual Arousal Disorder. *J. Women's Heal. Gender-Based Med.* **11**, (2002).

11. Siegel, S. *et al.* Five-Year Followup Results of a Prospective, Multicenter Study of Patients with Overactive Bladder Treated with Sacral Neuromodulation. *J. Urol.* **199**, 229–236 (2018).

12. Hull, T. *et al.* Long-term durability of sacral nerve stimulation therapy for chronic fecal incontinence. *Dis. Colon Rectum* **56**, 234–245 (2013).

13.   Yih, J. M., Killinger, K. A., Boura, J. A. & Peters, K. M. Changes in Sexual Functioning in Women

after Neuromodulation. *J Sex Med* (2013) doi:10.1111/jsm.12085.

14.   Banakhar, M., Gazwani, Y., El Kelini, M., Al-Shaiji, T. & Hassouna, M. Effect of sacral

neuromodulation on female sexual function and quality of life: Are they correlated? *J. Can. Urol.*

Assoc. **8**, e762–e767 (2014).

15.   Lombardi, G., Mondaini, N., Macchiarella, A., Cilotti, A. & Popolo, G. Del. Clinical female sexual

outcome after sacral neuromodulation implant for lower urinary tract symptom (LUTS). *J. Sex.*

Med. **5**, 1411–1417 (2008).

16.   Pauls, R. N. *et al.* Effects of sacral neuromodulation on female sexual function. *Int. Urogynecol. J.*

18, 391–395 (2007).

17.   Gill, B. C. *et al.* Improved sexual and urinary function in women with sacral nerve stimulation.

Neuromodulation **14**, 436–443 (2011).

18.   Van Balken, M. R., Verguns, H. & Bemelmans, B. L. H. Sexual functioning in patients with lower

urinary tract dysfunction improves after percutaneous tibial nerve stimulation. *Int. J. Impot. Res.*

18, 470–475 (2006).

19.   Martin-Garcia, M. & Crampton, J. A single-blind, randomized controlled trial to evaluate the

effectiveness of transcutaneous tibial nerve stimulation (TTNS) in Overactive Bladder symptoms in

women responders to percutaneous tibial nerve stimulation (PTNS). *Physiother. (United Kingdom)*

105, 469–475 (2019).

20.   Zimmerman, L. L. *et al.* Transcutaneous Electrical Nerve Stimulation to Improve Female Sexual

Dysfunction Symptoms: A Pilot Study. *Neuromodulation* **21**, 707–713 (2018).

21.   Berman, J. R. Physiology of female sexual function and dysfunction. *Int. J. Impot. Res.* **17**, 44–51

(2005).

22.   Zimmerman, L. L., Rice, I. C., Berger, M. B. & Bruns, T. M. Tibial Nerve Stimulation to Drive

Genital Sexual Arousal in an Anesthetized Female Rat. *J. Sex. Med.* **15**, 296–303 (2018).

23.   Cai, R. S., Alexander, M. S. & Marson, L. Activation of Somatosensory Afferents Elicit Changes in

Vaginal Blood Flow and the Urethrogenital Reflex Via Autonomic Efferents. *J. Urol.* **180**, 1167–

1172 (2008).

24.   Rice, I. C., Zimmerman, L. L., Ross, S. E., Berger, M. B. & Bruns, T. M. Time-Frequency Analysis

of Increases in Vaginal Blood Perfusion Elicited by Long-Duration Pudendal Neuromodulation in

Anesthetized Rats. *Neuromodulation* **20**, 807–815 (2017).

25.   Chang, H. Y., Cheng, C. L., Chen, J. J. J., Peng, C. W. & De Groat, W. C. Reflexes evoked by

electrical stimulation of afferent axons in the pudendal nerve under empty and distended bladder

conditions in urethane-anesthetized rats. *J. Neurosci. Methods* **150**, 80–89 (2006).

26.   George, A. T. *et al.* Pudendal nerve stimulation for bowel dysfunction in complete cauda equina

syndrome. *Ann. Surg.* **259**, 502–507 (2014).

27.   de Groat, W. C. & Tai, C. Impact of Bioelectronic Medicine on the Neural Regulation of Pelvic

Visceral Function. *Bioelectron. Med.* **2**, 25–36 (2015).

28.   Kovacevic, M. & Yoo, P. B. Reflex neuromodulation of bladder function elicited by posterior tibial

nerve stimulation in anesthetized rats. *Am. J. Physiol. - Ren. Physiol.* **308**, F320–F329 (2015).

29.   Booth, J. *et al.* A Feasibility Study of Transcutaneous Posterior Tibial Nerve Stimulation for

Bladder and Bowel Dysfunction in Elderly Adults in Residential Care. *J. Am. Med. Dir. Assoc.* **14**,

270–274 (2013).

30.   Hoang Roberts, L., Vollstedt, A., Volin, J., McCartney, T. & Peters, K. M. Initial experience using a

novel nerve stimulator for the management of pudendal neuralgia. *Neurourol. Urodyn.* **40**, 1670–

1677 (2021).

31.   Sipski, M. L., Alexander, C. J. & Rosen, R. Sexual arousal and orgasm in women: Effects of spinal

cord injury. *Ann. Neurol.* **49**, 35–44 (2001).

32.   Collinger, J. L. *et al.* Functional priorities, assistive technology, and brain-computer interfaces after

spinal cord injury. *J. Rehabil. Res. Dev.* **50**, 145–160 (2013).

33.   Anderson, K. D. Targeting recovery: Priorities of the spinal cord-injured population. *J.*

Neurotrauma **21**, 1371–1383 (2004).

34.   Giuliano, F., Rampin, O. & Allard, J. Neurophysiology and pharmacology of female genital sexual

response. *J. Sex Marital Ther.* **28**, 101–121 (2002).

35.   Woodard, T. L. & Diamond, M. P. Physiologic measures of sexual function in women: a review.

Fertil. Steril. **92**, 19–34 (2009).

36. Waxman, S. E. & Pukall, C. F. Laser doppler imaging of genital blood flow: A direct measure of

female sexual arousal. *J. Sex. Med.* **6**, 2278–2285 (2009).

37. Boyer, S. C., Bouchard, K. N. & Pukall, C. F. Laser Doppler Imaging as a Measure of Female

Sexual Arousal: Further Validation and Methodological Considerations. *Biol. Psychol.* **148**, 107741

(2019).

38. Basson, R. The female sexual response: A different model. *J. Sex Marital Ther.* **26**, 51–65 (2000).

39. Jabs, F. & Brotto, L. A. Identifying the disruptions in the sexual response cycles of women with

Sexual Interest/Arousal Disorder. *Can. J. Hum. Sex.* **27**, 123–132 (2018).

40. Cyr, M. P., Pinard, A., Dubois, O. & Morin, M. Reliability of vulvar blood perfusion in women with

provoked vestibulodynia using laser Doppler perfusion imaging and laser speckle imaging.

Microvasc. Res. **121**, 1–6 (2019).

41. Styles, S. J., MacLean, A. B., Reid, W. M. N. & Sultana, S. R. Laser Doppler perfusion imaging: A

method for measuring female sexual response. *BJOG An Int. J. Obstet. Gynaecol.* **113**, 599–601

(2006).

42. Giuliano, F. *et al.* Experimental Models for the Study of Female and Male Sexual Function. *J. Sex.*

Med. **7**, 2970–2995 (2010).

43. Allers, K. A. *et al.* I. Slow oscillations in vaginal blood flow: Alterations during sexual arousal in

rodents and humans. *J. Sex. Med.* **7**, 1074–1087 (2010).

44. Vachon, P., Simmerman, N., Zahran, A. R. & Carrier, S. Increases in clitoral and vaginal blood

flow following clitoral and pelvic plexus nerve stimulations in the female rat. *Int. J. Impot. Res.* **12**,

53–57 (2000).

45. Damaser, M. S., Broxton-King, C., Ferguson, C., Kim, F. J. & Kerns, J. M. Functional and

neuroanatomical effects of vaginal distention and pudendal nerve crush in the female rat. *J. Urol.*

170, 1027–1031 (2003).

46. Kimura, R., Miyazato, M., Ashikari, A., Oshiro, T. & Saito, S. Age-associated urethral dysfunction

in urethane-anesthetized rats. *Neurourol. Urodyn.* **37**, 1313–1319 (2018).

47. Goldman, J. M., Murr, A. S. & Cooper, R. L. The Rodent Estrous Cycle: Characterization of

Vaginal Cytology and Its Utility in Toxicological Studies. *Birth defects Res.* **80**, 83–97 (2007).

48.   McKenna, K. E. & Nadelhaft, I. The organization of the pudendal nerve in the male and female rat.

508   *J. Comp. Neurol.* **248**, 532–549 (1986).

509   49.   Pacheco, P., Martinez-Gomez, M., Whipple, B., Beyer, C. & Komisaruk, B. R. Somato-motor

510   components of the pelvic and pudendal nerves of the female rat. *Brain Res.* **490**, 85–94 (1989).

511   50.   Xu, J. J. *et al.* Tibial nerve stimulation increases vaginal blood perfusion and bone mineral density

512   and yield load in ovariectomized rat menopause model. *Int. Urogynecol. J.* (2022)

513   doi:10.1007/s00192-022-05125-5.

514   51.   Graziottin, A. & Gambini, D. *Anatomy and physiology of genital organs - women. Handbook of*

515   *Clinical Neurology* vol. 130 (Elsevier B.V., 2015).

516   52.   Hannan, J. L. *et al.* Characterization of the vasculature supply the genital tissues in female rats. *J.*

517   *Sex. Med.* **9**, 136–147 (2012).

518   53.   Badia, J., Pascual-Font, A., Vivó, M., Udina, E. & Navarro, X. Topographical distribution of motor

519   fascicles in the sciatic-tibial nerve of the rat. *Muscle and Nerve* **42**, 192–201 (2010).

520   54.   Stampas, A. *et al.* Bladder neuromodulation in acute spinal cord injury via transcutaneous tibial

521   nerve stimulation: Cystometrogram and autonomic nervous system evidence from a randomized

522   control pilot trial. *Front. Neurosci.* **13**, 119 (2019).

523   55.   Kamboonlert, K., Panyasriwanit, S., Tantisiriwat, N. & Kitisomprayoonkul, W. Effects of Bilateral

524   Transcutaneous Tibial Nerve Stimulation on Neurogenic Detrusor Overactivity in Spinal Cord

525   Injury: A Urodynamic Study. *Arch. Phys. Med. Rehabil.* **102**, 1165–1169 (2020).

526   56.   Georgakis, E. & Soames, R. Arterial supply to the sciatic nerve in the gluteal region. *Clin. Anat.* **21**,

527   62–65 (2008).

528