Title: Mycorrhizal fungi stabilize bacterial growth and diversity in water-limited soils

Running Title: Mycorrhizae stabilize bacteria in water-limited soils

Rachel Hestrin[1]*, Megan Kan[1], Marissa Lafler[1], Jessica Wollard[1], Jeffrey A. Kimbrel[1], Prasun Ray[2], Steven Blazewicz[1], Rhona Stuart[1], Kelly Craven[3], Mary Firestone[4], Erin Nuccio[1], Jennifer Pett-Ridge[1,5]*

[1]*Lawrence Livermore National Laboratory, Physical and Life Science Directorate, Livermore, CA 94550 USA*

[2]*Department of Natural Resources, University of Maryland Eastern Shore, Princess Anne, MD 21853 USA*

[3]*Noble Research Institute, Plant Biology Division, Ardmore, OK 73401 USA*

[4]*University of California Berkeley, Department of Environmental Science Policy and Management, Berkeley, CA 94720 USA*

[5]*University of California Merced, Life & Environmental Sciences Department, Merced, CA 95343 USA*

*Correspondence to:

Rachel Hestrin, 925-409-7954, hestrin1@llnl.gov

Jennifer Pett-Ridge, 925-337-7618, pettridge2@llnl.gov

Competing Interests: The authors declare no competing interests.

Abstract

Drought disrupts soil microbial activity and many biogeochemical processes. Although mycorrhizal fungi are known to impact plant functions and nutrient cycling during drought, their effects on drought-exposed soil microbial communities are not well resolved. We used $H_2^{18}O$ quantitative stable isotope probing (qSIP) to investigate microbial response to water limitation in the hyphospheres of two distinct mycorrhizal lineages (*Rhizophagus irregularis* and *Serendipita bescii*) grown with the bioenergy model grass *Panicum hallii*. In the absence of mycorrhizal inocula, water limitation resulted in significantly lower bacterial growth rates and lower diversity in the actively growing bacterial community. Water limitation also decoupled growth from CO_2 efflux, resulting in a threefold lower growth efficiency. In contrast, both mycorrhizal lineages appeared to protect bacterial communities exposed to water limitation: bacterial growth rates, growth efficiency, and the diversity of the active bacterial community were similar in water-replete and water-limited soils inoculated with either fungus. Several of the bacterial taxa that responded positively to mycorrhizal inocula in water-limited soil belong to lineages that are considered drought-susceptible. Together, these results indicate that mycorrhizal-bacterial interactions can stabilize bacterial communities in water-limited environments. As drought frequency and severity increase, these synergistic relationships may support ecosystem resilience to moisture stress.

Keywords: *Panicum hallii*, mycorrhizal fungi, bacteria, drought, quantitative stable isotope probing, plant-microbe interactions, [18]O-water

Introduction

Drought is a destabilizing stressor that alters plant productivity[1], microbial community composition[2-4], microbial biomass[5,6], greenhouse gas emissions[7,8], and many critical biogeochemical processes[9]. Plant-microbial mutualisms mitigate plant drought response and aid in post-drought recovery through a variety of mechanisms[10,11]. In particular, mycorrhizal fungi— which associate with the majority of terrestrial root systems[12]—can support plants during drought by facilitating water transport[13], soil aggregation[14,15], root growth[16], plant nutrient uptake[17], photosynthesis[18,19], and stomatal conductance[18-20]. While mycorrhizal fungi may also influence soil microbial communities in water-limited environments, multipartite feedbacks during and after drought remain poorly quantified[11].

There is increasing evidence that the mycorrhizal hyphosphere—the region of soil that surrounds mycorrhizal fungi—is a hotspot for multipartite interactions. These interactions may be synergistic or antagonistic, with significant effects on resource exchange[21,22], stress tolerance[23], and organismal abundance and activity[24,25]. Recent work indicates that specific bacterial lineages are more abundant in the presence of arbuscular mycorrhizal (AM) fungi[26-28] and that fungal traits shape the composition of mycorrhizal-associated bacterial communities[28-30]. Together, these observations suggest that mycorrhizal-bacterial interactions have important ecological functions. Still, relatively little is known about how mycorrhizal fungi influence bacterial communities exposed to environmental stressors such as moisture limitation. A better understanding of these relationships would help us predict microbial response to global change,

harness beneficial plant-microbial relations for sustainable agriculture, and assess other

applications of mycorrhizal-microbial mutualisms.

In situ investigation of mycorrhizal-bacterial interactions presents immense methodological

challenges. The mycorrhizal hyphosphere is small, difficult to sample, and may not exert a

detectable effect on microbial community composition or activity when assessed at a 'bulk'

scale[28]. Furthermore, a substantial proportion of soil DNA may represent inactive organisms or

extracellular 'relic' DNA[31-32]. Stable isotope probing (SIP) coupled with 16S rRNA gene

profiling can distinguish active organisms from inactive ones by tracing isotope incorporation

into newly-synthesized microbial DNA[33,34]. With quantitative SIP (qSIP), we can estimate

growth rates of amplicon sequence variants (ASVs) based on shifts in DNA buoyant density

caused by heavy isotope incorporation[35,36]. This enables sensitive detection of actively growing

taxa, even in samples where a substantial quantity of DNA belongs to dead or dormant

organisms. In challenging experimental systems like the mycorrhizal hyphosphere, qSIP has the

potential to provide novel insight into taxon-specific interactions.

In this study, we sought to investigate how two mycorrhizal lineages—the AM fungus

Rhizophagus irregularis and the Sebacinales fungus *Serendipita bescii*—mediate bacterial

response to water limitation in a marginal soil planted with the switchgrass bioenergy model

Panicum hallii (Hall's panicgrass). For the purpose of this study we define 'mycorrhiza' as a

symbiotic relationship between a fungus and a plant root, where the fungus acquires C from the

living plant in exchange for mineral nutrients or other resources[12]. Both AM and Sebacinales

fungi associate with a wide range of plant species, including switchgrass[37-39], and support plant

function during drought[40-42]. However, *R. irregularis* has a reduced genome and is dependent

upon host-provided C[43,44] and microbial transformation of nutrient sources into bioavailable

forms[45,46]. In contrast, *S. bescii* and other Serendipita lineages are facultative symbionts with a

broad enzymatic repertoire that enables direct resource acquisition from both living plants and

detritus[47-49]. We hypothesized that both fungi would mitigate moisture stress, but that distinct

bacterial communities would respond to each fungus. Our results demonstrate that mycorrhizal

fungi have a stabilizing effect on bacterial communities in water-limited soil.

Materials and Methods

Soil collection and characterization

Soil used for this study was a Pond Creek fine sandy loam, classified as a superactive, thermic

Pachic Argiustoll[50]. Soil was collected from a pasture in Caddo County, OK (35.072417/-

98.303667) where switchgrass is endemic, on traditional land of the Anadarko (Nadaco) tribe

and the Caddo Nation of Oklahoma. The soil is considered 'marginal' due to its high sand

content (69%), low C, N, and P content (< 0.4%, < 0.04%, and < 6 ppm, respectively); its pH is

~5 (ref. 51). Surface soil (0-20 cm) was collected in May 2019, transported to Livermore, CA,

sieved to 2 mm, and stored at 4 °C. A soil moisture retention curve was generated from air-dried

soil using a pressure plate apparatus (WP4C, METER Environment, Pullman, WA) and the

nonlinear fitting program SWRC-Fit to apply a Brooks and Corey model to the data[52].

Mycorrhizal inoculum

Spores of *R. irregularis* (formerly *Glomus intraradices*) isolate DAOM-197198 were purchased

from Premier Tech (Rivière-du-Loup, Quebec, Canada). *S. bescii* (sourced from the Noble

Research Institute) was grown in modified Melin-Norkran's (MMN) broth. Bentonite clay

particles were mixed with MMN broth, inoculated with *S. bescii,* and incubated at 24 °C for 8

weeks according methods described by Ray et al. (ref. 53). Uninoculated bentonite clay particles

were also mixed with MMN broth and incubated under the same conditions.

Greenhouse water limitation experiment

P. hallii seeds collected from the Edwards Plateau in central Texas were grown to seed, scarified,

surface sterilized, stratified at 4 °C for one week, and germinated in petri plates. After 5 days,

germinated seedlings were transferred into planting cones (Ray Leach Cone-tainers, Steuwe &

Sons, Tangent, OR, USA) filled with double-autoclaved sand. Five days after transfer into cones,

~500 *R. irregularis* spores or 500 µL *S. bescii* inoculum were injected into the sand to inoculate

the *P. hallii* seedlings. A subset of seedlings was left uninoculated. Eight weeks after inoculation,

seedlings were transplanted into 960 cm^3 containers (Anderson Plant Bands, Steuwe & Sons)

filled with a 50:50 (v:v) mixture of 'live' soil and double-autoclaved sand (CEMEX Lapis Lustre

Specialty Sand, #2/12) packed to 1.7 g mL^{-1} bulk density at 15% moisture. Additional inoculum

(consisting of 500 *R. irregularis* spores or 5 g bentonite clay particles coated with *S. bescii*) were

placed near the roots of seedlings as they were transplanted. Five g of uncoated bentonite clay

particles were placed near the roots of previously uninoculated plants and plants that had been

inoculated with *R. irregularis*. Each microcosm contained a 25 µm mesh hyphal ingrowth core

(2.3 cm diameter) filled with 75 g of the same 'live' soil (no sand) mixed with 0.5 g finely milled

switchgrass biomass as mycorrhizal bait. The multi-phase mycorrhizal inoculation procedure

was intended to give *R. irregularis* and *S. bescii* a colonization advantage over other native

mycorrhizal lineages or endophytes present in the soil.

After assembly, each microcosm was covered with 150 g double-autoclaved sand to prevent

cross-contamination of mycorrhizal inocula. All microcosms were watered with a total of 20 mL

ultrapure H_2O during the first week to facilitate plant and mycorrhizal establishment. Half of the

microcosms were watered on a weekly or bi-weekly schedule to maintain soil moisture at

approximately 15% (based on gravimetric measurements), a level that would not restrict plant

growth. The other half of the microcosms were not watered for the remainder of the experiment

and declined to 5% gravimetric soil moisture by 3 months. The moisture content of the soil

surrounding the roots (i.e., not directly within the hyphal ingrowth cores) in 1 microcosm per

treatment was monitored continuously with a volumetric water probe (ECH_2O EC-5, METER

Group, Inc., Pullman, WA, USA). Another microcosm per treatment was weighed weekly to

assess whole-microcosm gravimetric changes. Three replicate microcosms were maintained for

each of the 6 treatment combinations (3 mycorrhizal inocula x 2 moisture regimes). Average

daytime and nighttime temperatures were 27 °C and 24 °C, respectively, with a photoperiod of

16 h. After 3 months, the microcosms were destructively harvested. Soil from hyphal ingrowth

cores was homogenized, flash frozen in liquid N_2 and stored at -80 °C for DNA extractions or at

room temperature for qSIP assays (see Fig. 1a for experimental design).

Soil characteristics

Soil moisture content following the 3-month harvest was determined by mass difference after

drying soil for 48 h at 105 °C. Total soil C and N contents were measured with a Costech ECS

4010 Elemental Analyzer (Costech Analytical Technologies Inc., Valencia, CA).

Soil $H_2^{18}O$ qSIP assay

Three days after the 3-month harvest, soil from 3 hyphal ingrowth cores per treatment was

amended with either $H_2^{16}O$ or $H_2^{18}O$ (Fig. 1a). Each soil was divided into three subsamples: one

3.0 g dry weight equivalent sample for an initial $H_2^{16}O$ qSIP timepoint ("T0") and two 4.0 g dry

weight equivalent samples for $H_2^{16}O$ and $H_2^{18}O$ 7-day qSIP timepoint ("T7"). This resulted in a

total of 54 samples (18 T0 $H_2^{16}O$ samples, 18 T7 $H_2^{16}O$ samples, and 18 T7 $H_2^{18}O$ samples).

Each subsample was air-dried to 0.047 gravimetric water content in a biosafety cabinet and then

brought up to 22.13% soil moisture (60% field capacity) with either ultrapure water at natural

isotopic abundance or water enriched with ^{18}O (98.38 atom % $H_2^{18}O$, Isoflex, San Francisco,

CA, USA). The final enrichment of the samples containing $H_2^{18}O$ was 78.76 ^{18}O atom %.

Standardization of soil moisture across treatments was necessary for qSIP calculations. The T0

subsample (amended with $H_2^{16}O$) was immediately flash frozen in liquid N_2 and stored at -80 °C.

Each of the T7 soil samples was stored in the dark inside a separate 16 oz glass jar with a tight-

fitting lid containing a septum for gas sampling. After 7 days, the soils were flash frozen in

liquid N_2 and stored at -80 °C.

CO_2 efflux measurement

15 mL headspace gas samples were collected at the beginning and end of the qSIP assay. CO_2

concentrations were measured on a gas chromatograph equipped with a thermal conductivity

detector (GC-14A, Shimadzu, Columbia, MD).

DNA extraction and qPCR

DNA was extracted from each soil sample in quadruplicate with the Qiagen PowerSoil Pro kit

and then pooled per sample prior to downstream analysis. To assess the relative abundance of

mycorrhizal inocula, we quantified *R. irregularis* and *S. bescii* gene copy number with qPCR

primers designed specifically for each fungal lineage[53,54] (see Table S1 and SI text for additional

information). All sample and standard curve qPCR reactions were performed in triplicate. We

note that DNA can be unevenly distributed throughout fungal biomass and while not necessarily

indicative of total biomass or activity is a relative indicator of inoculation success[55,56].

Density gradient fractionation of DNA

To separate isotopically enriched DNA from unenriched DNA, samples were subjected to a

cesium chloride density gradient formed in an ultracentrifuge as previously described[57,58] with

the following minor modifications. For each sample, 5 µg of DNA in 150 µL 1xTE was mixed

with 1.00 mL gradient buffer, and 4.60 mL CsCl stock (1.885 g mL^{-1}) with a final average

density of 1.730 g mL^{-1}. Samples were loaded into 5.2 mL ultracentrifuge tubes and spun at 20

ºC for 108 hours at 176,284 RCF_{avg} in a Beckman Coulter Optima XE-90 ultracentrifuge using a

VTi65.2 rotor. Sample fractionation was automated using Lawrence Livermore National

Laboratory's high-throughput SIP pipeline, which automates the fractionation and clean-up tasks

for the density gradient SIP protocol. The content of each ultracentrifuge tube was fractionated

into 22 fractions (~236 µL each) using an Agilent Technologies 1260 isocratic pump to deliver

water at 0.25 mL min^{-1} through a 25G needle inserted through the top of the ultracentrifuge tube.

Each tube was mounted in a Beckman Coulter fraction recovery system with a side port needle

inserted through the bottom. The side port needle was routed to an Agilent 1260 Infinity fraction

collector, and fractions were collected in 96-well deep well plates. The density of each fraction

was measured using a Reichart AR200 digital refractometer fitted with a prism covering to

facilitate measurement from 5 µL, as previously described[59]. DNA in each fraction was purified

and concentrated using a Hamilton Microlab Star liquid handling system programmed to

automate glycogen/PEG precipitations[34]. Washed DNA pellets were suspended in 40 µL of 1xTE

and the DNA concentration of each fraction was quantified using a PicoGreen fluorescence

assay.

DNA sequencing

For each sample, we sequenced the 16S rRNA gene in unfractionated DNA as well as DNA

fractions containing > 1.0 ng µL^{-1}, resulting in 6-11 sequenced fractions per sample. The V4

region of the 16S rRNA gene was amplified with the 515F/806R primer pair[60], processed and

barcoded through the Illumina V2 PE150 sample preparation kit, and sequenced on an MiSeq v2

platform in 3 runs (Illumina, Inc., San Diego, CA; see SI text and Tables S2 & S3 for additional

information about taxonomic assignment and ASV recovery).

Quantitative stable isotope probing (qSIP) analysis

We calculated taxon-specific growth using the ^{18}O qSIP approach[33,35]. Briefly, the qSIP

mathematical model estimates taxon-specific ^{18}O incorporation into DNA based on the shift in

DNA density following exposure to natural abundance H_2O ($H_2^{16}O$) or 'heavy' $H_2^{18}O$. We

determined baseline densities for each taxon in the absence of an isotopic tracer (i.e., $H_2^{16}O$),

because even without tracer assimilation, DNA density varies by GC content[61]. For each ASV,

we calculated median ^{18}O atom percent excess (APE) and gross growth rate. To increase

confidence in these calculations, we first filtered out ASVs that were not recovered from all T0

and T7 replicates in a particular treatment.

Since ^{18}O incorporation could have occurred at any point during the qSIP assay, ^{18}O APE

represents an integrated measure of activity. We note that a taxon's ^{18}O APE value may be high

due to an increase in abundance (i.e., cells incorporate ^{18}O into DNA during replication; new cell

production increases total population size and average DNA ^{18}O content) or population turnover

(i.e., DNA from older unenriched cells degrades, while new cells grow and incorporate ^{18}O into

DNA). During turnover, DNA ^{18}O APE can increase while population-level abundance either

remains similar (if mortality matches growth rate) or declines (if mortality outpaces growth).

Therefore, a high ^{18}O APE value is not necessarily indicative of a net increase in abundance.

Statistical analyses

We performed all statistical analyses in R[62]. We assessed differences in mycorrhizal gene copy

number with a Wilcoxon rank sum test with a Benjamini-Hochberg correction for multiple

comparisons. We assessed differences in soil moisture, soil C and N content, Inverse Simpson's

diversity index, median ASV ^{18}O APE, and CO_2 efflux with a Tukey's HSD test comparing the

means of all treatments. First, we created a linear model with a fixed effect for treatment. If the

raw data did not meet the assumptions of normality, it was log-transformed prior to further

analysis. We calculated growth efficiency by dividing gross growth by CO_2 efflux per

treatment[63]. We used 'vegan' to perform a principal coordinates analysis and permutational

multivariate analysis of variance (PERMANOVA) on the weighted UniFrac distance between

communities present under different treatments and qSIP timepoints[64,65]. We used

'metagenomeSeq'[66] with a wrench normalization for sparse data to identify significant

differences in ASV relative abundance at T0 ($p < 0.05$ and $\log_2$ fold change > 2 or < -2). To

compare taxon-specific growth between two treatments, we calculated ^{18}O-based 'growth ratios'

by dividing an ASV's ^{18}O APE in one treatment by its ^{18}O APE in the second treatment. We

used a Wilcoxon signed rank test with a Benjamini-Hochberg correction for multiple

comparisons to assess whether mean phylum ratios were significantly greater than or less than 1.

Results

Mycorrhizal abundance and soil characteristics following 3-month water manipulation

After 3 months of *P. hallii* growth, *R. irregularis* and *S. bescii* abundances (measured with

strain-specific qPCR primers) were significantly higher in the hyphal ingrowth cores of

microcosms inoculated with each fungus, respectively (Fig. 1b; $p < 0.1$). Soil moisture was lower

in the hyphal ingrowth cores of water-limited versus water-replete microcosms (4.9% $\pm$ 0.85 and

15.2% $\pm$ 0.57, respectively, Fig. 1c; $p < 0.001$), and was not significantly associated with

mycorrhizal abundance ($p > 0.05$). Matric suction was also lower in water-limited hyphal

ingrowth core soils (Fig. S1). Total soil C and N did not vary significantly between treatments

(Fig. S2; $p > 0.05$).

Moisture and mycorrhizal effects on microbial community structure

We assessed the effects of moisture regime and mycorrhizal inoculum on microbial community structure with both traditional and qSIP-enabled 16S rRNA gene profiling (Tables S4-S6 & Fig. 2a,b). The differences we detected in mycorrhizal-inoculated samples represent the influence of *R. irregularis* or *S. bescii* in soils containing a background microbial community. A PERMANOVA of weighted UniFrac distances between the 'total' communities (i.e., 16S rRNA gene profiles sequenced from unfractionated DNA) suggests that moisture regime and mycorrhizal inoculum explained 29.5% of variation in community structure at the end of the qSIP assay (Table S4 & Fig. 2a; $p < 0.05$). Although there was a small shift in community structure during the qSIP assay (6.3% of variance explained; Table S5; $p < 0.001$), moisture and mycorrhizal effects persisted throughout the assay (Table S5; $p < 0.05$).

When we used qSIP to filter out inactive ASVs (i.e., those that did not incorporate significant quantities of ^{18}O) prior to conducting a PERMANOVA, we found that moisture regime, mycorrhizal inoculum, and their interactive effect explained over 75% of the variation in the structure of the actively growing community (20.4%, 27.4%, and 27.5%, respectively; Table S6 & Fig. 2b; $p < 0.001$). This is more than twice the effect size observed in the 'total' community identified through traditional 16S rRNA analysis. qSIP-based filtering also highlighted differences in alpha diversity. While there was no relationship between moisture regime or mycorrhizal inoculum and the alpha diversity of the 'total' communities sequenced (Fig. 2c; $p > 0.05$), water limitation was associated with lower alpha diversity of the actively growing community in uninoculated soils ($p < 0.01$), no difference in *R. irregularis*-inoculated soils ($p > 0.05$), and higher alpha diversity in soils inoculated with *S. bescii* ($p < 0.01$).

We conducted differential abundance analysis to evaluate how water limitation influenced ASV

relative abundance in the presence or absence of mycorrhizal inocula. Within each mycorrhizal

treatment, some ASVs were more abundant in water-limited soils while others were less

abundant (Fig. S3). In uninoculated soils, 56 ASVs were less abundant following water

limitation while only 24 were more abundant, contributing to a twofold difference in the

magnitude of cumulative $\log_2$ fold change (Fig. S3a,d). In contrast, within *R. irregularis*- and *S.*

bescii-inoculated soils, the number of ASVs that were more abundant following water limitation

was similar to the number that were less abundant, contributing to a similar magnitude of

cumulative $\log_2$ fold change (Fig. S3b,c,d).

Moisture and mycorrhizal effects on microbial growth and efficiency

We used $H_2^{18}O$ qSIP and CO_2 efflux measurements to quantify the effects of moisture regime

and mycorrhizal inoculum on ASV ^{18}O incorporation, gross growth rates, and growth efficiency.

Overall, DNA from soil amended with $H_2^{18}O$ was denser than DNA from soil amended with

$H_2^{16}O$ (Fig. S4), indicating that many organisms incorporated ^{18}O into newly-synthesized DNA.

Taxon-specific ^{18}O APE ranged from unenriched to 64.0 APE, with a median of 6.0. Between

35-59% of the total ASVs detected per treatment were significantly enriched (Table S3; lower

90% CI > 0); we refer to these as the 'actively growing' taxa.

In the absence of mycorrhizal inocula, microbial ASVs incorporated less ^{18}O in water-limited

soils: median ASV ^{18}O APE was 2.14 compared to 9.06 in water-replete soils (Fig. 3a; $p < 0.05$).

Gross growth was also lower in water-limited compared to water-replete soil (Fig. 3b). CO_2

efflux from uninoculated soils did not vary significantly between moisture regimes (Fig. 3c; p >

0.05). However, growth efficiency (gross growth divided by CO_2 efflux) was more than threefold

lower in water-limited soil: only 8.4 (+/- 0.24) µg DNA mg^{-1} CO_2 compared to 28.3 (+/- 2.16) in

water-replete soils (Table S7). In contrast, microbial communities in mycorrhizal-inoculated

soils were less affected by water limitation. Microbial ^{18}O APE was 6.56 compared to 7.45 in

water-limited versus water-replete *R. irregularis*-inoculated soils (Fig. 3a; p > 0.05), and 5.67

compared to 4.85 in water-limited versus water-replete *S. bescii*-inoculated soils (Fig. 3a; p >

0.05). Gross growth was also similar under both moisture regimes in mycorrhizal-inoculated

soils (Fig. 3b). CO_2 efflux from *S. bescii*-inoculated soils was higher than from *R. irregularis*-

inoculated soils, but the difference was only significant between water-replete soils (Fig. 3c; p <

0.05). Compared to uninoculated soils, moisture effects on growth efficiency in mycorrhizal-

inoculated soils were relatively small. Growth efficiency was 14.4 (+/- 0.8) µg DNA mg^{-1} CO_2

compared to 20.2 (+/- 1.25) in water-limited versus water-replete soil inoculated with *R.*

irregularis, and 12.5 (+/- 0.51) compared with 12.7 (+/- 0.75) in water-limited versus water-

replete soils inoculated with *S. bescii* (Table S7).

Taxon-specific response to moisture regime and mycorrhizal inocula

Since compensatory dynamics may result in the increased activity of some organisms while

others decline[67], we calculated ^{18}O-based 'growth ratios' for each actively growing ASV

detected in both water-limited and water-replete soil. A ratio of greater than 1.0 suggests that the

ASV sustained a higher growth rate in water-limited soil, a ratio of less than 1.0 suggests that the

ASV sustained a lower growth rate in water-limited soil, and a ratio indistinguishable from 1.0

suggests that the ASV maintained a similar growth rate under both conditions. In the absence of

mycorrhizal inocula, most ASVs sustained lower growth rates in water-limited soil (Fig. 4a).

When averaged at the phylum level, growth ratios were significantly less than 1 for 8 of the 10

most abundant phyla: Acidobacteria, Actinobacteria, Bacteroidetes, candidate division WPS-1,

Gemmatimonadetes, Planctomycetes, Proteobacteria, and Verrucomicrobia (Fig. 4a; adjusted p <

0.05). In contrast, bacterial response to water limitation in mycorrhizal-inoculated soils was

equivocal: many ASVs sustained lower growth rates, while many others sustained similar or

greater growth rates. When averaged at the phylum level, this resulted in almost no significant

effects of water limitation on microbial growth in mycorrhizal-inoculated soil (Fig 4b,c; adjusted

p > 0.05 for most phyla). An exception to this pattern were the Acidobacteria, which sustained

significantly higher growth rates in water-limited versus water-replete soils inoculated with *R.*

irregularis (Fig. 4b; adjusted p < 0.05).

We also calculated ^{18}O-based growth ratios for ASVs detected in both uninoculated and

mycorrhizal-inoculated water-limited soils. This allowed us to compare how mycorrhizal inocula

influenced the growth of individual ASVs in water-limited soil. Most ASVs sustained higher

growth rates in the presence of *R. irregularis* or *S. bescii* than in their absence (Fig. 5). Averaged

at the phylum level, populations belonging to the phyla Acidobacteria, Actinobacteria,

Bacteroidetes, candidate division WPS-1, Planctomycetes, Proteobacteria, and Verrucomicrobia

were significantly more active in *R. irregularis*-inoculated soil (Fig. 5a,c; adjusted p < 0.05). In

S. bescii-inoculated soils, populations from the phyla Actinobacteria and Proteobacteria were

significantly more active than in uninoculated soil (Fig. 5b,d; adjusted p < 0.05). Some ASVs

responded positively to both *R. irregularis* and *S. bescii*; others responded positively only to one

fungal lineage (Fig. 5c,d). Overall, the magnitude of positive growth response was greater in *R.*

irregularis- than in *S. bescii*-inoculated soils.

Positive bacterial response to mycorrhizal inocula was also apparent in ASVs that did not grow

under multiple conditions and therefore could not be compared via the ratios described above.

For example, there were no Firmicutes ASVs with detectable growth rates in uninoculated water-

limited soils (Fig. S5). However, several Firmicutes were active in mycorrhizal-inoculated soils

regardless of moisture regime. More than double the number of Firmicutes ASVs were active in

R. irregularis- compared to *S. bescii*-inoculated soils.

Discussion

Mycorrhizal fungi are well-known to improve plant drought tolerance[16-20,40-42], but their effects

on bacterial communities in droughted soils are poorly quantified. With $H_2^{18}O$ qSIP, we show

that mycorrhizal fungi protect bacterial communities from the suppressive and destabilizing

effects of water limitation.

Water limitation suppresses bacterial growth and efficiency

Our findings suggest that water limitation suppresses taxon-specific bacterial growth rates and

decouples growth from CO_2 efflux. Previous studies have shown that microbial carbon use

efficiency is lower in water-limited soils[68,69]. This is consistent with a microbial stress response,

wherein cells prioritize essential metabolic activities over replication[9,57]. Comparative

transcriptomics analysis of drought-selected microbial communities has shown that there are

trade-offs between stress tolerance and growth[70]. Slowed growth helps bacteria persist in the

presence of antibiotics[71,72], which can accumulate in dry soils as microorganisms compete for

limited resources[73]. There is also evidence that lower microbial growth efficiency in dry soils

may be due to reduced substrate diffusion[63,74]. Our observations indicate that water limitation

reduces microbial resource use efficiency and capacity to produce new biomass. Because

microbial biomass is an important precursor to soil organic matter[75], this may impact C storage

in droughted soils.

Growth suppression was apparent across a broad range of bacterial lineages present following

water limitation, including taxa that are considered to be drought tolerant. Several previous

studies have reported increases in monoderm (e.g., Actinobacteria, most Firmicutes, and some

Chloroflexi) compared to diderm (e.g., most Acidobacteria, Bacteroidetes, Proteobacteria, and

Verrucomicrobia) abundance and activity in dry soils[2,9,76-78]. Monoderm physiology (i.e., thick

cell walls and lack of an outer membrane) protects against oxidative damage under dry

conditions[9,76,77,79]. Monoderms can also outcompete other organisms through their capacity to

produce antibiotics[73,77,80] or utilize complex C substrates that remain available following water

limitation[4,73]. However, we did not find evidence that monoderm growth rates were higher than

those of diderms in water-limited soil. Taxa belonging to the monoderm phylum Actinobacteria

may remain metabolically active but replicate more slowly than other organisms[81]. This lifestyle

could explain the decoupling of growth and CO_2 efflux that we observed in water-limited soils.

Mycorrhizal fungi stabilize bacterial response to soil moisture

The AM fungus *R. irregularis* and the Sebacinales fungus *S. bescii* can substantially enhance

plant performance in water-limited soils[40-42], thereby stabilizing plant communities during stress.

We found that both fungi also appeared to protect bacterial communities exposed to water

limitation. Overall, bacterial growth and efficiency were similar in water-replete and water-

limited soils inoculated with either fungus, and always higher in mycorrhizal-inoculated soils

than in uninoculated soils. This 'protective' mycorrhizal effect extended broadly to the bacterial

community, including many taxa that are considered drought-susceptible. While previous studies

report a decrease in the relative abundances of Bacteroidetes, Planctomycetes, Verrucomicrobia,

and many Proteobacteria and Acidobacteria in droughted soils[2-4,82], we found that in the presence

of mycorrhizal inocula, many ASVs belonging to these phyla sustained similar growth rates in

water-replete and water-limited soils. Mycorrhizal fungi could help bacteria tolerate water

limitation through enhanced resource provisioning[21,83-85], soil structure[14,15], or biofilm

formation[86]. By supporting microbial diversity and activity in water-limited soil, mycorrhizal-

bacterial synergies may counteract the destabilizing effect of environmental stress[87] and improve

capacity for recovery.

In contrast to their synergistic effect in water-limited soils, *R. irregularis* and *S. bescii* appeared

to suppress bacterial growth and diversity in water-replete soils. Previous reports show that

mycorrhizal fungi can compete with other biota for N^{88} and $P^{24,25}$, that putative bacterial

predators are more abundant on extraradical mycorrhizal hyphae than in surrounding soil[28], and

that soil filtrates containing bacteria inhibit mycorrhizal proliferation[89]. Our results indicate that

there is a trade-off between mycorrhizal and bacterial activity. While this trade-off may reduce

maximum bacterial productivity under non-limiting conditions, it supports resilience under

limiting conditions. Similar context-dependency is well-documented in other mutualisms[90]. We

interpret the context-dependency of mycorrhizal-bacterial relationships as further indication of

their stabilizing ecological effect.

Magnitude of microbial response is mycorrhizal lineage-dependent

While both mycorrhizal lineages stabilized bacterial response to soil moisture, *R. irregularis*

elicited a stronger positive response than *S. bescii*—both with respect to the number of ASVs and

the magnitude of their response. Previous studies show that distinct microbial consortia associate

with different mycorrhizal lineages[28,29]. Although empirical evidence remains sparse, different

mycorrhizal exudate profiles, growth habits, and other functional traits may shape surrounding

microbial community composition and activity[45,91], a phenomenon documented more extensively

for root-microbial interactions[92-94]. Reduced bacterial growth rate and efficiency in *S. bescii*-

compared to *R. irregularis*-inoculated soils may be due to *S. bescii*'s wider enzymatic repertoire,

which could accelerate decomposition (as indicated by higher CO_2 efflux) or contribute to

greater competition with other soil biota. It is also possible that greater abundance of *R.*

irregularis compared to *S. bescii* contributed to the different magnitude of bacterial response.

Together, these findings indicate that different mycorrhizal types elicit similar, but distinct

bacterial response to water limitation, with significant effects on microbial community

composition and growth rates.

H$_2$^{18}O qSIP highlights microbial response

Previous studies report conflicting effects of drought on microbial biomass[5,6,95.96], activity[5,6,97,98],

and community composition[2-4,82]. These variable responses may be due to differences in drought

severity, biota examined, and experimental methods employed[10,99]. Previous drought exposure

can attenuate microbial response to additional water limitation[2]. With qSIP, we detected a strong

negative microbial response to water limitation in soils that had repeatedly experienced below-

average rainfall during the preceding decade[100]. This response was less apparent in our

traditional microbial community analyses. For example, alpha diversity seemed unaffected by

water limitation when assessed across 'total community' 16S rRNA gene profiles, but was

significantly reduced in the actively growing community identified by qSIP. Similarly, compared

to differential abundance analysis of 16S amplicons, qSIP identified tenfold more ASVs that

were suppressed by water limitation (56 and 579, respectively). We attribute these differences to

qSIP's capacity to distinguish actively growing organisms from the rest of the microbial

community.

Conclusion

As global precipitation patterns change and water limitation becomes more prevalent, it is

important to understand biotic response and conditions that might mitigate associated declines in

function. Mycorrhizal fungi support plant performance and alter biogeochemistry in droughted

soils, but their effect o`n soil bacterial response to water limitation is poorly explored. With

$H_2^{18}O$ qSIP, we show that mycorrhizal fungi protect bacterial communities from the suppressive

and destabilizing effects of water limitation. Both the AM fungus *R. irregularis* and the

Sebacinales fungus *S. bescii* facilitated greater bacterial growth rates, growth efficiency, and the

diversity of the active bacterial community in water-limited soil. While these divergent

mycorrhizal lineages stimulated responses of differing magnitude, broad microbial patterns were

similar, suggesting that the dominant underlying mechanisms are not limited to interactions

between a small number of taxa. Remarkably, both *R. irregularis* and *S. bescii* prompted a

positive bacterial response despite the presence of a background microbial community in

uninoculated soil. This finding is relevant for practical evaluation of mycorrhizal inoculants,

whose ability to persist and elicit a positive effect in natural settings is not well-established.

Together, our findings demonstrate that context-dependent mycorrhizal relationships support

biotic resilience to moisture stress and may promote recovery once moisture is restored.

Acknowledgements

We thank Christine Hawkes for *P. hallii* seeds; Eric Slessarev for soil collection; Megan Foley,

Anne Kakouridis, Benjamin Koch, and Alexa Nicolas for scientific discussions; Christina

Ramon for procurement; Katerina Estera-Molina, Don Herman, Ilexis Chu Jacoby, Aaron Chew,

Christina Fossum, Mengting Yuan, Tasnim Ahmed, Cynthia-Jeanette Mancilla, Melanie

Rodriguez-Fuentes, David Sanchez, Laura Adame, Emily Kline, Madeline Moore, Jack Hagen,

Anne Kakouridis, Ella Sieradzki, Nameer Baker, Sarah Baker, Alex Greenlon, Heejung Cho,

Alexa Nicolas, Peter Weber, Vanessa Brisson, Eric Slessarev, Gareth Trubl, Dinesh Adhikari,

Craig See, Maria Guerra, Rachel Neurath, and Evan Starr for their assistance with experimental

set up, maintenance, and harvests; and Tibisay Perez and Whendee Silver for assistance with GC

access and operation. This work was supported by the Genomic Science Program of the US

Department of Energy Office of Biological and Environmental Research as part of the LLNL

Biofuels SFA (SCW1039) and a sustainable bioenergy award (PI Firestone) to UC Berkeley

(DE-SC0014079), Noble Research Institute, University of Oklahoma, Lawrence Berkeley

National Laboratory, and Lawrence Livermore National Laboratory (SCW1555). Research at

LLNL was performed under the auspices of the U.S. Department of Energy at Lawrence

Livermore National Laboratory under Contract DE-AC52-07NA27344.

Competing Interests

499 The authors declare no competing interests.

References

1. Leng G, Hall J. Crop yield sensitivity of global major agricultural countries to droughts and

the projected changes in the future. Sci Total Environ 2019; 654: 811–821.

2. Bouskill NJ, Lim HC, Borglin S, Salve R, Wood TE, Silver WL, et al. Pre-exposure to

drought increases the resistance of tropical forest soil bacterial communities to extended

drought. ISME J 2013; 7: 384–394.

3. Acosta-Martinez V, Cotton J, Gardner T, Moore-Kucera J, Zak J, Wester D, et al.

Predominant bacterial and fungal assemblages in agricultural soils during a record

drought/heat wave and linkages to enzyme activities of biogeochemical cycling. Appl Soil

Ecol 2014; 84: 69–82.

4. Yuste JC, Fernandez-Gonzalez AJ, Fernandez-Lopez M, Ogaya R, Penuelas J, Sardans J, et

al. Strong functional stability of soil microbial communities under semiarid Mediterranean

conditions and subjected to long-term shifts in baseline precipitation. Soil Biol Biochem

2014; 69; 223–233.

5. Hueso S, García C, Hernández T. Severe drought conditions modify the microbial

community structure, size and activity in amended and unamended soils. Soil Biol Biochem

2012: 50: 167–173.

6. Alster CJ, German DP, Lu Y, Allison, SD. Microbial enzymatic responses to drought and to

nitrogen addition in a southern California grassland. Soil Biol Biochem 2013; 64: 68–79.

7. Intergovernmental Panel on Climate Change (IPCC), Global Warming of 1.5°C, V.

Masson☐Delmotte et al., Eds. (Cambridge Univ. Press, 2018).

8. O'Connell CS, Ruan L, Silver WL. Drought drives rapid shifts in tropical rainforest soil

biogeochemistry and greenhouse gas emissions. Nat Commun 2018; 9: 1348.

9. Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its

implications for ecosystem function. Ecology 2007; 88: 1386–1394.

10. Naylor D, Colemann-Derr D. Drought stress and root-associated bacterial communities.

Front Plant Sci 2018; 8: 2223.

11. de Vries FT, Griffiths RI, Knight CG, Nicolitch O, Williams A. Harnessing rhizosphere

microbiomes for drought-resilient crop production. Science 2020; 368: 270-274.

12. Smith SE, Read D. Mycorrhizal symbiosis. third ed. London: Academic Press; 2008. p. 145–

190.

13. Kakouridis A, Hagen JA, Kan MP, Mambelli S, Feldman LJ, Herman DJ, et al. Routes to

roots: direct evidence of water transport by arbuscular mycorrhizal fungi to host plants and

identification of a hyphal transport pathway. bioRxiv

https://doi.org/10.1101/2020.09.21.305409.

14. Rillig MC Mummey DL. Mycorrhizas and soil structure. New Phytol 2006; 171: 41-53.

15. Rillig MC, Lutgen ER, Ramsey PW, Klironomos JN, Gannon JE. Microbiota accompanying

different arbuscular mycorrhizal fungal isolates influence soil aggregation. Pedobiologia

2005; 49: 251–259.

16. Gong M, You X, Zhang Q. Effects of *Glomus intraradices* on the growth and reactive

oxygen metabolism of foxtail millet under drought. Ann Microbiol 2015; 65: 595–602.

17. Ruiz-Lozano JM. Arbuscular mycorrhizal symbiosis and alleviation of osmotic stress. New

perspectives for molecular studies. Mycorrhiza 2003; 13: 309-317.

18. Morte A, Lovisolo C, Schubert A. Effect of drought stress on growth and water relations of

the mycorrhizal association *Helianthemum almeriense–Terfezia claveryi*. Mycorrhiza 2000;

10: 115–119.

19. Birhane E, Sterck F, Fetene M, Bongers F, Kuyper T. Arbuscular mycorrhizal fungi enhance

photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed

water availability conditions. Oecologia 2012; 169: 895–904.

20. Duan X, Neuman DS, Reiber JM, Green CD, Saxton AM, Augé RM. Mycorrhizal influence

on hydraulic and hormonal factors implicated in the control of stomatal conductance during

drought. J Exp Bot 1996; 47: 1541–1550.

21. Kaiser C, Kilburn MR, Clode PL, Fuchslueger L, Koranda M, Cliff JP, et al. Exploring the

transfer of recent plant photosynthates to soil microbes: mycorrhizal pathway vs direct root

exudation. New Phytol 2015; 205: 1537-1551.

22. Hodge A, Campbell CD, Fitter AH. An arbuscular mycorrhizal fungus accelerates

decomposition and acquires nitrogen directly from organic matter. Nature 2001; 413: 297-

299.

23. Medina A, Probanza A, Gutierrez Mañero FJ, Azcón R. Interactions of arbuscular-

mycorrhizal fungi and *Bacillus* strains and their effects on plant growth, microbial

rhizosphere activity (thymidine and leucine incorporation) and fungal biomass (ergosterol

and chitin). Appl Soil Ecol. 2003; 22: 15–28.

24. Svenningsen NB, Watts-Williams SJ, Joner EJ. et al. Suppression of the activity of

arbuscular mycorrhizal fungi by the soil microbiota. ISME J 2018; 12: 1296–1307.

25. Cruz-Paredes C, Svenningsen NB, Nybroe O, Kjøller R, Frøslev TG, Jakobsen I. Suppression

of arbuscular mycorrhizal fungal activity in a diverse collection of non-cultivated soils.

FEMS Microbiol Ecol 2019; 95: fiz020.

26. Nuccio EE, Hodge A, Pett-Ridge J, Herman DJ, Weber PK, Firestone MK. An arbuscular

mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling

during litter decomposition. Environ Microbiol 2013; 15: 1870–81.

27. Zhang L, Shi N, Fan J, Wang F, George TS, Feng G. Arbuscular mycorrhizal fungi stimulate

organic phosphate mobilization associated with changing bacterial community structure

under field conditions. Environ Microbiol 2018a; 20: 2639–51.

28. Emmet BD, Levesque-Tremblay V, Harrison MJ. Conserved and reproducible bacterial

communities associate with extraradical hyphae of arbuscular mycorrhizal fungi. ISME J

2021; 15: 2276-2288.

29. Toljander JF, Artursson V, Paul LR, Jansson JK, Finlay RD. Attachment of different soil

bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal

vitality and fungal species. FEMS Microbiol Lett 2006; 254: 34–40.

30. Yuan MM, Kakouridis A, Starr E, Nguyen N, Shi S, Pett-Ridge J, et al. Fungal-bacterial co-

occurrence patterns differ between AMF and non-mycorrhizal fungi across space and time.

mBio 2021; 12: e03509-20.

31. Carini P, Marsden P, Leff J, Morgan E, Strickland M, Fierer N. Relic DNA is abundant in

soil and obscures estimates of soil microbial diversity. Nat Microbiol 2017; 2: 16242.

32. Lennon JT, Muscarella ME, Placella MA, Lehmkuhl BK. How, When, and Where Relic

DNA Affects Microbial Diversity. MBio 2018; 9: e00637-18.

33. Hungate BA, Mau RL, Schwartz E, Caporaso JG, Dijkstra P, van Gestel N, et al. Quantitative

microbial ecology through stable isotope probing. Appled Environ Microbiol 2015; 81:

7570–7581.

34. Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW, Murrell JC.

DNA stable-isotope probing. Nat Protoc 2007; 2: 860-866.

35. Koch BJ, McHugh TA, Hayer M, Schwartz E, Blazewicz SJ, Dijkstra P, et al. Estimating

taxon-specific population dynamics in diverse microbial communities. Ecosphere 2018; 9:

e02090–15.

36. Blazewicz SJ, Hungate BA, Koch BJ, Nuccio EE, Morrissey E, Brodie EL, Schwartz E, Pett-

Ridge J, Firestone MK. Taxon-specific microbial growth and mortality patterns reveal

distinct temporal population responses to rewetting in a California grassland soil. ISME J

2020; 14: 1520-1532.

37. Kilronomos JN. Host-specificity and functional diversity among arbuscular mycorrhizal

fungi. In: Bell, C.R., Brylinski, M. and Johnson-Green, P., Eds., Proceedings of the 8th

International Symposium on Microbial Ecology, Atlantic Canada Society from Microbial

Ecology, Halifax, 2000; 845-851.

38. Ray P, Guo Y, Chi MH, Krom N, Saha MC, Craven KD. *Serendipita bescii* promotes winter

wheat growth and modulates the host root transcriptome under phosphorus and nitrogen

starvation. Environ Microbiol 2021; 23: 1876–1888.

39. Lee MR, Hawkes CV. Widespread co-occurrence of Sebacinales and arbuscular mycorrhizal

fungi in switchgrass roots and soils has limited dependence on soil carbon or nutrients.

Plants, People, Planet 2021; 3: 614-626.

40. Ruiz-Lozano JM, Azcon R, Gomez M. Effects of arbuscular-mycorrhizal glomus species on

drought tolerance: physiological and nutritional plant responses. Appl Environ Microbiol

1995; 61: 456-460.

41. He F, Sheng M, Tang M. Effects of *Rhizophagus irregularis* on photosynthesis and

antioxidative enzymatic system in *Robinia pseudoacacia* L. under drought stress. Front Plant

Sci 2017; 8: 183.

42. Ghimire SR, Craven KD. Enhancement of switchgrass (*Panicum virgatum* L.) biomass

production under drought conditions by the ectomycorrhizal fungus *Sebacina vermifera*.

Appl Environ Microbiol 2011; 77: 7063-7067.

43. Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, et al. Genome of an

arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. Proc Natl

Acad Sci USA 2013; 110: 20117-20122.

44. Kamel L, Keller-Pearson M, Roux C, Ané JM. Biology and evolution of arbuscular

mycorrhizal symbiosis in the light of genomics. New Phytol 2017; 213: 531-536.

45. Zhang L, Feng G, Declerck S. Signal beyond nutrient, fructose, exuded by an arbuscular

mycorrhizal fungus triggers phytate mineralization by a phosphate solubilizing bacterium.

ISME J 2018; 12: 2339.

46. Hestrin R, Hammer EC, Mueller CW, Lehmann J. Synergies between mycorrhizal fungi and

soil microbial communities increase plant nitrogen acquisition. Commun Biol 2019; 2: 233.

47. Zuccaro A, Lahrmann U, Güldener U, Langen G, Pfiffi S, Biedenkopf D, et al. Endophytic

life strategies decoded by genome and transcriptome analyses of the mutualistic root

symbiont *Piriformospora indica*. PLoS Pathog. 2011; 7: e1002290.

48. Ray P, Chi MH, Guo Y, Chen C, Adam C, Kuo A, et al. **Genome sequence of the plant**

growth promoting fungus *Serendipita vermifera* subsp. *bescii*: The first native strain

from North America. Phytobiomes J 2018; 2: 62-63.

49. Dias T, Pimentel V, Cogo AJD, Costa R, Bertolazi AA, Miranda C, et al. The free-living

stage growth conditions of the endophytic fungus *Serendipita indica* may regulate its

potential as plant growth promoting microbe. Front Microbiol 2020; 11: 562238.

50. Moffatt HH. Soil Survey of Caddo County, Oklahoma. United States Department of 836

Agriculture Soil Conservation Service, Washington, D.C. 1973.

51. Sher Y, Baker NR, Herman NR, Fossum C, Hale L, Zhang XX, et al. Microbial extracellular

polysaccharide production and aggregate stability controlled by Switchgrass (*Panicum*

virgatum) root biomass and soil water potential. Soil Biol Biochem 2020; 143: 107742.

52. Seki K. SWRC fit - a nonlinear fitting program with a water retention curve for soils having

unimodal and bimodal pore structure. Hydrol Earth Syst Sci Discuss 2007; 4: 407-437.

53. Ray P, Ishiga T, Decker SR, Turner GB, Craven KD. A novel delivery system for the root

symbiotic fungus, *Sebacina vermifera*, and consequent biomass enhancement of low lignin

COMT switchgrass lines. BioEnerg Res 2015; 8: 922–933.

54. Badri DV, Chaparro JM, Zhang R, Shen Q, Vivanco JM. Application of natural blends of

phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that

phenolic-related compounds predominantly modulate the soil microbiome. J Biol Chem

2013; 288: 4502–4512.

55. Gamper HA, Young JP, Jones DL, Hodge A. Real-time PCR and microscopy: are the two

methods measuring the same unit of arbuscular mycorrhizal fungal abundance? Fungal Genet

Biol 2008; 45: 581-96.

56. Tellenbach C, Grünig CR, Sieber TN. Suitability of quantitative real-time PCR to estimate

the biomass of fungal root endophytes. Appl Environ Microbiol 2010; 76: 5764-5772.

57. Blazewicz SJ, Schwartz E, Firestone MK. Growth and death of bacteria and fungi underlie

rainfall-induced carbon dioxide pulses from seasonally dried soil. Ecology 2014; 95: 1162-

1172.

58. Sieradzki ET, Koch BJ, Greenlon A, Sachdeva R, Malmstrom RR, Mau RL, et al.

Measurement error and resolution in quantitative stable isotope probing: implications for

experimental design. mSystems 2020; 5: e00151-20.

59. Buckley DH, Huangyutitham V, Hsu SF, Nelson TA. Stable isotope probing with ^{15}N

achieved by disentangling the effects of genome G+C content and isotope enrichment on

DNA density. Appl Environ Microbiol 2007; 73: 3189-3195.

60. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.

Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc

Natl Acad Sci USA 2011; 108: 4516-4522.

61. Schildkraut CL, Marmur J, Doty P. Determination of the base composition of

deoxyribonucleic acid from its buoyant density in CsCl. J Mol Biol 1962; 4: 430–43.

62. R Core Team. R: a language and environment for statistical computing. Vienna: R

Foundation for Statistical Computing; 2019.

63. Manzoni S, Taylor P, Richter A, Porporato A, Ågren GI. Environmental and stoichiometric

controls on microbial carbon-use efficiency in soils. New Phytol 2012; 196: 79-91.

64. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, OHara RB, et al. vegan:

Community Ecology Package R package version 2.3-0. 2015. Available from

http://CRAN.R-project.org/package=vegan.

65. Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: an effective distance

metric for microbial community comparison. ISME J 2011; 5: 169–172.

66. Paulson JN, Stine OC, Bravo HC, Pop M. Differential abundance analysis for microbial

marker-gene surveys. Nat Meth 2013; 10: 1200-1202.

67. Wagg C, Dudenhöffer JH, Widmer F, van der Heijden MGA. Linking diversity, synchrony

and stability in soil microbial communities. Funct Ecol 2018; 32: 1280–1292.

68. Tiemann LK, Billings SA. Changes in variability of soil moisture alter microbial community

C and N resource use. Soil Biol Biochem 2011; 43: 1837–1847.

69. Domeignoz-Horta LA, Pold G, Liu XJA, Frey SD, Melillo JM, DeAngelis KM. Microbial

diversity drives carbon use efficiency in a model soil. Nat Commun 2020; 11: 3684.

70. Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD. Defining trait-

based microbial strategies with consequences for soil carbon cycling under climate change.

ISME J 2020; 14: 1–9.

71. Gefen O, Balaban NQ. The importance of being persistent: heterogeneity of bacterial

populations under antibiotic stress. FEMS Microbiol Rev 2009; 33: 704–717.

72. Fridman O, Goldberg O, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time

underlies antibiotic tolerance in evolved bacterial populations. Nature 2014; 513: 418–421.

73. Bouskill NJ, Wood TE, Baran R, Hao Z, Ye Z, Bowen BP, et al. Belowground response to

drought in a tropical forest soil. II. Change in microbial function impacts carbon

composition. Front Microbiol 2016; 7: 323.

74. Stark JM, Firestone MK. Mechanisms for soil moisture effects on activity of nitrifying bacteria. Appl Environ Microbiol 1995; 61: 218–221.

75. Liang C, Amelung W, Lehmann J, Kästner M. Quantitative assessment of microbial necromass contribution to soil organic matter. Glob Change Biol 2019; 25: 3578–3590.

76. Sutcliffe IC. A phylum level perspective on bacterial cell envelope architecture. Trends Microbiol 2010; 18: 464-470.

77. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, et al. Drought delays development of the sorghum root microbiome and enriches for monoderm bacteria. Proc Natl Acad Sci USA 2018; 115: E4284-E4293.

78. Santos-Medellín C, Liechty Z, Edwards J, Nguyen B, Huang B, Weimer BC, Sundaresan V. Prolonged drought imparts lasting compositional changes to the rice root microbiome. Nat Plants 2021; 7, 1065–1077.

79. Tocheva EI, Ortega DR, Jensen GJ. Sporulation, bacterial cell envelopes and the origin of life. Nat Rev Microbiol 2016; 14: 535–542.

80. Otoguro M, Yamamura H, Quintana ET. (2014) The Family Streptosporangiaceae. In:

Rosenberg E., DeLong E.F., Lory S., Stackebrandt E., Thompson F. (eds) The Prokaryotes.

Springer, Berlin, Heidelberg.

81. Nuccio EE, Starr E, Karaoz U, Brodie EL, Zhou J, Tringe SG, et al. Niche differentiation is

spatially and temporally regulated in the rhizosphere. ISME J 2020; 14: 999–1014.

82. Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal communities

to extreme desiccation and rewetting. ISME J 2013; 7: 2229–2241.

83. Guhr A, Borken W, Spohn M, Matzner E. Soil water redistribution by a saprotrophic fungus.

PNAS 2015; 112: 12647-14651.

84. Guhr A, Marzini C, Borken W, Poll C, Matzner E. Effect of water redistribution by two

distinct saprotrophic fungi on carbon mineralization and nitrogen translocation in dry soil.

Soil Biol Biochem 2016; 103: 380-387.

85. Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ, et al. Shifting carbon

flow from roots into associated microbial communities in response to elevated atmospheric

CO_2. Proc Natl Acad Sci USA 2010; 107: 10938–10942.

86. Cruz AF, Ishii T. Arbuscular mycorrhizal fungal spores host bacteria that affect nutrient

biodynamics and biocontrol of soil-borne plant pathogens. Biol Open 2012; 1: 52-57.

87. Hernandez, D.J., David, A.S., Menges, E.S. *et al.* Environmental stress destabilizes microbial

networks. ISME J 2021; 15: 1722–1734.

88. Bukovská P, Bonkowski M, Konvalinková T, Beskid O, Hujslová M, Püschel D et al.

Utilization of organic nitrogen by arbuscular mycorrhizal fungi-is there a specific role for

protists and ammonia oxidizers? Mycorrhiza 2018; 28: 269-283.

89. Leigh J, Fitter AH, Hodge A. Growth and symbiotic effectiveness of an arbuscular

mycorrhizal fungus in organic matter in competition with soil bacteria. FEMS Microbiol

Ecol 2011; 76: 428–438.

90. Bronstein JL. Conditional outcomes in mutualistic interactions. Trends Ecol Evol 1994; 9:

214–217.

91. Toljander JF, Lindahl BD, Paul LR, Elfstrand M, Finlay RD. Influence of arbuscular

mycorrhizal mycelial exudates on soil bacterial growth and community structure. FEMS

Microbiol Ecol 2007; 61: 295–304.

92. Zhalnina K, Louie KB, Hao Z, Mansoori N, Nunes da Rocha U, Shi S, et al. Dynamic root

exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial

community assembly. Nat Microbiol 2018; 3: 470–480.

93. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. Root

exudation of phytochemicals in *Arabidopsis* follows specific patterns that are

developmentally programmed and correlate with soil microbial functions. PLoS One 2013; 8:

e55731.

94. Shi S, Richardson AE, O'Callaghan M, DeAngelis KM, Jones EE, Stewart A, et al. Effects of

selected root exudate components on soil bacterial communities. FEMS Microbiol Ecol

2011; 77: 600–610.

95. Schaeffer SM, Homyak PM, Boot CM, Roux-Michollet D, Schimel JP. Soil carbon and

nitrogen dynamics throughout the summer drought in a California annual grassland. Soil Biol

Biochem 2017; 115: 54–62.

96. Fuchslueger L, Bahn M, Fritz K, Hasibeder R, Richter A. Experimental drought reduces the

transfer of recently fixed plant carbon to soil microbes and alters the bacterial community

composition in a mountain meadow. New Phytol 2014; 201: 916–927.

97. Davidson EA, Belk E, Boone RD. Soil water content and temperature as independent or

confounded factors controlling soil respiration in a temperate mixed hardwood forest. Glob

Change Biol 1998; 4: 217–227.

98. Herron PM, Stark JM, Holt C, Hooker T, Cardon ZG. Microbial growth efficiencies across a

soil moisture gradient assessed using 13C-acetic acid vapor and 15N-ammonia gas. Soil Biol

Biochem 2009; 41: 1262–69.

99. Schimel JP. Life in dry soils: Effects of drought on soil microbial communities and

processes. Annu Rev Ecol Evol Syst 2018; 49: 409–432.

100.   NOAA (National Oceanic and Atmospheric Administration). National Climatic Data

Center. 2021. https://www.ncdc.noaa.gov/

 Figures

Figure 1. Experimental design and effect of mycorrhizal inoculation and moisture

manipulation on mycorrhizal abundance and soil moisture after 3 months. Greenhouse

experiment with *Panicum hallii* (Hall's panicgrass) and two mycorrhizal inocula grown in a

marginal soil from central Oklahoma. **a** Illustration of microcosm experimental design and $H_2^{18}O$

stable isotope probing (SIP) incubation experiment conducted with soils collected from hyphal

ingrowth cores that excluded plant roots. *P. hallii* was inoculated with either *R. irregularis*, *S.*

bescii or left uninoculated (indicated in yellow, blue, or black, respectively) and grown for 3

months under water-replete or water-limited conditions (indicated in solid or dashed lines,

respectively; n = 3 replicates per treatment). Soils harvested at 3 months were mixed with either

enriched ($H_2^{18}O$) or natural abundance ($H_2^{16}O$) water and incubated for 7 days. **b** Abundance of

R. irregularis and *S. bescii* (DNA copies g^{-1} soil) in hyphal ingrowth cores after 3 months of

plant growth and moisture manipulation, measured with strain-specific qPCR primers. **c** Soil

moisture after 3-monts. Bold lines represent median value; whiskers represent upper and lower

quartiles (n = 3 replicates per treatment). Letters denote the results of a Wilcoxon rank sum test

performed separately for each qPCR plot (p < 0.1) and a Tukey's HSD test for soil moisture

comparisons (p < 0.001).

Figure 2. Structure and alpha diversity of total and active microbial communities based on

traditional and $H_2^{18}O$ qSIP-filtered 16S rRNA gene profiling. Principal coordinates analysis

(PCoA) of weighted UniFrac distances between 'total' microbial communities assessed with (**a**)

traditional 16S rRNA gene analysis and (**b**) $H_2^{18}O$ qSIP-filtered 'actively growing' communities

(i.e., ASVs that did not incorporate a significant quantity of ^{18}O were removed prior to analysis).

Moisture regime and mycorrhizal inoculum explained a total of 29.5% of the variation in 'total'

community structure ($p < 0.001$; n = 6 replicates) and 75.2% of variation in actively growing

community structure ($p < 0.001$; n = 6 replicates). Ellipses show treatment groupings in (**a**) and

(**b**). **c** Inverse Simpson's diversity index in total and actively growing communities. Letters

denote the results of a Tukey's HSD test (no significant differences between total communities; p

< 0.01 for comparisons between active communities; n = 6 replicates). Bold lines represent

median values; whiskers represent upper and lower quartiles. For all plots, uninoculated, *R.*

irregularis-inoculated, and *S. bescii*-inoculated soils are represented in black, yellow, and blue,

respectively. Water-replete and water-limited soils are represented in dark versus light color

shades, respectively.

Figure 3. H$_2$^{18}O qSIP-based microbial activity in hyphal ingrowth core soils exposed to different moisture regimes and mycorrhizal inocula. a Median ^{18}O APE of all bacterial and archaeal ASVs present under each treatment. Uninoculated, *R. irregularis*-inoculated, and *S. bescii*-inoculated soils are represented in black, yellow, and blue, respectively. Water-replete and water-limited soils are represented in solid and dashed lines, respectively. Letters denote the results of a Tukey's HSD test comparing the means of all treatments ($p < 0.01$; $n = 3$ replicates). **b** Taxon-specific gross growth rates (ng DNA g^{-1} soil day^{-1}) based on ASV ^{18}O incorporation,

920   summed by phylum for each treatment. **c** CO_2 efflux ($\mu g\ CO_2\ mg^{-1}$ soil C day^{-1}) from hyphal

921   ingrowth core soils during $H_2^{18}O$ qSIP assay. Bold lines represent median values; whiskers

922   represent upper and lower quartiles ($p < 0.05$; n = 6 replicates per treatment).

Figure 4. Comparison of taxon-specific activity in water-limited versus water-replete hyphal ingrowth core soils within each mycorrhizal treatment. ^{18}O atom percent excess (APE) 'active growth' ratios of ASVs present in both water-limited and water-replete soils that were either **(a)** uninoculated, **(b)** inoculated with *R. irregularis,* or **(c)** inoculated with *S. bescii*. Ratios were averaged at the phylum level. Ratios less than 1 (bars located to the left of the central line) indicate that water limitation suppressed growth. Ratios greater than 1 (bars located to the right of the central line) indicate that water limitation facilitated greater growth rates. Error bars represent the standard error. Asterisks denote phylum-level averages that are significantly

greater than or less than 1 (adjusted p < 0.05 based on Wilcoxon signed rank test and Benjamini-

Hochberg correction for multiple comparisons). Only taxa that incorporated significant ^{18}O

(lower 90% CI > 0) and belong to the 10 most abundant phyla are represented.

Figure 5. Comparison of taxon-specific activity in mycorrhizal-inoculated versus

uninoculated hyphal ingrowth core soils following water limitation. ^{18}O atom percent excess

(APE) 'active growth' ratios of bacterial ASVs present in water-limited soils that were either

uninoculated or inoculated with *R. irregularis* or *S. bescii*, averaged at the phylum level **(a,b)** or

displayed by individual ASV **(c,d)**. Ratios greater than 1 (to the right of the central line) indicate

that mycorrhizal fungi supported greater bacterial growth rates. For phylum level averages,

asterisks represent ^{18}O APE ratios that are significantly greater than 1 (adjusted $p < 0.05$ based

on Wilcoxon signed rank test and Benjamini-Hochberg correction for multiple comparisons).

Error bars represent the standard error within each bacterial phylum. Only taxa that incorporated

significant ^{18}O (lower 90% CI > 0) and belong to the 10 most abundant phyla are represented.

ASVs belonging to phyla that incorporated significantly more ^{18}O in mycorrhizal-inoculated

soils are represented in **c,d**.