Title: Plant-associated fungi support bacterial resilience following water limitation

Rachel Hestrin[1,2]*, Megan Kan[1], Marissa Lafler[1], Jessica Wollard[1], Jeffrey A. Kimbrel[1], Prasun Ray[3,4], Steven Blazewicz[1], Rhona Stuart[1], Kelly Craven[4,5], Mary Firestone[6], Erin Nuccio[1], Jennifer Pett-Ridge[1,7]*

[1]*Lawrence Livermore National Laboratory, Physical and Life Science Directorate, Livermore, CA 94550 USA*

[2]*University of Massachusetts, Stockbridge School of Agriculture, Amherst, MA 01003 USA*

[3]*University of Maryland Eastern Shore, Department of Natural Resources, Princess Anne, MD 21853 USA*

[4]*Noble Research Institute, Plant Biology Division, Ardmore, OK 73401 USA*

[5]*Oklahoma State University, Department of Biochemistry and Molecular Biology, Stillwater OK 74078 USA*

[6]*University of California Berkeley, Department of Environmental Science Policy and Management, Berkeley, CA 94720 USA*

[7]*University of California Merced, Life & Environmental Sciences Department, Merced, CA 95343 USA*

*Correspondence to:

Rachel Hestrin, 925-409-7954, rhestrin@umass.edu

Jennifer Pett-Ridge, 925-337-7618, pettridge2@llnl.gov

23   **Competing Interests:** The authors declare no competing interests.

Abstract

Drought disrupts soil microbial activity and many biogeochemical processes. Although plant-associated fungi can support plant performance and nutrient cycling during drought, their effects on nearby drought-exposed soil microbial communities are not well resolved. We used $H_2{}^{18}O$ quantitative stable isotope probing (qSIP) and 16S rRNA gene profiling to investigate bacterial community dynamics following water limitation in the hyphospheres of two distinct fungal lineages (*Rhizophagus irregularis* and *Serendipita bescii*) grown with the bioenergy model grass *Panicum hallii*. In uninoculated soil, a history of water limitation resulted in significantly lower bacterial growth potential and growth efficiency, as well as lower diversity in the actively growing bacterial community. In contrast, both fungal lineages had a protective effect on hyphosphere bacterial communities exposed to water limitation: bacterial growth potential, growth efficiency, and the diversity of the actively growing bacterial community were not suppressed by a history of water limitation in soils inoculated with either fungus. Despite their similar effects at the community level, the two fungal lineages did elicit different taxon-specific responses, and bacterial growth potential was greater in *R. irregularis*- compared in *S. bescii*-inoculated soils. Several of the bacterial taxa that responded positively to fungal inocula belong to lineages that are considered drought-susceptible. Overall, $H_2{}^{18}O$ qSIP highlighted treatment effects on bacterial community structure that were less pronounced using traditional 16S rRNA gene profiling. Together, these results indicate that fungal-bacterial synergies may support bacterial resilience to moisture limitation.

Keywords: *Panicum hallii*, mycorrhizal fungi, bacteria, drought, quantitative stable isotope probing, plant-microbe interactions, ${}^{18}O$-water

Introduction

Drought alters plant productivity[1], soil microbial biomass[2,3] and community composition[4,5], greenhouse gas emissions[6], and many critical biogeochemical processes[7]. Plant-microbial mutualisms mitigate plant drought response and may aid in post-drought recovery through a variety of mechanisms[8,9]. In particular, mutualistic root-associated fungi—such as mycorrhizal fungi, which form a symbiosis with most terrestrial plant families[10]—can support plants during drought by facilitating water transport[11], soil aggregation[12], root growth[13], plant nutrient uptake[14], photosynthesis[15,16], and stomatal conductance[15-17]. While mycorrhizal fungi may also influence the microbial communities that mediate nutrient cycling and other processes in drought-affected soil, multipartite plant-fungal-bacterial feedbacks remain poorly quantified[9].

The soil hyphosphere—the region that surrounds fungal hyphae—is a hotspot for fungal-bacterial interactions that influence microbial community composition[18-22], nutrient cycling[20-27], and plant growth[27,28]. Interactions between soil bacteria and plant-associated hyphae are likely shaped by resource dynamics. Plants share up to 20% of photosynthates with their mycorrhizal symbionts[29,30], which can rapidly transport these resources to surrounding bacteria[24,31]. Because fungi may explore a volume of soil that is two orders of magnitude greater than the area explored by plant roots[32], they may exert a substantial effect on soil microbiome structure and function. Distinct bacterial communities form in proximity to different fungal lineages[18,19,31]. However, we have a limited understanding of how different plant-associated fungi may shape the soil microbiome's response to environmental stress.

Investigation of fungal-bacterial interactions in hyphosphere soil is methodologically challenging

because the hyphosphere is small, dynamic, and may not exert a detectable effect on soil

microbial community composition or activity when assessed at a "bulk" scale[18]. Furthermore, a

substantial proportion of soil DNA may represent inactive organisms or extracellular "relic"

DNA[33,34]. Stable isotope probing (SIP) coupled with 16S rRNA gene profiling can distinguish

active organisms from those that are inactive by tracing isotope incorporation into newly

synthesized microbial DNA[35,36]. With quantitative SIP (qSIP), we can estimate taxon-specific

growth based on shifts in DNA buoyant density caused by heavy isotope incorporation[37,38]. This

enables sensitive detection of actively growing taxa, even in samples where a substantial quantity

of DNA belongs to dead or dormant organisms. In challenging experimental systems like the

hyphosphere, qSIP has the potential to provide novel insight into taxon-specific activity and

microbial growth potential following experimental treatments.

In this study, we investigated how two plant-associated fungal lineages—the arbuscular

mycorrhizal (AM) fungus *Rhizophagus irregularis* and the *Sebacinales* fungus *Serendipita*

bescii—mediate bacterial growth potential following water limitation in a marginal soil planted

with *Panicum hallii* (Hall's panicgrass), a model species closely related to the bioenergy crop

switchgrass. Both AM and *Sebacinales* fungi associate with a wide range of plant species,

including switchgrass[39-41], and support plant growth and nutrition during drought[42-44]. However,

R. irregularis has a reduced enzymatic repertoire[45,46] and depends largely upon host-provided C

and microbial transformation of nutrient sources into bioavailable forms[27,47,48]. In contrast, *S.*

bescii and other lineages of *Serendipita* are facultative symbionts with a broader enzymatic

repertoire that enables direct resource acquisition from both living plants and detritus[49-51]. We

hypothesized that both fungi would mitigate the effects of moisture limitation, but that bacterial

community composition and growth potential would be distinct in the soils colonized by each

fungus.

Materials and Methods

Soil collection and characterization

Soil used for this study was a Pond Creek fine sandy loam, classified as a superactive, thermic

Pachic Argiustoll[52]. Soil was collected from a pasture in Caddo County, OK (35.072417/-

98.303667) where switchgrass is endemic, on traditional land of the Anadarko (Nadaco) tribe

and the Caddo Nation of Oklahoma. Previous work has characterized this soil as "marginal" due

to its high sand content (69%), low pH (~5), and low C, N, and P content (< 0.4%, < 0.04%, and

< 6 ppm, respectively) (ref. 53). Surface soil (0-20 cm) was collected in May 2019, transported

to Livermore, CA, sieved to 2 mm, and stored at 4 °C. A soil moisture retention curve was

generated from air-dried soil using a pressure plate apparatus (WP4C, METER Environment,

Pullman, WA) and the nonlinear fitting program SWRC-Fit to apply a Brooks and Corey model

to the data[54].

Fungal inoculum

Spores of *R. irregularis* (formerly *Glomus intraradices*) isolate DAOM-197198 were purchased

from Premier Tech (Rivière-du-Loup, Quebec, Canada). *S. bescii* (sourced from the Noble

Research Institute) was grown in modified Melin-Norkran's (MMN) broth. Bentonite clay

particles were mixed with MMN broth, inoculated with *S. bescii,* and incubated at 24 °C for 8

weeks according methods described by Ray et al. (ref. 55). Uninoculated bentonite clay particles

were also mixed with MMN broth and incubated under the same conditions.

Greenhouse water limitation experiment

P. hallii seeds collected from the Edwards Plateau in central Texas were grown to seed, scarified,

surface sterilized, stratified at 4 °C for one week, and germinated in Petri plates. After five days,

germinated seedlings were transferred into planting cones (Ray Leach Cone-tainers, Steuwe &

Sons, Tangent, OR, USA) filled with double-autoclaved sand. Five days after transfer into cones,

~500 *R. irregularis* spores or 500 µL *S. bescii* inoculum were injected into the sand to inoculate

the *P. hallii* seedlings. A subset of seedlings was left uninoculated. Eight weeks after inoculation,

seedlings were transplanted into 960 cm^3 containers (Anderson Plant Bands, Steuwe & Sons)

filled with a 50:50 (v:v) mixture of live soil and double-autoclaved sand (CEMEX Lapis Lustre

Specialty Sand, #2/12) packed to 1.7 g mL^{-1} bulk density at 15% moisture. Additional inocula

(consisting of 500 *R. irregularis* spores or 5 g bentonite clay particles coated with *S. bescii*) were

placed near the roots of seedlings as they were transplanted. Five g of uncoated bentonite clay

particles were placed near the roots of previously uninoculated plants and plants that had been

inoculated with *R. irregularis*. Each microcosm contained a 25 µm mesh hyphal ingrowth core

(2.3 cm diameter) filled with 75 g of the same live soil (no sand) mixed with 0.5 g finely milled

switchgrass biomass as bait for the fungal inocula. The multi-phase fungal inoculation procedure

was intended to give *R. irregularis* and *S. bescii* a colonization advantage over native fungal

endophytes present in the soil.

After assembly, each microcosm was covered with 150 g double-autoclaved sand to inhibit

cross-contamination of fungal inocula. All microcosms were watered with a total of 20 mL

ultrapure H_2O during the first week to facilitate plant and fungal establishment. Half of the

microcosms were watered on a weekly or bi-weekly schedule to maintain soil moisture at

approximately 15% (based on gravimetric measurements), a level that would not restrict plant

growth. The other half of the microcosms were not watered for the remainder of the experiment

and declined to 5% soil moisture by three months. In a subset of microcosms, the moisture

content of the soil surrounding the roots (i.e., not within the hyphal ingrowth cores) was

monitored continuously with a volumetric water probe (ECH$_2$O EC-5, METER Group, Inc.,

Pullman, WA, USA). Another microcosm per treatment was weighed weekly to assess whole-

microcosm gravimetric changes. Three replicate microcosms were maintained for each of the six

treatment combinations (three fungal inoculum conditions x two moisture regimes). Average

daytime and nighttime temperatures were 27 °C and 24 °C, respectively, with a photoperiod of

16 h. After three months, the microcosms were destructively harvested. Soil from hyphal

ingrowth cores was homogenized, flash frozen in liquid N_2, and stored at -80 °C for DNA

extractions or at room temperature for qSIP assays (see Fig. 1a for experimental design).

Soil characteristics

Soil moisture content following the three-month harvest was determined by mass difference after

drying soil for 48 h at 105 °C. Total soil C and N contents were measured with a Costech ECS

4010 Elemental Analyzer (Costech Analytical Technologies Inc., Valencia, CA).

Soil H$_2$^{18}O qSIP assay

To determine bacterial growth potential following water limitation, we conducted an $H_2^{18}O$ qSIP

assay[35,37,38]. Three days after the three-month harvest, soil from three hyphal ingrowth cores per

treatment was amended with either $H_2^{16}O$ or $H_2^{18}O$ (Fig. 1a). Each soil sample was divided into

three subsamples: one 3.0 (+/- 0.2) g dry weight equivalent sample for an initial $H_2^{16}O$ qSIP

timepoint (T0) and two 4.0 (+/- 0.2) g dry weight equivalent samples for the $H_2^{16}O$ and $H_2^{18}O$

seven-day qSIP timepoint (T7). Two of the T0 samples contained ~2.0 g due to limited soil

availability. This resulted in a total of 54 samples (18 T0 $H_2^{16}O$ samples, 18 T7 $H_2^{16}O$ samples,

and 18 T7 $H_2^{18}O$ samples). Each subsample was air-dried to 4.7% gravimetric water content in a

biosafety cabinet and then brought up to 22.1% gravimetric water content (60% field capacity)

with either ultrapure water at natural isotopic abundance or water enriched with ^{18}O (98.38 atom

% $H_2^{18}O$, Isoflex, San Francisco, CA, USA). The final estimated enrichment of the samples

containing $H_2^{18}O$ was 78.76 ^{18}O atom %. We standardized the soil moisture and soil water

isotopic enrichment across treatments to minimize differences across fungal treatments during

the growth potential assay. The T0 subsample (amended with $H_2^{16}O$) was immediately flash

frozen in liquid N_2 and stored at -80 °C. Each of the T7 soil samples was stored in the dark inside

a separate 473.2 mL glass jar with a tight-fitting lid containing a septum for gas sampling. After

seven days, the soils were flash frozen in liquid N_2 and stored at -80 °C.

CO_2 efflux measurement

15 mL headspace gas samples were collected at the beginning and end of the qSIP assay. CO_2

concentrations were measured on a gas chromatograph equipped with a thermal conductivity

detector (GC-14A, Shimadzu, Columbia, MD).

DNA extraction and density gradient fractionation

DNA was extracted from each soil sample in quadruplicate with the DNeasy PowerSoil Pro kit (Qiagen, Germantown, MD, USA) and then pooled per sample prior to downstream analysis. To separate isotopically enriched DNA from unenriched DNA, samples were subjected to a cesium chloride density gradient formed in an ultracentrifuge as previously described[56] with the following minor modifications. For each sample, 5 µg of DNA in 150 µL 1xTE buffer was mixed with 1.00 mL gradient buffer, and 4.60 mL CsCl stock (1.885 g mL^{-1}) with a final average density of 1.730 g mL^{-1}. Samples were loaded into 5.2 mL ultracentrifuge tubes and spun at 20 ºC for 108 hours at 176,284 RCF$_{avg}$ (equivalent to 176,284 x g) in a Beckman Coulter Optima XE-90 ultracentrifuge using a VTi65.2 rotor. Sample fractionation was automated using Lawrence Livermore National Laboratory's high-throughput SIP pipeline[57], which automates the fractionation and clean-up tasks for the density gradient SIP protocol. The content of each ultracentrifuge tube was fractionated into 22 fractions (~236 µL each) using an Agilent Technologies 1260 isocratic pump to deliver water at 0.25 mL min^{-1} through a 25G needle inserted through the top of the ultracentrifuge tube. Each tube was mounted in a Beckman Coulter fraction recovery system with a side port needle inserted through the bottom. The side port needle was routed to an Agilent 1260 Infinity fraction collector, and fractions were collected in 96-well deep well plates. The density of each fraction was measured using a Reichart AR200 digital refractometer fitted with a prism covering to facilitate measurement from 5 µL, as previously described[58]. DNA in each fraction was purified and concentrated using a Hamilton Microlab Star liquid handling system programmed to automate glycogen/PEG precipitations[36] (GlycoBlue Coprecipitant, Invitrogen by Thermo Fisher Scientific, Waltham, MA, USA).

205 Washed DNA pellets were suspended in 40 µL of 1xTE buffer and the DNA concentration of

206 each fraction was quantified using a PicoGreen fluorescence assay.

Quantitative PCR (qPCR) to assess bacterial and fungal abundance

We conducted qPCR with "universal" bacterial primers[59,60] to measure 16S rRNA gene copy

number in each unfractionated and SIP-fractionated DNA sample (see Table S1 and SI text for

additional information). To assess the relative abundance of fungal inocula, we quantified *R.*

irregularis and *S. bescii* gene copy number in unfractionated DNA (extracted in triplicate prior

to the $H_2^{18}O$ qSIP assay) with qPCR primers designed specifically for each fungal lineage[55,61]

(see Table S1 and SI text for additional information). We note that DNA can be unevenly

distributed throughout fungal biomass. While fungal DNA copy number may not necessarily be

indicative of total biomass or activity, it is a relative indicator of inoculation success[62,63].

DNA sequencing

For each sample, we sequenced the 16S rRNA gene in unfractionated DNA as well as DNA

fractions containing > 1.0 ng $µL^{-1}$, resulting in 6-11 sequenced fractions per sample. The V4

region of the 16S rRNA gene was amplified with the 515F/806R primer pair[59,60], processed and

barcoded through the Illumina V2 PE150 sample preparation kit, and sequenced on a MiSeq v2

platform in three runs (Illumina, Inc., San Diego, CA; see SI text and Tables S2 & S3 for

additional information about taxonomic assignment and amplicon sequence variant (ASV)

recovery).

Quantitative stable isotope probing (qSIP) analysis

We calculated taxon-specific growth potential using the $H_2^{18}O$ qSIP approach[35,37,38]. Briefly, the

qSIP mathematical model estimates taxon-specific ^{18}O incorporation into DNA based on the shift

in DNA density following exposure to natural abundance H_2O ($H_2^{16}O$) or "heavy" $H_2^{18}O$. We

determined baseline densities for each taxon in the absence of an isotopic tracer (i.e., samples

amended with $H_2^{16}O$), because even without tracer assimilation, DNA density varies by GC

content[64]. For each ASV, we calculated median ^{18}O atom percent excess (APE) based on the

difference in the buoyant density of 16S rRNA gene profiles sequenced from soils amended with

$H_2^{16}O$ or $H_2^{18}O$. To calculate taxon-specific growth potential, we assumed linear population

growth and used the change in taxon-specific ^{18}O enrichment and 16S rRNA gene abundance

over the course of the qSIP assay (i.e., at T0 and at T7) to estimate taxon-specific growth

potential with the equation below. In this equation, absolute growth rate for taxon i is estimated

as b_i, where for each taxon i at the end of the qSIP assay (time t), $N_{TOTAL_{it}}$ and $N_{LIGHT_{it}}$ are the

total (unlabeled + ^{18}O-labeled) and unlabeled 16S rRNA gene abundances, respectively.

$$b_i = \frac{N_{TOTAL_{it}} - N_{LIGHT_{it}}}{t}$$

We estimated 16S rRNA gene abundance by multiplying the relative abundance of each ASV's

16S rRNA gene by the total 16S rRNA gene copy number measured in each DNA fraction with

qPCR. These calculations rest upon the assumptions that a) bacterial DNA synthesis is

proportional to cellular growth and division, b) for each ASV's population, all new bacterial

DNA is synthesized with an equal proportion of ^{18}O, and c) bacterial taxa have an average of six

16S rRNA gene copies per cell[37]. We note that qSIP-based growth estimates are approximations

due to several methodological challenges, including incomplete extraction of microbial DNA

from soil[65], inter-taxonomic variation in 16S rRNA gene copies per cell[66], and amplification and

sequencing biases[67,68]. To increase confidence in these estimates, we first filtered out ASVs that

were not recovered from all replicates of a particular treatment. This stringent filtering ensured

that our estimates were representative of the three independent biological replicates. To quantify

uncertainty around each estimate of taxon-specific ^{18}O APE and growth, we calculated 90%

confidence intervals using a bootstrapping procedure with 1000 iterations[37].

^{18}O APE represents an integrated measure of new DNA synthesis that occurred during the qSIP

assay. We note that while increases in a taxon's ^{18}O APE indicates new growth, the overall

population size might be increasing in abundance, remaining static, or declining depending on

whether or not growth outpaces mortality. During population turnover, DNA ^{18}O APE can

increase while population-level abundance either increases (if growth outpaces mortality),

remains similar (if growth matches mortality), or declines (if mortality outpaces growth).

Therefore, a high ^{18}O APE value is not necessarily indicative of a net increase in abundance. In

our study, because we maintained all soils at 22.1% moisture during the qSIP assay, growth

estimates represent bacterial growth potential immediately following twelve weeks of exposure

to contrasting moisture regimes, rather than actual growth rates under water-limited or water-

replete conditions.

We calculated a metric of growth efficiency by dividing gross bacterial growth by CO_2 efflux per

treatment. Microbial growth efficiency can be defined as the proportion of biomass synthesized

per unit C assimilated[69] or the proportion of C allocated towards growth rather than towards other

activities[70]. We consider the ratio of new DNA synthesis to CO_2 efflux to be a proxy for growth

efficiency since DNA synthesis is proportional to new microbial biomass production, and since

CO_2 efflux represents respiration from microbial activities (including both growth and non-

growth processes). Based on this metric, a higher proportion of non-growth activities (such as

metabolism, osmoregulation, motility, energy spilling reactions, O_2 stress responses, and other

maintenance activities) results in a decrease in growth efficiency.

Statistical analyses

We performed all statistical analyses in R[71]. We assessed differences in fungal inoculum gene

copy number with a non-parametric Kruskal-Wallis rank sum test. We assessed differences in

soil moisture, soil C and N content, DNA extraction efficiency, Inverse Simpson's diversity

index, median ASV ^{18}O APE, and CO_2 efflux with a Tukey's HSD test comparing the means of

all treatments. First, we created a linear model with a fixed effect for treatment. If the raw data

did not meet the assumptions of normality, it was log-transformed prior to further analysis.

We used the R package vegan to perform a principal coordinates analysis and permutational

multivariate analysis of variance (PERMANOVA) on the weighted UniFrac distance between

communities present following different moisture and fungal treatment combinations and at

different qSIP timepoints[72,73].

We used the 90% confidence interval around each ASV's median ^{18}O APE to identify ASVs that

incorporated significant quantities of ^{18}O into their DNA: only ASVs whose lower 90%

confidence interval did not overlap with zero were considered to be actively growing. To

compare taxon-specific growth potential between two treatments, we calculated pairwise ratios

of ^{18}O-based growth potential for taxa that were actively growing under both conditions. We

calculated these ratios by dividing an ASV's ^{18}O APE following one fungal*moisture treatment

by its ^{18}O APE following the other fungal*moisture treatment. For example:

$$\text{ASV 18O APE Ratio} = \frac{\text{ASV 18O APE in water} - \text{limited soil with } R.\ irregularis}{\text{ASV 18O APE in water} - \text{replete soil with } R.\ irregularis}$$

To assess whether these ratios were significantly greater than or less than one at the phylum level

(indicating a significant difference between growth potential in soils maintained under different

fungal or moisture conditions), we conducted a Wilcoxon signed rank test with a Benjamini-

Hochberg correction for multiple comparisons on the average of the ASV ^{18}O APE ratios within

each phylum.

Results

Plant growth, fungal abundance, and soil characteristics following three-month water

manipulation

After three months of growth, *P. hallii* plants reached similar mass across all fungal*moisture

treatments (Fig. S1; $p > 0.05$). *R. irregularis* and *S. bescii* abundances (measured with strain-

specific qPCR primers) were higher in the hyphal ingrowth cores of microcosms inoculated with

each fungus than in the ingrowth cores of microcosms that were not inoculated (Tables S4 & S5,

Fig. 1b,c; $p < 0.001$). Soil moisture within the water-replete hyphal ingrowth cores was more

than three times higher than the moisture content within the water-limited hyphal ingrowth cores

(Fig. 1d; $p < 0.001$). Matric suction was higher in water-limited soils, but did not reach a point

likely to cause a loss in cell turgor[74] (Fig. S2). Neither plant biomass nor fungal inoculum

abundance was associated with differences in soil moisture (Figs S1 & 1d; $p > 0.05$). Total soil

C, N, and DNA extracted did not vary between treatments (Figs S3 & S4a; $p > 0.05$).

Effects of water limitation and fungal inoculum on bacterial community structure

Moisture history and fungal inoculum shaped the structure of the bacterial communities present

in hyphal ingrowth core soils (Table S6 & Fig. 2a; $p < 0.001$, PERMANOVA of weighted

UniFrac distances). We detected only a small shift in bacterial community structure between the

beginning and end of the seven-day qSIP assay in the unfractionated samples (6.2% of variance

explained; Table S7; $p < 0.001$). With this traditional 16S rRNA gene profiling of unfractionated

DNA (sequenced from soils collected at the end of the qSIP assay), we found that moisture

regime and fungal inoculum explained only 30.0% of the variation between "total" bacterial

communities (i.e., bacteria identified in the total DNA pool, which many include dead and

dormant organisms).

We used $H_2{}^{18}O$ qSIP to filter out inactive ASVs (i.e., those that did not incorporate significant

quantities of ^{18}O into their DNA) and identify the actively growing subset of the total bacterial

community. In this subset, we found that moisture history, fungal inoculum, and their interactive

effect explained 86% of the variation in community structure (11.9%, 49.5%, and 24.6%,

respectively; Table S8 & Fig. 2b; $p < 0.001$, PERMANOVA). This is more than twice the effect

size observed through traditional 16S rRNA gene profiling. qSIP-based filtering also highlighted

differences in bacterial alpha diversity. There was no relationship between moisture history or

fungal inoculum and the alpha diversity of the total communities sequenced (Fig. 2c,d; $p > 0.05$).

However, for the actively growing communities, water limitation was associated with lower

alpha diversity in uninoculated soils ($p < 0.01$), no difference in *R. irregularis*-inoculated soils (p

> 0.05), and higher alpha diversity in *S. bescii*-inoculated soils ($p < 0.01$).

Overall, DNA from soil amended with $H_2{}^{18}O$ was denser than DNA from soil amended with

$H_2{}^{16}O$ (Fig. S4b), indicating that many organisms incorporated ^{18}O into newly synthesized DNA.

Our qSIP filtering parameters removed ASVs that were not identified in all replicates of a

particular treatment, resulting in 989-1,153 bacterial ASVs per treatment (Table S3). The ASVs

that remained following this stringent filtering step represent 62-73% of the ASVs sequenced per

treatment in unfractionated DNA and 21-25% of the ASVs sequenced per treatment in the SIP-

fractionated DNA (Tables S2 & S3). Of the ASVs that remained after qSIP filtering, taxon-

specific ^{18}O APE ranged from unenriched to 64.2 APE, with a median of 6.1. Between 30 and

60% of the total bacterial ASVs detected per treatment were enriched (Table S3; lower 90% CI >

0); we refer to these as actively growing taxa. We note that although the stringent filtering

parameters increase confidence in the taxon-specific APE estimates, these may result in an

overestimate of the proportion of actively growing taxa if less abundant taxa have lower growth

rates and are disproportionately filtered from the dataset.

Effects of water limitation and fungal inoculum on bacterial growth potential and growth

efficiency

We used $H_2{}^{18}O$ qSIP and CO_2 efflux measurements to quantify the effects of moisture history

and fungal inoculum on bacterial growth potential and growth efficiency. In uninoculated soils,

water limitation was associated with more than a 50% reduction in median ASV ^{18}O APE (Fig.

3a; $p < 0.05$). Similarly, bacterial gross growth potential in uninoculated water-limited soil was

less than one-third of gross growth potential in water-replete soil (Fig. 3b). In uninoculated soils,

moisture history did not result in a difference in total CO_2 efflux during the qSIP assay (Fig. 3c;

$p > 0.05$). Because bacterial growth potential was substantially lower following water limitation,

this translated into a 72% reduction in growth efficiency (gross growth potential divided by CO_2

efflux) in uninoculated soils (Table S9).

In fungal-inoculated soils, bacterial communities were less affected by water limitation than in

uninoculated soils. Moisture history was not associated with a difference in bacterial ASV ^{18}O

APE or potential gross growth in soil inoculated with either *R. irregularis* or *S. bescii* (Fig. 3a,b;

$p > 0.05$ for ^{18}O APE). CO_2 efflux from *S. bescii*-inoculated soils was higher than from *R.*

irregularis-inoculated soils, but the difference was only significant between the water-replete

soils (Fig. 3c; $p < 0.05$). Compared to uninoculated soils, water limitation in fungal-inoculated

soils had a less severe effect on potential bacterial growth efficiency, leading to a 17% reduction

in *R. irregularis*-inoculated soils and a 37% reduction in *S. bescii*-inoculated soils (Table S9).

Taxon-specific response to water limitation and fungal inoculum

Since compensatory dynamics may obscure taxon-specific response to different conditions[75], we

calculated ratios of ^{18}O APE for each actively growing ASV detected under two conditions. For

example, for ASVs detected in both water-limited and water-replete soil, a ratio of greater than

1.0 suggests that the ASV could sustain higher growth potential in water-limited soil, a ratio of

less than 1.0 suggests that the ASV could sustain higher growth potential in water-replete soil,

and a ratio indistinguishable from 1.0 suggests that the ASV could maintain similar growth

potential regardless of moisture history. In the absence of fungal inocula, most ASVs were

suppressed by water limitation (Fig. 4a). When averaged at the phylum level, ^{18}O APE ratios

were significantly less than 1.0 for six of the ten most abundant phyla: *Actinobacteria,*

Bacteroidetes, *Chloroflexi, Gemmatimonadetes, Proteobacteria*, and *Verrucomicrobia* (Fig. 4a;

adjusted $p < 0.05$). In contrast, bacterial growth potential in fungal-inoculated soils was less

strongly affected by moisture history: while water limitation was associated with lower growth

potential for many ASVs, many others sustained similar or greater growth potential. When

averaged at the phylum level, populations belonging to the phyla *Actinobacteria* and

Gemmatimonadetes were suppressed in water-limited soil inoculated with *R. irregularis* (Fig. 4b;

adjusted $p < 0.05$), while *Verrucomicrobia* were suppressed in water-limited soil inoculated with

S. bescii (Fig. 4c; adjusted $p < 0.05$). Only the *Acidobacteria* present in *R. irregularis*-inoculated

soil sustained higher growth potential following water limitation (Fig. 4b; adjusted $p < 0.05$).

To assess whether each fungus promoted or suppressed growth potential of specific bacteria in

water-limited soil, we calculated ^{18}O APE ratios for ASVs detected in both uninoculated and

fungal-inoculated soils following water limitation. Most ASVs sustained higher growth potential

in *R. irregularis-* or *S. bescii*-inoculated soil than in uninoculated soil (Figs 5 & S5). Averaged at

the phylum level, populations belonging to the phyla *Acidobacteria, Actinobacteria,*

Bacteroidetes, *Chloroflexi, Planctomycetes, Proteobacteria*, and *Verrucomicrobia* were

significantly more active in *R. irregularis*-inoculated soil (Fig. 5a; adjusted $p < 0.05$). In *S.*

bescii-inoculated soils, only the population belonging to the phylum *Actinobacteria* was

significantly more active than in uninoculated soil (Fig. 5b; adjusted $p < 0.05$). Some ASVs

responded positively to both *R. irregularis* and *S. bescii*; others responded positively only to one

fungal lineage (Fig. 5c-e). Overall, the magnitude of positive growth response was greater in *R.*

irregularis- relative to *S. bescii*-inoculated soils.

Several bacterial ASVs incorporated significant quantities of ^{18}O into their DNA under some

conditions, but not under others, as indicated by differences in gross growth potential measured

through the H_2^{18}O qSIP assay (Figs S6-S15). For any ASV that did not incorporate a significant

quantity of ^{18}O into its DNA under a particular moisture*fungal condition, it was not possible to

calculate an ^{18}O APE ratio to compare the ASV's growth potential under that moisture*fungal

condition to its growth potential under another condition. This was most apparent for the

Firmicutes in uninoculated soil: none of the *Firmicutes* ASVs incorporated significant quantities

of ^{18}O into their DNA following water limitation (Fig. S11). Therefore, the effect of the fungal

inocula on *Firmicutes* growth potential following water limitation is not represented in the ^{18}O

APE ratios (Fig. 5a,b; no *Firmicutes* bar shown). However, several *Firmicutes* were active in

fungal-inoculated soils following moisture limitation. More than double the number of

Firmicutes ASVs were active in *R. irregularis*- compared to *S. bescii*-inoculated soils.

Discussion

We found that plant-associated fungi have a protective effect on bacterial communities exposed

to water limitation, and that bacterial responses to different fungal lineages are distinct. Because

plant biomass was similar across all conditions investigated, we attribute differences in bacterial

community composition and growth potential to direct effects of moisture history and fungal

inocula, rather than to indirect effects mediated by plants. We found that H_2^{18}O qSIP highlighted

treatment differences that were not apparent through traditional 16S rRNA gene profiling. This

demonstrates the utility of DNA qSIP for investigation of the soil hyphosphere and other systems

in which it is difficult to discern a microbial signal above a complex background community.

Soil water limitation suppresses bacterial growth potential and growth efficiency

We observed a significant decrease in bacterial growth potential following three months of moisture limitation, but little effect of moisture history on respiration potential. Together, these responses resulted in a substantial reduction in bacterial growth efficiency. This may reflect a trade-off between microbial stress tolerance and growth in drought-affected soil[76]. Reduced microbial growth efficiency can occur when biota prioritize essential metabolic activities over cellular growth and replication[7,56,76-78]. Slowed growth also helps bacteria persist in the presence of antibiotics[79,80], which can accumulate in dry soils as microorganisms compete for limited resources[81].

We measured lower growth potential across a broad range of bacterial lineages present in our water-limited soils. Interestingly, this included several monoderm taxa (often referred to as Gram-positive organisms) belonging to the phyla *Actinobacteria*, *Firmicutes*, and *Chloroflexi*. Monoderm taxa have thick cell walls and lack an outer membrane, which can protect them against oxidative damage under dry conditions[7,82-85]. Several studies report that monoderms maintain greater abundance and activity in dry soils compared to diderms (also known as Gram-negative organisms, such as most of the taxa belonging to the phyla *Acidobacteria, Bacteroidetes, Proteobacteria*, and *Verrucomicrobia*)[7,84,85]. In addition to the protection conferred by the structure of their cellular envelope, monoderms may also be poised to outcompete other organisms in drought-affected soils through their capacity to produce antibiotics[81,84,86] or utilize complex C substrates that remain available following water limitation[2,81]. Our results suggest that even monoderm taxa that are considered relatively drought tolerant can be negatively affected by water limitation.

Plant-associated fungi support bacterial resilience in drought-affected soil, but fungal-bacterial relationships are context-dependent

While water limitation had a broad suppressive effect in uninoculated soils, many of the bacteria

in soils inoculated with either *R. irregularis* or *S. bescii* maintained similar growth potential

following cultivation under either water-replete or water-limited conditions. This protective

fungal effect extended throughout the bacterial community, and affected many taxa that are often

considered drought-susceptible. Relative abundances of *Bacteroidetes, Planctomycetes,*

Verrucomicrobia, and many *Proteobacteria* and *Acidobacteria* have been shown to decrease

following drought[4,5,87]. However, we found that many ASVs belonging to these phyla sustained

similar growth potential in fungal-inoculated soils, regardless of moisture treatment. This

suggests that *R. irregularis* and *S. bescii* modified edaphic conditions in some way that broadly

supported bacterial resilience to water limitation. Plant-associated fungi can exude plant-derived

C[24,29-32], promote biofilm formation[88], enhance soil aggregation through their interactions with

other soil biota[14,89], and facilitate bacterial transport through soil[90]. Together, these fungal-

mediated processes could help maintain soil connectivity, microbial activity, and nutrient cycling

under water-limited conditions, thereby preventing bacterial dormancy and death despite a

substantial decline in soil moisture. By supporting bacterial function in drought-affected soil,

plant-associated fungi may counteract the destabilizing effect of moisture stress[91] and improve

capacity for recovery once moisture is restored.

In contrast to their synergistic effects in water-limited soils, *R. irregularis* and *S. bescii* appeared

to suppress bacterial growth potential following water-replete conditions. This demonstrates that

the relationship between plant-associated fungi and hyphosphere bacteria is context-dependent,

and not entirely mutualistic. Plant-associated fungi are known to compete with other biota for

N[92] and P[20,21], and can suppress microbial decomposers[23,93]. Similarly, bacteria can inhibit

mycorrhizal proliferation[20,21,92]. Putative bacterial predators have also been found in greater

abundance on extraradical mycorrhizal hyphae than in surrounding soil[18]. We did not investigate

the potential mechanisms of suppressive interactions between bacteria and plant-associated

fungi. However, our observation that bacterial growth potential depends on moisture history in

fungal-inoculated soil indicates that there are trade-offs between fungal and bacterial growth. In

this trade-off, fungi may limit bacterial growth under resource-replete conditions, but promote

bacterial growth under resource-limited conditions. Similar context-dependency is well-

documented in other mutualisms[94]. In our system, context-dependency may indicate a stabilizing

ecological effect exerted by multipartite hyphosphere interactions.

Since *R. irregularis* and *S. bescii* were not actively associating with their plant hosts in our $H_2^{18}O$

qSIP assay, we attribute these results to fungal effects that had occurred during the preceding

three-month greenhouse experiment. Although the qSIP assay might have caused a nutrient flush

from the perturbed soil and biota, CO_2 efflux from the fungal-inoculated soils was not

significantly greater than from the uninoculated soils. Therefore, we conclude that differences in

bacterial growth potential, growth efficiency, and diversity of the actively growing community

present in fungal-inoculated soil were related to preceding fungal effects on the soil environment

rather than to bacterial decomposition of fungal necromass.

Magnitude of bacterial response is fungal lineage-dependent

While both fungal lineages supported bacterial resilience, *R. irregularis* elicited a stronger

positive response than *S. bescii*—both with respect to the number of ASVs and the magnitude of

individual responses. Distinct microbial consortia associate with different mycorrhizal

lineages[18,19,31,89]. Although empirical evidence remains sparse, different mycorrhizal exudate

profiles, growth habits, and other functional traits may shape the composition and activity of the

surrounding microbial community[48,89,95,96], a phenomenon that has been documented more

extensively for root-microbe interactions[97-99]. Lower bacterial growth potential and growth

efficiency in *S. bescii*- compared to *R. irregularis*-inoculated soils may be due to *S. bescii*'s

wider enzymatic repertoire, which could accelerate decomposition (as indicated by higher CO_2

efflux) or heighten competitive interactions with other soil biota. Additionally, higher gene copy

numbers of *R. irregularis* compared to *S. bescii* detected in hyphosphere soil suggest that *R.*

irregularis colonization levels were more robust. Greater fungal proliferation could be correlated

with greater resource distribution, enhanced soil structure, or other conditions that support

bacterial growth. Together, these findings suggest that diverse fungal lineages promote bacterial

resilience to water limitation, but that the individual taxa and magnitude of taxon-specific

response to each fungus is distinct.

Conclusion

As global precipitation patterns change, it is important to understand how drought influences

ecological functions and microbial interactions, both during and after water limitation. Plant-

associated fungi are known to support plant growth and nutrition in droughted soils, but their

simultaneous effects on the soil bacterial communities that mediate nutrient cycling and other

critical terrestrial processes remain poorly explored. With $H_2{}^{18}O$ qSIP, we show that plant-

associated fungi have a protective effect on bacterial communities exposed to water limitation.

Both the AM fungus *R. irregularis* and the *Sebacinales* fungus *S. bescii* facilitated greater

growth potential, growth efficiency, and diversity of actively growing hyphosphere bacteria in

drought-affected soil. While these divergent fungal lineages stimulated responses of differing

magnitude, the broad patterns were similar, suggesting that the dominant underlying mechanisms

may be conserved across a substantial portion of the bacterial community rather than limited to

interactions with a small number of bacterial taxa. Remarkably, both *R. irregularis* and *S. bescii*

had a protective effect on hyphosphere bacteria exposed to water limitation in a "live" soil,

which may have included functionally redundant fungal lineages. This finding is relevant for

practical evaluation of fungal inoculants, whose ability to persist and elicit a positive effect in

natural settings is not well-established. Additionally, our work demonstrates that $H_2^{18}O$ qSIP is a

useful approach in challenging systems such as the terrestrial hyphosphere, where microbial

dynamics may be difficult to detect with traditional 16S rRNA gene profiling. Together, our

findings demonstrate that context-dependent multipartite relationships support bacterial

resilience to water limitation and may promote post-drought recovery.

Acknowledgements

We thank Christine Hawkes for *P. hallii* seeds; Eric Slessarev for soil collection; Aaron Chew

for measuring soil water potential; Megan Foley, Anne Kakouridis, Benjamin Koch, and Alexa

Nicolas for scientific discussions; Christina Ramon for procurement; Katerina Estera-Molina,

Don Herman, Ilexis Chu Jacoby, Aaron Chew, Christina Fossum, Mengting Yuan, Tasnim

Ahmed, Cynthia-Jeanette Mancilla, Melanie Rodriguez-Fuentes, David Sanchez, Laura Adame,

Emily Kline, Madeline Moore, Jack Hagen, Anne Kakouridis, Ella Sieradzki, Nameer Baker,

Sarah Baker, Alex Greenlon, Heejung Cho, Alexa Nicolas, Peter Weber, Vanessa Brisson, Eric

Slessarev, Gareth Trubl, Dinesh Adhikari, Craig See, Maria Guerra, Rachel Neurath, and Evan

Starr for their assistance with experimental set up, maintenance, and harvests; and Tibisay Perez

and Whendee Silver for assistance with GC access and operation. This work was supported by

the Genomic Science Program of the US Department of Energy Office of Biological and

Environmental Research as part of the LLNL Biofuels SFA (SCW1039) and a sustainable

bioenergy award (PI Firestone) to UC Berkeley (DE-SC0014079), Noble Research Institute,

University of Oklahoma, Lawrence Berkeley National Laboratory, and Lawrence Livermore

National Laboratory (SCW1555). Research at LLNL was performed under the auspices of the

U.S. Department of Energy at Lawrence Livermore National Laboratory under Contract DE-

AC52-07NA27344.

Competing Interests

560   The authors declare no competing interests.

References

1. Leng G, Hall J. Crop yield sensitivity of global major agricultural countries to droughts and

the projected changes in the future. Sci Total Environ 2019; 654: 811–821.

2. Hueso S, García C, Hernández T. Severe drought conditions modify the microbial

community structure, size and activity in amended and unamended soils. Soil Biol Biochem

2012: 50: 167–173.

3. Alster CJ, German DP, Lu Y, Allison, SD. Microbial enzymatic responses to drought and to

nitrogen addition in a southern California grassland. Soil Biol Biochem 2013; 64: 68–79.

4. Bouskill NJ, Lim HC, Borglin S, Salve R, Wood TE, Silver WL, et al. Pre-exposure to

drought increases the resistance of tropical forest soil bacterial communities to extended

drought. ISME J 2013; 7: 384–394.

5. Acosta-Martinez V, Cotton J, Gardner T, Moore-Kucera J, Zak J, Wester D, et al.

Predominant bacterial and fungal assemblages in agricultural soils during a record

drought/heat wave and linkages to enzyme activities of biogeochemical cycling. Appl Soil

Ecol 2014; 84: 69–82.

6. O'Connell CS, Ruan L, Silver WL. Drought drives rapid shifts in tropical rainforest soil

biogeochemistry and greenhouse gas emissions. Nat Commun 2018; 9: 1348.

7. Schimel JP. Life in dry soils: Effects of drought on soil microbial communities and

processes. Annu Rev Ecol Evol Syst 2018; 49: 409–432.

8. Naylor D, Colemann-Derr D. Drought stress and root-associated bacterial communities.

Front Plant Sci 2018; 8: 2223.

9. de Vries FT, Griffiths RI, Knight CG, Nicolitch O, Williams A. Harnessing rhizosphere

microbiomes for drought-resilient crop production. Science 2020; 368: 270-274.

10. Smith SE, Read D. Mycorrhizal symbiosis. third ed. London: Academic Press; 2008. p. 145–

190.

11. Kakouridis A, Hagen JA, Kan MP, Mambelli S, Feldman LJ, Herman DJ, et al. Routes to

roots: direct evidence of water transport by arbuscular mycorrhizal fungi to host plants and

identification of a hyphal transport pathway. bioRxiv

https://doi.org/10.1101/2020.09.21.305409.

12. Rillig MC Mummey DL. Mycorrhizas and soil structure. New Phytol 2006; 171: 41-53.

13. Gong M, You X, Zhang Q. Effects of *Glomus intraradices* on the growth and reactive

oxygen metabolism of foxtail millet under drought. Ann Microbiol 2015; 65: 595–602.

14. Ruiz-Lozano JM. Arbuscular mycorrhizal symbiosis and alleviation of osmotic stress. New

perspectives for molecular studies. Mycorrhiza 2003; 13: 309-317.

15. Morte A, Lovisolo C, Schubert A. Effect of drought stress on growth and water relations of

the mycorrhizal association *Helianthemum almeriense–Terfezia claveryi*. Mycorrhiza 2000;

10: 115–119.

16. Birhane E, Sterck F, Fetene M, Bongers F, Kuyper T. Arbuscular mycorrhizal fungi enhance

photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed

water availability conditions. Oecologia 2012; 169: 895–904.

17. Duan X, Neuman DS, Reiber JM, Green CD, Saxton AM, Augé RM. Mycorrhizal influence

on hydraulic and hormonal factors implicated in the control of stomatal conductance during

drought. J Exp Bot 1996; 47: 1541-1550.

18. Emmett BD, Levesque-Tremblay V, Harrison MJ. Conserved and reproducible bacterial

communities associate with extraradical hyphae of arbuscular mycorrhizal fungi. ISME J

2021; 15: 2276–2288.

19. Toljander JF, Artursson V, Paul LR, Jansson JK, Finlay RD. Attachment of different soil

bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal

vitality and fungal species. FEMS Microbiol Lett 2006; 254: 34–40.

20. Svenningsen NB, Watts-Williams SJ, Joner EJ. et al. Suppression of the activity of

arbuscular mycorrhizal fungi by the soil microbiota. ISME J 2018; 12: 1296–1307.

21. Cruz-Paredes C, Svenningsen NB, Nybroe O, Kjøller R, Frøslev TG, Jakobsen I. Suppression

of arbuscular mycorrhizal fungal activity in a diverse collection of non-cultivated soils.

FEMS Microbiol Ecol 2019; 95: fiz020.

22. Nuccio EE, Hodge A, Pett-Ridge J, Herman DJ, Weber PK, Firestone MK. An arbuscular

mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling

during litter decomposition. Environ Microbiol 2013; 15: 1870–81.

23. Verbruggen E, Jansa J, Hammer EC, Rillig MC. Do arbuscular mycorrhizal fungi stabilize

litter-derived carbon in soil? J Ecol 2016; 104: 261-269.

24. Kaiser C, Kilburn MR, Clode PL, Fuchslueger L, Koranda M, Cliff JP, et al. Exploring the

transfer of recent plant photosynthates to soil microbes: mycorrhizal pathway vs direct root

exudation. New Phytol 2015; 205: 1537-1551.

25. Zhang L, Shi N, Fan J, Wang F, George TS, Feng G. Arbuscular mycorrhizal fungi stimulate

organic phosphate mobilization associated with changing bacterial community structure

under field conditions. Environ Microbiol 2018a; 20: 2639–51.

26. Hodge A, Campbell CD, Fitter AH. An arbuscular mycorrhizal fungus accelerates

decomposition and acquires nitrogen directly from organic matter. Nature 2001; 413: 297-

299.

27. Hestrin R, Hammer EC, Mueller CW, Lehmann J. Synergies between mycorrhizal fungi and

soil microbial communities increase plant nitrogen acquisition. Commun Biol 2019; 2: 233.

28. Medina A, Probanza A, Gutierrez Mañero FJ, Azcón R. Interactions of arbuscular-

mycorrhizal fungi and *Bacillus* strains and their effects on plant growth, microbial

rhizosphere activity (thymidine and leucine incorporation) and fungal biomass (ergosterol

and chitin). Appl Soil Ecol. 2003; 22: 15–28.

29. Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ, et al. Shifting carbon

flow from roots into associated microbial communities in response to elevated atmospheric

CO_2. Proc Natl Acad Sci USA 2010; 107: 10938–10942.

30. Jakobsen I, Rosenthal L. Carbon flow into soil and external hyphae from roots of

mycorrhizal cucumber plants. New Phytol 1990; 115: 77-83.

31. Zhou J, Chai X, Zhang L, George TS, Wang F, Feng G. Different arbuscular mycorrhizal

fungi cocolonizing on a single plant root system recruit distinct microbiomes. mSystems

2020; 5: e00929-20.

32. See CR, Keller AB, Hobbie SE, Kennedy PG, Weber PK, Pett-Ridge J. Hyphae move matter

and microbes to mineral microsites: Integrating the hyphosphere into conceptual models of

soil organic matter stabilization. Glob Change Biol 2022; 28: 2527-2540.

33. Carini P, Marsden P, Leff J, Morgan E, Strickland M, Fierer N. Relic DNA is abundant in

soil and obscures estimates of soil microbial diversity. Nat Microbiol 2017; 2: 16242.

34. Lennon JT, Muscarella ME, Placella MA, Lehmkuhl BK. How, when, and where relic DNA

affects microbial diversity. mBio 2018; 9: e00637-18.

35. Hungate BA, Mau RL, Schwartz E, Caporaso JG, Dijkstra P, van Gestel N, et al. Quantitative

microbial ecology through stable isotope probing. Appl Environ Microbiol 2015; 81: 7570–

7581.

36. Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW, Murrell JC.

DNA stable-isotope probing. Nat Protoc 2007; 2: 860-866.

37. Koch BJ, McHugh TA, Hayer M, Schwartz E, Blazewicz SJ, Dijkstra P, et al. Estimating

taxon-specific population dynamics in diverse microbial communities. Ecosphere 2018; 9:

e02090–15.

38. Blazewicz SJ, Hungate BA, Koch BJ, Nuccio EE, Morrissey E, Brodie EL, Schwartz E, Pett-

Ridge J, Firestone MK. Taxon-specific microbial growth and mortality patterns reveal

distinct temporal population responses to rewetting in a California grassland soil. ISME J

2020; 14: 1520-1532.

39. Kilronomos JN. Host-specificity and functional diversity among arbuscular mycorrhizal

fungi. In: Proceedings of the 8th International Symposium on Microbial Ecology. Eds. Bell

CR, Brylinski M, Johnson-Green P. Halifax: Atlantic Canada Society from Microbial

Ecology; 2000. p. 845-851.

40. Ray P, Guo Y, Chi MH, Krom N, Saha MC, Craven KD. *Serendipita bescii* promotes winter

wheat growth and modulates the host root transcriptome under phosphorus and nitrogen

starvation. Environ Microbiol 2021; 23: 1876–1888.

41. Lee MR, Hawkes CV. Widespread co-occurrence of *Sebacinales* and arbuscular mycorrhizal

fungi in switchgrass roots and soils has limited dependence on soil carbon or nutrients.

Plants, People, Planet 2021; 3: 614-626.

42. Ruiz-Lozano JM, Azcon R, Gomez M. Effects of arbuscular-mycorrhizal glomus species on

drought tolerance: physiological and nutritional plant responses. Appl Environ Microbiol

1995; 61: 456-460.

43. He F, Sheng M, Tang M. Effects of *Rhizophagus irregularis* on photosynthesis and

antioxidative enzymatic system in *Robinia pseudoacacia* L. under drought stress. Front Plant

Sci 2017; 8: 183.

44. Ghimire SR, Craven KD. Enhancement of switchgrass (*Panicum virgatum* L.) biomass

production under drought conditions by the ectomycorrhizal fungus *Sebacina vermifera*.

Appl Environ Microbiol 2011; 77: 7063-7067.

45. Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, et al. Genome of an

arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. Proc Natl

Acad Sci USA 2013; 110: 20117-20122.

46. Kamel L, Keller-Pearson M, Roux C, Ané JM. Biology and evolution of arbuscular

mycorrhizal symbiosis in the light of genomics. New Phytol 2017; 213: 531-536.

47. Bukovská P, Bonkowski M, Konvalinková T, Beskid O, Hujslová M, Püschel D et al.

Utilization of organic nitrogen by arbuscular mycorrhizal fungi-is there a specific role for

protists and ammonia oxidizers? Mycorrhiza 2018; 28: 269-283.

48. Zhang L, Feng G, Declerck S. Signal beyond nutrient, fructose, exuded by an arbuscular

mycorrhizal fungus triggers phytate mineralization by a phosphate solubilizing bacterium.

ISME J 2018; 12: 2339.

49. Zuccaro A, Lahrmann U, Güldener U, Langen G, Pfiffi S, Biedenkopf D, et al. Endophytic

life strategies decoded by genome and transcriptome analyses of the mutualistic root

symbiont *Piriformospora indica*. PLoS Pathog. 2011; 7: e1002290.

50. Ray P, Chi MH, Guo Y, Chen C, Adam C, Kuo A, et al. Genome sequence of the plant

growth promoting fungus *Serendipita vermifera* subsp. *bescii*: The first native strain from

North America**.** Phytobiomes J 2018; 2: 62-63.

51. Dias T, Pimentel V, Cogo AJD, Costa R, Bertolazi AA, Miranda C, et al. The free-living

stage growth conditions of the endophytic fungus *Serendipita indica* may regulate its

potential as plant growth promoting microbe. Front Microbiol 2020; 11: 562238.

52. Moffatt HH. Soil Survey of Caddo County, Oklahoma. United States Department of 836

Agriculture Soil Conservation Service, Washington, D.C. 1973.

53. Sher Y, Baker NR, Herman NR, Fossum C, Hale L, Zhang XX, et al. Microbial extracellular

polysaccharide production and aggregate stability controlled by Switchgrass (*Panicum

virgatum*) root biomass and soil water potential. Soil Biol Biochem 2020; 143: 107742.

54. Seki K. SWRC fit - a nonlinear fitting program with a water retention curve for soils having

unimodal and bimodal pore structure. Hydrol Earth Syst Sci Discuss 2007; 4: 407-437.

55. Ray P, Ishiga T, Decker SR, Turner GB, Craven KD. A novel delivery system for the root

symbiotic fungus, *Sebacina vermifera*, and consequent biomass enhancement of low lignin

COMT switchgrass lines. BioEnerg Res 2015; 8: 922–933.

56. Blazewicz SJ, Schwartz E, Firestone MK. Growth and death of bacteria and fungi underlie

rainfall-induced carbon dioxide pulses from seasonally dried soil. Ecology 2014; 95: 1162-

1172.

57. Nuccio EE, Blazewicz SJ, Lafler M, Campbell AN, Kakouridis A, Kimbrel JA, et al. HT-

SIP: A semi-automated Stable Isotope Probing pipeline identifies interactions in the

hyphosphere of arbuscular mycorrhizal fungi. bioRxiv.

https://biorxiv.org/cgi/content/short/2022.07.01.498377v1

58. Buckley DH, Huangyutitham V, Hsu SF, Nelson TA. Stable isotope probing with ^{15}N

achieved by disentangling the effects of genome G+C content and isotope enrichment on

DNA density. Appl Environ Microbiol 2007; 73: 3189-3195.

59. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA

primers for marine microbiomes with mock communities, time series and global field

samples. Environ Microbiol 2016; 18: 1403-1414.

60. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R

gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb Ecol

2015; 75: 129-137.

61. Badri A, Stefani FOP, Lachance G, Roy-Arcand L, Beaudet D, Vialle A, Hijri M. Molecular

diagnostic toolkit for *Rhizophagus irregularis* isolate DAOM-197198 using quantitative PCR

assay targeting the mitochondrial genome. Mycorrhiza 2016; 26: 721–733.

62. Gamper HA, Young JP, Jones DL, Hodge A. Real-time PCR and microscopy: are the two

methods measuring the same unit of arbuscular mycorrhizal fungal abundance? Fungal Genet

Biol 2008; 45: 581-96.

63. Tellenbach C, Grünig CR, Sieber TN. Suitability of quantitative real-time PCR to estimate

the biomass of fungal root endophytes. Appl Environ Microbiol 2010; 76: 5764-5772.

64. Schildkraut CL, Marmur J, Doty P. Determination of the base composition of

deoxyribonucleic acid from its buoyant density in CsCl. J Mol Biol 1962; 4: 430–43.

65. Martin-Laurent F, Phillipot L, Hallet S, Chaussod R, Germon JC, Soulas G, et al. DNA

extraction from soils: old bias for new microbial diversity analysis methods. Appl Environ

Microbiol 2001; 67: 2354–2359.

66. Louca S, Doebeli M, Parfrey LW. Correcting for 16S rRNA gene copy numbers in

microbiome surveys remains an unsolved problem. Microbiome 2018; 6: 41.

67. Kanagawa T. Bias and artifacts in multitemplate polymerase chain reactions (PCR). J Biosci

Bioeng 2003; 96: 317–323.

68. Kozarewa I, Ning Z, Quail MA, Sanders MJ, Berriman M, Turner DJ. Amplification-free

Illumina sequencing-library preparation facilitates improved mapping and assembly of

(G+C)-biased genomes. Nat Methods 2009; 6: 291–295.

69. Manzoni S, Taylor P, Richter A, Porporato A, Ågren GI. Environmental and stoichiometric

controls on microbial carbon-use efficiency in soils. New Phytol 2012; 196: 79-91.

70. Geyer KM, Dijkstra P, Sinsabaugh R, Frey SD. Clarifying the interpretation of carbon use

efficiency in soil through methods comparison. Soil Biol Biochem 2019; 128: 79-88.

71. R Core Team. R: a language and environment for statistical computing. Vienna: R

Foundation for Statistical Computing; 2019.

72. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, OHara RB, et al. vegan:

Community Ecology Package R package version 2.3-0. 2015. Available from

http://CRAN.R-project.org/package=vegan.

73. Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: an effective distance

metric for microbial community comparison. ISME J 2011; 5: 169–172.

74. Harris RF. Effect of water potential on microbial growth and activity. In: Water Potential

Relations in Soil Microbiology. Eds. JF Parr, WR Gardner, LF Elliott. Madison, WI: Am Soc

Agron; 1981. p. 23–95.

75. Wagg C, Dudenhöffer JH, Widmer F, van der Heijden MGA. Linking diversity, synchrony

and stability in soil microbial communities. Funct Ecol 2018; 32: 1280–1292.

76. Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD. Defining trait-

based microbial strategies with consequences for soil carbon cycling under climate change.

ISME J 2020; 14: 1–9.

77. Tiemann LK, Billings SA. Changes in variability of soil moisture alter microbial community

C and N resource use. Soil Biol Biochem 2011; 43: 1837–1847.

78. Domeignoz-Horta LA, Pold G, Liu XJA, Frey SD, Melillo JM, DeAngelis KM. Microbial

diversity drives carbon use efficiency in a model soil. Nat Commun 2020; 11: 3684.

79. Gefen O, Balaban NQ. The importance of being persistent: heterogeneity of bacterial

populations under antibiotic stress. FEMS Microbiol Rev 2009; 33: 704–717.

80. Fridman O, Goldberg O, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time

underlies antibiotic tolerance in evolved bacterial populations. Nature 2014; 513: 418–421.

81. Bouskill NJ, Wood TE, Baran R, Hao Z, Ye Z, Bowen BP, et al. Belowground response to

drought in a tropical forest soil. II. Change in microbial function impacts carbon

composition. Front Microbiol 2016; 7: 323.

82. Sutcliffe IC. A phylum level perspective on bacterial cell envelope architecture. Trends

Microbiol 2010; 18: 464-470.

862 83. Tocheva EI, Ortega DR, Jensen GJ. Sporulation, bacterial cell envelopes and the origin of

863   life. Nat Rev Microbiol 2016; 14: 535–542.

865 84. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, et al. Drought delays

866   development of the sorghum root microbiome and enriches for monoderm bacteria. Proc Natl

867   Acad Sci USA 2018; 115: E4284-E4293.

869 85. Santos-Medellín C, Liechty Z, Edwards J, Nguyen B, Huang B, Weimer BC, Sundaresan V.

870   Prolonged drought imparts lasting compositional changes to the rice root microbiome. Nat

871   Plants 2021; 7: 1065–1077.

873 86. Otoguro M, Yamamura H, Quintana ET. The Family *Streptosporangiaceae*. In: The

874   Prokaryotes. Eds. Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F. Berlin,

875   Heidelberg: Springer; 2104. p. 1011-1045.

877 87. Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal communities

878   to extreme desiccation and rewetting. ISME J 2013; 7: 2229–2241.

880 88. Cruz AF, Ishii T. Arbuscular mycorrhizal fungal spores host bacteria that affect nutrient

881   biodynamics and biocontrol of soil-borne plant pathogens. Biol Open 2012; 1: 52-57.

89. Rillig MC, Lutgen ER, Ramsey PW, Klironomos JN, Gannon JE. Microbiota accompanying

different arbuscular mycorrhizal fungal isolates influence soil aggregation. Pedobiologia

2005; 49: 251–259.

90. Jiang F, Zhang L, Zhou J, George TS, Feng G. Arbuscular mycorrhizal fungi enhance

mineralisation of organic phosphorus by carrying bacteria along their extraradical hyphae.

New Phytol 2021; 230: 304-315.

91. Hernandez DJ, David AS, Menges ES, Searcy CA, Afkhami ME. Environmental stress

destabilizes microbial networks. ISME J 2021; 15: 1722–1734.

92. Leigh J, Fitter AH, Hodge A. Growth and symbiotic effectiveness of an arbuscular

mycorrhizal fungus in organic matter in competition with soil bacteria. FEMS Microbiol

Ecol 2011; 76: 428–438.

93. Leifheit EF, Verbruggen E, Rillig MC. Arbuscular mycorrhizal fungi reduce decomposition

of woody plant litter while increasing soil aggregation. Soil Biol Biochem 2015; 81: 323-

328.

94. Bronstein JL. Conditional outcomes in mutualistic interactions. Trends Ecol Evol 1994; 9:

214–217.

95. Toljander JF, Lindahl BD, Paul LR, Elfstrand M, Finlay RD. Influence of arbuscular

mycorrhizal mycelial exudates on soil bacterial growth and community structure. FEMS

Microbiol Ecol 2007; 61: 295–304.

96. Zhang L, Zhou J, George, TS, Limpens E, Feng G. Arbuscular mycorrhizal fungi conducting

the hyphosphere bacterial orchestra. Trends Plant Sci 2022; 27: 402-411.

97. Zhalnina K, Louie KB, Hao Z, Mansoori N, Nunes da Rocha U, Shi S, et al. Dynamic root

exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial

community assembly. Nat Microbiol 2018; 3: 470–480.

98. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. Root

exudation of phytochemicals in *Arabidopsis* follows specific patterns that are

developmentally programmed and correlate with soil microbial functions. PLoS One 2013; 8:

e55731.

99. Shi S, Richardson AE, O'Callaghan M, DeAngelis KM, Jones EE, Stewart A, et al. Effects of

selected root exudate components on soil bacterial communities. FEMS Microbiol Ecol

2011; 77: 600–610.

     Figure Legends

Figure 1. Experimental design, fungal inoculum abundance, and soil moisture. a Microcosm

design for quantitative $H_2^{18}O$ stable isotope probing (qSIP) assay of root-free hyphosphere soils.

P. hallii plants were inoculated with either *R. irregularis*, *S. bescii*, or left uninoculated

(indicated in yellow, blue, or black, respectively) and grown under water-replete or water-limited

conditions (indicated in dark or light shades and solid or dashed lines, respectively; n = 3

replicates per treatment). After three months, hyphosphere soils were amended with enriched

($H_2^{18}O$) or natural abundance ($H_2^{16}O$) water and incubated for seven days. **b** Abundance of *R.*

irregularis and (**c**) *S. bescii* (DNA copies g^{-1} soil) measured with strain-specific qPCR primers

after three months. **d** Soil moisture after three months. Bold lines represent median value;

whiskers represent upper and lower quartiles (n = 6 replicates per fungal*moisture treatment

combination). Asterisks denote the results of a nonparametric Kruskal-Wallis rank sum test

performed separately for each qPCR plot ($p < 0.001$). Letters denote a Tukey's HSD test for soil

moisture comparisons ($p < 0.001$).

Figure 2. Structure and alpha diversity of total and actively growing bacterial communities based on traditional and H$_2$^{18}O qSIP-filtered 16S rRNA gene profiling. Principal coordinates analysis (PCoA) of weighted UniFrac distances between bacterial communities assessed with (**a**) traditional 16S rRNA gene profiles and (**b**) H$_2$^{18}O qSIP-filtered 16S rRNA gene profiles representing the actively growing communities (i.e., ASVs that did not incorporate a significant quantity of ^{18}O were removed prior to analysis). Moisture history and fungal inoculum explained a total of 30% of the variation in total community structure ($p < 0.001$; n = 6 replicates) and 86% of variation in actively growing community structure ($p < 0.001$; n = 6 replicates). Ellipses show

treatment groupings in **(a)** and **(b)**. **c** Inverse Simpson's diversity index in total and actively

growing communities. Letters denote the results of a Tukey's HSD test (no significant

differences between total communities; $p < 0.01$ for comparisons between actively growing

communities; n = 6 replicates). Bold lines represent median values; whiskers represent upper and

lower quartiles. For all plots, uninoculated, *R. irregularis*-inoculated, and *S. bescii*-inoculated

soils are represented in black, yellow, and blue, respectively. Water-replete and water-limited

soils are represented in dark or light color shades and solid or dashed boxplot outlines,

respectively.

959
Figure 3. H$_2$^{18}O qSIP-based bacterial growth potential, gross growth, and CO$_2$ efflux from

hyphal ingrowth core soils following exposure to different moisture regimes and fungal

inocula. a Median ^{18}O atom percent excess (APE) of bacterial ASVs. Uninoculated, *R.*

irregularis-inoculated, and *S. bescii*-inoculated soils are represented in black, yellow, and blue,

respectively. Water-replete and water-limited soils are represented in dark or light shades and

solid or dashed outlines, respectively. Letters denote the results of a Tukey's HSD test

comparing the means of all treatments ($p < 0.01$; n = 3 replicates). **b** Taxon-specific gross growth

(16S rRNA gene copies g^{-1} soil) based on ASV ^{18}O incorporation, summed by phylum for each

treatment. **c** CO_2 efflux (μg CO_2 mg^{-1} soil C) from hyphal ingrowth core soils during $H_2^{18}O$ qSIP

assay. Bold lines represent median values; whiskers represent upper and lower quartiles ($p <$

0.05; n = 6 replicates per treatment).

Figure 4. Comparison of taxon-specific bacterial growth potential following water-limited versus water-replete conditions in uninoculated or fungal-inoculated hyphal ingrowth core soils. ^{18}O atom percent excess (APE) active growth ratios of ASVs present in both water-limited and water-replete soils that were either **(a)** uninoculated, **(b)** inoculated with *R. irregularis,* or **(c)** inoculated with *S. bescii*. Ratios were averaged at the phylum level. Ratios less than 1.0 (bars located to the left of the central vertical line) indicate that water limitation suppressed growth potential. Ratios greater than 1.0 (bars located to the right of the central line) indicate that water

979 limitation facilitated greater growth potential. Error bars represent the standard error. Asterisks

980 denote phylum-level averages that are significantly greater than or less than 1.0 (adjusted $p <$

981 0.05 based on Wilcoxon signed rank test and Benjamini-Hochberg correction for multiple

982 comparisons). Only taxa that incorporated a significant quantity of ^{18}O (lower 90% CI > 0) and

983 belong to the ten most abundant phyla are represented.

984
Figure 5. Comparison of taxon-specific bacterial growth potential in fungal-inoculated

versus uninoculated hyphal ingrowth core soils following water limitation. ^{18}O atom percent

excess (APE) active growth ratios of bacterial ASVs present in water-limited soils that were

either uninoculated or inoculated with *R. irregularis* or *S. bescii*, averaged at the phylum level

(a,b) or displayed by individual ASV grouped within class for the phyla *Acidobacteria* **(c)**,

Verrucomicrobia **(d)**, and *Actinobacteria* **(e)**. Ratios greater than 1.0 (to the right of the vertical

central line) indicate that fungal inocula supported greater bacterial growth potential. For phylum

level averages, asterisks represent ^{18}O APE ratios that are significantly greater than 1.0 (adjusted

$p < 0.05$ based on Wilcoxon signed rank test and Benjamini-Hochberg correction for multiple

comparisons). Error bars represent the standard error within each bacterial phylum. Only taxa

that incorporated a significant quantity of ^{18}O (lower 90% CI > 0) and belong to the ten most

abundant phyla are represented.