Chromosomal integration of *bla*CTX-M genes in diverse *Escherichia*

***coli* isolates recovered from river water in Japan**

Ryota Gomi[a*], Masaki Yamamoto[b], Michio Tanaka[b], Yasufumi Matsumura[b]

[a]Department of Environmental Engineering, Graduate School of Engineering, Kyoto

University, Katsura, Nishikyo-ku, 615-8540, Kyoto, Japan

[b]Department of Clinical Laboratory Medicine, Graduate School of Medicine, Kyoto

University, 54 Shogoin-kawahara-cho, Sakyo-ku, 606-8507, Kyoto, Japan

[*]Corresponding author

E-mail address: gomi.ryota.34v@kyoto-u.jp; Tel: +81-75-383-3354; Fax: +81-75-383-3358

Abstract

Occurrence of extended-spectrum β-lactamase (ESBL)-producing *Escherichia coli* (ESBLEC)

in environmental waters is of great concern. However, unlike clinical ESBLEC, their genetic

characteristics, in particular the genetic contexts of ESBL genes, are not well understood. In

this study, we sequenced and analyzed the genomes of CTX-M-producing *E. coli* isolates

recovered from river water to fully characterize the genetic contexts of bla_{CTX-M} genes. Among

the 14 isolates with completed genomes, bla_{CTX-M} genes were detected on the chromosome in

nine isolates. All but one chromosomal bla_{CTX-M} genes were associated with IS*Ecp1* and were

carried on different transposition units ranging in size from 2,855 bp to 11,093 bp; the

exception, $bla_{CTX-M-2}$, was associated with IS*CR1*. The remaining five isolates carried bla_{CTX-M}

genes on epidemic IncI1 plasmids of different sequence types (STs) (ST3, ST16, ST113, and

ST167) (n = 4) or on an IncB/O/K/Z plasmid (n = 1). This study revealed that environmental

E. coli carry bla_{CTX-M} genes in diverse genetic contexts. Apparent high prevalence of

chromosomal bla_{CTX-M} potentially indicates that some *E. coli* can stably maintain bla_{CTX-M} genes

in environmental waters, though further studies are needed to confirm this.

Keywords

Escherichia coli, bla_{CTX-M}, genome sequencing, antibiotic resistance, river water, one health

1. Introduction

Occurrence of antibiotic-resistant bacteria (ARB) in surface waters is a public health concern. Among ARB, extended-spectrum-β-lactamase (ESBL)-producing *Escherichia coli* (ESBLEC) represent a major threat to public health, because ESBLEC can be important causes of community-onset infections for which treatment options are limited (Pitout, 2010; Pitout and Laupland, 2008). ESBL enzymes are divided into several families (e.g., CTX-M and ESBL variants of TEM and SHV), among which CTX-M enzymes have been recognized as the most common ESBL type (Castanheira et al., 2021). Previous studies have reported the predominance of CTX-M ESBLs among environmental *E. coli*. For example, CTX-M-14(-like) was prevalent among ESBLEC in surface waters in China, Japan, and Korea, and CTX-M-15 was prevalent among ESBLEC in surface waters in France, Germany, and Switzerland (Falgenhauer et al., 2021; Girlich et al., 2020; Jang et al., 2013; Liu et al., 2018; Miyagi and Hirai, 2019; Zurfluh et al., 2013).

ESBL genes are often found on plasmids, which facilitates their horizontal spread (Castanheira et al., 2021). However, some previous studies also detected ESBL genes on chromosomes and suggested that integration of ESBL genes into chromosomes could contribute to the stabilization and maintenance of ESBL genes in the absence of antibiotic selection pressure (Hirai et al., 2013; Nair et al., 2021; Rodriguez et al., 2014). Thus, information on the location (chromosome or plasmid) of ESBL genes is important to predict the mobility and stability of

these genes. Although, previous studies have reported the occurrence of ESBLEC in

environmental waters, limited information exists regarding the locations and genetic contexts

of ESBL genes in environmental ESBLEC (Falgenhauer et al., 2021; Girlich et al., 2020; Liu

et al., 2018). Here, we completely sequenced the genomes of CTX-M-producing *E. coli* isolates

recovered from river water in Japan, aiming to elucidate the locations and genetic contexts of

the $bla_{\text{CTX-M}}$ genes in environmental ESBLEC. The genetic contexts of chromosomal $bla_{\text{CTX-M}}$

genes in the genomes of environmental *E. coli* retrieved from the NCBI Reference Sequence

Database (RefSeq) were also characterized.

2. Materials and methods

2.1. Bacterial isolates.

We previously collected 531 *E. coli* isolates from the Yamato River in Japan between 2011

and 2013 and detected 18 isolates with $bla_{\text{CTX-M}}$ genes (Gomi et al., 2014; Gomi et al., 2017).

Briefly, we collected a total of 27 water samples from 10 sites in the Yamato River in August

and November of 2011, February of 2012, and October of 2013 (note that water samples were

not collected from all sampling sites on each sampling occasion). The samples were processed

using the membrane filter method with XM-G agar (Nissui, Tokyo, Japan) to obtain *E. coli*

isolates. In total, 531 *E. coli* isolates were obtained, and a subset of isolates (n = 155) was

selected and subjected to Illumina short-read sequencing. Among the sequenced isolates, we

detected 18 isolates that carried bla_{CTX-M} genes. These 18 isolates were analyzed as described

below.

2.2. Antibiotic susceptibility testing.

Antibiotic susceptibility was assessed by microdilution using the dry plate Eiken assay (Eiken,

Tokyo, Japan) as described previously (Gomi et al., 2017) (see **Table S1** for the antimicrobial

agents used). The results were interpreted according to CLSI criteria (CLSI, 2020) and

EUCAST epidemiological cutoff (ECOFF) values (http://www.eucast.org). ESBL production

was confirmed following the CLSI guidelines (CLSI, 2020). A detailed description of antibiotic

susceptibility testing is provided in the Supplementary Materials and methods.

2.3. Genome sequencing and assembly.

DNA was extracted from each isolate using a DNeasy PowerSoil Pro Kit (Qiagen, Hilden,

Germany). The short-read sequencing library was prepared using the Illumina DNA Prep

(Illumina, San Diego, CA), and the library was sequenced on an Illumina NovaSeq 6000 system

with 250-bp paired-end sequencing (although we obtained Illumina reads for the isolates in our

previous study (Gomi et al., 2017), the Illumina reads had low depth/uneven coverage and

hence we sequenced the isolates again with Illumina technology in the present study). The

long-read sequencing library was prepared using the SQK-LSK109 kit (Oxford Nanopore

Technologies, Oxford, UK), and the prepared library was sequenced on the MinION with a

FLO-MIN106 flow cell.

Illumina short reads were subsampled using seqtk (v1.3, https://github.com/lh3/seqtk), and the

subsampled reads were trimmed with fastp (v0.20.0) (Chen et al., 2018). Oxford Nanopore

Technologies (ONT) long reads were filtered with Filtlong (v0.2.0,

https://github.com/rrwick/Filtlong) to remove reads with low quality or short length. Hybrid

assembly of Illumina short reads and ONT long reads was performed by employing a short-

read-first approach, i.e., Unicycler (v0.4.8) with the --no_correct option (Wick et al., 2017).

Hybrid assembly was also performed using a long-read-first approach, i.e., long-read assembly

by Flye (v2.9) (Kolmogorov et al., 2019) followed by long-read polishing using Medaka

(v1.4.4, https://github.com/nanoporetech/medaka) and short-read polishing using Polypolish

(v0.4.3) (Wick and Holt, 2022). Assembly graphs of both hybrid assemblies were visualized

using Bandage (v0.8.1) for each isolate (Wick et al., 2015), and a better assembly was chosen

as the final assembly.

2.4. Genomic analysis.

Genomes were annotated using the RAST server (Aziz et al., 2008), ISfinder (Siguier et al.,

2006), ResFinder 4.1 (Bortolaia et al., 2020), and the BLAST algorithm

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). Plasmid replicons were detected using

PlasmidFinder 2.1 (Carattoli et al., 2014). Relaxase types, mate-pair formation types, and

transferability of plasmids were determined using MOB-typer (Robertson and Nash, 2018).

IncI1 plasmids were annotated using plasmids R64, ColIb-P9, and R621a as references

(Carattoli et al., 2021).

2.5. Characterization of environmental *E. coli* genomes with *bla*$_{CTX-M}$ genes in RefSeq.

RefSeq *E. coli* genomes with an assembly level of "complete genome" (n = 1903) were

downloaded in January 2022. Metadata information, such as collection date and isolation

source, was extracted from the GenBank files using an in-house Python script, and genomes of

E. coli from environmental waters were retrieved (n = 53). Antibiotic resistance genes were

detected using ABRicate (https://github.com/tseemann/abricate), and genomes carrying *bla*$_{CTX-}$

$_M$ on the chromosome were analyzed as described above.

2.6. Accession numbers.

The complete genomes and sequence read files were deposited in GenBank and the NCBI SRA

under BioProject PRJNA800231 (also see **Table S2** for the accession number of each replicon

for the completed genomes and **Table S1** for SRA accession numbers of the remaining

genomes).

3. Results and discussion

3.1. Basic characteristics of *E. coli* isolates sequenced in this study.

All 18 *bla*$_{CTX-M}$-carrying isolates characterized in the present study were non-susceptible/non-

wild type to ampicillin, aztreonam, cefazolin, cefepime, cefotaxime, and cefpodoxime, while

all were susceptible/wild type to amoxicillin-clavulanic acid, piperacillin-tazobactam,

imipenem, meropenem, amikacin, and colistin. Non-susceptibility/non-wild type rates for the

remaining antibiotics ranged from 6% (fosfomycin) to 89% (ampicillin-sulbactam) (**Table S1**).

ESBL production was confirmed in all the isolates. The 18 isolates were highly diverse,

comprising 16 different sequence types (STs), namely ST38 (n = 2), ST69 (n = 2), and a single

isolate from each of ST10, ST46, ST117, ST216, ST354, ST457, ST744, ST1193, ST2003,

ST3107, ST6214, ST6215, ST6216, and ST6220 (**Table S1**). The bla_{CTX-M} genes detected were

also diverse, namely $bla_{CTX-M-14}$ (n = 8), $bla_{CTX-M-2}$ (n = 2), $bla_{CTX-M-15}$ (n = 2), $bla_{CTX-M-24}$ (n = 2),

$bla_{CTX-M-55}$ (n = 2), $bla_{CTX-M-1}$ (n = 1), $bla_{CTX-M-3}$ (n = 1), and $bla_{CTX-M-8}$ (n = 1). One isolate

(KTa008) carried two copies of the $bla_{CTX-M-14}$ gene. In Japan, ESBLEC are frequently detected

among *E. coli* infections. For example, 22% of extraintestinal pathogenic *E. coli* isolates were

positive for one or more ESBL genes in a study conducted at 10 acute-care hospitals

(Matsumura et al., 2017a). Like in other countries, these ESBLEC infections in Japan have

been shown to be mainly due to CTX-M-producing ST131 (Matsumura et al., 2017b). However,

the ESBLEC isolates analyzed in the present study belonged to a wide variety of STs and none

belonged to ST131, which indicates that the clonal composition of ESBLEC between

environmental settings and clinical settings may be different in Japan.

Of the 18 isolates, the genomes of 14 could be completed. These genomes carried 0 to 8

plasmids (see **Table S2** for details). The genomes of the remaining four isolates could not be

completed, probably due to the presence of long repeats in the genomes. The location and

genetic contexts of the bla_{CTX-M} genes were determined for the 14 isolates with completed

genomes. Of these 14, nine carried bla_{CTX-M} genes on chromosomes and five carried bla_{CTX-M}

genes on plasmids.

3.2. Genetic contexts of chromosomal bla_{CTX-M} genes.

In eight of the nine isolates which carried bla_{CTX-M} genes on chromosomes, the bla_{CTX-M} genes

were associated with IS$Ecp1$ and carried on transposition units (TPU) of different sizes (2,855

bp to 11,093 bp) (**Figure 1**). These TPU were flanked by AT-rich 5-bp target site duplications

(TSD). None of the TPU were identical, and the insertion sites of the TPU were different in all

cases, indicating that different transposition events contributed to the chromosomal integration

of each TPU. Notably, the TPU in KFu023 contained multiple resistance genes, including

$bla_{CTX-M-15}$, $aac(6')$-Ib-cr, bla_{OXA-1}, and $\Delta catB3$. This seems to highlight the ability of IS$Ecp1$ to

capture sequences of different lengths in different transposition events and thus simultaneously

mobilize antibiotic resistance genes of different origins (Partridge et al., 2018). In most cases,

these TPU were directly inserted into chromosomes. However, in KKa019, the TPU was part

of a 35,625 bp multiresistance region bracketed by two copies of IS26 (annotated as IS$15DI$ in

the ISFinder database), and the whole multiresistance region was flanked by 8-bp TSD

(CGTTGCCG) in the chromosome (**Figure 2a**). KTa008 carried two copies of $bla_{CTX-M-14}$,

which were separated by ~1,169 kbp in the chromosome. Analysis of the genetic contexts of

these $bla_{CTX-M-14}$ genes indicated transposition of an already chromosomally-located TPU and

an adjacent region into another chromosomal location (**Figure S1**). This type of 'recurrent'

transposition was previously reported in clinical ESBLEC, though the TPU and insertion sites

were different (Hamamoto and Hirai, 2019).

The above results indicate that analysis of sequences immediately upstream and downstream

of bla_{CTX-M} (e.g., identification of IS$Ecp1$ upstream and $orf477$ downstream of bla_{CTX-M}) is not

sufficient to track the movement and evolution of bla_{CTX-M}, since TPU can be very long

(sometimes > 10,000 bp) or can be inserted into a large multiresistance region. This can be

overcome, for example, by determining the complete genomes as shown in the present study.

We searched for *E. coli* genomes in the public database which carry the same TPU in the same

chromosomal location as those identified in the present study. BLASTN searches were

performed against the nucleotide collection (nr/nt) database in January 2022 using the TPU and

surrounding sequences identified in the present study as query sequences. Three of the nine

TPU/insertion site combinations were found in the database, while the remaining six were not

found, potentially corresponding to novel integration events (**Table S3**).

In KOr019, $bla_{CTX-M-2}$ was associated with IS$CR1$ and situated within a complex class 1 integron,

which is congruent with previous studies (Canton et al., 2012; Partridge et al., 2018). This

complex class 1 integron was closely related to a recently reported transposon Tn7337 (**Figure**

2b) (Coppola et al., 2022). KOr019 carried the IRt (inverted repeat at *tni* end)-IS6100-IRt

structure, but this structure was truncated by IS*26* in Tn*7337*. The 17,747 bp structure from IRi

(inverted repeat at *intI*1 end) to IRt was inserted into the chromosome of KOr019, creating 5-

bp TSD of AAATG. This insertion seems to have been mediated by Tni proteins provided *in*

trans, since the *tni* genes in the complex class 1 integron were absent (Partridge et al., 2018).

3.3. Genetic contexts of *bla*$_{CTX-M}$ genes on plasmids.

The five completed plasmids with *bla*$_{CTX-M}$ genes belonged to IncI1 (n = 4) and IncB/O/K/Z (n

= 1) (**Figure 3a**). IncI1 plasmids were typed as four different STs, namely ST3, ST16, ST113,

and ST167, by IncI1 pMLST (Carattoli et al., 2014). TPU were inserted into plasmid backbones

in the IncI1/ST3, IncI1/ST16, and IncI1/ST167 plasmids. Interestingly, the IncI1/ST16 and

IncI1/ST167 plasmids carried the same TPU (except for insertion of IS*1* in the IncI1/ST16

plasmid) inserted into the same location. IncI1/ST167 is a single-locus variant of IncI1/ST16,

which indicates the TPU was inserted into the plasmid backbone before the ST diversification.

In the IncI1/ST113 plasmid, an IS26-ΔIS*10R*-*bla*$_{CTX-M-8}$-IS*26* structure was inserted into the

plasmid. BLASTN analysis identified IncI1 plasmids belonging to the same IncI1-STs and

carrying the same TPU/IS*26* structures inserted into the same locations from different countries

(e.g., MH847453 from France for *bla*$_{CTX-M-1}$-carrrying IncI1/ST3, CP056650 from the UK for

bla$_{CTX-M-55}$-carrrying IncI1/ST16, KY964068 from Portugal for *bla*$_{CTX-M-8}$-carrrying

IncI1/ST113, CP021208 from China for *bla*$_{CTX-M-55}$-carrrying IncI1/ST167), which indicates

circulation of epidemic CTX-M IncI1 plasmids in geographically distant countries.

Comparison of IncI1 plasmids revealed the replication region, the leading region, and the

transfer regions in the IncI1 reference plasmid R64 were well conserved in all IncI1 plasmids

detected in this study, while many genes in the stability region were lacking (**Figure 3b**). In

the IncB/O/K/Z plasmid, $bla_{CTX-M-14}$ was linked to ΔTn*1721* and a class 1 integron. Similar

genetic contexts were previously reported (e.g., JF701188).

3.4. Chromosomal location of bla_{CTX-M} in the RefSeq assemblies.

Among the 53 RefSeq environmental *E. coli* genomes, 20 carried bla_{CTX-M}. Sixteen carried

bla_{CTX-M} on plasmids and four carried bla_{CTX-M} on chromosomes (**Table S4**). None of the 20

RefSeq genomes with bla_{CTX-M} belonged to ST131, again indicating a potential difference in

clonal composition of ESBLEC in environmental settings and clinical settings. The four

genomes with chromosomal bla_{CTX-M} belonged to ST38 (n = 2), ST130 (n = 1) and ST648 (n =

1) and originated from various countries (Japan, New Zealand, Norway, and Switzerland),

which indicates that diverse *E. coli* isolates carrying chromosomal bla_{CTX-M} are also present in

environmental waters in other countries. In all four genomes, bla_{CTX-M} was associated with TPU

ranging from 2,971 bp to 13,552 bp. A TPU containing multiple resistance genes besides

bla_{CTX-M} and a TPU inserted into a multiresistance region were detected, again highlighting the

importance of these mechanisms in the emergence of multidrug resistant *E. coli* in the

environment (see **Figure S2** for the analysis of the genetic contexts of bla_{CTX-M} in these

genomes).

ST38, which was detected in two RefSeq genomes containing chromosomal $bla_{\text{CTX-M}}$, was also

detected in two isolates with chromosomal $bla_{\text{CTX-M}}$ sequenced in this study. Interestingly, ST38

was also found to be prevalent in some previous studies reporting chromosomal $bla_{\text{CTX-M}}$

(Guenther et al., 2017; Guiral et al., 2011; Rodriguez et al., 2014). The reason for the

prevalence of ST38 among *E. coli* with chromosomal $bla_{\text{CTX-M}}$ is unknown. However, multiple

integration events seem to have been involved in the acquisition of chromosomal $bla_{\text{CTX-M}}$ in

ST38 (see **Figure 1**, **Figure 2a**, and **Figure S2**), which potentially indicates ST38 has genetic

mechanisms that can promote chromosomal integration of $bla_{\text{CTX-M}}$.

3.5. Study limitations.

This study has some limitations. The *E. coli* isolates analyzed in the present study were

collected between 2011 and 2013 (i.e., ~10 years ago). Additionally, a limited number of

complete genomes were determined (n = 14), and the isolates were collected regionally, not

nationwide. However, we also included RefSeq environmental *E. coli* genomes from various

countries with various collection dates in our analysis, which might mitigate these limitations.

4. Conclusions.

Here, we demonstrated that the genetic contexts of $bla_{\text{CTX-M}}$ genes in environmental *E. coli* are

highly diverse in terms of the associated mobile elements (various TPU, IS*CR1*, and IS*26*) and

integration sites. This study also revealed that the chromosomal locations of $bla_{\text{CTX-M}}$ genes

were unexpectedly frequent in environmental ESBLEC in Japan. Chromosomal integration of

bla$_{\text{CTX-M}}$ might allow *E. coli* to stably maintain *bla*$_{\text{CTX-M}}$ in environmental waters, though further

studies are needed to confirm this.

Acknowledgements

We acknowledge the NGS core facility of the Genome Information Research Center at the

Research Institute for Microbial Diseases of Osaka University for the support in DNA

sequencing. Computations were partially performed on the NIG supercomputer at ROIS

National Institute of Genetics. This work was supported by the Environment Research and

Technology Development Fund (JPMEERF20205006) of the Ministry of the Environment,

Japan. This study was also supported in-part by the Program for the Development of Next-

generation Leading Scientists with Global Insight (L-INSIGHT), sponsored by the Ministry of

Education, Culture, Sports, Science and Technology (MEXT), Japan.

References

Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid

annotations using subsystems technology. BMC Genomics 2008; 9: 75.

Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0 for

predictions of phenotypes from genotypes. J Antimicrob Chemother 2020; 75: 3491-

3500.

Canton R, Gonzalez-Alba JM, Galan JC. CTX-M Enzymes: Origin and Diffusion. Front

Microbiol 2012; 3: 110.

Carattoli A, Villa L, Fortini D, Garcia-Fernandez A. Contemporary IncI1 plasmids involved in

the transmission and spread of antimicrobial resistance in Enterobacteriaceae. Plasmid

2021; 118: 102392.

Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, et al. In silico

detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence

typing. Antimicrob Agents Chemother 2014; 58: 3895-903.

Castanheira M, Simner PJ, Bradford PA. Extended-spectrum beta-lactamases: an update on

their characteristics, epidemiology and detection. JAC Antimicrob Resist 2021; 3:

dlab092.

Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics 2018; 34: i884-i890.

CLSI. *Performance Standards for Antimicrobial Susceptibility Testing—Thirtieth Edition:*

M100.   2020.

Coppola N, Cordeiro NF, Trenchi G, Esposito F, Fuga B, Fuentes-Castillo D, et al. Imported

One-Day-Old Chicks as Trojan Horses for Multidrug-Resistant Priority Pathogens

Harboring *mcr-9*, *rmtG*, and Extended-Spectrum beta-Lactamase Genes. Appl Environ

Microbiol 2022; 88: e0167521.

Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene gain,

loss and rearrangement. PLoS One 2010; 5: e11147.

Falgenhauer L, Zur Nieden A, Harpel S, Falgenhauer J, Domann E. Clonal CTX-M-15-

Producing *Escherichia coli* ST-949 Are Present in German Surface Water. Front

Microbiol 2021; 12: 617349.

Girlich D, Bonnin RA, Naas T. Occurrence and Diversity of CTX-M-Producing *Escherichia*

coli From the Seine River. Front Microbiol 2020; 11: 603578.

Gomi R, Matsuda T, Matsui Y, Yoneda M. Fecal source tracking in water by next-generation

sequencing technologies using host-specific *Escherichia coli* genetic markers. Environ

Sci Technol 2014; 48: 9616-23.

Gomi R, Matsuda T, Matsumura Y, Yamamoto M, Tanaka M, Ichiyama S, et al. Whole-

Genome Analysis of Antimicrobial-Resistant and Extraintestinal Pathogenic

Escherichia coli in River Water. Appl Environ Microbiol 2017; 83: e02703-16.

Guenther S, Semmler T, Stubbe A, Stubbe M, Wieler LH, Schaufler K. Chromosomally

encoded ESBL genes in *Escherichia coli* of ST38 from Mongolian wild birds. J

Antimicrob Chemother 2017; 72: 1310-1313.

Guiral E, Mendez-Arancibia E, Soto SM, Salvador P, Fabrega A, Gascon J, et al. CTX-M-15-

producing enteroaggregative *Escherichia coli* as cause of travelers' diarrhea. Emerg

Infect Dis 2011; 17: 1950-3.

Hamamoto K, Hirai I. Characterisation of chromosomally-located bla_{CTX-M} and its surrounding

sequence in CTX-M-type extended-spectrum beta-lactamase-producing *Escherichia*

coli isolates. J Glob Antimicrob Resist 2019; 17: 53-57.

Hirai I, Fukui N, Taguchi M, Yamauchi K, Nakamura T, Okano S, et al. Detection of

chromosomal $bla_{CTX-M-15}$ in *Escherichia coli* O25b-B2-ST131 isolates from the Kinki

region of Japan. Int J Antimicrob Agents 2013; 42: 500-6.

Jang J, Suh YS, Di DY, Unno T, Sadowsky MJ, Hur HG. Pathogenic *Escherichia coli* strains

producing extended-spectrum beta-lactamases in the Yeongsan River basin of South

Korea. Environ Sci Technol 2013; 47: 1128-36.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat

graphs. Nat Biotechnol 2019; 37: 540-546.

Liu H, Zhou H, Li Q, Peng Q, Zhao Q, Wang J, et al. Molecular characteristics of extended-

spectrum beta-lactamase-producing *Escherichia coli* isolated from the rivers and lakes

in Northwest China. BMC Microbiol 2018; 18: 125.

Matsumura Y, Noguchi T, Tanaka M, Kanahashi T, Yamamoto M, Nagao M, et al. Population

structure of Japanese extraintestinal pathogenic *Escherichia coli* and its relationship

with antimicrobial resistance. J Antimicrob Chemother 2017a; 72: 1040-1049.

Matsumura Y, Pitout JDD, Peirano G, DeVinney R, Noguchi T, Yamamoto M, et al. Rapid

Identification of Different *Escherichia coli* Sequence Type 131 Clades. Antimicrob

Agents Chemother 2017b; 61.

Miyagi K, Hirai I. A survey of extended-spectrum beta-lactamase-producing

Enterobacteriaceae in environmental water in Okinawa Prefecture of Japan and

relationship with indicator organisms. Environ Sci Pollut Res Int 2019; 26: 7697-7710.

Nair S, Chattaway M, Langridge GC, Gentle A, Day M, Ainsworth EV, et al. ESBL-producing

strains isolated from imported cases of enteric fever in England and Wales reveal

multiple chromosomal integrations of $bla_{CTX-M-15}$ in XDR *Salmonella* Typhi. J

Antimicrob Chemother 2021; 76: 1459-1466.

Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile Genetic Elements Associated with

Antimicrobial Resistance. Clin Microbiol Rev 2018; 31.

Pitout JD. Infections with extended-spectrum beta-lactamase-producing enterobacteriaceae:

changing epidemiology and drug treatment choices. Drugs 2010; 70: 313-33.

Pitout JD, Laupland KB. Extended-spectrum beta-lactamase-producing Enterobacteriaceae: an

emerging public-health concern. Lancet Infect Dis 2008; 8: 159-66.

Robertson J, Nash JHE. MOB-suite: software tools for clustering, reconstruction and typing of

plasmids from draft assemblies. Microb Genom 2018; 4: e000206.

Rodriguez I, Thomas K, Van Essen A, Schink AK, Day M, Chattaway M, et al. Chromosomal

location of bla_{CTX-M} genes in clinical isolates of *Escherichia coli* from Germany, The

Netherlands and the UK. Int J Antimicrob Agents 2014; 43: 553-7.

Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre for

bacterial insertion sequences. Nucleic Acids Res 2006; 34: D32-6.

Wick RR, Holt KE. Polypolish: Short-read polishing of long-read bacterial genome assemblies.

PLoS Comput Biol 2022; 18: e1009802.

Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies

from short and long sequencing reads. PLoS Comput Biol 2017; 13: e1005595.

Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo genome

assemblies. Bioinformatics 2015; 31: 3350-2.

Zurfluh K, Hachler H, Nuesch-Inderbinen M, Stephan R. Characteristics of extended-spectrum

346   beta-lactamase- and carbapenemase-producing Enterobacteriaceae Isolates from rivers

347   and lakes in Switzerland. Appl Environ Microbiol 2013; 79: 3021-6.

 FIGURES

352 **Figure 1**. Genetic contexts of chromosomal bla_{CTX-M} genes. Red arrows indicate antibiotic

353 resistance genes, gray arrows indicate mobile elements, and white arrows indicate other

354 genes. The blue shaded box indicates the regions with 100% nucleotide identity. The entire

355 resistance regions of KKa019 and KOr019 are shown in **Figure 2**. 5-bp TSD are shown next

356 to each TPU. hp: hypothetical protein.

Figure 2. (a) A multiresistance region on the chromosome of KKa019. Insertion of the

resistance region into *flhA* generated 8-bp TSD of CGTTGCCG. A BLASTN search against

the nucleotide collection (nr/nt) database using the 35,625 bp multiresistance region as a

query sequence identified some related structures, but none showed >90% coverage. (b) A

multiresistance region on the chromosome of KOr019. Tn*7337* is shown for the purpose of

comparison. The 17,747 bp structure from IRi to IRt was inserted into an intergenic region in

the KOr019 chromosome, generating 5-bp TSD of AAATG. The light blue shaded box

indicates regions with >99% nucleotide identity.

Figure 3. (a) Genetic contexts of bla_{CTX-M} genes on plasmids. (b) Mauve comparison of IncI1

plasmids (Darling et al., 2010). IncI1 reference plasmid R64 is shown at the top. Four major

regions in R64 (replication, stability, leading, and conjugative transfer) are indicated.

Homologous segments are shown as colored blocks. Locations of bla_{CTX-M} genes are indicated

with arrows.