Transcriptional circuitry of NKX2-1 and SOX1 defines an unrecognized lineage subtype of small cell lung cancer

Ranran Kong[1,2,3], Ayushi S. Patel[2,3,4], Takashi Sato[2,3,5], Seungyeul Yoo[6,7], Li Bao[8], Abhilasha Sinha[2,3], Feng Jiang[2,3], Yang Tian[2,3], Maya Fridrikh[2,3], Shuhui Liu[9], Jie Feng[10], Xijing He[11,12], Jiantao Jiang[1], Yuefeng Ma[1], Karina Grullon[2,3], Dawei Yang[2,3,13], Charles A. Powell[2,3], Mary Beth Beasley[14], Jun Zhu[3,6,7], Eric L. Snyder[15,16], Shaomin Li[1]*, and Hideo Watanabe[2,3,6]*

[1]Department of Thoracic Surgery, The Second Affiliated Hospital of Medical School, Xi'an Jiaotong University, Xi'an, Shaanxi, 710004, China;

[2]Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[3]Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[4]Division of Hematology & Medical Oncology, Laura and Isaac Perlmutter Cancer Center, New York University Langone Medical Center, New York, NY, USA;

[5]Department of Respiratory Medicine, Kitasato University School of Medicine, Sagamihara, Japan; Division of Pulmonary Medicine, Department of Medicine, Keio University School of Medicine, Tokyo, Japan;

[6]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[7]Sema4, a Mount Sinai venture, Stamford, CT 06902, USA;

[8]Ningxia People's Hospital, Yinchuan, Ningxia Province, 750001, China;

[9]Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[10]Department of Nephrology, The First Affiliated Hospital of Medical School, Xi'an Jiaotong University, Xi'an, Shaanxi, 710061, China;

[11]Department of Orthopedics, The Second Affiliated Hospital of Medical School, Xi'an Jiaotong University,

Xi'an, Shaanxi, 710004, China;

[12]Xi'an International Medical Center, Xi'an, Shaanxi, 710126, China;

[13]Department of Pulmonary and Critical Care Medicine, Zhongshan Hospital Fudan University, Shanghai,

200032, China;

[14]Department of Pathology and Laboratory Medicine, Icahn School of Medicine at Mount Sinai, New York,

NY 10029, USA;

[15]Department of Pathology, University of Utah, Salt Lake City, UT 84112; Department of Oncological

Sciences, University of Utah, Salt Lake City, UT 84112, USA;

[16]Huntsman Cancer Institute, University of Utah, Salt Lake City, UT 84112

* Both authors share senior authorship.

H.W. has been supported by ATS Foundation Unrestricted Grant (ATS-2017-24), the American

Lung Association of the Northeast Lung Cancer Discovery Award (LCD-504985), Department of

Defense (W81XWH-19-1-0613) and NIH (R01CA240342). R.K. is supported by Shaanxi

Provincial Natural Science Foundation, China (2017JM8046). T. S. is supported by the Japanese

Respiratory Society the 6th Lilly Oncology Fellowship Program and the Uehara Memorial

Foundation. J. F. is supported by National Natural Science Foundation of China (NSFC) (No.

82100718). D. Y. is supported by Shanghai Pujiang Program (20PJ1402400). C.A.P. and H. W.

are supported by NIH (R01CA163772). E.L.S. is supported by the NIH (R01CA212415,

R01CA240317) and the American Lung Association Lung Cancer Discovery Award (LCD-

821670). This work was supported in part through the computational resources and staff expertise

provided by Scientific Computing at the Icahn School of Medicine at Mount Sinai.

Correspondence and requests for reprints should be addressed to Hideo Watanabe, Division of

Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, Icahn School of Medicine

at Mount Sinai, New York, NY 10029, USA. Phone: 212-241-9242; Fax: 212-876-5519; Email:

hideo.watanabe@mssm.edu.

Author contributions: Conceptualization & Methodology, R.K., and H.W.; Data Collection: R.K.,

A.S.P., T.S., S.Y., L.B., F.J., M.F., S.L., J.F., X.H., J.J., Y.M., K.G., D.Y., E.L.S., S.L., L.S., and

H.W.; Resources, E.L.S., M.B.B., C.A.P., S.L., and H.W.; Data Analysis, R.K., S.Y., A.S.P., S.L.,

M.B.B., E.L.S., and H.W.; Writing – Original Draft, R.K., A.S.P. and H.W.; Writing-Review &

Editing, R.K., A.S.P., E.L.S., and H.W.; Supervision, C.A.P., J.Z. S.L., and H.W.; Project

Administration & Funding Acquisition, H.W.

Running title: Collaboration of NKX2-1 and SOX1 defines SCLC-Aα subtype

Descriptor number: 9.26 Lung Cancer: Cellular and Molecular Aspects

The total word count for the body of the manuscript: 3,554

This article has an online data supplement, which is accessible from this issue's table of content

online at www.atsjournals.org

Abstract:

Rationale: The current molecular classification of small cell lung cancer (SCLC) based on

expression of four lineage transcription factors still leaves its major subtype SCLC-A as a

heterogeneous group, necessitating more precise characterization of lineage subclasses.

Objectives: To refine the current SCLC classification with epigenomic profiles and to identify

features of the re-defined SCLC subtypes.

Methods: We performed unsupervised clustering of epigenomic profiles on 25 SCLC cell lines.

Functional significance was evaluated by cell growth, apoptosis and xenograft using CRISPR-

Cas9-mediated deletion. The specific cistromic profiles by ChIP-seq and its functional

transcriptional partners using co-immunoprecipitation followed by mass spectrometry were

determined. $Rb1^{fl/fl}Trp53^{fl/fl}$ and $Rb1^{fl/fl}Trp53^{fl/fl}Nkx2\text{-}1^{fl/fl}$ mouse models were engineered to

explore the function of *Nkx2-1* in tumor initiation and differentiation. H3K27ac profiles were

analyzed to reveal 6 human SCLC specimen and 20 mice tumors epigenomic landscapes.

Measurements and Main Results: We identified an epigenomic subclusters of the major SCLC-

A subtype, named SCLC-Aα and SCLC-Aσ. SCLC-Aα was characterized by the presence of a

super-enhancer at the *NKX2-1* locus, which was observed in human SCLC specimens and a murine

SCLC model. We found NKX2-1, a dual lung and neural lineage factor, is uniquely relevant in

SCLC-Aα. We further found maintenance of this neural identity in SCLC-Aα is mediated by

collaborative transcriptional activity with another neuronal transcriptional factor SOX1.

Conclusions: We comprehensively describe an additional epigenomic heterogeneity of the major

SCLC-A subtype, and define SCLC-Aα subtype by the core regulatory circuitry representing

NKX2-1 and SOX1 super-enhancers and their functional collaborations to maintain neuronal

linage state. (Word count 250)

Keywords: small cell lung cancer (SCLC); *NKX2-1/TTF-1*; *SOX1*; Super-enhancer

INTRODUCTION

Small cell lung cancer (SCLC) is an aggressive, lethal, and rapidly progressing neuroendocrine (NE) malignancy, accounting for ~15% of lung cancer incidence and 250,000 deaths worldwide annually (1). The five-year survival rate for patients with SCLC is only 6% compared to 19% for non-SCLC. The lack of progress in SCLC treatment is largely attributable to the inadequate molecular understanding. Advances in genomic technology have enabled new molecular characterization of SCLCs. Genome-wide methylation and transcriptomes of human SCLCs and PDX models identified NE and non-NE subtypes. NE subtype is defined by the expression of neuronal lineage transcription factors (TFs) ASCL1 and/or NeuroD1 (2). Two recent studies described the non-NE SCLCs, one signified by YAP1 expression and another by POU2F3 expression (3, 4). In light of these findings, a consortium proposed a new molecular classification of SCLCs that distinguishes them into four subtypes based on the expression of four lineage-defining transcription factors: ASCL1, NeuroD1, YAP1 and POU2F3 referred to as SCLC-A, SCLC-N, SCLC-Y and SCLC-P, respectively (5, 6). The classification, while marking significant advance, is still a work in progress. After that report, a newer study proposed an inflamed SCLC subtype (SCLC-I) (7). In addition, not all SCLCs clearly fall into the four clusters as they often express multiple factors at variable levels, especially in the SCLC-A subtype which account for 70% of SCLCs (8). This wide range of heterogeneity warrants novel approaches to develop a more precise classification of SCLC based on lineage subtype.

Lineage-specific transcription factors and the presence of SE at their gene locus are critical determinants of cellular identity (9). In our study, we applied epigenomic approaches to investigate the differential enhancer landscapes in SCLC cell lines, human SCLC samples and autochthonous murine SCLCs from a genetic engineered mouse model (GEMM) to reveal additional intertumoral

heterogeneity, providing comprehensive evidence of distinct histone modification profiles.

Furthermore, we determined a novel mechanism of subtype-specific transcriptional regulation in

SCLC, integrating the core transcriptional networks and their collaborative regulation of NE target

genes.

METHODS

For a detailed description of the methods, please see the online supplement.

Cell lines

Cell lines used in this study are listed in Table E1.

Chromatin Immunoprecipitation followed by sequencing (ChIP-seq)

ChIP-seq was performed as described previously (10). Protein G magnetic beads (Dynabeads: Life

Technologies) were pre-incubated with anti-H3K27ac (Abcam, ab4729) or anti-NKX2-1 (Bethyl

Laboratories, A300-BL4000) or anti-SOX1 (R&D, AF3369) antibodies. Up to 10 ng of DNA was

used for the library construction using NEBNext Ultra II DNA Library Prep Kit (NEB, E7645).

Sequencing was performed on NextSeq 500 (Illumina).

RNA-seq

The procedure was performed as previously described (8, 10).

Lentiviral transduction of genes and CRISPR-Cas9 genome editing

FLAG-NKX2-1 or *FLAG-GFP* open reading frame (ORF) was cloned into pLEX_306 (a gift from

David Root, Addgene plasmid #41391). Cells stably expressing Cas9 were generated by infection

with the lentiCas9-Blast plasmid (Addgene # 52962, a gift from Feng Zhang). sgRNAs targeting

NKX2-1 or *SOX1* were selected from Brunello library (Table E2) (11). Additional details are

provided in the online supplement.

Immunohistochemistry (IHC)

Immunohistochemical analyses were performed on xenograft tumor specimens, human primary

SCLC tumor specimens and genetic engineered mouse model (GEMM) tumor specimens with

anti-NKX2-1 (1:100; Santa Cruz, sc-53136), anti-SOX1 (R&D, AF3369) or anti-ASCL1 (1:100,

BD, 556604) antibody and detected with the Vectastain ABC kit.

Co-immunoprecipitation and LC-MS/MS

Chromatin fraction was prepared and co-immunoprecipitated with the methods described

previously (12) with modifications. Peptides were detected, isolated, and fragmented to produce a

tandem mass-spectrum of specific fragment ions for each peptide with LTQ Orbitrap Velos Pro

ion-trap mass spectrometer (Thermo Fisher Scientific, Waltham, MA).

Cell proliferation assay

AlamarBlue Cell Viability Reagent was used to analyze cell proliferation. Cells were plated at a

density of 3,000 cells/well with five replicates in a 96-well plate.

Xenograft model

The procedure was performed as previously described (ref). All xenograft studies were approved

by the institutional Animal Care and Use Committee at Icahn School of Medicine at Mount Sinai

(IACUC-2018-0021).

Annexin-V staining

Alexa Fluor® 488 annexin V/Dead Cell Apoptosis Kit (Thermo Fisher Scientific) was used

according to manufacturers' recommendations. Fluorescence was measured by FACS Canto II

(BD Bioscience).

In situ proximity ligation assay (in situ PLA)

In situ PLA was performed with DuoLink in Situ Reagents (Sigma) according to the instructions.

Images were acquired with a ZEISS Axiocam 503 mono confocal microscope.

Mice and tumor initiation

Mice harboring Rb1^{flox}, Trp53^{flox} and Nkx2-1^{flox} alleles have been previously described (13, 14).

Tumors were generated by intratracheal delivery of Ad5CMV-Cre adenovirus (University of Iowa,

Gene Transfer Vector Core) as described (15). All studies were approved by the Committee for

Animal Care at MIT (A-3125–01).

Data analysis

Details are provided in the online supplement.

Statistical Analysis

Details of specific analyses are provided in the figure legends.

Data and materials availability

SCLC cell lines, human primary SCLC tumors, RP and RPN tumors H3K27ac ChIP-seq, Cell lines

and RP and RPN tumors RNA-seq, NKX2-1 and SOX1 ChIP-seq are deposited in NCBI GEO:

GSE183373.

RESULTS

Epigenomic profiling subdivided SCLC-A by differential super-enhancers

Genome-scale chromatin organization of the cell more faithfully reflects their lineage state than

transcriptomics (16). To refine current SCLC classification, we profiled genome-wide H3K27ac

to define SE regions across 16 SCLC cell lines. Unsupervised hierarchical clustering on SEs on

genes for transcriptional regulation with additional public data (4) revealed four subclasses (Figure

1A), supported by principal component analysis (PCA) (see Figure E1A in the online data

supplement). Our clusters recapitulated aspects of the expression classification. Cluster III

corresponds to SCLC-N (Figure 1A; purple). Cluster IV represents SCLC-P and SCLC-Y (Figure

1A; blue). Notably, epigenomic clustering distinguished SCLC-A into two distinct Clusters I and

II (Figure 1A; red and green) referred to as SCLC-Aα and SCLC-Aσ, while *ASCL1* SE is present

in 92.9% (13/14) of SCLC-A (Figure E1B).

To understand the difference underpinning SCLC-Aα and SCLC-Aσ, we identified 140

differentially enriched SEs. Among 110 SEs enriched in SCLC-Aα was *NKX2-1* (Figure 1B),

where H3K27ac signal was largely absent in other subclasses (Figure 1C). The 30 SEs enriched in

SCLC-Aσ included *TCF4*, a transcription factor that regulates synaptic plasticity (17) (Figure 1B

and Figure E1B). NKX2-1, also known as *TTF1*, a lineage transcription factor essential for the

development and maintenance of lungs, brain, and thyroid, is a frequently amplified lineage-

specific oncogene and a well-known marker to diagnose lung adenocarcinoma (LADC) (18) (19,

20). However, *NKX2-1* is not significantly amplified in human SCLCs (21) nor in the SCLC lines

used (Figure E1C). Nonetheless, SCLC-Aα cell lines harbor *NKX2-1* SE, whereas SCLC-Aσ cell

lines do not (Figure 1D and Figure E1D). Local H3K27ac enrichment pattern at the *NKX2-1* locus

in SCLC-Aα was almost identical to LADCs with *NKX2-1* amplification (Figure E1E). In 6 human

SCLC specimens, we found one SCLC-P tumor with *POU2F3* SE, and threes mixed SCLC-A/N

tumors with SEs near *ASCL1* and *NEUROD1* loci (Figure E1F). Notably, we observed two tumors

representing SCLC-Aα with high signals at *ASCL1* and *NKX2-1* loci (Figure 1E).

By immunohistochemistry (IHC) on another 34 human SCLC cohort, 65% (22/34) of cases

were NKX2-1-positive (Figure E2A), consistent with prior studies (22-26). In cell lines, SCLC-

Aα expressed substantially higher levels of NKX2-1 protein than other SCLC subtypes and

equivalent to *NKX2-1*-amplified LADCs (Figure 1F). Furthermore, RNA-seq datasets (5, 27)

confirmed *NKX2-1* is highly expressed in SCLC (Figure E2B), particularly in SCLC-A (Figure

1G), comparable to the levels in LADC (Figure E2C), consistent with a previous report (22).

Nonetheless, non-bimodal distribution of *NKX2-1* expression suggests that the distinction between

SCLC-Aα and Aσ is not discernible from expression analysis, perhaps analogous to the findings

that *NKX2-1* is almost universally expressed in LADC yet only amplified in 10-15%, which we

previously characterized as a distinct biological subgroup (19, 28).

NKX2-1 **survival dependency is unique to SCLC-Aα**

A recent study implicated that NKX2-1 promotes SCLC growth and regulates NE differentiation

and anti-apoptosis (29). We deleted *NKX2-1* in SCLC-Aα cell lines (Figure 2A) and found it

significantly suppressed cell growth and increased annexin-V-positive populations (Figure 2C and

E3A). By contrast, ablating NKX2-1 in SCLC cells from other clusters (Figure 2B) had little effect.

Given that acute removal of a master lineage factor in established cancer often leads to a halt in

cell growth (30), these data suggest NKX2-1 does not play a general oncogenic role in all lineage

states but its function is uniquely relevant to SCLC-Aα.

We further found a significant decrease in tumor volume for NKX2-1-deleted SCLC-Aα

cells in xenograft (Figure 2D and Figure E3B), confirming its essentiality *in vivo*. No obvious

histological differences were observed in the cells that formed small tumors which escaped from

NKX2-1 deletion (Figure E3C). In a short period after NKX2-1 deletion *in vitro*, expression of NE

markers was not significantly altered (Figure E3D). Overall, these data suggest that NKX2-1 is

functionally critical for SCLC-Aα *in vitro* and *in vivo*.

Genomic occupancy of *NKX2-1* in SCLC-Aα cells distinct from that in LADC cells

227   In LADC, NKX2-1 regulates gene expression and shares its cistromes with normal lung

(19); however, SCLCs typically do not express canonical NKX2-1 targets. Given NKX2-1's roles

in subsets of cortical, striatal, and pallidal neurons (31), we speculated its potential role for

neuronal differentiation in SCLC-Aα. Therefore, we profiled NKX2-1 cistromes by ChIP-seq in 5

SCLC-Aα cell lines and compared them to that from 3 LADC cell lines. Pairwise correlation

coefficient of all NKX2-1-bound sites and PCA revealed histology-specific binding profiles

(Figure 3A and E4A). Regions 4 (low-intensity signals) and 1 (high-intensity signals) are common

between SCLC-Aα and LADC (Figure 3B), which included the *RUNX1* locus (Figure 3C). Region

3 was unique to LADC that included *LMO3,* an essential mediator of NKX2-1 functions in LADCs

(19). Region 2 was unique to SCLC-Aα that included *DCC,* a receptor for netrin required for axon

guidance (32) (Figure 3C). These data suggest NKX2-1 regulates distinct transcriptional programs

across these two lineage states.

239   Ontologies on 661 SCLC-Aα-unique regions (Figure 3D and E4B) were enriched for

neuronal development, differentiation, and structure (Figure 3E). By contrast, genes regulating

neuronal biology were absent in LADC-unique regions (Figure E4C). Motif analysis revealed

SCLC-Aα-specific sites are enriched with ASCL2, Olig2, and SOX motifs, suggesting more

association with neuronal TFs (Figure E4D). In contrast, AP-1, Ets, Runx, and Tead-binding

motifs are enriched in LADC-specific regions, consistent with our previous report (19). These data

indicate that contribution of NKX2-1 in regulating pathways for maintenance of neural

differentiation state of SCLC-Aα, distinct from its role in normal lung epithelium and LADCs.

NKX2-1 interactome in chromatin of SCLC-Aα cells includes SOX1

NKX2-1 interacts with FOXA1 to co-regulate the expression of a set of genes for lineage control

in LADC (19, 20). Therefore, to identify unique partners that interact with NKX2-1 to exert distinct

transcriptional program in SCLC-Aα, we performed co-immunoprecipitation (co-IP) followed by

liquid chromatography tandem mass-spectrometry (LC-MS/MS) in the chromatin fraction of two

SCLC-Aα cell lines ectopically expressing FLAG-NKX2-1 (Figure 4A and Figure E5A). We

found 27 overlapping NKX2-1-interacting TFs in both cell lines (Figure 4B).

We further identified core regulatory circuitries (CRCs) (33) for SCLC-Aα and other SCLC

clusters from our H3K27ac data. Common CRCs included NEUROD1 for SCLC-N, POU2F3 and

ASCL2 for SCLC-P/Y, and TCF4 in SCLC-Aσ, consistent with our differential SE analysis

(Figure 1B and E5B). Common CRCs in SCLC-Aα included NKX2-1 (Figure 4C) along with

SOX1, NFIB, TCF12 and PBX1 also found in the SCLC-Aα NKX2-1-interactome. Of those,

H3K27ac signal at *SOX1* was highly enriched in SCLC-Aα (Figure 1B and Figure 4D), whereas

NFIB, *TCF12* and *PBX1* SEs were present across SCLC subtypes (Figure E5C). *SOX1* is highly

expressed in SCLC-A (Figure E5D) and positively correlated with *NKX2-1* expression in 185

SCLC tumors and cell lines (Figure E5E). SOX1 protein was also highly expressed specifically in

SCLC-Aα cell lines (Figure E5F).

The interaction between NKX2-1 and SOX1 was confirmed by co-IP followed by western

blot in two SCLC-Aα cell lines (Figure 4E), and further by proximity ligation assay (PLA) in 4

SCLC-Aα cell lines (Figure 4F and Figure E5G). These data suggest robust protein-protein

interaction between NKX2-1 and SOX1 uniquely in SCLC-Aα.

NKX2-1 and SOX1 co-occupy the SCLC-Aα genome to collaborate in neuronal gene

regulation

Next, we profiled genome-wide binding profiles of NKX2-1 and SOX1 in four SCLC-Aα cell lines

and found the majority of SOX1 peaks were bound by NKX2-1 (48.6-67.8%) (Figure 5A). The

2,652 regions co-bound by NKX2-1 and SOX1 in NCI-H187 were enriched for functions in neuron

differentiation and development, consistent with the aggregate analysis (Figure E6A and E6B).

We defined high-confidence NKX2-1 (5,663) and SOX1 (1,101) binding sites observed in ≧3 cell

lines, and found 604 peaks (54.9% of SOX1 peaks) are overlapped (Figure 5B), which included

the *NKX2-1* locus as well as neuronal genes *NRXN3* (34, 35) and *YWHAZ* (36) (Figure 5C) and

ontologically enriched for functions in neuron differentiation and development (Figure 5D). These

data suggest the interaction between NKX2-1 and SOX1 elicits specific cellular identity of SCLC-

Aα.

Collaboration of NKX2-1 and SOX1 is required to control the neuronal state in SCLC-Aα

We evaluated the transcriptional effect of depletion of *SOX1* or *NKX2-1* in NCI-H187 by RNA-

seq. After the abrogation of NKX2-1 or SOX1, 1,121 and 2,499 genes were significantly

downregulated respectively, with 873 genes commonly downregulated, suggesting a substantial

overlap in their transcriptional outputs (Figure 6A). We then queried the differentially co-

downregulated genes with the 2,652 genes with transcriptional start sites (TSSs) within 50 kb of

SOX1-NKX2-1 co-occupied peaks in NCI-H187, which were significantly enriched for NKX2-1

or SOX1 downregulated genes by ssGSEA analysis (Figure 6B). Leading edge analysis on these

genes revealed 289 overlaps, which we found are enriched for neuron differentiation and

development (Figure 6C). Together, these data suggest that NKX2-1 and SOX1 are specifically

co-localized on the SCLC-Aα genome and collaboratively contribute to controlling neuronal

differentiation state. We also found that SOX1 deletion, while not affecting the NKX2-1

expression, led to reduced cell growth (Figure E7A) with increased apoptosis (Figure E7B).

We previously found that expression of Nkx2-1 was critical for global Foxa1/2 binding

pattern in the LADC genome (20). To test if SOX1 is required for NKX2-1-binding to its target

genes in SCLC-Aα, we performed NKX2-1 ChIP-seq in NCI-H187 cells upon SOX1 depletion.

Pairwise correlation coefficient of NKX2-1 peaks across cells deleted for SOX1 or control

revealed strong correlation amongst each group (Figure 6D), supported by PCA analysis (Figure

E7C). Differential analysis on NKX2-1-bound sites found 2,067 peaks decreased upon SOX1

deletion (Figure E7D). k-means clustering of the differential sites found relatively uniform

reduction of NKX2-1 binding after SOX1 deletion (Figure 6E), where we also found decreased

H3K27ac modification compared to other sites (Figure 6F), suggesting that these NKX2-1-lost

sites are no longer in active state, similar to a recent finding on Sox2 in esophageal squamous cell

cancer model for Klf5 (37). The data suggest that loss of SOX1 releases NKX2-1 from co-occupied

loci in SCLC-Aα.

Nkx2-1 is essential for development of SCLC-Aα tumors in an autochthonous mouse model

The $Rb1^{fl/fl}Trp53^{fl/fl}$ (RP) is commonly used SCLC models that recapitulates the biology of

ASCL1-positive/classic human SCLCs (13) (Figure 7A). We first profiled H3K27ac to identify

SEs in mSCLC tumors from RP model, where Nkx2-1 is diffusely expressed. We found all 10

tumors from 2 different mice harbored high-level H3K27ac signal around the *Ascl1* locus, but none

harbored SEs at the *Neurod1*, *Yap1* or *Pou2f3* locus, suggesting SCLC-A subtype (Figure 7B and

Figure E8A) consistent with requirement of *Ascl1* for tumor formation in this model (38, 39). We

found 3 of 10 tumors simultaneously had high-level H3K27ac signals at the *Nkx2-1* and *Sox1* loci,

reminiscent of SCLC-Aα, while other 7 tumors have neither. This suggests that RP can adopt an

epigenetic state resembling either SCLC-Aα or SCLC-Aσ. Consistent but ambiguous expression

levels by IHC suggested epigenetic profiles distinguish these lineage classes more clearly (Figure

7C).

To explore whether NKX2-1 contributes to mSCLC differentiation, we generated

Rb1^{fl/fl}Trp53^{fl/fl}Nkx2-1^{fl/fl} (RPN) mice. We confirmed Nkx2-1 expression was absent in their tumors,

while 80% of them are Ascl1-positive suggesting that they remain representing SCLC-A. Sox1

staining was only weakly positive in 6/10 cases while 4 other cases, including the two Ascl1-

negative tumors, were negative (Figure 7C). We observed no significant difference in overall

survival between the two genotypes (Figure E8B). Morphologically, all but one mSCLC tumor

from the RP model had classical SCLC morphology (scant cytoplasm, nuclei with fine chromatin

and inconspicuous nucleoli), whereas a subset of tumors in RPN mice (15% in RPN vs. 4% in RP)

exhibited a 'non-classical' histology defined by higher grade/vesicular nuclei by two expert

pathologists (E.L.S. and M.B) (Figure 7D). These suggest that loss of *Nkx2*-1 leads to stochastic

propensity to form non-classical morphology, the while majority of the tumors retain classical

histology representing SCLC-A.

Next, we profiled H3K27ac in 10 mSCLC tumors from four RPN mice (11-13 months

post-initiation, including five classical and five non-classical). None of the tumors from the RPN

mice had high-level H3K27ac signal at the *Neurod1*, *Yap1* or *Pou2f3* loci, whereas most of RPN

tumors (7/10) harbored high-level H3K27ac signals around the *Ascl1* locus (Figure 7B and Figure

E8A). Strikingly, we found signals at the *Nkx2-1* and the *Sox1* loci were absent in all RPN tumors,

suggesting SCLC-Aσ subtype (Figure 7B). These data suggest that while the RP model can

develop both SCLC-Aα and SCLC-Aσ, *Nkx2-1* deletion at tumor initiation precludes SCLC-Aα

development.

Differential transcriptomic analysis from the tumors in RPN and RP mice also revealed

significantly downregulated genes in RPN tumors enriched for nervous system development

(Figure 7E), suggesting that SCLC-Aσ is hierarchically closer to the SCLC-N and SCLC-Y/P

subtypes, also supported by the data that variant morphology is more frequent in RPN. To address

whether the absence of Nkx2-1 resulted in distinct tumor evolution in RPN, we profiled copy

numbers of those mSCLCs and found that *Nfib* and *Mycl* loci are frequently amplified in RP model

(6/10) consistent with previous reports; however, these amplifications were absent in all tumors in

RPN model, suggesting the lack of Nkx2-1 predisposed those RPN tumors to undergo a distinct

path leading to a different lineage class of tumors (Figure 7F and Figure E8C).

DISCUSSION

In the current classification, SCLC-A constitutes 70% of SCLCs, while emerging evidence

suggests its heterogeneity. Our study revealed distinctive epigenetic landscapes within SCLC-A

that we designated SCLC-Aα and SCLC-Aσ, characterized by the presence of *NKX2-1* and *TCF4*

SEs. NKX2-1 is required for the lung development and later specification of the peripheral lung,

but also of bronchial NE cells (40). NKX2-1 is a well-known marker to diagnose LADCs (41), but

it is also frequently expressed in SCLC. As NKX2-1 plays key roles in brain development (42-45),

it has been speculated that NKX2-1 is involved in NE pathways, based on its expression enriched

in SCLC-A (46) and its physical interaction with ASCL1 (29). In our study, while most SCLC-A

expresses NKX2-1 at variable levels, only 48% had *NKX2-1* SE. Importantly, *NKX2-1* deletion in

SCLC-Aσ and SCLC-N cells, where NKX2-1 protein is still expressed, had little cell growth

effects, suggesting its relevancy only in SCLC-Aα.

Further, our data suggest this function is mediated through interaction with SOX1.

Combinations of master TFs that assemble at tissue-specific cis-regulatory sites control distinct

cellular functions essential for maintaining lineage states in many contexts. Therefore, while the

role of SOX1 had never been described in the lung, it is plausible that their interaction is critical

to retain unique NE state and it further suggests biological plasticity between glandular cell lineage

and pulmonary NE lineage.

The roles of NKX2-1 in carcinogenesis are complex and remain poorly understood. Nkx2-

1 deletion in neoplastic lungs causes loss of pulmonary identity and reversion to a foregut lineage.

Our data suggest that Nkx2-1 contributes to maintaining a specific neural lineage state that have

classical SCLC morphology. The observation that *Nkx2-1* deletion is tolerated at tumor initiation

in the GEMM, but not in established human SCLC-Aα cell lines, provokes a few potential

interpretations. First, the timing of the deletion might be critical in determining the outcome, as

evidenced in other studies on lineage factors (47, 48). Second, as CMV-Cre targets multiple cell

types, *Nkx2-1* deletion may favor origination from certain cell types, thus influencing tumor

evolution, as reflected by the lack of amplification at *Nfib* and *Mycl*. Third, early/in situ RPN

lesions that would have become SCLC-Aα might simply evolve to an alternative identity due to

lack of *Nkx2-1*.

Together, we subdivided the major SCLC-A subtype into SCLC-Aα and SCLC-Aσ by

revealing distinct epigenetic landscape in SCLC cell lines, human SCLC tumors, and GEMM.

Collaborative NKX2-1/SOX1 regulation defines the unique lineage state for SCLC-Aα. This study

will provide foundation for future identification of vulnerabilities and enable us to develop higher

level of personalized therapeutic approaches to benefit SCLC patients.

Acknowledgements

We thank Hong Cai, Yi Ge, Ying Liu, Peipei Guo, Yifei Sun, Xiaokun Liu, Korey Kam, Mingzhe

Li, Megan Vetter, Tyler Jacks and Karen Yee for helpful discussions and technical assistance,

Aleksandra Wroblewska and Brian D. Brown for providing a lentiviral vector for cloning sgRNAs;

NextSeq Sequencing Facility of the Department of Oncological Sciences at Icahn School of

Medicine at Mount Sinai (ISMMS), Saboor Hekmaty, Gayatri Panda and Ravi Sachidanandam for

sequencing assistance. The authors also thank Taplin Mass Spectrometry Facility at Harvard

Medical School, the Biorepository and Pathology Core Facility at ISMMS and Center for

Comparative Medicine and Surgery at the Icahn School of Medicine at Mount Sinai. Graphical

Abstract was created with BioRender.com.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; 69: 7-34.
2. Poirier JT, Gardner EE, Connis N, Moreira AL, de Stanchina E, Hann CL, Rudin CM. DNA methylation in
small cell lung cancer defines distinct disease subtypes and correlates with high expression of
EZH2. *Oncogene* 2015; 34: 5869-5878.
3. McColl K, Wildey G, Sakre N, Lipka MB, Behtaj M, Kresak A, Chen Y, Yang M, Velcheti V, Fu P, Dowlati A.
Reciprocal expression of INSM1 and YAP1 defines subgroups in small cell lung cancer. *Oncotarget*
2017; 8: 73745-73756.
4. Huang YH, Klingbeil O, He XY, Wu XLS, Arun G, Lu B, Somerville TDD, Milazzo JP, Wilkinson JE, Demerdash
OE, Spector DL, Egeblad M, Shi JW, Vakoc CR. POU2F3 is a master regulator of a tuft cell-like
variant of small cell lung cancer. *Gene Dev* 2018; 32: 915-928.
5. George J, Lim JS, Jang SJ, Cun Y, Ozretic L, Kong G, Leenders F, Lu X, Fernandez-Cuesta L, Bosco G, Muller
C, Dahmen I, Jahchan NS, Park KS, Yang D, Karnezis AN, Vaka D, Torres A, Wang MS, Korbel JO,
Menon R, Chun SM, Kim D, Wilkerson M, Hayes N, Engelmann D, Putzer B, Bos M, Michels S, Vlasic
I, Seidel D, Pinther B, Schaub P, Becker C, Altmuller J, Yokota J, Kohno T, Iwakawa R, Tsuta K,
Noguchi M, Muley T, Hoffmann H, Schnabel PA, Petersen I, Chen Y, Soltermann A, Tischler V, Choi
CM, Kim YH, Massion PP, Zou Y, Jovanovic D, Kontic M, Wright GM, Russell PA, Solomon B, Koch
I, Lindner M, Muscarella LA, la Torre A, Field JK, Jakopovic M, Knezevic J, Castanos-Velez E, Roz L,
Pastorino U, Brustugun OT, Lund-Iversen M, Thunnissen E, Kohler J, Schuler M, Botling J, Sandelin
M, Sanchez-Cespedes M, Salvesen HB, Achter V, Lang U, Bogus M, Schneider PM, Zander T, Ansen
S, Hallek M, Wolf J, Vingron M, Yatabe Y, Travis WD, Nurnberg P, Reinhardt C, Perner S, Heukamp
L, Buttner R, Haas SA, Brambilla E, Peifer M, Sage J, Thomas RK. Comprehensive genomic profiles
of small cell lung cancer. *Nature* 2015; 524: 47-53.
6. Rudin CM, Poirier JT, Byers LA, Dive C, Dowlati A, George J, Heymach JV, Johnson JE, Lehman JM,
MacPherson D, Massion PP, Minna JD, Oliver TG, Quaranta V, Sage J, Thomas RK, Vakoc CR, Gazdar
AF. Molecular subtypes of small cell lung cancer: a synthesis of human and mouse model data.
Nat Rev Cancer 2019; 19: 289-297.
7. Gay CM, Stewart CA, Park EM, Diao L, Groves SM, Heeke S, Nabet BY, Fujimoto J, Solis LM, Lu W, Xi Y,
Cardnell RJ, Wang Q, Fabbri G, Cargill KR, Vokes NI, Ramkumar K, Zhang B, Della Corte CM, Robson
P, Swisher SG, Roth JA, Glisson BS, Shames DS, Wistuba, II, Wang J, Quaranta V, Minna J, Heymach
JV, Byers LA. Patterns of transcription factor programs and immune pathway activation define
four major subtypes of SCLC with distinct therapeutic vulnerabilities. *Cancer Cell* 2021; 39: 346-
360 e347.
8. Patel AS, Yoo S, Kong R, Sato T, Sinha A, Karam S, Bao L, Fridrikh M, Emoto K, Nudelman G, Powell CA,
Beasley MB, Zhu J, Watanabe H. Prototypical oncogene family Myc defines unappreciated distinct
lineage states of small cell lung cancer. *Sci Adv* 2021; 7.
9. Iwafuchi-Doi M, Zaret KS. Pioneer transcription factors in cell reprogramming. *Genes Dev* 2014; 28:
2679-2692.
10. Sato T, Yoo S, Kong R, Sinha A, Chandramani-Shivalingappa P, Patel A, Fridrikh M, Nagano O, Masuko
T, Beasley MB, Powell CA, Zhu J, Watanabe H. Epigenomic profiling discovers trans-lineage SOX2
partnerships driving tumor heterogeneity in lung squamous cell carcinoma. *Cancer Res* 2019.
11. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova Z, Wilen C,
Orchard R, Virgin HW, Listgarten J, Root DE. Optimized sgRNA design to maximize activity and
minimize off-target effects of CRISPR-Cas9. *Nat Biotechnol* 2016; 34: 184-191.
12. Watanabe H, Ma Q, Peng S, Adelmant G, Swain D, Song W, Fox C, Francis JM, Pedamallu CS, DeLuca
DS, Brooks AN, Wang S, Que J, Rustgi AK, Wong KK, Ligon KL, Liu XS, Marto JA, Meyerson M, Bass

AJ. SOX2 and p63 colocalize at genetic loci in squamous cell carcinomas. *J Clin Invest* 2014; 124: 1636-1645.

13. Meuwissen R, Linn SC, Linnoila RI, Zevenhoven J, Mooi WJ, Berns A. Induction of small cell lung cancer by somatic inactivation of both Trp53 and Rb1 in a conditional mouse model. *Cancer Cell* 2003; 4: 181-189.

14. Kusakabe T, Kawaguchi A, Hoshi N, Kawaguchi R, Hoshi S, Kimura S. Thyroid-specific enhancer-binding protein/NKX2.1 is required for the maintenance of ordered architecture and function of the differentiated thyroid. *Mol Endocrinol* 2006; 20: 1796-1809.

15. DuPage M, Dooley AL, Jacks T. Conditional mouse lung cancer models using adenoviral or lentiviral delivery of Cre recombinase. *Nature protocols* 2009; 4: 1064-1072.

16. Zheng H, Xie W. The role of 3D genome organization in development and cell differentiation. *Nat Rev Mol Cell Biol* 2019; 20: 535-550.

17. Kennedy AJ, Rahn EJ, Paulukaitis BS, Savell KE, Kordasiewicz HB, Wang J, Lewis JW, Posey J, Strange SK, Guzman-Karlsson MC, Phillips SE, Decker K, Motley ST, Swayze EE, Ecker DJ, Michael TP, Day JJ, Sweatt JD. Tcf4 Regulates Synaptic Plasticity, DNA Methylation, and Memory Function. *Cell Rep* 2016; 16: 2666-2685.

18. Campbell JD, Alexandrov A, Kim J, Wala J, Berger AH, Pedamallu CS, Shukla SA, Guo G, Brooks AN, Murray BA, Imielinski M, Hu X, Ling S, Akbani R, Rosenberg M, Cibulskis C, Ramachandran A, Collisson EA, Kwiatkowski DJ, Lawrence MS, Weinstein JN, Verhaak RG, Wu CJ, Hammerman PS, Cherniack AD, Getz G, Cancer Genome Atlas Research N, Artyomov MN, Schreiber R, Govindan R, Meyerson M. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat Genet* 2016; 48: 607-616.

19. Watanabe H, Francis JM, Woo MS, Etemad B, Lin W, Fries DF, Peng S, Snyder EL, Tata PR, Izzo F, Schinzel AC, Cho J, Hammerman PS, Verhaak RG, Hahn WC, Rajagopal J, Jacks T, Meyerson M. Integrated cistromic and expression analysis of amplified NKX2-1 in lung adenocarcinoma identifies LMO3 as a functional transcriptional target. *Genes Dev* 2013; 27: 197-210.

20. Snyder EL, Watanabe H, Magendantz M, Hoersch S, Chen TA, Wang DG, Crowley D, Whittaker CA, Meyerson M, Kimura S, Jacks T. Nkx2-1 represses a latent gastric differentiation program in lung adenocarcinoma. *Mol Cell* 2013; 50: 185-199.

21. Peifer M, Fernandez-Cuesta L, Sos ML, George J, Seidel D, Kasper LH, Plenker D, Leenders F, Sun R, Zander T, Menon R, Koker M, Dahmen I, Muller C, Di Cerbo V, Schildhaus HU, Altmuller J, Baessmann I, Becker C, de Wilde B, Vandesompele J, Bohm D, Ansen S, Gabler F, Wilkening I, Heynck S, Heuckmann JM, Lu X, Carter SL, Cibulskis K, Banerji S, Getz G, Park KS, Rauh D, Grutter C, Fischer M, Pasqualucci L, Wright G, Wainer Z, Russell P, Petersen I, Chen Y, Stoelben E, Ludwig C, Schnabel P, Hoffmann H, Muley T, Brockmann M, Engel-Riedel W, Muscarella LA, Fazio VM, Groen H, Timens W, Sietsma H, Thunnissen E, Smit E, Heideman DA, Snijders PJ, Cappuzzo F, Ligorio C, Damiani S, Field J, Solberg S, Brustugun OT, Lund-Iversen M, Sanger J, Clement JH, Soltermann A, Moch H, Weder W, Solomon B, Soria JC, Validire P, Besse B, Brambilla E, Brambilla C, Lantuejoul S, Lorimier P, Schneider PM, Hallek M, Pao W, Meyerson M, Sage J, Shendure J, Schneider R, Buttner R, Wolf J, Nurnberg P, Perner S, Heukamp LC, Brindle PK, Haas S, Thomas RK. Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat Genet* 2012; 44: 1104-1110.

22. Horie M, Miyashita N, Mattsson JSM, Mikami Y, Sandelin M, Brunnstrom H, Micke P, Nagase T, Saito A. An integrative transcriptome analysis reveals a functional role for thyroid transcription factor-1 in small cell lung cancer. *J Pathol* 2018; 246: 154-165.

23. Misch D, Blum T, Boch C, Weiss T, Crolow C, Griff S, Mairinger T, Bauer TT, Kollmeier J. Value of thyroid transcription factor (TTF)-1 for diagnosis and prognosis of patients with locally advanced or metastatic small cell lung cancer. *Diagn Pathol* 2015; 10: 21.

484 24. Miyauchi E, Motoi N, Ono H, Ninomiya H, Ohyanagi F, Nishio M, Okumura S, Ichinose M, Ishikawa Y.
485    Distinct Characteristics of Small Cell Lung Cancer Correlate With Central or Peripheral Origin:
486    Subtyping Based on Location and Expression of Transcription Factor TTF-1. *Medicine (Baltimore)*
487    2015; 94: e2324.
488 25. Wu M, Wang B, Gil J, Sabo E, Miller L, Gan L, Burstein DE. p63 and TTF-1 immunostaining. A useful
489    marker panel for distinguishing small cell carcinoma of lung from poorly differentiated squamous
490    cell carcinoma of lung. *Am J Clin Pathol* 2003; 119: 696-702.
491 26. Ordonez NG. Value of thyroid transcription factor-1 immunostaining in distinguishing small cell lung
492    carcinomas from other small cell carcinomas. *Am J Surg Pathol* 2000; 24: 1217-1223.
493 27. Jiang L, Huang J, Higgs BW, Hu Z, Xiao Z, Yao X, Conley S, Zhong H, Liu Z, Brohawn P, Shen D, Wu S, Ge
494    X, Jiang Y, Zhao Y, Lou Y, Morehouse C, Zhu W, Sebastian Y, Czapiga M, Oganesyan V, Fu H, Niu Y,
495    Zhang W, Streicher K, Tice D, Zhao H, Zhu M, Xu L, Herbst R, Su X, Gu Y, Li S, Huang L, Gu J, Han B,
496    Jallal B, Shen H, Yao Y. Genomic Landscape Survey Identifies SRSF1 as a Key Oncodriver in Small
497    Cell Lung Cancer. *PLoS Genet* 2016; 12: e1005895.
498 28. Weir BA, Woo MS, Getz G, Perner S, Ding L, Beroukhim R, Lin WM, Province MA, Kraja A, Johnson LA,
499    Shah K, Sato M, Thomas RK, Barletta JA, Borecki IB, Broderick S, Chang AC, Chiang DY, Chirieac LR,
500    Cho J, Fujii Y, Gazdar AF, Giordano T, Greulich H, Hanna M, Johnson BE, Kris MG, Lash A, Lin L,
501    Lindeman N, Mardis ER, McPherson JD, Minna JD, Morgan MB, Nadel M, Orringer MB, Osborne
502    JR, Ozenberger B, Ramos AH, Robinson J, Roth JA, Rusch V, Sasaki H, Shepherd F, Sougnez C, Spitz
503    MR, Tsao MS, Twomey D, Verhaak RG, Weinstock GM, Wheeler DA, Winckler W, Yoshizawa A, Yu
504    S, Zakowski MF, Zhang Q, Beer DG, Wistuba, II, Watson MA, Garraway LA, Ladanyi M, Travis WD,
505    Pao W, Rubin MA, Gabriel SB, Gibbs RA, Varmus HE, Wilson RK, Lander ES, Meyerson M.
506    Characterizing the cancer genome in lung adenocarcinoma. *Nature* 2007; 450: 893-898.
507 29. Hokari S, Tamura Y, Kaneda A, Katsura A, Morikawa M, Murai F, Ehata S, Tsutsumi S, Ishikawa Y,
508    Aburatani H, Kikuchi T, Miyazono K, Koinuma D. Comparative analysis of TTF-1-binding DNA
509    regions in small cell lung cancer and non-small cell lung cancer. *Mol Oncol* 2019.
510 30. Heinz S, Romanoski CE, Benner C, Glass CK. The selection and function of cell type-specific enhancers.
511    *Nat Rev Mol Cell Biol* 2015; 16: 144-154.
512 31. Magno L, Barry C, Schmidt-Hieber C, Theodotou P, Hausser M, Kessaris N. NKX2-1 Is Required in the
513    Embryonic Septum for Cholinergic System Development, Learning, and Memory. *Cell Rep* 2017;
514    20: 1572-1584.
515 32. Liu Y, Bhowmick T, Liu Y, Gao X, Mertens HDT, Svergun DI, Xiao J, Zhang Y, Wang JH, Meijers R.
516    Structural Basis for Draxin-Modulated Axon Guidance and Fasciculation by Netrin-1 through DCC.
517    *Neuron* 2018; 97: 1261-1267 e1264.
518 33. Saint-Andre V, Federation AJ, Lin CY, Abraham BJ, Reddy J, Lee TI, Bradner JE, Young RA. Models of
519    human core transcriptional regulatory circuitries. *Genome Res* 2016; 26: 385-396.
520 34. Reissner C, Runkel F, Missler M. Neurexins. *Genome Biol* 2013; 14: 213.
521 35. Sudhof TC. Neuroligins and neurexins link synaptic function to cognitive disease. *Nature* 2008; 455:
522    903-911.
523 36. Toma C, Torrico B, Hervas A, Valdes-Mas R, Tristan-Noguero A, Padillo V, Maristany M, Salgado M,
524    Arenas C, Puente XS, Bayes M, Cormand B. Exome sequencing in multiplex autism families
525    suggests a major role for heterozygous truncating mutations. *Mol Psychiatry* 2014; 19: 784-790.
526 37. Wu Z, Zhou J, Zhang X, Zhang Z, Xie Y, Liu JB, Ho ZV, Panda A, Qiu X, Cejas P, Canadas I, Akarca FG,
527    McFarland JM, Nagaraja AK, Goss LB, Kesten N, Si L, Lim K, Liu Y, Zhang Y, Baek JY, Liu Y, Patil DT,
528    Katz JP, Hai J, Bao C, Stachler M, Qi J, Ishizuka JJ, Nakagawa H, Rustgi AK, Wong KK, Meyerson M,
529    Barbie DA, Brown M, Long H, Bass AJ. Reprogramming of the esophageal squamous carcinoma
530    epigenome by SOX2 promotes ADAR1 dependence. *Nat Genet* 2021; 53: 881-894.

38. Borromeo MD, Savage TK, Kollipara RK, He M, Augustyn A, Osborne JK, Girard L, Minna JD, Gazdar AF,
Cobb MH, Johnson JE. ASCL1 and NEUROD1 Reveal Heterogeneity in Pulmonary Neuroendocrine
Tumors and Regulate Distinct Genetic Programs. *Cell Rep* 2016; 16: 1259-1272.
39. Olsen RR, Ireland AS, Kastner DW, Groves SM, Spainhower KB, Pozo K, Kelenis DP, Whitney CP, Guthrie
MR, Wait SJ, Soltero D, Witt BL, Quaranta V, Johnson JE, Oliver TG. ASCL1 represses a SOX9(+)
neural crest stem-like state in small cell lung cancer. *Genes Dev* 2021; 35: 847-869.
40. La Rosa S, Chiaravalli AM, Placidi C, Papanikolaou N, Cerati M, Capella C. TTF1 expression in normal
lung neuroendocrine cells and related tumors: immunohistochemical study comparing two
different monoclonal antibodies. *Virchows Arch* 2010; 457: 497-507.
41. Yamaguchi T, Hosono Y, Yanagisawa K, Takahashi T. NKX2-1/TTF-1: an enigmatic oncogene that
functions as a double-edged sword for cancer cell survival and progression. *Cancer Cell* 2013; 23:
718-723.
42. Puelles L, Kuwana E, Puelles E, Bulfone A, Shimamura K, Keleher J, Smiga S, Rubenstein JL. Pallial and
subpallial derivatives in the embryonic chick and mouse telencephalon, traced by the expression
of the genes Dlx-2, Emx-1, Nkx-2.1, Pax-6, and Tbr-1. *J Comp Neurol* 2000; 424: 409-438.
43. Flames N, Pla R, Gelman DM, Rubenstein JL, Puelles L, Marin O. Delineation of multiple subpallial
progenitor domains by the combinatorial expression of transcriptional codes. *J Neurosci* 2007; 27:
9682-9695.
44. Sussel L, Marin O, Kimura S, Rubenstein JL. Loss of Nkx2.1 homeobox gene function results in a ventral
to dorsal molecular respecification within the basal telencephalon: evidence for a transformation
of the pallidum into the striatum. *Development* 1999; 126: 3359-3370.
45. Minocha S, Valloton D, Ypsilanti AR, Fiumelli H, Allen EA, Yanagawa Y, Marin O, Chedotal A, Hornung
JP, Lebrand C. Nkx2.1-derived astrocytes and neurons together with Slit2 are indispensable for
anterior commissure formation. *Nat Commun* 2015; 6: 6887.
46. Cardnell RJ, Li L, Sen T, Bara R, Tong P, Fujimoto J, Ireland AS, Guthrie MR, Bheddah S, Banerjee U, Kalu
NN, Fan YH, Dylla SJ, Johnson FM, Wistuba, II, Oliver TG, Heymach JV, Glisson BS, Wang J, Byers
LA. Protein expression of TTF1 and cMYC define distinct molecular subgroups of small cell lung
cancer with unique vulnerabilities to aurora kinase inhibition, DLL3 targeting, and other targeted
therapies. *Oncotarget* 2017; 8: 73419-73432.
47. Camolotto SA, Pattabiraman S, Mosbruger TL, Jones A, Belova VK, Orstad G, Streiff M, Salmond L,
Stubben C, Kaestner KH, Snyder EL. FoxA1 and FoxA2 drive gastric differentiation and suppress
squamous identity in NKX2-1-negative lung cancer. *Elife* 2018; 7.
48. Zewdu R, Mehrabad EM, Ingram K, Fang P, Gillis KL, Camolotto SA, Orstad G, Jones A, Mendoza MC,
Spike BT, Snyder EL. An NKX2-1/ERK/WNT feedback loop modulates gastric identity and response
to targeted therapy in lung adenocarcinoma. *Elife* 2021; 10.

Figure legends

Figure 1. Epigenomic profiling subdivided SCLC-A by differential super-enhancers

(**A**). Unsupervised hierarchical clustering of 25 SCLC cell lines using SE signals near

transcriptional regulator genes. The color scale indicates the intensity of SE signals normalized to

linear scores using the sum of squares of the values. Each row represents a SE locus (747 loci).

The color shades on the dendrogram indicate four novel SE clusters, the top red columns display

the presence of SE near the *NKX2-1* locus, and the bottom column colors denote the consensus

expression clusters. (**B**). Volcano plot showing 140 differentially enriched SEs (110 enriched in

SCLC-Aα and 30 enriched in SCLC-Aσ) at thresholds of absolute fold change ≥ 5 and FDR ≤ 0.25

depicted as dotted lines. (**C**). Genome view tracks of H3K27ac signal at the *NKX2-1* locus in four

clusters of SCLC cell lines. Normalized scales are shown in Figure E1B. (**D**). SE plots with all

enhancers rank-ordered by H3K27ac signals in DMS 79, NCI-H187, NCI-H209 and NCI-H345.

Dotted lines depict tangential cutoffs to define SEs. (**E**). Genome view tracks of H3K27ac signals

at the *ASCL1*, *NEUROD1*, *POU2F3*, *YAP1* and *NKX2-1* loci in 2 human SCLC specimens of

SCLC-A subtype. (**F**). Protein expression of NKX2-1, ASCL1 and vinculin as a loading control in

SCLC cell lines from different clusters. *NKX2-1*-amplified adenocarcinoma cell lines NCI-H3122

and NCI-H2087 are shown as reference. (**G**). Box and jitter plots showing the distribution of

NKX2-1 mRNA expression in SCLC-A, SCLC-N, SCLC-P, and SCLC-Y subtypes of 135 primary

SCLC tumors.

Figure 2. *NKX2-1* survival dependency is unique to SCLC-Aα

(**A**). *(Top)* Immunoblots showing protein expression of NKX2-1 and vinculin as a loading control

in SCLC-Aα cell lines with three *NKX2-1* sgRNAs, or two non-target (NT) sgRNAs. *(Bottom)*

Cell growth curve for SCLC-Aα cell lines infected with *NKX2-1* sgRNAs or NT sgRNAs. Data

are shown as mean ± SD (n=3); *P < 0.0001 by two-sided two-way ANOVA. (**B**). *(Top)*

Immunoblots showing protein expression of NKX2-1 and vinculin as a loading control in SCLC-

Aσ (NCI-H2171, NCI-H69) and SCLC-N (NCI-H1963) cell lines with *NKX2-1* sgRNAs, or non-

target (NT) sgRNAs. *(Bottom)* Cell growth curve for cell lines infected with *NKX2-1* sgRNA or

NT sgRNAs. Data are shown as mean ± SD (n=3); *P < 0.0001 by two-sided two-way ANOVA.

(**C**). Bar plots showing percentage of apoptotic NCI-H187 cells infected with NT sgRNA and

NKX2-1 sgRNA, measured by annexin V staining. Means ± SD of three biological replicates. *P

< 0.0001, t test. Representative data for NT sg1 and NKX2-1 sg1 can be found in Figure E2A. (**D**).

(*Left*) Tumor growth curve for xenograft tumors of NCI-H187 cells infected with *NKX2-1* sg1 or

NT sg1 *in vivo*. Mean±SD of four biological replicates. *P < 0.0001, t test. Xenograft tumors were

resected at 3 weeks after inoculation. (*Right*) Images of resected xenograft tumors of NCI-H209

cells infected with NT sg1 or *NKX2-1* sg1 *in vivo* at day 23.

Figure 3. Genomic occupancy of *NKX2-1* in SCLC-Aα cells distinct from that in LADC cells

(**A**). Correlation matrix depicting pairwise comparisons of identified NKX2-1 binding peaks in the

5 SCLC-Aα cell lines and in 3 LADC cell lines. Color scale represents degree of Pearson's

correlation coefficient. (**B**). Heatmap showing signals from NKX2-1 ChIP-seq data in the 5 SCLC-

Aα cell lines and 3 LUAC cell lines. ChIP signal intensity is shown by red shading. Y-axis contains

all peaks that are bound by NKX2-1 in any of SCLC-Aα and LUAC (n=37,198). Peaks are divided

into 4 regions by *k-means* (k=4) clustering. (**C**). NKX2-1 ChIP-seq signals in SCLC-Aα and

LADC cell lines at representative loci (*RUNX1*; cluster 1, *LMO3*; cluster 3 and *DCC*; cluster 2).

Red: SCLC-Aα cell lines. Black: LADC cell lines. (**D**). Heatmap showing 2,671 differentially

NKX2-1 bound regions (fold change $\geq 2^5$ and FDR $\leq 1\times10^{-5}$), 2,010 enriched in LADC and 661

enriched in SCLC-Aα, between 5 SCLC-Aα cell lines and 3 LUAC cell lines. (**E**). Enriched

ontology by GREAT analyses for regions differentially bound by NKX2-1 in 5 SCLC-Aα

compared to LADCs.

Figure 4. NKX2-1 interactome in chromatin of SCLC-Aα cells includes SOX1

(**A**). Experimental strategy to identify binding partners of NKX2-1 in SCLC-Aα. Chromatin

prepared from the nuclear fraction of SCLC-Aα cell lines NCI-H187 and NCI-H1436 stably

expressing FLAG-NKX2-1 was solubilized using micrococcal nuclease digestion. Co-

immunoprecipitated proteins with anti-FLAG antibody from solubilized chromatin were identified

by LC-MS/MS (see Data file S2 for the complete list). (**B**). Venn diagram showing the number of

transcription factors identified in each of the 2 SCLC-Aα cell lines and the 27 overlaps found in

both cell lines. The proteins indicated in yellow color were also found in CRC analysis in Figure

4C. (**C**). Core transcriptional regulatory circuitries (CRCs) commonly identified in 8 SCLC-Aα

cell lines. The genes in yellow color were also found in LC-MS/MS analysis in Figure 4B. (**D**).

Genome view tracks of H3K27ac signal at the *SOX1* locus in four clusters of SCLC cell lines. (**E**).

Immunoblots with anti-SOX1 antibody on endogenous co-immunoprecipitates by anti-NKX2-1

antibody in SCLC-Aα (DMS 79 and NCI-H187) cells. (**F**). *In situ* proximity ligation assay signals

(red fluorescence) targeting endogenous SOX1 and NKX2-1 in NCI-H841, NCI-H187, NCI-H209

and DMS 79 cells. Nuclei are counterstained with DAPI (blue). See Figure E4G for additional

control.

Figure 5. NKX2-1 and SOX1 co-occupy the SCLC-Aα genome to collaborate in gene

regulation

(**A**). Heatmap showing k-means clustering (*k*=4) of signals at the common 4,051 loci from NKX2-

1 (red) and SOX1 (blue) ChIP-seq data in the 4 SCLC-Aα cell lines. ChIP signal intensity is shown

by shading for all NKX2-1 and SOX1 occupied loci. (**B**). Venn diagram showing the overlap of

NKX2-1 common binding sites and SOX1 common binding sites present in at least 3 SCLC-Aα

cell lines. (**C**). Genome view tracks of NKX2-1 (red) and SOX1 (blue) ChIP-seq signals in four

SCLC-Aα (DMS 79, NCI-H187, NCI-H209 and NCI-H345) cell lines and 3 LADC cell lines at

the representative NKX2-1-SOX1 co-occupied loci (*NRXN3* and *YWHAZ*; SCLC-Aα unique*,*

NKX2-1; SCLC-Aα/LADC common). (**D**). Enriched ontology by GREAT analyses for regions in

604 overlapping high-confidence NKX2-1-SOX1 binding sites.

Figure 6. Collaboration of NKX2-1 and SOX1 is required to control the neuronal state in

SCLC-Aα

(**A**). Venn diagram showing the overlap of downregulated genes after depletion of NKX2-1 (fold

change > 1.4 and adjusted P < 0.05) and SOX1 (fold change > 2 and adjusted P < 0.05) in NCI-

H187 cells. (**B**). Gene Set Enrichment Analysis (GSEA) of genes associated with 2,652 NKX2-1-

SOX1 co-occupied loci for expression data upon repression of NKX2-1 (*left*) or SOX1 (*right*) in

NCI-H187 cells, with their leading-edge genes shaded. Enrichment was significant for both (NES=

-2.29, P < 0.001 for NKX2-1 and NES= -2.54, P < 0.001 for SOX1). (**C**). Enriched gene ontologies

(MSigDB, biological process) for 289 common genes in leading-edge genes for both NKX2-1 and

SOX1 deletions. (**D**). Correlation matrix depicting pairwise comparisons of identified NKX2-1

binding regions in three *SOX1* sgRNAs and three control cells (two non-target (NT) sgRNAs, one

parental cell) of NCI-H187 cell line. Color scale represents degree of Pearson's correlation

coefficient. (**E**). Heatmap showing k-means clustering (*k*=4) of 2,067 NKX2-1 ChIP-seq loci in

NCI-H187 sgSOX1 compared to control cells with reduced signal intensity. ChIP-seq signal

intensity is shown by color shading. (**F**). Average H3K27ac ChIP-seq signals on NKX2-1 reduced

binding sites and NKX2-1 non-reduced binding sites defined in Figure E6D for NCI-H187 parental,

NT sg1, SOX1 sg2 and SOX1 sg3 cells.

Figure 7. *Nkx2-1* is essential for development of SCLC-Aα tumors in an autochthonous

mouse model

(**A**). Genetically engineered mouse models of SCLCs. *Rb1*$^{flox/flox}$; *Trp53*$^{flox/flox}$ (RP) mice or

Rb1$^{flox/flox}$; *Trp53*$^{flox/flox}$, *Nkx2-1*$^{flox/flox}$ (RPN) mice were transduced with Adeno-CMV-*Cre* to

initiate SCLC. (**B**). Genome view tracks at the Nkx2-1, Sox1, and Ascl1 loci for H3K27ac ChIP-

seq signals in 10 SCLC tumors from RP mice and 10 SCLC tumors from RPN mice. * indicates 3

tumors that have high-level signal at both *Nkx2-1* and *Sox1* loci. The arrowheads on *Nkx2-1* locus

indicate the loxP sites in RPN mice**. (C)**. Representative immune-staining images for Nkx2-1,

Sox1, and Ascl1 on SCLC tumors from RP and RPN mice. Scale bars, 25μm. (**D**). Representative

hematoxylin and eosin (H&E) images of SCLC tumors from RP and RPN mice. Scale bars, 2.5mm

and 50μm, respectively. (**E**). Enriched gene ontologies (MSigDB, biological process) for the

differentially downregulated genes in RPN tumors compared to RP tumors. (**F**). Average copy

number profiles in RP and RPN models. Arrows indicate the *Nfib* and *Mycl* loci that are

specifically amplified in RP tumors (individual copy number profiles at those loci are available in

Figure E8C).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

Adeno CMV-Cre → Intratracheal instillation

*Rb1^flox/flox^; Trp53^flox/flox^* (RP)

Adeno CMV-Cre → Intratracheal instillation

*Rb1^flox/flox^; Trp53^flox/flox^; Nkx2-1^flox/flox^* (RPN)

B

H3K27ac

chr12   chr8   chr10

Nkx2-1   *Sox1*   *Ascl1*

C

RP (Tumor 5)   RPN (Tumor 5)

Nkx2-1

Sox1

Ascl1

D

H&E

RP

RPN

E

Downregulated genes in RPN

- Central Nervous System Development
- Cell Cell Signaling
- Nuclear Chromosome
- Animal Organ Morphogenesis
- Postsynapse
- Epithelium Development
- Behavior
- Locomotory Behavior
- Renal System Development
- Chromosome

$-log_{10}$ (p-Value)

F

Nfib   *Mycl*

RP

RPN

chr 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

Log2