Transcriptional circuitry of NKX2-1 and SOX1 defines an unrecognized lineage subtype of small cell lung cancer

Ranran Kong[1,2,3], Ayushi S. Patel[2,3,4], Takashi Sato[2,3,5], Seungyeul Yoo[6,7], Li Bao[8], Abhilasha Sinha[2,3], Feng Jiang[2,3], Yang Tian[2,3], Maya Fridrikh[2,3], Shuhui Liu[9], Jie Feng[10], Xijing He[11,12], Jiantao Jiang[1], Yuefeng Ma[1], Karina Grullon[2,3], Dawei Yang[2,3,13], Charles A. Powell[2,3], Mary Beth Beasley[14], Jun Zhu[3,6,7], Eric L. Snyder[15,16], Shaomin Li[1]*, and Hideo Watanabe[2,3,6]*

[1]Department of Thoracic Surgery, The Second Affiliated Hospital of Medical School, Xi'an Jiaotong University, Xi'an, Shaanxi, 710004, China;

[2]Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[3]Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[4]Division of Hematology & Medical Oncology, Laura and Isaac Perlmutter Cancer Center, New York University Langone Medical Center, New York, NY, USA;

[5]Department of Respiratory Medicine, Kitasato University School of Medicine, Sagamihara, Japan; Division of Pulmonary Medicine, Department of Medicine, Keio University School of Medicine, Tokyo, Japan;

[6]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[7]Sema4, a Mount Sinai venture, Stamford, CT 06902, USA;

[8]Ningxia People's Hospital, Yinchuan, Ningxia Province, 750001, China;

[9]Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA;

[10]Department of Nephrology, The First Affiliated Hospital of Medical School, Xi'an Jiaotong University, Xi'an, Shaanxi, 710061, China;

[11]Department of Orthopedics, The Second Affiliated Hospital of Medical School, Xi'an Jiaotong

University, Xi'an, Shaanxi, 710004, China;

[12]Xi'an International Medical Center, Xi'an, Shaanxi, 710126, China;

[13]Department of Pulmonary and Critical Care Medicine, Zhongshan Hospital Fudan University, Shanghai,

200032, China;

[14]Department of Pathology and Laboratory Medicine, Icahn School of Medicine at Mount Sinai, New

York, NY 10029, USA;

[15]Department of Pathology, University of Utah, Salt Lake City, UT 84112; Department of Oncological

Sciences, University of Utah, Salt Lake City, UT 84112, USA;

[16]Huntsman Cancer Institute, University of Utah, Salt Lake City, UT 84112

* Both authors share senior authorship.

H.W. has been supported by ATS Foundation Unrestricted Grant (ATS-2017-24), the American

Lung Association of the Northeast Lung Cancer Discovery Award (LCD-504985), Department

of Defense (W81XWH-19-1-0613) and NIH (R01CA240342). R.K. is supported by Shaanxi

Provincial Natural Science Foundation, China (2017JM8046). T. S. is supported by the Japanese

Respiratory Society, the 6th Lilly Oncology Fellowship Program, and the Uehara Memorial

Foundation. J. F. is supported by National Natural Science Foundation of China (NSFC) (No.

82100718). D. Y. is supported by Shanghai Pujiang Program (20PJ1402400). C.A.P. and H. W.

are supported by NIH (R01CA163772). E.L.S. is supported by the NIH (R01CA212415,

R01CA240317) and the American Lung Association Lung Cancer Discovery Award (LCD-

821670). This work was supported in part through the computational resources and staff

expertise provided by Scientific Computing at the Icahn School of Medicine at Mount Sinai.

Correspondence and requests for reprints should be addressed to Hideo Watanabe, Division of

Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, Icahn School of

Medicine at Mount Sinai, New York, NY 10029, USA. Phone: 212-241-9242; Fax: 212-876-

5519; Email: hideo.watanabe@mssm.edu.

Author contributions: Conceptualization & Methodology, R.K., and H.W.; Data Collection:

R.K., A.S.P., T.S., S.Y., L.B., F.J., M.F., S.L., J.F., X.H., J.J., Y.M., K.G., D.Y., E.L.S., S.L.,

L.S., and H.W.; Resources, E.L.S., M.B.B., C.A.P., S.L., and H.W.; Data Analysis, R.K., S.Y.,

A.S.P., S.L., M.B.B., E.L.S., and H.W.; Writing – Original Draft, R.K., A.S.P. and H.W.;

Writing-Review & Editing, R.K., A.S.P., E.L.S., and H.W.; Supervision, C.A.P., J.Z. S.L., and

H.W.; Project Administration & Funding Acquisition, H.W.

Running title: Collaboration of NKX2-1 and SOX1 defines SCLC-A1 subtype

Descriptor number: 9.26 Lung Cancer: Cellular and Molecular Aspects

The total word count for the body of the manuscript: 3,500

This article has an online data supplement, which is accessible from this issue's table of content

online   at www.atsjournals.org

Abstract:

Rationale: The current molecular classification of small cell lung cancer (SCLC) based on

expression of four lineage transcription factors still leaves its major subtype SCLC-A as a

heterogeneous group, necessitating more precise characterization of lineage subclasses.

Objectives: To refine the current SCLC classification with epigenomic profiles and to identify

features of the re-defined SCLC subtypes.

Methods: We performed unsupervised clustering of epigenomic profiles on 25 SCLC cell lines.

Functional significance was evaluated by cell growth, apoptosis and xenograft using CRISPR-

Cas9-mediated deletion. The specific cistromic profiles by ChIP-seq and its functional

transcriptional partners using co-immunoprecipitation followed by mass spectrometry were

determined. $Rb1^{fl/fl}Trp53^{fl/fl}$ and $Rb1^{fl/fl}Trp53^{fl/fl}Nkx2\text{-}1^{fl/fl}$ mouse models were engineered to

explore the function of $Nkx2\text{-}1$ in tumor initiation and differentiation. H3K27ac profiles were

analyzed to reveal 6 human SCLC specimen and 20 mice tumors epigenomic landscapes.

Measurements and Main Results: We identified an epigenomic subclusters of the major SCLC-

A subtype, named SCLC-A1 and SCLC-A2. SCLC-A1 was characterized by the presence of a

super-enhancer at the $NKX2\text{-}1$ locus, which was observed in human SCLC specimens and a

murine SCLC model. We found NKX2-1, a dual lung and neural lineage factor, is uniquely

relevant in SCLC-A1. We further found maintenance of this neural identity in SCLC-A1 is

mediated by collaborative transcriptional activity with another neuronal transcriptional factor

SOX1.

Conclusions: We comprehensively describe an additional epigenomic heterogeneity of the major

SCLC-A subtype, and define SCLC-A1 subtype by the core regulatory circuitry representing

NKX2-1 and SOX1 super-enhancers and their functional collaborations to maintain neuronal

linage state. (Word count 250)

Keywords: small cell lung cancer (SCLC); *NKX2-1/TTF-1*; *SOX1*; Super-enhancer

INTRODUCTION

Small cell lung cancer (SCLC) is an aggressive, lethal, and rapidly progressing neuroendocrine

(NE) malignancy, accounting for ~15% of lung cancer incidence and 250,000 deaths worldwide

annually (1). The five-year survival rate for patients with SCLC is only 6% compared to 19% for

non-SCLC. The lack of progress in SCLC treatment is largely attributable to the inadequate

molecular understanding. Advances in genomic technology have enabled new molecular

characterization of SCLCs. Genome-wide methylation and transcriptomes of human SCLCs and

PDX models identified NE and non-NE subtypes. NE subtype is defined by the expression of

neuronal lineage transcription factors (TFs) ASCL1 and/or NeuroD1 (2). Two recent studies

described the non-NE SCLCs, one signified by YAP1 expression and another by POU2F3

expression (3, 4). In light of these findings, a consortium proposed a new molecular

classification of SCLCs that distinguishes them into four subtypes based on the expression of

four lineage-defining transcription factors: ASCL1, NeuroD1, YAP1 and POU2F3 referred to as

SCLC-A, SCLC-N, SCLC-Y and SCLC-P, respectively (5, 6). The classification, while marking

significant advance, is still a work in progress. After that report, a newer study proposed an

inflamed SCLC subtype (SCLC-I) (7). In addition, not all SCLCs clearly fall into the four

clusters as they often express multiple factors at variable levels, especially in the SCLC-A

subtype which account for 70% of SCLCs (8). This wide range of heterogeneity warrants novel

approaches to develop a more precise classification of SCLC based on lineage subtype.

Lineage-specific transcription factors and the presence of SE at their gene locus are

critical determinants of cellular identity (9). In our study, we applied epigenomic approaches to

investigate the differential enhancer landscapes in SCLC cell lines, human SCLC samples and

autochthonous murine SCLCs from a genetic engineered mouse model (GEMM) to reveal

additional intertumoral heterogeneity, providing comprehensive evidence of distinct histone

modification profiles. Furthermore, we determined a novel mechanism of subtype-specific

transcriptional regulation in SCLC, integrating the core transcriptional networks and their

collaborative   regulation   of   NE   target   genes.

METHODS

For a detailed description of the methods, please see the online supplement.

Cell lines

Cell lines used in this study are listed in Table E1.

Chromatin Immunoprecipitation followed by sequencing (ChIP-seq)

ChIP-seq was performed as described previously (10). Protein G magnetic beads (Dynabeads:

Life Technologies) were pre-incubated with anti-H3K27ac (Abcam, ab4729) or anti-NKX2-1

(Bethyl Laboratories, A300-BL4000) or anti-SOX1 (R&D, AF3369) antibodies. Up to 10 ng of

DNA was used for the library construction using NEBNext Ultra II DNA Library Prep Kit (NEB,

E7645). Sequencing was performed on NextSeq 500 (Illumina).

RNA-seq

The procedure was performed as previously described (8, 10).

Lentiviral transduction of genes and CRISPR-Cas9 genome editing

FLAG-NKX2-1 or *FLAG-GFP* open reading frame (ORF) was cloned into pLEX_306 (a gift

from David Root, Addgene plasmid #41391). Cells stably expressing Cas9 were generated by

infection with the lentiCas9-Blast plasmid (Addgene # 52962, a gift from Feng Zhang). sgRNAs

targeting *NKX2-1* or *SOX1* were selected from Brunello library (Table E2) (11). Additional

details are provided in the online supplement.

Immunohistochemistry (IHC)

Immunohistochemical analyses were performed on xenograft tumor specimens, human primary

SCLC tumor specimens and genetic engineered mouse model (GEMM) tumor specimens with

anti-NKX2-1 (1:100; Santa Cruz, sc-53136), anti-SOX1 (R&D, AF3369) or anti-ASCL1 (1:100,

BD, 556604) antibody and detected with the Vectastain ABC kit.

Co-immunoprecipitation and LC-MS/MS

Chromatin fraction was prepared and co-immunoprecipitated with the methods described

previously (12) with modifications. Peptides were detected, isolated, and fragmented to produce

a tandem mass-spectrum of specific fragment ions for each peptide with LTQ Orbitrap Velos Pro

ion-trap mass spectrometer (Thermo Fisher Scientific, Waltham, MA).

Cell proliferation assay

Details are provided in the online supplement.

Xenograft model

The procedure was performed as previously described (ref). All xenograft studies were approved

by the institutional Animal Care and Use Committee at Icahn School of Medicine at Mount Sinai

(IACUC-2018-0021).

Annexin-V staining

Details are provided in the online supplement.

In situ proximity ligation assay (in situ PLA)

In situ PLA was performed with DuoLink in Situ Reagents (Sigma) according to the instructions.

Images were acquired with a ZEISS Axiocam 503 mono confocal microscope.

Mice and tumor initiation

Mice harboring Rb1^{flox}, Trp53^{flox} and Nkx2-1^{flox} alleles have been previously described (13, 14).

Tumors were generated by intratracheal delivery of Ad5CMV-Cre adenovirus (University of

Iowa, Gene Transfer Vector Core) as described (15). All studies were approved by the

Committee for Animal Care at MIT (A-3125–01).

Data analysis

Details are provided in the online supplement.

Statistical Analysis

Details of specific analyses are provided in the figure legends.

Data and materials availability

SCLC cell lines, human primary SCLC tumors, RP and RPN tumors H3K27ac ChIP-seq, Cell

lines and RP and RPN tumors RNA-seq, NKX2-1 and SOX1 ChIP-seq are deposited in NCBI

GEO: GSE183373.

RESULTS

Epigenomic profiling subdivided SCLC-A by differential super-enhancers

Genome-scale chromatin organization of the cell more faithfully reflects their lineage state than

transcriptomics (16). To refine current SCLC classification, we profiled genome-wide H3K27ac

to define SE regions across 16 SCLC cell lines. Unsupervised hierarchical clustering on SEs on

genes for transcriptional regulation with additional public data (4) revealed four subclasses

(Figure 1A), supported by principal component analysis (PCA) (see Figure E1A in the online

data supplement). Our clusters recapitulated aspects of the expression classification. Cluster III

corresponds to SCLC-N (Figure 1A; purple). Cluster IV represents SCLC-P and SCLC-Y

(Figure 1A; blue). Notably, epigenomic clustering distinguished SCLC-A into two distinct

Clusters I and II (Figure 1A; red and green) referred to as SCLC-A1 and SCLC-A2, while

ASCL1 SE is present in 92.9% (13/14) of SCLC-A (Figure E1B).

To understand the difference underpinning SCLC-A1 and SCLC-A2, we identified 140

differentially enriched SEs. Among 110 SEs enriched in SCLC-A1 was *NKX2-1* (Figure 1B),

where H3K27ac signal was largely absent in other subclasses (Figure 1C). The 30 SEs enriched

in SCLC-A2 included *TCF4*, a transcription factor that regulates synaptic plasticity (17) (Figure

1B and Figure E1B). NKX2-1, also known as *TTF1*, a lineage transcription factor essential for

the development and maintenance of lungs, brain, and thyroid, is a frequently amplified lineage-

specific oncogene and a well-known marker to diagnose lung adenocarcinoma (LADC) (18) (19,

20). However, *NKX2-1* is not significantly amplified in human SCLCs (21) nor in the SCLC

lines used (Figure E1C). Nonetheless, SCLC-A1 cell lines harbor *NKX2-1* SE, whereas SCLC-

A2 cell lines do not (Figure 1D and Figure E1D). Local H3K27ac enrichment pattern at the

NKX2-1 locus in SCLC-A1 was almost identical to LADCs with *NKX2-1* amplification (Figure

E1E). In 6 human SCLC specimens, we found one SCLC-P tumor with *POU2F3* SE, and threes

mixed SCLC-A/N tumors with SEs near *ASCL1* and *NEUROD1* loci (Figure E1F). Notably, we

observed two tumors representing SCLC-A1 with high signals at *ASCL1* and *NKX2-1* loci

(Figure 1E).

By immunohistochemistry (IHC) on another 34 human SCLC cohort, 65% (22/34) of

cases were NKX2-1-positive (Figure E2A), consistent with prior studies (22-26). In cell lines,

SCLC-A1 expressed substantially higher levels of NKX2-1 protein than other SCLC subtypes

and equivalent to *NKX2-1*-amplified LADCs (Figure 1F). Furthermore, RNA-seq datasets (5, 27)

confirmed *NKX2-1* is highly expressed in SCLC (Figure E2B), particularly in SCLC-A (Figure

1G), comparable to the levels in LADC (Figure E2C), consistent with a previous report (22).

Nonetheless, non-bimodal distribution of *NKX2-1* expression suggests that the distinction

between SCLC-A1 and A2 is not discernible from expression analysis, perhaps analogous to the

findings that *NKX2-1* is almost universally expressed in LADC yet only amplified in 10-15%,

which we previously characterized as a distinct biological subgroup (19, 28).

NKX2-1 survival dependency is unique to SCLC-A1

A recent study implicated that NKX2-1 promotes SCLC growth and regulates NE differentiation

and anti-apoptosis (29). We deleted *NKX2-1* in SCLC-A1 cell lines (Figure 2A) and found it

significantly suppressed cell growth and increased annexin-V-positive populations (Figure 2C

and E3A). By contrast, ablating NKX2-1 in SCLC cells from other clusters (Figure 2B) had little

effect. Given that acute removal of a master lineage factor in established cancer often leads to a

halt in cell growth (30), these data suggest NKX2-1 does not play a general oncogenic role in all

lineage states but its function is uniquely relevant to SCLC-A1.

We further found a significant decrease in tumor volume for NKX2-1-deleted SCLC-A1

cells in xenograft (Figure 2D and Figure E3B), confirming its essentiality *in vivo*. No obvious

histological differences were observed in the cells that formed small tumors which escaped from

NKX2-1 deletion (Figure E3C). In a short period after NKX2-1 deletion *in vitro*, expression of

NE markers was not significantly altered (Figure E3D). Overall, these data suggest that NKX2-1

is functionally critical for SCLC-A1 *in vitro* and *in vivo*.

Genomic occupancy of *NKX2-1* in SCLC-A1 cells distinct from that in LADC cells

In LADC, NKX2-1 regulates gene expression and shares its cistromes with normal lung

(19); however, SCLCs typically do not express canonical NKX2-1 targets. Given NKX2-1's

roles in subsets of cortical, striatal, and pallidal neurons (31), we speculated its potential role for

neuronal differentiation in SCLC-A1. Therefore, we profiled NKX2-1 cistromes by ChIP-seq in

5 SCLC-A1 cell lines and compared them to that from 3 LADC cell lines. Pairwise correlation

coefficient of all NKX2-1-bound sites and PCA revealed histology-specific binding profiles

(Figure 3A and E4A). Regions 4 (low-intensity signals) and 1 (high-intensity signals) are

common between SCLC-A1 and LADC (Figure 3B), which included the *RUNX1* locus (Figure

3C). Region 3 was unique to LADC that included *LMO3,* an essential mediator of NKX2-1

functions in LADCs (19). Region 2 was unique to SCLC-A1 that included *DCC,* a receptor for

netrin required for axon guidance (32) (Figure 3C). These data suggest NKX2-1 regulates

distinct transcriptional programs across these two lineage states.

Ontologies on 661 SCLC-A1-unique regions (Figure 3D and E4B) were enriched for

neuronal development, differentiation, and structure (Figure 3E). By contrast, genes regulating

neuronal biology were absent in LADC-unique regions (Figure E4C). Motif analysis revealed

SCLC-A1-specific sites are enriched with ASCL2, Olig2, and SOX motifs, suggesting more

association with neuronal TFs (Figure E4D). In contrast, AP-1, Ets, Runx, and Tead-binding

motifs are enriched in LADC-specific regions, consistent with our previous report (19). These

data indicate that contribution of NKX2-1 in regulating pathways for maintenance of neural

differentiation state of SCLC-A1, distinct from its role in normal lung epithelium and LADCs.

NKX2-1 interactome in chromatin of SCLC-A1 cells includes SOX1

NKX2-1 interacts with FOXA1 to co-regulate the expression of a set of genes for lineage control

in LADC (19, 20). Therefore, to identify unique partners that interact with NKX2-1 to exert

distinct transcriptional program in SCLC-A1, we performed co-immunoprecipitation (co-IP)

followed by liquid chromatography tandem mass-spectrometry (LC-MS/MS) in the chromatin

fraction of two SCLC-A1 cell lines ectopically expressing FLAG-NKX2-1 (Figure 4A and

Figure E5A). We found 27 overlapping NKX2-1-interacting TFs in both cell lines (Figure 4B).

We further identified core regulatory circuitries (CRCs) (33) for SCLC-A1 and other

SCLC clusters from our H3K27ac data. Common CRCs included NEUROD1 for SCLC-N,

POU2F3 and ASCL2 for SCLC-P/Y, and TCF4 in SCLC-A2, consistent with our differential SE

analysis (Figure 1B and E5B). Common CRCs in SCLC-A1 included NKX2-1 (Figure 4C) along

with SOX1, NFIB, TCF12 and PBX1 also found in the SCLC-A1 NKX2-1-interactome. Of those,

H3K27ac signal at *SOX1* was highly enriched in SCLC-A1 (Figure 1B and Figure 4D), whereas

NFIB, *TCF12* and *PBX1* SEs were present across SCLC subtypes (Figure E5C). *SOX1* is highly

expressed in SCLC-A (Figure E5D) and positively correlated with *NKX2-1* expression in 185

SCLC tumors and cell lines (Figure E5E). SOX1 protein was also highly expressed specifically

in SCLC-A1 cell lines (Figure E5F).

The interaction between NKX2-1 and SOX1 was confirmed by co-IP followed by

western blot in two SCLC-A1 cell lines (Figure 4E), and further by proximity ligation assay

(PLA) in 4 SCLC-A1 cell lines (Figure 4F and Figure E5G). These data suggest robust protein-

protein interaction between NKX2-1 and SOX1 uniquely in SCLC-A1.

NKX2-1 and SOX1 co-occupy the SCLC-A1 genome to collaborate in neuronal gene

regulation

Next, we profiled genome-wide binding profiles of NKX2-1 and SOX1 in four SCLC-A1 cell

lines and found the majority of SOX1 peaks were bound by NKX2-1 (48.6-67.8%) (Figure 5A).

The 2,652 regions co-bound by NKX2-1 and SOX1 in NCI-H187 were enriched for functions in

neuron differentiation and development, consistent with the aggregate analysis (Figure E6A and

E6B). We defined high-confidence NKX2-1 (5,663) and SOX1 (1,101) binding sites observed in

$\geqq$3 cell lines, and found 604 peaks (54.9% of SOX1 peaks) are overlapped (Figure 5B), which

included the *NKX2-1* locus as well as neuronal genes *NRXN3* (34, 35) and *YWHAZ* (36) (Figure

5C) and ontologically enriched for functions in neuron differentiation and development (Figure

5D). These data suggest the interaction between NKX2-1 and SOX1 elicits specific cellular

identity of SCLC-A1.

Collaboration of NKX2-1 and SOX1 is required to control the neuronal state in SCLC-A1

We evaluated the transcriptional effect of depletion of *SOX1* or *NKX2-1* in NCI-H187 by RNA-

seq. After the abrogation of NKX2-1 or SOX1, 1,121 and 2,499 genes were significantly

downregulated respectively, with 873 genes commonly downregulated, suggesting a substantial

overlap in their transcriptional outputs (Figure 6A). We then queried the differentially co-

downregulated genes with the 2,652 genes with transcriptional start sites (TSSs) within 50 kb of

SOX1-NKX2-1 co-occupied peaks in NCI-H187, which were significantly enriched for NKX2-1

or SOX1 downregulated genes by ssGSEA analysis (Figure 6B). Leading edge analysis on these

genes revealed 289 overlaps, which we found are enriched for neuron differentiation and

development (Figure 6C). Together, these data suggest that NKX2-1 and SOX1 are specifically

co-localized on the SCLC-A1 genome and collaboratively contribute to controlling neuronal

differentiation state. We also found that SOX1 deletion, while not affecting the NKX2-1

expression, led to reduced cell growth (Figure E7A) with increased apoptosis (Figure E7B).

We previously found that expression of Nkx2-1 was critical for global Foxa1/2 binding

pattern in the LADC genome (20). To test if SOX1 is required for NKX2-1-binding to its target

genes in SCLC-A1, we performed NKX2-1 ChIP-seq in NCI-H187 cells upon SOX1 depletion.

Pairwise correlation coefficient of NKX2-1 peaks across cells deleted for SOX1 or control

revealed strong correlation amongst each group (Figure 6D), supported by PCA analysis (Figure

E7C). Differential analysis on NKX2-1-bound sites found 2,067 peaks decreased upon SOX1

deletion (Figure E7D). *k*-means clustering of the differential sites found relatively uniform

reduction of NKX2-1 binding after SOX1 deletion (Figure 6E), where we also found decreased

H3K27ac modification compared to other sites (Figure 6F), suggesting that these NKX2-1-lost

sites are no longer in active state, similar to a recent finding on Sox2 in esophageal squamous

cell cancer model for Klf5 (37). The data suggest that loss of SOX1 releases NKX2-1 from co-

occupied loci in SCLC-A1.

***Nkx2-1* is essential for development of SCLC-A1 tumors in an autochthonous mouse model**

The *Rb1^{fl/fl}Trp53^{fl/fl}* (RP) is commonly used SCLC models that recapitulates the biology of

ASCL1-positive/classic human SCLCs (13) (Figure 7A). We first profiled H3K27ac to identify

SEs in mSCLC tumors from RP model, where Nkx2-1 is diffusely expressed. We found all 10

tumors from 2 different mice harbored high-level H3K27ac signal around the *Ascl1* locus, but

none harbored SEs at the *Neurod1*, *Yap1* or *Pou2f3* locus, suggesting SCLC-A subtype (Figure

7B and Figure E8A) consistent with requirement of *Ascl1* for tumor formation in this model (38,

39). We found 3 of 10 tumors simultaneously had high-level H3K27ac signals at the *Nkx2-1* and

Sox1 loci, reminiscent of SCLC-A1, while other 7 tumors have neither. This suggests that RP can

adopt an epigenetic state resembling either SCLC-A1 or SCLC-A2. Consistent but ambiguous

expression levels by IHC suggested epigenetic profiles distinguish these lineage classes more

clearly (Figure 7C).

To explore whether NKX2-1 contributes to mSCLC differentiation, we generated

Rb1^{fl/fl}Trp53^{fl/fl}Nkx2-1^{fl/fl} (RPN) mice. We confirmed Nkx2-1 expression was absent in their

tumors, while 80% of them are Ascl1-positive suggesting that they remain representing SCLC-A.

Sox1 staining was only weakly positive in 6/10 cases while 4 other cases, including the two

Ascl1-negative tumors, were negative (Figure 7C). We observed no significant difference in

overall survival between the two genotypes (Figure E8B). Morphologically, all but one mSCLC

tumor from the RP model had classical SCLC morphology (scant cytoplasm, nuclei with fine

chromatin and inconspicuous nucleoli), whereas a subset of tumors in RPN mice (15% in RPN

vs. 4% in RP) exhibited a 'non-classical' histology defined by higher grade/vesicular nuclei by

two expert pathologists (E.L.S. and M.B) (Figure 7D). These suggest that loss of *Nkx2*-1 leads to

stochastic propensity to form non-classical morphology, the while majority of the tumors retain

classical histology representing SCLC-A.

Next, we profiled H3K27ac in 10 mSCLC tumors from four RPN mice (11-13 months

post-initiation, including five classical and five non-classical). None of the tumors from the RPN

mice had high-level H3K27ac signal at the *Neurod1*, *Yap1* or *Pou2f3* loci, whereas most of RPN

tumors (7/10) harbored high-level H3K27ac signals around the *Ascl1* locus (Figure 7B and

Figure E8A). Strikingly, we found signals at the *Nkx2-1* and the *Sox1* loci were absent in all RPN

tumors, suggesting SCLC-A2 subtype (Figure 7B). These data suggest that while the RP model

can develop both SCLC-A1 and SCLC-A2, *Nkx2-1* deletion at tumor initiation precludes SCLC-

A1 development.

Differential transcriptomic analysis from the tumors in RPN and RP mice also revealed

significantly downregulated genes in RPN tumors enriched for nervous system development

(Figure 7E), suggesting that SCLC-A2 is hierarchically closer to the SCLC-N and SCLC-Y/P

subtypes, also supported by the data that variant morphology is more frequent in RPN. To

address whether the absence of Nkx2-1 resulted in distinct tumor evolution in RPN, we profiled

copy numbers of those mSCLCs and found that *Nfib* and *Mycl* loci are frequently amplified in

RP model (6/10) consistent with previous reports; however, these amplifications were absent in

all tumors in RPN model, suggesting the lack of Nkx2-1 predisposed those RPN tumors to

undergo a distinct path leading to a different lineage class of tumors (Figure 7F and Figure E8C).

DISCUSSION

In the current classification, SCLC-A constitutes 70% of SCLCs, while emerging evidence

suggests its heterogeneity. Our study revealed distinctive epigenetic landscapes within SCLC-A

that we designated SCLC-A1 and SCLC-A2, characterized by the presence of *NKX2-1* and *TCF4*

SEs. NKX2-1 is required for the lung development and later specification of the peripheral lung,

but also of bronchial NE cells (40). NKX2-1 is a well-known marker to diagnose LADCs (41),

but it is also frequently expressed in SCLC. As NKX2-1 plays key roles in brain development

(42-45), it has been speculated that NKX2-1 is involved in NE pathways, based on its expression

enriched in SCLC-A (46) and its physical interaction with ASCL1 (29). In our study, while most

SCLC-A expresses NKX2-1 at variable levels, only 48% had *NKX2-1* SE. Importantly, *NKX2-1*

deletion in SCLC-A2 and SCLC-N cells, where NKX2-1 protein is still expressed, had little cell

growth effects, suggesting its relevancy only in SCLC-A1.

Further, our data suggest this function is mediated through interaction with SOX1.

Combinations of master TFs that assemble at tissue-specific cis-regulatory sites control distinct

cellular functions essential for maintaining lineage states in many contexts. Therefore, while the

role of SOX1 had never been described in the lung, it is plausible that their interaction is critical

to retain unique NE state and it further suggests biological plasticity between glandular cell

lineage and pulmonary NE lineage.

The roles of NKX2-1 in carcinogenesis are complex and remain poorly understood.

Nkx2-1 deletion in neoplastic lungs causes loss of pulmonary identity and reversion to a foregut

lineage. Our data suggest that Nkx2-1 contributes to maintaining a specific neural lineage state

that have classical SCLC morphology. The observation that *Nkx2-1* deletion is tolerated at tumor

initiation in the GEMM, but not in established human SCLC-A1 cell lines, provokes a few

potential interpretations. First, the timing of the deletion might be critical in determining the

outcome, as evidenced in other studies on lineage factors (47, 48). Second, as CMV-Cre targets

multiple cell types, *Nkx2-1* deletion may favor origination from certain cell types, thus

influencing tumor evolution, as reflected by the lack of amplification at *Nfib* and *Mycl*. Third,

early/in situ RPN lesions that would have become SCLC-A1 might simply evolve to an

alternative identity due to lack of *Nkx2-1*.

Together, we subdivided the major SCLC-A subtype into SCLC-A1 and SCLC-A2 by

revealing distinct epigenetic landscape in SCLC cell lines, human SCLC tumors, and GEMM.

Collaborative NKX2-1/SOX1 regulation defines the unique lineage state for SCLC-A1. This

study will provide foundation for future identification of vulnerabilities and enable us to develop

higher level of personalized therapeutic approaches to benefit SCLC patients.

Acknowledgements

We thank Hong Cai, Yi Ge, Ying Liu, Peipei Guo, Yifei Sun, Xiaokun Liu, Korey Kam,

Mingzhe Li, Megan Vetter, Tyler Jacks and Karen Yee for helpful discussions and technical

assistance, Aleksandra Wroblewska and Brian D. Brown for providing a lentiviral vector for

cloning sgRNAs; NextSeq Sequencing Facility of the Department of Oncological Sciences at

Icahn School of Medicine at Mount Sinai (ISMMS), Saboor Hekmaty, Gayatri Panda and Ravi

Sachidanandam for sequencing assistance. The authors thank Taplin Mass Spectrometry Facility

at Harvard Medical School, the Biorepository and Pathology Core Facility and Center for

Comparative Medicine and Surgery at ISMMS.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; 69: 7-34.

2. Poirier JT, Gardner EE, Connis N, Moreira AL, de Stanchina E, Hann CL, Rudin CM. DNA methylation in small cell lung cancer defines distinct disease subtypes and correlates with high expression of EZH2. *Oncogene* 2015; 34: 5869-5878.

3. McColl K, Wildey G, Sakre N, Lipka MB, Behtaj M, Kresak A, Chen Y, Yang M, Velcheti V, Fu P, Dowlati A. Reciprocal expression of INSM1 and YAP1 defines subgroups in small cell lung cancer. *Oncotarget* 2017; 8: 73745-73756.

4. Huang YH, Klingbeil O, He XY, Wu XLS, Arun G, Lu B, Somerville TDD, Milazzo JP, Wilkinson JE, Demerdash OE, Spector DL, Egeblad M, Shi JW, Vakoc CR. POU2F3 is a master regulator of a tuft cell-like variant of small cell lung cancer. *Gene Dev* 2018; 32: 915-928.

5. George J, Lim JS, Jang SJ, Cun Y, Ozretic L, Kong G, Leenders F, Lu X, Fernandez-Cuesta L, Bosco G, Muller C, Dahmen I, Jahchan NS, Park KS, Yang D, Karnezis AN, Vaka D, Torres A, Wang MS, Korbel JO, Menon R, Chun SM, Kim D, Wilkerson M, Hayes N, Engelmann D, Putzer B, Bos M, Michels S, Vlasic I, Seidel D, Pinther B, Schaub P, Becker C, Altmuller J, Yokota J, Kohno T, Iwakawa R, Tsuta K, Noguchi M, Muley T, Hoffmann H, Schnabel PA, Petersen I, Chen Y, Soltermann A, Tischler V, Choi CM, Kim YH, Massion PP, Zou Y, Jovanovic D, Kontic M, Wright GM, Russell PA, Solomon B, Koch I, Lindner M, Muscarella LA, la Torre A, Field JK, Jakopovic M, Knezevic J, Castanos-Velez E, Roz L, Pastorino U, Brustugun OT, Lund-Iversen M, Thunnissen E, Kohler J, Schuler M, Botling J, Sandelin M, Sanchez-Cespedes M, Salvesen HB, Achter V, Lang U, Bogus M, Schneider PM, Zander T, Ansen S, Hallek M, Wolf J, Vingron M, Yatabe Y, Travis WD, Nurnberg P, Reinhardt C, Perner S, Heukamp L, Buttner R, Haas SA, Brambilla E, Peifer M, Sage J, Thomas RK. Comprehensive genomic profiles of small cell lung cancer. *Nature* 2015; 524: 47-53.

6. Rudin CM, Poirier JT, Byers LA, Dive C, Dowlati A, George J, Heymach JV, Johnson JE, Lehman JM, MacPherson D, Massion PP, Minna JD, Oliver TG, Quaranta V, Sage J, Thomas RK, Vakoc CR, Gazdar AF. Molecular subtypes of small cell lung cancer: a synthesis of human and mouse model data. *Nat Rev Cancer* 2019; 19: 289-297.

7. Gay CM, Stewart CA, Park EM, Diao L, Groves SM, Heeke S, Nabet BY, Fujimoto J, Solis LM, Lu W, Xi Y, Cardnell RJ, Wang Q, Fabbri G, Cargill KR, Vokes NI, Ramkumar K, Zhang B, Della Corte CM, Robson P, Swisher SG, Roth JA, Glisson BS, Shames DS, Wistuba, II, Wang J, Quaranta V, Minna J, Heymach JV, Byers LA. Patterns of transcription factor programs and immune pathway activation define four major subtypes of SCLC with distinct therapeutic vulnerabilities. *Cancer Cell* 2021; 39: 346-360 e347.

8. Patel AS, Yoo S, Kong R, Sato T, Sinha A, Karam S, Bao L, Fridrikh M, Emoto K, Nudelman G, Powell CA, Beasley MB, Zhu J, Watanabe H. Prototypical oncogene family Myc defines unappreciated distinct lineage states of small cell lung cancer. *Sci Adv* 2021; 7.

9. Iwafuchi-Doi M, Zaret KS. Pioneer transcription factors in cell reprogramming. *Genes Dev* 2014; 28: 2679-2692.

10. Sato T, Yoo S, Kong R, Sinha A, Chandramani-Shivalingappa P, Patel A, Fridrikh M, Nagano O, Masuko T, Beasley MB, Powell CA, Zhu J, Watanabe H. Epigenomic profiling discovers trans-lineage SOX2 partnerships driving tumor heterogeneity in lung squamous cell carcinoma. *Cancer Res* 2019.

11. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat Biotechnol* 2016; 34: 184-191.

12. Watanabe H, Ma Q, Peng S, Adelmant G, Swain D, Song W, Fox C, Francis JM, Pedamallu CS, DeLuca DS, Brooks AN, Wang S, Que J, Rustgi AK, Wong KK, Ligon KL, Liu XS, Marto JA, Meyerson M, Bass

AJ. SOX2 and p63 colocalize at genetic loci in squamous cell carcinomas. *J Clin Invest* 2014; 124: 1636-1645.

13. Meuwissen R, Linn SC, Linnoila RI, Zevenhoven J, Mooi WJ, Berns A. Induction of small cell lung cancer by somatic inactivation of both Trp53 and Rb1 in a conditional mouse model. *Cancer Cell* 2003; 4: 181-189.

14. Kusakabe T, Kawaguchi A, Hoshi N, Kawaguchi R, Hoshi S, Kimura S. Thyroid-specific enhancer-binding protein/NKX2.1 is required for the maintenance of ordered architecture and function of the differentiated thyroid. *Mol Endocrinol* 2006; 20: 1796-1809.

15. DuPage M, Dooley AL, Jacks T. Conditional mouse lung cancer models using adenoviral or lentiviral delivery of Cre recombinase. *Nature protocols* 2009; 4: 1064-1072.

16. Zheng H, Xie W. The role of 3D genome organization in development and cell differentiation. *Nat Rev Mol Cell Biol* 2019; 20: 535-550.

17. Kennedy AJ, Rahn EJ, Paulukaitis BS, Savell KE, Kordasiewicz HB, Wang J, Lewis JW, Posey J, Strange SK, Guzman-Karlsson MC, Phillips SE, Decker K, Motley ST, Swayze EE, Ecker DJ, Michael TP, Day JJ, Sweatt JD. Tcf4 Regulates Synaptic Plasticity, DNA Methylation, and Memory Function. *Cell Rep* 2016; 16: 2666-2685.

18. Campbell JD, Alexandrov A, Kim J, Wala J, Berger AH, Pedamallu CS, Shukla SA, Guo G, Brooks AN, Murray BA, Imielinski M, Hu X, Ling S, Akbani R, Rosenberg M, Cibulskis C, Ramachandran A, Collisson EA, Kwiatkowski DJ, Lawrence MS, Weinstein JN, Verhaak RG, Wu CJ, Hammerman PS, Cherniack AD, Getz G, Cancer Genome Atlas Research N, Artyomov MN, Schreiber R, Govindan R, Meyerson M. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat Genet* 2016; 48: 607-616.

19. Watanabe H, Francis JM, Woo MS, Etemad B, Lin W, Fries DF, Peng S, Snyder EL, Tata PR, Izzo F, Schinzel AC, Cho J, Hammerman PS, Verhaak RG, Hahn WC, Rajagopal J, Jacks T, Meyerson M. Integrated cistromic and expression analysis of amplified NKX2-1 in lung adenocarcinoma identifies LMO3 as a functional transcriptional target. *Genes Dev* 2013; 27: 197-210.

20. Snyder EL, Watanabe H, Magendantz M, Hoersch S, Chen TA, Wang DG, Crowley D, Whittaker CA, Meyerson M, Kimura S, Jacks T. Nkx2-1 represses a latent gastric differentiation program in lung adenocarcinoma. *Mol Cell* 2013; 50: 185-199.

21. Peifer M, Fernandez-Cuesta L, Sos ML, George J, Seidel D, Kasper LH, Plenker D, Leenders F, Sun R, Zander T, Menon R, Koker M, Dahmen I, Muller C, Di Cerbo V, Schildhaus HU, Altmuller J, Baessmann I, Becker C, de Wilde B, Vandesompele J, Bohm D, Ansen S, Gabler F, Wilkening I, Heynck S, Heuckmann JM, Lu X, Carter SL, Cibulskis K, Banerji S, Getz G, Park KS, Rauh D, Grutter C, Fischer M, Pasqualucci L, Wright G, Wainer Z, Russell P, Petersen I, Chen Y, Stoelben E, Ludwig C, Schnabel P, Hoffmann H, Muley T, Brockmann M, Engel-Riedel W, Muscarella LA, Fazio VM, Groen H, Timens W, Sietsma H, Thunnissen E, Smit E, Heideman DA, Snijders PJ, Cappuzzo F, Ligorio C, Damiani S, Field J, Solberg S, Brustugun OT, Lund-Iversen M, Sanger J, Clement JH, Soltermann A, Moch H, Weder W, Solomon B, Soria JC, Validire P, Besse B, Brambilla E, Brambilla C, Lantuejoul S, Lorimier P, Schneider PM, Hallek M, Pao W, Meyerson M, Sage J, Shendure J, Schneider R, Buttner R, Wolf J, Nurnberg P, Perner S, Heukamp LC, Brindle PK, Haas S, Thomas RK. Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat Genet* 2012; 44: 1104-1110.

22. Horie M, Miyashita N, Mattsson JSM, Mikami Y, Sandelin M, Brunnstrom H, Micke P, Nagase T, Saito A. An integrative transcriptome analysis reveals a functional role for thyroid transcription factor-1 in small cell lung cancer. *J Pathol* 2018; 246: 154-165.

23. Misch D, Blum T, Boch C, Weiss T, Crolow C, Griff S, Mairinger T, Bauer TT, Kollmeier J. Value of thyroid transcription factor (TTF)-1 for diagnosis and prognosis of patients with locally advanced or metastatic small cell lung cancer. *Diagn Pathol* 2015; 10: 21.

24. Miyauchi E, Motoi N, Ono H, Ninomiya H, Ohyanagi F, Nishio M, Okumura S, Ichinose M, Ishikawa Y.
Distinct Characteristics of Small Cell Lung Cancer Correlate With Central or Peripheral Origin:
Subtyping Based on Location and Expression of Transcription Factor TTF-1. *Medicine (Baltimore)*
2015; 94: e2324.
25. Wu M, Wang B, Gil J, Sabo E, Miller L, Gan L, Burstein DE. p63 and TTF-1 immunostaining. A useful
marker panel for distinguishing small cell carcinoma of lung from poorly differentiated
squamous cell carcinoma of lung. *Am J Clin Pathol* 2003; 119: 696-702.
26. Ordonez NG. Value of thyroid transcription factor-1 immunostaining in distinguishing small cell lung
carcinomas from other small cell carcinomas. *Am J Surg Pathol* 2000; 24: 1217-1223.
27. Jiang L, Huang J, Higgs BW, Hu Z, Xiao Z, Yao X, Conley S, Zhong H, Liu Z, Brohawn P, Shen D, Wu S,
Ge X, Jiang Y, Zhao Y, Lou Y, Morehouse C, Zhu W, Sebastian Y, Czapiga M, Oganesyan V, Fu H,
Niu Y, Zhang W, Streicher K, Tice D, Zhao H, Zhu M, Xu L, Herbst R, Su X, Gu Y, Li S, Huang L, Gu J,
Han B, Jallal B, Shen H, Yao Y. Genomic Landscape Survey Identifies SRSF1 as a Key Oncodriver in
Small Cell Lung Cancer. *PLoS Genet* 2016; 12: e1005895.
28. Weir BA, Woo MS, Getz G, Perner S, Ding L, Beroukhim R, Lin WM, Province MA, Kraja A, Johnson LA,
Shah K, Sato M, Thomas RK, Barletta JA, Borecki IB, Broderick S, Chang AC, Chiang DY, Chirieac
LR, Cho J, Fujii Y, Gazdar AF, Giordano T, Greulich H, Hanna M, Johnson BE, Kris MG, Lash A, Lin L,
Lindeman N, Mardis ER, McPherson JD, Minna JD, Morgan MB, Nadel M, Orringer MB, Osborne
JR, Ozenberger B, Ramos AH, Robinson J, Roth JA, Rusch V, Sasaki H, Shepherd F, Sougnez C,
Spitz MR, Tsao MS, Twomey D, Verhaak RG, Weinstock GM, Wheeler DA, Winckler W, Yoshizawa
A, Yu S, Zakowski MF, Zhang Q, Beer DG, Wistuba, II, Watson MA, Garraway LA, Ladanyi M,
Travis WD, Pao W, Rubin MA, Gabriel SB, Gibbs RA, Varmus HE, Wilson RK, Lander ES, Meyerson
M. Characterizing the cancer genome in lung adenocarcinoma. *Nature* 2007; 450: 893-898.
29. Hokari S, Tamura Y, Kaneda A, Katsura A, Morikawa M, Murai F, Ehata S, Tsutsumi S, Ishikawa Y,
Aburatani H, Kikuchi T, Miyazono K, Koinuma D. Comparative analysis of TTF-1-binding DNA
regions in small cell lung cancer and non-small cell lung cancer. *Mol Oncol* 2019.
30. Heinz S, Romanoski CE, Benner C, Glass CK. The selection and function of cell type-specific enhancers.
Nat Rev Mol Cell Biol 2015; 16: 144-154.
31. Magno L, Barry C, Schmidt-Hieber C, Theodotou P, Hausser M, Kessaris N. NKX2-1 Is Required in the
Embryonic Septum for Cholinergic System Development, Learning, and Memory. *Cell Rep* 2017;
20: 1572-1584.
32. Liu Y, Bhowmick T, Liu Y, Gao X, Mertens HDT, Svergun DI, Xiao J, Zhang Y, Wang JH, Meijers R.
Structural Basis for Draxin-Modulated Axon Guidance and Fasciculation by Netrin-1 through DCC.
Neuron 2018; 97: 1261-1267 e1264.
33. Saint-Andre V, Federation AJ, Lin CY, Abraham BJ, Reddy J, Lee TI, Bradner JE, Young RA. Models of
human core transcriptional regulatory circuitries. *Genome Res* 2016; 26: 385-396.
34. Reissner C, Runkel F, Missler M. Neurexins. *Genome Biol* 2013; 14: 213.
35. Sudhof TC. Neuroligins and neurexins link synaptic function to cognitive disease. *Nature* 2008; 455:
903-911.
36. Toma C, Torrico B, Hervas A, Valdes-Mas R, Tristan-Noguero A, Padillo V, Maristany M, Salgado M,
Arenas C, Puente XS, Bayes M, Cormand B. Exome sequencing in multiplex autism families
suggests a major role for heterozygous truncating mutations. *Mol Psychiatry* 2014; 19: 784-790.
37. Wu Z, Zhou J, Zhang X, Zhang Z, Xie Y, Liu JB, Ho ZV, Panda A, Qiu X, Cejas P, Canadas I, Akarca FG,
McFarland JM, Nagaraja AK, Goss LB, Kesten N, Si L, Lim K, Liu Y, Zhang Y, Baek JY, Liu Y, Patil DT,
Katz JP, Hai J, Bao C, Stachler M, Qi J, Ishizuka JJ, Nakagawa H, Rustgi AK, Wong KK, Meyerson M,
Barbie DA, Brown M, Long H, Bass AJ. Reprogramming of the esophageal squamous carcinoma
epigenome by SOX2 promotes ADAR1 dependence. *Nat Genet* 2021; 53: 881-894.

38. Borromeo MD, Savage TK, Kollipara RK, He M, Augustyn A, Osborne JK, Girard L, Minna JD, Gazdar AF,
Cobb MH, Johnson JE. ASCL1 and NEUROD1 Reveal Heterogeneity in Pulmonary Neuroendocrine
Tumors and Regulate Distinct Genetic Programs. *Cell Rep* 2016; 16: 1259-1272.
39. Olsen RR, Ireland AS, Kastner DW, Groves SM, Spainhower KB, Pozo K, Kelenis DP, Whitney CP,
Guthrie MR, Wait SJ, Soltero D, Witt BL, Quaranta V, Johnson JE, Oliver TG. ASCL1 represses a
SOX9(+) neural crest stem-like state in small cell lung cancer. *Genes Dev* 2021; 35: 847-869.
40. La Rosa S, Chiaravalli AM, Placidi C, Papanikolaou N, Cerati M, Capella C. TTF1 expression in normal
lung neuroendocrine cells and related tumors: immunohistochemical study comparing two
different monoclonal antibodies. *Virchows Arch* 2010; 457: 497-507.
41. Yamaguchi T, Hosono Y, Yanagisawa K, Takahashi T. NKX2-1/TTF-1: an enigmatic oncogene that
functions as a double-edged sword for cancer cell survival and progression. *Cancer Cell* 2013; 23:
718-723.
42. Puelles L, Kuwana E, Puelles E, Bulfone A, Shimamura K, Keleher J, Smiga S, Rubenstein JL. Pallial and
subpallial derivatives in the embryonic chick and mouse telencephalon, traced by the expression
of the genes Dlx-2, Emx-1, Nkx-2.1, Pax-6, and Tbr-1. *J Comp Neurol* 2000; 424: 409-438.
43. Flames N, Pla R, Gelman DM, Rubenstein JL, Puelles L, Marin O. Delineation of multiple subpallial
progenitor domains by the combinatorial expression of transcriptional codes. *J Neurosci* 2007;
27: 9682-9695.
44. Sussel L, Marin O, Kimura S, Rubenstein JL. Loss of Nkx2.1 homeobox gene function results in a
ventral to dorsal molecular respecification within the basal telencephalon: evidence for a
transformation of the pallidum into the striatum. *Development* 1999; 126: 3359-3370.
45. Minocha S, Valloton D, Ypsilanti AR, Fiumelli H, Allen EA, Yanagawa Y, Marin O, Chedotal A, Hornung
JP, Lebrand C. Nkx2.1-derived astrocytes and neurons together with Slit2 are indispensable for
anterior commissure formation. *Nat Commun* 2015; 6: 6887.
46. Cardnell RJ, Li L, Sen T, Bara R, Tong P, Fujimoto J, Ireland AS, Guthrie MR, Bheddah S, Banerjee U,
Kalu NN, Fan YH, Dylla SJ, Johnson FM, Wistuba, II, Oliver TG, Heymach JV, Glisson BS, Wang J,
Byers LA. Protein expression of TTF1 and cMYC define distinct molecular subgroups of small cell
lung cancer with unique vulnerabilities to aurora kinase inhibition, DLL3 targeting, and other
targeted therapies. *Oncotarget* 2017; 8: 73419-73432.
47. Camolotto SA, Pattabiraman S, Mosbruger TL, Jones A, Belova VK, Orstad G, Streiff M, Salmond L,
Stubben C, Kaestner KH, Snyder EL. FoxA1 and FoxA2 drive gastric differentiation and suppress
squamous identity in NKX2-1-negative lung cancer. *Elife* 2018; 7.
48. Zewdu R, Mehrabad EM, Ingram K, Fang P, Gillis KL, Camolotto SA, Orstad G, Jones A, Mendoza MC,
Spike BT, Snyder EL. An NKX2-1/ERK/WNT feedback loop modulates gastric identity and
response to targeted therapy in lung adenocarcinoma. *Elife* 2021; 10.

Figure legends

Figure 1. Epigenomic profiling subdivided SCLC-A by differential super-enhancers

(**A**). Unsupervised hierarchical clustering of 25 SCLC cell lines using SE signals near

transcriptional regulator genes. The color scale indicates the intensity of SE signals normalized

to linear scores using the sum of squares of the values. Each row represents a SE locus (747 loci).

The color shades on the dendrogram indicate four novel SE clusters, the top red columns display

the presence of SE near the *NKX2-1* locus, and the bottom column colors denote the consensus

expression clusters. (**B**). Volcano plot showing 140 differentially enriched SEs (110 enriched in

SCLC-A1 and 30 enriched in SCLC-A2) at thresholds of absolute fold change ≥ 5 and FDR $\leq$

0.25 depicted as dotted lines. (**C**). Genome view tracks of H3K27ac signal at the *NKX2-1* locus

in four clusters of SCLC cell lines. Normalized scales are shown in Figure E1B. (**D**). SE plots

with all enhancers rank-ordered by H3K27ac signals in DMS 79, NCI-H187, NCI-H209 and

NCI-H345. Dotted lines depict tangential cutoffs to define SEs. (**E**). Genome view tracks of

H3K27ac signals at the *ASCL1*, *NEUROD1*, *POU2F3*, *YAP1* and *NKX2-1* loci in 2 human SCLC

specimens of SCLC-A subtype. (**F**). Protein expression of NKX2-1, ASCL1 and vinculin as a

loading control in SCLC cell lines from different clusters. *NKX2-1*-amplified adenocarcinoma

cell lines NCI-H3122 and NCI-H2087 are shown as reference. (**G**). Box and jitter plots showing

the distribution of *NKX2-1* mRNA expression in SCLC-A, SCLC-N, SCLC-P, and SCLC-Y

subtypes of 135 primary SCLC tumors.

Figure 2. *NKX2-1* survival dependency is unique to SCLC-A1

(**A**). *(Top)* Immunoblots showing protein expression of NKX2-1 and vinculin as a loading

control in SCLC-A1 cell lines with three *NKX2-1* sgRNAs, or two non-target (NT) sgRNAs.

P a g e | 23

(Bottom) Cell growth curve for SCLC-A1 cell lines infected with *NKX2-1* sgRNAs or NT

sgRNAs. Data are shown as mean ± SD (n=3); *P < 0.0001 by two-sided two-way ANOVA. (**B**).

(Top) Immunoblots showing protein expression of NKX2-1 and vinculin as a loading control in

SCLC-A2 (NCI-H2171, NCI-H69) and SCLC-N (NCI-H1963) cell lines with *NKX2-1* sgRNAs,

or non-target (NT) sgRNAs. *(Bottom)* Cell growth curve for cell lines infected with *NKX2-1*

sgRNA or NT sgRNAs. Data are shown as mean ± SD (n=3); *P < 0.0001 by two-sided two-way

ANOVA. (**C**). Bar plots showing percentage of apoptotic NCI-H187 cells infected with NT

sgRNA and *NKX2-1* sgRNA, measured by annexin V staining. Means ± SD of three biological

replicates. *P < 0.0001, t test. Representative data for NT sg1 and NKX2-1 sg1 can be found in

Figure E2A. (**D**). (*Left*) Tumor growth curve for xenograft tumors of NCI-H187 cells infected

with *NKX2-1* sg1 or NT sg1 *in vivo*. Mean±SD of four biological replicates. *P < 0.0001, t test.

Xenograft tumors were resected at 3 weeks after inoculation. (*Right*) Images of resected

xenograft tumors of NCI-H209 cells infected with NT sg1 or *NKX2-1* sg1 *in vivo* at day 23.

Figure 3. Genomic occupancy of *NKX2-1* in SCLC-A1 cells distinct from that in LADC

cells

(**A**). Correlation matrix depicting pairwise comparisons of identified NKX2-1 binding peaks in

the 5 SCLC-A1 cell lines and in 3 LADC cell lines. Color scale represents degree of Pearson's

correlation coefficient. (**B**). Heatmap showing signals from NKX2-1 ChIP-seq data in the 5

SCLC-A1 cell lines and 3 LUAC cell lines. ChIP signal intensity is shown by red shading. Y-

axis contains all peaks that are bound by NKX2-1 in any of SCLC-A1 and LUAC (n=37,198).

Peaks are divided into 4 regions by *k-means* (k=4) clustering. (**C**). NKX2-1 ChIP-seq signals in

SCLC-A1 and LADC cell lines at representative loci (*RUNX1*; cluster 1, *LMO3*; cluster 3 and

DCC; cluster 2). Red: SCLC-A1 cell lines. Black: LADC cell lines. (**D**). Heatmap showing 2,671

differentially NKX2-1 bound regions (fold change $\geq 2^5$ and FDR $\leq 1x10^{-5}$), 2,010 enriched in

LADC and 661 enriched in SCLC-A1, between 5 SCLC-A1 cell lines and 3 LUAC cell lines. (**E**).

Enriched ontology by GREAT analyses for regions differentially bound by NKX2-1 in 5 SCLC-

A1 compared to LADCs.

Figure 4. NKX2-1 interactome in chromatin of SCLC-A1 cells includes SOX1

(**A**). Experimental strategy to identify binding partners of NKX2-1 in SCLC-A1. Chromatin

prepared from the nuclear fraction of SCLC-A1 cell lines NCI-H187 and NCI-H1436 stably

expressing FLAG-NKX2-1 was solubilized using micrococcal nuclease digestion. Co-

immunoprecipitated proteins with anti-FLAG antibody from solubilized chromatin were

identified by LC-MS/MS (see Data file S2 for the complete list). (**B**). Venn diagram showing the

number of transcription factors identified in each of the 2 SCLC-A1 cell lines and the 27

overlaps found in both cell lines. The proteins indicated in yellow color were also found in CRC

analysis in Figure 4C. (**C**). Core transcriptional regulatory circuitries (CRCs) commonly

identified in 8 SCLC-A1 cell lines. The genes in yellow color were also found in LC-MS/MS

analysis in Figure 4B. (**D**). Genome view tracks of H3K27ac signal at the *SOX1* locus in four

clusters of SCLC cell lines. (**E**). Immunoblots with anti-SOX1 antibody on endogenous co-

immunoprecipitates by anti-NKX2-1 antibody in SCLC-A1 (DMS 79 and NCI-H187) cells. (**F**).

In situ proximity ligation assay signals (red fluorescence) targeting endogenous SOX1 and

NKX2-1 in NCI-H841, NCI-H187, NCI-H209 and DMS 79 cells. Nuclei are counterstained with

DAPI (blue). See Figure E4G for additional control.

Figure 5. NKX2-1 and SOX1 co-occupy the SCLC-A1 genome to collaborate in gene

regulation

(**A**). Heatmap showing k-means clustering (*k*=4) of signals at the common 4,051 loci from

NKX2-1 (red) and SOX1 (blue) ChIP-seq data in the 4 SCLC-A1 cell lines. ChIP signal intensity

is shown by shading for all NKX2-1 and SOX1 occupied loci. (**B**). Venn diagram showing the

overlap of NKX2-1 common binding sites and SOX1 common binding sites present in at least 3

SCLC-A1 cell lines. (**C**). Genome view tracks of NKX2-1 (red) and SOX1 (blue) ChIP-seq

signals in four SCLC-A1 (DMS 79, NCI-H187, NCI-H209 and NCI-H345) cell lines and 3

LADC cell lines at the representative NKX2-1-SOX1 co-occupied loci (*NRXN3* and *YWHAZ*;

SCLC-A1 unique*, NKX2-1*; SCLC-A1/LADC common). (**D**). Enriched ontology by GREAT

analyses for regions in 604 overlapping high-confidence NKX2-1-SOX1 binding sites.

Figure 6. Collaboration of NKX2-1 and SOX1 is required to control the neuronal state in

SCLC-A1

(**A**). Venn diagram showing the overlap of downregulated genes after depletion of NKX2-1 (fold

change > 1.4 and adjusted P < 0.05) and SOX1 (fold change > 2 and adjusted P < 0.05) in NCI-

H187 cells. (**B**). Gene Set Enrichment Analysis (GSEA) of genes associated with 2,652 NKX2-

1-SOX1 co-occupied loci for expression data upon repression of NKX2-1 (*left*) or SOX1 (*right*)

in NCI-H187 cells, with their leading-edge genes shaded. Enrichment was significant for both

(NES= -2.29, P < 0.001 for NKX2-1 and NES= -2.54, P < 0.001 for SOX1). (**C**). Enriched gene

ontologies (MSigDB, biological process) for 289 common genes in leading-edge genes for both

NKX2-1 and SOX1 deletions. (**D**). Correlation matrix depicting pairwise comparisons of

identified NKX2-1 binding regions in three *SOX1* sgRNAs and three control cells (two non-

target (NT) sgRNAs, one parental cell) of NCI-H187 cell line. Color scale represents degree of

Pearson's correlation coefficient. (**E**). Heatmap showing k-means clustering (*k*=4) of 2,067

NKX2-1 ChIP-seq loci in NCI-H187 sgSOX1 compared to control cells with reduced signal

intensity. ChIP-seq signal intensity is shown by color shading. (**F**). Average H3K27ac ChIP-seq

signals on NKX2-1 reduced binding sites and NKX2-1 non-reduced binding sites defined in

Figure E6D for NCI-H187 parental, NT sg1, SOX1 sg2 and SOX1 sg3 cells.

Figure 7. *Nkx2-1* is essential for development of SCLC-A1 tumors in an autochthonous

mouse model

(**A**). Genetically engineered mouse models of SCLCs. $Rb1^{flox/flox}$; $Trp53^{flox/flox}$ (RP) mice or

$Rb1^{flox/flox}$; $Trp53^{flox/flox}$, $Nkx2\text{-}1^{flox/flox}$ (RPN) mice were transduced with Adeno-CMV-*Cre* to

initiate SCLC. (**B**). Genome view tracks at the Nkx2-1, Sox1, and Ascl1 loci for H3K27ac ChIP-

seq signals in 10 SCLC tumors from RP mice and 10 SCLC tumors from RPN mice. * indicates

3 tumors that have high-level signal at both *Nkx2-1* and *Sox1* loci. The arrowheads on *Nkx2-*

1 locus indicate the loxP sites in RPN mice**. (C**). Representative immune-staining images for

Nkx2-1, Sox1, and Ascl1 on SCLC tumors from RP and RPN mice. Scale bars, 25μm. (**D**).

Representative hematoxylin and eosin (H&E) images of SCLC tumors from RP and RPN mice.

Scale bars, 2.5mm and 50μm, respectively. (**E**). Enriched gene ontologies (MSigDB, biological

process) for the differentially downregulated genes in RPN tumors compared to RP tumors. (**F**).

Average copy number profiles in RP and RPN models. Arrows indicate the *Nfib* and *Mycl* loci

that are specifically amplified in RP tumors (individual copy number profiles at those loci are

available in Figure E8C).

Figure 1

Figure 2

Figure 3

Figure 4

A

SCLC-A1

FLAG-NKX2-1

Nuclear Preparation

Chromatin Preparation

Micrococcal Nuclease Digestion

Immunoprecipitation

LC-MS/MS analysis

B

LC-MS/MS analysis

H187   H1436

48   27   50

SOX1, ADNP2, NFIB, ZBTB25, ZNF148
AHDC1, KDM5B, ZNF292, CREB1, ST18
ZBTB33, ZBTB40, RREB1, KDM2A
CUX1, TCF12, CASZ1, ZNF644, AKAP8
HMBOX1, ZNF516, KDM2B, ARID2
PBX1, TCF20, SKI, ZNF592

C

SCLC-A1
Core Transcriptional Regulatory Circuitry (CRC)

SE	FOXJ3
SE	NKX2-1
SE	FOXA1
SE	ELF3
SE	SOX13
SE	ZBTB7C
SE	SOX1
SE	ETV6
SE	TCF12
SE	HES1
SE	SP3
SE	ETS2
SE	NFIB
SE	TGIF1
SE	HOXD8
SE	ZNF524
SE	PBX1
SE	TP73
SE	RUNX1
SE	ZNF219
SE	TEAD1
SE	POU3F2
SE	ATF4
SE	RFX2
SE	BCL6
SE	NR2F2
SE	E2F2

D

Chr13   H3K27ac

112,700kb   112,720kb   112,740kb

Cluster I (SCLC-A1): H1436, COR-L88, H510A, H209, H128, H345, DMS 79, H187

Cluster II (SCLC-A2): SHP-77, H146, H69, H2171, H2081, H889

Cluster III (SCLC-N): H1963, HCC33, H446, H524, H82

Cluster IV (SCLC-P/Y): H1048, H196, H841, H211, COR-L311, H526

SOX1

E

Input   IP

IgG   NKX2-1

DMS 79   H187   DMS 79   H187   DMS 79   H187

NKX2-1

SOX1

F

| SCLC-P/Y H841 | SCLC-A1 DMS 79 | SCLC-A1 H209 | SCLC-A1 H187 |

SOX1-NKX2-1 Interaction (PLA)

Merged with DAPI

Figure 5

Figure 6

Figure 7

A

Adeno CMV-Cre → Intratracheal instillation

Rb1flox/flox; Trp53flox/flox (RP)

Adeno CMV-Cre → Intratracheal instillation

Rb1flox/flox; Trp53flox/flox; Nkx2-1flox/flox (RPN)

B H3K27ac

chr12 | chr8 | chr10

C

RP (Tumor 5) | RPN (Tumor 5)

Nkx2-1

Sox1

Ascl1

D H&E

RP

RPN

E Downregulated genes in RPN

Central Nervous System Development
Cell Cell Signaling
Nuclear Chromosome
Animal Organ Morphogenesis
Postsynapse
Epithelium Development
Behavior
Locomotory Behavior
Renal System Development
Chromosome

-log$_{10}$ (p-Value)

F

Nfib *Mycl*

RP

RPN

chr 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

Log2