1 **Climate biogeography of *Arabidopsis thaliana:* linking distribution models and individual**
2 **variation**
3

4 Christina Yim[1], Emily S. Bellis[1,2], Victoria L. DeLeo[1], Diana Gamba[1], Robert Muscarella[3],
5 Jesse R. Lasky[1,*]
6
7 [1]Department of Biology
8 Pennsylvania State University
9 University Park
10 USA
11
12 [2]Department of Computer Science
13 Arkansas State University
14 Jonesboro
15 USA
16
17 [3]Plant Ecology and Evolution
18 Evolutionary Biology Centre
19 Uppsala University
20 Uppsala
21 Sweden
22
23 *Corresponding Author
24 lasky@psu.edu

25
26 **Biosketch**
27 The research team is interested in understanding how individual-level processes influence
28 evolutionary and ecological dynamics at continental scales.
29
30 **Author contributions**
31 All authors contributed to study design. CY, ESB, VLD, DG, and JRL conducted analyses. CY
32 and JRL led the writing with contributions from all authors.

Abstract

AIM

Patterns of individual variation are key to testing hypotheses about the mechanisms underlying biogeographic patterns. However, it is challenging to gather data on individual-level variation at large spatial scales. Model organisms are potentially important systems for biogeographical studies, given the available range-wide natural history collections, and the importance of providing biogeographical context to their genetic and phenotypic diversity.

LOCATION
Global

TAXON
Arabidopsis thaliana ("Arabidopsis")

METHODS
We fit occurrence records to climate data, and then projected the distribution of Arabidopsis under last glacial maximum, current, and future climates. We confronted model predictions with individual performance measured on 2,194 herbarium specimens, and we asked whether predicted suitability was associated with life-history and genomic variation measured on ~900 natural accessions.

RESULTS
The most important climate variables constraining the Arabidopsis distribution were winter cold in northern and high elevation regions and summer heat in southern regions. Herbarium specimens from regions with lower habitat suitability in both northern and southern regions were smaller, supporting the hypothesis that the distribution of Arabidopsis is constrained by climate-associated factors. Climate anomalies partly explained interannual variation in herbarium specimen size, but these did not closely correspond to local limiting factors identified in the distribution model. Late-flowering genotypes were absent from the lowest suitability regions, suggesting slower life histories are only viable closer to the center of the realized niche. We identified glacial refugia farther north than previously recognized, as well as refugia concordant with previous population genetic findings. Lower latitude populations, known to be genetically distinct, are most threatened by future climate change. The recently colonized range of Arabidopsis was well-predicted by our native-range model applied to certain regions but not others, suggesting it has colonized novel climates.

MAIN CONCLUSIONS
Integration of distribution models with performance data from vast natural history collections is a route forward for testing biogeographical hypotheses about species distributions and their relationship with evolutionary fitness across large scales.

Introduction

Biogeographic patterns emerge from processes that act on many individuals and different local populations. Individuals die or survive in response to fluctuating environments, individuals migrate, and population allele frequencies change over time. While theory has often described how processes acting on individuals and local populations can generate biogeographic patterns (Keitt et al., 2001; Kirkpatrick & Barton, 1997; Pulliam, 2000) empirical study is challenged by the logistics required to measure individual-level variation across entire species ranges. Additionally, there is a growing call to connect range-wide, large scale model predictions with information on molecular, individual, and population-level processes (Kearney & Porter, 2009; Lasky et al., 2020). Thus there is a need for greater integration of biogeography with organismal biology to test hypotheses about the organismal and population mechanisms controlling distributions, and to understand how distributions and biogeographical history leave their imprints on individuals and populations.

In particular, a major goal in biology and biogeography is to understand how environmental conditions limit individual performance and species distributions. First, the environment-distribution relationship can help project species distributions under past climates to understand their history (Forester et al., 2013). Second, the environment-distribution relationship can identify regions with currently suitable habitat but that are unoccupied for other reasons (Elith et al., 2010). Third, the environment-distribution relationship can predict how distributions will shift under future environments (Thomas et al., 2004).

Environment-distribution relationships fundamentally arise from processes acting on individuals (Clark, 2010). On average, transplant experiments show that individual performance declines outside species' natural range (Hargreaves et al., 2014) and efforts to integrate information at the individual level into distribution models are emerging (Buckley et al., 2011; Elith et al., 2010; Lasky et al., 2020; Merow et al., 2014; Samis & Eckert, 2007). Where populations inhabit harsh environments (e.g. at range margins), local adaptations can emerge, such as life history changes to tolerate or escape harsh periods (Bontrager et al., 2021). For example, in Arabidopsis there is evidence that local adaptation to abiotic environment involves genetic changes in life history (e.g. flowering time) (Lovell et al., 2013; Martínez-Berdeja et al., 2020).

A vast resource of individual-level information can be found in natural history collections, and advances in the digitization of museum specimens are rapidly expanding the available data on range-wide variation (Heberling, 2021; Lopez et al., 2019). For example, Bontrager & Angert (2015) showed with herbarium specimens of *Clarkia* that fecundity decreased with drier summers, and toward one range margin both summer precipitation and individual fecundity declined, suggesting a limiting mechanism. In Arabidopsis, DeLeo et al. (2020) found decadal shifts in traits of herbarium specimens. For many species, seed banks (a type of natural history collection) house great diversity from across their ranges that can also be used to study their biogeography (Estarague et al., 2021; Scholl et al., 2000). These approaches are potentially valuable in model organisms, where researchers can link detailed information on genetics and organismal biology with population and community processes (Rudman et al., 2019; Takou et al., 2019). In this spirit we focus on *Arabidopsis thaliana,* a small annual plant (hereafter referred to as "Arabidopsis") (Koornneef & Meinke, 2010). Arabidopsis has been key to linking molecular biology, physiology, evolution, and ecology, but past biogeographic studies lacked statistical

inference (Hoffmann, 2002) or overlooked large parts of its range (Banta et al., 2012; Zou et al.,
2017). Distribution models fit to occurrence data are explicit quantitative statement of
environment-distribution relationships and allow inference of the environmental factors limiting
distributions (Elith et al., 2010; G. Li et al., 2015).
Arabidopsis is native across Eurasia and Africa and with human assistance has
colonized the Americas and Australia. Other species in the genus *Arabidopsis* are more
restricted to cool temperate climates, with Arabidopsis having expanded to a broader range, *e.g.*
Mediterranean habitats (Hoffmann, 2005). Arabidopsis can behave as a spring annual with a
rapid life cycle, germinating in the spring and flowering in the late spring and summer. However,
many individuals are longer-lived winter annuals, germinating in the fall and flowering in early
spring (Wilczek et al., 2009). Experiments have shown that winter cold is a major factor limiting
performance (Ågren & Schemske, 2012; Korves et al., 2007). Additionally, Arabidopsis lacks
physiological traits for dealing with severe water deficit so drought likely limits individual
performance in nature (Clauw et al., 2015). Studies of large-scale environmental response in
Arabidopsis have focused on the role of local adaptation in genetic diversity in the species (e.g.
Hancock et al., 2011; Lasky et al., 2018; Martínez-Berdeja et al., 2020; Toledo et al., 2020), but
less is known about the determinants of the species' distribution. The last overview of the
climate biogeography of Arabidopsis was Hoffmann (2002), who considered Arabidopsis native
to western Eurasia, but recently introduced in China and much of Africa. However, studies show
these latter populations are old, being genetically diverse, and with many unique genetic
variants (Durvasula et al., 2017; Zou et al., 2017).
Distributions are dynamic through time due to environmental change, dispersal, and
demographic stochasticity. Studies have used genetic data to infer Arabidopsis refugia during
the last glacial maximum (LGM), where populations (sometimes referred to as "relicts")
persisted locally before subsequent admixture with an expanding, now widely distributed "non-
relict" lineage (Lee et al., 2017). Whether these refugia corresponded to suitable climates is less
clear. Existing projections of future climate impacts on Arabidopsis have focused on *relative*
climate impacts on different genotypes (Exposito-Alonso et al., 2018; Fournier-Level et al.,
2016), rather than distribution dynamics. Additionally, many species have colonized new regions
due to human introduction, sometimes exhibiting traits different from native range populations,
potentially in response to new environments (Turner et al., 2015). Arabidopsis has colonized
many regions, but it is unclear to what degree these represent novel environments.
To evaluate biogeographic hypotheses linked to organismal and population processes,
we used recently developed modeling approaches and updated climate and occurrence data.
We combined occurrences (including many outside Europe poorly represented in previous
work) with climate data to build distribution models, and then test how model predictions
correspond to performance estimated from herbarium specimens and genetic variation in
natural accessions. We ask the following questions:
1. What climate factors constrain the distribution of Arabidopsis? We hypothesize that
159   winter cold and summer drought are the most important constraints, depending on
160   region.
161   2. Are individual performance and genetic variation associated with model-estimated
162   habitat suitability? We hypothesize that individuals reach larger sizes in regions with

greater suitability and that populations adapt along gradients in suitability through
changes in traits such as flowering time, a key component of life history.

3. Are occurrences outside the native range predicted by a native range model, suggesting
stable realized niches following colonization? Or is there evidence Arabidopsis has
colonized novel environments?

4. Where did Arabidopsis persist during the Last Glacial Maximum (LGM)? And where will
Arabidopsis move in future climates?

Methods

Occurrence data

We developed a set of high-quality occurrence data (*i.e.* species ID verified and location
checked, N=4,024) from published research (Durvasula et al., 2017; Hsu et al., 2019;
Mandáková et al., 2017; Zacchello et al., 2022; Zeng et al., 2017; Zou et al., 2017), publicly
available herbarium and germplasm accessions with known collection locations (Alonso-Blanco
et al., 2016; DeLeo et al., 2020), and some of our recent field collections in East Africa (Gamba
et al., 2022). These span a period of 1794 - 2018. The herbarium specimens and new
collections include little-studied populations in Saudi Arabia, Somalia, Djibouti, Eritrea, Rwanda,
Ethiopia, Uganda, Sudan, and Nepal. Duplicate occurrence points were eliminated (samples are
often split and sent to different herbaria). For model fitting, we excluded occurrences from
putative non-native regions (the Americas, New Zealand, Japan).

We also used occurrence data (with coordinates and without flagged problems
N=115,226) from the Global Biodiversity Information Facility (GBIF) to test model predictions in
regions outside of the native range of Arabidopsis (downloaded 30 Dec 2020, Gbif.Org, 2020).
We deem these occurrences as lower quality given that many have not had the species identity
and location checked (DeLeo et al., 2020).

Environmental data

Climate data were extracted from CHELSA (Climatologies at High resolution for the Earth's
Land Surface Areas) v1.2 at 30 arc second spatial resolution (Karger et al., 2017). Current
conditions are the average of 1979-2013 estimates. We selected the following climate variables
based on hypothesized importance (Gienapp et al., 2017; Hancock et al., 2011; Lasky et al.,
2014, 2018) and relatively low inter-correlation (Pearson correlation coefficients among
variables at occurrences < 0.75): isothermality (Bio3), minimum temperature of coldest month
(Bio6), temperature annual range (Bio7), mean temperature of wettest quarter (Bio8), mean
temperature of the warmest quarter (Bio10), precipitation seasonality (Bio15), precipitation of
wettest quarter (Bio16), and precipitation of driest quarter (Bio17). We also included elevation
from Tozer et al., (2019).

For projecting past distributions, we obtained climate estimates from the last glacial
maximum (LGM) at 21k yrs before present from CHELSA PMIP3 (Karger et al., 2017). We used
the global elevation and bathymetry map with 15 arc second resolution from Tozer et al. (2019)
and we adjusted elevations to make sea level 134 m lower than present (Lambeck et al., 2014)
to project potential suitable habitat at the LGM on land in areas currently submerged. For
projecting future distributions, we used climate projections from five divergent global climate
models for 2050 from CHELSA v1.2 using the RCP 4.5 emissions scenario (Karger et al., 2017).

We also show RCP 6.0 in the supplement for context (Figure S13), though it is highly similar to
4.5 in the target time period.
To characterize temporal variation in climate (climate anomalies), we used the Climate
Research Unit (CRU) TS 4.01 dataset, providing a global time series of monthly temperature
and precipitation for the period 1900-2010 at a 0.5⁰ resolution (Harris et al., 2014). From the
CRU data we calculated the same bioclimatic parameters that we used from CHELSA, but in
the CRU data these bioclimatic variables were specific to each herbarium specimen in the time
period it was collected (Supplemental Methods). We then calculated local anomalies for each of
these variables by taking the observed value, subtracting mean across the entire time series,
and dividing by the standard deviation (DeLeo et al., 2020).
217
Performance estimates from herbarium specimens
We estimated fecundity on a subset of herbarium specimens using two traits. First, we
measured the length of the longest inflorescence, reasoning that longer inflorescences would
have more fruits and seeds. Second, we measured maximum rosette leaf length, reasoning that
larger rosettes would support later reproductive investment if these collected plants were
allowed to continue growth *in situ* (see Supplement for a validation). We used ImageJ on 2,194
specimen images to estimate the tallest point of each inflorescence (N=2,188) and the
maximum rosette leaf length (N=1,264; see Supplement).
226
Range-wide genetic variation in life history
Many late-flowering Arabidopsis genotypes require cold cues (vernalization) to flower and also
show slower growth and more stress tolerance, delineating a life history axis (Lovell et al., 2013;
Vasseur et al., 2018). To assess life history variation, we used published experimental data on
flowering time for 898 whole-genome resequenced accessions from the native range with
reliable geographic coordinates grown at 10 and 16⁰C (The 1001 Genomes Consortium 2016).
To estimate vernalization sensitivity, we calculated the difference between flowering time at 10
and 16⁰C.
235
Species Distribution Modeling
We thinned the original 4,024 high quality occurrence points to one sample per 1 degree grid
cell to reduce sampling bias (N=764) using the 'sp' package (Bivand et al., 2008). To
characterize potentially inhabited sites, we generated pseudoabsence background points using
the 'dismo' package (Hijmans & Elith, 2013) by sampling 10,000 random points within a 500 km
buffer around occurrence points. We limited this buffer to 500 km to avoid including low sampled
regions in Africa and Asia that might result in biased model fit. We present a result with 1,000
km buffers in the supplement for reference.
We used Maxent version 3.4.0 to generate a species distribution model (Phillips et al.,
2006). MaxEnt was implemented with the R package 'ENMeval' v2, and parameters were
optimized using the 'checkerboard2' method for cross validation (Muscarella et al., 2014).
Among the tested settings (ENMeval defaults), we chose the model with lowest AICc value and
used this to project habitat suitability under recent conditions across the globe. For all models
we used the logistic output of MaxEnt that scales suitability from zero to unity.

We used permutation importances to determine which climatic factors drove predictions
in the distribution model. We also used the 'limiting' function in the R package 'rmaxent' to
determine the most limiting climatic factors in each location, defined as the environmental
covariate that has the largest decrease in suitability at a given location relative to the suitability
achieved if the covariate had its value equal to the global mean (Baumgartner et al., 2017; Elith
et al., 2010). State another way, the local limiting factor is the environmental condition most
limiting suitability, compared to an alternative scenario where that condition takes its global
mean.
In non-native regions, we evaluated whether Arabidopsis is limited from further
expansion at range edges by climate, *i.e.* whether there were no more unoccupied suitable
environments near existing populations. To do so, we calculated suitability in a zone 50-100 km
from existing GBIF occurrences and compared occupancy in these buffers in the native range to
invaded regions.
We also projected the MaxEnt model using past (LGM) and future climate conditions.
For future conditions, we calculated the mean and standard deviation of habitat suitability
projected for the five climate models. To assess whether model predictions were extrapolating
into poorly characterized or novel climates, we compared the present-day model training
climates to each predicted climate conditions, calculating multivariate environmental similarity
surfaces (MESS) following Elith et al. (2010). Higher values on a MESS map indicate conditions
in a location (or point in time) are similar to the reference environmental conditions used to fit
the model. Negative values indicate that at least some variables are outside the range of
environments used to fit the model, signifying extrapolation into novel environments (Elith et al.,
2010).
273
Performance and habitat suitability
We asked whether suitability corresponded to plant size. We first tested these relationships with
Pearson's and Spearman's correlations. We also fit Generalized Additive Models (GAMs) where
herbarium specimen sizes were the response variable. The model included covariates for
suitability at the collection location (square-root transformed to reduce the lower-tail influence)
and (as a nuisance variable) the year of collection (scaled to mean zero and unit variance) to
account for potential changes in size over time. We used GAMs to allow smooth spatial
variation in parameters, allowing us to capture regional variation in environmental response (for
an extensive description of GAMs in biogeography see Guisan et al., 2002). The model can be
represented as:
284
285    $$\text{Phenotype}_{i,year} = \beta_0 + \beta_{i,year}\text{Year} + \beta_{i,HS}\text{Habitat Suitability}_i + \square_{i,year}$$
286
where *i* represents the specimen location. Errors were modeled as normally distributed.
Spatially-varying parameters were constrained to smooth spatial variation in the GAMs using
the 'mgcv' package for R, which we used with REML to fit all GAMs (Wood, 2011). We
considered covariates to be significant at a given location if their 95% confidence interval (CI)
excluded 0, and the Moran's I (Cooper, 2021) for model residuals did not indicate spatial
autocorrelation.

We also asked whether local limiting factors corresponded to climate variables with
significant effects on plant performance. We hypothesized that for a given region, temporal
fluctuations in local limiting factors would be associated with temporal variation in plant size in
herbarium specimens. To address this hypothesis, we tested if specimen sizes were correlated
with anomalies in climate conditions corresponding to the climate variables used in the MaxEnt
model. We used the yearly climate anomalies we calculated from CRU in GAMs with spatially
varying coefficients, where for a phenotype measured at location i:
300
301         $\text{Phenotype}_{i,year} = \beta_{i,0} + \beta_{i,year}\text{Year} + \beta_{i,climate}\text{BioClim Variable Anomaly} + \square_{i,year}$
302
303   We then qualitatively assessed whether local limiting factors inferred by MaxEnt had anomalies
304   that were correlated with plant size (local $\beta_{i,climate}$ coefficients in the GAMs).
305
Genetic variation and habitat suitability
We tested how several aspects of genetic variation are related to suitability. First, we assessed
whether there was a genetic signature to genotypes currently found in regions that were also
highly suitable during the LGM. Lee et al. (2017) identified outlier haplotypes across the genome
that were highly differentiated, hypothesizing that these were genomic regions deriving from
long isolated "relict" populations that persisted through glacial cycles. We used bivariate
correlations and Fisher's exact test to test if higher LGM suitability was associated with a
greater number of outlier genomic regions.
Next, we focused on genetic variation in a key life history trait, flowering time. We
estimated linear models relating suitability and flowering time. To account for population
structure and neutral processes that can affect spatial variation in flowering time, we also tested
whether suitability was associated with flowering time after using random effects for genome-
wide similarity among accessions. A significant suitability-flowering time association in this
model would suggest selection linked to suitability acts on flowering time. This test is akin to
Q_{ST}-F_{ST} contrasts (Whitlock & Guillaume, 2009), except that an explicit environmental gradient is
tested (suitability). We implemented this mixed model the function 'lmekin' from the R package
'coxme' (Therneau & Therneau, 2015), along with 'kinship2' (Sinnwell et al., 2014). The kinship
matrix was obtained from 2,027,463 published whole genome resequencing SNPs (Alonso-
Blanco et al., 2016). Additionally, to test for geographic variation in suitability-flowering time
relationships, we fitted GAMs of flowering time with spatially varying coefficients:
326
327         $\text{FloweringTime}_{i,year} = \beta_0 + \beta_{i,HS}\text{Habitat Suitability} + \square_i$.
328
Our last strategy was to scan the Arabidopsis genome for genes where different alleles
were found in high versus low suitability locations. This approach could be considered a novel
application of a genotype-environment association (GEA) in an environmental genome wide
association study (GWAS) (Lasky et al., 2023). Such a change in allele frequency across
gradients in suitability would suggest that this genetic variation was involved in local adaptation
to low versus high suitability environments. We used univariate linear mixed-effects models in
GEMMA (v 0.98.3) (Zhou & Stephens, 2012) to perform the environmental GWAS in a set of
2,053,939 SNPs filtered for MAF=0.05 from 1003 native-range ecotypes part of the 1001

genomes panel. GEMMA includes random effects that are correlated according to genome-wide
similarity between samples ("kinship"), allowing it to account for genomic background effects.
339
340
341   **Results**
342
Q1. Climate constraints on the distribution of Arabidopsis
Our optimization of MaxEnt models (with AICc) resulted in a selection of a model with linear and
quadratic effects, with the regularization multiplier of 0.5. The training AUC value was 0.79 and
the average test AUC with checkerboard2 cross validation was 0.78. The omission rate of the
10th percentile of suitability for training points was 0.10, suggesting our models were not overfit
as they were able to predict low probability occurrences as well as expected (Fielding & Bell,
1997; Peterson et al., 2011). The areas of high suitability overall correspond well to the
documented Arabidopsis distribution, with one notable exception being tropical lowland sites
(Congo basin) in sub-Saharan Africa (Figure 1A). This region was near zero multivariate
environmental similarity to training data, indicating the model may have been poorly constrained
there, while most of the regions of high predicted suitability where Arabidopsis is documented
have positive similarity (MESS, Figure S2).
355
The permutation importance of the model covariates across the entire native range revealed
that the minimum temperature of the coldest month (PI = 38%) and the mean temperature of the
warmest quarter (PI = 31%) were the two most important variables, suggesting winter cold
stress and summer heat stress are most important in constraining Arabidopsis's distribution
(Table S1). The next most important variable was isothermality (PI =15%), as Arabidopsis tends
to be found in regions with low isothermality (e.g. most temperate zones).
To identify spatial variation in climate constraints, we also identified local limiting factors
(Figure 1B). Across northern Eurasia, minimum temperature of the coldest month limited habitat
suitability. Across the Mediterranean and tropical/subtropical regions, the temperature of the
warmest quarter was limiting. Across Eastern Europe and Central Asia, winter cold was limiting
adjacent to other regions where summer heat was limiting, highlighting the multiple stressors in
this region.
We hypothesized that for a given region, temporal fluctuations in local limiting factors
would be associated with temporal variation in plant size in herbarium specimens. We found
that years with warmer minimum winter temperatures were associated with significantly taller
inflorescences from Turkey to central Asia (Figure S3, panel B). Concordantly, in much of this
region the Maxent model indicated that minimum temperature of the coldest of the month was
the most limiting factor (namely in the Caucusus and from Kazakhstan to northern India, Figure
1B). We also found that years with greater seasonality of precipitation were associated with
shorter inflorescences in central Asia, but taller inflorescences in central Europe (Figure S3,
panel F). Partly consistent, in the southern part of this central Asian region the MaxEnt model
indicated precipitation seasonality was limiting (Figure 1B). Years with higher isothermality were
positively associated with inflorescence height in Eastern Europe but this was not a limiting
factor in this region (Figure S3, panel A). Other climate anomalies were not significantly
associated with temporal variation in plant size (Figures S3 and S4).

Q2. Declining individual performance and prevalence of stress escape life history in low
suitability regions

As habitat suitability increased, so did the inflorescence height of individual plants in herbarium
specimens (Pearson's r = 0.12, p < 10^{-8}; Spearman's rho = 0.10, p < 10^{-6}; n = 2169). The
relationship with suitability was even stronger for maximum rosette leaf length (r = 0.25, p < 10^{-16}
; rho = 0.26, p < 10^{-16}; n = 1253, Figure 2), which was also more correlated with total fruit +
flower number (Supplement), suggesting fecundity was greater in regions of high predicted
suitability. We tested size-suitability relationships using GAMs with spatially varying suitability
coefficients (but not including climate anomalies – distinct from the previous section Q1). We
found a consistently positive relationship between suitability and size that was significant for
rosette leaf length in western and central Europe and for inflorescence height across most of
Eurasia (Figs S5 and S6). Unexpectedly, in this model where year of collection was considered
primarily as a nuisance variable, inflorescence height significantly declined over time in
northwest Europe (Fig S6; e.g. in Scandinavia and Scotland, maximum rosette leaf length vs
year, Spearman's rho = -0.14).

We compared habitat suitability with published data on genetic variation in flowering
time. We found that flowering time at 10ºC and flowering time plasticity (flowering time at 10ºC –
time at 16ºC) were significantly positively associated with suitability (r=0.19, p<10^{-8} and r=0.13,
p=0.0002, respectively), but not for days to flower at 16 ºC (r=0.05, p=0.1264, Figures 2 & S7).
In GAMs with spatially-varying suitability coefficients, the suitability-flowering time pattern were
largely consistent across Eurasia (Figure S8). The suitability-flowering time association was
significant even when accounting for genomic similarity among accessions, suggesting selection
associated with suitability could maintain geographic clines in flowering time. Specifically, linear
mixed-effects models found a positive association for flowering time at 10ºC (p=0.0004, n=884),
and also at 16ºC, the latter of which may have been obscured by population structure that was
unaccounted for in the simple linear regression model (p=0.02, n=852, Tables S2-S3). The
suitability association was not significant for plasticity in flowering time when accounting for
genomic similarity (p=0.77, n=852) (Table S4). Consistent with this potential obscuring of
population genetic structure, we found that 10 ADMIXTURE genetic clusters in Arabidopsis
(Alonso-Blanco et al., 2016) were significantly different in their habitat suitability
(F(9,993)=201.6, p<10^{-16}, Table S5, Figure S9).

We scanned the genome for genes that showed allele frequency correlations with
suitability. The most strongly associated SNP was in the putative promoter region (134 bp from
the start, linear mixed model, p < 10^{-16}) of ERF53 (AT2G20880), a transcription factor that
regulates response to drought, salt, and heat (Figure S11) (Cheng et al., 2012; B. Li et al.,
2019). This SNP showed a strong allele frequency cline from Europe to Asia, where the
alternate allele was nearly fixed in accessions east of the Ural Mountains, which is a region of
low estimated suitability (Fig S10). Furthermore, the alternate allele was associated with lower
expression of ERF53 (Wilcox test, p = 0.0013) in published transcriptome data (Kawakatsu et
al., 2016), suggesting a *cis*-regulatory variant is locally adapted to low suitability parts of Asia.
Relatedly, Jiang et al., (2022) showed that knockouts of ERF53 have temperature-dependent
effects on fecundity.

425
Q3. Using the native range model to predict outside the native range
Australia and New Zealand occurrences were largely in areas predicted to be highly suitable
(Figure 3). By contrast, North America includes occurrences in highly suitable areas (Pacific
Northwest) but many in low suitability areas (interior east, where summer heat was predicted
limiting, Figure S12). The eastern US also has near zero similarity on the MESS map,
suggesting a novel set of conditions compared to the native range (Figure S2). Similarly, the
regions where Arabidopsis occurred in Korea and Japan were of lower suitability than in the
native range, also with summer heat as the predicted limiting factor (Figure S12), suggesting
Arabidopsis in eastern North America and East Asia inhabits climates with distinctly warm
summers.
436   We investigated whether non-native Arabidopsis occupy all available habitat in their
regions or whether suitable habitat remains unoccupied. We found that locales 50-100 km from
occurrences included some areas of high suitability in North America, suggesting suitable sites
remain unoccupied (4.3% of grid cells in these 50-100 km regions had suitability > 0.6). There
were extensive areas of high suitability along the Pacific coast of North America to around 60ºN,
but no occurrences north of 50º, which we confirmed with an expert botanist (pers comm.
Matthew Carlson). The timing of initial colonization is likely not a factor given that occurrences
from near Vancouver date at least to 1939. This region has near zero similarity on the MESS
map, potentially indicating that the native range model is not well constrained there (Figure S2).
Similarly, the southern coast of Australia is highly suitable but Arabidopsis is apparently absent
(35.0% of these 50-100 km regions with suitability > 0.6). Records are restricted to the
southeast and southwest (confirmed by botanists, Shelley James, Tim Entwistle, Neville Walsh
pers. comm.), even though records date at least to 1959 in SE Australia. In South America there
are a few Arabidopsis records from the Southern Cone, and it appears to be rare in the region
(pers. comm. Diego Salariato), while apparently suitable environments occur throughout
Patagonia and high elevation Andean sites that are apparently unoccupied (pers comm.
Gwendolyn Peyre, Santiago Madriñán). Large areas of southern Australia and South America
show positive similarity on the MESS map suggesting the model is well constrained in those
regions. By contrast in Korea and Japan there are very few sites expected to be highly suitable
that are not already occupied by Arabidopsis (2.4% of these 50-100 km regions with suitability >
0.6).
457
Q4. The distribution of Arabidopsis during the Last Glacial Maximum and in the future
We projected suitability onto the climate at the LGM and found several putative refugia, where
past environmental conditions could have supported Arabidopsis persistence. In particular, the
Mediterranean/Caucuses/south Caspian Sea, larger portions of Africa > 1000 m (current) asl,
and China and SE Asia appear as refugia (Figure 4). North Africa, the Atlantic European coast,
and the islands of Sicily, Corsica, and Sardinia also appear as highly suitable potential refugia.
464   We projected climate suitability for Arabidopsis in the year 2050. Some current high
suitability regions will remain so, such as in Northern Europe. Nevertheless, we found poleward
and up-elevation shifts in regions of high suitability and retreats at lower latitudes and
elevations. In the native range, all African sites show declining suitability, as did most
Mediterranean sites, highlighting vulnerability of these populations. By contrast, Arctic Europe,

and the mountains of central Asia/Tibetan plateau show increased suitability, highlighting
potential range expansions (Figure 5).
We found concordance between our LGM predictions and the frequency of outlier
genomic regions in modern genotypes (Lee et al., 2017). The suitability during the LGM at the
site of genotype origin was positively correlated with the number of outlier haplotypes they
carried (N = 933, Pearson's r = 0.22, p<10^{-12}; Spearman's rho = 0.14, p<10^{-5}). This relationship
was partly due to the fact that >90% of the distinct relict genotypes defined by Lee et al. (2017)
were located in regions where the LGM suitability was over 0.25, versus only 48% for non-relicts
(Fisher's exact test p<10^{-5}, 0.25 approximately defining range limits in current climates, but this
test was highly significant for suitability thresholds from 0.1 to 0.5). However, even among non-
relicts, the higher the LGM suitability the greater the frequency of outlier haplotypes (N = 911,
Pearson's r = 0.20, p<10^{-10}; Spearman's rho = 0.10, p=0.0017).
481
482
483 **Discussion**
484
We used the model plant Arabidopsis in a case study of integrative climate biogeography of a
species' past, present, and future distributions. This species is key to a large body of plant
biology research, and an in-depth study connecting its biogeography to genetic and phenotypic
variation provides important context for the biology of Arabidopsis. The size of individual plants
in herbarium specimens, as well as genetic variation in flowering time and a gene controlling
abiotic stress response, suggested that regions with lower predicted suitability harbored
populations with altered life history and physiology. The consistency between model predictions
and individual variation provides partial model validation, bolstering our confidence in
conclusions from model projections. Using these model projections, we identified new glacial
refugia, and looking to the future, we found that genetically distinct lower latitude populations
are most threatened by climate change.
496
The current distribution of Arabidopsis and its limiting factors
The fitted native-range model largely corresponded to occurrences and indicates Arabidopsis
broadly distributed across Europe, moister regions of central and eastern Asia, and mountains
across Africa. Our results advance beyond the most recent climate biogeographical study of
Arabidopsis by Hoffmann (2002), who used monthly climate data (but notably no synthetic
bioclimatic variables) from Leemans & Cramer (1991) mostly to qualitatively describe the
regions occupied by Arabidopsis. Our study included populations in sub-Saharan Africa and
Asia that were overlooked or considered non-native by Hoffmann (2002). We also included
occurrences from regions apparently missed by Hoffmann (2002). Zou et al. (2017) more
recently built models of Arabidopsis's distribution using Eurasian populations and default
MaxEnt settings, but did not include occurrences from Arabia, sub-Saharan Africa, and much of
the Himalayas, Russia, and central Asia, and did not subsample occurrence to reduce bias.
Likely as a result of these issues, the predictions of Zou et al. (2017) show a pronounced peak
in suitability in Germany, which was densely sampled in the genomics studies used for
occurrences by Zou et al. (2017), but predicted low suitability in much of the core European

range and near zero suitability in sub-Saharan Africa, Arabia, and most of the Russian part of
the species range.
The two dominant limiting factors in our model were winter cold (at higher latitudes and
elevations) and summer heat (at lower latitudes and elevations). Winter cold is recognized to
limit Arabidopsis performance, in particular winter cold appears to be a dominant force in local
adaptation of Arabidopsis (Ågren & Schemske, 2012; Gienapp et al., 2017; Monroe et al.,
2016). In southern Europe, where summer heat was inferred to be limiting, Arabidopsis flowers
in early spring (DeLeo et al., 2020) and thus summer heat is not usually directly experienced. In
spring in these regions, warm temperatures might not reach consistently stressful levels (e.g. to
induce fruit abortion) (Warner & Erwin, 2005) but it may be that moisture deficit driven by
evaporative demand is directly limiting in late spring. Where suitability was highest for
Arabidopsis, including Britain and the Atlantic coast of France, high isothermality and high
precipitation of the driest quarter were identified as limiting. However, interpreting limiting factors
in an area of extremely high suitability (near one) is not meaningful given that suitability can
scarcely be increased.
Despite inference of winter cold and summer heat as primarily limiting, these were only
partly reflected by temporal fluctuations in individual plant performance from herbarium
specimens. Winter cold and precipitation variability anomalies were significantly associated with
specimen size in much of Asia, where these were also the MaxEnt modeled limiting factors from
Iran and Kazakhstan to Afghanistan and the Himalayas, suggesting these climate factors truly
limit Arabidopsis populations. However, there were discrepancies between individual
performance and limiting factors, likely for several reasons. First, the size of specimens is an
imperfect fitness proxy. Usually only reproductive individuals are collected in herbaria, and
individuals that would die before reproduction are excluded. Second, individuals are not
randomly sampled from populations (Daru et al., 2018). Third, MaxEnt models face limitations
due to misspecification, problems with occurrence data, or a mismatch between covariates and
the true ecological factors limiting populations. Nonetheless, the limiting factors we identified
here fit well with our knowledge of Arabidopsis ecophysiology and natural history.
540
Habitat suitability, individual performance, and life history
We found a decrease in plant size with decreasing habitat suitability across the range of
Arabidopsis, suggesting that our model suitability captured a substantial part of environmental
effects on individual fitness. Similar to our findings, Karasov et al., (2022) found declines in the
size of natural Arabidopsis plants at latitudinal extremes of the European range. Given the great
variability in individual plant performance within populations obvious to casual observers, it is
unsurprising that suitability here explained a minority of total variation in individual size, leaving
most variation unexplained (Figure 2). Furthermore, fecundity and fitness response to
treatments often have low heritability in many species (Price & Schluter, 1991), even in
controlled experiments (Lasky et al., 2015).
There have still been few studies of individual performance with the geographic scope
that allows inference across a species range (Angert & Schemske, 2005; Csergő et al., 2017;
Greiser et al., 2020; Samis & Eckert, 2007). In a synthesis of studies of 40 species, Lee-Yaw et
al. (2016) found that, on average, individual performance and distribution-model inferred
suitability decline beyond range margins. From 42 studies Lee-Yaw et al. (2021) found that 38%

identified some predictive ability of distribution models for individual performance. However,
many previous studies relied on intensive observations of a small number of populations, while
our estimates of performance from thousands of herbarium specimens allowed us to cover most
of the species range. The increased availability of digitized museum specimens with trait data
indicates an opportunity to estimate performance across distributions for many species
(Bontrager & Angert, 2015).
We found that low suitability regions had earlier flowering genotypes. This relationship
was noisy, with early flowering genotypes frequent in all levels of suitability but later flowering
genotypes lacking from the least suitable regions. The suitability-flowering time associations
were significant when accounting for genome-wide similarity between accessions, suggesting
they reflect selection associated with suitability. We interpret the direction of the relationship as
indicating that when suitability is low, Arabidopsis employs stress escape strategies, *i.e.* a rapid
life cycle during favorable conditions (Ludlow, 1989). Nevertheless, our findings suggest that
low suitability central Asian populations have some stress tolerating mechanisms, as they
harbor distinct allele at a transcription factor (ERF53, AT2G20880) that regulates response to
abiotic stressors (Cheng et al., 2012; B. Li et al., 2019) including performance response to
temperature (Jiang et al., 2022). The restriction of late flowering genotypes to more suitable
regions is counterintuitive given physiological work showing these are more stress tolerant
(Lovell et al., 2013). However, it may be true that more favorable conditions make possible a
slow growing, slow flowering, freezing-tolerant winter annual strategy.
576
The distribution of Arabidopsis outside its native range
Arabidopsis has spread across the globe, largely to climates well-predicted by native range
models (western North America, Australia, and New Zealand), suggesting some stability in
climate niche, but also to some regions predicted to be less suitable (eastern North America,
Korea, and Japan). Whether these are truly of lower suitability is unknown without performance
data. It may be that Arabidopsis has colonized novel environments in these regions, or that the
low suitability is only a model artefact. Studies often find limited transferability among native and
introduced range models (Early & Sax, 2014; Liu et al., 2020).
585
The distribution of Arabidopsis at the last glacial maximum
Lee et al. (2017) hypothesized five glacial refugia for Eurasia to be in Iberia, Sicily, Balkans, the
Levant, and Turkmenistan. These are consistent with our range reconstruction, although we do
not find clear barriers of unsuitable climates that would have isolated Italian, Balkan, and Levant
LGM populations. Furthermore, we note that Sardinia/Corsica appears as a refugium, and we
note that narrow, highly suitable areas along the southern and eastern Caspian Sea could be
the location of the hypothesized Turkmenistan refugium. Lee et al. (2017) further hypothesized
that the bulk of current European genotypes derive from an expansion originating on the
northwest coast of the Black Sea, where we found a strip of highly suitable conditions for
Arabidopsis during the LGM. Additional genetic studies hypothesized a long history of existence
in sub-Saharan Africa (Durvasula et al., 2017) and conditions during the LGM suggest the
species could have been more widespread than currently. Concordantly, Chala et al., (2017)
used models of the distribution of Afroalpine habitat generally during the LGM to find expanded

areas in East Africa compared to present day, though some locations such as Jebel Marra,
Sudan were still surrounded by landscapes of unsuitable habitat even at the LGM.
Our findings also may help explain the history of Cape Verde Island populations, which
Fulgione et al., (2022) estimated were bottlenecked and split from a source population 5-10 kya.
The source population appears to be extinct and but is most similar to current Moroccan
populations (Fulgione et al., 2022). Currently there are no known African Atlantic coastal, low
elevation populations that would have been sources. However, we found that during the LGM
and presumably for some time after, low elevation coastal high suitability habitat extended as far
south as Ras Nouadhibo on the African coast, suggesting a source for the Cape Verde
immigrants.
We found that the North Atlantic European coast was highly suitable (including areas
currently inhabited) during the LGM, but current British Isle and French populations do not show
genetic signatures of refugia (Lee et al., 2017). In both cases, local, distinct genotypes that
survived the LGM may have gone extinct following expansion by the now dominant European
genetic cluster ("non-relicts") (Fulgione & Hancock, 2018; Lee et al., 2017). The previous
Arabidopsis distribution modeling of the LGM (Zou et al., 2017) did not identify France and NE
Iberia as highly suitable during the LGM and did not mask the Caspian Sea (thus overinflating
that refugium). Zou et al. (2017) also did not identify the African refugia (except for a small
region in SE Africa) and potential refugia in mountains of SE Asia.
618
The future distribution of Arabidopsis
Over the next decades conditions are predicted to worsen for Arabidopsis across large areas of
its range. Central Spain and mountains in Africa and Arabia may see the worst changes.
Impacts may already be emerging: our intensive field search in the Dai Forest of Djibouti in
2018 failed to yield any Arabidopsis, despite the presence of abundant (but likely more drought
tolerant) annual mustards (Brassicaceae) *Erucastrum arabicum* and *Sisymbrium erysimoides* in
otherwise appropriate habitat for Arabidopsis. Arabidopsis was collected in this juniper
woodland in 1954 but recent decades have seen increased drought and tree mortality (Witsen,
2012), suggesting potential extinction of this isolated marginal Arabidopsis population. Given the
unique genetic diversity of the model plant Arabidopsis housed in its lower latitude populations
(Durvasula et al., 2017; Gamba et al., 2022; Hsu et al., 2019; Lee et al., 2017; Zou et al., 2017)
the conservation of these populations could benefit plant biology research.
By contrast, conditions are expected to improve for the northernmost populations in
Europe, suggesting a potential current colonization front of Arabidopsis, in addition to higher
elevation locations in Tibet, the Caucuses, Ural, Alps, and Hengduan mountains adjacent to
currently inhabited regions. Similar currently unoccupied higher elevations are available in many
African mountains though these are usually small mountaintop areas and may be poorly
characterized by CHELSA climate data (Karger et al., 2017). Future distribution models would
benefit from improved environmental data for these high elevation tropical sites.
638
639 *Conclusion*
640 Biogeographic patterns emerge from processes affecting individuals and populations. Species
641 distribution models provide potentially powerful windows into past, present, and future
642 macroecology, but they are rarely confronted with range-wide individual performance data. Here

we showed that lower suitability habitats inferred from distribution models had smaller plants with distinct life history, suggesting a stress escape strategy. While the relationships were noisy it may still be remarkable that they emerge against the many microsite contributors to individual level variation in a habitat generalist annual plant. Arabidopsis populations are distributed across diverse climates, but genetically distinct populations in lower latitudes that are potentially valuable for research are also highly threatened by anthropogenic climate change in the next few decades. We believe that combining distribution models with individual data on genotype and phenotype across a species range can be a useful approach to dissect the organismal mechanisms underlying distributions and to validate distribution models.

Acknowledgments

This manuscript benefited from the comments of three anonymous reviewers. Victoria Meagher and Jaden Hill measured morphology on herbarium specimens. Patrick Herné assisted in digitizing many herbarium specimens and Eugene Shakirov translated many Russian specimen labels. Data from the MNHN in Paris were obtained from the participatory science program 'Les Herbonautes' (MNHN/Tela Botanica) which is part of Infrastructure Nationale e-RECOLNAT: ANR-11-INBS-0004. This work was supported by NIH award R35 GM138300 to JRL. RM was supported by grant 2019-03758 from Vetenskapsrådet. We also thank volunteers for digitization and georeferencing through DigiVol, a website developed by the Australian Museum in collaboration with the Atlas of Living Australia.

Conflict of interest

The authors declare no conflict of interest.

Data availability

MaxEnt model projections, occurrence data, and herbarium specimen data will be stored in a Dryad repository upon acceptance of this manuscript. 1001 Genomes data were obtained from 1001genomes.org. Current and paleoclimate data were obtained from chelsa-climate.org. Time series climate data were obtained from data.ceda.ac.uk.

References

Ågren, J., & Schemske, D. W. (2012). Reciprocal transplants demonstrate strong adaptive differentiation of the model organism Arabidopsis thaliana in its native range. *New Phytologist*, *194*(4), 1112–1122. https://doi.org/10.1111/j.1469-8137.2012.04112.x

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M., Cao, J., Chae, E., Dezwaan, T. M., Ding, W., Ecker, J. R., Exposito-Alonso, M., Farlow, A., Fitz, J., Gan, X., Grimm, D. G., Hancock, A. M., Henz, S. R., Holm, S., … Zhou, X. (2016). 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell*, *166*, 481–491. https://doi.org/10.1016/j.cell.2016.05.063

Angert, A. L., & Schemske, D. W. (2005). The Evolution of Species' Distributions: Reciprocal Transplants Across the Elevation Ranges of Mimulus Cardinalis and M. Lewisii. *Evolution*, *59*(8), 1671–1684. https://doi.org/10.1111/j.0014-3820.2005.tb01817.x

Banta, J. A., Ehrenreich, I. M., Gerard, S., Chou, L., Wilczek, A., Schmitt, J., Kover, P. X., & Purugganan, M. D. (2012). Climate envelope modelling reveals intraspecific relationships among flowering phenology, niche breadth and potential range size in Arabidopsis thaliana. *Ecology Letters*, *15*(8), 769–777. https://doi.org/10.1111/j.1461-0248.2012.01796.x

Baumgartner, J., Wilson, P., & Esperón-Rodríguez, M. (2017). rmaxent: Tools for working with Maxent in R. *R Package Version 0.8*, *3*.

Bivand, R. S., Pebesma, E. J., Gómez-Rubio, V., & Pebesma, E. J. (2008). *Applied spatial data analysis with R* (Vol. 747248717). Springer.

Bontrager, M., & Angert, A. L. (2015). Effects of range-wide variation in climate and isolation on floral traits and reproductive output of Clarkia pulchella. *American Journal of Botany*, *103*, 10–21. https://doi.org/10.3732/ajb.1500091

Bontrager, M., Usui, T., Lee-Yaw, J. A., Anstett, D. N., Branch, H. A., Hargreaves, A. L., Muir, C. D., & Angert, A. L. (2021). Adaptation across geographic ranges is consistent with strong selection in marginal climates and legacies of range expansion. *Evolution*, *75*(6), 1316–1333. https://doi.org/10.1111/evo.14231

Buckley, L. B., Waaser, S. A., MacLean, H. J., & Fox, R. (2011). Does including physiology improve species distribution model predictions of responses to recent climate change? *Ecology*, *92*(12), 2214–2221. https://doi.org/10.1890/11-0066.1

Chala, D., Zimmermann, N. E., Brochmann, C., & Bakkestuen, V. (2017). Migration corridors for alpine plants among the 'sky islands' of eastern Africa: Do they, or did they exist? *Alpine Botany*, *127*(2), 133–144. https://doi.org/10.1007/s00035-017-0184-z

Cheng, M.-C., Hsieh, E.-J., Chen, J.-H., Chen, H.-Y., & Lin, T.-P. (2012). Arabidopsis RGLG2, Functioning as a RING E3 Ligase, Interacts with AtERF53 and Negatively Regulates the Plant Drought Stress Response. *Plant Physiology*, *158*(1), 363–375. https://doi.org/10.1104/pp.111.189738

Clark, J. S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest Trees. *Science*, *327*(5969), 1129–1132.

Clauw, P., Coppens, F., De Beuf, K., Dhondt, S., Van Daele, T., Maleux, K., Storme, V., Clement, L., Gonzalez, N., & Inzé, D. (2015). Leaf Responses to Mild Drought Stress in Natural Variants of Arabidopsis. *Plant Physiology*, *167*(3), 800–816. https://doi.org/10.1104/pp.114.254284

Cooper, M. (2021). *moranfast: Calculate Moran's I quickly with low memory footprint for Big Data.* [C++]. https://github.com/mcooper/moranfast

Csergő, A. M., Salguero-Gómez, R., Broennimann, O., Coutts, S. R., Guisan, A., Angert, A. L., Welk, E., Stott, I., Enquist, B. J., McGill, B., Svenning, J.-C., Violle, C., & Buckley, Y. M. (2017). Less favourable climates constrain demographic strategies in plants. *Ecology Letters*, *20*(8), 969–980. https://doi.org/10.1111/ele.12794

Daru, B. H., Park, D. S., Primack, R. B., Willis, C. G., Barrington, D. S., Whitfeld, T. J. S.,
Seidler, T. G., Sweeney, P. W., Foster, D. R., Ellison, A. M., & Davis, C. C. (2018).
Widespread sampling biases in herbaria revealed from large-scale digitization. *New*
Phytologist, *217*(2), 939–955. https://doi.org/10.1111/nph.14855
DeLeo, V. L., Menge, D. N. L., Hanks, E. M., Juenger, T. E., & Lasky, J. R. (2020). Effects of
two centuries of global environmental variation on phenology and physiology of
Arabidopsis thaliana. *Global Change Biology*, *26*(2), 523–538.
https://doi.org/10.1111/gcb.14880
Durvasula, A., Fulgione, A., Gutaker, R. M., Alacakaptan, S. I., Flood, P. J., Neto, C.,
Tsuchimatsu, T., Burbano, H. A., Picó, F. X., Alonso-Blanco, C., & Hancock, A. M.
(2017). African genomes illuminate the early history and transition to selfing in
Arabidopsis thaliana. *Proceedings of the National Academy of Sciences*, 201616736.
https://doi.org/10.1073/pnas.1616736114
Early, R., & Sax, D. F. (2014). Climatic niche shifts between species' native and naturalized
ranges raise concern for ecological forecasts during invasions and climate change.
Global Ecology and Biogeography, *23*(12), 1356–1365.
https://doi.org/10.1111/geb.12208
Elith, J., Kearney, M., & Phillips, S. (2010). The art of modelling range-shifting species. *Methods*
in Ecology and Evolution, *1*(4), 330–342. https://doi.org/10.1111/j.2041-
210X.2010.00036.x
Estarague, A., Vasseur, F., Sartori, K., Bastias, C., Cornet, D., Rouan, L., Beurier, G., Exposito-
Alonso, M., Herbette, S., Bresson, J., Vile, D., & Violle, C. (2021). *Into the range: A*
latitudinal gradient or a center-margins differentiation of ecological strategies in
Arabidopsis thaliana? (p. 2021.10.15.461205).
https://doi.org/10.1101/2021.10.15.461205
Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A., & Weigel, D. (2018).
Genomic basis and evolutionary potential for extreme drought adaptation in Arabidopsis
thaliana. *Nature Ecology & Evolution*, *2*(2), 352. https://doi.org/10.1038/s41559-017-
0423-0
Fielding, A. H., & Bell, J. F. (1997). A review of methods for the assessment of prediction errors
in conservation presence/absence models. *Environmental Conservation*, *24*(1), 38–49.
https://doi.org/10.1017/S0376892997000088
Forester, B. R., DeChaine, E. G., & Bunn, A. G. (2013). Integrating ensemble species
distribution modelling and statistical phylogeography to inform projections of climate
change impacts on species distributions. *Diversity and Distributions*, *19*(12), 1480–1495.
https://doi.org/10.1111/ddi.12098
Fournier-Level, A., Perry, E. O., Wang, J. A., Braun, P. T., Migneault, A., Cooper, M. D.,
Metcalf, C. J. E., & Schmitt, J. (2016). Predicting the evolutionary dynamics of seasonal
adaptation to novel climates in Arabidopsis thaliana. *Proceedings of the National*
Academy of Sciences, *113*(20), E2812–E2821.
https://doi.org/10.1073/pnas.1517456113
Fulgione, A., & Hancock, A. M. (2018). Archaic lineages broaden our view on the history of
Arabidopsis thaliana. *New Phytologist*, *219*(4), 1194–1198.
https://doi.org/10.1111/nph.15244
Fulgione, A., Neto, C., Elfarargi, A. F., Tergemina, E., Ansari, S., Göktay, M., Dinis, H., Döring,
N., Flood, P. J., Rodriguez-Pacheco, S., Walden, N., Koch, M. A., Roux, F., Hermisson,
J., & Hancock, A. M. (2022). Parallel reduction in flowering time from de novo mutations
enable evolutionary rescue in colonizing lineages. *Nature Communications*, *13*(1), Article
1. https://doi.org/10.1038/s41467-022-28800-z
Gamba, D., Lorts, C., Haile, A., Sahay, S., Lopez, L., Xia, T., Kulesza, E., Elango, D., Kerby, J.,
Yifru, M., Bulafu, C. E., Wondimu, T., Glowacka, K., & Lasky, J. (2022). *The genomics*

774	*and physiology of abiotic stressors associated with global elevation gradients in*
775	*Arabidopsis thaliana* (p. 2022.03.22.485410). bioRxiv.
776	https://doi.org/10.1101/2022.03.22.485410
777	Gbif.Org. (2020). *Occurrence Download*. The Global Biodiversity Information Facility.
778	https://doi.org/10.15468/DL.H95GHR
779	Gienapp, P., Fior, S., Guillaume, F., Lasky, J. R., Sork, V. L., & Csilléry, K. (2017). Genomic
780	Quantitative Genetics to Study Evolution in the Wild. *Trends in Ecology & Evolution*,
781	*32*(12), 897–908. https://doi.org/10.1016/j.tree.2017.09.004
782	Greiser, C., Hylander, K., Meineri, E., Luoto, M., & Ehrlén, J. (2020). Climate limitation at the
783	cold edge: Contrasting perspectives from species distribution modelling and a transplant
784	experiment. *Ecography*, *43*(5), 637–647. https://doi.org/10.1111/ecog.04490
785	Guisan, A., Edwards, T. C., & Hastie, T. (2002). Generalized linear and generalized additive
786	models in studies of species distributions: Setting the scene. *Ecological Modelling*,
787	*157*(2), 89–100. https://doi.org/10.1016/S0304-3800(02)00204-1
788	Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G.,
789	Toomajian, C., Roux, F., & Bergelson, J. (2011). Adaptation to Climate Across the
790	Arabidopsis thaliana Genome. *Science*, *334*(6052), 83–86.
791	Hargreaves, A. L., Samis, K. E., Eckert, C. G., Schmitz, A. E. O. J., & Bronstein, E. J. L. (2014).
792	Are Species' Range Limits Simply Niche Limits Writ Large? A Review of Transplant
793	Experiments beyond the Range. *The American Naturalist*, *183*(2), 157–173.
794	https://doi.org/10.1086/674525
795	Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of
796	monthly climatic observations – the CRU TS3.10 Dataset. *International Journal of*
797	*Climatology*, *34*(3), 623–642. https://doi.org/10.1002/joc.3711
798	Heberling, J. M. (2021). Herbaria as Big Data Sources of Plant Traits. *International Journal of*
799	*Plant Sciences*. https://doi.org/10.1086/717623
800	Hijmans, R. J., & Elith, J. (2013). Species distribution modeling with R. *R CRAN Project*.
801	Hoffmann, M. H. (2002). Biogeography of Arabidopsis thaliana (L.) Heynh. (Brassicaceae).
802	*Journal of Biogeography*, *29*(1), 125–134.
803	Hoffmann, M. H. (2005). EVOLUTION OF THE REALIZED CLIMATIC NICHE IN THE GENUS:
804	ARABIDOPSIS (BRASSICACEAE). *Evolution*, *59*(7), 1425-1436.
805	Hsu, C.-W., Lo, C.-Y., & Lee, C.-R. (2019). On the postglacial spread of human commensal
806	Arabidopsis thaliana: Journey to the East. *New Phytologist*, *222*(3), 1447–1457.
807	https://doi.org/10.1111/nph.15682
808	Jiang, Y., Wang, Z., Du, H., Dong, R., Yuan, Y., & Hua, J. (2022). Assessment of functional
809	relevance of genes associated with local temperature variables in Arabidopsis thaliana.
810	*Plant, Cell & Environment*, *45*(11), 3290–3304. https://doi.org/10.1111/pce.14417
811	Johnson, T. C., Scholz, C. A., Talbot, M. R., Kelts, K., Ricketts, R. D., Ngobi, G., Beuning, K.,
812	Ssemmanda, I., & McGill, J. W. (1996). Late Pleistocene Desiccation of Lake Victoria
813	and Rapid Evolution of Cichlid Fishes. *Science*, *273*(5278), 1091–1093.
814	https://doi.org/10.1126/science.273.5278.1091
815	Karasov, T. L., Neumann, M., Shirsekar, G., Monroe, G., Team, P., Weigel, D., & Schwab, R.
816	(2022). *Drought selection on Arabidopsis populations and their microbiomes* (p.
817	2022.04.08.487684). bioRxiv. https://doi.org/10.1101/2022.04.08.487684
818	Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N.
819	E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's
820	land surface areas. *Scientific Data*, *4*(1), Article 1.
821	https://doi.org/10.1038/sdata.2017.122
822	Kawakatsu, T., Huang, S. C., Jupe, F., Sasaki, E., Schmitz, R. J., Urich, M. A., Castanon, R.,
823	Nery, J. R., Barragan, C., He, Y., Chen, H., Dubin, M., Lee, C.-R., Wang, C., Bemm, F.,
824	Becker, C., O'Neil, R., O'Malley, R. C., Quarless, D. X., … Ecker, J. R. (2016).

Epigenomic Diversity in a Global Collection of Arabidopsis thaliana Accessions. *Cell*, *166*(2), 492–505. https://doi.org/10.1016/j.cell.2016.06.044

Kearney, M., & Porter, W. (2009). Mechanistic niche modelling: Combining physiological and spatial data to predict species' ranges. *Ecology Letters*, *12*(4), 334–350. https://doi.org/10.1111/j.1461-0248.2008.01277.x

Keitt, T. H., Lewis, M. A., & Holt, R. D. (2001). Allee effects, invasion pinning, and species' borders. *The American Naturalist*, *157*(2), 203–216. https://doi.org/10.1086/318633

Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a Species' Range. *The American Naturalist*, *150*(1), 1–23.

Koornneef, M., & Meinke, D. (2010). The development of Arabidopsis as a model plant. *The Plant Journal*, *61*(6), 909–921. https://doi.org/10.1111/j.1365-313X.2009.04086.x

Korves, T. M., Schmid, K. J., Caicedo, A. L., Mays, C., Stinchcombe, J. R., Purugganan, M. D., & Schmitt, J. (2007). Fitness Effects Associated with the Major Flowering Time Gene FRIGIDA in Arabidopsis thaliana in the Field. *The American Naturalist*, *169*(5), E141–E157.

Lambeck, K., Rouby, H., Purcell, A., Sun, Y., & Sambridge, M. (2014). Sea level and global ice volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National Academy of Sciences*, *111*(43), 15296–15303. https://doi.org/10.1073/pnas.1411762111

Lasky, J. R., Des Marais, D. L., Lowry, D. B., Povolotskaya, I., McKay, J. K., Richards, J. H., Keitt, T. H., & Juenger, T. E. (2014). Natural variation in abiotic stress responsive gene expression and local adaptation to climate in Arabidopsis thaliana. *Molecular Biology and Evolution*, *31*(9), 2283–2296. https://doi.org/10.1093/molbev/msu170

Lasky, J. R., Forester, B. R., & Reimherr, M. (2018). Coherent synthesis of genomic associations with phenotypes and home environments. *Molecular Ecology Resources*, *18*(1), 91–106. https://doi.org/10.1101/051110

Lasky, J. R., Hooten, M. B., & Adler, P. B. (2020). What processes must we understand to forecast regional-scale population dynamics? *Proceedings of the Royal Society B: Biological Sciences*, *287*(1940), 20202219. https://doi.org/10.1098/rspb.2020.2219

Lasky, J. R., Upadhyaya, H. D., Ramu, P., Deshpande, S., Hash, C. T., Bonnette, J., Juenger, T. E., Hyma, K., Acharya, C., Mitchell, S. E., Buckler, E. S., Brenton, Z., Kresovich, S., & Morris, G. P. (2015). Genome-environment associations in sorghum landraces predict adaptive traits. *Science Advances*, *1*(6), e1400218. https://doi.org/10.1126/sciadv.1400218

Lee, C.-R., Svardal, H., Farlow, A., Exposito-Alonso, M., Ding, W., Novikova, P., Alonso-Blanco, C., Weigel, D., & Nordborg, M. (2017). On the post-glacial spread of human commensal Arabidopsis thaliana. *Nature Communications*, *8*(1), 14458. https://doi.org/10.1038/ncomms14458

Leemans, R., & Cramer, W. P. (1991). The CLIMATE data base of monthly normals for globally gridded climatic variables (Version 2.1) [Data set]. In *Potsdam Institute for Climate Impact Research*. PANGAEA. https://doi.org/10.1594/PANGAEA.759491

Lee-Yaw, J. A., Kharouba, H. M., Bontrager, M., Mahony, C., Csergő, A. M., Noreen, A. M. E., Li, Q., Schuster, R., & Angert, A. L. (2016). A synthesis of transplant experiments and ecological niche models suggests that range limits are often niche limits. *Ecology Letters*, *19*(6), 710–722. https://doi.org/10.1111/ele.12604

Lee-Yaw, J. A., McCune, J., Pironon, S., & Sheth, S. N. (2021). Species distribution models rarely predict the biology of real populations. *Ecography*, *n/a*(n/a). https://doi.org/10.1111/ecog.05877

Li, B., Gao, Z., Liu, X., Sun, D., & Tang, W. (2019). Transcriptional Profiling Reveals a Time-of-Day-Specific Role of REVEILLE 4/8 in Regulating the First Wave of Heat Shock–Induced Gene Expression in Arabidopsis. *The Plant Cell*, *31*(10), 2353–2369. https://doi.org/10.1105/tpc.19.00519

Li, G., Du, S., & Guo, K. (2015). Evaluation of Limiting Climatic Factors and Simulation of a
Climatically Suitable Habitat for Chinese Sea Buckthorn. *PLoS ONE, 10*(7), e0131659.
https://doi.org/10.1371/journal.pone.0131659

Liu, C., Wolter, C., Xian, W., & Jeschke, J. M. (2020). Species distribution models have limited
spatial transferability for invasive species. *Ecology Letters*, *23*(11), 1682–1692.
https://doi.org/10.1111/ele.13577

Lopez, L., Bellis, E. S., Wafula, E., Hearne, S. J., Honaas, L., Ralph, P. E., Timko, M. P.,
Unachukwu, N., dePamphilis, C. W., & Lasky, J. R. (2019). Transcriptomics of host-
specific interactions in natural populations of the parasitic plant purple witchweed (Striga
hermonthica). *Weed Science*, *67*(4), 397–411. https://doi.org/10.1017/wsc.2019.20

Lovell, J. T., Juenger, T. E., Michaels, S. D., Lasky, J. R., Platt, A., Richards, J. H., Yu, X.,
Easlon, H. M., Sen, S., & McKay, J. K. (2013). Pleiotropy of FRIGIDA enhances the
potential for multivariate adaptation. *Proceedings of the Royal Society B: Biological*
Sciences, *280*(1763), 20131043. https://doi.org/10.1098/rspb.2013.1043

Ludlow, M. M. (1989). Strategies of response to water stress. In K. H. Kreeb, H. Richter, & T. M.
Hinckley (Eds.), *Structural and Functional Responses to Environmental Stresses* (pp.
269–281). SPB Academic.

Mandáková, T., Thorbjörnsson, H., Pisupati, R., Reichardt, I., Lysak, M. A., & Anamthawat-
Jónsson, K. (2017). Icelandic accession of Arabidopsis thaliana confirmed with
cytogenetic markers and its orign inferred from whole-genome sequencing. *Icelandic*
Agricultural Sciences, *30*, 29–38. https://doi.org/10.16886/IAS.2017.03

Martínez-Berdeja, A., Stitzer, M. C., Taylor, M. A., Okada, M., Ezcurra, E., Runcie, D. E., &
Schmitt, J. (2020). Functional variants of DOG1 control seed chilling responses and
variation in seasonal life-history strategies in Arabidopsis thaliana. *Proceedings of the*
National Academy of Sciences, *117*(5), 2526–2534.
https://doi.org/10.1073/pnas.1912451117

Merow, C., Latimer, A. M., Wilson, A. M., McMahon, S. M., Rebelo, A. G., & Silander Jr, J. A.
(2014). On using integral projection models to generate demographically driven
predictions of species' distributions: Development and validation using sparse data.
Ecography, *37*(12), 1167–1183. https://doi.org/10.1111/ecog.00839

Monroe, J. G., McGovern, C., Lasky, J. R., Grogan, K., Beck, J., & McKay, J. K. (2016).
Adaptation to warmer climates by parallel functional evolution of CBF genes in
Arabidopsis thaliana. *Molecular Ecology*, *25*(15), 3632–3644.
https://doi.org/10.1111/mec.13711

Muscarella, R., Galante, P. J., Soley-Guardia, M., Boria, R. A., Kass, J. M., Uriarte, M., &
Anderson, R. P. (2014). ENMeval: An R package for conducting spatially independent
evaluations and estimating optimal model complexity for Maxent ecological niche
models. *Methods in Ecology and Evolution*, *5*(11), 1198–1205.
https://doi.org/10.1111/2041-210X.12261

Peterson, A., Soberón, J., Pearson, R., Anderson, R., Martínez-Meyer, E., Nakamura, M., &
Araújo, M. (2011). Ecological Niches and Geographic Distributions. In *Monographs in*
Population Biology (Vol. 49). https://doi.org/10.1515/9781400840670

Pheloung, P. (1999). Preventing the introduction of potential new weeds to Australia. *Practical*
Weed Management: Protecting Agriculture and the Environment. 10th Biennial Noxious
Weeds Conference, Ballina, Australia, 20th-22nd July 1999., 18–24.

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species
geographic distributions. *Ecological Modelling*, *190*(3), 231–259.
https://doi.org/10.1016/j.ecolmodel.2005.03.026

Prentice, I. C., Sykes, M. T., Lautenschlager, M., Harrison, S. P., Denissenko, O., & Bartlein, P.
J. (1993). Modelling Global Vegetation Patterns and Terrestrial Carbon Storage at the

Last Glacial Maximum. *Global Ecology and Biogeography Letters*, *3*(3), 67–76.
https://doi.org/10.2307/2997548
Price, T., & Schluter, D. (1991). On the Low Heritability of Life-History Traits. *Evolution*, *45*(4),
853–861. https://doi.org/10.2307/2409693
Pulliam, H. r. (2000). On the relationship between niche and distribution. *Ecology Letters*, *3*(4),
349–361. https://doi.org/10.1046/j.1461-0248.2000.00143.x
Rudman, S. M., Greenblum, S., Hughes, R. C., Rajpurohit, S., Kiratli, O., Lowder, D. B.,
Lemmon, S. G., Petrov, D. A., Chaston, J. M., & Schmidt, P. (2019). Microbiome
composition shapes rapid genomic adaptation of Drosophila melanogaster. *Proceedings*
of the National Academy of Sciences, *116*(40), 20025–20032.
https://doi.org/10.1073/pnas.1907787116
Samis, K. E., & Eckert, C. G. (2007). Testing the abundant center model using range-wide
demographic surveys of two coastal dune plants. *Ecology*, *88*(7), 1747–1758.
https://doi.org/10.1890/06-1153.1
Scholl, R. L., May, S. T., & Ware, D. H. (2000). Seed and Molecular Resources for Arabidopsis.
Plant Physiology, *124*(4), 1477–1480. https://doi.org/10.1104/pp.124.4.1477
Sinnwell, J. P., Therneau, T. M., & Schaid, D. J. (2014). The kinship2 R Package for Pedigree
Data. *Human Heredity*, *78*(2), 91–93. https://doi.org/10.1159/000363105
Takou, M., Wieters, B., Kopriva, S., Coupland, G., Linstädter, A., & De Meaux, J. (2019). Linking
genes with ecological strategies in Arabidopsis thaliana. *Journal of Experimental Botany*,
70(4), 1141–1151. https://doi.org/10.1093/jxb/ery447
Therneau, T. M., & Therneau, M. T. M. (2015). Package 'coxme.' *R Package Version*, *2*(5).
Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C.,
Erasmus, B. F. N., de Siqueira, M. F., Grainger, A., Hannah, L., Hughes, L., Huntley, B.,
van Jaarsveld, A. S., Midgley, G. F., Miles, L., Ortega-Huerta, M. A., Townsend
Peterson, A., Phillips, O. L., & Williams, S. E. (2004). Extinction risk from climate
change. *Nature*, *427*(6970), 145–148. https://doi.org/10.1038/nature02121
Toledo, B., Marcer, A., Méndez-Vigo, B., Alonso-Blanco, C., & Picó, F. X. (2020). An ecological
history of the relict genetic lineage of Arabidopsis thaliana. *Environmental and*
Experimental Botany, *170*, 103800. https://doi.org/10.1016/j.envexpbot.2019.103800
Tozer, B., Sandwell, D. T., Smith, W. H. F., Olson, C., Beale, J. R., & Wessel, P. (2019). Global
Bathymetry and Topography at 15 Arc Sec: SRTM15+. *Earth and Space Science*, *6*(10),
1847–1864. https://doi.org/10.1029/2019EA000658
Turner, K. G., Fréville, H., & Rieseberg, L. H. (2015). Adaptive plasticity and niche expansion in
an invasive thistle. *Ecology and Evolution*, *5*(15), 3183–3197.
https://doi.org/10.1002/ece3.1599
Vasseur, F., Sartori, K., Baron, E., Fort, F., Kazakou, E., Segrestin, J., Garnier, E., Vile, D., &
Violle, C. (2018). Climate as a driver of adaptive variations in ecological strategies in
Arabidopsis thaliana. *Annals of Botany*, *122*(6), 935–945.
https://doi.org/10.1093/aob/mcy165
Warner, R. M., & Erwin, J. E. (2005). Naturally occurring variation in high temperature induced
floral bud abortion across Arabidopsis thaliana accessions. *Plant, Cell & Environment*,
28(10), 1255–1266. https://doi.org/10.1111/j.1365-3040.2005.01361.x
Whitlock, M. C., & Guillaume, F. (2009). Testing for Spatially Divergent Selection: Comparing
QST to FST. *Genetics*, *183*(3), 1055–1063. https://doi.org/10.1534/genetics.108.099812
Wilczek, A. M., Roe, J. L., Knapp, M. C., Cooper, M. D., Lopez-Gallego, C., Martin, L. J., Muir,
C. D., Sim, S., Walker, A., Anderson, J., Egan, J. F., Moyers, B. T., Petipas, R.,
Giakountis, A., Charbit, E., Coupland, G., Welch, S. M., & Schmitt, J. (2009). Effects of
Genetic Perturbation on Seasonal Life History Plasticity. *Science*, *323*(5916), 930–934.

Witsen, D. (2012). *Djibouti Biodiversity and Tropical Forests: 118/119 Assessment* (p. 56).
United States Agency for International Development.
https://usaidgems.org/Documents/FAA&Regs/FAA118119/Djibouti2012.pdf
Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. *Journal of the Royal Statistical*
Society: Series B (Statistical Methodology), *73*(1), 3–36. https://doi.org/10.1111/j.1467-
9868.2010.00749.x
Zacchello, G., Bomers, S., Böhme, C., Postma, F. M., & Ågren, J. (2022). Seed dormancy
varies widely among Arabidopsis thaliana populations both between and within
Fennoscandia and Italy. *Ecology and Evolution*, *12*(3), e8670.
https://doi.org/10.1002/ece3.8670
Zeng, L., Gu, Z., Xu, M., Zhao, N., Zhu, W., Yonezawa, T., Liu, T., Qiong, L., Tersing, T., Xu, L.,
Zhang, Y., Xu, R., Sun, N., Huang, Y., Lei, J., Zhang, L., Xie, F., Zhang, F., Gu, H., …
Zhong, Y. (2017). Discovery of A high-altitude ecotype and ancient lineage of
Arabidopsis thaliana from Tibet. *Science Bulletin*, *62*(24), 1628–1630.
https://doi.org/10.1016/j.scib.2017.10.007
Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association
studies. *Nat Genet*, *44*(7), 821–824. https://doi.org/10.1038/ng.2310
Zou, Y.-P., Hou, X.-H., Wu, Q., Chen, J.-F., Li, Z.-W., Han, T.-S., Niu, X.-M., Yang, L., Xu, Y.-C.,
Zhang, J., Zhang, F.-M., Tan, D., Tian, Z., Gu, H., & Guo, Y.-L. (2017). Adaptation of
Arabidopsis thaliana to the Yangtze River basin. *Genome Biology*, *18*(1), 239.
https://doi.org/10.1186/s13059-017-1378-9

[single column]
Figure 1. (A**)** Habitat suitability (green) under current climate conditions with thinned
occurrences (n = 662) used in fitting shown as black circles and (B) limiting factors from the
MaxEnt model fit to current climates. Regions far (>500 km) from known occurrence have a gray
mask in (A). For reference, regions of very low suitability (less than 0.2) in (B) are marked with
gray 'x' symbols. Equal Earth projection was used. Abbreviations in (B) as follows: precipitation
of the driest quarter "Prec driest Q", precipitation of the wettest quarter "Prec wettest Q",
precipitation seasonality "Prec seasonality", mean temperature of the warmest quarter "Mean
temp wrm Q", mean temperature of the wettest quarter "Mean temp wettest Q", temperature
annual range "Temp ann range", minimum temperature of the coldest month "min temp coldest
month.
A.

1014
B.

[single column]
Figure 2. Predicted current habitat suitability compared with individual plant size (A & B) and
genetic variation in a measure of life history (C). Size measures include inflorescence height (A,
relationship with suitability: Pearson's r = 0.12, Spearman's rho = 0.10) and maximum rosette
leaf length (B, r = 0.25, rho = 0.26) from herbarium specimens. Flowering time of natural
accessions (C, r = 0.21) was taken from published data on a growth chamber experiment at
10ºC (Alonso-Blanco et al., 2016). Linear model fits are shown. N = 2053 for inflorescence
height, N = 1179 for maximum rosette leaf length, and N=953 for flowering time.
1024

1025
1026

1027 [single column]
1028 **Figure 3.** Distribution of predicted habitat suitability (based on our native-range MaxEnt model,
1029 underlying map surfaces in green) for GBIF occurrences in various regions (black circles). We
1030 do not include a histogram for South America because there are too few occurrences. Equal
1031 Earth projection was used.

1032
1033

[single column]
Figure 4. (A) Predicted distribution during the Last Glacial Maximum and (B) areas in black with
suitability > 0.25 during both the LGM and current conditions as well as within 500 km of known
current occurrences. Because of their greater level at the LGM, in (A) we masked the LGM
Caspian and Aral Seas (including regions of high putative suitability) from the map (Prentice et
al., 1993). Lake Victoria was left unmasked as it was likely very low during the LGM (Johnson et
al., 1996). Equal Earth projection was used. For (B), the threshold of 0.25 was chosen for
reference as it approximately matches the current distribution of Arabidopsis; the alternate
thresholds for visualization of 0.2 and 0.3 and be found in Figures S14 and S15.

1043
1044

[single column]
Figure 5. Future predicted suitability for Arabidopsis (A), change in suitability from future
compared with present (blue indicates improving suitability and red decreasing, B), regions of
potential colonization (blue) continued occupancy (gray), and extinction (red) based on a
threshold suitability of 0.25 for occupancy (C), and the standard deviation in suitability among
the 5 tested climate models giving uncertainty (D). Equal Earth projection is used. RCP 4.5
emissions scenario is shown, see Fig S13 for highly similar patterns under RCP 6.0. For (C), the
threshold of 0.25 was chosen for reference as it approximately matches the current distribution
of Arabidopsis; the alternate thresholds for visualization of 0.2 and 0.3 and be found in Figure
S16 and S17.

1055
1056
1057
1058