1 **Climate biogeography of *Arabidopsis thaliana:* linking distribution models and individual**
2 **variation**
3

4 Christina Yim[1], Emily S. Bellis[1,2], Victoria L. DeLeo[1], Diana Gamba[1], Robert Muscarella[3],
5 Jesse R. Lasky[1,*]
6
7 [1]Department of Biology
8 Pennsylvania State University
9 University Park
10 USA
11
12 [2]Department of Computer Science
13 Arkansas State University
14 Jonesboro
15 USA
16
17 [3]Plant Ecology and Evolution
18 Evolutionary Biology Centre
19 Uppsala University
20 Uppsala
21 Sweden
22
23 *Corresponding Author
24 lasky@psu.edu

25
26 **Biosketch**
27 The research team is interested in understanding how individual-level processes influence
28 evolutionary and ecological dynamics at continental scales.
29
30 **Author contributions**
31 All authors contributed to study design. CY, ESB, VLD, DG, and JRL conducted analyses. CY
32 and JRL led the writing with contributions from all authors.

Abstract

AIM

The role of environmental conditions in limiting species distributions is often hypothesized in biogeography, but it is challenging to gather large-scale data to demonstrate environmental impacts on individual performance. The past and present biogeography of model organisms is key context to understanding how environment shapes species' genetic and phenotypic diversity.

LOCATION

Global

TAXON

Arabidopsis thaliana ("Arabidopsis")

METHODS

We fit occurrence records to climate data, and then projected the distribution of Arabidopsis under last glacial maximum, current, and future climates. We confronted model predictions with individual performance measured on 2,194 herbarium specimens, and we asked whether predicted suitability was associated with life-history and genomic variation measured on 898 natural accessions.

RESULTS

The most important climate variables constraining the Arabidopsis distribution were winter cold in northern and high elevation regions and summer heat in southern regions. Herbarium specimens from regions with lower habitat suitability in both northern and southern regions were smaller, supporting the hypothesis that the distribution of Arabidopsis is constrained by climate-associated factors. Climate anomalies partly explained interannual variation in herbarium specimen size, but these did not closely correspond to local limiting factors identified in the distribution model. Late-flowering genotypes were absent from the lowest suitability regions, suggesting slower life histories are only viable closer to the center of the realized niche. We identified glacial refugia farther north than previously recognized, as well as refugia concordant with previous population genetic findings. Lower latitude populations, known to be genetically distinct, are most threatened by future climate change. The recently colonized range of Arabidopsis was well-predicted by our native-range model applied to certain regions but not others, suggesting it has colonized novel climates.

MAIN CONCLUSIONS

Integration of distribution models with performance data from vast natural history collections is a route forward for testing biogeographical hypotheses about species distributions and their relationship with evolutionary fitness across large scales.

Introduction

A major goal in biology and biogeography is to understand how environmental conditions limit the performance of individuals and species distributions. First, the environment-distribution relationship can help project species distributions under past climates to understand their ecological and evolutionary history (Forester et al., 2013). Second, the environment-distribution relationship can identify regions with currently suitable habitat but that are unoccupied for a variety of reasons (e.g. dispersal limitation) (Elith et al., 2010). Third, the environment-distribution relationship can predict how distributions will shift under future environments (Thomas et al., 2004), which is an urgent task due to rapid anthropogenic climate change.

One avenue to advance these goals in future biogeographic studies is through greater integration of biogeography with organismal biology to test hypotheses about the organismal and population mechanisms controlling distributions. For example, it is becoming possible to collect molecular genetic and phenotypic data on large numbers of organisms across a species range. These approaches are particularly accessible (and valuable) in studies on model organisms like *Arabidopsis thaliana*. With model systems researchers can link detailed information on genetics and organismal biology with population and community processes (Rudman et al., 2019; Takou et al., 2019) to contribute ecological context and advance integrative biology. At the same time, advances in the organization and digitization of museum specimens are rapidly expanding the available data on range-wide variation among individuals from nature. In this spirit we focus here on *Arabidopsis thaliana,* a small annual plant (hereafter referred to as "Arabidopsis") (Koornneef & Meinke, 2010). Arabidopsis has been key to understanding how molecular biology and physiology are linked to ecology, but past biogeographic studies did not use newer distribution modeling tools (Hoffmann, 2002) or overlooked large parts of its range (Banta et al., 2012; Zou et al., 2017).

Environment-distribution relationships fundamentally arise from processes acting on individuals (Clark, 2010). In general, transplant experiments show that individual performance tends to decline outside species' natural geographic range (Hargreaves et al., 2014) and efforts to integrate information at the individual level into distribution models are emerging (Buckley et al., 2011; Elith et al., 2010; Lasky et al., 2020; Merow et al., 2014; Samis & Eckert, 2007). Where populations inhabit harsh environments (e.g. at range margins), local adaptations can emerge, such as life history changes to tolerate or escape harsh periods (Bontrager et al., 2021). In Arabidopsis, there is evidence that local adaptation to environment involves genetic changes in life history (e.g. flowering time) (Lovell et al., 2013; Martínez-Berdeja et al., 2020). However, it is challenging to determine how environment shapes individual performance and life history variation across the ranges of broadly distributed species, as well as the consequences for distributions.

A vast resource of individual-level information can be found in natural history collections (Heberling, 2021; Lopez et al., 2019). For example, Bontrager & Angert (2015) showed with herbarium specimens of *Clarkia* that fecundity decreased with drier summers, and toward the western species range margin both summer precipitation and individual fecundity declined, suggesting a mechanism limiting its distribution. In Arabidopsis, DeLeo et al. (2020) found decadal shifts in traits of herbarium specimens. For many species, seed banks house great diversity from across their ranges that can also be used to study their biogeography (Estarague

et al., 2021; Scholl et al., 2000). Here we estimate performance and life history variation from
museum and seed bank accessions to interrogate predictions from distribution models.
Arabidopsis is native across Eurasia and Africa and with human assistance has
colonized the Americas and Australia. Other species in the genus *Arabidopsis* are more
restricted to cool temperate climates, with Arabidopsis having expanded to a broader range, *e.g.*
Mediterranean habitats (Hoffmann, 2005). Arabidopsis can behave as a spring annual with a
rapid life cycle, germinating in the spring and flowering in the late spring and summer. However,
many individuals are longer-lived winter annuals, germinating in the fall and flowering in early
spring (Wilczek et al., 2009). Studies of large-scale environmental response in Arabidopsis have
focused on the role of local adaptation in genetic diversity in the species (e.g. Hancock et al.,
2011; Lasky et al., 2018; Martínez-Berdeja et al., 2020; Toledo et al., 2020), but less is known
about the determinants of the species' distribution. The last overview of the climate
biogeography of Arabidopsis was Hoffmann (2002), who considered Arabidopsis native to
western and central Eurasia, but non-native in China and sub-Saharan Africa. However, studies
show these latter populations are likely native, being genetically diverse, and with many unique
genetic variants (Durvasula et al., 2017; Zou et al., 2017). Advances in distribution modeling
(Elith et al., 2010; Muscarella et al., 2014) and updates to climate and occurrence datasets
highlight the need to update to our understanding of Arabidopsis biogeography.
Studies suggest that multiple climatic factors limit performance and distribution of
Arabidopsis. Experiments have shown that winter cold is a major factor limiting performance
(Ågren & Schemske, 2012; Korves et al., 2007). Additionally, Arabidopsis lacks physiological
traits for dealing with severe water deficit so it is likely individual performance in nature is limited
by drought (Clauw et al., 2015). Based on occurrence data, Hoffmann (2002) concluded that
spring and fall cold limited Arabidopsis in northern Europe, winter temperatures in Russia, heat
and drought toward the southern range margins, and a lack of snow in central Asia. However,
these conclusions were based on visual inspection of climate isotherms and range margins, not
statistical inference. Distribution models fit to occurrence data allow an explicit quantitative
statement of environment-distribution relationships and the potential to infer the environmental
factors limiting distributions (Elith et al., 2010; G. Li et al., 2015).
Distributions are dynamic through time due to environmental change, dispersal, and
demographic stochasticity. Studies have used genetic data to infer Arabidopsis refugia during
the last glacial maximum (LGM), where subpopulations (sometimes referred to as "relicts")
persisted locally before subsequent admixture with an expanding, now widely distributed "non-
relict" lineage (Lee et al., 2017). Whether these refugia corresponded to suitable climates is less
clear. Future climate change projected impacts on Arabidopsis have focused on *relative* climate
impacts on different genotypes (Exposito-Alonso et al., 2018; Fournier-Level et al., 2016), rather
than distribution dynamics. Additionally, many species, have colonized new regions due to
human introduction, sometimes exhibiting traits that appear distinct from native range
populations, potentially due to plastic or genetic responses to new environments (Turner et al.,
2015). Arabidopsis has colonized many regions, but it is unclear to what degree these places
represent novel environments.
Here we sought to demonstrate an approach to integrate distribution modeling with
organismal biology and intraspecific variation, and to revise our understanding of Arabidopsis's
biogeography. To do so, we combined occurrences (including many outside Europe poorly

represented in previous work) with climate data to build distribution models, and then test how
model predictions correspond to performance estimated from herbarium specimens and genetic
variation in natural accessions. We ask the following questions:

1. What climate factors constrain the distribution of Arabidopsis? We hypothesize that winter cold and summer drought are the most important constraints, depending on region.
2. Are individual performance and genetic variation associated with model-estimated habitat suitability? We hypothesize that individuals reach larger sizes in regions with greater suitability and that populations adapt along gradients in suitability through changes in traits such as flowering time, a key component of life history.
3. Are occurrences outside the native range predicted by a native range model, suggesting stable realized niches following colonization? Or is there evidence Arabidopsis has colonized novel environments?
4. Where did Arabidopsis persist during the Last Glacial Maximum (LGM)? And where will Arabidopsis move in future climates?

Methods

Occurrence data

We developed a set of high-quality occurrence data (*i.e.* species ID verified and location checked, N=4,024) from published research (Durvasula et al., 2017; Hsu et al., 2019; Mandáková et al., 2017; Zeng et al., 2017; Zou et al., 2017), publicly available herbarium and germplasm accessions with known collection locations (Alonso-Blanco et al., 2016; DeLeo et al., 2020), and some of our recent field collections in East Africa (Gamba et al., 2022). These span a period of 1794 - 2018. The herbarium specimens and new collections include little-studied populations in Saudi Arabia, Somalia, Djibouti, Eritrea, Rwanda, Ethiopia, Uganda, Sudan, and Nepal. Duplicate occurrence points were eliminated (samples are often split and sent to different herbaria). For model fitting, we excluded occurrences from putative non-native regions (the Americas, New Zealand, Japan).

We also used occurrence data (with coordinates and without flagged problems N=115,226) from the Global Biodiversity Information Facility (GBIF) to test model predictions in regions outside of the native range of Arabidopsis (downloaded 30 Dec 2020, Gbif.Org, 2020). We deem these occurrences as lower quality given that many have not had the species identity and location checked (DeLeo et al., 2020).

Environmental data

Climate data were extracted from CHELSA (Climatologies at High resolution for the Earth's Land Surface Areas) v1.2 at 30 arc second spatial resolution (Karger et al., 2017). Current conditions are the average of 1979-2013 estimates. We selected the following climate variables based on hypothesized importance (Gienapp et al., 2017; Hancock et al., 2011; Lasky et al., 2014, 2018) and relatively low inter-correlation (Pearson correlation coefficients among variables at occurrences < 0.75): isothermality (Bio3), minimum temperature of coldest month (Bio6), temperature annual range (Bio7), mean temperature of wettest quarter (Bio8), mean temperature of the warmest quarter (Bio10), precipitation seasonality (Bio15), precipitation of

wettest quarter (Bio16), and precipitation of driest quarter (Bio17). We also included altitude
from Hijmans et al. (2005).
For projecting past distributions, we obtained climate estimates from the last glacial
maximum (LGM) at 21k yrs before present from CHELSA PMIP3 (Karger et al., 2017). We used
the global altitude and bathymetry map with 15 arc second resolution from Tozer et al. (2019)
with a sea level 134 m lower than present (Lambeck et al., 2014) to project potential suitable
habitat at the LGM on land in areas currently submerged. For projecting future distributions, we
used climate projections five divergent global climate models for 2050 from CHELSA v1.2 using
the RCP 4.5 emissions scenario (Karger et al., 2017). We also show RCP 6.0 in the supplement
for context (Figure S13), though it is highly similar to 4.5 in the target time period.
To characterize temporal variation in climate (climate anomalies), we used the Climate
Research Unit (CRU) TS 4.01 dataset, providing a global time series of monthly temperature
and precipitation for the period 1900-2010 at a 0.5º resolution (Harris et al., 2014). From the
CRU data we calculated the same bioclimatic parameters that we used from CHELSA, but in
the CRU data these bioclimatic variables were specific to each herbarium specimen in the time
period it was collected (Supplemental Methods). We then calculated local anomalies for each of
these variables by taking the observed value, subtracting mean across the entire time series,
and dividing by the standard deviation (DeLeo et al., 2020).
225
Performance estimates from herbarium specimens
We estimated fecundity on a subset of herbarium specimens using two traits. First, we
measured the length of the longest inflorescence, reasoning that longer inflorescences would
have more fruits and seeds. Second, we measured maximum rosette leaf length, reasoning that
larger rosettes would support later reproductive investment if these collected plants were
allowed to continue growth *in situ* (see Supplement for a validation). We used ImageJ on 2,194
specimen images to estimate the tallest point of each inflorescence (N=2,188) and the
maximum rosette leaf length (N=1,264; see Supplement).
234
Range-wide genetic variation in life history
Many late-flowering Arabidopsis genotypes require cold cues (vernalization) to flower and also
show slower growth and more stress tolerance, delineating a life history axis (Lovell et al., 2013;
Vasseur et al., 2018). To assess life history variation, we used published experimental data on
flowering time for 898 whole-genome resequenced accessions from the native range with
reliable geographic coordinates grown at 10 and 16ºC (The 1001 Genomes Consortium 2016).
To estimate vernalization sensitivity, we calculated the difference between flowering time at 10
and 16ºC.
243
Species Distribution Modeling
We thinned the original 4,024 high quality occurrence points to one sample per 1 km grid cell to
reduce sampling bias (N=662) using the 'sp' package (Bivand et al., 2008). To characterize
potentially inhabited sites, we generated pseudoabsence background points using the 'dismo'
package (Hijmans & Elith, 2013) by sampling 10,000 random points within a 500 km buffer
around occurrence points.

We used Maxent version 3.4.0 to generate a species distribution model (Phillips et al.,
2006). MaxEnt was implemented with the R package 'ENMeval' v2, and parameters were
optimized using the 'checkerboard2' method for cross validation (Muscarella et al., 2014).
Among the tested settings (ENMeval defaults), we chose the model with lowest AICc value and
used this to project habitat suitability under recent conditions across the globe. For all models
we used the logistic output of MaxEnt that scales suitability from zero to unity.
We used permutation importances to determine which climatic factors drove predictions
in the distribution model. We also used the 'limiting' function in the R package 'rmaxent' to
determine the most limiting climatic factors in each location, defined as the environmental
covariate that has the largest decrease in suitability at a given location relative to the suitability
achieved if the covariate had its value equal to the global mean (Baumgartner et al., 2017; Elith
et al., 2010). State another way, the local limiting factor is the environmental condition most
limiting suitability, compared to an alternative scenario where that condition takes its global
mean.
In non-native regions, we evaluated whether Arabidopsis is limited from further
expansion at range edges by climate, *i.e.* whether there were no more unoccupied suitable
environments near existing populations. To do so, we calculated suitability in a zone 50-100 km
from existing GBIF occurrences and compared occupancy in these buffers in the native range to
invaded regions.
We also projected the MaxEnt model using past (LGM) and future climate conditions.
For future conditions, we calculated the mean and standard deviation of habitat suitability
projected for the five climate models. To assess whether model predictions were extrapolating
into poorly characterized or novel climates, we compared the present-day model training
climates to each predicted climate conditions, calculating multivariate environmental similarity
surfaces (MESS) following Elith et al. (2010). Higher values on a MESS map indicate conditions
in a location (or point in time) are similar to the reference environmental conditions used to fit
the model. Negative values indicate that at least some variables are outside the range of
environments used to fit the model, signifying extrapolation into novel environments (Elith et al.,
2010).
279
Performance and habitat suitability
We asked whether suitability corresponded to plant size. We first tested these relationships with
Pearson's and Spearman's correlations. We also fit Generalized Additive Models (GAMs) where
herbarium specimen sizes were the response variable. The model included covariates for
suitability at the collection location (square-root transformed to reduce the lower-tail influence)
and (as a nuisance variable) the year of collection (scaled to mean zero and unit variance) to
account for potential changes in size over time. We used GAMs to allow smooth spatial
variation in parameters, allowing us to capture any regional variation. The model can be
represented as:
289
290        $\text{Phenotype}_{i,year} = \beta_0 + \beta_{i,year}\text{Year} + \beta_{i,HS}\text{Habitat Suitability}_i + \varepsilon_{i,year}$
291
where *i* represents the specimen location. Errors were modeled as normally distributed.
Spatially-varying parameters were constrained to smooth spatial variation using the 'mgcv'

package for R, which we used with REML to fit all GAMs (Wood, 2011). We considered
covariates to be significant at a given location if their 95% confidence interval (CI) excluded 0,
and the Moran's I (Cooper, 2021) for model residuals did not indicate spatial autocorrelation.
We also asked whether local limiting factors corresponded to climate variables with
significant effects on plant performance. We hypothesized that for a given region, temporal
fluctuations in local limiting factors would be associated with temporal variation in plant size in
herbarium specimens. To address this hypothesis, we tested if specimen sizes were correlated
with anomalies in climate conditions corresponding to the climate variables used in the MaxEnt
model. We used the yearly climate anomalies we calculated from CRU in GAMs with spatially
varying coefficients, where for a phenotype measured at location i:
304
305                      $\text{Phenotype}_{i,year} = \beta_{i,0} + \beta_{i,year}\text{Year} + \beta_{i,climate}\text{BioClim Variable Anomaly} + \varepsilon_{i,year}$
306
307   We then qualitatively assessed whether local limiting factors inferred by MaxEnt had anomalies
308   that were correlated with plant size (local $\beta_{i,climate}$ coefficients in the GAMs).
309
Genetic variation and habitat suitability
We used two strategies to test how genetic variation in Arabidopsis varies with suitability. First,
we focused on genetic variation in a key life history trait, flowering time. We estimated linear
models relating suitability and flowering time. To account for population structure and neutral
processes that can affect spatial variation in flowering time, we also tested whether suitability
was associated with flowering time after using random effects for genome-wide similarity among
accessions. A significant suitability-flowering time association in this model would suggest
selection linked to suitability acts on flowering time. This test is akin to Q_{ST}-F_{ST} contrasts
(Whitlock & Guillaume, 2009), except that an explicit environmental gradient is tested
(suitability). We implemented this mixed model the function 'lmekin' from the R package 'coxme'
(Therneau & Therneau, 2015), along with 'kinship2' (Sinnwell et al., 2014). The kinship matrix
was obtained from 2,027,463 published whole genome resequencing SNPs (Alonso-Blanco et
al., 2016). Additionally, to test for geographic variation in suitability-flowering time relationships,
we fitted GAMs of flowering time with spatially varying coefficients:
324
325                      $\text{FloweringTime}_{i,year} = \beta_0 + \beta_{i,HS}\text{Habitat Suitability} + \varepsilon_i$.
326
Our second strategy was to scan the Arabidopsis genome for genes where different
alleles were found in high versus low suitability locations. Such a change in allele frequency
across gradients in suitability would suggest that this genetic variation was involved in local
adaptation to low versus high suitability environments. We used univariate linear mixed-effects
models in GEMMA (v 0.98.3) (Zhou & Stephens, 2012) to perform genome wide association
studies (GWAS) in a set of 2,053,939 SNPs filtered for MAF=0.05 from 1003 native-range
ecotypes part of the 1001 genomes panel.
334
335
336   **Results**
337

Q1. Climate constraints on the distribution of Arabidopsis
Our optimization of MaxEnt models (with AICc) resulted in a selection of a model with linear and
quadratic effects, with the regularization multiplier of 0.5. The training AUC value was 0.78 and
the average test AUC with checkerboard2 cross validation was 0.78. The omission rate of the
10th percentile of suitability for training points was 0.10, suggesting our models were not overfit
as they were able to predict low probability occurrences as well as expected (Fielding & Bell,
1997; Peterson et al., 2011). The areas of high suitability overall correspond well to the
documented Arabidopsis distribution, with one notable exception being tropical lowland sites
(Congo basin) in sub-Saharan Africa (Figure 1A). This region was near zero multivariate
environmental similarity to training data, indicating the model may have been poorly constrained
there, while most of the regions of high predicted suitability where Arabidopsis is documented
have positive similarity (MESS, Figure S2).
350
351

Figure 1. (A) Habitat suitability (green) under current climate conditions with thinned
occurrences (n = 662) used in fitting shown as black circles and (B) limiting factors from the
MaxEnt model fit to current climates. Regions far (>500 km) from known occurrence have a gray
mask in (A). Regions of low suitability (less than 0.25) in (B) are marked with gray 'x' symbols.
Equal Earth projection was used. Abbreviations in (B) as follows: precipitation of the driest
quarter "Prec driest Q", precipitation of the wettest quarter "Prec wettest Q", precipitation
seasonality "Prec seas", mean temperature of the warmest quarter "Mean temp wrm Q", mean
temperature of the wettest quarter "Mean temp wettest Q", temperature annual range "Temp
ann range", minimum temperature of the coldest month "min temp coldest month.
A.

380 **B.**

The permutation importance of the model covariates across the entire native range revealed
that the minimum temperature of the coldest month (PI = 40%) and the mean temperature of the
warmest quarter (PI = 32%) were the two most important variables, suggesting winter cold
stress and summer heat stress are most important in constraining Arabidopsis's distribution
(Table S1). The next most important variable was isothermality (PI =14%), as Arabidopsis tends
to be found in regions with low isothermality (e.g. most temperate zones).
To identify spatial variation in climate constraints, we also identified local limiting factors
(Figure 1B). Across northern Eurasia, minimum temperature of the coldest month limited habitat
suitability. Across the Mediterranean and tropical/subtropical regions, the temperature of the
warmest quarter was limiting. Across Eastern Europe and Central Asia, winter cold was limiting
adjacent to other regions where summer heat was limiting, highlighting the multiple stressors in
this region.
We hypothesized that for a given region, temporal fluctuations in local limiting factors
would be associated with temporal variation in plant size in herbarium specimens. We found
that years with warmer minimum winter temperatures were associated with significantly taller
inflorescences from Turkey to central Asia (Figure S3). Concordantly, in much of this region the
Maxent model indicated that minimum temperature of the coldest of the month was the most
limiting factor (namely in the Caucuses and from Kazakhstan to northern India, Figure 1B). We
also found that years with greater seasonality of precipitation were associated with shorter
inflorescences in central Asia, but taller inflorescences in central Europe (Figure S3). Partly
consistent, in the southern part of this central Asian region the MaxEnt model indicated
precipitation seasonality was limiting (Figure 1B). Years with higher isothermality were positively
associated with inflorescence height in Eastern Europe but this was not a limiting factor in this
region (Figure S3). Other climate anomalies were not significantly associated with temporal
variation in plant size.
411
Q2. Declining fitness and stress escape life history in regions of low predicted habitat suitability
As habitat suitability increased, so did the inflorescence height of individual plants in herbarium
specimens (Pearson's r = 0.11, p < 10^{-7}; Spearman's rho = 0.09, p < 10^{-5}; n = 2053). The
relationship with suitability was even stronger for maximum rosette leaf length (r = 0.23, p < 10^{-16}; rho = 0.24, p < 10^{-16}; n = 1179, Figure 2), which was also more correlated with total fruit +
flower number (Supplement), suggesting fecundity was greater in regions of high predicted
suitability. We tested size-suitability relationships using GAMs with spatially varying suitability
coefficients (but not including climate anomalies – distinct from the previous section Q1). We
found a consistently positive relationship between suitability and size that was significant for
rosette leaf length in western Europe and for inflorescence height across most of Eurasia (Figs
S5 and S6). Unexpectedly, in these models where year of collection was considered primarily
as a nuisance variable, plant size significantly declined over time in northwest Europe (Figs S5
and S6; e.g. in Scandinavia, maximum rosette leaf length vs year, Spearman's rho = -0.17).
425
Figure 2. Predicted current habitat suitability compared with individual plant size (A & B) and
genetic variation in a measure of life history (C). Size measures include inflorescence height (A,
relationship with suitability: Pearson's r = 0.11, Spearman's rho = 0.09) and maximum rosette
leaf length (B, r = 0.23, rho = 0.24) from herbarium specimens. Flowering time of natural

accessions (C, r = 0.21) was taken from published data on a growth chamber experiment at
10ºC (Alonso-Blanco et al., 2016). Linear model fits are shown. N = 2053 for inflorescence
height, N = 1179 for maximum rosette leaf length, and N=953 for flowering time.
433

434
We compared habitat suitability with published data on genetic variation in flowering
time. We found that flowering time at 10ºC and flowering time plasticity were significantly
positively associated with suitability (r=0.21, p<10[-11] and r=0.10, p=0.0021, respectively), and
also for days to flower at 16 ºC (r=0.08, p=0.0166, Figures 2 & S7). In GAMs with spatially-
varying suitability coefficients, the suitability-flowering time pattern were largely consistent
across Eurasia (Figure S8). The suitability-flowering time association was significant even when
accounting for genomic similarity among accessions, suggesting selection associated with
suitability could maintain geographic clines in flowering time. Specifically, linear mixed-effects
models found a positive association for flowering time at 10ºC (p<10[-6], n=953), and also at
16ºC, the latter of which may have been obscured by population structure that was unaccounted
for in the simple linear regression model (p<10[-5], n=920). The suitability association was not
significant for plasticity in flowering time when accounting for genomic similarity (p=0.38, n=920)
(Tables S1-S3). Consistent with this potential obscuring of population genetic structure, we
found that 10 ADMIXTURE genetic clusters in Arabidopsis (Alonso-Blanco et al., 2016) were
significantly different in their habitat suitability (F(9,993)=201.6, p<10[-16], Table S5, Figure S9).
We scanned the genome for genes that showed allele frequency correlations with
suitability. The most strongly associated SNP was in the putative promoter region (878 bp from
the start) of ERF53 (AT2G20880), a transcription factor that regulates response to drought, salt,
and heat (Figure S11) (Cheng et al., 2012; B. Li et al., 2019). This SNP showed a strong allele
frequency cline from Europe to Asia, where the alternate allele was nearly fixed in accessions
east of the Ural Mountains, which is a region of low estimated suitability (Fig S10). Furthermore,
we found that this alternate SNP allele was associated with higher expression of ERF53 (Wilcox
test, p = 0.0071) in published transcriptome data (Kawakatsu et al., 2016), suggesting a *cis*-
regulatory variant is locally adapted to low suitability parts of Asia.
459
460
Q3. Using the native range model to predict outside the native range

Australia and New Zealand occurrences were largely in areas predicted to be highly suitable
(Figure 3). By contrast, North America includes occurrences in highly suitable areas (Pacific
Northwest) but many in low suitability areas (interior east, where summer heat was predicted
limiting, Figure S12). The eastern US also has near zero similarity on the MESS map,
suggesting a novel set of conditions compared to the native range (Figure S2). Similarly, the
regions where Arabidopsis occurred in Korea and Japan were of lower suitability than in the
native range, also with summer heat as the predicted limiting factor (Figure S12), suggesting
Arabidopsis in eastern North America and East Asia inhabits climates with distinctly warm
summers.
471
472 **Figure 3.** Distribution of predicted habitat suitability (based on our native-range MaxEnt model,
473 underlying map surfaces in green) for GBIF occurrences in various regions (black circles). We
474 do not include a histogram for South America because there are too few occurrences. Equal
475 Earth projection was used.

476
We investigated whether non-native Arabidopsis occupy all available habitat in their
regions or whether suitable habitat remains unoccupied. We found that locales 50-100 km from
occurrences included many areas of high suitability in North America, suggesting suitable sites
remain unoccupied (11.8% of grid cells in these 50-100 km regions had suitability > 0.6). In
comparison there were fewer such sites in the native range (5.1% of these 50-100 km regions
had suitability > 0.6). There were extensive areas of high suitability along the Pacific coast of
North America to around 60ºN, but no occurrences north of 50º, which we confirmed with an
expert botanist (pers comm. Matthew Carlson). The timing of the invasion is likely not a factor
given that occurrences from near Vancouver date at least to 1939. This region has near zero
similarity on the MESS map, potentially indicating that the native range model is not well
constrained there (Figure S2). Similarly, the southern coast of Australia is highly suitable but
Arabidopsis is apparently absent (31.9% of these 50-100 km regions with suitability > 0.8).

Records are restricted to the southeast and southwest (confirmed by botanists, Shelley James,
Tim Entwistle, Neville Walsh pers. comm.), even though records date at least to 1959 in SE
Australia. In South America there are a few Arabidopsis records from the Southern Cone, and it
appears to be rare in the region (pers. comm. Diego Salariato), while apparently suitable
environments occur throughout Patagonia and high elevation Andean sites that are apparently
unoccupied (pers comm. Gwendolyn Peyre, Santiago Madriñán). Large areas of southern
Australia and South America show positive similarity on the MESS map suggesting the model is
well constrained in those regions. By contrast in Korea and Japan there are very few sites
expected to be highly suitable that are not already occupied by Arabidopsis (4.9% of these 50-
100 km regions with suitability > 0.6).
499
Q4. The distribution of Arabidopsis during the Last Glacial Maximum and in the future
We projected suitability onto the climate at the LGM and found several putative refugia, where
past environmental conditions could have supported Arabidopsis persistence. In particular, the
Mediterranean/Caucuses/south Caspian Sea, much of sub-Sharan Africa > 1000 m (current)
asl, and China and SE Asia appear as refugia (Figure 4). North Africa, the Atlantic European
coast, and the islands of Sicily, Corsica, and Sardinia also appear as highly suitable potential
refugia.
507
508

Figure 4. (A) Predicted distribution during the Last Glacial Maximum and (B) areas in black with
suitability > 0.3 during both the LGM and current conditions as well as within 500 km of known
current occurrences. Because of their greater level at the LGM, in (A) we masked the LGM
Caspian and Aral Seas (including regions of high putative suitability) from the map (Prentice et
al., 1993). Lake Victoria was left unmasked as it was likely very low during the LGM (Johnson et
al., 1996). Equal Earth projection was used.

515
516
We projected climate suitability for Arabidopsis in the year 2050. Some current high suitability
regions will remain so, such as in Northern Europe. Nevertheless, we found poleward and up-
elevation shifts in regions of high suitability and retreats at lower latitudes and elevations. In the

native range, all African sites show declining suitability, as did most Mediterranean sites,
highlighting vulnerability of these populations. By contrast, Arctic Europe, and the mountains of
central Asia/Tibetan plateau show increased suitability, highlighting potential range expansions
(Figure 5).
524
Figure 5. Future predicted suitability for Arabidopsis (A), change in suitability from future
compared with present (blue indicates improving suitability and red decreasing, B), regions of
potential colonization (blue) continued occupancy (gray), and extinction (red) based on a
threshold suitability of 0.25 for occupancy (C), and the standard deviation in suitability among
the 5 tested climate models giving uncertainty (D). Equal Earth projection is used. RCP 4.5
emissions scenario is shown, see Fig S13 for highly similar patterns under RCP 6.0.

531
532
533
534
535   **Discussion**
536
We used the model plant Arabidopsis in a case study of integrative climate biogeography of a
species' past, present, and future distributions. This species is key to a large body of plant
biology research, and an in-depth study connecting its biogeography to genetic and phenotypic
variation provides important context for the biology of Arabidopsis. The size of individual plants

in herbarium specimens, as well as genetic variation in flowering time, suggested that regions
with lower predicted suitability harbored populations with reduced fitness and altered life history.
The consistency between model predictions and individual variation provides partial model
validation, bolstering our confidence in conclusions from model projections. Using these model
projections, we identified new glacial refugia, and looking to the future, we found that genetically
distinct lower latitude populations are most threatened by climate change.
547
The current distribution of Arabidopsis and its limiting factors
The fitted native-range model largely corresponded to occurrences and indicates Arabidopsis
broadly distributed across Europe, moister regions of central and eastern Asia, and mountains
across Africa. Our results advance beyond the most recent climate biogeographical study of
Arabidopsis by Hoffmann (2002), who used monthly climate data (but notably no synthetic
bioclimatic variables) from Leemans & Cramer (1991) mostly to qualitatively describe the
regions occupied by Arabidopsis. Our study included populations in sub-Saharan Africa and
Asia that were overlooked or considered non-native by Hoffmann (2002). We also included
occurrences from regions apparently missed by Hoffmann (2002). Zou et al. (2017) more
recently built models of Arabidopsis's distribution using Eurasian populations and default
MaxEnt settings, but did not include occurrences from Arabia, sub-Saharan Africa, and much of
the Himalayas, Russia, and central Asia, and did not subsample occurrence to reduce bias.
Likely as a result of these issues, the predictions of Zou et al. (2017) show a pronounced peak
in suitability in Germany, which was densely sampled in the genomics studies used for
occurrences by Zou et al. (2017), but predicted low suitability in much of the core European
range and near zero suitability in sub-Saharan Africa, Arabia, and most of the Russian part of
the species range.
The two dominant limiting factors in our model were winter cold (at lower latitudes and
elevations) and summer heat (at higher latitudes and elevations). Winter cold is recognized to
limit Arabidopsis performance, in particular winter cold appears to be a dominant force in local
adaptation of Arabidopsis (Ågren & Schemske, 2012; Gienapp et al., 2017; Monroe et al.,
2016). In southern Europe, where summer heat was inferred to be limiting, Arabidopsis flowers
in early spring (DeLeo et al., 2020) and thus summer heat is not usually directly experienced. In
spring in these regions, warm temperatures might not reach consistently stressful levels (e.g. to
induce fruit abortion) (Warner & Erwin, 2005) but it may be that moisture deficit driven by
evaporative demand is directly limiting in late spring. Where suitability was highest for
Arabidopsis, including the British Isles and a belt along the coast from France to Poland,
temperature annual range was identified as limiting. However, interpreting limiting factors in an
area of extremely high suitability (near one) is not meaningful given that suitability can scarcely
be increased.
Despite inference of winter cold and summer heat as primarily limiting, these were only
partly reflected by temporal fluctuations in individual plant performance from herbarium
specimens. Winter cold and precipitation variability anomalies were significantly associated with
specimen size in much of Asia, where these were also the MaxEnt modeled limiting factors from
Iran and Kazakhstan to Afghanistan and the Himalayas, suggesting these climate factors truly
limit Arabidopsis populations. However, there were discrepancies between individual
performance and limiting factors, likely for several reasons. First, the size of specimens is an

imperfect fitness proxy. Usually only reproductive individuals are collected in herbaria, and
individuals that would die before reproduction are excluded. Second, individuals are not
randomly sampled from populations (Daru et al., 2018). Third, MaxEnt models face limitations
due to misspecification, problems with occurrence data, or a mismatch between covariates and
the true ecological factors limiting populations. Nonetheless, the limiting factors we identified
here fit well with our knowledge of Arabidopsis ecophysiology and natural history.
591
Habitat suitability, individual performance, and life history
We found a decrease in plant size with decreasing habitat suitability across the range of
Arabidopsis, suggesting that our model suitability captured a substantial part of environmental
effects on individual fitness. Given the great variability in individual plant performance within
populations obvious to casual observers, it is unsurprising that suitability explains a minority of
total variation in individual size, leaving most variation unexplained (Figure 2). Furthermore,
fecundity and fecundity response to treatments often have low heritability in many species
(Price & Schluter, 1991), even in controlled experiments (Lasky et al., 2015).
600   There have still been few studies of individual performance with the geographic scope
that allows inference across a species range (Angert & Schemske, 2005; Csergő et al., 2017;
Greiser et al., 2020; Samis & Eckert, 2007). In a synthesis of studies of 40 species, Lee-Yaw et
al. (2016) found that, on average, individual performance and distribution-model inferred
suitability decline beyond range margins. From 42 studies Lee-Yaw et al. (2021) found that 38%
identified some predictive ability of distribution models for individual performance. However,
many previous studies relied on intensive observations of a small number of populations, while
our estimates of performance from thousands of herbarium specimens allowed us to cover most
of the species range. The increased availability of digitized museum specimens with trait data
indicates an opportunity to estimate performance across distributions for many species
(Bontrager & Angert, 2015).
611   We found that low suitability regions had earlier flowering genotypes. This relationship
was noisy, with early flowering genotypes frequent in all levels of suitability, but later flowering
genotypes lacking from the least suitable regions. The suitability-flowering time associations
were significant when accounting for genome-wide similarity between accessions, suggesting
they reflect selection associated with suitability. We interpret the direction of the relationship as
indicating that when suitability is low, Arabidopsis employs stress escape strategies, *i.e.* a rapid
life cycle during favorable conditions (Ludlow, 1989). Nevertheless, our findings suggest that
low suitability central Asian populations have some stress tolerating mechanisms, as they
harbor distinct allele at a transcription factor (ERF53, AT2G20880) that regulates response to
abiotic stressors (Cheng et al., 2012; B. Li et al., 2019). The restriction of late flowering
genotypes to more suitable regions is counterintuitive given physiological work showing these
are more stress tolerant (Lovell et al., 2013). However, it may be true that more favorable
conditions make possible a slow growing, slow flowering, freezing-tolerant winter annual
strategy.
625
The distribution of Arabidopsis outside its native range
Arabidopsis has spread across the globe, largely to climates well-predicted by native range
models, but to some regions predicted to be less suitable. Populations in western North

America, Australia, and New Zealand occur in climates well-predicted by the native range,
suggesting stable realized niches following colonization. But in eastern North America, Korea,
and Japan, these climates tend to be of low predicted suitability. Whether these are truly of
lower suitability is unknown without performance data. It may be that Arabidopsis has colonized
novel environments in these regions, or that the low suitability is only a model artefact. Studies
often find limited transferability among native and introduced range models (Early & Sax, 2014;
Liu et al., 2020).
In southern Australia, the Southern Cone, and Alaska there are widespread seemingly
suitable habitats where Arabidopsis is absent. The lack of more populations may partly owe to
the strict quarantine on imports to Australia to avoid species invasions (Pheloung, 1999).
Additionally, a lack of human disturbance in some regions (e.g. Alaska) may limit the potential
for colonization given Arabidopsis often is found in disturbed sites.
641
The distribution of Arabidopsis at the last glacial maximum
Using population genetic patterns, Lee et al. (2017) hypothesized five glacial refugia for Eurasia
to be in Iberia, Sicily, Balkans, the Levant, and Turkmenistan. These are consistent with our
range reconstruction, although we do not find clear barriers of unsuitable climates that would
have isolated Italian, Balkan, and Levant LGM populations. Furthermore, we note that
Sardinia/Corsica appears as a refugium, and we note that narrow, highly suitable areas along
the southern and eastern Caspian Sea could be the location of the hypothesized Turkmenistan
refugium. Lee et al. (2017) further hypothesized that the bulk of current European genotypes
derive from an expansion originating on the northwest coast of the Black Sea, where we found a
strip of highly suitable conditions for Arabidopsis during the LGM. Additional genetic studies
hypothesized a long history of existence in sub-Saharan Africa (Durvasula et al., 2017) and
conditions during the LGM suggest the species could have been much more widespread than
currently. Concordantly, Chala et al., (2017) used models of the distribution of Afroalpine habitat
generally during the LGM to find expanded areas in East Africa compared to present day,
though some locations such as Jebel Marra, Sudan were still surrounded by landscapes of
unsuitable habitat even at the LGM.
We found that the North Atlantic European coast was highly suitable (including areas
currently inhabited) during the LGM, but current British Isle and French populations do not show
genetic signatures of refugia (Lee et al., 2017). In both cases, local, distinct genotypes that
survived the LGM may have gone extinct following expansion by the now dominant European
genetic cluster ("non-relicts") (Fulgione & Hancock, 2018; Lee et al., 2017). The previous
Arabidopsis distribution modeling of the LGM (Zou et al., 2017) did not identify France and NE
Iberia as highly suitable during the LGM and did not mask the Caspian Sea (thus overinflating
that refugium). Zou et al. (2017) also did not identify the African refugia (except for a small
region in SE Africa) and potential refugia in mountains of SE Asia.
667
The future distribution of Arabidopsis
Over the next decades climate conditions are predicted to worsen for Arabidopsis across large
areas of its range. Central Spain and mountains in north and east Africa and Arabia may see
the most worsening. These climate change effects may already be emerging: our intensive field
search in the Dai Forest of Djibouti in 2018 failed to yield any Arabidopsis, despite the presence

of abundant (but likely more drought tolerant) annual mustards (Brassicaceae) *Erucastrum*
arabicum and *Sisymbrium erysimoides* in otherwise appropriate habitat for Arabidopsis*.*
Arabidopsis was collected in this juniper woodland in 1954 but recent decades have seen
increased drought and tree mortality (Witsen, 2012), and the potential extinction of this isolated
marginal Arabidopsis population. Given the unique genetic diversity of the model plant
Arabidopsis housed in its lower latitude populations (Durvasula et al., 2017; Hsu et al., 2019;
Lee et al., 2017; Zou et al., 2017) the conservation of these populations could benefit plant
biology research.
By contrast, conditions are expected to improve for the northernmost populations in
Europe, suggesting a potential current colonization front of Arabidopsis, in addition to higher
elevation locations in Tibet, the Caucuses, Ural, Alps, and Hengduan mountains adjacent to
currently inhabited regions. Similar currently unoccupied higher elevations are available in many
African mountains though these are usually small mountaintop areas and may be poorly
characterized by CHELSA climate data (Karger et al., 2017). Future distribution models would
benefit from improved environmental data for these high elevation tropical sites.
688
Conclusion
Species distribution models provide potentially powerful windows into past, present, and future
macroecology, but they are rarely confronted with individual performance data. Here we showed
that lower suitability habitats inferred from distribution models had smaller plants with distinct life
history, suggesting a stress escape strategy. While the relationships were noisy it may still be
remarkable that they emerge against the many microsite contributors to individual level variation
in a habitat generalist annual plant. Arabidopsis populations are distributed across diverse
climates, but genetically distinct populations in lower latitudes that are potentially valuable for
research are also highly threatened by anthropogenic climate change in the next few decades.
We believe that combining distribution models with individual data on traits and genotypes
across a species range can be a useful approach to validate distribution models and to dissect
the organismal mechanisms underlying distributions.
701

Acknowledgments
Victoria Meagher and Jaden Hill measured morphology on herbarium specimens. Patrick Herné
assisted in digitizing many herbarium specimens and Eugene Shakirov translated many
Russian specimen labels. Data from the MNHN in Paris were obtained from the participatory
science program 'Les Herbonautes' (MNHN/Tela Botanica) which is part of Infrastructure
Nationale e-RECOLNAT: ANR-11-INBS-0004. This work was supported by NIH award R35
GM138300 to JRL. RM was supported by grant 2019-03758 from Vetenskapsrådet. We also
thank volunteers for digitization and georeferencing through DigiVol, a website developed by the
Australian Museum in collaboration with the Atlas of Living Australia.

711
712

References

Ågren, J., & Schemske, D. W. (2012). Reciprocal transplants demonstrate strong adaptive

differentiation of the model organism Arabidopsis thaliana in its native range. *New*

Phytologist, *194*(4), 1112–1122. https://doi.org/10.1111/j.1469-8137.2012.04112.x

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M., Cao, J.,

Chae, E., Dezwaan, T. M., Ding, W., Ecker, J. R., Exposito-Alonso, M., Farlow, A., Fitz,

J., Gan, X., Grimm, D. G., Hancock, A. M., Henz, S. R., Holm, S., … Zhou, X. (2016).

1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana.

Cell, *166*, 481–491. https://doi.org/10.1016/j.cell.2016.05.063

Angert, A. L., & Schemske, D. W. (2005). The Evolution of Species' Distributions: Reciprocal

Transplants Across the Elevation Ranges of Mimulus Cardinalis and M. Lewisii.

Evolution, *59*(8), 1671–1684. https://doi.org/10.1111/j.0014-3820.2005.tb01817.x

Banta, J. A., Ehrenreich, I. M., Gerard, S., Chou, L., Wilczek, A., Schmitt, J., Kover, P. X., &

Purugganan, M. D. (2012). Climate envelope modelling reveals intraspecific

relationships among flowering phenology, niche breadth and potential range size in

Arabidopsis thaliana. *Ecology Letters*, *15*(8), 769–777. https://doi.org/10.1111/j.1461-

0248.2012.01796.x

Baumgartner, J., Wilson, P., & Esperón-Rodríguez, M. (2017). rmaxent: Tools for working with

Maxent in R. *R Package Version 0.8*, *3*.

Bivand, R. S., Pebesma, E. J., Gómez-Rubio, V., & Pebesma, E. J. (2008). *Applied spatial data*

analysis with R (Vol. 747248717). Springer.

Bontrager, M., & Angert, A. L. (2015). Effects of range-wide variation in climate and isolation on

floral traits and reproductive output of Clarkia pulchella. *American Journal of Botany*,

103, 10–21. https://doi.org/10.3732/ajb.1500091

Bontrager, M., Usui, T., Lee-Yaw, J. A., Anstett, D. N., Branch, H. A., Hargreaves, A. L., Muir, C.

D., & Angert, A. L. (2021). Adaptation across geographic ranges is consistent with strong

selection in marginal climates and legacies of range expansion. *Evolution*, *75*(6), 1316–

1333. https://doi.org/10.1111/evo.14231

Buckley, L. B., Waaser, S. A., MacLean, H. J., & Fox, R. (2011). Does including physiology

improve species distribution model predictions of responses to recent climate change?

Ecology, *92*(12), 2214–2221. https://doi.org/10.1890/11-0066.1

Chala, D., Zimmermann, N. E., Brochmann, C., & Bakkestuen, V. (2017). Migration corridors for

alpine plants among the 'sky islands' of eastern Africa: Do they, or did they exist? *Alpine*

Botany, *127*(2), 133–144. https://doi.org/10.1007/s00035-017-0184-z

Cheng, M.-C., Hsieh, E.-J., Chen, J.-H., Chen, H.-Y., & Lin, T.-P. (2012). Arabidopsis RGLG2,

Functioning as a RING E3 Ligase, Interacts with AtERF53 and Negatively Regulates the

Plant Drought Stress Response. *Plant Physiology*, *158*(1), 363–375.

https://doi.org/10.1104/pp.111.189738

Clark, J. S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest

Trees. *Science*, *327*(5969), 1129–1132.

Clauw, P., Coppens, F., Beuf, K. D., Dhondt, S., Daele, T. V., & Maleux, K. (2015). Leaf

responses to mild drought stress in natural variants of Arabidopsis. *Plant Physiol.*, *167*.

https://doi.org/10.1104/pp.114.254284

Cooper, M. (2021). *moranfast: Calculate Moran's I quickly with low memory footprint for Big*

Data. [C++]. https://github.com/mcooper/moranfast

Csergő, A. M., Salguero-Gómez, R., Broennimann, O., Coutts, S. R., Guisan, A., Angert, A. L.,

Welk, E., Stott, I., Enquist, B. J., McGill, B., Svenning, J.-C., Violle, C., & Buckley, Y. M.

(2017). Less favourable climates constrain demographic strategies in plants. *Ecology*

Letters, *20*(8), 969–980. https://doi.org/10.1111/ele.12794

Daru, B. H., Park, D. S., Primack, R. B., Willis, C. G., Barrington, D. S., Whitfeld, T. J. S.,

764     Seidler, T. G., Sweeney, P. W., Foster, D. R., Ellison, A. M., & Davis, C. C. (2018).

765     Widespread sampling biases in herbaria revealed from large-scale digitization. *New*

766     *Phytologist*, *217*(2), 939–955. https://doi.org/10.1111/nph.14855

767 DeLeo, V. L., Menge, D. N. L., Hanks, E. M., Juenger, T. E., & Lasky, J. R. (2020). Effects of

768     two centuries of global environmental variation on phenology and physiology of

769     Arabidopsis thaliana. *Global Change Biology*, *26*(2), 523–538.

770     https://doi.org/10.1111/gcb.14880

771 Durvasula, A., Fulgione, A., Gutaker, R. M., Alacakaptan, S. I., Flood, P. J., Neto, C.,

772     Tsuchimatsu, T., Burbano, H. A., Picó, F. X., Alonso-Blanco, C., & Hancock, A. M.

773     (2017). African genomes illuminate the early history and transition to selfing in

774     Arabidopsis thaliana. *Proceedings of the National Academy of Sciences*, 201616736.

775     https://doi.org/10.1073/pnas.1616736114

776 Early, R., & Sax, D. F. (2014). Climatic niche shifts between species' native and naturalized

777     ranges raise concern for ecological forecasts during invasions and climate change.

778     *Global Ecology and Biogeography*, *23*(12), 1356–1365.

779     https://doi.org/10.1111/geb.12208

780 Elith, J., Kearney, M., & Phillips, S. (2010). The art of modelling range-shifting species. *Methods*

781     *in Ecology and Evolution*, *1*(4), 330–342. https://doi.org/10.1111/j.2041-

782     210X.2010.00036.x

783 Estarague, A., Vasseur, F., Sartori, K., Bastias, C., Cornet, D., Rouan, L., Beurier, G., Exposito-

784     Alonso, M., Herbette, S., Bresson, J., Vile, D., & Violle, C. (2021). *Into the range: A*

785     *latitudinal gradient or a center-margins differentiation of ecological strategies in*

786     *Arabidopsis thaliana?* (p. 2021.10.15.461205).

787     https://doi.org/10.1101/2021.10.15.461205

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A., & Weigel, D. (2018).

Genomic basis and evolutionary potential for extreme drought adaptation in Arabidopsis

thaliana. *Nature Ecology & Evolution*, *2*(2), 352. https://doi.org/10.1038/s41559-017-

0423-0

Fielding, A. H., & Bell, J. F. (1997). A review of methods for the assessment of prediction errors

in conservation presence/absence models. *Environmental Conservation*, *24*(1), 38–49.

https://doi.org/10.1017/S0376892997000088

Forester, B. R., DeChaine, E. G., & Bunn, A. G. (2013). Integrating ensemble species

distribution modelling and statistical phylogeography to inform projections of climate

change impacts on species distributions. *Diversity and Distributions*, *19*(12), 1480–1495.

https://doi.org/10.1111/ddi.12098

Fournier-Level, A., Perry, E. O., Wang, J. A., Braun, P. T., Migneault, A., Cooper, M. D.,

Metcalf, C. J. E., & Schmitt, J. (2016). Predicting the evolutionary dynamics of seasonal

adaptation to novel climates in Arabidopsis thaliana. *Proceedings of the National*

Academy of Sciences, *113*(20), E2812–E2821.

https://doi.org/10.1073/pnas.1517456113

Fulgione, A., & Hancock, A. M. (2018). Archaic lineages broaden our view on the history of

Arabidopsis thaliana. *New Phytologist*, *219*(4), 1194–1198.

https://doi.org/10.1111/nph.15244

Gamba, D., Lorts, C., Haile, A., Sahay, S., Lopez, L., Xia, T., Kulesza, E., Elango, D., Kerby, J.,

Yifru, M., Bulafu, C. E., Wondimu, T., Glowacka, K., & Lasky, J. (2022). *The genomics*

and physiology of abiotic stressors associated with global elevation gradients in

Arabidopsis thaliana (p. 2022.03.22.485410). bioRxiv.

https://doi.org/10.1101/2022.03.22.485410

Gbif.Org. (2020). *Occurrence Download*. The Global Biodiversity Information Facility.

https://doi.org/10.15468/DL.H95GHR

Gienapp, P., Fior, S., Guillaume, F., Lasky, J. R., Sork, V. L., & Csilléry, K. (2017). Genomic

Quantitative Genetics to Study Evolution in the Wild. *Trends in Ecology & Evolution*,

32(12), 897–908. https://doi.org/10.1016/j.tree.2017.09.004

Greiser, C., Hylander, K., Meineri, E., Luoto, M., & Ehrlén, J. (2020). Climate limitation at the

cold edge: Contrasting perspectives from species distribution modelling and a transplant

experiment. *Ecography*, *43*(5), 637–647. https://doi.org/10.1111/ecog.04490

Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G.,

Toomajian, C., Roux, F., & Bergelson, J. (2011). Adaptation to Climate Across the

Arabidopsis thaliana Genome. *Science*, *334*(6052), 83–86.

Hargreaves, A. L., Samis, K. E., Eckert, C. G., Schmitz, A. E. O. J., & Bronstein, E. J. L. (2014).

Are Species' Range Limits Simply Niche Limits Writ Large? A Review of Transplant

Experiments beyond the Range. *The American Naturalist*, *183*(2), 157–173.

https://doi.org/10.1086/674525

Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of

monthly climatic observations – the CRU TS3.10 Dataset. *International Journal of*

Climatology, *34*(3), 623–642. https://doi.org/10.1002/joc.3711

Heberling, J. M. (2021). Herbaria as Big Data Sources of Plant Traits. *International Journal of*

Plant Sciences. https://doi.org/10.1086/717623

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high

resolution interpolated climate surfaces for global land areas. *International Journal of*

Climatology, *25*(15), 1965–1978.

Hijmans, R. J., & Elith, J. (2013). Species distribution modeling with R. *R CRAN Project*.

Hoffmann, M. H. (2002). Biogeography of Arabidopsis thaliana (L.) Heynh. (Brassicaceae).

Journal of Biogeography, *29*(1), 125–134.

Hoffmann, M. H. (2005). EVOLUTION OF THE REALIZED CLIMATIC NICHE IN THE GENUS:

ARABIDOPSIS (BRASSICACEAE). *Evolution*, *59*(7), 1425-1436.

840 Hsu, C.-W., Lo, C.-Y., & Lee, C.-R. (2019). On the postglacial spread of human commensal

841   Arabidopsis thaliana: Journey to the East. *New Phytologist*, *222*(3), 1447–1457.

842   https://doi.org/10.1111/nph.15682

843 Johnson, T. C., Scholz, C. A., Talbot, M. R., Kelts, K., Ricketts, R. D., Ngobi, G., Beuning, K.,

844   Ssemmanda, I., & McGill, J. W. (1996). Late Pleistocene Desiccation of Lake Victoria

845   and Rapid Evolution of Cichlid Fishes. *Science*, *273*(5278), 1091–1093.

846   https://doi.org/10.1126/science.273.5278.1091

847 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N.

848   E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's

849   land surface areas. *Scientific Data*, *4*(1), 1–20. https://doi.org/10.1038/sdata.2017.122

850 Kawakatsu, T., Huang, S. C., Jupe, F., Sasaki, E., Schmitz, R. J., Urich, M. A., Castanon, R.,

851   Nery, J. R., Barragan, C., He, Y., Chen, H., Dubin, M., Lee, C.-R., Wang, C., Bemm, F.,

852   Becker, C., O'Neil, R., O'Malley, R. C., Quarless, D. X., … Ecker, J. R. (2016).

853   Epigenomic Diversity in a Global Collection of Arabidopsis thaliana Accessions. *Cell*,

854   *166*(2), 492–505. https://doi.org/10.1016/j.cell.2016.06.044

855 Koornneef, M., & Meinke, D. (2010). The development of Arabidopsis as a model plant. *The*

856   *Plant Journal*, *61*(6), 909–921. https://doi.org/10.1111/j.1365-313X.2009.04086.x

857 Korves, T. M., Schmid, K. J., Caicedo, A. L., Mays, C., Stinchcombe, J. R., Purugganan, M. D.,

858   & Schmitt, J. (2007). Fitness Effects Associated with the Major Flowering Time Gene

859   FRIGIDA in Arabidopsis thaliana in the Field. *The American Naturalist*, *169*(5), E141–

860   E157.

861 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., & Sambridge, M. (2014). Sea level and global ice

862   volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National*

863   *Academy of Sciences*, *111*(43), 15296–15303. https://doi.org/10.1073/pnas.1411762111

864 Lasky, J. R., Des Marais, D. L., Lowry, D. B., Povolotskaya, I., McKay, J. K., Richards, J. H.,

865   Keitt, T. H., & Juenger, T. E. (2014). Natural variation in abiotic stress responsive gene

866  expression and local adaptation to climate in Arabidopsis thaliana. *Molecular Biology*

867  *and Evolution*, *31*(9), 2283–2296. https://doi.org/10.1093/molbev/msu170

868 Lasky, J. R., Forester, B. R., & Reimherr, M. (2018). Coherent synthesis of genomic

869  associations with phenotypes and home environments. *Molecular Ecology Resources*,

870  *18*(1), 91–106. https://doi.org/10.1101/051110

871 Lasky, J. R., Hooten, M. B., & Adler, P. B. (2020). What processes must we understand to

872  forecast regional-scale population dynamics? *Proceedings of the Royal Society B:*

873  *Biological Sciences*, *287*(1940), 20202219. https://doi.org/10.1098/rspb.2020.2219

874 Lasky, J. R., Upadhyaya, H. D., Ramu, P., Deshpande, S., Hash, C. T., Bonnette, J., Juenger,

875  T. E., Hyma, K., Acharya, C., Mitchell, S. E., Buckler, E. S., Brenton, Z., Kresovich, S., &

876  Morris, G. P. (2015). Genome-environment associations in sorghum landraces predict

877  adaptive traits. *Science Advances*, *1*(6), e1400218.

878  https://doi.org/10.1126/sciadv.1400218

879 Lee, C.-R., Svardal, H., Farlow, A., Exposito-Alonso, M., Ding, W., Novikova, P., Alonso-Blanco,

880  C., Weigel, D., & Nordborg, M. (2017). On the post-glacial spread of human commensal

881  Arabidopsis thaliana. *Nature Communications*, *8*(1), 14458.

882  https://doi.org/10.1038/ncomms14458

883 Leemans, R., & Cramer, W. P. (1991). The CLIMATE data base of monthly normals for globally

884  gridded climatic variables (Version 2.1) [Data set]. In *Potsdam Institute for Climate*

885  *Impact Research*. PANGAEA. https://doi.org/10.1594/PANGAEA.759491

886 Lee-Yaw, J. A., Kharouba, H. M., Bontrager, M., Mahony, C., Csergő, A. M., Noreen, A. M. E.,

887  Li, Q., Schuster, R., & Angert, A. L. (2016). A synthesis of transplant experiments and

888  ecological niche models suggests that range limits are often niche limits. *Ecology*

889  *Letters*, *19*(6), 710–722. https://doi.org/10.1111/ele.12604

Lee-Yaw, J. A., McCune, J., Pironon, S., & Sheth, S. N. (2021). Species distribution models

rarely predict the biology of real populations. *Ecography*, *n/a*(n/a).

https://doi.org/10.1111/ecog.05877

Li, B., Gao, Z., Liu, X., Sun, D., & Tang, W. (2019). Transcriptional Profiling Reveals a Time-of-

Day-Specific Role of REVEILLE 4/8 in Regulating the First Wave of Heat Shock–

Induced Gene Expression in Arabidopsis. *The Plant Cell*, *31*(10), 2353–2369.

https://doi.org/10.1105/tpc.19.00519

Li, G., Du, S., & Guo, K. (2015). Evaluation of Limiting Climatic Factors and Simulation of a

Climatically Suitable Habitat for Chinese Sea Buckthorn. *PLoS ONE*, *10*(7), e0131659.

https://doi.org/10.1371/journal.pone.0131659

Liu, C., Wolter, C., Xian, W., & Jeschke, J. M. (2020). Species distribution models have limited

spatial transferability for invasive species. *Ecology Letters*, *23*(11), 1682–1692.

https://doi.org/10.1111/ele.13577

Lopez, L., Bellis, E. S., Wafula, E., Hearne, S. J., Honaas, L., Ralph, P. E., Timko, M. P.,

Unachukwu, N., dePamphilis, C. W., & Lasky, J. R. (2019). Transcriptomics of host-

specific interactions in natural populations of the parasitic plant purple witchweed (Striga

hermonthica). *Weed Science*, *67*(4), 397–411. https://doi.org/10.1017/wsc.2019.20

Lovell, J. T., Juenger, T. E., Michaels, S. D., Lasky, J. R., Platt, A., Richards, J. H., Yu, X.,

Easlon, H. M., Sen, S., & McKay, J. K. (2013). Pleiotropy of FRIGIDA enhances the

potential for multivariate adaptation. *Proceedings of the Royal Society B: Biological*

Sciences, *280*(1763), 20131043. https://doi.org/10.1098/rspb.2013.1043

Ludlow, M. M. (1989). Strategies of response to water stress. In K. H. Kreeb, H. Richter, & T. M.

Hinckley (Eds.), *Structural and Functional Responses to Environmental Stresses* (pp.

269–281). SPB Academic.

Mandáková, T., Thorbjörnsson, H., Pisupati, R., Reichardt, I., Lysak, M. A., & Anamthawat-

Jónsson, K. (2017). Icelandic accession of Arabidopsis thaliana confirmed with

cytogenetic markers and its orign inferred from whole-genome sequencing. *Icelandic*

Agricultural Sciences, *30*, 29–38. https://doi.org/10.16886/IAS.2017.03

Martínez-Berdeja, A., Stitzer, M. C., Taylor, M. A., Okada, M., Ezcurra, E., Runcie, D. E., &

Schmitt, J. (2020). Functional variants of DOG1 control seed chilling responses and

variation in seasonal life-history strategies in Arabidopsis thaliana. *Proceedings of the*

National Academy of Sciences, *117*(5), 2526–2534.

https://doi.org/10.1073/pnas.1912451117

Merow, C., Latimer, A. M., Wilson, A. M., McMahon, S. M., Rebelo, A. G., & Silander Jr, J. A.

(2014). On using integral projection models to generate demographically driven

predictions of species' distributions: Development and validation using sparse data.

Ecography, *37*(12), 1167–1183. https://doi.org/10.1111/ecog.00839

Monroe, J. G., McGovern, C., Lasky, J. R., Grogan, K., Beck, J., & McKay, J. K. (2016).

Adaptation to warmer climates by parallel functional evolution of CBF genes in

Arabidopsis thaliana. *Molecular Ecology*, *25*(15), 3632–3644.

https://doi.org/10.1111/mec.13711

Muscarella, R., Galante, P. J., Soley-Guardia, M., Boria, R. A., Kass, J. M., Uriarte, M., &

Anderson, R. P. (2014). ENMeval: An R package for conducting spatially independent

evaluations and estimating optimal model complexity for Maxent ecological niche

models. *Methods in Ecology and Evolution*, *5*(11), 1198–1205.

https://doi.org/10.1111/2041-210X.12261

Peterson, A., Soberón, J., Pearson, R., Anderson, R., Martínez-Meyer, E., Nakamura, M., &

Araújo, M. (2011). Ecological Niches and Geographic Distributions. In *Monographs in*

Population Biology (Vol. 49). https://doi.org/10.1515/9781400840670

Pheloung, P. (1999). Preventing the introduction of potential new weeds to Australia. *Practical*

Weed Management: Protecting Agriculture and the Environment. 10th Biennial Noxious

Weeds Conference, Ballina, Australia, 20th-22nd July 1999., 18–24.

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species

geographic distributions. *Ecological Modelling*, *190*(3), 231–259.

https://doi.org/10.1016/j.ecolmodel.2005.03.026

Prentice, I. C., Sykes, M. T., Lautenschlager, M., Harrison, S. P., Denissenko, O., & Bartlein, P.

J. (1993). Modelling Global Vegetation Patterns and Terrestrial Carbon Storage at the

Last Glacial Maximum. *Global Ecology and Biogeography Letters*, *3*(3), 67–76.

https://doi.org/10.2307/2997548

Price, T., & Schluter, D. (1991). On the Low Heritability of Life-History Traits. *Evolution*, *45*(4),

853–861. https://doi.org/10.2307/2409693

Rudman, S. M., Greenblum, S., Hughes, R. C., Rajpurohit, S., Kiratli, O., Lowder, D. B.,

Lemmon, S. G., Petrov, D. A., Chaston, J. M., & Schmidt, P. (2019). Microbiome

composition shapes rapid genomic adaptation of Drosophila melanogaster. *Proceedings*

of the National Academy of Sciences, *116*(40), 20025–20032.

https://doi.org/10.1073/pnas.1907787116

Samis, K. E., & Eckert, C. G. (2007). Testing the abundant center model using range-wide

demographic surveys of two coastal dune plants. *Ecology*, *88*(7), 1747–1758.

https://doi.org/10.1890/06-1153.1

Scholl, R. L., May, S. T., & Ware, D. H. (2000). Seed and Molecular Resources for Arabidopsis.

Plant Physiology, *124*(4), 1477–1480. https://doi.org/10.1104/pp.124.4.1477

Sinnwell, J. P., Therneau, T. M., & Schaid, D. J. (2014). The kinship2 R Package for Pedigree

Data. *Human Heredity*, *78*(2), 91–93. https://doi.org/10.1159/000363105

Takou, M., Wieters, B., Kopriva, S., Coupland, G., Linstädter, A., & De Meaux, J. (2019). Linking

genes with ecological strategies in Arabidopsis thaliana. *Journal of Experimental Botany*,

70(4), 1141–1151. https://doi.org/10.1093/jxb/ery447

Therneau, T. M., & Therneau, M. T. M. (2015). Package 'coxme.' *R Package Version*, *2*(5).

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C.,

Erasmus, B. F. N., de Siqueira, M. F., Grainger, A., Hannah, L., Hughes, L., Huntley, B.,

van Jaarsveld, A. S., Midgley, G. F., Miles, L., Ortega-Huerta, M. A., Townsend

Peterson, A., Phillips, O. L., & Williams, S. E. (2004). Extinction risk from climate

change. *Nature*, *427*(6970), 145–148. https://doi.org/10.1038/nature02121

Toledo, B., Marcer, A., Méndez-Vigo, B., Alonso-Blanco, C., & Picó, F. X. (2020). An ecological

history of the relict genetic lineage of Arabidopsis thaliana. *Environmental and*

Experimental Botany, *170*, 103800. https://doi.org/10.1016/j.envexpbot.2019.103800

Tozer, B., Sandwell, D. T., Smith, W. H. F., Olson, C., Beale, J. R., & Wessel, P. (2019). Global

Bathymetry and Topography at 15 Arc Sec: SRTM15+. *Earth and Space Science*, *6*(10),

1847–1864. https://doi.org/10.1029/2019EA000658

Turner, K. G., Fréville, H., & Rieseberg, L. H. (2015). Adaptive plasticity and niche expansion in

an invasive thistle. *Ecology and Evolution*, *5*(15), 3183–3197.

https://doi.org/10.1002/ece3.1599

Vasseur, F., Sartori, K., Baron, E., Fort, F., Kazakou, E., Segrestin, J., Garnier, E., Vile, D., &

Violle, C. (2018). Climate as a driver of adaptive variations in ecological strategies in

Arabidopsis thaliana. *Annals of Botany*, *122*(6), 935–945.

https://doi.org/10.1093/aob/mcy165

Warner, R. M., & Erwin, J. E. (2005). Naturally occurring variation in high temperature induced

floral bud abortion across Arabidopsis thaliana accessions. *Plant, Cell & Environment*,

28(10), 1255–1266. https://doi.org/10.1111/j.1365-3040.2005.01361.x

Whitlock, M. C., & Guillaume, F. (2009). Testing for Spatially Divergent Selection: Comparing

QST to FST. *Genetics*, *183*(3), 1055–1063. https://doi.org/10.1534/genetics.108.099812

Wilczek, A. M., Roe, J. L., Knapp, M. C., Cooper, M. D., Lopez-Gallego, C., Martin, L. J., Muir,

C. D., Sim, S., Walker, A., Anderson, J., Egan, J. F., Moyers, B. T., Petipas, R.,

Giakountis, A., Charbit, E., Coupland, G., Welch, S. M., & Schmitt, J. (2009). Effects of

Genetic Perturbation on Seasonal Life History Plasticity. *Science*, *323*(5916), 930–934.

Witsen, D. (2012). *Djibouti Biodiversity and Tropical Forests: 118/119 Assessment* (p. 56).

United States Agency for International Development.

https://usaidgems.org/Documents/FAA&Regs/FAA118119/Djibouti2012.pdf

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. *Journal of the Royal Statistical*

Society: Series B (Statistical Methodology), *73*(1), 3–36. https://doi.org/10.1111/j.1467-

9868.2010.00749.x

Zeng, L., Gu, Z., Xu, M., Zhao, N., Zhu, W., Yonezawa, T., Liu, T., Qiong, L., Tersing, T., Xu, L.,

Zhang, Y., Xu, R., Sun, N., Huang, Y., Lei, J., Zhang, L., Xie, F., Zhang, F., Gu, H., …

Zhong, Y. (2017). Discovery of A high-altitude ecotype and ancient lineage of

Arabidopsis thaliana from Tibet. *Science Bulletin*, *62*(24), 1628–1630.

https://doi.org/10.1016/j.scib.2017.10.007

Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association

studies. *Nat Genet*, *44*(7), 821–824. https://doi.org/10.1038/ng.2310

Zou, Y.-P., Hou, X.-H., Wu, Q., Chen, J.-F., Li, Z.-W., Han, T.-S., Niu, X.-M., Yang, L., Xu, Y.-C.,

Zhang, J., Zhang, F.-M., Tan, D., Tian, Z., Gu, H., & Guo, Y.-L. (2017). Adaptation of

Arabidopsis thaliana to the Yangtze River basin. *Genome Biology*, *18*(1), 239.

https://doi.org/10.1186/s13059-017-1378-9

1012