Skeletal Muscle TFEB Signaling Promotes Central Nervous System Function and

Reduces Neuroinflammation during Aging and Neurodegenerative Disease

Ian Matthews [1,2]

Allison Birnbaum [1,2]

Anastasia Gromova [3]

Amy W. Huang [1,2]

Kailin Liu [1,2]

Eleanor A. Liu [1,2]

Kristen Coutinho [5]

Megan McGraw [6]

Dalton C. Patterson [6]

Macy T. Banks [6]

Amber C. Nobles [7]

Nhat Nguyen [3]

Gennifer E. Merrihew [8]

Lu Wang [9]

Eric Baeuerle [10-12]

Elizabeth Fernandez [10-12]

Nicolas Musi [13]

Michael J. MacCoss [8]

Helen C. Miranda [14-16]

* Albert R. La Spada [3,4] (co-corresponding author)*

Constanza J. Cortes [1,2] (corresponding author)

[1] Leonard Davis School of Gerontology, [2] University of Southern California, [3] Department of
Pathology and Laboratory Medicine, University of California, Irvine, [4] Department of Neurology
and Department of Biological Chemistry, UCI Institute for Neurotherapeutics, University of
California, Irvine, [5] Comprehensive Diabetes Center, University of Alabama at Birmingham, [6]
Department of Cell, Developmental and Integrative Biology, University of Alabama at
Birmingham, [7] Department of Neurology, University of Alabama at Birmingham, [8] Department of
Genome Sciences, University of Washington , [9] Department of Environmental and Occupational
Health Sciences, University of Washington, [10] Department of Pharmacology, [11] Barshop Institute
for Longevity and Aging Studies, University of Texas Health San Antonio; [12] Geriatric Research,
Education and Clinical Center, South Texas Veterans Health Care Network, San Antonio TX
USA [13] Department of Medicine, Cedars-Sinai Medical Center, [14] Department of Genetics and
Genome Sciences, School of Medicine, Case Western University, [15] Department of
Neurosciences, School of Medicine, Case Western Reserve University, [16] RNA Center, School
of Medicine, Case Western Reserve University,

Funding: NIH R01 AG077536 (CJC), University of Washington Nathan Shock Center P30
AG013280 (Pilot Grant to CJC), San Antonio Nathan Shock Center of Excellence in the Basic
Biology of Aging P30AG013319 (Pilot Grant to CJC), NIH R35 122140 (ALS).

Acknowledgements: We thank members of the lab past and present, as well as former and
current colleagues and collaborators for their helpful contributions.

Summary

Skeletal muscle has recently arisen as a novel regulator of Central Nervous System (CNS) function and aging, secreting bioactive molecules known as myokines with metabolism-modifying functions in targeted tissues, including the CNS. Here we report the generation of a novel transgenic mouse with enhanced skeletal muscle lysosomal and mitochondrial function via targeted overexpression of Transcription Factor E-B (TFEB). We have discovered that the resulting geroprotective effects in skeletal muscle reduce neuroinflammation and the accumulation of tau-associated pathological hallmarks in a mouse model of tau pathology. Muscle-specific TFEB overexpression also significantly ameliorates proteotoxicity, reduces neuroinflammation and promotes transcriptional remodeling of the aged CNS, preserving cognition and memory in aged mice. Our results implicate the maintenance of skeletal muscle function throughout aging to direct regulation of CNS health and disease, and suggest that skeletal-muscle originating factors may act as novel therapeutic targets against age-associated neurodegenerative disorders.

INTRODUCTION

Aging is associated with an organism-wide progressive loss of tissue form and function, broadly characterized by the 'hallmarks of aging' [1]. In particular, the aging central nervous system (CNS) [1,2] exhibits a global loss in protein homeostasis (proteostasis), impaired neuroplasticity and resilience, and an increase in neuroinflammation [2,3]. These alterations are believed to render the aging CNS vulnerable to age-associated dysfunction and the development of neurodegenerative disease [2].

Over the last ten years, there has been growing evidence that suggests prominent contributions of the periphery to the etiology of age-associated neurodegenerative diseases [4-8]. In particular, manipulation of skeletal muscle protein quality control pathways yields important benefits to the invertebrate CNS, protecting against the accumulation of aggregation-prone neurodegenerative disease proteins in the brain and the retina [5,8,9]. Although the mechanisms responsible for these benefits remain poorly understood, some of these effects are mediated by secreted factors that communicate metabolic and inflammatory signals between tissues [5,7,9]. In agreement with this, blood circulating factors have recently arisen as potent regulators of mammalian CNS aging and metabolism [10-13]. For example, exposure to a young plasma environment (either through heterochronic parabiosis or plasma transfers) can rescue function in the aging CNS [12-14] by decreasing neuroinflammation [15] and enhancing neurogenesis [10]. Studies have further demonstrated that increasing the levels of these individual factors in peripheral circulation is sufficient to rejuvenate the aged CNS [12,16,17], supporting the existence of geroprotective circulating factors with CNS targeting effects. Although the source and identity of these neuroprotective circulating cytokines are unclear, several of them are expressed in, and are known to be secreted from, skeletal muscle [5,18-20]. In fact, skeletal muscle acts as an endocrine organ, secreting a myriad of bioactive factors that induce metabolic changes in a non-cell autonomous manner in distant tissues like liver [21], adipose tissue [22], and the CNS [19].

Recently, this muscle-to-brain signaling axis has been proposed to have important implications for CNS aging and age-associated neurodegenerative disease [19,23,24]. Indeed, skeletal muscle function has arisen as a key predictor for phenotypic and clinical outcomes in age-associated neurodegenerative diseases, including Alzheimer's Disease (AD) and Parkinson's Disease (PD) [25,26]. Consistent with this, we were the first to uncover a novel mechanistic basis for motor neuron disease in the polyglutamine disease Spinal and Bulbar Muscular Atrophy (SBMA), where disruption of skeletal muscle autophagy initiates the pathogenic cascade that culminates

in neuronal toxicity and death [27]. Furthermore, we demonstrated that rescuing proteotoxicity

specifically in skeletal muscle was sufficient to rescue motor neuron degeneration and extend

lifespan in this model, despite robust misfolded protein burden in the brain [28]. We also

demonstrated the feasibility and efficacy of peripheral delivery of antisense oligonucleotides

reducing skeletal muscle expression of the misfolded protein in rescuing neurological

phenotypes [29], confirming that therapeutic targeting of skeletal muscle can have important

benefits to the CNS.

Skeletal muscle metabolism is regulated in part by Transcription Factor E-B (TFEB), a master

regulator of a lysosomal-to-nucleus signaling axis that integrates cellular metabolism and

lysosomal function [30-32]. TFEB expression and function are strongly induced in skeletal muscle

in response to interventions with documented neuroprotective effects on the aging [33,34] and

neurodegenerative disease-afflicted CNS, including low nutrient conditions [35] and exercise [36].

Indeed, TFEB controls metabolic flexibility in muscle during exercise, inducing the expression of

genes involved in mitochondrial biogenesis, fatty acid oxidation, and oxidative phosphorylation

[36]. This coordinated action optimizes mitochondrial substrate utilization, enhancing ATP

production and exercise capacity [36]. These findings identify TFEB as a critical mediator of the

beneficial effects of exercise on metabolism. However, the precise role of TFEB-dependent

signaling in skeletal muscle during aging, and its impact on the muscle-to-brain axis, remains

largely unknown.

Here we report the generation of a novel transgenic mouse with enhanced muscle metabolism

via overexpression of TFEB. We have discovered that the resulting enhanced skeletal muscle

lysosomal function is maintained throughout the lifespan, protecting against the onset of age-

associated mitochondrial dysfunction in aging skeletal muscle. Furthermore, we report that

overexpression of TFEB in skeletal muscle significantly reduces hippocampal accumulation of

neuropathological hallmarks and reduces neuroinflammation in a mouse model of tau

pathology, despite no exogenous activation of the transgene in the CNS. Muscle TFEB

overexpression also ameliorates proteotoxicity, reduces neuroinflammation, and promotes

transcriptional remodeling of the aged CNS without neurodegenerative disease burden,

preserving cognitive performance in healthy aging mice. Our results implicate, for the first time

in mammals, maintenance of skeletal muscle function to direct regulation of CNS health, and

suggest that skeletal-muscle originating factors may act as novel therapeutic targets against

age-associated neurodegenerative diseases.

RESULTS

TFEB overexpression maintains fiber type composition, function, and metabolism-associated networks in aging skeletal muscle

We derived a novel line of conditional transgenic mice carrying the β-actin promoter in combination with the CMV enhancer (CAGGS) and with a floxed 3x-FLAG-eGFP STOP cassette placed just 5' to a 3x-FLAG-human *TFEB* transgene (i.e. fxSTOP-TFEB transgenic mice) (**Figure 1A**). In the presence of Cre-recombinase, the STOP codon cassette will be excised out, allowing expression of the human TFEB transgene, also known as a Cre-loxP ON system. To drive overexpression of TFEB specifically in skeletal muscle, we crossed male fxSTOP-TFEB mice with female Human Skeletal Actin (HSA)-Cre mice to achieve widespread expression of Cre-recombinase in cells of the myogenic lineage [37].

 The resulting cTFEB;HSACre bigenic mice of both sexes exhibited a strong induction of human *TFEB mRNA* expression in skeletal muscle tissue (**Supplemental Figure 1A, quadriceps**). This exogenous *TFEB* expression was still prominent at 24 months of age in cTFEB;HSACre transgenic skeletal muscle (**Supplemental Figure 1B, quadriceps)** suggesting that overexpression of the transgene remains active even at very old ages. Immunoblotting of quadricep skeletal muscle confirmed robust expression of 3x-FLAG-TFEB protein in cTFEB;HSACre bigenic muscle lysates of both sexes (**Figure 1B**). Human TFEB protein levels in male cTFEB;HSACre mice amounted to about 2-fold higher levels than their female littermates at both ages examined. Importantly, and in agreement with our qRT-PCR studies, human TFEB was still robustly expressed at 24 months of age in both sexes, suggesting life-long expression of our transgene upon embryonic recombination (**Figure 1B**). As TFEB is a transcription factor, we confirmed nuclear localization of human TFEB in cTFEB;HSA-Cre skeletal muscle myonuclei via immunofluorescence (**Supplemental Figure 1C, in white**).

Next, we confirmed TFEB protein overexpression exclusively in skeletal muscle of cTFEB;HSACre bigenic mice by multiple approaches. First, we confirmed expression of the biologically inert 3x-FLAG-eGFP cassette upstream of the STOP codon (**Figure 1A**) and no exogenous expression of 3x-FLAG-TFEB in other highly metabolic tissues of non-myogenic lineages in cTFEB;HSACre transgenic mice, including brown adipose tissue (**Supplemental Figure 1D**) and liver (**Supplemental Figure 1E**). We detected a relatively mild expression of human *TFEB* mRNA levels in the heart of cTFEB;HSACre transgenic mice, confirming previous

reports indicating low-level cross-over into cardiac tissue in the HSA-Cre model. However, this

expression amounted to about ~400-fold less in young animals and ~15-fold less in aged

animals than that detected in skeletal muscle at the same ages (**Supplemental Figure 1A,**

ventricles). Importantly, we could not detect 3x-FLAG-TFEB protein via immunoblotting in

cTFEB;HSACre heart ventricle lysates (**Supplemental Figure 1F**). This suggests that despite

some HSA-Cre 'leakiness' into cardiac muscle, synthesis of exogenous TFEB protein in heart

tissue is below detectable levels, confirming the high skeletal muscle tropism of HSA-Cre driven

expression. Finally, we also did not detect expression of exogenous human TFEB mRNA in the

hippocampus (**Supplemental Figure 1A & 1B, hippocampus**) or hemibrain protein lysates

(**Supplemental Figure 1G**). To further confirm no CNS expression of transgenes associated

with the HSA-Cre driver line, we directly assessed the Cre-mediated recombination of an

additional transgene reporter in TdTomato Ai9;HSACre mice. We confirmed robust tdTomato

fluorescence in skeletal muscle (**Supplemental Figure 1H**) but no detectable TdTomato

fluorescence in the CNS (**Supplemental Figure 1I**), including the hippocampus (**Supplemental**

Figure 1I, insets), of TdTomato Ai9;HSACre mice, consistent with the muscle-restricted

specificity of this HSA-Cre-loxP-ON approach [38]. Overall, these data confirm previous reports of

striated muscle-specific Cre-mediated recombination using the HSA-Cre transgenic driver line

[37], and demonstrate detectable expression of human TFEB protein only in skeletal muscle of

cTFEB;HSACre bigenic mice. This establishes our newly developed inducible fxSTOP-TFEB

mouse as an effective model to generate spatiotemporally controlled transgenic expression of

human TFEB.

Analysis of skeletal muscle wet weight, a proxy for skeletal muscle size, revealed trends

towards larger muscle wet weights in young cTFEB;HSACre animals, and a robust and

significant preservation of quadriceps and gastrocnemius skeletal muscle wet weight in aged

(22–24 month-old) transgenic mice of both sexes (**Figure 1C-D**). Indeed, while control animals

lost around 20-30% of their quadriceps wet weight during aging, there was no significant

differences between young or aged cTFEB;HSACre skeletal muscle weight (**Figure 1C-D**).

Interestingly, this maintenance of muscle wet weight was not associated with changes in total

body weight at either age examined (**Figure 1E**). Consistent with these findings, we determined

that overexpression of TFEB significantly decreases the number of small fibers (<1500 μm^2) and

increases the number of large fibers (>1500 μm^2) in young gastrocnemius skeletal muscle of

both sexes (**Figure 1F-G**). This was accompanied by an increase in abundance of type 1 (~1%

more) and type 2A (~9% more) fibers (**Figure 1H-K**) and an increase in the cross-sectional area

(CSA) of Type 2B (~200 um^2 larger) and 2X (~300 um^2 larger) fibers with no changes in Feret's

minimum diameter (**Supplemental Figure 2A-H)**, which was prevalent in male mice and

trending in female mice. Importantly, while control male gastrocnemius muscle lost a significant

portion of their large fibers during aging, they were not altered in aged male cTFEB:HSACre

transgenic muscle (**Figure 1F-G**). The abundance of Type 1 and Type 2A fibers was also still

prevalent in aged male cTFEB:HSACre transgenic muscle (**Figure 1H-K and Supplemental**

Figure 2G-H), suggesting that TFEB overexpression remodels both fast-twitch glycolytic and

oxidative fiber fibers and that these changes are maintained during aging**.** To determine the

physiological effects of this fiber type switch on skeletal muscle function, we examined exercise

performance in all of our cohorts. We determined that at 6 months of age, muscle-TFEB

overexpressing mice had a 20% (males) and 40% (females) longer maximum running distance

to exhaustion (**Figure 1L**) and engaged in more total work (**Figure 1M**) than their control

littermates. Importantly, this increased enhanced physical performance was still present in aged

female cTFEB;HSACre transgenic mice (**Figure 1M**). Taken together, these results suggest that

lifelong-expression of TFEB prevents fiber type atrophy and alters fiber type switching in aging

skeletal muscle, ultimately preserving muscle size and running performance during aging.

Given the principal role of TFEB-regulated signaling (i.e. lysosomal and mitochondrial function)

to skeletal muscle health [39,40], as well as the profound geroprotective effects of TFEB

overexpression on muscle fiber types, muscle size and exercise performance in aging muscle

(**Figure 1**), we sought to examine the overall changes in the skeletal muscle protein landscape

in response to TFEB overexpression throughout aging. We performed proteome profiling of

young and aged, control and cTFEB;HSACre male quadricep skeletal muscle. Differential

abundance analysis of species revealed 909 proteins associated with TFEB overexpression in

young (6-month-old) skeletal muscle (**Figure 2A**). Of these, 857 (~94%) were significantly

overrepresented and 52 (~6%) were underrepresented compared to their littermate controls. At

18 months of age, a time point where age-associated dysfunction is prevalent in murine skeletal

muscle [41], the same proteomics approach identified 351 differentially expressed proteins in

TFEB-expressing skeletal muscle (**Figure 2B**). Of these, 283 proteins (~81%) were significantly

enriched and 68 proteins (~19%) were significantly decreased in cTFEB;HSACre bigenic

muscle, with only 53 of these proteins being uniquely expressed in older mice. Importantly, at

both ages examined, TFEB remained significantly overexpressed (**Figure 2A & 2B, boxes**),

confirming our previous finding that exogenous skeletal muscle TFEB expression is maintained

across the lifespan of our cTFEB;HSACre model (**Figure 1B**).

To visualize the key networks of differentially expressed proteins associated with TFEB-

overexpression, we performed a functional enrichment analysis on all overexpressed proteins

using the STRING protein-protein interaction network [42] (**Figure 2C**). Using TFEB as our 'bait'

for functional connectivity nodes, we identified two clear interaction networks: the first one was

associated with lysosomal biogenesis and function (**Figure 2C, blue**), and included Lamp1 (an

essential lysosomal structural protein), Grn (progranulin) and Psap (prosaposin) as well as the

lysosomal proteases: Ctsb (Cathepsin B), Ctsd (Cathepsin D), and Ctsl (Cathepsin L). The

second interaction network is associated with key skeletal muscle proteostasis and metabolic

signaling pathways including mTOR, Akt and GSK3β (**Figure 2C, green**). Both of these nodes

are key elements of the signaling network controlled by TFEB-mediated transcription [31,32], were

previously reported to be transcriptionally elevated in skeletal muscle after viral-mediated TFEB

overexpression [36], and are central regulators of aging [1].

To further understand the functions of all the upregulated proteins in TFEB-overexpressing

skeletal muscle, KEGG enrichment analysis was performed. As expected, central proteostasis

categories were significantly upregulated in young TFEB-overexpressing skeletal muscle:

including Hypoxia-inducible factor 1 (HIF-1) signaling, proteasome, and autophagy/lysosomal

function categories (**Figure 2D, orange**). Importantly, this enrichment in proteostasis categories

and TFEB-regulated nodes of signaling was also present in our aging cohorts (**Figure 2D, top**).

We confirmed functional activation of TFEB-dependent transcription in cTFEB:HSA-Cre bigenic

muscle, using our previously reported TFEB-responsive muscle gene targets [27] including

Lamp1, *Atg5* (a key regulator of autophagy initiation), *Ctsd* and *Ctsb* (**Figure 2E**).

Immunoblotting studies confirmed significant increases in Lamp1 and Ctsd protein in

cTFEB;HSACre bigenic muscle (**Figure 2F**). Consistent with previous studies demonstrating

TFEB overexpression increases the number and size of LAMP1-positive lysosomes in multiple

tissues [30,35,43], we also detected significant increases in the number of LAMP1+ puncta and total

LAMP1+ area in cross-sections of young female (and similar trends in male) cTFEB;HSACre

skeletal muscle (**Figure 2G**). There were no significant changes to LC3II/LCI ratios or total

polyubiquitinated protein levels in cTFEB;HSACre transgenic mice (**Supplemental Figure 2I**).

These findings confirm physiologically significant increases in TFEB-dependent transcription [36]

and lysosomal biogenesis [32,36] in skeletal muscle of cTFEB:HSA-Cre mice.

Additional functional categories enriched in our proteomics analysis related to cellular

metabolism and mitochondrial function, including thermogenesis, oxidative phosphorylation and

pyruvate metabolism (**Figure 2D, gold**). We also detected central fatty acid metabolism

pathways, including PPAR signaling and metabolism, and degradation and biosynthesis of fatty
acids (**Figure 2D, green**). This is consistent with previously published reports suggesting that
acute TFEB expression in young skeletal muscle induces the expression of genes involved in
mitochondrial biogenesis and oxidative phosphorylation [36]. Here we confirm for the first time that
these changes in cTFEB;HSACre muscle are maintained throughout aging, particularly those
associated with mitochondrial metabolism and fatty acid metabolism (**Figure 2D, top, gold and**
green).

We assessed whether these proteomic profiles associated with mitochondrial function translated
into mitochondrial function differences using high-resolution respirometry with simultaneous
measurements ROS production assays [44]. As reported previously for soleus skeletal muscle [44],
oxygen consumption rates decrease with aging across all mitochondrial complexes in semi-
permeabilized soleus muscle of control mice of both sexes (6 vs. 18 months of age, gold icons)
(**Figure 2H-2I**). Whereas there was no difference in mitochondrial respiration rates between
control and TFEB-overexpressing skeletal muscle at young ages (closed icons), respiration
rates of aged cTFEB;HSACre transgenic muscle (open icons) remained comparable to their
young counterparts across all mitochondrial complexes examined (**Figure2 H-2I**), particularly in
female muscle. Together with our functional data showing increases in exercise capacity in
cTFEB;HSACre mice (**Figure 1L-M**), these results suggest that muscle-TFEB overexpression
promotes maintenance of mitochondrial function and bioenergetic reserves, preventing the age-
associated decline of mitochondrial respiration commonly observed in aging skeletal muscle [1,44].

Finally, proteomics KEGG analysis also showed enrichment in cTFEB;HSACre bigenic muscle
for classical signaling pathways shown to decline during aging, including the Insulin-IGF1-
signaling (IIS) and AMPK signaling [1] (**Figure 2D, blue**). We also detected multiple hits for
amino acid metabolism pathways, such as catabolism of branched-chain amino acids (BCAAs),
including valine, leucine and isoleucine metabolism (**Figure 2D, purple**). Importantly, these
changes were still evident in our aged cohorts (**Figure 2D, top**), consistent with our functional
studies suggesting that TFEB overexpression throughout the lifespan has multiple benefits in
skeletal muscle, targeting established hallmarks of aging [1].

The skeletal muscle 'secretome' (the totality of released organic and inorganic molecules
released from muscle resident cells) is highly dynamic, responding to physiological and
pathophysiological stimuli [45,46] and potentially also changing with age [46]. To directly examine any
changes to the skeletal muscle protein-based secretome associated with TFEB-overexpression,

we pursued additional *in silico* analysis through the Vertebrate Secretome Database (VerSeDa)

on our proteomics cohorts to identify unique age- and genotype-associated signatures of

potentially secreted proteins [47] differentially expressed in cTFEB;HSACre skeletal muscle. We

identified multiple proteins with predicted secreted profiles to be enriched in young and aged

TFEB-expressing skeletal muscle (**Figure 3A**). Some secreted proteins exhibited upregulation

(e.g., nucleobindin-1, angiopoietin-related protein 1) or down-regulation (collagen alpha-1 (I)

chain), only in young cTFEB;HSACre muscle (**Figure 3A, green**). Others remained up-

regulated at both ages in cTFEB;HSACre muscle, (prosaposin, mammalian ependymin-related

protein 1 (MERP1) and dehydrogenase/ reductase SDR family member 7C, D/R SDR7C)

(**Figure 3A, overlap**), or were only differentially expressed in aged muscle (biglycan-1, PG-S1,

Figure 3A, blue), highlighting the dynamic nature of the skeletal muscle secretome throughout

aging. Cathepsin B (Ctsb), a previously documented muscle-originating secreted factor with

known CNS-targeting effects [20] and a known target of TFEB-dependent transcription (**Figure

2E**) [32] was our top up-regulated hit at both ages examined. We validated significant increases in

pro-enzyme levels of Ctsb in both sexes and at both ages examined (**Figure 3B**), as well as

significant increases in mature (secreted) Ctsb isoforms in aged cTFEB;HSACre transgenic

muscle (**Figure 3B**). We also detected trends towards increased levels of circulating CTSB in

the serum of female cTFEB;HSACre mice (**Figure 3E**). Given the potent remodeling of the

lysosomal network associated with activation of TFEB-mediated transcription (**Figure 2**) [30-32],

our data suggests that overexpression of TFEB in skeletal muscle drives increased expression

and secretion of mature Ctsb, a CNS targeting-factor, into circulation.

Enhanced skeletal muscle TFEB-signaling reduces accumulation of

hyperphosphorylated tau and microglial activation in a mouse model of tau pathology.

To assess any potential neuroprotective effects of enhancing skeletal muscle TFEB-signaling in

the context of age-associated neurodegenerative disease pathologies, we derived

cTFEB;HSACre transgenic mice on the MAPT P301S (PS19) background. This is a well-known

model of neurofibrillary tangle toxicity, a hallmark of AD and related tauopathies [48], and is

characterized by prominent accumulation of hippocampal hyperphosphorylated human mutant

MAPT protein and pronounced neuroinflammation, including microgliosis and astrocyte

reactivity [48].

We first confirmed muscle-restricted 3x-FLAG-TFEB overexpression in skeletal muscle lysates

from MAPT P301S;cTFEB;HSACre mice, as well as the presence of human Tau in hippocampal

lysates of both groups via immunoblotting (**Figure 4A**). As previously reported [48], at 9 months of age there is robust accumulation of Ser202/Thr205 hyperphosphorylated tau (residues abnormally phosphorylated in paired helical filaments during AD progression) in the dentate gyrus of the hippocampus of MAPT P301S mice of both sexes [48] (**Figure 4B, middle panel**). Conversely, we noted a significant reduction in the total fluorescence counts of AT8-positive phosphorylated tau in the dentate gyrus of MAPT P301S;cTFEB;HSACre mice (**Figure 4B**, **purple**) compared with MAPT P301S littermate controls (**Figure 4B**, **teal**). More so, this reduction was particularly evident in the levels of 'intracellular tau' accumulating around hippocampal cell bodies (**Figure 4B, insets**), suggesting reduced accumulation of toxic tau species and aggregates [49] in the hippocampus of MAPT P301S;cTFEB;HSACre mice. We confirmed similar but less striking reductions in Thr231 phospho-tau (another primary site of tau phosphorylation associated with neurotoxic tau accumulation) immunostaining (**Supplemental Figure 3A**), suggesting overall reductions in tau hyperphosphorylation in the hippocampus of MAPT P301S transgenic mice with TFEB overexpression in skeletal muscle.

Astrocytes and microglia are key mediators of neuroinflammation, and undergo key morphological and functional alterations during neurodegenerative disease, including changes in ramification and cellular process complexity [50], reduced immune surveillance activity [51] and increased reactivity phenotypes [52]. We confirmed significant increases in the staining for GFAP-positive structures (a marker of astrocyte reactivity) in the dentate gyrus of 9-month-old MAPT P301S mice of both sexes (**Figure 4C, teal**) compared to their MAPT-negative littermate controls (**Figure 4C, gold**). We also detected significant increases in the volume of IBA1+-positive microglia in the same brain region (**Figure 4C, teal**), consistent with previous reports in the MAPT P301S model indicating high levels of neuroinflammation starting as early as 4 months of age [48,53]. Strikingly, we noted significant reductions in both of these pro-inflammatory glial morphometric parameters in the hippocampus of MAPT P301S;cTFEB;HSACre littermates (**Figure 4C, purple**), suggesting overall reduced hippocampal neuroinflammation during symptomatic disease stages through activation of skeletal muscle TFEB-expression.

Enhancing TFEB-signaling in skeletal muscle promotes neurotrophic signaling and reduces hippocampal neuroinflammation in MAPT P301S transgenic mice.

To gain more precise insights into disease-relevant transcriptional changes in the hippocampus of MAPT P301S mice with enhanced muscle-TFEB expression, we used the Nanostring nCounter® Mouse Alzheimer's Disease panel. This platform directly assesses the expression of

770 targeted genes curated from human and pre-clinical models of AD and robustly captures disease-relevant signatures and their modifications after pre-clinical interventions [54]. Using this approach, we found 24 individual differentially expressed genes in MAPT P301S;cTFEB;HSACre hippocampi compared to their littermate MAPT P301S+ controls between both sexes (**Supplemental Figure 3B-C**). Interestingly, most differentially regulated genes were downregulated (18 of 24), consistent with their identity as transcriptional drivers of disease [54]. Many of these transcripts are associated with microglial activation (*Rhoc, Clic1*) markers of neuronal function (*Gdap1l1, Amot, Map3k9),* phospholipid and cholesterol remodeling (*Ano6, Cyp27a1*), growth factor signaling (*Vgf*), endothelial cell migration and matrix adhesion (*Rras, Cdc42Ep1*) and guanine nucleotide exchange factors (*Dock3*). Multiple down-regulated genes were associated with novel variants and loci associated with increased polygenic neurodegenerative disease risk (*Prkd3, Arhgap31)* as well as predictors of the rate of cognitive decline in AD (*Hecw1* [55]). *Pbxip1,* another significantly downregulated marker, has been found to be associated with phosphorylated-tau and $A\beta_{1-40}$ levels in the human temporal cortex[56]. Additional transcriptional changes of significance include genes of unknown function but are classified as transcriptional identifiers of AD progression *(Ptprn, Cpne2,* and *Eri2)[54]*. Up-regulated transcripts included gene repressors (*Zbtb33*), modulators of neuroinflammation (*Pecam1*) and neurodegeneration (*Scna*), as well as a carbohydrate transmembrane transporter (*Tmem144*) and a GTP-ase nucleotide exchange factor (*Rcc1*).

Annotation of the differentially co-expressed/co-functional clusters included in this panel revealed significant reductions in the nCounter AD microglial activation module (composed of 63 genes associated with microglial activation during AD progression) (**Figure 4D**) in the hippocampi of 6 month old female (but not male) MAPT P301S;cTFEB;HSACre mice compared with MAPT P301S littermates. This is consistent with reduced neuroinflammation in the hippocampus of MAPT P301S mice with enhanced muscle-TFEB expression, as suggested by our glial morphometric analysis (**Figure 4C**). Interestingly, we also detected a sex-bias in the trophic factor functional cluster (composed of genes *Arhgdib, Bcl2, Calm3, Camk4, Gab1, Mapk9, Mapkapk2, Ntrk2, Pik3r1* and *Psen1*), with a significant improvement back to baseline expression in the hippocampus of male (but not female) MAPT P301S;cTFEB;HSACre mice (**Figure 4E**). We confirmed activation of neurotrophic signaling as early as 6 months of age in the hippocampus of male MAPT P301S;cTFEB;HSACre mice by targeted qRT-PCR analysis, which showed corrections of declining expression levels of neurotrophic factors *Bdnf* and *Fndc5* back to control littermate levels (**Figure 4F-G**) [23,57]. Importantly, there was also a significant

extension (38%) in the lifespan of male MAPT P301S;cTFEB;HSACre mice (**Figure 4H**). While

the median survival of male MAPT P301S mice was 352 days (**Figure 4H, teal curve**), male

MAPT P301S;cTFEB;HSACre mice median survival was 489 days (**Figure 4H, purple curve).**

Although we did not detect a similar lifespan extension in female MAPT P301S;cTFEB;HSACre

mice (**Figure 4I**), it is worth noting that in our hands, female MAPT P301S transgenic mice

already live significantly longer than their male littermate controls (median survival of 468.5

days), perhaps suggesting they have reached maximal lifespan in this model. Finally, we did not

detect any differences in ataxia progression [58] (**Supplemental Figure 3D**) or weight loss

(**Supplemental Figure 3E**) between our groups. Altogether, these provocative results suggest

that enhanced skeletal muscle-TFEB expression modifies the accumulation of pathogenic tau

isoforms, reduces neuroinflammation and re-activates neurotrophic signaling in the

hippocampus of MAPT P301S mice in a sex-dependent manner. Importantly, this occurs in the

absence of detectable exogenous human TFEB expression in the CNS (**Supplemental Figure**

1), suggesting an independent mechanism of muscle-to-brain communication underlying these

effects.

Skeletal muscle over-expression of TFEB decreases neuroinflammation markers and

lipofuscin accumulation in the healthy aged CNS.

To examine the effect of increased muscle-TFEB expression on the healthy aging brain without

neurodegenerative disease burden, we assessed the transcriptional and functional status of

known inflammatory markers in the CNS of healthy aged (20+ months old) control and

cTFEB;HSACre mice, an age when global decreases in proteostasis and increases in pro-

inflammatory signaling are detectable in most murine tissues [1,59], including the CNS [1,2]. Total

hippocampal mRNA levels of *Ccl2* and *Nfκb*, pro-inflammatory cytokines previously reported to

mediate microglial responses to inflammation, were significantly reduced in the hippocampus of

aging female (but not male) cTFEB;HSACre bigenic mice (**Figure 5A-B**). Interestingly, the

expression of *Il6* (interleukin 6) was significantly elevated in the same groups (**Figure 5C**). We

also detected significantly higher levels of expression of *Bdnf* (Brain Derived Neurotrophic

Factor) in hippocampal lysates from aged male (but not female) mice (**Figure 5D**), similar to the

upregulation of hippocampal BDNF we observed in our male MAPT P301;cTFEB;HSACre

model (**Figure 4F**).

Using similar morphometric analysis as before to quantify glial cell shape and morphology, we

did not detect any differences in microglia (IBA1+) or astrocyte (GFAP+) number, volume or

ramification state in the dentate gyrus of aged cTFEB;HSACre bigenic mice compared to their littermate controls (**Figure 5E-F**). This may suggest that the expression differences observed via transcriptional approaches above may either originate in non-glial cells (neurons or endothelial cells, for example) or not translate into detectable morphological changes at the ages examined. It is important to note that age-associated changes in neuroinflammation in an otherwise healthy brain are not as pronounced as those seen in the context of neurodegenerative disease (such as in the MAPT P301S model, for example). Overall, these results suggest a shift in the cytokine transcriptional landscape of the aging female hippocampus with upregulated skeletal muscle TFEB-expression towards reduced inflammatory signaling, consistent with what we observed in our MAPT P301S studies (**Figure 4**).

Accumulation of lipofuscin, a non-degradable intracellular auto-fluorescent polymer, becomes prominent in the aging brain, likely reflecting an age-associated decline in basal CNS autophagy [1]. We examined the clearance capabilities of the aging CNS by indirect immunofluorescence of lipofuscin granules in brain sections of aged (21-24 months old) control and cTFEB;HSACre mice. While we detected striking lipofuscin granule accumulation in several brain regions of aged control mice, including the hippocampus (**Figure 5G, top**), there was a significant decrease in lipofuscin deposition in the dentate gyrus of aged cTFEB;HSACre females (**Figure 5G, bottom),** with a similar non-significant trend in males. To determine any changes in proteostasis associated signaling in this brain region, we next examined the transcriptional levels of multiple autophagy and lysosomal associated genes. Consistent with previous reports, we confirmed significant age-associated increases in the expression of autophagy lysosomal markers *Atg8/Lc3*, *Lamp1, Sqstm1,* and *Ctsd* in the male hippocampus, which has been proposed to occur as a compensation for age-associated decline in proteostasis function (**Figure 5H-K**). Interestingly, these differences were not as prominent in the aging control female hippocampus. Furthermore, some of these transcriptional increases were reduced in the hippocampus of aged male cTFEB;HSACre bigenic mice, suggesting that enhancing skeletal muscle TFEB-signaling promotes maintenance of hippocampal proteostasis during aging. Our data thus indicate a potential sex-bias in activation of neuroprotective signaling in the hippocampus of aging cTFEB:HSACre bigenic mice, with females showing decreased neuroinflammation (**Figure 5A-C**) and males showing increased neurotrophin expression levels (**Figure 5D**). Altogether, our results suggest that overexpression of TFEB in skeletal muscle throughout the lifespan promotes maintenance of protein quality control

444　regulation in the CNS, ultimately reducing proteostatic burden and neuroinflammation in the

445　aging hippocampus, similar to what we observed in our MAPT P301S studies.

446　Improved performance in neurocognitive testing of aging cTFEB;HSACre mice.

We next pursued neurocognitive testing in aged (16-18 months old) control

and cTFEB;HSACre mice. First, we evaluated spatial learning and memory using the Barnes

maze task, a hippocampal working memory test known to be sensitive to aging in mice. We

documented a significant decrease in the number of errors per trial (**Figure 5L**) and significantly

faster escape times (**Figure 5M**) in aged cTFEB;HSACre mice of both sexes in comparison to

controls. By the last trial, cTFEB;HSACre mice escaped the maze twice as quickly as control

mice. In the novel object recognition task – an independent behavioral test of hippocampal

recognition memory – we also found that aged cTFEB;HSACre mice of both sexes exhibited a

significantly greater number of contacts with the novel object compared to controls during the

test phase (**Figure 5N**). Indeed, aged cTFEB;HSACre mice have around a ~15% higher

preference for the novel object relative to their age-matched littermate controls (**Figure**

5O). Importantly, we also confirmed no differences in visual performance and motor activity (not

shown) between aged control and cTFEB;HSACre cohorts. Hence, these results provide

exciting evidence for a pronounced improvement in neural function in the aging brain

of cTFEB;HSACre mice.

462　Enhancing skeletal muscle TFEB-signaling transcriptionally remodels the hippocampus

463　of cTFEB;HSACre bigenic mice throughout the lifespan in a sex-dependent manner.

To determine the molecular basis of the benefits of enhancing skeletal muscle TFEB-signaling

on the aging CNS, we performed unbiased transcriptome analysis on hippocampal RNAs

isolated from young (6 months old) and aged (22-24 months old) mice (**Figures 6-7**). By this

time point, aging control animals have increased neuroinflammation (**Figure 5A-B**), have

evident CNS proteostasis dysfunction (**Figure 5G**), and display deficits in cognitive function

(**Figure 5L-O**), phenotypes that are significantly improved in cTFEB;HSACre mice.

Bulk RNA-Seq analysis of hippocampal RNA lysates revealed robust changes in gene

expression in young cTFEB;HSACre bigenic mice compared to their age- and sex-matched

littermate controls (**Figure 6, Figure 7, Supplemental Figure 4, and Supplemental Figure 5**).

In young female cTFEB;HSACre hippocampi, we detected 618 differentially expressed genes,

(DEGs, 435 up-regulated and 165 down-regulated) (**Figure 6A**). Functional GO Enrichment

analysis showed these DEGs mapped to key categories associated with synaptic function,

including ion- and voltage-gated channel activity, transporter complexes, as well as synapse

and synaptic membrane categories (**Figure 6C, boxes**). Enrichment analysis of differential

gene expression (KEGG) revealed additional critical categories associated with neural function,

including oxytocin signaling, cAMP signaling, Hippo signaling and gap junctions (**Figure 6E**).

Furthermore, metabolic and functional pathway analysis (Reactome) also identified key

categories known to regulate cognitive plasticity, including GPCR ligand binding, voltage gated

potassium channels, and transmission across chemical synapses, among others

(**Supplemental Figure 4A**).

Consistent with our biochemical and functional findings suggesting increased neuroprotective

benefits in aging cTFEB;HSACre mice (**Figure 5**), we found that these transcriptional changes

modulating synaptic function pathways are still present (although not as prominent) in the aging

hippocampus of aged female cTFEB;HSACre mice (**Figure 6B, Supplemental Figure 4B&D**).

Out of a total of 1765 DEGs, (541 up-regulated and 1224 down-regulated), KEGG functional

pathways enriched in our aged female hippocampus included key categories associated with

preservation of neuronal activity and cognitive function, including axon guidance, HIF-1

signaling, AMPK signaling and PI3-Akt signaling (**Figure 6F**). We confirmed the age-associated

differential expression of a subset of our top-hit genes in a second group of hippocampal RNA

lysates using targeted qRT-PCR (**Supplemental Figure 4E**). Altogether, this suggests that

overexpression of TFEB in skeletal muscle activates multiple neuroprotective gene networks

associated with synaptic function and signaling in the young and aged female hippocampus.

Conversely, in the young male cTFEB;HSACre hippocampus mice we detected 471 DEGs with

a 1.25-fold expression change (216 up-regulated and 255 down-regulated), which GO-term

analysis associated mostly with multiple mitochondria or ribosomal/transcription targeting

pathways (**Figure 7A and C, boxes**). This included mitochondrial respiratory chain,

mitochondrial protein complexes, electron transport chain and ATP metabolic processes

(**Figure 7C**). Additional enriched metabolic pathways in the hippocampus of young

cTFEB;HSACre males included non-sense mediated decay, metabolism of RNA and translation

(**Supplemental Figure 5A**). Consistent with this, KEGG analysis revealed the most enriched

functional pathways were ribosomes and oxidative phosphorylation (**Figure 7E**). As before, we

confirmed the differential expression of a random sampling of our top-hit genes using targeted

qRT-PCR (**Supplemental Figure 5E).** When we performed similar analysis on the male aging

hippocampus, we detected 539 DEGs in the hippocampus of aged male cTFEB;HSACre mice

(278 up-regulated, 261 down-regulated) (**Figure 7B**), with only mild significance on functional

pathways (**Figure 7D, Figure 7F, Supplemental Figure 5B, and Supplemental Figure 5D**).

Further analysis also revealed that most DEGs appeared to be sex-specific. Out of all DEGs in

young cTFEB;HSACre hippocampi (618 in females and 471 in males), there were only 25

shared DEGs, with 16 of these changing in the same direction in both sexes (**Supplemental**

Figure 6A). A similar pattern was found in the aged cTFEB;HSACre hippocampus (1765 total

DEGs in females and 539 DEGs in males), with 39 shared DEGs and only 3 genes changing in

the same direction in both sexes (**Supplemental Figure 6B**). Our results thus strongly suggest

a sex bias in the hippocampal transcriptional responses associated with muscle-TFEB

overexpression, with the female hippocampi up-regulating neuronal signaling and synaptic

function and the male hippocampi upregulating transcriptional and mitochondrial responses.

Finally, many of these changes are preserved up until late stages in the lifespan of our model

(24 months of age), suggesting that these transcriptional changes may underlie the

neurocognitive benefits observed in our aging cTFEB;HSACre cohorts.

DISCUSSION

Skeletal muscle comprises roughly 40% of the total body mass in a healthy young adult human

and is highly susceptible to age-associated metabolic and functional decline [60,61]. Skeletal

muscle health has recently been linked to multiple chronic age-associated conditions, including

neurodegenerative diseases [24,26], but the role of the so-called muscle-to-brain axis on the

pathology of these conditions remains largely unknown. Here, and for the first time in mammals,

we provide direct evidence demonstrating the CNS benefits of enhancing muscle-TFEB

expression, both during healthy aging and in the context of neurodegenerative proteinopathy.

We found that lifelong overexpression of TFEB, a master regulator of lysosomal and

mitochondrial function, reduces age-associated functional and biochemical hallmarks of aging in

skeletal muscle [1] **(Figure 1-2)**. We confirmed previous reports showing that in young skeletal

muscle, acute TFEB-overexpression induces the expression of genes involved in mitochondrial

biogenesis, fatty acid oxidation, and oxidative phosphorylation, optimizing mitochondrial

substrate utilization and enhancing exercise capacity [36]. Furthermore, we extend these findings

and report that life-long overexpression of muscle-TFEB counteracts age-associated

mitochondrial dysfunction in skeletal muscle (**Figure 2**) [1,44], ultimately preserving skeletal

muscle mass and preventing fiber type switching during aging (**Figure 1**). Indeed, young TFEB-

overexpressing muscle had significant increases in the abundance of Type 1 and Type 2A

fibers, which are considered to be highly reliant on oxidative phosphorylation and thus more

resistant to fatigue. This is consistent with the observed increases in exercise performance of

cTFEB;HSACre transgenic mice at both ages (**Figure 1**). Importantly, although we focused our

analysis on two key hallmarks of aging (lysosomal and mitochondrial function), our proteomics

studies also detected important changes in branched chain amino acid signaling, which have

been recently linked to lifespan and healthspan extension in rodents [62], indicating that TFEB

overexpression may have pan-geroprotective effects on aging skeletal muscle.

The hippocampus is a critical structure in cognitive aging, and hippocampal atrophy and

dysfunction are key features of both aging and neurodegenerative diseases [63,64]. In addition to

rescuing age-associated hallmarks in skeletal muscle, we also discovered significant

improvements in multiple disease-associated biochemical hallmarks in the hippocampus of

cTFEB;HSACre transgenic mice. Indeed, we observed robust benefits on morphological and

transcriptional markers of neuroinflammation in the MAPT P301S model of tau toxicity, as well

as a significant extension on the lifespan of male MAPT P301S;cTFEB;HSACre transgenic mice

(**Figure 4**). We also report important reductions in proteotoxicity and neuroinflammation and

improved neurocognitive performance in healthy aging (22-24 month old) cTFEB;HSACre mice

(**Figure 5**). Importantly, we observed these benefits without any detectable expression of our

transgene in the CNS (**Supplemental Figure 1**). Although previous work examining the muscle-

to-brain axis has demonstrated important benefits on the accumulation of misfolded protein

inclusions in the CNS [5,8,19,28], similar to what we observed in the MAPT P301S;cTFEB;HSACre

hippocampus, the profound reductions in neuroinflammation observed in our model suggest

modulation of skeletal muscle TFEB-signaling may be a novel target to modulate

neuroinflammatory signaling in vivo in the context of aging and/or neurodegenerative diseases.

Previous transcriptional studies of the healthy aging hippocampus have identified dysregulation

of ion homeostasis, disruption of neurotransmission, and ribosome biogenesis as key biological

process exhibiting differential regulation with aging [2,65]. Specifically, genes related to synaptic

transmission and plasticity appear to be particularly sensitive to age-associated transcriptional

dysregulation (reviewed here [3]). Many of these gene networks are also significantly altered in

AD and other associated tauopathies [57]. It was quite interesting that similar networks were

transcriptionally remodeled in the healthy aging cTFEB;HSACre hippocampus, suggesting that

these gene networks may be uniquely sensitive to both neurotoxic (aging, neurodegeneration)

and neurotrophic (muscle-TFEB overexpression) signaling. Indeed, manipulation of skeletal

muscle TFEB modulated expression of similar functional enrichment categories (i.e. trophic

factor signaling, neuroinflammation) in the aging and tau-afflicted hippocampus of our

cTFEB;HSACre mice in a sex-dependent manner. When we compared KEGG pathways of the

differentially expressed genes from the nCounter AD panel in MAPT P301S;cTFEB;HSACre

hippocampus with those enriched in the hippocampus of our 24-month old cTFEB;HSAcre mice,

we see shared regulation of pathways involved in axon guidance, MAPK signaling, cAMP

signaling, and Neuroactive ligand-receptor interactions, suggesting potential converging

neurotrophic benefits associated with muscle-TFEB expression.

One of our most exciting (and unexpected) findings is the sex-bias in our CNS and skeletal

muscle responses to muscle-TFEB overexpression. In general, we struggled to identify previous

literature thoroughly describing female-specific changes to many of these hallmarks of aging on

either tissue [1]. To our surprise, while we were able to validate 'classical' age-associated findings

such as fiber type switching in aging control males, these phenotypes were less robust or not

present in age-matched control females (**Figure 1**), rendering interpretation of the sex-bias of

our CNS-targeting benefits difficult. While we ultimately observed neuroprotective effects on

both male and female cTFEB;HSACre transgenic mice, it appears that the mechanisms

underlying these benefits are not the same. For example, whereas muscle-TFEB

overexpression in the MAPT P301S model reduces the accumulation of hyperphosphorylated

tau in both sexes, it appears to drive the transcriptional silencing of pro-inflammatory microglial

signaling in females, and activate trophic factor signaling in males instead. We also detected

similar sex-differences in our healthy aging transcriptional studies, with the female

cTFEB;HSACre hippocampus activating pathways associated with synaptic function, while the

male cTFEB;HSACre hippocampus activated ribosomal and mitochondrial signaling instead.

One potential explanation could be the sex-bias in TFEB expression in skeletal muscle, which is

~50% higher in male mice, suggesting a threshold effect for local remodeling of skeletal muscle.

However, the fact both sexes do display similar exercise endurance at young ages and

comparable protection of mitochondrial respiration rates at old ages again suggest converging

outcomes of TFEB-mediated geroprotection of aging skeletal muscle. Another consideration is

that in our healthy aging studies, both sexes are well into the stage of reproductive senescence,

which has been reported to have significant negative impacts on neural function [66]. While we did

not measure circulating levels of sex hormones in our models, there is plenty of evidence for de

novo synthesis on neuroactive steroids in the brain having neurotrophic and anti-inflammatory

benefits, the phenotypes we also see being modified in our cTFEB;HSACre groups.

Interestingly, CNS estrogen appears to control similar signaling pathways as those we detected

as differentially expressed in the hippocampus of muscle-TFEB overexpressing female (i.e.

synapse function, neurotransmitter signaling) and male (i.e. mitochondrial function) mice [66].

While it is unclear whether these differential responses between male and female hippocampi

reflect local differences in the CNS itself (as recently published here [67]) or peripheral differences

in the expression, secretion or trafficking of circulating signals (such as Ctsb, for example), they

underscore the fundamental need to examine sex differences contributing to the biology of

aging and neurodegenerative diseases [68].

Recent evidence has shown that muscle-originating circulating factors (myokines) appear to

play central roles in regulating CNS health and function. Ctsb is secreted from skeletal muscle

into circulation in response to exercise, and is required for the full manifestation of exercise-

associated benefits on the CNS, including increases in hippocampal Bdnf and the activation of

neurogenesis [20]. Circulating Ctsb has also been proposed to cross the blood-brain barrier [20],

and appears to remodel the extracellular matrix [69], enhancing axonal outgrowth through

degradation of chondroitin sulfate [70,71]. Furthermore, high Ctsb expression levels in the

hippocampus have been reported in low-anxiety mouse lines [72,73], suggesting that Ctsb may

play important roles in maintaining neuronal homeostasis in brain regions with high relevance

for both aging and age-associated neurodegenerative disease. Through our proteomics in silico

analysis, we identified Ctsb as our highest expressed secreted factor in cTFEB;HSACre skeletal

muscle, and we confirmed higher levels of circulating Ctsb in the serum of cTFEB;HSACre

transgenic mice. Our results imply that additional research into the role and function of

circulating proteases and their ability to remodel the CNS may be warranted. Interestingly,

recent evidence demonstrates pronounced involvement of the (macro)autophagy molecular

machinery in cellular secretion [74]. Indeed, key signaling molecules and cytokines associated

with inflammation, such as IL1B (interleukin 1 beta) and IL6 (interleukin 6), are released into the

extracellular space via autophagy-dependent secretion. There is also a growing body of

evidence that age [46] and exercise [75], known modifiers of skeletal muscle metabolism, alter the

composition of the skeletal muscle secretome. This suggests that activation of TFEB-

overexpression may be modifying the skeletal muscle secretome, potentially enriching it with

neurotrophic factors or depleting it from neurotoxic factors, although this hypothesis remains to

be tested.

Over the last ten years, there has been growing evidence that suggests prominent contributions

of the periphery to the etiology of neurodegenerative diseases. Our discovery that skeletal

muscle TFEB-signaling can be a crucial site for regulation of CNS health and function provides

compelling evidence for a new therapeutic delivery avenue into the brain, providing a currently

unexplored new diagnostic and therapeutic intervention site (i.e. skeletal muscle) for preserving

brain health.

Authorship Contributions

IM, CJC, AG, HM, AB, AH, KL, EL, KC, MM, DP, MB, AN, NN, GM, LW, MM, NM, EB and EF:

Performed the experimental work. Reviewed and edited the manuscript.

CJC and IM: Wrote and edited the manuscript.

CJC and ALS: Conceptualized the experiments outlined here and obtained all associated

funding.

Figure Legends

Figure 1: Skeletal muscle overexpression of TFEB preserves muscle size, fiber typing and increases exercise endurance during aging.

(A) Schematic depicting 3x-FLAG human TFEB transgene. Cassettes are arranged from 5' (left) to 3' (right). Note that the 3x-FLAG-EGFP-STOP cassette (flanked by loxP sites) prevents expression of the 3x-FLAG human TFEB transgene unless in the presence of Cre recombinase.

(B) Immunoblot of control and cTFEB;HSACre quadriceps lysates confirming 3x-FLAG-TFEB protein expression in young (5-6 months) and aged (22-24 months) mice of both sexes. C=control, T=cTFEB;HSACre individuals. Transgenic 3x-FLAG-TFEB protein weighs ~75 kDa. Total Protein Stain Free lanes are shown below as a loading control. Densitometry quantification relative to average of young male control individuals to the right.

(C-E) Skeletal muscle (quadriceps, C and gastrocnemius, D) wet weight and body weight (E) in young (6 months) and aged (22-24 months) control and cTFEB;HSACre transgenic mice of both sexes.

(F) Representative images of fibertyping stains of cross-section of gastrocnemius muscle of young (7-9 months) and aged (22-24 months) control and cTFEB;HSACre transgenic mice of both sexes. MyHC1 (Fiber Type 1) in red, MyHC2A (Fiber Type 2A) in green, MyHC2B (Fiber Type 2B) in blue, MyHC2X (Fiber Type 2x) unstained, and laminin in white. Scale bars = 500 μm.

(G) Fiber cross sectional area distribution curves from (F). *, statistical difference between control and cTFEB;HSACre. #, statistical difference between young and aged controls. Each data point represents the average of 3 separate images collected from 3 sections/individual.

(H-K) Quantifications of each fiber type percentage from (F). Each data point represents the average of 3 separate images collected from 3 sections/individual.

(L-M) High-intensity exhaustive treadmill exercise of young (6 months) and aged (18 months) control and cTFEB;HSACre transgenic mice of both sexes showing maximum distance run till exhaustion (L) and total work (body weight/distance run, M).

Each point represents the average of all data from a single individual. Controls are age-matched littermates. Unless otherwise stated, data are presented as mean ± SEM. */# $p<0.05$, **/## $p<0.01$, ****/#### $p<0.0001$ ANOVA and post-hoc multiple comparisons for each sex. Lack of annotation indicates comparisons were not significant.

Figure 2: Muscle TFEB increases lysosomal network size and preserves mitochondrial function during aging.

(A-B) Volcano plot of proteomics analysis depicting differentially regulated proteins in young (A, 6 months old) and aged (B, 18 months old) cTFEB;HSACre quadriceps muscle relative to controls. Overexpressed proteins in red, underexpressed proteins in blue. TFEB (rectangles) is highly expressed in both aged and young muscle.

(C) STRING predicted protein-protein interactions of overexpressed proteins from (A) showing interacting networks converging on TFEB (blue) and mTOR (green).

(D) KEGG enrichment analysis clusters overexpressed proteins from (A-B) into categories associated with proteostasis and autophagy (orange), mitochondrial metabolism (gold), fatty acid metabolism (green), age-associated signaling (blue), and amino acid metabolism (purple).

(E) qRT-PCR for *Lamp1*, *Atg5*, *Ctsd*, and *CtsB* expression in quadriceps mRNA lysates from young (6 months old) male cTFEB;HSACre mice and controls.

(F) Immunoblot analysis for LAMP1 and Cathepsin D protein from young (5-6 months) and aged (22-24 months) control and cTFEB;HSACre quadriceps lysates mice of both sexes. Total Protein Stain Free lanes are shown as a loading control. Marker densitometry quantification relative to Protein Stain Free densitometry shown below.

(G) Representative images of gastrocnemius cross section from young (6-7 months old) control and cTFEB;HSACre female mice stained for LAMP1 (green), nuclei/Hoechst (blue). Quantification of LAMP1+ puncta and LAMP1+ area/muscle section for both sexes are shown on the right. Scale bars = 100 μm.

(H-I) Mitochondrial function measured by oxygen flux normalized to citrate synthase activity in soleus muscle from young (6 months old) and aged (18 months old) male (H) and female (I) control and cTFEB;HSACre semi-permeabilized soleus muscle. CI_P is complex I and CI+CII_P is complex I and II linked respiration coupled to ATP production. CI+CII_E is uncoupled complex I and II respiration or maximum respiration. CII_E is complex II respiration in the uncoupled state.

Each point represents the average of all data collected from an individual. Controls are age-matched littermates. All data were analyzed by 2-way between-subject (age × genotype) ANOVA with post hoc multiple comparisons, or one-way ANOVA and multiple comparisons post-hoc for each sex. Data are presented as mean ± SEM. * $p<0.05$, ** $p<0.01$. Lack of annotation indicates comparisons were not significant.

Figure 3: Increased levels of known CNS-targeting myokines in TFEB-expressing skeletal muscle.

(A) VerSeDa-predicted secreted proteins identified as up-regulated or down-regulated in young (green) and aged (blue) cTFEB;HSACre transgenic muscle, from proteomics studies from Figure 2A-B.

(B) Immunoblot analysis for Cathepsin B protein in quadriceps protein lysates from young (5-6 months) and aged (22-24 months) control and cTFEB;HSACre quadriceps lysates mice of both sexes. Note that the mature fragment of Cathepsin B is the secreted form. Total Protein Stain Free lanes are shown as a loading control. Marker densitometry quantification relative to Protein Stain Free densitometry shown below.

(C) Levels of cathepsin B in serum from young (5-6 months) and aged (22-24 months) control and cTFEB;HSACre mice of both sexes.

Each point represents the average of technical replicates from an individual. Controls are age-matched littermates. All data were analyzed by 2-way between-subject (age × genotype) ANOVA with post hoc multiple comparisons, data are presented as mean ± SEM. * $p<0.05$, ** $p<0.01$. Lack of annotation indicates comparisons were not significant.

Figure 4: Skeletal muscle TFEB-overexpression reduces accumulation of hyperphosphorylated tau and microglial activation in a mouse model of tau pathology.

(A) Immunoblot of 3x-FLAG TFEB protein in quadriceps muscle lysates from 9-month old MAPT P301S;cTFEB;HSACre and MAPT P301S mice (top). Immunoblot for total Tau protein (T6 antibody) in hippocampal lysates from same individuals (bottom). GAPDH is shown below as a loading control.

(B) Representative merged images of the hippocampus dentate gyrus stained for phosphorylated tau (AT8, white) and nuclei/Hoechst (blue) in 9-month old control (top, in gold), MAPT P301S (middle, in teal), and MAPT P301S;cTFEB;HSACre mice (bottom, in purple). Insets depicting perinuclear AT8 staining are 5X zooms of areas demarcated by yellow squares. Representative images are from male mice. Quantification of total phosphotau staining/section (right, top) and for intracellular phosphotau staining, (right, bottom). Each point represents the average of all sections containing the dentate gyrus (2-3 sections) for an individual animal. Scale bars = 100 μm.

(C) Representative merged images of the dentate gyrus stained for astrocytes (GFAP, green), microglia (IBA1, red) and nuclei (Hoechst, blue) in 9-month old control (top, gold), MAPT P301S (middle, teal), and MAPT P301S;cTFEB;HSACre mice (bottom, purple). Insets depicting glia

morphology are 5X zooms of areas demarcated by yellow squares. Representative images are from male mice. Quantification of IBA1+ microglia volume (top, right) and GFAP+ astrocyte staining amounts per section (bottom, right). Each point represents the average of all sections containing the dentate gyrus (2-3 sections) for an individual animal. Scale bars = 100 µm.

(D-E) Nanostring nCounter AD panel cluster analysis of differentially expressed genes for gene clusters associated with microglial activation (D) and with trophic factors (E) in P301S MAPT hippocampi (teal) compared to MAPT P301S;cTFEB;HSACre (purple) age-matched littermates of both sexes. Analysis done via nSolver (Nanostring) differential gene expression analysis software. Each point represents cluster scoring for a single individual hippocampal mRNA lysate.

(F-G) qRT-PCR for *Bdnf* (F) and *Fndc5* (G) expression in hippocampal RNA lysates from 9-month old male MAPT P301S (teal) and MAPT P301S;cTFEB;HSACre mice (purple) relative to controls (not shown).

(H-I) Survival curves for male (H) and female (I) control (gold), MAPT P301S (teal), and MAPT P301S;cTFEB;HSACre mice (purple).

Controls are littermates. Unless otherwise noted, data are represented as mean ± SEM. * p<0.05, ** p<0.01,*** p<0.001, One-way ANOVA, post-hoc multiple comparisons (B,C), t-test (D-G) and log-rank (Mantel-Cox) test (H-I). Lack of annotation indicates comparisons were not significant.

Figure 5: Skeletal muscle TFEB-overexpression decreases neuroinflammation, lipofuscin accumulation, and improves cognitive performance in aged mice.

(A-D) qRT-PCR for *Ccl2* (A), *Nfkb* (B), *Il6* (C), and *Bdnf* (D) expression in hippocampal mRNA lysates from 18-month old cTFEB;HSACre mice and controls.

(E-F) Representative images dentate gyrus of the hippocampus stained for Hoechst (blue), IBA1 (E, red, representative images are from females), and GFAP (F, green, representative images are from males) in 18-22-month old control (top) and cTFEB;HSACre (bottom) mice. Insets depicting glia morphology are 5X zooms of areas demarcated by yellow squares. Quantification of IBA1+ microglial volume (E, top right) and IBA1+ branching index (E, bottom right). Average number of GFAP+ astrocytes pixels per section (F, top) and GFAP+ astrocyte branching index (F, bottom). Each point represents the average of all sections containing the dentate gyrus from an individual. Scale bars = 100 µm.

(G) Representative merged images of the dentate gyrus of 18-22-month old control (top) and cTFEB;HSACre (middle) mice stained for Hoechst (blue), and lipofuscin puncta

(autofluorescence, green, representative images are from females). Insets depicting

autofluorescent puncta within the dentate gyrus are 5X zooms of areas demarcated by yellow

squares. Scale bars = 100 μm.

(H-K) qRT-PCR for *Atg8* (H), *Lamp1* (I), *Sqstm1* (J), and *Ctsd* (K) expression in hippocampal

lysates from young (6 months old, closed diamonds) and aged (21 months old, open diamonds)

cTFEB;HSACre mice (purple diamonds) and controls (gold diamonds).

(L-Q) Neurocognitive phenotyping of aged (18 months old) cTFEB;HSACre and control animals.

Barnes maze (L,M), novel object recognition (N,O).

Controls are age-matched littermates. Each point represents the average of all data collected

from an individual. Data are represented as mean ± SEM. * p<0.05, ** p<0.01, One-way

ANOVA or two-way RMANOVA, post-hoc-tests. Lack of annotation indicates comparisons were

not significant.

Figure 6: Transcriptome changes in the hippocampi of female cTFEB;HSACre transgenic

mice suggest synaptic remodeling and improved synaptic function.

(A-B) Volcano plot of differentially expressed genes in hippocampal lysates from female young

(A, 6 months old) and aged (B, 24 months old) cTFEB;HSACre mice relative to controls.

(C-D) GO term analysis of differentially expressed genes from A and B including gene counts

and adjusted p values. Boxes highlight neurotrophic signaling pathways.

(E-F) KEGG analysis of differentially expressed genes from A and B. Number of differentially

expressed genes (DE) are shown above each category. Boxes highlight categories associated

with neural function.

Figure 7: Transcriptome changes in the hippocampi of male cTFEB;HSACre transgenic

mice suggest activation of mitochondrial and ribosomal signaling.

(A-B) Volcano plot of differentially expressed genes in hippocampal lysates from male young (A,

6 months old, 2 fold change) and aged (B, 24 months old, 1.25 fold change) cTFEB;HSACre

mice relative to controls.

(C-D) GO term analysis of differentially expressed genes from A and B including gene counts

and adjusted p values. Boxes highlight ribosomal and mitochondrial pathways.

(E-F) KEGG analysis of differentially expressed genes from A and B. Number of differentially

expressed genes (DE) are shown above each category. Boxes highlight ribosomal and

mitochondrial pathways.

Supplemental Figures

Supplemental Figure 1: Skeletal muscle-restricted expression of human TFEB protein.

(A-B) qRT-PCR analysis for human *TFEB* mRNA expression in 6-month old (A) and 21-24 month old (B) control and cTFEB;HSACre quadriceps skeletal muscle, heart ventricles and hippocampus.

(C) Representative images of gastrocnemius cross sections stained for Hoechst (blue) and Human TFEB (white). Insets of nuclear hTFEB staining are 5x zooms of the areas denoted by the yellow squares. Scale bars = 100 μm.

(D-G) Immunoblot analysis for 3x-FLAG TFEB protein expression from lysates of brown adipose (D), liver (E), heart ventricles (F), and brain (G) tissue from 6-month old female cTFEB;HSACre mice and control littermates. cTFEB;HSACre skeletal muscle lysates are shown in the first lane of each blot as a positive control for 3x-FLAG TFEB expression. FLAG-eGFP protein cassette (arrow) as well as non-specific bands (asterisks) are indicated.

(H) Representative images of quadriceps and gastrocnemius cross sections from 7-week old male TdTomato;HSACre transgenic mice stained for Hoechst (blue), TdTomato (red), and Laminin (white). Scale bars = 100 μm.

(I) Representative images of hemibrain sections from 7-week old male TdTomato;HSACre transgenic mice stained for Hoechst (blue), TdTomato (red). Insets of hippocampal regions are 5x zooms of the areas denoted by the yellow squares. Scale bars = 1 cm.

Controls are age-matched littermates. Each point represents the average of all data collected from an individual. Data are represented as mean ± SEM. One-way ANOVA, post-hoc multiple comparisons * $p<0.05$, ** $p<0.01$, *** $p<0.001$. Lack of annotation indicates comparisons were not significant.

Supplemental Figure 2: Fiber typing metrics in cTFEB;HSACre transgenic muscle.

(A-H) Feret's minimum diameter (A-D) and fiber cross sectional area (E-H) of all 4 fiber types in gastrocnemius muscle of young (7-9 months) and aged (22-24 months) control and cTFEB;HSACre transgenic mice of both sexes.

(I) Immunoblot analysis for LC3 and total ubiquitin in quadriceps protein lysates from young (5-6 months) and aged (22-24 months) control and cTFEB;HSACre mice of both sexes. Total Protein Stain Free lanes are shown as a loading control. Marker densitometry quantification relative to Protein Stain Free densitometry shown below.

Controls are age-matched littermates. Each point represents the average of all data collected

from an individual. Data are presented as mean ± SEM. * p<0.05, ** p<0.01, **** p<0.0001

ANOVA and interactions of sex x age post-hoc. Lack of annotation indicates comparisons were

not significant.

Supplemental Figure 3: Muscle-TFEB overexpression reduces hyperphosphorylated tau

accumulation and modifies expression of AD-associated genes in the hippocampus of

MAPT P301S transgenic mice.

(A) Representative images of the hippocampal dentate gyrus from 9-month old control (top),

MAPT P301S (middle), and MAPT P301S;cTFEB;HSACre (bottom) mice stained for Hoechst

(blue) and AT180 PhosphoTau (white). Insets of hippocampal regions are 5x zooms of the

areas denoted by the yellow squares. Representative images are from males. Scale bars= 100

μm.

(B-C) Volcano plot of differentially expressed genes in the hippocampi of 9-month old male (B)

and female (C) MAPT P301S;cTFEB:HSACre animals relative to age-matched P301S MAPT

littermates using the Nanostring nCounter AD panel. Down-regulated genes in blue, up-

regulated genes in red.

(D) Cumulative ataxia score of 9-month old control, MAPT P301S, and MAPT

P301S;cTFEB;HSACre mice of both sexes.

(E) Weight (in grams) of 11-month old control MAPT P301S, and MAPT P301S;cTFEB;HSACre

mice of both sexes.

Controls are age-matched littermates. Each point represents the average of all data collected

from an individual. Data are represented as mean ± SEM. One-way ANOVA, post-hoc multiple

comparisons * p<0.05, ** p<0.01. Lack of annotation indicates comparisons were not significant.

Supplemental Figure 4: Functional analysis of differentially expressed genes in the

hippocampus of female cTFEB;HSACre transgenic mice.

(A-B) Reactome analysis of differentially expressed genes in hippocampal mRNA lysates from

female young (6 months old, A) and aged (24 months old, B) cTFEB;HSACre mice. Number of

differentially expressed genes (DE) are shown above each category.

(C-D) GO enrichment analysis of differentially expressed genes from A-B. Number of

differentially expressed genes (DE) are shown above each category.

(E) qRT-PCR analysis for select DE genes in hippocampal mRNA lysates from female young (6 months old, solid circles) and aged (24 months old, open circles) control (gold) and cTFEB;HSACre (purple) mice. Expression levels normalized to young controls. Controls are age-matched littermates. Each point represents the average of all data collected from an individual. Data are represented as mean ± SEM, O One-way ANOVA, post-hoc multiple comparisons, * $p < 0.05$, ** $p < 0.01$. Lack of annotation indicates comparisons were not significant.

Supplemental Figure 5: Functional analysis of differentially expressed genes in the hippocampus of male cTFEB;HSACre transgenic mice

(A-B) Reactome analysis of differentially expressed genes in hippocampal mRNA lysates from male young (6 months old, A) and aged (24 months old, B) cTFEB;HSACre mice. Number of differentially expressed genes (DE) are shown above each category.

(C-D) GO enrichment analysis of differentially expressed genes from A-B. Number of differentially expressed genes (DE) are shown above each category.

(E) qRT-PCR analysis for select DE genes in hippocampal mRNA lysates from female young (6 months old, solid circles) and aged (24 months old, open circles) control (gold) and cTFEB;HSACre (purple) mice. Expression levels normalized to young controls. Controls are age-matched littermates. Each point represents the average of all data collected from an individual. Data are represented as mean ± SEM, One-way ANOVA, post-hoc multiple comparisons, * $p < 0.05$, ** $p < 0.01$. Lack of annotation indicates comparisons were not significant.

Supplemental Figure 6: Overlap of differentially expressed genes confirms sex-specific transcriptional remodeling in the hippocampus of TFEB;HSACre transgenic mice.

(A) Overlap of differentially expressed genes (DEGs) in hippocampal mRNA lysates from young (6 months old) cTFEB;HSACre mice relative to their age-matched controls. Number of up- or down-regulated DEGs are shown next to each unique and shared category.

(B) Overlap of differentially expressed genes (DEGs) in hippocampal mRNA lysates from aged (24 months old) cTFEB;HSACre mice relative to their age-matched controls. Number of up- or down-regulated DEGs are shown next to each unique and shared category.

912 **STAR★Methods**

913 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-FLAG® M2	Sigma-Aldrich	RRID: AB_262044
Rabbit anti-TFEB	Cell Signaling Technology	RRID: AB_11220225
Rabbit monoclonal anti-LC3A/B	Cell Signaling Technology	RRID: AB_2617131
Mouse IgG1 monoclonal anti-GAPDH	Invitrogen	Cat #: AM4300
Rat IgG2a monoclonal anti-LAMP1	DSHB	Cat #: 1D4B
Rabbit anti-SQSTM1/p62	Cell Signaling Technology	Cat #: 5114T
Mouse monoclonal anti-Ubiquitin	Santa Cruz Biotechnology	RRID: AB_628423
Rabbit polyclonal anti-Cathepsin B (H190)	Cell Signaling Technology	RRID: AB_2292384
Goat polyclonal anti-Cathepsin D (C-20)	Santa Cruz	RRID: AB_637896
Chicken polyclonal anti-GFAP	Abcam	RRID: AB_304558
Rabbit polyclonal anti-IBA1	FUJIFILM Wako Shibayagi	RRID: AB_839504
Mouse monoclonal anti-Phospho-Tau (Thr231) (AT180)	Thermo Fisher Scientific	RRID: AB_223649
Mouse monoclonal anti-Phospho-Tau (Ser202, Thr205) (AT8)	Thermo Fisher Scientific	RRID: AB_223647
Rabbit polyclonal anti-Laminin	Abcam	RRID: AB_298179
Rabbit polyclonal anti-LC3B	Cell Signaling Technology	RRID: AB_915950
Guinea Pig polyclonal anti-p62	Progen	RRID: AB_2687531
Mouse monoclonal anti-Myosin Heavy Chain Type I	DSHB	RRID: AB_2235587
Mouse monoclonal anti-Myosin Heavy Chain Type IIA	DSHB	RRID: AB_2147165
Mouse monoclonal anti-Myosin Heavy Chain Type IIB	DSHB	RRID: AB_2266724
Bacterial and virus strains		
pCMV-human TFEB (NM_007162) Human Untagged Clone	Origene	CAT#: SC122773
Chemicals, peptides, and recombinant proteins		
2,2,2-Tribromoethanol ≥99%	VWR	Cat #: BT145585
2-Methyl-2-butanol 99%	VWR	Cat #: TS16662-0010
TRIzol	Invitrogen	Cat #: 15596026
Turbo DNase	Thermo Fisher	Cat #: AM2238
Sybrgreen	Life Technologies	Cat #: 4309155

RIPA Lysis and Extraction Buffer	Invitrogen	Cat #: 89900
1X Halt™ Protease and Phosphatase Inhibitor Cocktail	Invitrogen	Cat #: 78442
iScript™ cDNA Synthesis Kit	BioRad	Cat #: 1708891
SYBR™ Green PCR Master Mix	Life Technologies	Cat #: 4309155
Critical Commercial Assays		
Pierce™ BCA Protein Assay Kits	Invitrogen	Cat #: 23225
Mouse Cathepsin B ELISA Kit	Abcam	Cat #: ab303743
Deposited data		
TFEB Muscle proteomics	Panorama Public	ID: PXD045241
Hippocampal Bulk RNA-seq data	GEO	GEO: GSE242363
Experimental models: Organisms/strains		
Mouse: fxSTOP-TFEB	This paper	N/A
Mouse: HSA-Cre (FVB.Cg-Tg(ACTA1-cre)79Jme/J)	JAX Judith Melki	RRID: IMSR_JAX:006139
Mouse: TdTomato (B6.Cg-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J	JAX Philippe Soriano	RRID: IMSR_JAX:007909
Mouse: MAPT P301S (B6;C3-Tg(Prnp-MAPT*P301S)PS19Vle/J)	JAX Virginia Lee	RRID: IMSR_JAX:008169
Oligonucleotides		
Actb	IDT	Mm.PT.39a.22214843.g
Lamp1	Biorad	10025636
Atg5	IDT	Mm.PT.58.29236732
CTSD	IDT	Mm.PT.58.7639164
CTSB	IDT	237299388
Fndc5	IDT	Mm.PT.58.31333197
BDNF	IDT	Mm.PT.58.28814650
Ccl2	IDT	287573375
NFkB1	IDT	287573349
IL6	IDT	Mm.pt.58.10005566
Atg8 -LC3/ Map1lc3b		Mm.PT.58.29764292
Sqstm1	IDT	Mm.PT.58.5854953
Human TFEB	IDT	Hs.PT.58.50432039
Oprd1	IDT	Mm.PT.58.41742748
CCL3/MIP-1alpha	IDT	Mm.PT.58.29283216
Hgf	IDT	Mm.PT.58.43661644.g

Gpr88	IDT	Mm.PT.58.4366164 4.g
Stap1	IDT	Mm.PT.58.3163513 49
Gabrr2	IDT	Mm.PT.58.6928777 7
Pomc	IDT	Mm.PT.58.5917321 1
Fosb	IDT	Mm.PT.58.1099087 8
S100a8	IDT	Mm.PT.58.4400340 02.gs
S100a9	IDT	Mm.PT.58.4178757 62
Gnb3	IDT	Mm.PT.58.3303540 04
Tlr8	IDT	Mm.PT.58.1602115 50
Romo1	IDT	Mm.PT.56a.1113838 846
Software and algorithms		
FIJI	Schindelin et al.[76]	https://imagej.net/s oftware/fiji/
QuantStudio 3	Applied Biosystems	RRID:SCR_018712
STRING Version 12.0	STRING Consortium	RRID: SCR_005223
NanoString nCounter Analysis System	NanoString	RRID:SCR_021712

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be
fulfilled by the lead contact, Constanza Cortes (cjc_149@usc.edu).

Materials availability

The fxSTOp-TFEB line of transgenic mice is available for sharing with academic institutions
free-of-charge. For-profit institutions will adhere to UCSD's policy regarding copyrighted
material.

Data and code availability

- Bulk RNA-seq data have been deposited at GEO and are publicly available as of the date of publication. Accession numbers are listed in the key resources table. Muscle proteomics data have been deposited at Panorama Public and are publicly available as of the date of publication. ID numbers are listed in the key resources table. Microscopy data reported in this paper will be shared by the lead contact upon request.
- This paper does not report original code.
- Any additional information required to reanalyze the data reported in this work paper is available from the lead contact upon request.

Experimental model details

Mouse models used were the novel fxSTOP-TFEB transgenic line generated for this publication, the HSA-Cre transgenic mice (JAX Strain No: 006139 | HSA-Cre79) obtained from JAX, TdTomato mice (B6.Cg-Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze/J}$, JAX Strain #:007909) also from JAX, and the MAPT P301S line (Stock No: 008169) which originated in the laboratory of Dr. Virginia Lee. All animals have the C57BL/6J genetic background, both males and females were used wherever possible.

Method details

Generation of fxSTOP-TFEB transgenic mice.

We used a pCMV-human TFEB expressing vector from Origene (clone # sc122773), used previously [27]. We cloned a 3x-FLAG fragment via insertion into the Acc65I and Pac1 restriction sites of the targeting vector. The fxSTOP-TFEB vector was generated in multiple steps, as follows: (1) 5′ and 3′ targeting arms were inserted by PCR onto the ends of a minimal 1.8 kb, chloramphenicol-resistant vector; (2) an ampicillin resistance cassette was cloned into the MluI, XmaI sites of the targeting vector; (3) an 1.1 kb fragment encoding the actin promoter, a loxP site, a 3x-FLAG tag fused to EGFP was cloned into the NheI, PacI restriction sites of the targeting vector; (4) a 2.8 kb fragment containing the second loxP site, followed by 3x-FLAG-human TFEB sequence. The final fxSTOP-FLAG-human TFEB vector was microinjected into C57BL/6J:C3H/HeJ F1 hybrid oocytes. Of six founders identified, two gave rise to fxSTOP-FLAG-human TFEB lines with comparable phenotypes. Of these lines, we chose to focus our

studies principally on line 419. The presence of the floxed 3x-FLAG-eGFP STOP-3x-FLAG-

human TFEB cassette is verified by allele dependent qPCR genotyping analysis with each

generation (Transnetyx), and expression of the 3x-FLAG-eGFP sequence (upstream of the

STOP codon) in single transgenic fxSTOP-TFEB mice is assessed every other generation. We

confirmed excision of 3x-FLAG-EGFP-STOP cassette in the presence of Cre-recombinase by

crossing fxSTOP-TFEB transgenic female mice with HSA-Cre transgenic mice (JAX Strain No:

006139 | HSA-Cre79), and corresponds to the HSA-Cre driver mouse line from the laboratory of

Dr. Judith Melki. TdTomato mice (B6.Cg-Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze/J}$, JAX Strain

#:007909) express robust tdTomato fluorescence following Cre-mediated recombination, is

congenic on the C57BL/6J genetic background and were also obtained from JAX. The MAPT

P301S line (Stock No: 008169) originated in the laboratory of Dr. Virginia Lee.

cTFEB;HSACre;MAPT P301S transgenic mice were generated by crossing fxSTOP-TFEB

females to MAPT P301S males. Double transgenic offspring was then crossed with

heterozygote HSA-Cre transgenic individuals. All commercially available lines were obtained

directly from JAX. All lines have been maintained in a C57/B6 background for over ten

generations in our lab. Mice were housed in a temperature- and humidity-controlled

environment on a 12-hour light/dark cycle with food and water ad libitum. Mice of both sexes

were used for all experiments in equal numbers, or as close to equal numbers as possible. For

all MAPT P301 experiments, control animals are MAPT P301S;fxSTOP-TFEB-;HSACre- and/or

MAPT P301S;fxSTOP-TFEB+;HSACre- littermates. For all aging experiments, control animals

are fxSTOP-TFEB-;HSACre- and/or fxSTOP-TFEB+;HSACre- littermates. In our hands, we did

not detect any differences on tau or aging-associated CNS phenotypes on fxSTOP-TFEB+

animals compared to the wild-type controls. All animal experimentation adhered to NIH

guidelines and was approved by and performed in accordance with the University of California,

San Diego, Duke University, University of Alabama at Birmingham, and the University of

Southern California Institutional Animal Care and Use.

Tissue Collections.

Animals were anesthetized with 3.8% Avertin Solution. All animals received a transcardial

perfusion with 60 mLs of cold 1x PBS. Half of CNS tissue (right/left hemi-brain) was post-fixed in

4% PFA for less than 24 hours before being transferred to 30% sucrose for another 24 hours

before cryo-sectioning and staining the other half was micro-dissected into cortex and

hippocampus and flash-frozen in liquid nitrogen for RNA and protein analyses. Half of skeletal

muscle tissue (right/left limb muscle) was fresh frozen in OCT following dissection, while the

other half was flash-frozen in liquid nitrogen for RNA and protein analyses.

RT-PCR Analysis.

Flash frozen perfused isolated mouse tissues (quadriceps, hippocampus) were placed in plastic

tubes with silica beads (MP Biomedical, 116913100) and homogenized in the FastPrep-24™ 5G

bead beating grinder and lysis system (MP Biomedical, 116005500) in TRIzol (Invitrogen,

15596026). RNA was extracted via standard phenol-chloroform procedures followed by DNase

digestion (ThermoFisher, AM2238). cDNA was generated using iScript (Bio-Rad Technologies,

1708891). Quantitative PCR was run using Sybrgreen (Life Technologies, 4309155) for

chemical detection (Applied Biosystems; QuantStudio 3). Enrichment was determined based on

double-delta CT value. Primers were ordered from IDT Predesigned qPCR Assays unless

otherwise specified. Primers used are listed in the Key Resources Table.

Protein Extraction and ImmunoBlot Analysis.

Protein lysates from muscle tissue was prepared as previously described [27,28]. In short, Flash

frozen perfused isolated mouse tissues (quadriceps, hippocampus) were placed in plastic tubes

with silica beads (MP Biomedical, 116913100) and homogenized in the FastPrep-24™ 5G bead

grinder and lysis system (MP Biomedical, 116005500) in RIPA Lysis and Extraction Buffer

(Invitrogen, 89900), 1X Halt™ Protease and Phosphatase Inhibitor Cocktail (Invitrogen, 78442),

and 1% SDS. Protein concentration was quantified using a Pierce™ BCA Protein Assay

(23227). For brain tissue, Fifty or 25 µg of homogenized proteins were loaded per lane, and

after running Any KD, 10%, or 4-15% Mini-PROTEAN TGX Gels (BioRad, 4568124, 4561034,

and 4561084), samples were transferred to 0.45 µm PVDF membranes (BioRad, 1704275),

which were blocked in 5% BSA in PBS at RT for 1 hr. For skeletal muscle, proteins were

separated on 4-20% Stain-Free TGX gels (Bio-Rad, 5678092), separated to PVDF, and blocked

in 5% non-fat dry milk in TBST (1X TBS with 0.05% Tween-20) for 1 hour at RT. Membranes

were incubated with anti-FLAG antibody (Sigma, M2, F1804, 1:1000), anti-TFEB antibody (Cell

Signaling, 4240,1:1000), anti-Lamp1 antibody (DSHB, 1DB4, 1:1000), anti-LC3A/B antibody

(Cell Signaling, 12741, 1:1000), anti-P62 antibody (Cell signaling, 5114T, 1:1000), anti-ubiquitin

antibody (Santa Cruz, sc-8017, 1:1000), anti-Cathepsin B antibody (Cell signaling, 3383S,

1:1000), anti-Cathepsin D antibody (Santa Cruz, sc-6486, 1:1000),or anti-GAPDH (Invitrogen,

AM4300, 1:5000) in PBS-T with 5% BSA at 4°C overnight. The primary antibody was visualized

with horseradish-peroxidase conjugated anti-rabbit at (Cell Signaling, 7074P2, 1:5,000) and
enhanced chemiluminescence (BioRad, 170-5060) or goat-anti-mouse IgG 680 (Invitrogen,
A21058, 1:10,000). Densitometry analysis was performed using the BioRad Image Lab 6.1
software application.

Immunofluorescence Studies.

Tissue was embedded in OCT (TissueTek, 4583), frozen utilizing 2-methylbutane (VWR,
103525-278) submerged in liquid nitrogen and stored at -80 C until used. All samples were
sectioned on a standing cryostat (Leica). Brain sections were 20 microns thick, while muscle
sections were 15 microns thick. For immunohistochemistry, sections were permeabilized with
.25% Triton for 15 mins and blocked with 4% BSA for 30 minutes-1 hour. Brain sections were
then incubated with anti-GFAP (ab4674, Abcam, 1:200), anti-IBA1 (Wako, 019-19741, 1:200),
anti-AT180 TAU (ThermoFisher, MN1040, 1:100) and anti-AT8 TAU (ThermoFisher, MN1020,
1:100) while muscle sections were incubated with anti-laminin (Abcam, ab11575, 1:200), anti-
FLAG (Sigma, F1804, 1:1000), anti-LC3B (Cell Signaling, 2775S, 1:200), anti-LAMP1 (Novus,
NB100-77685, 1:200), anti-p62 (Progen, GP62-C, 1:200), anti-MHC Type 1 (DSHB, BD-D5,
1:100), anti-MHC Type 2A (DSHB, SC-71, 1:100), anti-MHC Type 2B (DSHB, BF-F3, 1:100),
and anti-hTFFEB (CST, 4240, 1:200) overnight at 4°C and incubated with secondary antibodies
at RT for 1 hour (both diluted in 4% BSA). Next the slides were washed with Hoescht
(ThermoFisher, 62249, 1:5000) and mounted with prolong glass (Invitrogen, P36984). All slides
were washed with 1X PBS three times for 5 minutes each between steps.

Slides with brain sections stained for TAU and glia markers were imaged in the UAB HIRF HRIF
Confocal/Light Microscopy Core utilizing the 10x objective on the Nikon A15R/SIM Confocal
microscope. Z-stacks of the entire hippocampus area/section were collected, and max intensity
projections were generated using FIJI. Lipofuscin imaging was performed using an
epifluorescent Nikon light microscope with a 20x objective. For TdTomato imaging, stitched
whole-section z-stacks were acquired using a Nikon A1R HD25 confocal microscope of nuclei in
the blue channel and native TdTomato fluorescence in the red channel. Muscle sections were
imaged with an ECHO Revolution epifluorescent microscope.

For quantification of astrocyte and/or microglia parameters, NIS elements was used to collect
morphometric information and numbers. A size exclusion parameter was used for GFAP
positive objects under 10 microns, IBA-1 positive objects under 15 microns, and any object

larger than 5,000 microns were all excluded due to standard assessments of cell sizes. FIJI was

used for quantification of lipofuscin and LAMP1 puncta. Muscle fibertype composition, number,

and size were quantified using Myosoft [77], a FIJI plugin.

Quantitative Proteomics Sample Preparation and Data-independent Acquisition Mass

Spectrometry.

Lysis/Digestion

We identified distinct protein species from crude total extractions of whole quadricep muscle by

mass spectrometry. Quadricep muscle powder lysate in 50 mM ammonium bicarbonate with

0.1% RapiGest (Waters, Cat# 186001861) were processed and digested using Qiagen Tissue

Lyser II (Qiagen) following the manufacturer's protocol. Protein concentrations were determined

using Pierce BCA Protein Assay Kit (ThermoFisher, PI23227). 150 µg of lysate was reduced

with DTT (Sigma, Cat # D0632), alkylated with IAA (Sigma, Cat # D0632) and digested at 1 µg

trypsin (Thermo, Cat # PI-90057) to 50 µg protein ratio for 16 hours at 37°C. Digests were

acidified with 200 mM HCl (Fisher, Cat # A144), cleaned with MCX columns (Waters, Cat#

186000782), dried with a vacuum concentrator, and reconstituted in 0.1% formic acid (Fisher,

Cat# A117) in water.

Liquid Chromatography and Mass Spectrometry

One µg of each sample with 150 fmol of Pierce Retention Time Calibrant (PRTC, Thermo, Cat #

PI-88321) was loaded onto a 30 cm fused silica picofrit 75 µm column (New Objective, Woburn,

MA, Cat # PF3607510N5) and 3.5 cm 150 µm fused silica Kasil1 (PQ Corporation) frit trap

loaded with 3 µm Reprosil-Pur C18 (ESI Source Solutions, Cat # r13.aq.0001) reverse-phase

resin analyzed with a Thermo Easy nano-LC 1000 coupled to a Thermo Orbitrap Fusion Lumos

Mass Spectrometer. Data-independent (DIA) mass spectrometry with gas-phase fractionation

was performed as described previously [78,79].

Data Analysis

Thermo DIA MS/MS RAW files were converted to mzML format using Proteowizard (version

3.0.19045) [80] using vendor peak picking and demultiplexing. Chromatogram spectral libraries

were created using default settings (10 ppm tolerances, trypsin digestion, HCD b- and y-ions) of

Walnut in EncyclopeDIA (version 0.6.14) [81] using the Uniprot mouse canonical FASTA [82].

Quantitative spectral libraries were created by mapping spectral to the chromatogram spectral

library using EncyclopeDIA requiring a minimum of 3 quantitative ions and filtering peptides at a

1% FDR using Percolator 3.01 [83]. The quantitative spectral library is imported into Skyline-daily

4.2.1.19058 [84] with the Uniprot mouse canonical FASTA as the background proteome to map

peptides to proteins. The mzML data is imported and all data is TIC normalized.

Data and code availability

The Skyline document and raw files for the proteomics DIA MS/MS data are available at

Panorama Public [85]. ProteomeXchange ID: PXD045241. Access URL:

https://panoramaweb.org/mouse-muscle-TFEB-proteomics.url. All original data has been

deposited at Panorama and is publicly available as of the date of publication. Any additional

information required to reanalyze the data reported in this paper is available from the lead

contact upon request.

High-Resolution Respirometry and ROS Production.

High-resolution respirometry (HRR) was conducted on a G-model Oxygraph-2k machine

(Oroboros Instruments, Innsbruck, Austria). Reagents, muscle fiber preparation, and Substrate

Uncoupler Inhibitor Titration (SUIT) protocols used for HRR have been previously described [86].

Antimycin A was sufficient to block complex 3 activity; therefore, myxothiazol was not added in

our SUIT protocol. ROS production was detected using the O2k-Fluo (Oroboros Instruments)

and a modified Amplex Red assay. After daily background correction, chambers were closed

and 20 µM final concentration Amplex Red ultra (Life Technologies, A36006), 1 U/mL

horseradish peroxidase (Sigma-P8250), and 5 U/mL superoxide dismutase (Sigma-S8409)

were added to the wells. After the amp-slope stabilized 0.1 µM H_2O_2 (Sigma-95321) was added

for calibration. Throughout the entire assay, repeated injections of 0.1 µM H_2O_2 were used for

calibration including one injection after the muscle tissue was added to correct for

autofluorescence. All subsequent 0.1 µM H_2O_2 injections were used to control for changing

concentrations of Amplex Red during the assay. Samples for HRR aere conducted in duplicate

and means were used for the final analyses. To eliminate confounds from tissue damage or

hyperpermeabilization, only samples with stable oxygen flux and Raw-amps were included in

the final analysis.

Gene Expression Analysis.

RNA Sequencing and Bioinformatics Analysis

RNA was extracted as mentioned above from flash frozen hippocampal tissue for four biological replicates (sex/genotype) and checked for quality by bioAnalyzer by Novogene (Novogene Corporation INC, Sacramento, CA, United States). mRNA libraries were prepared by Novogene using poly-T oligo-attached magnetic beads. cDNA was synthesized with random hexamer primers and dUTP or dTTP depending on library directionality. For the both non-directional and directional library prep; end repair, A-tailing, adapter ligation, size selection, amplification, and purification were performed. For the directional library, end repair, USER enzyme digestion was also performed. Libraries were checked via Qubit and qPCR for quantity and by bioanalyzer for size. Quantified libraries were pooled and bulk RNA-seq analysis was performed via NovaSeq PE150 (Illumina, San Diego, CA, United States) high throughput sequencing. Bioinformatics analysis for differentially expressed genes was performed by Novogene. Raw reads were cleaned to remove adapters, poly-N and low qualities reads from the data. The reference index was built using Hisat2 v2.05 and pair-end cleaned reads were aligned to the reference genome GRCm39 through Hisat2 v2.05. reads were counted using featureCounts v1.5.0-p3. FPKM of each gene was calculated based on the length of the gene and read counts mapped. Differential expression analysis was performed using the DEseq2 R package (1.20.0). Gene Ontology (GO) and KEGG enrichment analysis of differentially expressed genes was implemented by the clusterProfiler R package. For all analysis methods and graphic representations, a p-value less than 0.05 was considered significantly enriched by differentially expressed genes. For young (6-month) animals, a 2-fold change was used to determine differentially expressed genes. A 1.2 fold change was used for aged (24-month) animals. Volcano plots were generated in RStudio using ggplot2 using these criteria.

Nanostring nCounter AD panel

Assays were performed with 100 ng aliquots of RNA, extracted using the PureLink RNA extraction kit (Invitrogen) and run via the NanoString nCounter Analysis system (NanoString Technologies, Seattle, WA, USA) at the UAB Nanostring Core, following previously described and established protocols[54]. After codeset hybridization overnight, the samples were washed and immobilized to a cartridge using the NanoString nCounter Prep Station. Cartridges were scanned in the nCounter Digital Analyzer at 555 fields of view for the maximum level of sensitivity. Raw reads and the Nanostring Mm_AD30_v1.1 probe annotation codeset were imported into the Nanostring proprietary software, nSolver 4.0, and QC was performed following

company protocol guidelines. Counts for target genes were normalized to house-keeping genes included in the panel (Cltc, Gapdh, Gusb, Hprt, Pgk1, and Tubb5). Specifically, background correction was performed using the negative control at the cutoff of mean + 2 standard deviation. Housekeeping genes were used for normalization based on geometric mean, Gene expression was normalized using NanoStringNorm R package. All p-values were adjusted using a false discovery rate (FDR) correction of 1% for multiple comparisons. Differential expression and pathway scoring analysis, as well as graphical QC representation were performed in the nSolver Advanced Analysis Software 2.0 using a custom analysis to utilize generated normalized counts. Differential gene expression values were presented asMAPT P301S baseline compared to the MAPT P301S;cTFEB;HSACre cohort for each sex, and a p value of 0.05 or less was described as significant. Volcano plots were generated with ggplot2 in Rstudio using significant genes and a cutoff of 1.2 or greater fold change. The pathway scoring module summarizes a single score from the data from a pathway's genes [87].

Data and code availability

Bulk RNA-seq data have been deposited at GEO and are publicly available as of the date of publication. Accession numbers (GSE242363) are listed in the key resources table. No new code was generated or used in the analysis of this dataset. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Cathepsin B ELISA Analysis.

Fifty µl of serum (collected via blood cardiac puncture and isolated via standard coagulation/centrifugation protocols) was diluted 1:1 and processed with the Cathepsin B ELISA kit (Abcam, ab119585). Diluted samples were processed following kit instructions, and imaged at 450 nm (Tecan, Infinite M plex). Amount of cathepsin B was determined using a standard dilution curve of known concentration.

Mouse Phenotyping and Behavioral Studies.

Barnes Maze

The Barnes maze apparatus is an opaque Plexiglas disc 75 cm in diameter elevated 58 cm above the floor by a tripod. Twenty holes, 5 cm in diameter, are located 5 cm from the perimeter, and a black Plexiglas escape box (19 x 8 x 7 cm) is placed under one of the holes.

Distinct spatial cues are located all around the maze and are kept constant throughout the

study. On the first day of testing, a training session was performed, which consists of placing the

mouse in the escape box for one minute. After the one-minute habituation period, the first

session was started. At the beginning of each session, the mouse was placed in the middle of

the maze in a 10 cm high cylindrical black start chamber. After 10 seconds the start chamber

was removed, a buzzer (80 dB) and a light (400 lux) were turned on, and the mouse was set

free to explore the maze. The session ended when the mouse entered the escape tunnel or

after 3 min elapsed. When the mouse entered the escape tunnel, the buzzer was turned off and

the mouse was allowed to remain in the dark for one minute. When the mouse did not enter the

tunnel by itself it was gently put in the escape box for one minute. The tunnel was always

located underneath the same hole (stable within the spatial environment), which was randomly

determined for each mouse. Mice were tested once a day for 4 days for the acquisition portion

of the study. For the 5th test (probe test), the escape tunnel was removed and the mouse was

allowed to freely explore the maze for 3 min. The time spent in each quadrant was determined

and the percent time spent in the target quadrant (the one originally containing the escape box)

was compared with the average percent time in the other three quadrants. This was a direct test

of spatial memory as there was no potential for local cues to be used in the mouse's behavioral

decision. Two weeks later the mice were tested again with the escape box in the original

position (retention test). This allows for the examination of long-term memory. Finally, on the

day after this test, the escape tunnel was moved to a new location (90 degrees from the original

position) and the behavior of the mouse was recorded. This is called the reversal test and it

allows for the examination of perseveration at the old hole as well as the working memory

strategies the mice adopted to locate the new tunnel location. Each session was videotaped and

scored by an experimenter blind to the genotype of the mouse. Measures recorded include the

latency to enter the escape box and the number of errors made per session. Errors are defined

as nose pokes and head deflections over any hole that did not have the tunnel beneath it. The

probe data were analyzed using Noldus Ethovision software to determine time spent in each

quadrant of the maze as well as to assess activity measures.

Novel Object Recognition

Mice were individually habituated to a 51cm x 51cm x 39cm open field for 5 min. Mice were then

tested with two identical objects placed in the field (either two 250 ml amber bottles or two clear

plastic cylinders 6x6x16cm half filled with glass marbles). An individual animal was allowed to

explore for 5 min, now with the objects present. After two such trials (each separated by 1

minute in a holding cage), the mouse was tested in the object novelty recognition test in which a

novel object replaced one of the familiar objects (for example, an amber bottle if the cylinders

were initially used). All objects and the arena were thoroughly cleaned with 70% ethanol

between trials to remove odors. Behavior was video recorded and then scored for number

contacts (touching with nose or nose pointing at object and within 0.5 cm of object) and/or for

time contacting the objects. Habituation to the objects across the familiarization trials

(decreased contacts) was an initial measure of learning and then renewed interest (increased

contacts) in the new object indicated successful object memory. Recognition indexes were

calculated using the following formula: (# contacts during test)/(# contacts in last familiarization

trial + # contacts during test). Values greater than 0.5 indicate increased interest, whereas

values less than 0.5 indicate decreased interest in the object during the test relative to the final

familiarization trial. quadrant of the maze as well as to assess activity measures.

Quantification and statistical analysis

All data were analyzed by t-test, 1-way, 2-way or 3-way between-subject ANOVA with post hoc

comparisons depending on the number of variables and groups in each analysis. For ANOVA, if

statistical significance ($p < 0.05$) was achieved, we performed post hoc analysis to account for

multiple comparisons. The level of significance (α) was always set at 0.05. Survival curves were

analyzed using Log-rank (Mantel-Cox) Test. Data were analyzed using Prism 7 (GraphPad

Software, La Jolla, CA) and are represented as means and standard error of the means. All

experiments and data analyses were conducted in a blinded fashion. All data were prepared for

analysis with standard spread sheet software (Microsoft Excel).

1224

1225

REFERENCES

Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging: An expanding universe. *Cell* **186**, 243-278, doi:10.1016/j.cell.2022.11.001 (2023).

Mattson, M. P. & Arumugam, T. V. Hallmarks of Brain Aging: Adaptive and Pathological Modification by Metabolic States. *Cell Metab* **27**, 1176-1199, doi:10.1016/j.cmet.2018.05.011 (2018).

Ham, S. & Lee, S. V. Advances in transcriptome analysis of human brain aging. *Exp Mol Med* **52**, 1787-1797, doi:10.1038/s12276-020-00522-6 (2020).

Demontis, F., Patel, V. K., Swindell, W. R. & Perrimon, N. Intertissue control of the nucleolus via a myokine-dependent longevity pathway. *Cell Rep* **7**, 1481-1494, doi:10.1016/j.celrep.2014.05.001 (2014).

Rai, M. *et al.* Proteasome stress in skeletal muscle mounts a long-range protective response that delays retinal and brain aging. *Cell Metab* **33**, 1137-1154.e1139, doi:10.1016/j.cmet.2021.03.005 (2021).

Durieux, J., Wolff, S. & Dillin, A. The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell* **144**, 79-91, doi:10.1016/j.cell.2010.12.016 (2011).

Taylor, R. C. & Dillin, A. XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. *Cell* **153**, 1435-1447, doi:10.1016/j.cell.2013.05.042 (2013).

Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in Drosophila muscles regulates organism-wide proteostasis during aging. *Cell* **143**, 813-825, doi:10.1016/j.cell.2010.10.007 (2010).

van Oosten-Hawle, P., Porter, R. S. & Morimoto, R. I. Regulation of organismal proteostasis by transcellular chaperone signaling. *Cell* **153**, 1366-1378, doi:10.1016/j.cell.2013.05.015 (2013).

Villeda, S. A. *et al.* The ageing systemic milieu negatively regulates neurogenesis and cognitive function. *Nature* **477**, 90-94, doi:10.1038/nature10357 (2011).

Conboy, I. M. *et al.* Rejuvenation of aged progenitor cells by exposure to a young systemic environment. *Nature* **433**, 760-764, doi:10.1038/nature03260 (2005).

Castellano, J. M. *et al.* Human umbilical cord plasma proteins revitalize hippocampal function in aged mice. *Nature* **544**, 488-492, doi:10.1038/nature22067 (2017).

Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature* **583**, 596-602, doi:10.1038/s41586-020-2499-y (2020).

Villeda, S. A. *et al.* Young blood reverses age-related impairments in cognitive function and synaptic plasticity in mice. *Nat Med* **20**, 659-663, doi:10.1038/nm.3569 (2014).

De Miguel, Z. *et al.* Exercise plasma boosts memory and dampens brain inflammation via clusterin. *Nature*, doi:10.1038/s41586-021-04183-x (2021).

Horowitz, A. M. *et al.* Blood factors transfer beneficial effects of exercise on neurogenesis and cognition to the aged brain. *Science* **369**, 167-173, doi:10.1126/science.aaw2622 (2020).

Ozek, C., Krolewski, R. C., Buchanan, S. M. & Rubin, L. L. Growth Differentiation Factor 11 treatment leads to neuronal and vascular improvements in the hippocampus of aged mice. *Sci Rep* **8**, 17293, doi:10.1038/s41598-018-35716-6 (2018).

Wrann, C. D. *et al.* Exercise induces hippocampal BDNF through a PGC-1α/FNDC5 pathway. *Cell Metab* **18**, 649-659, doi:10.1016/j.cmet.2013.09.008 (2013).

Islam, M. R. *et al.* Exercise hormone irisin is a critical regulator of cognitive function. *Nature Metabolism* **3**, 1058-1070, doi:10.1038/s42255-021-00438-z (2021).

Moon, H. Y. *et al.* Running-Induced Systemic Cathepsin B Secretion Is Associated with Memory Function. *Cell Metab* **24**, 332-340, doi:10.1016/j.cmet.2016.05.025 (2016).

White, P. J. *et al.* Protectin DX alleviates insulin resistance by activating a myokine-liver glucoregulatory axis. *Nature Medicine* **20**, 664-669, doi:10.1038/nm.3549 (2014).

22   Quinn, L. S., Anderson, B. G., Strait-Bodey, L., Stroud, A. M. & Argilés, J. M. Oversecretion of
interleukin-15 from skeletal muscle reduces adiposity. *American Journal of Physiology-*
Endocrinology and Metabolism **296**, E191-E202 (2009).
23   Lourenco, M. V. *et al.* Exercise-linked FNDC5/irisin rescues synaptic plasticity and memory
defects in Alzheimer's models. *Nat Med* **25**, 165-175, doi:10.1038/s41591-018-0275-4 (2019).
24   Gupta, R., Khan, R. & Cortes, C. J. Forgot to Exercise? Exercise Derived Circulating Myokines in
Alzheimer's Disease: A Perspective. *Front Neurol* **12**, 649452, doi:10.3389/fneur.2021.649452
(2021).
25   Boyle, P. A., Buchman, A. S., Wilson, R. S., Leurgans, S. E. & Bennett, D. A. Association of muscle
strength with the risk of Alzheimer disease and the rate of cognitive decline in community-
dwelling older persons. *Arch Neurol* **66**, 1339-1344, doi:10.1001/archneurol.2009.240 (2009).
26   Kim, J. *et al.* Association of muscle and visceral adipose tissues with the probability of
Alzheimer's disease in healthy subjects. *Sci Rep* **9**, 949, doi:10.1038/s41598-018-37244-9 (2019).
27   Cortes, C. J. *et al.* Polyglutamine-expanded androgen receptor interferes with TFEB to elicit
autophagy defects in SBMA. *Nat Neurosci* **17**, 1180-1189, doi:10.1038/nn.3787 (2014).
28   Cortes, C. J. *et al.* Muscle expression of mutant androgen receptor accounts for systemic and
motor neuron disease phenotypes in spinal and bulbar muscular atrophy. *Neuron* **82**, 295-307,
doi:10.1016/j.neuron.2014.03.001 (2014).
29   Lieberman, A. P. *et al.* Peripheral androgen receptor gene suppression rescues disease in mouse
models of spinal and bulbar muscular atrophy. *Cell reports* **7**, 774-784,
doi:10.1016/j.celrep.2014.02.008 (2014).
30   Settembre, C. *et al.* TFEB links autophagy to lysosomal biogenesis. *Science* **332**, 1429-1433,
doi:10.1126/science.1204592 (2011).
31   Palmieri, M. *et al.* Characterization of the CLEAR network reveals an integrated control of
cellular clearance pathways. *Hum Mol Genet* **20**, 3852-3866, doi:10.1093/hmg/ddr306 (2011).
32   Sardiello, M. *et al.* A gene network regulating lysosomal biogenesis and function. *Science* **325**,
473-477, doi:10.1126/science.1174447 (2009).
33   Morgan, T. E. *et al.* The mosaic of brain glial hyperactivity during normal ageing and its
attenuation by food restriction. *Neuroscience* **89**, 687-699, doi:10.1016/s0306-4522(98)00334-0
(1999).
34   van Praag, H., Shubert, T., Zhao, C. & Gage, F. H. Exercise enhances learning and hippocampal
neurogenesis in aged mice. *J Neurosci* **25**, 8680-8685, doi:10.1523/jneurosci.1731-05.2005
(2005).
35   Settembre, C. *et al.* TFEB controls cellular lipid metabolism through a starvation-induced
autoregulatory loop. *Nat Cell Biol* **15**, 647-658, doi:10.1038/ncb2718 (2013).
36   Mansueto, G. *et al.* Transcription Factor EB Controls Metabolic Flexibility during Exercise. *Cell*
Metab **25**, 182-196, doi:10.1016/j.cmet.2016.11.003 (2017).
37   Leu, M. *et al.* Erbb2 regulates neuromuscular synapse formation and is essential for muscle
spindle development. *Development* **130**, 2291-2301 (2003).
38   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for
the whole mouse brain. *Nat Neurosci* **13**, 133-140, doi:10.1038/nn.2467 (2010).
39   Masiero, E. *et al.* Autophagy is required to maintain muscle mass. *Cell Metab* **10**, 507-515,
doi:10.1016/j.cmet.2009.10.008 (2009).
40   Kustermann, M. *et al.* Loss of the novel Vcp (valosin containing protein) interactor Washc4
interferes with autophagy-mediated proteostasis in striated muscle and leads to myopathy in
vivo. *Autophagy* **14**, 1911-1927, doi:10.1080/15548627.2018.1491491 (2018).
41   Giacomello, E. *et al.* Age Dependent Modification of the Metabolic Profile of the Tibialis Anterior
Muscle Fibers in C57BL/6J Mice. *Int J Mol Sci* **21**, doi:10.3390/ijms21113923 (2020).

42   Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein-protein networks, and
1322   functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res* **49**,
1323   D605-D612, doi:10.1093/nar/gkaa1074 (2021).
1324   43   Spampanato, C. *et al.* Transcription factor EB (TFEB) is a new therapeutic target for Pompe
1325   disease. *EMBO Mol Med* **5**, 691-706, doi:10.1002/emmm.201202176 (2013).
1326   44   Valentine, J. M. *et al.* NFkappaB Regulates Muscle Development and Mitochondrial Function. *J
1327   Gerontol A Biol Sci Med Sci* **75**, 647-653, doi:10.1093/gerona/gly262 (2020).
1328   45   Rai, M. & Demontis, F. Systemic Nutrient and Stress Signaling via Myokines and Myometabolites.
1329   *Annu Rev Physiol* **78**, 85-107, doi:10.1146/annurev-physiol-021115-105305 (2016).
1330   46   Mancinelli, R. *et al.* Biological Aspects of Selected Myokines in Skeletal Muscle: Focus on Aging.
1331   *Int J Mol Sci* **22**, doi:10.3390/ijms22168520 (2021).
1332   47   Cortazar, A. R., Oguiza, J. A., Aransay, A. M. & Lavin, J. L. VerSeDa: vertebrate secretome
1333   database. *Database (Oxford)* **2017**, doi:10.1093/database/baw171 (2017).
1334   48   Yoshiyama, Y. *et al.* Synapse loss and microglial activation precede tangles in a P301S tauopathy
1335   mouse model. *Neuron* **53**, 337-351, doi:10.1016/j.neuron.2007.01.010 (2007).
1336   49   Moloney, C. M., Lowe, V. J. & Murray, M. E. Visualization of neurofibrillary tangle maturity in
1337   Alzheimer's disease: A clinicopathologic perspective for biomarker research. *Alzheimers Dement*
1338   **17**, 1554-1574, doi:10.1002/alz.12321 (2021).
1339   50   Jurga, A. M., Paleczna, M. & Kuter, K. Z. Overview of General and Discriminating Markers of
1340   Differential Microglia Phenotypes. *Front Cell Neurosci* **14**, 198, doi:10.3389/fncel.2020.00198
1341   (2020).
1342   51   Rozovsky, I., Finch, C. E. & Morgan, T. E. Age-related activation of microglia and astrocytes: in
1343   vitro studies show persistent phenotypes of aging, increased proliferation, and resistance to
1344   down-regulation. *Neurobiol Aging* **19**, 97-103, doi:10.1016/s0197-4580(97)00169-3 (1998).
1345   52   Godbout, J. P. *et al.* Exaggerated neuroinflammation and sickness behavior in aged mice
1346   following activation of the peripheral innate immune system. *FASEB J* **19**, 1329-1331,
1347   doi:10.1096/fj.05-3776fje (2005).
1348   53   Sidoryk-Wegrzynowicz, M. *et al.* Astrocytes in mouse models of tauopathies acquire early
1349   deficits and lose neurosupportive functions. *Acta Neuropathol Commun* **5**, 89,
1350   doi:10.1186/s40478-017-0478-9 (2017).
1351   54   Preuss, C. *et al.* A novel systems biology approach to evaluate mouse models of late-onset
1352   Alzheimer's disease. *Mol Neurodegener* **15**, 67, doi:10.1186/s13024-020-00412-5 (2020).
1353   55   Sherva, R. *et al.* Genome-wide association study of the rate of cognitive decline in Alzheimer's
1354   disease. *Alzheimers Dement* **10**, 45-52, doi:10.1016/j.jalz.2013.01.008 (2014).
1355   56   Simon, M. J. *et al.* Transcriptional network analysis of human astrocytic endfoot genes reveals
1356   region-specific associations with dementia status and tau pathology. *Sci Rep* **8**, 12389,
1357   doi:10.1038/s41598-018-30779-x (2018).
1358   57   Knopman, D. S. *et al.* Alzheimer disease. *Nat Rev Dis Primers* **7**, 33, doi:10.1038/s41572-021-
1359   00269-y (2021).
1360   58   Guyenet, S. J. *et al.* A simple composite phenotype scoring system for evaluating mouse models
1361   of cerebellar ataxia. *J Vis Exp*, doi:10.3791/1787 (2010).
1362   59   Lehallier, B. *et al.* Undulating changes in human plasma proteome profiles across the lifespan.
1363   *Nature Medicine* **25**, 1843-1850, doi:10.1038/s41591-019-0673-2 (2019).
1364   60   Jiao, J. & Demontis, F. Skeletal muscle autophagy and its role in sarcopenia and organismal
1365   aging. *Curr Opin Pharmacol* **34**, 1-6, doi:10.1016/j.coph.2017.03.009 (2017).
1366   61   Wohlgemuth, S. E., Seo, A. Y., Marzetti, E., Lees, H. A. & Leeuwenburgh, C. Skeletal muscle
1367   autophagy and apoptosis during aging: effects of calorie restriction and life-long exercise.
1368   *Experimental gerontology* **45**, 138-148, doi:10.1016/j.exger.2009.11.002 (2010).

62 Yu, D. *et al.* The adverse metabolic effects of branched-chain amino acids are mediated by
isoleucine and valine. *Cell Metab* **33**, 905-922 e906, doi:10.1016/j.cmet.2021.03.025 (2021).
63 Adler, D. H. *et al.* Characterizing the human hippocampus in aging and Alzheimer's disease using
a computational atlas derived from ex vivo MRI and histology. *Proc Natl Acad Sci U S A* **115**,
4252-4257, doi:10.1073/pnas.1801093115 (2018).
64 Ivanisevic, J. *et al.* Metabolic drift in the aging brain. *Aging (Albany NY)* **8**, 1000-1020,
doi:10.18632/aging.100961 (2016).
65 Ximerakis, M. *et al.* Single-cell transcriptomic profiling of the aging mouse brain. *Nat Neurosci*
22, 1696-1708, doi:10.1038/s41593-019-0491-3 (2019).
66 Zarate, S., Stevnsner, T. & Gredilla, R. Role of Estrogen and Other Sex Hormones in Brain Aging.
Neuroprotection and DNA Repair. *Front Aging Neurosci* **9**, 430, doi:10.3389/fnagi.2017.00430
(2017).
67 Hahn, O. *et al.* Atlas of the aging mouse brain reveals white matter as vulnerable foci. *Cell*,
doi:10.1016/j.cell.2023.07.027 (2023).
68 Cortes, C. J. & De Miguel, Z. Precision Exercise Medicine: Sex Specific Differences in Immune and
CNS Responses to Physical Activity. *Brain Plasticity*, pp. 1-13, doi:10.3233/BPL-220139 (2022).
69 Vizovisek, M., Fonovic, M. & Turk, B. Cysteine cathepsins in extracellular matrix remodeling:
Extracellular matrix degradation and beyond. *Matrix Biol* **75-76**, 141-159,
doi:10.1016/j.matbio.2018.01.024 (2019).
70 Tran, A. P., Sundar, S., Yu, M., Lang, B. T. & Silver, J. Modulation of Receptor Protein Tyrosine
Phosphatase Sigma Increases Chondroitin Sulfate Proteoglycan Degradation through Cathepsin
B Secretion to Enhance Axon Outgrowth. *J Neurosci* **38**, 5399-5414,
doi:10.1523/JNEUROSCI.3214-17.2018 (2018).
71 Saini, M. G. & Bix, G. J. Oxygen-glucose deprivation (OGD) and interleukin-1 (IL-1) differentially
modulate cathepsin B/L mediated generation of neuroprotective perlecan LG3 by neurons. *Brain*
Res **1438**, 65-74, doi:10.1016/j.brainres.2011.12.027 (2012).
72 Muigg, P. *et al.* Differential stress-induced neuronal activation patterns in mouse lines
selectively bred for high, normal or low anxiety. *PLoS One* **4**, e5346,
doi:10.1371/journal.pone.0005346 (2009).
73 Czibere, L. *et al.* Profiling trait anxiety: transcriptome analysis reveals cathepsin B (Ctsb) as a
novel candidate gene for emotionality in mice. *PLoS One* **6**, e23604,
doi:10.1371/journal.pone.0023604 (2011).
74 New, J. & Thomas, S. M. Autophagy-dependent secretion: mechanism, factors secreted, and
disease implications. *Autophagy* **15**, 1682-1693, doi:10.1080/15548627.2019.1596479 (2019).
75 Chow, L. S. *et al.* Exerkines in health, resilience and disease. *Nat Rev Endocrinol* **18**, 273-289,
doi:10.1038/s41574-022-00641-2 (2022).
76 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**,
676-682, doi:10.1038/nmeth.2019 (2012).
77 Encarnacion-Rivera, L., Foltz, S., Hartzell, H. C. & Choo, H. Myosoft: An automated muscle
histology analysis tool using machine learning algorithm utilizing FIJI/ImageJ software. *PLoS One*
15, e0229041, doi:10.1371/journal.pone.0229041 (2020).
78 Herrera, J. J. *et al.* Acarbose has sex-dependent and -independent effects on age-related
physical function, cardiac health, and lipid biology. *JCI Insight* **5**, doi:10.1172/jci.insight.137474
(2020).
79 Amodei, D. *et al.* Improving Precursor Selectivity in Data-Independent Acquisition Using
Overlapping Windows. *J Am Soc Mass Spectrom* **30**, 669-684, doi:10.1007/s13361-018-2122-8
(2019).

80   Adusumilli, R. & Mallick, P. Data Conversion with ProteoWizard msConvert. *Methods Mol Biol*
1550, 339-368, doi:10.1007/978-1-4939-6747-6_23 (2017).

81   Searle, B. C. *et al.* Chromatogram libraries improve peptide detection and quantification by data
independent acquisition mass spectrometry. *Nat Commun* **9**, 5128, doi:10.1038/s41467-018-
07454-w (2018).

82   Wu, C. H. *et al.* The Universal Protein Resource (UniProt): an expanding universe of protein
information. *Nucleic Acids Res* **34**, D187-191, doi:10.1093/nar/gkj161 (2006).

83   Kall, L., Canterbury, J. D., Weston, J., Noble, W. S. & MacCoss, M. J. Semi-supervised learning for
peptide identification from shotgun proteomics datasets. *Nat Methods* **4**, 923-925,
doi:10.1038/nmeth1113 (2007).

84   MacLean, B. *et al.* Skyline: an open source document editor for creating and analyzing targeted
proteomics experiments. *Bioinformatics* **26**, 966-968, doi:10.1093/bioinformatics/btq054
(2010).

85   Sharma, V. *et al.* Panorama Public: A Public Repository for Quantitative Data Sets Processed in
Skyline. *Mol Cell Proteomics* **17**, 1239-1244, doi:10.1074/mcp.RA117.000543 (2018).

86   Pesta, D. *et al.* Similar qualitative and quantitative changes of mitochondrial respiration
following strength and endurance training in normoxia and hypoxia in sedentary humans. *Am J*
Physiol Regul Integr Comp Physiol **301**, R1078-1087, doi:10.1152/ajpregu.00285.2011 (2011).

87   Tomfohr, J., Lu, J. & Kepler, T. B. Pathway level analysis of gene expression using singular value
decomposition. *BMC Bioinformatics* **6**, 225, doi:10.1186/1471-2105-6-225 (2005).

1436

Figure 1

Figure 2

Figure 3

A.

Predicted Secreted
Differentially Expressed Proteins
in cTFEB;HSACre Skeletal Muscle

Age: 6 Months

Up-regulated
- Nucleobindin-1
- Thrombospondin 1
- Myocilin
- Angiopoietin-related protein 1

Down-Regulated
- Collagen alpha-1 (I) chain

Up-regulated
- **cathepsin B**
- Prosaposin
- MERP1
- D/R SDR 7C

Up-regulated
none

Down-Regulated
- Biglycan (PG-S1)

Age: 18 Months

B.

Females   Males
Young   Aged   Young   Aged

Group ID: C C T T C T C T C C T T T T C T C C C C T T T T   kDa

Pro-CTSB ➤   – 40
Mature ➤   – 35
CTSB ➤   – 25

Total Protein
(StainFree)

Legend:
◆ Controls
◆ cTFEB;HSACre
◇ Controls
◇ cTFEB;HSACre

Pro-CTSB Levels
(norm. to StainFree Lane)

Males: Young, Aged   Females: Young, Aged

Mature CTSB Levels
(norm. to StainFree Lane)

Males: Young, Aged   Females: Young, Aged

C.

Legend:
◆ Controls
◆ cTFEB;HSACre
◇ Controls
◇ cTFEB;HSACre

CTSB Serum
Levels (pg/µL)

Males: Young, Aged   Females: Young, Aged

Figure 4

Figure 5

Figure 6

A.

B.

C.

D.

E.

F.

Figure 7

A. Males
Age : 6 months

B. Males
Age : 24 months

C. GO Term Description

D. GO Term Description

E. KEGG Pathway

F. KEGG Pathway