Skeletal Muscle Proteostasis Promotes Central Nervous System Function and Reduces Neuroinflammation during Aging and Neurodegenerative Disease

Ian Matthews [1,2]

Allison Birnbaum [1,2]

Anastasia Gromova [3]

Kristen Coutinho [5]

Megan McGraw [6]

Dalton C. Patterson [6]

Macy T. Banks [6]

Amber Nobles [7]

Nhat Nguyen [3]

Helen C. Miranda [8-10]

Albert R. La Spada [3,4] (co-corresponding author)

Constanza J. Cortes [1,2] (corresponding author)

[1] Leonard Davis School of Gerontology, [2] University of Southern California, [3] Department of Pathology and Laboratory Medicine, University of California, Irvine, [4] Department of Neurology and Department of Biological Chemistry, UCI Institute for Neurotherapeutics, University of California, Irvine, [5] Comprehensive Diabetes Center, University of Alabama at Birmingham, [6] Department of Cell, Developmental and Integrative Biology, University of Alabama at Birmingham, [7] Department of Neurology, University of Alabama at Birmingham, [8] Department of Genetics and Genome Sciences, School of Medicine, Case Western University, [9] Department of Neurosciences, School of Medicine, Case Western Reserve University, [10] RNA Center, School of Medicine, Case Western Reserve University,

Funding: NIH R01 AG077536 (CJC), University of Washington Nathan Shock Center Pilot

Grant (CJC), NIH R35 122140 (ALS).

Acknowledgements: We thank members of the lab past and present, as well as former and

current colleagues and collaborators for their helpful contributions.

Authorship Attributions: IM, CJC, AG, HM, AB, KC, MM, DP, MB, AN, and NN: Performed the

experimental work. Reviewed and edited the manuscript. CJC and IM: Wrote and edited the

manuscript. CJC and ALS: Conceptualized the experiments outlined here.

ABSTRACT

Skeletal muscle has recently arisen as a novel regulator of Central Nervous System (CNS)

function and aging, secreting bioactive molecules known as myokines with proteostasis and

metabolism-modifying functions in targeted tissues, including the CNS. Here we report the

generation of a novel transgenic mouse with enhanced skeletal muscle proteostasis via targeted

overexpression of Transcription Factor E-B, a master regulator of proteostasis and lysosomal

function. We have discovered that the resulting enhanced muscle proteostasis function

significantly ameliorates proteotoxicity, reduces neuroinflammation and promotes transcriptional

remodeling of the aging CNS, preserving cognition and memory in aging mice. Enhancing

skeletal muscle proteostasis also reduces neuroinflammation and accumulation of tau-

associated pathological hallmarks in a mouse model of tau pathology. Our results implicate the

maintenance of skeletal muscle proteostasis throughout aging to direct regulation of the CNS

metabolism and function, and suggest that skeletal-muscle originating factors may act as novel

therapeutic targets against age-associated neurodegenerative diseases.

INTRODUCTION

Aging is associated with an organism-wide progressive loss of tissue form and function, broadly characterized by the 'hallmarks of aging' [1]. In particular, the aging central nervous system (CNS) [1,2] exhibits a global loss in protein homeostasis (proteostasis), impaired neuroplasticity and resilience, and an increase in neuroinflammation [2,3]. These alterations are believed to render the aging CNS vulnerable to age-associated dysfunction and the development of neurodegenerative disease [2].

Over the last ten years, there has been growing evidence that suggests prominent contributions of the periphery to the etiology of age-associated neurodegenerative diseases [4-8]. In particular, manipulation of skeletal muscle protein quality control pathways yields important benefits to the invertebrate CNS by activating proteostasis and protecting against the accumulation of aggregation-prone neurodegenerative disease proteins in the brain and the retina [5,8,9]. Although the mechanisms responsible for these benefits remain poorly understood, some of these effects are mediated by secreted factors that communicate important metabolic, proteostatic and inflammatory changes between tissues [5,7,9]. Indeed, blood circulating factors have recently arisen as potent regulators of mammalian CNS aging and metabolism [10-13]. For example, exposure to a young plasma environment (either through heterochronic parabiosis or plasma transfers) can rescue function in the aging CNS [12,14] by decreasing neuroinflammation [15] and enhancing neurogenesis [10]. Studies have further demonstrated that increasing the levels of these individual factors in peripheral circulation is sufficient to rejuvenate the aged CNS [12,16,17], supporting the existence of geroprotective circulating factors with CNS targeting effects. Although the source and identity of these neuroprotective circulating cytokines are unclear, several of them are expressed in, and are known to be secreted from, skeletal muscle [5,18-20]. In fact, skeletal muscle acts as an endocrine organ secreting a myriad of bioactive factors that induce metabolic changes in a cell non-autonomous manners in distant tissues like liver [21], adipose tissue [22], and even the CNS [19].

Recently, this muscle-to-brain signaling axis has been proposed to have important implications for CNS aging and age-associated neurodegenerative disease [19,23,24]. Skeletal muscle function has arisen as a key predictor for phenotypic and clinical outcomes in age-associated neurodegenerative diseases, including Alzheimer's Disease (AD) and Parkinson's Disease (PD) [25,26]. Consistent with this, we were the first to uncover a novel mechanistic basis for motor neuron disease in the polyglutamine disease Spinal and Bulbar Muscular Atrophy (SBMA),

where disruption of skeletal muscle autophagy [27] initiates the pathogenic cascade that

culminates in neuronal toxicity and death [28,29]. Furthermore, we demonstrated that rescuing

proteotoxicity specifically in skeletal muscle was sufficient to rescue motor neuron degeneration

and extend lifespan in this model, despite robust misfolded protein burden in the brain [29].

Importantly, we also demonstrated the feasibility and efficacy of peripheral delivery of antisense

oligonucleotides targeting skeletal muscle expression of the misfolded protein in rescuing

neurological phenotypes [30], confirming that therapeutic reduction of misfolded protein

expression in skeletal muscle can have important benefits to the CNS.

A central player in proteostasis is macroautophagy (hereafter referred to as autophagy), which

degrades and recycles intracellular material such as mitochondria, protein aggregates, and

other large subcellular components too large to be processed through the proteasome. As with

most other aging tissues, autophagy activity steadily declines with age in skeletal muscle [31],

leading to a chronic accumulation of damaged macromolecules and organelles [31-33] and

contributing to the onset of age-associated decline [1,31,34-37]. In accordance with this, genetic

disruption of autophagy is sufficient to drive cellular and biochemical hallmarks of aging in

young skeletal muscle, including accumulation of protein aggregates [34,36-39], mitochondrial

dysfunction [40-43] and compromised metabolic signaling [40,44,45]. Autophagy dysfunction has thus

been proposed to act as a primary regulator of biological aging for skeletal muscle [1,31].

Interestingly, well-known interventions that maintain or re-activate global and skeletal muscle

autophagy via genetic [46-48], behavioral [49] and pharmacological [50,51] manipulation extend lifespan

and/or markers of cognitive health span in murine and primate models. Altogether, this suggests

that maintenance of skeletal muscle proteostasis may promote neurotrophic signaling in the

brain, providing systemic benefits against disease and age-associated CNS decline.

Skeletal muscle autophagy is regulated in part by Transcription Factor E-B (TFEB), a master

regulator of a lysosomal-to-nucleus signaling axis that integrates cellular metabolism and

autophagy [52-54]. TFEB expression and function are strongly induced in skeletal muscle in

response to interventions with documented neuroprotective effects on the aging [55,56] and

neurodegenerative disease-afflicted CNS, including low nutrient conditions [57] and exercise [40].

However, the precise role of TFEB-dependent signaling in skeletal muscle during aging, and its

impact on the muscle-to-brain axis remains largely unknown. Here we report the generation of a

novel transgenic mouse with enhanced muscle proteostasis via overexpression of TFEB. We

have discovered that the resulting enhanced skeletal muscle proteostasis function is maintained

throughout the lifespan, protecting against the onset age-associated phenotypes in aging

skeletal muscle. Furthermore, we report that enhancing skeletal muscle proteostasis

significantly reduces hippocampal accumulation of neuropathological hallmarks and reduces

neuroinflammation in a mouse model of tau pathology, despite no exogenous activation of our

transgene in the CNS. Enhancing skeletal muscle proteostasis via TFEB overexpression across

the lifespan also ameliorates proteotoxicity, reduces neuroinflammation and promotes

transcriptional remodeling of the aging CNS without neurodegenerative disease burden,

preserving cognitive performance in aging mice. Our results implicate, for the first time in

mammals, maintenance of skeletal muscle proteostasis to direct regulation of CNS health and

function, and suggest that skeletal-muscle originating factors may act as novel therapeutic

targets against age-associated neurodegenerative diseases.

RESULTS

TFEB overexpression preserves proteostasis-related and metabolism-associated
networks in aging skeletal muscle

To allow for tissue-specific expression of TFEB under controlled conditions, we took advantage

of the Cre-loxP system for mammalian gene editing. We derived a novel line of conditional

transgenic mice carrying the β-actin promoter in combination with the CMV enhancer (CAGGS)

and with a floxed 3x-FLAG-eGFP STOP cassette placed just 5' to a 3x-FLAG-human *TFEB*

transgene (i.e. fxSTOP-TFEB transgenic mice) (**Figure 1A**). In the presence of Cre-

recombinase, the STOP codon cassette will be excised out, allowing expression of the human

TFEB transgene, also known as a Cre-loxP ON system. To drive overexpression of TFEB

specifically in skeletal muscle, we crossed male fxSTOP-TFEB mice with female Human

Skeletal Actin (HSA)-Cre mice to achieve widespread expression of Cre-recombinase only in

cells of the myogenic lineage [58].

The resulting cTFEB;HSACre bigenic mice of both sexes exhibited a strong induction of human

TFEB expression in skeletal muscle tissue (**Supplemental Figure 1A, left**). This exogenous

human *TFEB* expression was still quite prominent at 24 months of age in cTFEB;HSACre

transgenic skeletal muscle (**Supplemental Figure 1B, left)** suggesting that overexpression of

the transgene remains active even at very old ages. Immunoblotting of quadricep skeletal

muscle confirmed robust expression of 3x-FLAG-TFEB protein in cTFEB;HSACre bigenic

muscle lysates (**Figure 1B**). As TFEB is a transcriptional factor, we confirmed nuclear

localization of human TFEB in cTFEB;HSA-Cre skeletal muscle myonuclei via

immunofluorescence (**Supplemental Figure 1C, in white**).

Next, we confirmed TFEB overexpression exclusively in skeletal muscle of cTFEB;HSACre

bigenic mice by multiple approaches. First, we confirmed expression of the biologically inert 3x-

FLAG-eGFP cassette upstream of the STOP codon (**Figure 1A**) and no exogenous expression

of 3x-FLAG-TFEB in other highly metabolic tissues of non-myogenic lineages in cTFEB;HSACre

transgenic mice, including brown adipose tissue (**Supplemental Figure 1D**) and liver

(**Supplemental Figure 1E**). Consistent with these findings, we also did not detect expression of

exogenous 3x-FLAG-TFEB in hemibrain protein lysates (**Supplemental Figure 1F**). Indeed, at

all ages examined, we detected no differences in *TFEB* transcription in hippocampal RNA

lysates regardless of their genotype or age (**Supplemental Figure 1A & B, right**). To directly

assess the Cre-mediated recombination of an additional transgene reporter, we generated

TdTomato Ai9;HSACre mice and confirmed robust tdTomato fluorescence in skeletal muscle of

bigenic individuals (**Supplemental Figure 1G**). Importantly, and consistent with our results in

our cTFEB;HSACre model, there was no detectable TdTomato fluorescence in the CNS

(**Supplemental Figure 1H**), including the hippocampus (**Supplemental Figure 1H, insets**) of

TdTomato Ai9;HSACre mice, consistent with the muscle-restricted specificity of this HSA-Cre-

loxP-ON approach [59]. Overall, these data confirm previous reports of striated muscle-specific

Cre-mediated recombination using the HSA-Cre transgenic driver line [58], and demonstrate

expression of exogenous *TFEB* only in skeletal muscle of cTFEB;HSACre bigenic mice. This

establishes our newly developed inducible fxSTOP-TFEB mouse as an effective model to

generate spatiotemporally controlled transgenic expression of human *TFEB*.

TFEB directly regulates transcription of multiple lysosomal and autophagy-associated genes

[52,54], promoting lysosomal function and autophagy activation, as well as mitochondrial

biogenesis and lipid metabolism in vivo [40,57] and in vitro [52,54]. We confirmed functional activation

of TFEB-dependent transcription in cTFEB:HSA-Cre bigenic muscle, using our previously

reported TFEB-responsive muscle gene targets [27] including *Lamp1* (an essential lysosomal

structural protein), *Atg5* (a key regulator of autophagy initiation), *Ctsd* (Cathepsin D) and *Ctsb*

(Cathepsin B, both lysosomal proteases) (**Figure 1C-F**). Immunoblotting studies confirmed

significant increases of TFEB transcriptional targets LAMP1 and CTSD at the protein level in

cTFEB;HSACre bigenic muscle (**Figure 1G**).Consistent with previous studies demonstrating

TFEB overexpression increases the number and size of LAMP1-positive lysosomes in multiple

tissues [52,57,60], we also detected significant increases in the number of LAMP1+ puncta and total

LAMP1+ area in cross-sections of female (and similar trends in male) cTFEB;HSACre skeletal

muscle (**Figure 1H**). These findings confirm physiologically significant increases in TFEB-

dependent transcription [40] and lysosomal biogenesis [40,54] in skeletal muscle of cTFEB:HSA-Cre

mice.

Given the principal role of autophagy function to skeletal muscle health [35,45], and the predicted

contributions of age-associated autophagy dysfunction to muscle wasting and sarcopenia during

aging [61], we sought to examine the overall changes in the skeletal muscle protein landscape in

response to TFEB overexpression throughout aging. We performed proteome profiling of young

and aged, control and cTFEB;HSACre male quadricep skeletal muscle. Differential abundance

analysis of species revealed around 1000 proteins associated with TFEB overexpression in

young (6-month-old) skeletal muscle (**Figure 1I**). Of these, 650 (65%) were significantly

overrepresented and 350 (35%) were underrepresented in TFEB-overexpressing muscle

compared to their littermate controls. At 18 months of age, a time point where age-associated

proteostasis dysfunction is prevalent in murine skeletal muscle [62], the same proteomics

approach identified around 400 differentially expressed proteins in TFEB-expressing skeletal

muscle (**Figure 1J**). Of these, 250 proteins (83%) were significantly enriched and 50 proteins

(17%) were significantly decreased in cTFEB;HSACre bigenic muscle. Importantly, at both ages

examined, TFEB remained significantly overexpressed (**Figure 1I-J, boxes**), confirming our

previous finding of maintenance of exogenous skeletal muscle TFEB expression across the

lifespan of our cTFEB;HSACre model (**Supplemental Figure 1A-B**).

To visualize the key networks of differentially expressed proteins associated with TFEB-

overexpression, we performed a functional enrichment analysis on all overexpressed proteins

using the STRING protein-protein interaction network [63] (**Figure 1K**). Using TFEB as our 'bait'

for functional connectivity nodes, we identified two clear interaction networks: one associated

with lysosomal biogenesis and function (**Figure 1K, blue**) and a second one associated with

key skeletal muscle proteostasis and metabolic signaling pathways including mTOR, AKT and

GSK3β (**Figure 1K, green**). Both of these nodes are key elements of the CLEAR signaling

network controlled by TFEB-mediated transcription [53,54], were previously reported to be

transcriptionally elevated in skeletal muscle after viral-mediated TFEB overexpression [40], and

are central regulators of aging [1].

To further understand the functions of all the upregulated proteins in TFEB-overexpressing

skeletal muscle, KEGG enrichment analysis was performed. As expected, central proteostasis

categories were significantly upregulated in TFEB-overexpressing skeletal muscle: including

Hypoxia-inducible factor 1 (HIF-1) signaling, proteasome, and autophagy/lysosomal function

categories (**Figure 1L, orange**), consistent with our targeted immunoblot studies (**Figure 1G**)

and our STRING analysis (**Figure 1K**). Importantly, this enrichment in proteostasis categories

and TFEB-regulated nodes of signaling was still present in our aging cohorts (**Figure 1L, top**),

suggesting that overexpression of TFEB throughout the lifespan preserves proteostasis function

in aging skeletal muscle.

TFEB overexpression has been reported to reactivate declining proteostasis associated with

age- or disease-associated impaired lysosomal function [27,60,64]. We wondered if the enrichment

in proteostasis pathways observed in aged skeletal muscle with TFEB-overexpression would

result in functional outcomes reflective of proteostasis maintenance. Analysis of autophagy

marker LC3, which is cleaved and post-translationally modified into its mature form (LC3-II)

upon autophagy induction [65], revealed a significant increase in LC3-II:LC3-I ratio in quadricep

muscle of 12 month-old cTFEB:HSA-Cre mice (**Figure 1M**). This suggests that in addition to

activating lysosomal biology and function, expression of TFEB in skeletal muscle results in

functional activation of autophagy in vivo, as reported in other TFEB-overexpression studies in

skeletal muscle, liver, and brain [57,60,64].

Interestingly, additional functional categories enriched in our proteomics analysis related to

cellular metabolism and mitochondrial function, including thermogenesis, oxidative

phosphorylation and pyruvate metabolism (**Figure 1L, gold**). We also detected central fatty acid

metabolism pathways, including PPAR signaling and metabolism, and degradation and

biosynthesis of fatty acids (**Figure 1L, green**). This is consistent with previously published

reports suggesting that exogenous TFEB expression in young skeletal muscle induces the

expression of genes involved in mitochondrial biogenesis, fatty acid oxidation, and oxidative

phosphorylation [40]. Here we confirm for the first time that these changes in cTFEB;HSACre

muscle are maintained throughout aging, particularly those associated with mitochondrial

metabolism and fatty acid metabolism (**Figure 1L, top, gold and green**).

KEGG analysis also showed enrichment in cTFEB;HSACre bigenic muscle for classical

signaling pathways shown to decline during aging, including members of the Insulin-IGF1-

signaling (IIS) and AMPK signaling [1] (**Figure 1L, blue**). Finally, we also detected multiple hits

for amino acid metabolism pathways, such as catabolism of branched-chain amino acids

(BCAAs), including valine, leucine and isoleucine metabolism (**Figure 1L, purple**). Importantly,

these changes were still evident in our aged cohorts (**Figure 1L, top**), suggesting a potential

geroprotective effect of TFEB-associated signaling in aging skeletal muscle. In agreement with

this, analysis of skeletal muscle of 12-month-old control mice showed marked accumulation of

age-associated histopathological markers, including fiber decompaction, intrafibrillary myonuclei

and multilamellar structure accumulation (**Figure 1N, top**). Remarkably, these features were

mostly absent in skeletal muscle sections from age- and littermate-matched cTFEB:HSA-Cre

mice (**Figure 1N, bottom**). Instead, aging cTFEB;HSACre muscle retained compact fiber

organization and perifibrillar myonuclei and did not exhibit any multilammelar structures. Indeed,

the appearance of the aging cTFEB;HSACre skeletal muscle was much more consistent with

their younger counterparts. In agreement with this, we also detected reduced expression of

muscle stress factor *Mstn* (myostatin) [66] (**Figure 1O**) and increased expression of metabolic

modulator *Fndc5* [66] in aged cTFEB;HSACre skeletal muscle (**Figure 1P**). Altogether, our results

suggest maintenance of functional proteostasis and mitochondrial function via TFEB

overexpression throughout aging has multiple benefits in skeletal muscle, targeting established

hallmarks of aging.

Enhanced skeletal muscle proteostasis reduces accumulation of hyperphosphorylated

tau and microglial activation in a mouse model of tau pathology.

To assess the potential neuroprotective effects of enhancing skeletal muscle proteostasis in the

context of age-associated neurodegenerative disease pathologies, we derived cTFEB;HSACre

transgenic mice in the MAPT P301S (PS19) background. This is a well-known model of

neurofibrillary tangle toxicity, a hallmark of AD and related tauopathies [67], and is characterized

by prominent accumulation of hippocampal hyperphosphorylated human mutant MAPT protein

and pronounced neuroinflammation, including microgliosis and astrocyte reactivity [67].

We first confirmed muscle-restricted 3x-FLAG-TFEB overexpression in skeletal muscle lysates

from MAPT P301S;cTFEB;HSACre mice, as well as the presence of human Tau in hippocampal

lysates of both groups via immunoblotting (**Figure 2A**). As previously reported [67], at 9 months of

age there is robust accumulation of Ser202/Thr205 hyperphosphorylated tau (residues

abnormally phosphorylated in paired helical filaments during AD progression) in the dentate

gyrus of the hippocampus of MAPT P301S mice of both sexes [67] (**Figure 2B, middle panel**).

Conversely, we noted a significant reduction in the total fluorescence counts of AT8-positive

phosphorylated tau in the dentate gyrus of MAPT P301S;cTFEB;HSACre mice (**Figure 2B**,

purple) compared with MAPT P301S littermate controls (**Figure 2B**, **teal**). More so, this

reduction was particularly evident in the levels of 'intracellular tau' accumulating around

hippocampal cell bodies (**Figure 2B, insets**), suggesting reduced accumulation of toxic tau

species and aggregates [68] in the hippocampus of MAPT P301S;cTFEB;HSACre mice. We

confirmed similar but less striking reductions in Thr231 phospho-tau (another primary site of tau

phosphorylation associated with neurotoxic tau accumulation) immunostaining (**Supplemental**

Figure 2A), suggesting overall reductions in tau hyperphosphorylation in the hippocampus of

MAPT P301S transgenic mice with TFEB overexpression in skeletal muscle.

Astrocytes and microglia are key mediators of neuroinflammation, and undergo key

morphological and functional alterations during neurodegenerative disease, including changes

in ramification and cellular process complexity [69], reduced immune surveillance activity [70] and

increased reactivity phenotypes [71]. We confirmed significant increases in the staining for GFAP-

positive structures (a marker of astrocyte reactivity) in the dentate gyrus of 9-month-old MAPT

P301S mice of both sexes (**Figure 2C, teal**) compared to their MAPT-negative littermate

controls (**Figure 2C, gold**). We also detected significant increases in the volume of IBA1+-

positive microglia in the same brain region (**Figure 2C, teal**), consistent with previous reports in

the MAPT P301S model indicating high levels of neuroinflammation starting as early as 4

months of age [67,72]. Strikingly, we noted significant reductions in both of these pro-inflammatory

glial morphometric parameters in the hippocampus of MAPT P301S;cTFEB;HSACre littermates

(**Figure 2C, purple**), suggesting overall reduced hippocampal neuroinflammation during

symptomatic disease stages through activation of skeletal muscle proteostasis.

Enhancing skeletal muscle proteostasis promotes neurotrophic signaling and reduces

hippocampal neuroinflammation in MAPT P301S transgenic mice.

To gain more precise insights into disease-relevant transcriptional changes in the hippocampus

of MAPT P301S mice with enhanced skeletal muscle proteostasis, we used the Nanostring

nCounter® Alzheimer's Disease panel. This platform directly assesses the expression of 770

genes curated from human and pre-clinical models of AD and robustly captures disease-

relevant signatures and their modifications after pre-clinical interventions [73]. Using this

approach, we found a total of 79 differentially expressed genes in MAPT P301S;cTFEB;HSACre

hippocampi compared to their littermate MAPT P301S+ controls (**Figure 2D-E**). Interestingly,

most differentially regulated genes were downregulated (76 of 79), consistent with their identity

as transcriptional drivers of disease [73]. Many of these transcripts are associated with microglial

activation (including *Rhoc, Bcar3, and Alcam*), markers of neuronal function (*Gdap1l1),*

phospholipid remodeling (*Ano6*), growth factor signaling (*Vgf*), endothelial cell migration and

matrix adhesion (*Rras*) and guanine nucleotide exchange factors (*Dock3*). Interestingly, multiple

down-regulated genes were associated with novel variants and loci associated with increased

polygenic neurodegenerative disease risk (*Prkd3* [74], *Bcar3* [75], *Nfia* [76], and *Elovl7* [77]) as well as

predictors of the rate of cognitive decline in AD (*Hecw1* [78]). Interestingly, expression of *Arhgdib,*

recently identified as a potential novel biomarker for tau neurofibrillary tangle accumulation and

neuropathological Braak stage in human entorhinal cortex [79], was also significantly reduced,

consistent with our reduced hyperphosphorylated tau staining in the hippocampus of MAPT

P301S ;cTFEB;HSACre transgenic mice (**Figure 2B**). *Pbxip1,* another significantly

downregulated marker, has been found to be associated with phosphorylated-tau and $A\beta_{1-40}$

levels in the human temporal cortex [80]. Additional transcriptional reductions of significance

include genes of unknown function but classified as transcriptional identifiers of AD progression

(Ptprn, Cpne2, and *Eri2)* [73]. Functional annotation of the differentially expressed co-

expression/functional clusters revealed significant reductions in the nCounter AD microglial

activation module (composed of 63 genes associated with microglial activation during AD

progression) (**Figure 2F**) in the hippocampi of 9 month old female (but not male) MAPT

P301S;cTFEB;HSACre mice compared with MAPT P301S littermates. This is consistent with

increased neuroprotection and reduced neuroinflammation in the hippocampus of MAPT P301S

mice with enhanced skeletal muscle proteostasis, as suggested by our glail morphometric

analysis (**Figure 2C**). Interestingly, we also detected a sex-bias in the trophic factor functional

cluster (composed of genes *Arhgdib, Bcl2, Calm3, Camk4, Gab1, Mapk9, Mapkapk2, Ntrk2,*

Pik3r1 and *Psen1*), with a significant improvement back to baseline expression in the

hippocampus of male (but not female) MAPT P301S;cTFEB;HSACre mice (**Figure 2G**). We

confirmed activation of neurotrophic signaling as early as 6 months of age in the hippocampus

of male MAPT P301S;cTFEB;HSACre mice by targeted qRT-PCR analysis, which showed

corrections of declining expression levels of neurotrophic factors *Bdnf* and *Fndc5* back to control

littermate levels (**Figure 2H-I**) [23,81]. Importantly, there was also a significant extension (38%) in

the lifespan of male MAPT P301S;cTFEB;HSACre mice (**Supplemental Figure 2D**). While the

median survival of male MAPT P301S mice was 352 days (**Supplemental Figure 2D, teal**

curve), male MAPT P301S;cTFEB;HSACre mice median survival was 489 days (**Supplemental**

Figure 2D, purple curve). Although we did not detect a similar lifespan extension in female

MAPT P301S;cTFEB;HSACre mice (**Supplemental Figure 2E**), it is worth noting that in our

hands, female MAPT P301S transgenic mice already live significantly longer than their male

littermate controls (median survival of 468.5 days), perhaps suggesting they have reached

maximal lifespan in this model. Finally, we did not detect any differences in ataxia progression [82]

(**Supplemental Figure 2B**) or weight loss (**Supplemental Figure 2C**) between our groups.

Altogether, these provocative results suggest that enhanced skeletal muscle proteostasis

modifies the accumulation of pathogenic tau isoforms and reduces neuroinflammation in the

hippocampus of MAPT P301S mice in a sex-dependent manner via transcriptional remodeling

and activation of neurotrophic signaling. Importantly, this occurs in the absence of detectable

exogenous human TFEB expression in the CNS (**Supplemental Figure 1**), suggesting an

independent mechanism of muscle-to-brain communication underlying these effects.

Skeletal muscle over-expression of TFEB decreases neuroinflammation markers and

lipofuscin accumulation in the healthy aged CNS.

To examine the effect of increased skeletal muscle proteostasis on the healthy aging brain

without neurodegenerative disease burden, we assessed the transcriptional and functional

status of known inflammatory markers in the CNS of healthy aged (20+ months old) control and

cTFEB;HSACre mice, an age when global decreases in proteostasis and increases in pro-

inflammatory signaling are detectable in most murine tissues [1,83], including the CNS [1,2]. Total

hippocampal mRNA levels of *Ccl2* and *Nfκb*, pro-inflammatory cytokines previously reported to

mediate microglial responses to inflammation, were significantly reduced in the hippocampus of

aging female (but not male) cTFEB;HSACre bigenic mice (**Figure 3A-B**). Interestingly, the

expression of *Il6* (interleukin 6) was significantly elevated in the same groups (**Figure 3C**). We

also detected significantly higher levels of expression of *Bdnf* (Brain Derived Neurotrophic

Factor) in hippocampal lysates from aged male (but not female) mice (**Figure 3D**), similar to

what we observed in our MAPT P301;cTFEB;HSACre model (**Figure 2H**).

Using similar morphometric analysis as before to quantify glial cell shape and morphology, we

did not detect any differences in microglia (IBA1+) or astrocyte (GFAP+) number, volume or

ramification state in the dentate gyrus of aged cTFEB;HSACre bigenic mice compared to their

littermate controls (**Figure 3E-F**). This may suggest that the expression differences observed

via transcriptional approaches above may either originate in non-glial cells (neurons or

endothelial cells, for example) or not translate into detectable morphological changes at the

ages examined. It is important to note that age-associated changes in neuroinflammation in an

otherwise healthy brain are not as pronounced as those seen in the context of

neurodegenerative disease (such as in the MAPT P301S model, for example). Overall, these

results suggest a shift in the cytokine transcriptional landscape of the aging hippocampus with

upregulated skeletal muscle proteostasis towards reduced inflammatory signaling in female

mice, consistent with what we observed in our MAPT P301S studies (**Figure 2**).

Accumulation of lipofuscin, a non-degradable intracellular auto-fluorescent polymer, becomes

prominent in the aging brain, likely reflecting an age-associated decline in basal CNS autophagy

[1]. We examined the clearance capabilities of the aging CNS by indirect immunofluorescence of

lipofuscin granules in brain sections of aged (21-24 months old) control and

cTFEB;HSACre mice. While we detected striking lipofuscin granule accumulation in several

brain regions of aged control mice, including the hippocampus (**Figure 3G, top**), there was a

significant decrease in lipofuscin deposition in the dentate gyrus of cTFEB;HSACre females

(**Figure 3G, bottom),** with a similar non-significant trend in males. To determine any changes in

proteostasis associated signaling in this brain region, we next examined the transcriptional

levels of multiple autophagy and lysosomal associated genes. Consistent with previous reports,

we confirmed significant age-associated increases in the expression of autophagy lysosomal

markers *Atg8/Lc3*, *Lamp1, Sqstm1,* and *Ctsd* in the male hippocampus, which has been

proposed to occur as a compensation for age-associated decline in proteostasis function

(**Figure 3H-K**). Interestingly, these differences were not as prominent in the aging female

hippocampus. Furthermore, some of these transcriptional increases were reduced in the

hippocampus of aged male cTFEB;HSACre bigenic mice, suggesting that enhancing skeletal

muscle proteostasis prevents the age-associated hippocampal proteostasis dysfunction

associated with age. Our data thus indicates a potential sex-bias in activation of neuroprotective

signaling in the hippocampus of aging cTFEB:HSACre bigenic mice, with females decreasing

neuroinflammation (**Figure 3A-C**) and males increasing neurotrophin expression levels (**Figure

3D**). Altogether, our results suggest that overexpression of TFEB in skeletal muscle throughout

the lifespan promotes maintenance of protein quality control regulation in the CNS, ultimately

reducing proteostatic burden and neuroinflammation in the aging hippocampus, similar to what

we observed in our MAPT P301S studies.

Improved performance in neurocognitive testing of aging cTFEB;HSACre mice.

Given the maintenance of proteostasis, reductions in neuroinflammation markers and increases

in *Bdnf* expression observed in the aging cTFEB;HSACre hippocampi, we pursued

neurocognitive testing in aged (16-18 months old) control and cTFEB;HSACre mice. First, we

evaluated spatial learning and memory using the Barnes maze task, a hippocampal working

memory test known to be sensitive to aging in mice. We documented significantly faster escape

times (**Figure 3L**) and a significant decrease in the number of errors per trial (**Figure 3M**) in

aged cTFEB;HSACre mice of both sexes in comparison to controls. Indeed, by the last

trial, cTFEB;HSACre mice escaped the maze twice as quickly as control mice. In the novel

object recognition task – an independent behavioral test of hippocampal recognition memory –

we also found that aged cTFEB;HSACre mice of both sexes exhibited a significantly greater

number of contacts with the novel object compared to controls during the test

phase (**Figure 3N**). Indeed, aged cTFEB;HSACre mice have around a ~15% higher preference

for the novel object relative to their age-matched littermate controls (**Figure 3O**). Importantly, we

also confirmed no differences in body weight (**Figure 3P**), visual performance, and motor

activity (not shown) between aged control and cTFEB;HSACre cohorts. We did detect a

significant increase in grip strength in aged male (and a similar trend in female) cTFEB;HSACre

bigenic mice (**Figure 3Q**), consistent with our findings suggesting that overexpression of TFEB

in skeletal muscle can also improve muscle function during aging (**Figure 1**).

Hence, these results provide exciting evidence for a pronounced improvement in neural function

in the aging brain of cTFEB;HSACre mice with enhanced muscle proteostasis.

Enhancing skeletal muscle proteostasis transcriptionally remodels the hippocampus of

cTFEB;HSACre bigenic mice throughout the lifespan in a sex-dependent manner.

To determine the molecular basis of the benefits of enhancing skeletal muscle proteostasis on

the aging CNS, we performed unbiased transcriptome analysis on hippocampal RNAs isolated

from young (6 months old) and aged (22-24 months old) mice (**Figures 4-5**). By this time point,

aging control animals have increased neuroinflammation (**Figure 3A-B**), have evident CNS

proteostasis dysfunction (**Figure 3G**), and display deficits in cognitive function (**Figure 3L-O**),

phenotypes that are significantly improved in cTFEB;HSACre mice.

Bulk RNA-Seq analysis of hippocampal RNA lysates revealed robust changes in gene

expression in young cTFEB;HSACre bigenic mice compared to their age- and sex-matched

littermate controls (**Figure 4, Figure 5, Supplemental Figure 3, and Supplemental Figure 4**).

In young female cTFEB;HSACre hippocampi, we detected 1194 differentially expressed genes,

(385 up-regulated and 809 down-regulated) (**Figure 4A**). To identify specific neural signaling

functions associated with these differentially expressed genes, we performed GO Enrichment

analysis. We found that key categories associated with synaptic function, including ion- and

voltage-gated channel activity, transporter complexes, as well as synapse and synaptic

membrane categories, were significantly enriched in the hippocampus of young female
cTFEB;HSACre mice (**Figure 4C**). Enrichment analysis of differential gene expression (KEGG)
revealed additional critical categories associated with neural function, including oxytocin
signaling, cAMP signaling, Hippo signaling and gap junctions (**Figure 4E**). Furthermore,
metabolic and functional pathway analysis (Reactome) of differentially expressed genes also
identified key categories known to regulate cognitive plasticity, including GPCR ligand binding,
voltage gated potassium channels, and transmission across chemical synapses, among others
(**Supplemental Figure 3A**).

Consistent with our biochemical and functional findings suggesting increased neuroprotective
benefits in aging cTFEB;HSACre mice (**Figure 3**), we found that these transcriptional changes
modulating synaptic function pathways are still present (although not as prominent) in the aging
hippocampus of aged female cTFEB;HSACre mice (**Figure 4B, Figure 4D, Supplemental**
Figure 3B, and Supplemental Figure 3D). Indeed, KEGG functional pathways enriched in our
aged female hippocampus included key categories associated with preservation of neuronal
activity and cognitive function, including axon guidance, HIF-1 signaling, AMPK signaling and
PI3-Akt signaling (**Figure 4F**). We confirmed the age-associated differential expression of a
subset of our top-hit genes in a second group of hippocampal RNA lysates using targeted qRT-
PCR (**Supplemental Figure 3E**). Altogether, this suggests that overexpression of TFEB in
skeletal muscle activates multiple neuroprotective gene networks associated with synaptic
function and signaling in the female hippocampus, and that this transcriptional remodeling is
preserved through aging.

Analysis of the functional enrichment of top differentially expressed genes in the hippocampus
of young male cTFEB;HSACre mice revealed a different pattern. We detected 1797 differentially
expressed genes (928 up-regulated and 869 down-regulated), which GO-term analysis
associated mostly with multiple mitochondria or ribosomal/transcription targeting pathways
(**Figure 5A and C**). This included mitochondrial respiratory chain, mitochondrial protein
complexes, electron transport chain and ATP metabolic processes (**Figure 5C**). Additional
enriched metabolic pathways in the hippocampus of young cTFEB;HSACre males included non-
sense mediated decay, metabolism of RNA and translation (**Supplemental Figure 4A**).
Consistent with this, KEGG analysis revealed the most enriched functional pathways were
ribosomes and oxidative phosphorylation (**Figure 5E**). As before, we confirmed the differential
expression of a random sampling of our top-hit genes using targeted qRT-PCR (**Supplemental**
Figure 4E). When we performed similar analysis on the male aging hippocampus, we only

detected 539 differentially expressed genes in the hippocampus of aged male cTFEB;HSACre

mice (279 up-regulated, 261 down-regulated) (**Figure 5B**), with only mild significance on

functional pathways (**Figure 5D, Figure 5F, Supplemental Figure 4B, and Supplemental**

Figure 4D). Our results thus strongly suggest a sex bias in the hippocampal transcriptional

responses associated with muscle-TFEB overexpression, with the female hippocampi up-

regulating neuronal signaling and synaptic function and the male hippocampi upregulating

transcriptional and mitochondrial responses. Finally, many of these changes are preserved up

until late stages in the lifespan of our model (24 months of age), suggesting that these

transcriptional changes may underlie the neurocognitive benefits observed in our aging

cTFEB;HSACre cohorts.

DISCUSSION

Skeletal muscle comprises roughly 40% of the total body mass in a healthy young adult human

and is highly susceptible to age-associated proteostatic decline [32,33]. Skeletal muscle autophagy

facilitates metabolic reprogramming and mitochondrial quality control during conditions of high

metabolic demand [34,57,84] and is also required to maintain muscle health and muscle mass [35,85]

[86]. Skeletal muscle health has been linked to multiple chronic age-associated conditions,

including diabetes [87], cardiovascular disease [88] and, more recently, neurodegenerative diseases

[24-26]. Here, and for the first time in mammals, we provide direct evidence demonstrating the

benefits of enhanced skeletal muscle proteostasis on the brain throughout aging and in the

context of neurodegenerative proteinopathy. We find that life-long overexpression of TFEB (a

master regulator of proteostasis and metabolism) reduces the development of age-associated

biochemical hallmarks in skeletal muscle. Indeed, we detected robust up-regulation of multiple

fundamental aging pathways [89], including proteostasis and mitochondrial function (**Figure 1**).

We also detected important changes in branched-chain amino acid signaling, which have been

recently linked to lifespan and healthspan extension in rodents [90], indicating that maintenance of

proteostasis through TFEB overexpression has pan-geroprotective effects on aging skeletal

muscle.

The hippocampus is a critical structure in cognitive aging, and hippocampal atrophy and

dysfunction are key features of both aging and neurodegenerative diseases [91,92]. In addition to

rescuing age-associated hallmarks in skeletal muscle, we also discovered significant

improvements in multiple disease-associated biochemical hallmarks in the aging hippocampus

of cTFEB;HSACre transgenic mice, ultimately preserving cognitive function in the aging brain.

Importantly, we observed these benefits without any detectable expression of our transgene in

the CNS (**Supplemental Figure 1**), suggesting that the observed reductions in proteotoxicity

and neuroinflammation (**Figure 2**), as well as the improved neurocognitive performance of aged

cTFEB;HSACre mice (**Figure 3**) are due solely to the overexpression of TFEB in the skeletal

muscle.

Tauopathies are clinically, biochemically and morphologically heterogeneous neurodegenerative

diseases characterized by the deposition of abnormal aggregates of tau in the brain. The MAPT

P301S model recapitulates multiple aspects of tauopathies, including microglial activation,

filamentous tau inclusions and synapse loss [67]. Our studies indicate that enhancing skeletal

muscle proteostasis has significant benefits on multiple markers of gliosis in this model of tau

toxicity, reducing astrocytic GFAP expression and the volume of microglia present in the

hippocampus of MAPT P301S mice (**Figure 2C**). This suggests that enhancing muscle

proteostasis in MAPT P301S mice reduces microglial and astrocyte activation in the

hippocampus, ultimately reducing neuroinflammation. Indeed, we confirmed significantly

reduced transcriptional activation of the nCounter AD microglial activation panel (**Figure 2F**), a

proxy for pro-inflammatory signaling in the context of AD-associated pathologies [81]. This

highlights both the robust pro-inflammatory state associated with tau pathogenesis, as well as

the intriguing possibility that manipulation of skeletal muscle proteostasis pathways may

represent a novel target for modulation of inflammation in the CNS.

Previous transcriptional studies of the healthy aging hippocampus have identified dysregulation

of ion homeostasis, disruption of neurotransmission, and ribosome biogenesis as key biological

process exhibiting differential regulation with aging [2,93]. Specifically, genes related to synaptic

transmission and plasticity appear to be particularly sensitive to age-associated transcriptional

dysregulation (reviewed here [3]). Many of these gene networks are also significantly altered in

AD and other associated tauopathies [81]. Importantly, similar networks were transcriptionally

remodeled in the healthy aging cTFEB;HSACre hippocampus, underscoring potential new

mechanisms of neuroprotective benefits. Further studies to identify specific responders in the

CNS (during healthy aging or in the context of neurodegenerative disease) may yield fascinating

new insights into avenues for therapeutic development.

Manipulation of skeletal muscle proteostasis modulates expression of similar functional

enrichment categories (ie. Trophic factor signaling, neuroinflammation) in the aging and tau-

afflicted hippocampus of our cTFEB;HSACre mice in a sex-dependent manner. Interestingly, we

detected significant reductions in the accumulation of hyperphosphorylated tau in both sexes in
the MAPT P301S model, suggesting differential mechanisms may underlie the neuroprotective
benefits mediated by enhancing skeletal muscle proteostasis. While it is unclear whether these
differential responses between male and female hippocampi to the enhancement of skeletal
muscle proteostasis reflect local differences in the CNS itself or peripheral differences in the
expression, secretion or trafficking of circulating signals, they underscore the fundamental need
to examine sex differences contributing to the biology of aging and neurodegenerative diseases.

Recent evidence has shown that muscle-originating circulating factors (myokines) appear to
play central roles in regulating CNS health and function. For example, Amyrel amylase, a
skeletal muscle secreted protein with maltose producing function, was identified as a key
proteostasis stress-induced myokine with CNS targeting effects in *Drosophila Melanogaster*[5].
Consistent with our own findings here, muscle-specific *amyrel* overexpression improves
proteostasis and prevents neurodegeneration induced by aggregation-prone disease-associated
pathogenic proteins in the fly brain and retina[5]. Another example is Fibronectin-domain III
containing 5 (FNDC5), a glycosylated type 1 membrane protein that is highly enriched in
skeletal muscle. FNDC5 can be proteolytically cleaved into an N-terminal fragment known as
irisin, which is then released into circulation[87]. Recent studies have demonstrated that
peripheral delivery of irisin reduces neuroinflammation and rescues cognitive decline in two
separate mouse models of AD, and is a central modulator of cognitive function[19]. Likewise, the
lysosomal protease CTSB is secreted from skeletal muscle into circulation in response to
exercise and is required for the full manifestation of exercise-associated benefits on the CNS,
including increases in the levels of hippocampal BDNF and the activation of neurogenesis[20].
Interestingly, recent evidence demonstrates pronounced involvement of the
macroautophagic/autophagic molecular machinery in cellular secretion[94]. Indeed, key signaling
molecules and cytokines associated with inflammation, such as IL1B (interleukin 1 beta) and IL6
(interleukin 6), are released into the extracellular space via autophagy-dependent secretion[94].
There is also a growing body of evidence that age[66] and exercise[95], known modifiers of
skeletal muscle proteostasis and metabolism, alter the composition of the totality of factors
produced and secreted by skeletal muscle. This suggests that activation of muscle proteostasis
via TFEB-overexpression may, in addition to the local geroprotective benefits in muscle tissue,
also be modifying the skeletal muscle 'secretome', potentially enriching it with neurotrophic
factors or depleting it from neurotoxic factors, although this hypothesis remains to be tested.

Importantly, our work indicates for the first time that one potential mechanism underlying the

neuroprotective benefits associated with enhancing skeletal muscle proteostasis is a reduction

in neuroinflammation, particularly in the female hippocampus. Although previous work

examining the muscle-to-brain axis has demonstrated important benefits on the accumulation of

misfolded protein inclusions in the CNS [5,8,19] (similar to what we observed in the MAPT

P301S;cTFEB;HSACre hippocampus), the profound reductions in both morphological and

transcriptional markers of neuroinflammation observed in this model suggest modulation of

skeletal muscle proteostasis may be a novel target to modulate neuroinflammatory signaling in

vivo in the context of aging and/or neurodegenerative diseases.

Over the last ten years, there has been growing evidence that suggests prominent contributions

of the periphery to the etiology of neurodegenerative diseases. Our discovery that skeletal

muscle proteostasis can be a crucial site for regulation of CNS health and function provides

compelling evidence for a new therapeutic delivery avenue into the brain, providing a currently

unexplored new diagnostic and therapeutic intervention site (i.e. skeletal muscle) for preserving

brain health.

METHODS

Generation of fxSTOP-TFEB transgenic mice.

We used a pCMV-human TFEB expressing vector from Origene (clone # sc122773), used previously [27]. We cloned a 3x-FLAG fragment via insertion into the Acc65I and Pac1 restriction sites of the targeting vector. The fxSTOP-TFEB vector was generated in multiple steps, as follows: (1) 5′ and 3′ targeting arms were inserted by PCR onto the ends of a minimal 1.8 kb, chloramphenicol-resistant vector; (2) an ampicillin resistance cassette was cloned into the MluI, XmaI sites of the targeting vector; (3) an 1.1 kb fragment encoding the actin promoter, a loxP site, a 3x-FLAG tag fused to EGFP was cloned into the NheI, PacI restriction sites of the targeting vector; (4) a 2.8 kb fragment containing the second loxP site, followed by 3x-FLAG-human TFEB sequence. The final fxSTOP-FLAG-human TFEB vector was microinjected into C57BL/6J:C3H/HeJ F1 hybrid oocytes. Of six founders identified, two gave rise to fxSTOP-FLAG-human TFEB lines with comparable phenotypes. Of these lines, we chose to focus our studies principally on line 419. The presence of the floxed 3x-FLAG-eGFP STOP-3x-FLAG-human TFEB cassette is verified by allele dependent qPCR genotyping analysis with each generation (Transnetyx), and expression of the 3x-FLAG-eGFP sequence (upstream of the STOP codon) in single transgenic fxSTOP-TFEB mice is assessed every other generation. We confirmed excision of 3x-FLAG-EGFP-STOP cassette in the presence of Cre-recombinase by crossing fxSTOP-TFEB transgenic female mice with HSA-Cre transgenic mice (JAX Strain No: 006139 | HSA-Cre79), and corresponds to the HSA-Cre driver mouse line from the laboratory of Dr. Judith Melki. TdTomato mice (B6.Cg-Gt(ROSA)26Sor^tm9(CAG-tdTomato)Hze/J, JAX Strain #:007909) express robust tdTomato fluorescence following Cre-mediated recombination, is congenic on the C57BL/6J genetic background and were also obtained from JAX. The MAPT P301S line (Stock No: 008169) originated in the laboratory of Dr. Virginia Lee. cTFEB;HSACre;MAPT P301S transgenic mice were generated by crossing fxSTOP-TFEB females to MAPT P301S males. Female double transgenic offspring was then crossed with heterozygote HSA-Cre transgenic males. All commercially available lines were obtained directly from JAX. All lines have been maintained in a C57/B6 background for over ten generations in our lab. Mice were housed in a temperature- and humidity-controlled environment on a 12-hour light/dark cycle with food and water ad libitum. Mice of both sexes were used for all experiments in equal numbers, or as close to equal numbers as possible. For all MAPT P301 experiments, control animals are MAPT P301S-;fxSTOP-TFEB-;HSACre- and/or MAPT P301S-;fxSTOP-TFEB+;HSACre- littermates. For all aging experiments, control animals are fxSTOP-TFEB-

;HSACre- and/or fxSTOP-TFEB+;HSACre- littermates. In our hands, we did not detect any

differences on tau or aging-associated CNS phenotypes on fxSTOP-TFEB+ animals compared

to the wild-type controls. All animal experimentation adhered to NIH guidelines and was

approved by and performed in accordance with the University of California, San Diego, Duke

University and University of Alabama at Birmingham Institutional Animal Care and Use. The

fxSTOp-TFEB line of transgenic mice is available for sharing with academic institutions free-of-

charge. For-profit institutions will adhere to UCSD's policy regarding copyrighted material.

Tissue Collections

Animals were anesthetized with 3.8% Avertin Solution. All animals received a transcardial

perfusion with 60 mLs of cold 1x PBS. Half of CNS tissue (right/left hemi-brain) was post-fixed in

4% PFA for less than 24 hours before being transferred to 30% sucrose for another 24 hours

before cryo-sectioning and staining the other half was micro-dissected into cortex and

hippocampus and flash-frozen in liquid nitrogen for RNA and protein analyses. Half of skeletal

muscle tissue (right/left limb muscle) was fresh frozen in OCT following dissection, while the

other half was flash-frozen in liquid nitrogen for RNA and protein analyses.

RT-PCR Analysis

Flash frozen perfused isolated mouse tissues (quadriceps, hippocampus) were placed in plastic

tubes with silica beads (MP Biomedical, 116913100) and homogenized in TRIzol (Invitrogen,

15596026). RNA was extracted via standard phenol-chloroform procedures followed by DNase

digestion (Thermo Fisher, AM2238). cDNA was generated using iScript (Bio-Rad Technologies,

1708891). Quantitative PCR was run using Sybrgreen (Life Technologies, 4309155) for

chemical detection (Applied Biosystems; QuantStudio 3). Enrichment was determined based on

double-delta CT value. Primers were ordered from IDT Predesigned qPCR Assays unless

otherwise specified. Primers used are listed in Primer Table 1.

ImmunoBlot Analysis

Protein lysates from muscle tissue was prepared as previously described [27,29]. Protein

concentration was quantified using a Pierce™ BCA Protein Assay (23227). Fifty or 25 µg of

homogenized proteins were loaded per lane, and after running Any KD, 10%, or 4-15% Mini-

PROTEAN TGX Gels (BioRad, 4568124, 4561034, and 4561084), samples were transferred to

0.2 µm (for LC3) or 0.45 µm (for everything else) PVDF membranes (BioRad, 162-0175 and

1704275), which were blocked in 5% BSA in PBS at RT for 1 hr. Membranes were incubated

with anti-FLAG antibody (Sigma, M2, 1:1000), anti-TFEB antibody (Cell Signaling,

4240,1:1000), anti-Lamp1 antibody (Novus Bio, NB600-956, 1:1000), anti-LC3 antibody (Cell

Signaling, 2775S, 1:1000) or anti-GAPDH (Invitrogen, AM4300, 1:5000) in PBS-T with 5% BSA

at 4°C overnight. The primary antibody was visualized with horseradish-peroxidase conjugated

anti-rabbit at (Cell Signaling, 7074P2, 1:5,000) and enhanced chemiluminescence (BioRad,

170-5060) or goat-anti-mouse IgG 680 (Invitrogen, A21058, 1:10,000). Densitometry analysis

was performed using the BioRad Image Lab 6.1 software application.

Immunoflourescence Studies

Tissue was embedded in OCT (TissueTek, 4583), frozen utilizing 2-methylbutane (VWR,

103525-278) submerged in liquid nitrogen and stored at -80C until used. All samples were

sectioned on a standing cryostat (Leica). Sections of brain were 20 microns, while muscle tissue

was 15 microns thick. For immunohistochemistry, sections were permeabilized with .25% Triton

for 15 mins and blocked with 4% BSA for 30 minutes-1 hour. Brain sections were then

incubated with anti-GFAP (ab4674, Abcam, 1:200), anti-IBA1 (Wako, 019-19741, 1:200), anti-

AT180 TAU (Thermo, MN1040, 1:100) and anti-AT8 TAU (Invitrogen, MN1020, 1:100) while

muscle sections were incubated with anti-laminin (Sigma, L9393, 1:200), anti-FLAG (Sigma,

F1804, 1:1000), anti-LC3 (Cell Signaling, 2775S, 1:200), anti-LAMP1 (DHSB, 1D4B, 1:100) and

anti-hTFFEB (CST, 4240, 1:200) overnight at 4°C and incubated with secondary antibodies at

RT for 1 hour (both diluted in 4% BSA). Next the slides were washed with Hoescht (Thermo

Scientific, 62249, 1:5000) and mounted with prolong glass (Invitrogen, P36984). All slides were

washed with 1X PBS three times for 5 minutes each between steps.

Slides were imaged in the UAB HIRF HRIF Confocal/Light Microscopy Core utilizing the 10x

objective on the Nikon A15R/SIM Confocal microscope. Z-stacks of the entire hippocampus

area/section were collected, and max intensity projections were generated using FIJI. Lipofuscin

imaging was performed using an epifluorescent Nikon light microscope with a 20x objective. For

TdTomato imaging, stitched whole-section z-stacks were acquired using a Nikon A1R HD25

confocal microscope of nuclei in the blue channel and native TdTomato fluorescence in the red

channel.

For quantification of astrocyte and/or microglia parameters, raw ND2 files were uploaded to the

NIS elements platform and thresholds were set to eliminate background in each channel in

order to delineate each object. Morphometric information was collected in addition to the

automated counting of objects. A size exclusion parameter was used for GFAP positive objects

under 10 microns, IBA-1 positive objects under 15 microns, and any object larger than 5,000

microns were all excluded due to standard assessments of cell sizes. FIJI was used for

quantification of lipofuscin and LAMP1 puncta.

Quantitative proteomics sample preparation and data-independent acquisition mass
spectrometry.

We identified distinct protein species from crude total extractions of whole quadricep muscle by

mass spectrometry after trypsin digestion, in collaboration with the University of Washington

Nathan Shock Center Protein Phenotypes of Aging Proteomics Core. Quadricep muscle powder

samples were processed and digested using S-Trap Micro Spin Column (Protifi, C02-micro-80)

following the manufacturer's protocol. SDS (5%) lysis and solubilization buffer were added to

liver powder samples and homogenized on ice using a sonicator probe. Protein concentrations

were determined using Pierce BCA Protein Assay Kit (Thermo Fisher Scientific, PI23227).

Proteins were digested at 1 ug trypsin to 25 ug protein ratio on S-Trap at 47°C for 1 h. Peptides

extracted from S-Trap were dried with a vacuum concentrator and reconstituted in 0.1% formic

acid in water prior to mass spectrometry acquisition. Data were acquired using data-

independent acquisition (DIA) on a Thermo EASY-nLC 1000 and a Thermo Q-Exactive HF

orbitrap mass spectrometer. Peptides were separated using PicoFrit Self-Packed Columns (360

µm OD x 75 µm ID; New Objective, Woburn, MA) packed to 30 cm with 3 µm ReproSil-Pur C18

beads (ESI Source Solutions, r13.aq.0001). The trap column was packed to 2 cm with same

C18 beads using 360 µm OD x 150 µm ID fused silica capillary fritted with Kasil on one end.

Solvent A was 0.1% formic acid in water, and solvent B was 0.1% formic acid in 80%

acetonitrile. For each injection, 1 µg of sample was loaded and eluted using a 90-min gradient

from 5% to 45% B at 250 nL/min. DIA methods followed the chromatogram library workflow as

described in detail previously [96]. All muscle samples were pooled in equal ratio to create a

muscle library sample. The library samples were used to generate chromatogram libraries using

the strategy described before [97]. Thermo Q Exactive HF MS was used to acquire six gas phase

fractionated runs spanning a total mass range of 400 to 1000 m/z, each with staggered 4 m/z

narrow precursor isolation windows [98]. EncyclopeDIA software suite was used to generate a

chromatogram library [96] and retention time models for all peptides detected at 1% FDR

determined by Percolator [99]. For quantitative DIA run, equal amounts of protein were used for

each sample. A 24 × 24 m/z staggered window from 400.43190 to 1000.7048 m/z was used as

described previously [98]. EncyclopeDIA software suite was used to search the chromatogram

library with calibrated retention time and quantify peptides and proteins in each sample.

Proteomics Analysis

A linear model was fit to the protein level data using the Bioconductor limma package [100], and

the estimated SVs were adjusted as covariates in our model. The limma package uses empirical

Bayes moderated statistics, which improves power by 'borrowing strength' between proteins in

order to moderate the residual variance [101]. Adjusted P value was calculated with Benjamini-

Hochberg method across groups (young transgenic, young control, old transgenic, and old

control). Proteins were sorted for significant differential expression using a cutoff of less than

5% FDR and greater than 1.5 foldchange in young and old cohorts as a change due to skeletal

muscle overexpression of humanized TFEB. Volcano plots were created in R studio with ggplot

and ggrepel. Significantly upregulated proteins in cTFEB;HSA-Cre mice were queried using the

KEGG pathway analyzer, which is a feature of STRING's web tool [63]. Bar graphs were created

in Rstudio with the ggplot2 packages.

Significantly upregulated proteins, up to two degrees from TFEB, were entered into STRINGdb

web tool for functional network analysis. A confidence interaction score of 0.70 and MCL

clustering was applied to the networks. The networks were exported from STRING and imported

into Cytoscape for further style editing [102].

Gene Expression Analysis.

RNA Sequencing and Bioinformatics Analysis

RNA was extracted as mentioned above from flash frozen hippocampal tissue for four biological

replicates (sex/genotype) and checked for quality. mRNA libraries were prepared by Novogene

(Novogene Corporation INC, Sacramento, CA, United States) and bulk RNA-seq analysis was

performed via NovaSeq PE150 (Illumina, San Diego, CA, United States) high throughput

sequencing. Bioinformatics analysis for differentially expressed genes was performed by

Novogene.

Nanostring nCounter AD panel

Assays were performed with 100 ng aliquots of RNA using the NanoString nCounter Analysis

system (NanoString Technologies, Seattle, WA, USA) at the UAB Nanostring Core, following

previously described and established protocols [73]. Counts for target genes were normalized to

house-keeping genes included in the panel (*Cltc, Gapdh, Gusb, Hprt, Pgk1,* and *Tubb5*). After

codeset hybridization overnight, the samples were washed and immobilized to a cartridge using

the NanoString nCounter Prep Station. Cartridges were scanned in the nCounter Digital

Analyzer at 555 fields of view for the maximum level of sensitivity. Gene expression was

normalized using NanoStringNorm R package. Specifically, background correction was

performed using the negative control at the cutoff of mean + 2 standard deviation. All p values

were adjusted using a false discovery rate (FDR) correction of 1% for multiple comparisons.

Housekeeping genes were used to for normalization based on geometric mean. Data and heat

analyses were performed in the nSolver Analysis Software 2.0. Gene expression values were

presented as the percentage of the MAPT P301S group for comparison of MAPT P301S

pathogenesis to the MAPT P301S;cTFEB;HSACre cohort.

Mouse Phenotyping and Behavioral Studies.

Barnes Maze

The Barnes maze apparatus is an opaque Plexiglas disc 75 cm in diameter elevated 58 cm

above the floor by a tripod. Twenty holes, 5 cm in diameter, are located 5 cm from the

perimeter, and a black Plexiglas escape box (19 x 8 x 7 cm) is placed under one of the holes.

Distinct spatial cues are located all around the maze and are kept constant throughout the

study. On the first day of testing, a training session was performed, which consists of placing the

mouse in the escape box for one minute. After the one-minute habituation period, the first

session was started. At the beginning of each session, the mouse was placed in the middle of

the maze in a 10 cm high cylindrical black start chamber. After 10 seconds the start chamber

was removed, a buzzer (80 dB) and a light (400 lux) were turned on, and the mouse was set

free to explore the maze. The session ended when the mouse entered the escape tunnel or

after 3 min elapsed. When the mouse entered the escape tunnel, the buzzer was turned off and

the mouse was allowed to remain in the dark for one minute. When the mouse did not enter the

tunnel by itself it was gently put in the escape box for one minute. The tunnel was always

located underneath the same hole (stable within the spatial environment), which was randomly

determined for each mouse. Mice were tested once a day for 4 days for the acquisition portion

of the study. For the 5th test (probe test), the escape tunnel was removed and the mouse was

allowed to freely explore the maze for 3 min. The time spent in each quadrant was determined

and the percent time spent in the target quadrant (the one originally containing the escape box)

was compared with the average percent time in the other three quadrants. This was a direct test

of spatial memory as there was no potential for local cues to be used in the mouse's behavioral

decision. Two weeks later the mice were tested again with the escape box in the original

position (retention test). This allows for the examination of long term memory. Finally, on the

day after this test, the escape tunnel was moved to a new location (90 degrees from the original

position) and the behavior of the mouse was recorded. This is called the reversal test and it

allows for the examination of perseveration at the old hole as well as the working memory

strategies the mice adopted to locate the new tunnel location. Each session was videotaped and

scored by an experimenter blind to the genotype of the mouse. Measures recorded include the

latency to enter the escape box and the number of errors made per session. Errors are defined

as nose pokes and head deflections over any hole that did not have the tunnel beneath it. The

probe data were analyzed using Noldus Ethovision software to determine time spent in each

quadrant of the maze as well as to assess activity measures.

Novel Object Recognition

Mice were individually habituated to a 51cm x 51cm x 39cm open field for 5 min. Mice were then

tested with two identical objects placed in the field (either two 250 ml amber bottles or two clear

plastic cylinders 6x6x16cm half filled with glass marbles). An individual animal was allowed to

explore for 5 min, now with the objects present. After two such trials (each separated by 1

minute in a holding cage), the mouse was tested in the object novelty recognition test in which a

novel object replaced one of the familiar objects (for example, an amber bottle if the cylinders

were initially used). All objects and the arena were thoroughly cleaned with 70% ethanol

between trials to remove odors. Behavior was video recorded and then scored for number

contacts (touching with nose or nose pointing at object and within 0.5 cm of object) and/or for

time contacting the objects. Habituation to the objects across the familiarization trials

(decreased contacts) was an initial measure of learning and then renewed interest (increased

contacts) in the new object indicated successful object memory. Recognition indexes were

calculated using the following formula: (# contacts during test)/(# contacts in last familiarization

trial + # contacts during test). Values greater than 0.5 indicate increased interest, whereas

values less than 0.5 indicate decreased interest in the object during the test relative to the final

familiarization trial.

Statistical Analysis

All data were analyzed by t-test, 1-way, 2-way or 3-way between-subject ANOVA with post hoc

comparisons depending on the number of variables and groups in each analysis. For ANOVA, if

statistical significance ($p < 0.05$) was achieved, we performed post hoc analysis to account for

multiple comparisons. The level of significance (α) was always set at 0.05. Survival curves were

analyzed using Log-rank (Mantel-Cox) Test. Data were analyzed using Prism 7 (GraphPad

Software, La Jolla, CA) and are represented as means and standard error of the means. All

experiments and data analyses were conducted in a blinded fashion. All data were prepared for

analysis with standard spread sheet software (Microsoft Excel).

Figure Legends

Figure 1: Autophagy and lysosomal activation in skeletal muscle protects against age-associated phenotypes.

(A) Schematic depicting 3x-FLAG human TFEB transgene. Cassettes are arranged from 5' (left) to 3' (right). Note that the 3x-FLAG-EGFP-STOP cassette (flanked by loxP sites) prevents expression of the 3x-FLAG human TFEB transgene unless in the presence of Cre recombinase β-actin promoter.

(B) Immunoblot of control (left) and cTFEB;HSACre (right) quadriceps lysates. Transgenic 3x-FLAG-TFEB protein weighs ~50 kDa (arrow). Non-specific band is shown below as a loading control (asterisk).

(C-F) qRT-PCR for *Lamp1* (C), *Atg5* (D), *Ctsd* (E), and *CtsB* (F) expression in quadriceps mRNA lysates from young (6 months old) male cTFEB;HSACre mice and controls.

(G) Immunoblot analysis for LAMP1 and Cathepsin D protein in quadriceps protein lysates from young (6 months old) female cTFEB;HSACre mice and controls. Densitometry quantification relative to average of control individuals shown below.

(H) Representative images of gastrocnemius from young (6-7 months old) control and cTFEB;HSACre mice stained for LAMP1 (green), nuclei/Hoescht (blue). Quantification of LAMP1+ puncta and LAMP1+ area/muscle section are shown on the right. Each data point represents the average of 10 separate images collected from 3-5 sections/individual. Scale bars = 100 μm.

(I-J) Volcano plot of proteomics analysis depicting differentially regulated proteins in young (I, 6 months old, I) and aged (J, 18 months old) cTFEB;HSACre quadriceps muscle relative to controls. Overexpressed proteins in red, underexpressed proteins in blue. Note that TFEB itself (rectangles) is highly expressed in both aged and young muscle.

(K) STRING predicted protein-protein interactions of overexpressed proteins from (I) showing predicted interacting networks converging on TFEB (blue) and mTOR (green).

(L) KEGG enrichment analysis clusters overexpressed proteins from (I-J) into categories associated with proteostasis and autophagy (orange), mitochondrial metabolism (gold), fatty acid metabolism (green), age-associated signaling (blue), and amino acid metabolism (purple).

(M) Immunoblot for LC3 protein levels in quadriceps lysates from young (6 months old) female cTFEB;HSACre and controls. Densitometry quantification of LC3II/LC3I Ratio (normalized to tubulin) shown below.

(N) Representative images of tibialis anterior cross sections stained for Hematoxylin/Eosin (left)

and Trichrome (right) in aged (16 months old) control and cTFEB;HSACre transgenic mice.

Intrafibrillary myonuclei (yellow, arrows) and multilaminar structures (yellow, asterisks) are

indicated. Scale bars = 20 μm.

(O-P) qRT-PCR for *Mstn* (O) and *Fndc5* (P) expression in quadriceps mRNA lysates from aged

(20-22 months old) cTFEB;HSACre and controls.

Each point represents the average of all sections from an individual. Controls are age-matched

littermates. Unless otherwise stated, data are presented as mean ± SEM * p<0.05, ** p<0.01, T-

test.

Figure 2: Skeletal muscle TFEB-overexpression reduces accumulation of

hyperphosphorylated tau and microglial activation in a mouse model of tau pathology.

(A) Immunoblot of 3x-FLAG TFEB protein in quadriceps muscle lysates from 9-month old MAPT

P301S;cTFEB;HSACre and MAPT P301S mice (top). Immunoblot for total Tau protein (T6

antibody) in hippocampal lysates from same individuals (bottom). GAPDH is shown below as a

loading control.

(B) Representative merged images of the hippocampus dentate gyrus stained for

phosphorylated tau (AT8, white) and nuclei/Hoescht (blue) in 9-month old control (top, in gold),

MAPT P301S (middle, in teal), and MAPT P301S;cTFEB;HSACre mice (bottom, in purple).

Insets depicting perinuclear AT8 staining are 5X zooms of areas demarcated by yellow squares.

Quantification of total phosphotau staining/section (right, top) and for intracellular phosphotau

staining, (right, bottom). Each point represents the average of all sections containing the dentate

gyrus (2-3 sections) for an individual animal. Scale bars = 100 μm.

(C) Representative merged images of the dentate gyrus stained for astrocytes (GFAP, green),

microglia (IBA1, red) and nuclei (Hoescht, blue) in 9-month old control (top, gold), MAPT P301S

(middle, teal), and MAPT P301S;cTFEB;HSACre mice (bottom, purple). Insets depicting glia

morphology are 5X zooms of areas demarcated by yellow squares. Quantification of IBA1+

microglia volume (top, right) and GFAP+ astrocyte staining amounts per section (bottom, right).

Each point represents the average of all sections containing the dentate gyrus (2-3 sections) for

an individual animal. Scale bars = 100 μm.

(D-E) Volcano plot of differentially expressed genes in the hippocampi of 9-month old male (D)

and female (E) MAPT P301S;cTFEB:HSACre animals relative to age-matched P301S MAPT

littermates using the Nanostring nCounter AD panel. Down-regulated genes in blue, up-

regulated genes in red.

(F-G) Nanostring nCounter AD panel cluster analysis of differentially expressed genes from D-E
for gene clusters associated with microglial activation (F) and with trophic factors (G) in MAPT
P301S;cTFEB;HSACre hippocampi (purple) compared to P301S MAPT (teal) age-matched
littermates. Analysis done via nSolver (Nanostring) differential gene expression analysis
software. Each point represents a single individual hippocampal mRNA lysate.
(H-I) qRT-PCR for *Bdnf* (H) and *Fndc5* (I) expression in hippocampal RNA lysates from 9-month
old male MAPT P301S (teal) and MAPT P301S;cTFEB;HSACre mice (purple) relative to
controls (not shown).
Controls are littermates. Unless otherwise noted, data are represented as mean ± SEM *
p<0.05, ** p<0.01,*** p<0.001, One-way ANOVA, post-hoc T-test (B,C) and t-test (F-I).
881
Figure 3: Skeletal muscle TFEB-overexpression decreases neuroinflammation, lipofuscin
accumulation, and improves cognitive performance in aged mice.
(A-D) qRT-PCR for *Ccl2* (A), *Nfkb* (B), *Il6* (C), and *Bdnf* (D) expression in hippocampal mRNA
lysates from 18-month old cTFEB;HSACre mice and controls.
(E-F) Representative images dentate gyrus of the hippocampus stained for Hoescht (blue),
IBA1 (E, red), and GFAP (F, green) in 18-22-month old control (top) and cTFEB;HSACre
(bottom) mice. Insets depicting glia morphology are 5X zooms of areas demarcated by yellow
squares. Quantification of IBA1+ microglial volume (E, top right) and IBA1+ branching index (E,
bottom right). Average number of GFAP+ astrocytes pixels per section (F, top) and GFAP+
astrocyte branching index (F, bottom). Each point represents the average of all sections
containing the dentate gyrus from an individual mouse. Scale bars = 100 µm.
(G) Representative merged images of the dentate gyrus of 18-22-month old control (top) and
cTFEB;HSACre (middle) mice stained for Hoescht (blue), and lipofuscin puncta
(autofluorescence, green). Insets depicting autofluorescent puncta within the dentate gyrus are
5X zooms of areas demarcated by yellow squares. Scale bars = 100 µm.
(H-K) qRT-PCR for *Atg8/LC3* (H), *Lamp1* (I), *Sqstm1* (J), and *Ctsd* (K) expression in
hippocampal lysates from young (6 months old, closed diamonds) and aged (21 months old,
open diamonds) cTFEB;HSACre mice (purple diamonds) and controls (gold diamonds).
(L-Q) Neurocognitive phenotyping of aged (18 months old) cTFEB;HSACre and control animals.
Barnes maze (L,M), novel object recognition (N,O) total body weight (P) and grip strength (Q).
Controls are age-matched littermates. Data are represented as mean ± SEM * p<0.05, **
p<0.01, One-way ANOVA or two-way RMANOVA, post-hoc T-test.
904

Figure 4: Transcriptome changes in the hippocampi of female cTFEB;HSACre transgenic mice suggest synaptic remodeling and improved synaptic function.

(A-B) Volcano plot of differentially expressed genes in hippocampal lysates from female young (A, 6 months old) and aged (B, 24 months old) cTFEB;HSACre mice relative to controls.

(C-D) GO term analysis of differentially expressed genes from A and B including gene counts and adjusted p values. Boxes highlight neurotrophic signaling pathways.

(E-F) KEGG analysis of differentially expressed genes from A and B. Number of differentially expressed genes (DE) are shown above each category. Arrows highlight categories associated with neural function.

Figure 5: Transcriptome changes in the hippocampi of male cTFEB;HSACre transgenic mice suggest activation of mitochondrial and ribosomal signaling.

(A-B) Volcano plot of differentially expressed genes in hippocampal lysates from male young (A, 6 months old) and aged (B, 24 months old) cTFEB;HSACre mice relative to controls.

(C-D) GO term analysis of differentially expressed genes from A and B including gene counts and adjusted p values. Boxes highlight ribosomal and mitochondrial pathways.

(E-F) KEGG analysis of differentially expressed genes from A and B. Number of differentially expressed genes (DE) are shown above each category. Arrows highlight ribosomal and mitochondrial pathways.

Supplemental Figures

Supplemental Figure 1

(A-B) qRT-PCR analysis for human *TFEB* mRNA expression in 6-month old (A) and 21-month old (B) cTFEB;HSACre quadriceps skeletal muscle (left) and hippocampus (right) compared to controls.

(C) Representative images of gastrocnemius cross sections stained for Hoescht (blue) and Human TFEB (white). Insets of nuclear hTFEB staining are 5x zooms of the areas denoted by the yellow squares. Scale bars = 100 μm.

(D-F) Immunoblot analysis for 3x-FLAG TFEB protein expression from lysates of brown adipose (D), liver (E), and brain (F) tissue from 6-month old cTFEB;HSACre mice and age-matched control littermates. cTFEB;HSACre skeletal muscle lysates are shown in the first lane of each

blot as a positive control for 3x-FLAG TFEB expression. FLAG-eGFP protein cassette (arrow)

as well as non-specific bands (asterisks) are indicated.

(G) Representative images of quadriceps and gastrocnemius cross sections from 7-week old

TdTomato;HSACre transgenic mice stained for Hoescht (blue), TdTomato (red), and Laminin

(white). Scale bars = 100 μm.

(H) Representative images of hemibrain sections from TdTomato;HSACre transgenic mice

stained for Hoescht (blue), TdTomato (red). Insets of hippocampal regions are 5x zooms of the

areas denoted by the yellow squares. Scale bars = 1 cm.

Controls are age-matched littermates. Data are represented as mean ± SEM, T-test, * p<0.05,

** p<0.01, *** p<0.001.

Supplemental Figure 2

(A) Representative images of the hippocampal dentate gyrus from 9-month old control (top),

MAPT P301S (middle), and MAPT P301S;cTFEB;HSACre (bottom) mice stained for Hoescht

(blue) and AT180 PhosphoTau (white). Insets of hippocampal regions are 5x zooms of the

areas denoted by the yellow squares. Scale bars= 100 μm.

(B) Cumulative ataxia score of 9-month old control, MAPT P301S, and MAPT

P301S;cTFEB;HSACre mice.

(C) Weight (in grams) of 11-month old control MAPT P301S, and MAPT P301S;cTFEB;HSACre

mice.

(D-E) Survival curves for male (D) and female (E) control, MAPT P301S, and MAPT

P301S;cTFEB;HSACre mice.

Controls are age-matched littermates. Data are represented as mean ± SEM, One-way ANOVA

post-hoc T-test or Log-rank (Mantel-Cox) test * p<0.05, ** p<0.01.

Supplemental Figure 3

(A-B) Reactome analysis of differentially expressed genes in hippocampal mRNA lysates from

female young (6 months old, A) and aged (24 months old, B) cTFEB;HSACre mice. Number of

differentially expressed genes (DE) are shown above each category.

(C-D) GO enrichment analysis of differentially expressed genes from A-B. Number of

differentially expressed genes (DE) are shown above each category.

(E) qRT-PCR analysis for select DE genes in hippocampal mRNA lysates from female young (6

months old, solid circles) and aged (24 months old, open circles) control (gold) and

cTFEB;HSACre (purple) mice. Expression levels normalized to young controls.

Controls are age-matched littermates. Data are represented as mean ± SEM, One-way ANOVA

post-hoc T-test * p<0.05, ** p<0.01.

Supplemental Figure 4

(A-B) Reactome analysis of differentially expressed genes in hippocampal mRNA lysates from

male young (6 months old, A) and aged (24 months old, B) cTFEB;HSACremice. Number of

differentially expressed genes (DE) are shown above each category.

(C-D) GO enrichment analysis of differentially expressed genes from A-B. Number of

differentially expressed genes (DE) are shown above each category.

(E) qRT-PCR analysis for select DE genes in hippocampal mRNA lysates from female young (6

months old, solid circles) and aged (24 months old, open circles) control (gold) and

cTFEB;HSACre (purple) mice. Expression levels normalized to young controls.

Controls are age-matched littermates. Data are represented as mean ± SEM, One-way ANOVA

post-hoc T-test * p<0.05, ** p<0.01.

REFERENCES

1   Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging.
Cell **153**, 1194-1217, doi:10.1016/j.cell.2013.05.039 (2013).

2   Mattson, M. P. & Arumugam, T. V. Hallmarks of Brain Aging: Adaptive and Pathological
Modification by Metabolic States. *Cell Metab* **27**, 1176-1199, doi:10.1016/j.cmet.2018.05.011
(2018).

3   Ham, S. & Lee, S. V. Advances in transcriptome analysis of human brain aging. *Exp Mol Med* **52**,
1787-1797, doi:10.1038/s12276-020-00522-6 (2020).

4   Demontis, F., Patel, V. K., Swindell, W. R. & Perrimon, N. Intertissue control of the nucleolus via
a myokine-dependent longevity pathway. *Cell Rep* **7**, 1481-1494,
doi:10.1016/j.celrep.2014.05.001 (2014).

5   Rai, M. *et al.* Proteasome stress in skeletal muscle mounts a long-range protective response that
delays retinal and brain aging. *Cell Metab* **33**, 1137-1154 e1139, doi:10.1016/j.cmet.2021.03.005
(2021).

6   Durieux, J., Wolff, S. & Dillin, A. The cell-non-autonomous nature of electron transport chain-
mediated longevity. *Cell* **144**, 79-91, doi:10.1016/j.cell.2010.12.016 (2011).

7   Taylor, R. C. & Dillin, A. XBP-1 is a cell-nonautonomous regulator of stress resistance and
longevity. *Cell* **153**, 1435-1447, doi:10.1016/j.cell.2013.05.042 (2013).

8   Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in Drosophila muscles regulates organism-
wide proteostasis during aging. *Cell* **143**, 813-825, doi:10.1016/j.cell.2010.10.007S0092-
8674(10)01143-8 [pii] (2010).

9   van Oosten-Hawle, P., Porter, R. S. & Morimoto, R. I. Regulation of organismal proteostasis by
transcellular chaperone signaling. *Cell* **153**, 1366-1378, doi:10.1016/j.cell.2013.05.015 (2013).

10   Villeda, S. A. *et al.* The ageing systemic milieu negatively regulates neurogenesis and cognitive
function. *Nature* **477**, 90-94, doi:10.1038/nature10357 (2011).

11   Conboy, I. M. *et al.* Rejuvenation of aged progenitor cells by exposure to a young systemic
environment. *Nature* **433**, 760-764, doi:10.1038/nature03260 (2005).

12   Castellano, J. M. *et al.* Human umbilical cord plasma proteins revitalize hippocampal function in
aged mice. *Nature* **544**, 488-492, doi:10.1038/nature22067 (2017).

13   Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature* **583**, 596-
602, doi:10.1038/s41586-020-2499-y (2020).

14   Villeda, S. A. *et al.* Young blood reverses age-related impairments in cognitive function and

synaptic plasticity in mice. *Nat Med* **20**, 659-663, doi:10.1038/nm.3569 (2014).

15   De Miguel, Z. *et al.* Exercise plasma boosts memory and dampens brain inflammation via

clusterin. *Nature*, doi:10.1038/s41586-021-04183-x (2021).

16   Horowitz, A. M. *et al.* Blood factors transfer beneficial effects of exercise on neurogenesis and

cognition to the aged brain. *Science* **369**, 167-173, doi:10.1126/science.aaw2622 (2020).

17   Ozek, C., Krolewski, R. C., Buchanan, S. M. & Rubin, L. L. Growth Differentiation Factor 11

treatment leads to neuronal and vascular improvements in the hippocampus of aged mice. *Sci*

Rep **8**, 17293, doi:10.1038/s41598-018-35716-6 (2018).

18   Wrann, C. D. *et al.* Exercise induces hippocampal BDNF through a PGC-1α/FNDC5 pathway. *Cell*

Metab **18**, 649-659, doi:10.1016/j.cmet.2013.09.008 (2013).

19   Islam, M. R. *et al.* Exercise hormone irisin is a critical regulator of cognitive function. *Nat Metab*

3, 1058-1070, doi:10.1038/s42255-021-00438-z (2021).

20   Moon, H. Y. *et al.* Running-Induced Systemic Cathepsin B Secretion Is Associated with Memory

Function. *Cell Metab* **24**, 332-340, doi:10.1016/j.cmet.2016.05.025 (2016).

21   White, P. J. *et al.* Protectin DX alleviates insulin resistance by activating a myokine-liver

glucoregulatory axis. *Nat Med* **20**, 664-669, doi:10.1038/nm.3549 (2014).

22   Quinn, L. S., Anderson, B. G., Strait-Bodey, L., Stroud, A. M. & Argilés, J. M. Oversecretion of

interleukin-15 from skeletal muscle reduces adiposity. *American Journal of Physiology-*

Endocrinology and Metabolism **296**, E191-E202 (2009).

23   Lourenco, M. V. *et al.* Exercise-linked FNDC5/irisin rescues synaptic plasticity and memory

defects in Alzheimer's models. *Nat Med* **25**, 165-175, doi:10.1038/s41591-018-0275-4 (2019).

24   Gupta, R., Khan, R. & Cortes, C. J. Forgot to Exercise? Exercise Derived Circulating Myokines in

Alzheimer's Disease: A Perspective. *Front Neurol* **12**, 649452, doi:10.3389/fneur.2021.649452

(2021).

25   Boyle, P. A., Buchman, A. S., Wilson, R. S., Leurgans, S. E. & Bennett, D. A. Association of muscle

strength with the risk of Alzheimer disease and the rate of cognitive decline in community-

dwelling older persons. *Arch Neurol* **66**, 1339-1344, doi:10.1001/archneurol.2009.240 (2009).

26   Kim, J. *et al.* Association of muscle and visceral adipose tissues with the probability of

Alzheimer's disease in healthy subjects. *Sci Rep* **9**, 949, doi:10.1038/s41598-018-37244-9 (2019).

27   Cortes, C. J. *et al.* Polyglutamine-expanded androgen receptor interferes with TFEB to elicit

autophagy defects in SBMA. *Nature neuroscience* **17**, 1180-1189, doi:10.1038/nn.3787 (2014).

28   Cortes, C. J. & La Spada, A. R. Motor neuron degeneration in spinal and Bulbar Muscular Atrophy
is a skeletal muscle-driven process: Relevance to therapy development and implications for
related motor neuron diseases. *Rare Dis* **2**, e962402, doi:10.4161/2167549X.2014.962402
(2014).

29   Cortes, C. J. *et al.* Muscle expression of mutant androgen receptor accounts for systemic and
motor neuron disease phenotypes in spinal and bulbar muscular atrophy. *Neuron* **82**, 295-307,
doi:10.1016/j.neuron.2014.03.001 (2014).

30   Lieberman, A. P. *et al.* Peripheral androgen receptor gene suppression rescues disease in mouse
models of spinal and bulbar muscular atrophy. *Cell Rep* **7**, 774-784,
doi:10.1016/j.celrep.2014.02.008 (2014).

31   Aman, Y. *et al.* Autophagy in healthy aging and disease. *Nat Aging* **1**, 634-650,
doi:10.1038/s43587-021-00098-4 (2021).

32   Jiao, J. & Demontis, F. Skeletal muscle autophagy and its role in sarcopenia and organismal
aging. *Curr Opin Pharmacol* **34**, 1-6, doi:10.1016/j.coph.2017.03.009 (2017).

33   Wohlgemuth, S. E., Seo, A. Y., Marzetti, E., Lees, H. A. & Leeuwenburgh, C. Skeletal muscle
autophagy and apoptosis during aging: effects of calorie restriction and life-long exercise. *Exp*
Gerontol **45**, 138-148, doi:10.1016/j.exger.2009.11.002 (2010).

34   Grumati, P. *et al.* Autophagy is defective in collagen VI muscular dystrophies, and its reactivation
rescues myofiber degeneration. *Nat Med* **16**, 1313-1320, doi:10.1038/nm.2247 (2010).

35   Masiero, E. *et al.* Autophagy is required to maintain muscle mass. *Cell Metab* **10**, 507-515,
doi:10.1016/j.cmet.2009.10.008 (2009).

36   Masiero, E. & Sandri, M. Autophagy inhibition induces atrophy and myopathy in adult skeletal
muscles. *Autophagy* **6**, 307-309 (2010).

37   Nascimbeni, A. C., Fanin, M., Masiero, E., Angelini, C. & Sandri, M. The role of autophagy in the
pathogenesis of glycogen storage disease type II (GSDII). *Cell death and differentiation* **19**, 1698-
1708, doi:10.1038/cdd.2012.52 (2012).

38   Hara, T. *et al.* Suppression of basal autophagy in neural cells causes neurodegenerative disease
in mice. *Nature* **441**, 885-889, doi:nature04724 [pii]10.1038/nature04724 (2006).

39   Komatsu, M. *et al.* Loss of autophagy in the central nervous system causes neurodegeneration in
mice. *Nature* **441**, 880-884, doi:nature04723 [pii]10.1038/nature04723 (2006).

40   Mansueto, G. *et al.* Transcription Factor EB Controls Metabolic Flexibility during Exercise. *Cell*
Metab **25**, 182-196, doi:10.1016/j.cmet.2016.11.003 (2017).

41   Mammucari, C. *et al.* FoxO3 controls autophagy in skeletal muscle in vivo. *Cell Metab* **6**, 458-
471, doi:10.1016/j.cmet.2007.11.001 (2007).

42   Romanello, V. & Sandri, M. Mitochondrial Quality Control and Muscle Mass Maintenance. *Front*
Physiol **6**, 422, doi:10.3389/fphys.2015.00422 (2015).

43   Sandri, M. *et al.* PGC-1alpha protects skeletal muscle from atrophy by suppressing FoxO3 action
and atrophy-specific gene transcription. *Proc Natl Acad Sci U S A* **103**, 16260-16265,
doi:10.1073/pnas.0607795103 (2006).

44   Kim, K. H. *et al.* Autophagy deficiency leads to protection from obesity and insulin resistance by
inducing Fgf21 as a mitokine. *Nat Med* **19**, 83-92, doi:10.1038/nm.3014 (2013).

45   Kustermann, M. *et al.* Loss of the novel Vcp (valosin containing protein) interactor Washc4
interferes with autophagy-mediated proteostasis in striated muscle and leads to myopathy in
vivo. *Autophagy* **14**, 1911-1927, doi:10.1080/15548627.2018.1491491 (2018).

46   Lapierre, L. R. *et al.* The TFEB orthologue HLH-30 regulates autophagy and modulates longevity
in Caenorhabditis elegans. *Nature communications* **4**, 2267, doi:10.1038/ncomms3267 (2013).

47   Pyo, J. O. *et al.* Overexpression of Atg5 in mice activates autophagy and extends lifespan. *Nature*
communications **4**, 2300, doi:10.1038/ncomms3300 (2013).

48   Simonsen, A. *et al.* Promoting basal levels of autophagy in the nervous system enhances
longevity and oxidant resistance in adult Drosophila. *Autophagy* **4**, 176-184, doi:5269 [pii]
(2008).

49   Colman, R. J. *et al.* Caloric restriction delays disease onset and mortality in rhesus monkeys.
Science **325**, 201-204, doi:10.1126/science.1173635325/5937/201 [pii] (2009).

50   Eisenberg, T. *et al.* Induction of autophagy by spermidine promotes longevity. *Nat Cell Biol* **11**,
1305-1314, doi:10.1038/ncb1975ncb1975 [pii] (2009).

51   Harrison, D. E. *et al.* Rapamycin fed late in life extends lifespan in genetically heterogeneous
mice. *Nature* **460**, 392-395, doi:10.1038/nature08221nature08221 [pii] (2009).

52   Settembre, C. *et al.* TFEB links autophagy to lysosomal biogenesis. *Science* **332**, 1429-1433,
doi:10.1126/science.1204592 (2011).

53   Palmieri, M. *et al.* Characterization of the CLEAR network reveals an integrated control of
cellular clearance pathways. *Human molecular genetics* **20**, 3852-3866, doi:ddr306
[pii]10.1093/hmg/ddr306 (2011).

54   Sardiello, M. *et al.* A gene network regulating lysosomal biogenesis and function. *Science* **325**,
473-477, doi:1174447 [pii]10.1126/science.1174447 (2009).

55   Morgan, T. E. *et al.* The mosaic of brain glial hyperactivity during normal ageing and its
attenuation by food restriction. *Neuroscience* **89**, 687-699, doi:10.1016/s0306-4522(98)00334-0
(1999).

56   van Praag, H., Shubert, T., Zhao, C. & Gage, F. H. Exercise enhances learning and hippocampal
neurogenesis in aged mice. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **25**, 8680-8685, doi:10.1523/jneurosci.1731-05.2005 (2005).

57   Settembre, C. *et al.* TFEB controls cellular lipid metabolism through a starvation-induced
autoregulatory loop. *Nature cell biology* **15**, 647-658, doi:10.1038/ncb2718 (2013).

58   Leu, M. *et al.* Erbb2 regulates neuromuscular synapse formation and is essential for muscle
spindle development. *Development* **130**, 2291-2301 (2003).

59   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for
the whole mouse brain. *Nat Neurosci* **13**, 133-140, doi:10.1038/nn.2467 (2010).

60   Spampanato, C. *et al.* Transcription factor EB (TFEB) is a new therapeutic target for Pompe
disease. *EMBO Mol Med* **5**, 691-706, doi:10.1002/emmm.201202176 (2013).

61   Lee, D. E., Bareja, A., Bartlett, D. B. & White, J. P. Autophagy as a Therapeutic Target to Enhance
Aged Muscle Regeneration. *Cells* **8**, doi:10.3390/cells8020183 (2019).

62   Giacomello, E. *et al.* Age Dependent Modification of the Metabolic Profile of the Tibialis Anterior
Muscle Fibers in C57BL/6J Mice. *Int J Mol Sci* **21**, doi:10.3390/ijms21113923 (2020).

63   Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein-protein networks, and
functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res* **49**,
D605-D612, doi:10.1093/nar/gkaa1074 (2021).

64   Decressac, M. *et al.* TFEB-mediated autophagy rescues midbrain dopamine neurons from alpha-
synuclein toxicity. *Proc Natl Acad Sci U S A* **110**, E1817-1826, doi:10.1073/pnas.1305623110
(2013).

65   Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy
(3rd edition). *Autophagy* **12**, 1-222, doi:10.1080/15548627.2015.1100356 (2016).

66   Mancinelli, R. *et al.* Biological Aspects of Selected Myokines in Skeletal Muscle: Focus on Aging.
Int J Mol Sci **22**, doi:10.3390/ijms22168520 (2021).

67   Yoshiyama, Y. *et al.* Synapse loss and microglial activation precede tangles in a P301S tauopathy
mouse model. *Neuron* **53**, 337-351, doi:10.1016/j.neuron.2007.01.010 (2007).

68   Moloney, C. M., Lowe, V. J. & Murray, M. E. Visualization of neurofibrillary tangle maturity in
Alzheimer's disease: A clinicopathologic perspective for biomarker research. *Alzheimers Dement*
17, 1554-1574, doi:10.1002/alz.12321 (2021).

69   Jurga, A. M., Paleczna, M. & Kuter, K. Z. Overview of General and Discriminating Markers of
Differential Microglia Phenotypes. *Front Cell Neurosci* **14**, 198, doi:10.3389/fncel.2020.00198
(2020).

70   Rozovsky, I., Finch, C. E. & Morgan, T. E. Age-related activation of microglia and astrocytes: in
vitro studies show persistent phenotypes of aging, increased proliferation, and resistance to
down-regulation. *Neurobiol Aging* **19**, 97-103, doi:10.1016/s0197-4580(97)00169-3 (1998).

71   Godbout, J. P. *et al.* Exaggerated neuroinflammation and sickness behavior in aged mice
following activation of the peripheral innate immune system. *FASEB J* **19**, 1329-1331,
doi:10.1096/fj.05-3776fje (2005).

72   Sidoryk-Wegrzynowicz, M. *et al.* Astrocytes in mouse models of tauopathies acquire early
deficits and lose neurosupportive functions. *Acta Neuropathol Commun* **5**, 89,
doi:10.1186/s40478-017-0478-9 (2017).

73   Preuss, C. *et al.* A novel systems biology approach to evaluate mouse models of late-onset
Alzheimer's disease. *Mol Neurodegener* **15**, 67, doi:10.1186/s13024-020-00412-5 (2020).

74   de Rojas, I. *et al.* Common variants in Alzheimer's disease and risk stratification by polygenic risk
scores. *Nature communications* **12**, 3417, doi:10.1038/s41467-021-22491-8 (2021).

75   Scelsi, M. A. *et al.* Genetic study of multimodal imaging Alzheimer's disease progression score
implicates novel loci. *Brain* **141**, 2167-2180, doi:10.1093/brain/awy141 (2018).

76   Kim, B. H., Nho, K., Lee, J. M. & Alzheimer's Disease Neuroimaging, I. Genome-wide association
study identifies susceptibility loci of brain atrophy to NFIA and ST18 in Alzheimer's disease.
Neurobiol Aging **102**, 200 e201-200 e211, doi:10.1016/j.neurobiolaging.2021.01.021 (2021).

77   Li, G. *et al.* Association of GALC, ZNF184, IL1R2 and ELOVL7 With Parkinson's Disease in Southern
Chinese. *Front Aging Neurosci* **10**, 402, doi:10.3389/fnagi.2018.00402 (2018).

78   Sherva, R. *et al.* Genome-wide association study of the rate of cognitive decline in Alzheimer's
disease. *Alzheimers Dement* **10**, 45-52, doi:10.1016/j.jalz.2013.01.008 (2014).

79   Chai K, L. J., Zhang X, Gu H, Cao P, Ye W, Tang H, Liu J, Chen S, Wan F, Liu GL, Shen D. ARHGDIB
Plays a Novel Role in the Braak Stages of Alzheimer's Diseases via the Immune Response
Mediated by Microglia *Pre-print: Research Square*, doi:10.21203/rs.3.rs-474315/v1 (2021).

80   Simon, M. J. *et al.* Transcriptional network analysis of human astrocytic endfoot genes reveals
region-specific associations with dementia status and tau pathology. *Sci Rep* **8**, 12389,
doi:10.1038/s41598-018-30779-x (2018).

81   Knopman, D. S. *et al.* Alzheimer disease. *Nat Rev Dis Primers* **7**, 33, doi:10.1038/s41572-021-
00269-y (2021).

82   Guyenet, S. J. *et al.* A simple composite phenotype scoring system for evaluating mouse models
of cerebellar ataxia. *J Vis Exp*, doi:10.3791/17871787 [pii] (2010).

83   Lehallier, B. *et al.* Undulating changes in human plasma proteome profiles across the lifespan.
Nat Med **25**, 1843-1850, doi:10.1038/s41591-019-0673-2 (2019).

84   He, C., Sumpter, R., Jr. & Levine, B. Exercise induces autophagy in peripheral tissues and in the
brain. *Autophagy* **8**, 1548-1551, doi:10.4161/auto.21327 (2012).

85   Raben, N. *et al.* Suppression of autophagy in skeletal muscle uncovers the accumulation of
ubiquitinated proteins and their potential role in muscle damage in Pompe disease. *Human*
molecular genetics **17**, 3897-3908, doi:10.1093/hmg/ddn292 (2008).

86   Vainshtein, A., Grumati, P., Sandri, M. & Bonaldo, P. Skeletal muscle, autophagy, and physical
activity: the menage a trois of metabolic regulation in health and disease. *J Mol Med (Berl)* **92**,
127-137, doi:10.1007/s00109-013-1096-z (2014).

87   Bostrom, P. *et al.* A PGC1-alpha-dependent myokine that drives brown-fat-like development of
white fat and thermogenesis. *Nature* **481**, 463-468, doi:10.1038/nature10777 (2012).

88   Otaka, N. *et al.* Myonectin Is an Exercise-Induced Myokine That Protects the Heart From
Ischemia-Reperfusion Injury. *Circ Res* **123**, 1326-1338, doi:10.1161/CIRCRESAHA.118.313777
(2018).

89   Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging: An
expanding universe. *Cell* **186**, 243-278, doi:10.1016/j.cell.2022.11.001 (2023).

90   Yu, D. *et al.* The adverse metabolic effects of branched-chain amino acids are mediated by
isoleucine and valine. *Cell Metab* **33**, 905-922 e906, doi:10.1016/j.cmet.2021.03.025 (2021).

91   Adler, D. H. *et al.* Characterizing the human hippocampus in aging and Alzheimer's disease using
a computational atlas derived from ex vivo MRI and histology. *Proc Natl Acad Sci U S A* **115**,
4252-4257, doi:10.1073/pnas.1801093115 (2018).

92   Ivanisevic, J. *et al.* Metabolic drift in the aging brain. *Aging (Albany NY)* **8**, 1000-1020,
doi:10.18632/aging.100961 (2016).

93   Ximerakis, M. *et al.* Single-cell transcriptomic profiling of the aging mouse brain. *Nat Neurosci*
22, 1696-1708, doi:10.1038/s41593-019-0491-3 (2019).

94   New, J. & Thomas, S. M. Autophagy-dependent secretion: mechanism, factors secreted, and
disease implications. *Autophagy* **15**, 1682-1693, doi:10.1080/15548627.2019.1596479 (2019).

95   Chow, L. S. *et al.* Exerkines in health, resilience and disease. *Nat Rev Endocrinol* **18**, 273-289,
doi:10.1038/s41574-022-00641-2 (2022).

96   Searle, B. C. *et al.* Chromatogram libraries improve peptide detection and quantification by data
independent acquisition mass spectrometry. *Nature communications* **9**, 5128,
doi:10.1038/s41467-018-07454-w (2018).

97   Pino, L. K., Just, S. C., MacCoss, M. J. & Searle, B. C. Acquiring and Analyzing Data Independent
Acquisition Proteomics Experiments without Spectrum Libraries. *Mol Cell Proteomics* **19**, 1088-
1103, doi:10.1074/mcp.P119.001913 (2020).

98   Amodei, D. *et al.* Improving Precursor Selectivity in Data-Independent Acquisition Using
Overlapping Windows. *J Am Soc Mass Spectrom* **30**, 669-684, doi:10.1007/s13361-018-2122-8
(2019).

99   Kall, L., Canterbury, J. D., Weston, J., Noble, W. S. & MacCoss, M. J. Semi-supervised learning for
peptide identification from shotgun proteomics datasets. *Nat Methods* **4**, 923-925,
doi:10.1038/nmeth1113 (2007).

100   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47, doi:10.1093/nar/gkv007 (2015).

101   Smyth, G. K. Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3, doi:10.2202/1544-6115.1027
(2004).

102   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res* **13**, 2498-2504, doi:10.1101/gr.1239303 (2003).

1230

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5