Diverse regulatory pathways modulate "bet hedging" of competence induction in

epigenetically-differentiated phase variants of *Streptococcus pneumoniae*

Min Jung Kwun[1], Alexandru V. Ion[1], Marco R. Oggioni[2,3], Stephen D. Bentley[4], Nicholas J.

Croucher[1,*]

[1] MRC Centre for Global Infectious Disease Analysis, Sir Michael Uren Hub, White City

Campus, Imperial College London, London, W12 0BZ, UK

[2] Department of Genetics, University of Leicester, University Road, Leicester, LE1 7RH

[3] Dipartimento di Farmacia e Biotecnologie, Università di Bologna, Via Irnerio 42, 40126

Bologna, Italy

[4] Parasites & Microbes, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton,

Cambridge, CB10 1SA, UK

Abstract

Despite enabling *Streptococcus pneumoniae* to acquire antibiotic resistance and evade

vaccine-induced immunity, transformation occurs at variable rates across pneumococci.

Phase variants of isolate RMV7, distinguished by altered methylation patterns driven by the

translocating variable restriction-modification (*tvr*) locus, differed significantly in their

transformation efficiencies and biofilm thicknesses. These differences were replicated when

the corresponding *tvr* alleles were introduced into an RMV7 derivative lacking the locus.

RNA-seq identified differential expression of the type 1 pilus, causing the variation in biofilm

formation, and inhibition of competence induction in the less transformable variant,

RMV7$_{domi}$. This was partly attributable to lower expression of ManLMN in RMV7$_{domi}$, which

promoted competence induction through importing *N*-acetylglucosamine. This effect was

potentiated by orthologues of the gram-negative competence regulatory machinery.

Furthermore, a phage-related chromosomal island was more active in RMV7$_{domi}$, which

inhibited transformation by increasing expression of the stress response proteins ClpP and

HrcA. However, HrcA increased competence induction in the other variant, with its effects

depending on Ca^{2+} supplementation or heat shock. Hence the heterogeneity in

transformation efficiency likely reflects the diverse signalling pathways by which it is

affected. This regulatory complexity will modulate population-wide responses to

synchronising quorum sensing signals to produce co-ordinated yet stochastic "bet hedging"

behaviour.

Competence for natural transformation was first identified in *Streptococcus pneumoniae* (the pneumococcus) in the early 20[th] century (1). Cells can be "transformed" to express a new phenotype through the acquisition of exogenous DNA, which is integrated into their genome through homologous recombination following its import through the specialised cell-encoded competence machinery (2). Transformation has played a key role in the emergence of antibiotic-resistant *S. pneumoniae*, both through generating "mosaic" alleles of core loci (3–5) and the acquisition of specialised resistance genes (6). It has also enabled vaccine evasion through recombinations affecting the capsule polysaccharide synthesis (*cps*) locus altering surface antigens (7, 8).

Despite the ability of transformation to accelerate such adaptive evolution in *S. pneumoniae*, considerable variation in the rate of diversification of strains through this mechanism persists across the species (9). Epidemiological studies have found the *r*/*m* ratio of base substitutions introduced through homologous recombination, relative to those occurring through point mutation, varies from well over 10 (7, 9) to below 0.1 (9, 10) across the species. Similarly, *in vitro* assays have identified >100-fold differences in the transformation efficiency of *S. pneumoniae* genotypes, with substantial variation even between isolates of the same serotype or strain (11–14). Many isolates are routinely found not to be transformable under standard conditions (11). This is often the consequence of integrative mobile genetic elements (MGEs) disrupting genes necessary for transformation (6, 15–17), selfishly preventing themselves from being eliminated from the chromosome (17). Yet in other non-transformable isolates, the highly-conserved competence machinery is intact (11, 18). This suggests the variation in transformation rates also reflects differences in regulation of the competence system.

The best-characterised signal inducing transformation in *S. pneumoniae* is the competence stimulating peptide (CSP) pheromone, which acts as a quorum-sensing system that signals through the ComDE two-component regulator (19). This activates early competence genes

after about ten minutes (20). These include *comX*, encoding an alternative sigma factor (21). ComX enables the RNA polymerase to recognise late competence genes (20), which feature a "combox" signal in their promoters (22, 23). This results in pneumococci entering a transient competent state around 20 minutes post-CSP induction, after which the relevant machinery is degraded (24), and the cells become temporarily refractory to induction (21).

Transformation efficiency is known to vary between isogenic pneumococci through phase variation in capsule production. Transparent colony variants produce less capsule, and are less virulent, and more transformable, than opaque colony variants (25). This short-term variation has been linked to rapid changes at the phase-variable inverting variable restriction (*ivr*) locus, encoding the conserved Type I *Spn*III restriction-modification system (RMS) and the IvrR recombinase that drives sequence inversions within the locus (26–31). These rearrangements switch the target recognition domains (TRDs) within the active HsdS specificity protein, which determines the DNA motif that is targeted by both the methylase and endonuclease activities of the system. Consequently, changes at this single locus have pleiotropic effects through altering genome-wide methylation patterns (27). These phase-variable RMSs can thereby maintain phenotypic heterogeneity within a genetically near-homogenous population (32), resulting in "bet hedging" that can increase the chances of a species surviving a changing environment (33, 34).

The second pneumococcal phase-variable Type I restriction-modification system (named *Spn*IV) (27), encoded by the translocating variable restriction (*tvr*) locus (28), varies through excision-reintegration mediated by the recombinase TvrR (35). This locus is active in almost all pneumococci, but the complement of TRDs varies between isolates, increasing the diversity of HsdS proteins, and possible methylation patterns, across the species (28, 35). This locus is inactive in the R6 laboratory isolate that is typically used to study pneumococcal competence (28, 35, 36). Here, we characterised clinical isolates in which the

tvr locus is intact, to understand how phase variation in *Spn*IV activity might contribute to

phenotypic heterogeneity in clonally-derived populations.

Results

Phase variants of *S. pneumoniae* RMV7 differ in their transformation efficiency

A previous screen for variation at the *tvr* locus identified a diverse panel of restriction-

modification variants (RMVs) (35). Four RMV isolates underwent sufficiently rapid phase

variation in culture to enable the isolation of pairs of genotypes that differed in the motif

targeted by their *Spn*IV systems, which is determined by the *hsdS* gene nearest the 5' end

of the locus (35), but were isogenic across the rest of the genome. The differences in the

arrangement of the *tvr* loci were "locked" in each variant through *tvrR* being knocked out, or

disrupted, by a selectable and counter-selectable Janus cassette marker (35, 37). These

pairs of otherwise-isogenic locked phase variants were screened for differences in their

transformation efficiency (Fig. 1A). In the RMV6 and RMV7 pairs, the variant in which the

active *tvr* locus HsdS comprised the TRDs III-i (directing the *Spn*IV system to target the

bipartite motif TGAN$_7$TATC) was found to have significantly higher transformation efficiency,

following induction by exogenous CSP, than their counterpart. These genotypes both

originate from the multidrug-resistant GPSC1 strain (38). The RMV6 variants were found to

be distinguished by a mutation in *dltD*, which commonly occurs during *in vitro* culturing (39)

and affects cell wall biology. By contrast, the serotype 19F RMV7 variants exhibited a ~100-

fold difference in their transformation rate (Fig. 1A), despite having identical *dlt* operons, and

were therefore characterised in greater detail.

This substantial difference reflected the RMV7 variant carrying the alternative form of the *tvr*

locus, with an active HsdS comprising the TRDs III-iii (which directs the *Spn*IV system to

target the motif TGAN$_7$TCC; Fig. 1B and S1), having an almost-undetectable transformation

efficiency. Culturing the wild-type isolate (RMV7$_{wt}$) over successive days demonstrated large

changes in the relative frequency of these *tvr* variants, although the less transformable

variant (RMV7 *tvrR*::Janus; henceforth, RMV7$_{domi}$, carrying *tvr*$_{domi}$) was typically dominant in

prevalence relative to the rarer, more transformable variant (RMV7 Δ*tvrR*; henceforth,

RMV7$_{rare}$, carrying *tvr*$_{rare}$; Fig. 1C). The unmodified RMV7$_{wt}$ had a transformation efficiency

similar to RMV7$_{domi}$, concordant with the relative proportions of the variants observed *in vitro*,

whereas that of RMV7$_{rare}$ was confirmed to be ~50-fold higher (Fig. 1D). This difference

could not be explained by high spontaneous mutation rates, nor by changes in the speed at

which competence for transformation was activated (Fig. S2). Therefore, RMV7$_{domi}$ and

RMV7$_{rare}$ exhibited distinctive phenotypes that correlated with their *tvr* locus arrangements.

RMV7$_{rare}$ was also significantly more adhesive to an abiotic surface (Fig. 1E), which can be

considered a proxy for biofilm formation (40). These differences in both adhesion and

transformation could be explained by RMV7$_{rare}$ being enriched for transparent phase

variants. However, microscopy found no clear difference in colony morphology between the

variants (Fig. S3). An alternative explanation for the phenotypic differences could be

mutations that occurred during genetic manipulation of the isolates (41). Alignment of the

two variants' assemblies found them to be distinguished by seven non-synonymous single

nucleotide polymorphisms and two premature stop codons outside the *tvr* locus, none of

which were within genes known to directly affect the competence machinery (Table S1).

Nevertheless, we tested how transformation was affected by mutations in RMV7$_{rare}$ that were

absent from both the RMV7$_{domi}$ and RMV7$_{wt}$ sequences. Neither a non-synonymous change

in *phoB*, encoding a phosphate-sensitive response regulator (42), nor disruption of the

transporter gene *pstS*, affect by a premature stop codon in RMV7$_{rare}$, explained the

contrasting transformation efficiencies of the variants (Fig. S4). Hence the differences

between RMV7 variants could not be explained by point mutations or alterations in

encapsulation.

To test whether the phenotypic differences were causatively associated with variation in the

*Spn*IV RMS, the *tvr* locus of RMV7$_{wt}$ was replaced with a chloramphenicol acetyltransferase

(*cat*) resistance marker to generate RMV7$_{wt}$ *tvr*::*cat*. The *tvr* loci of RMV7$_{domi}$ and RMV7$_{rare}$

(both modified by a Janus cassette within *tvrR*; see Methods) were separately introduced

into RMV7$_{wt}$ *tvr*::*cat*, generating the otherwise isogenic knock-in recombinants RMV7

tvr$_{domi}$::Janus and RMV7 *tvr*$_{rare}$::Janus, carrying the two different locked *tvr* loci (Fig. S5). An

initial characterisation of these genotypes demonstrated RMV7$_{wt}$ *tvr*::*cat* was substantially

more transformable than RMV7$_{wt}$, suggesting methylation at *tvr*$_{domi}$ motifs caused the

repression of competence induction (Fig. S5). The *tvr*$_{domi}$::Janus and *tvr*$_{rare}$::Janus mutants

reproduced the phenotypic divergence between RMV7$_{domi}$ and RMV7$_{rare}$ in both biofilm

formation and transformation (Fig. S5), providing further evidence that these differences

were driven by epigenetic variation.

The ~17-fold difference in transformation efficiency between the *tvr*$_{domi}$::Janus and

tvr$_{rare}$::Janus variants was nevertheless smaller than that measured between RMV7$_{domi}$ and

RMV7$_{rare}$. Hence the transformability of these genotypes was assayed over five independent

six-day passages, to test whether the difference between them would rise, in case any

consequences of DNA methylation were slow to emerge. However, the observed disparity in

the transformation efficiency of the *tvr*$_{domi}$::Janus and *tvr*$_{rare}$::Janus genotypes was stable

(Fig. 1F). This smaller difference may represent changes in the expression of the introduced

tvr loci, the effect of mutations outside the *tvr* locus in RMV7$_{domi}$ or RMV7$_{rare}$, else suggest

that the effect of methylation on gene expression may be indirectly mediated via effects on

nucleoid organisation. Nevertheless, both naturally-isolated and genetically-engineered pairs

of RMV7 *tvr*$_{rare}$ and *tvr*$_{domi}$ variants replicated a significant and reproducible difference in

biofilm formation and transformation efficiency.

Epigenetic effects on the induction of competence genes

To understand how the *tvr* loci caused a difference in transformation, RNA-seq was used to

quantify patterns of transcription in the recombinants RMV7 *tvr*$_{domi}$::Janus and RMV7

tvr$_{rare}$::Janus. Samples were taken pre-CSP, 10 minutes post-CSP, and 20 minutes post-

CSP from each of three biological replicates (Fig. 2). The 18 RNA samples were sequenced

as 200 nt paired-end multiplexed libraries on a single Illumina HiSeq 4000 lane. After

alignment to the RMV7$_{domi}$ genome, analysis of the RNA-seq data found the fragment length

distributions (Fig. S6) and gene expression densities (Fig. S7) were consistent across

samples (Table S2). Q-Q plots suggested a Benjamini-Hochberg corrected *q* value of 10^{-3}

was an appropriate threshold for identifying significant transcriptional variation (Fig. S8-9).

This identified 154 genes that significantly differed in their expression between the two

genotypes prior to CSP exposure, or between the pre- and post-CSP samples from the

same genotype (Fig. 2; Table S3).

Comparing the overall transcriptional patterns among datasets found the biggest separation

distinguished the post-CSP RMV7 *tvr*$_{rare}$::Janus transcriptomes from the others (Fig. S10),

suggesting a major difference in the induction of the competence system between the

variants. In RMV7 *tvr*$_{rare}$::Janus, the early competence genes showed elevated expression

10 minutes post-CSP, including a >20-fold induction of *comCDE* (Fig. S11). The late

competence genes exhibited more variable patterns of transcription (Fig. S12). Multiple

nucleotide metabolism and transporter genes (*purA*, *tadA*, *dut*, *ribF*, *adeQ*; Fig. S13) were

upregulated, and transcription of the competence-induced biofilm formation signal gene *briC*

(43) rose >10-fold (Fig. S11). Transcription of a three gene cluster encoding another

transporter, induced by CSP (44) and antimicrobial peptides (45), rose >3 fold, and

consequently was named *pieABC* (peptide-induced exporter; CDSs IONPJBJN_01324-6 in

RMV7$_{domi}$, corresponding to SP_0785-787 in TIGR4; Fig. S13, Table S3).

By contrast, CSP significantly upregulated just five genes in RMV7 *tvr*$_{domi}$::Janus: 5.7-fold

and 6.5-fold increases in the transcription of the quorum sensing genes *comCDE* and *briC*

respectively (Fig. S11), and a 2.9-fold increase in transcription of the stress response gene *csbD*. However, there was no sign of late competence genes being activated, which requires the alternative sigma factor ComX. However, expression of the *comX* gene itself can be difficult to determine through RNA-seq (Fig. S11), owing to the presence of two near-identical paralogues in pneumococcal genomes (46).

As an independent test of these transcriptional differences, qRT-PCR experiments were undertaken on the original RMV7$_{domi}$ and RMV7$_{rare}$ variants, and the control genotypes RMV7$_{wt}$ and RMV7$_{wt}$ *tvr*::*cat*. The qRT-PCR data showed genotypes of both lower (RMV7$_{wt}$ and RMV7$_{domi}$) and higher (RMV7$_{rare}$ and RMV7$_{wt}$ *tvr*::Janus) transformation efficiency up-regulated the early competence genes *comD* and *comX* in response to CSP (Fig. S14). However, the late competence genes *comEA* and *comYC* were only significantly upregulated in the more transformable genotypes. Hence the difference in transformability between the RMV7 variants was a consequence of late competence gene activation being blocked in RMV7$_{domi}$ through effects of *tvr*$_{domi}$ expression.

Pre-CSP expression differences associate elevated intracellular stress with *tvr*$_{domi}$

This difference in competence induction was likely caused by the 53 genes that significantly differed in their expression between RMV7 *tvr*$_{domi}$::Janus and *tvr*$_{rare}$::Janus prior to CSP exposure (Fig. 2; Table S3). These did not include any *cps* locus genes, which were non-significantly more highly expressed in RMV7 *tvr*$_{rare}$::Janus (Fig. S15), confirming that the elevated transformation efficiency of this genotype did not reflect an enrichment of transparent phase variants (27, 47). An analysis of the distances from protein coding sequence start codons to the nearest upstream methylation site found no general relationship between differential expression and proximal methylation for either *tvr*$_{domi}$ or *tvr*$_{rare}$ motifs (Fig. S1 and S16). Of the four cases of significantly differentially-expressed genes being close to variable methylation sites, the modification was only likely to modulate transcription initiation at the *piuABCD* operon (Fig. S17). This encodes an iron transporter,

and was >5-fold more highly expressed in RMV7$_{rare}$ (Fig. 2B; Fig. S13). However, knocking

out *piuA* did not reduce the transformation efficiency of RMV7$_{rare}$, suggesting this change

was independent of those affecting competence (Fig. S18).

Hence the differences in the pre-CSP transcriptomes are unlikely to be attributable to a

small number of promoters that are strongly affected by direct modification, consistent with

other genome-wide studies (27, 48). Instead, the differences likely represent the

consequences of chromosome-level changes in DNA conformation or nucleoid interactions

at particularly sensitive promoters (49). One notable aspect of the overall distribution of the

*Spn*IV target motifs was that the *tvr*$_{rare}$ motifs were uniformly distributed, whereas the *tvr*$_{domi}$

motifs were enriched in one segment of the chromosome (Fig. S19). Such an uneven

distribution could accentuate the effects of shifting patterns of modification.

Correspondingly, three transcriptional patterns suggested the *tvr*$_{domi}$ methylation pattern was

associated with dysregulation and increased intracellular stress. The first was the activation

of heat shock responses. Transcription of the chaperone gene *prsA* and chaperone regulator

gene *hrcA* were 3.4-fold and 4.1-fold higher in *tvr*$_{domi}$::Janus, respectively. Correspondingly,

the *groES-groEL* and *grpE-dnaK*-IONPJBJN_02152-*dnaJ* operons of the *hrcA* regulon were

also more highly expressed in *tvr*$_{domi}$::Janus, although the difference was only significant for

IONPJBJN_02152 (Fig. S21; Table S3). The *accBC-yqhY-amaP-csbD* cell wall stress

operon (50) (IONPJBJN_01032-6) was also upregulated in RMV7 *tvr*$_{domi}$::Janus (Fig. S20).

This was likely a consequence of increased expression of the *mgrA* gene, encoding a

regulator, in RMV7 *tvr*$_{domi}$::Janus, albeit this difference did not reach genome-wide

significance (Fig. S20). Consistent with this activity of MgrA, the *rlrA* islet, encoding the type

1 pilus, was approximately five-fold lower in *tvr*$_{domi}$::Janus (51).

The second indicator of stress in in RMV7 *tvr*$_{domi}$::Janus was the 3.7-fold higher transcription

of *ciaRH*, encoding a two-component system that enables cells to survive lysis-inducing

conditions (52), and is known to inhibit competence (53). The third was the increased ~1.5-

fold higher transcription of a phage-related chromosomal island (PRCI; also known as a

phage-inducible chromosomal island, or PICI), integrated adjacent to *dnaN* (PRCI*dnaN*; Fig.

S22; Table S3). This is one of two PRCIs associated with GPSC1 (7, 28), the other being

integrated near *uvrA* (Fig. 2). The regulatory mechanisms of these elements are not

thoroughly characterised (54), and in the absence of a helper prophage, it is unclear exactly

what signal may have triggered this increased activity. Given integrative MGEs are likely

under selection to reduce host cell transformability (17), the increased activity of PRCI*dnaN*

could drive inhibition of the competence machinery. Hence PRCI*dnaN* and *ciaRH* were the

primary candidates for causing the observed difference in transformability between the

RMV7 variants.

ManLMN links competence induction to carbon source metabolism

The higher expression of the *ciaRH* genes in the poorly-transformable genotypes RMV7domi

and RMV7wt, relative to the more transformable genotype RMV7rare, was confirmed by qRT-

PCR (Fig S15). CiaRH binds at least 15 promoter sequences, five of which drive the

expression of *cia*-dependent small RNAs (csRNA) that suppress the induction of

competence by inhibiting CSP production (55). Their expression is difficult to determine

using conventional RNA-seq approaches due to their size (56), but they are unlikely to affect

competence induced by exogenous CSP (57). The remaining ten drive the expression of

protein coding sequences (55) (CDSs), which were more highly expressed in RMV7

*tvr*domi::Janus compared to RMV7 *tvr*rare::Janus pre-CSP (Fig. S23). These included the

extracytoplasmic chaperone and serine protease HtrA (55, 58, 59), which can block

competence induction at low CSP concentrations through degrading extracellular CSP (58,

60). In agreement with some previous studies, elimination of *htrA* further increased the

transformability of RMV7rare (58), but the same mutation had no significant effect in RMV7wt

(Fig S24). This suggests HtrA inhibits the induction of competence, but is unlikely to explain

much of the difference between these variants. Similarly, knock out of *ciaRH* itself had little

effect in either genotype (Fig. S24), suggesting its expression was correlated with, rather

than causative of, the difference in transformation efficiency between the variants.

Among the CiaRH regulon, the most significant difference in expression between RMV7$_{rare}$

and RMV7$_{domi}$ was the ManLMN carbon source importer operon (Table S3). The *manLMN*

genes were more highly expressed in RMV7$_{rare}$, as CiaRH binds promoter sequences within

the operon (36, 55, 61, 62) and acts as a repressor (55, 63). Disrupting the *manLMN* locus

reduced the transformation efficiency of RMV7$_{rare}$ by >5-fold, while having little effect in

RMV7$_{wt}$ (Fig. 3A-B, S25). To test whether the observed transformation differences reflected

a growth defect, *manLMN*::Janus mutants of both RMV7$_{wt}$ and RMV7$_{rare}$ were cultured in the

same rich media (Fig. S26; Table S4). RMV7$_{domi}$ grew more slowly than RMV7$_{rare}$, consistent

with the former suffering greater intracellular stress. However, removal of *manLMN* had little

effect on growth in either variant, suggesting the transporter's effect on transformation was

through regulation rather than proliferation. Hence the lower expression of *manLMN* in

RMV7$_{domi}$ accounts for much of its reduced transformation efficiency relative to RMV7$_{rare}$.

ManLMN is a phosphotransferase system (PTS) transporter that can facilitate the import of

glucose, mannose, galactose, fructose, aminoglucose and *N*-acetylglucosamine (GlcNAc)

(64). Supplementation of liquid media with these carbon sources did not substantially affect

growth profiles (Fig. S26), apart from the addition of glucose causing small ManLMN-

dependent increases in the replication rate of both phase variants (Fig. S27). However,

GlcNAc supplementation increased transformation efficiency ~10-fold in RMV7$_{wt}$ (Fig. 3A)

and ~2-fold in RMV7$_{rare}$ (Fig. 3B). In both variants, this effect was dependent upon *manLMN*,

as was confirmed through disrupting, and then restoring, *manL* (Fig. S28). This potentiation

of the induction of competence by GlcNAc was also observed in the other RMV isolates (Fig.

3E). Therefore, ManLMN links carbon source availability to pneumococcal transformability.

***N*-acetylglucosamine activates competence through TfoX and YjbK**

Competence in *Vibrio cholerae* is induced GlcNAc disaccharides (65), thought to be

generated from degradation of chitin (66). This is mediated through the Transformation

Factor X (TfoX) protein (67, 68). An orthologue of this protein (TfoX$_{Hflu}$, also called Sxy) is

also central to induction of competence in *Haemophilus influenzae* (69) by 3',5'-cyclic AMP

(cAMP) signalling (70, 71). Intracellular concentrations of 3',5'-cAMP rise in many

Proteobacteria when the primary glucose PTS transporter is inactive, as the accumulation of

phosphorylated EIIA PTS subunits stimulates adenylate cyclase activity (72). Search were

undertaken for analogues of either of these pathways in *S. pneumoniae*.

An orthologue of the N terminal domain of TfoX$_{Hflu}$ was annotated, but not described, in *S.*

pneumoniae ATCC 700669 (73). In RMV7, this gene (*tfoX*$_{Spn}$; IONPJBJN_02097) is

conserved in the same genomic location as in ATCC 706669, two genes upstream of the

comEA competence operon (Fig. S29). The TfoX$_{Spn}$ amino acid sequence was predicted to

form a four-strand beta sheet flanked by alpha helices (Fig. S30), resembling the N-terminal

domain of gram-negative orthologues (Fig. S31). Disruption of *tfoX*$_{Spn}$ in RMV7$_{rare}$ both

decreased the pneumococcus' basal transformation efficiency in the absence of

supplements (Fig. S25), and eliminated the GlcNAc-induced elevation in this rate (Fig. 3C).

Restoring *tfoX* rescued RMV7$_{rare}$'s responsiveness to GlcNAc.

A gene encoding a candidate adenylate cyclase, *yjbK*, was also identified in RMV7

(IONPJBJN_01639). This protein is predicted to have a β-barrel structure (Fig. S32), as

observed for orthologous enzymes synthesising 3',5'-cAMP (Fig. S33). The *yjbK* gene could

be both disrupted, and restored, in RMV7$_{rare}$. Transformation assays with these genotypes

demonstrated the loss of YjbK reduced the transformation efficiency of RMV7$_{rare}$ both in the

absence of supplements, and following the addition of GlcNAc (Fig. 3D, S25). However, no

3',5'-cAMP signalling pathway is known in Firmicutes (74). Correspondingly, an ELISA assay

demonstrated 3',5'-cAMP levels in *S. pneumoniae* were close to the lower detection

threshold, far below those of *Escherichia coli*, and unaffected by *yjbK* disruption (Fig. S34).

Additionally, exogenous 3',5'-cAMP had no effect on transformation efficiencies in any

RMV7 genotypes (Fig. S35). Therefore, it is unlikely that YjbK's regulatory role is mediated

through 3',5'-cAMP production.

To test whether the effects of GlcNAc were the consequence of it acting as a signal, or as a

metabolic substrate, $nagA$ was disrupted in RMV7$_{wt}$ and RMV7$_{rare}$ (Fig. S36, S37). The

import of GlcNAc by ManLMN generates intracellular GlcNAc-6-phosphate, which can be

used in cell wall synthesis, or converted to glucosamine-6-phosphate by NagA.

Glucosamine-6-phosphate is also generated by the import of aminoglucose by ManLMN,

and must be converted to the glycolytic substrate fructose-6-phosphate, as both GlcNAc-6-

phosphate and glucosamine-6-phosphate are cytotoxic (75). The $nagA$::Janus mutant in the

faster-growing RMV7$_{rare}$ variant exhibited a growth defect that was exacerbated by GlcNAc

supplementation, demonstrating NagA was necessary for processing imported GlcNAc.

Similarly, a $nagA$::Janus mutant in the more GlcNAc-responsive RMV7$_{wt}$ variant could not be

transformed in the presence of GlcNAc (Fig. S37). However, exogenous aminoglucose did

not increase growth or transformation in these $nagA^-$ mutants, nor in any $nagA^+$ genotypes

(Fig. 3), confirming the effect of GlcNAc on transformation was not caused by faster growth

or glycolysis. By contrast, RMV7$_{rare}$ $tfoX$::Janus and $yjbK$::Janus mutants grew faster than

RMV7$_{rare}$ (Fig. S36), and this difference was enhanced when media were supplemented with

GlcNAc (Fig. S37), demonstrating their reduced transformation efficiencies did not reflect a

growth defect. This heightened replication rate occurred despite the disruption of $tfoX$ and

$yjbK$ not affecting $nagA$ transcription, and decreasing the expression of $manL$ slightly.

Therefore TfoX and YjbK appear to be regulators, rather than metabolic enzymes. Hence

the differential effects of GlcNAc and aminoglucose on cells is likely to reflect signalling

effects specific to GlcNAc involving TfoX, YjbK, and other proteins.

GlcNAc and ManLMN regulate pneumococcal physiology through multiple pathways

Disruption of *manLMN*, *tfoX* and *yjbK* in the laboratory genotype R6 was also found to

reduce transformation efficiency, although detecting these effects required culturing in a low-

sugar chemically-defined medium (see Methods; Fig. S38). However, disrupting *yjbK* and

tfoX in RMV7$_{wt}$ did not affect the GlcNAc-associated increase in transformation efficiency,

although transformation was notably reduced in the *yjbK*::Janus mutant (Fig. 3E). These

results suggested ManLMN affected competence through at least two pathways, at least one

of which was dependent upon TfoX and YjbK.

To test this proposed arrangement of pathways, genotypes were constructed by combining

pairs of mutations in *manLMN*, *tfoX* and *yjbK*. Transformation assays demonstrated the

double mutants in which *manLMN* was disrupted behaved similarly to the *manLMN*::Janus

single mutant (Fig. 3D). Disruption of both *tfoX* and *yjbK* did not cause a substantially

greater effect than observed in either of the corresponding single mutants. This suggested

both operated intracellularly within the same pathway (Fig. 3D), and their activity depended

on the import of molecules by ManLMN. This suggests ManLMN affects transformation

efficiency through multiple pathways, one of which involves both TfoX and YjbK, consistent

with the *tfoX*::Janus and *yjbK*::Janus mutants exhibiting similar phenotypes (Fig. 3 and S36,

S37). Hence ManLMN is a key pleiotropic regulator.

To test whether differences in ManLMN activity may also explain the difference in biofilm

formation between the RMV7 variants, adhesion of RMV7$_{rare}$ and derived *manLMN*::Janus

mutants to an abiotic surface was quantified in the presence of different carbon sources (Fig.

3F). Although disruption of *manLMN* resulted in changes to biofilm thicknesses in response

to different carbon source supplements, there was little difference in unsupplemented media.

Instead, we hypothesised that the difference was caused by the type 1 pilus, as the *rlrA* pilus

islet was more highly expressed in RMV7 *tvr*$_{rare}$::Janus, which replicated the thicker biofilm

phenotype of RMV7$_{rare}$ (Fig. 2). Disruption of the pilus structural genes (*rrgABC*), or their

activator gene (*rlrA*), reduced the surface adherence of RMV7$_{rare}$ to that of RMV7$_{wt.}$

However, the same mutation in the RMV7$_{wt}$ background had little effect (Fig. S39). Restoring

the pilus in RMV7$_{rare}$ rescued the biofilm thickness phenotype (Fig. 3C). Hence multiple

regulatory pathways underlie the individual phenotypic differences between the phase

variants.

Mobile element activation represses transformation by increasing intracellular stress

As additional regulatory pathways were likely to affect competence induction, we tested the

hypothesis that the increased activity of PRCI$_{dnaN}$ in RMV7 tvr_{domi}::Janus may also inhibit the

competence of the host cell, in order to prevent the MGE being deleted through homologous

recombination (17). The entire element was removed, either with (RMV7$_{wt}$

PRCI$_{dnaN+att}$::Janus) or without (RMV7$_{wt}$ PRCI$_{dnaN}$::Janus) the flanking *att* sites. In both cases,

a ~5-fold increase in transformation rates was observed (Fig. 4A). This implied the PRCI

inhibited the activation of the competence system. To if this was caused by a specific locus

within the MGE, four mutations were generated within the PRCI: one removing the

regulatory genes; one removing the regulatory genes and IONPJBJN_00496, a gene that

encoded a protein similar to DNA damage-inducible protein D (DinD), which inhibits RecA

activity in *E. coli* (76); one removing the replication genes; and one removing the integration,

regulatory and replication genes. However, none of these mutations had such a large effect

on transformation rates as the elimination of the entire element (Fig. 4A-B). This suggested

the inhibition of competence induction was not the consequence of a single gene product,

but instead the activity of the MGE itself.

A qRT-PCR assay was employed to test whether the deletion of PRCI$_{dnaN}$ could affect

transformation through disrupting the expression of other loci. Neither *ciaR* nor *manL*

expression was altered when the PRCI was removed (Fig. 4C), suggesting an alternative

pathway was involved. As PRCI$_{dnaN}$ caused a growth defect in RMV7$_{domi}$ (Fig. S40), it was

hypothesised that the element's activity might drive the higher expression of the stress

response proteins in tvr_{domi}::Janus (Fig. S21). Correspondingly, transcription of both the

chaperone regulator HrcA and chaperone-protease ClpP was approximately halved

following deletion of $PRCI_{dnaN}$ in $RMV7_{wt}$ (Fig. 4C). These genes were confirmed to more

highly expressed in $RMV7_{domi}$ than $RMV7_{rare}$ by qRT-PCR (Fig. S21, S41). Yet the

transcription of multiple other chaperones that were significantly more highly expressed in

$RMV7_{domi}$ (Table S3) were unperturbed by the loss of $PRCI_{dnaN}$, suggesting the mobile

element is one of multiple causes of increased intracellular stress in this variant.

The ClpP protease is known to inhibit the induction of competence (77), suggesting the

increased expression of this protein may link PRCI activation with reducing transformation.

Correspondingly, disrupting *clpP* significantly increased the transformation efficiency of

$RMV7_{wt}$ (Fig. 4B). Removing $PRCI_{dnaN}$ in this *clpP*⁻ background did not elevate the

transformation efficiency further, consistent with the effects of mobile element activation

being mediated through this protease. Disruption of *hrcA* caused a smaller, but still

significant, rise in transformation efficiency, with limited evidence of a further increase in the

double mutants *hrcA*::*cat* $PRCI_{dnaN}$::Janus and *hrcA*::Janus *clpP*::*cat* (Fig. 4B). This suggests

the inhibition of transformation driven by $PRCI_{dnaN}$ primarily reflected its stimulation of

increased ClpP activity, with HrcA also likely contributing independently. This is consistent

with the *hrcA*::Janus *clpP*::*cat* double mutant exhibiting a stronger growth defect than either

single mutant, indicating the proteins have non-redundant functions (Fig. S41). Hence

increased intracellular stress driven by MGE activity represses competence through at least

one chaperone-mediated pathway.

Activation and repression of competence by the chaperone regulator HrcA

The lower activity of $PRCI_{dnaN}$ in $RMV7_{rare}$ suggested that ClpP would be less active in this

variant. Correspondingly, disruption of *clpP* caused a smaller growth defect in this variant

(Fig. S42), and only a three-fold increase in transformation efficiency, as compared to the

>100-fold increase observed in $RMV7_{wt}$ (Fig. 5A). Yet transformation efficiency decreased in

$RMV7_{rare}$ following the disruption of *hrcA*, contrasting with the same mutation reproducibly

causing increased transformation efficiency in RMV7$_{wt}$ (Fig. 4B, 5A). Furthermore, the

RMV7$_{rare}$ *hrcA*::Janus *clpP*::*cat* double mutant also exhibited an elevated transformation

efficiency, suggesting this effect of HrcA dominated that of ClpP in this phase variant.

HrcA is unusual in having two conformations (78, 79), only one of which binds the CIRCE

DNA motif, enabling autoregulation through repressing the chaperone-encoding gene cluster

that includes *hrcA* (80). In *S. pneumoniae*, HrcA binding of CIRCE is reduced at elevated

temperatures, relieving its repression of the *dnaK* and *groEL* operons, enabling a heat shock

response (81). By contrast, Ca^{2+} ions facilitate HrcA-CIRCE motif binding, inhibiting

chaperone expression (79). Both low Ca^{2+} concentrations and extreme temperatures inhibit

the induction of competence (82). Hence the divergent effects in the phase variants could

reflect the two conformations of HrcA having different effects on the regulation of

transformation.

The DNA-binding conformation of HrcA was active in RMV$_{rare}$, as *dnaK* and *groEL*

expression was increased in the *hrcA*::Janus genotype (Fig. S43). There was no change in

transcription of *clpP*, which is outside the HrcA regulon (Fig. S41). Increasing the proportion

of the DNA-binding conformation through supplementation with CaCl$_2$ increased the

transformation efficiency of RMV7$_{rare}$ in a dose-dependent manner (Fig. 5B). This response

was lost in the *hrcA^{-}* genotype, and regained when *hrcA* was restored (Fig. 5B). These

effects could be reproduced following gene disruption and restoration in *S. pneumoniae* R6

(Fig. S44). Hence HrcA aids the activation of transformation when adopting the DNA-binding

conformation facilitated by Ca^{2+} ions.

As Ca^{2+} increases HrcA's DNA binding ability, the regulation of transformation seemed likely

to occur through altering transcription. Quantifying the expression of *hrcA* post-CSP with and

without CaCl$_2$ supplementation confirmed the ion concentrations added altered HrcA's

autorepressive activities (Fig. 5C), and decreased expression of the regulon representative

dnaK (Fig. S45). In contrast to normal growth, the reduction in *hrcA* transcription was

associated with slightly increased *clpP* expression, as a potential compensatory mechanism

(Fig. S41). However, these changes were primarily manifest 20-60 min post-CSP

administration, suggesting the transcriptional effects were too slow to affect competence

induction. Correspondingly, neither the early competence gene *comX*, nor the late

competence gene *comEA*, showed a decreased response to CSP in the *hrcA*::Janus mutant

(Fig. 5D). In contrast, strong suppression of *comEA* transcription was evident in the

manLMN mutant. This suggests HrcA regulates competence through protein-protein

interactions.

As both ClpP and HrcA have roles in thermotolerance, to test whether they caused the

reduction in transformation efficiency associated with elevated temperatures, the

transformation efficiencies of RMV7$_{rare}$ and RMV7$_{wt}$ *hrcA*$^-$ and *clpP*$^-$ mutants were compared

at 40 °C (Fig. 5A). This heat shock caused a growth defect (Fig. S42), and decreased

transformation efficiency, in both RMV7$_{rare}$ and RMV7$_{wt}$. The disruption of *clpP* had no

significant effect on transformation efficiency in either variant under these conditions,

whereas the disruption of *hrcA* significantly increased transformation efficiency in both phase

variants after the heat shock (Fig. 5A). Restoration of *hrcA* reversed the phenotypic change

caused by the gene disruption (Fig. S46). This is consistent with HrcA repressing the

induction of competence when intracellular stresses shift the protein away from its DNA-

binding conformation. This could also explain the phenotypes of the double mutants. The

more stressed RMV7$_{wt}$ *hrcA*::Janus *clpP*::*cat* genotype was not transformable, as it exhibited

a severe growth defect. By contrast, the RMV7$_{rare}$ *hrcA*::Janus *clpP::cat* double mutant

transformed and grew at similar rates to the *hrcA*::Janus single mutant, suggesting a lower

dependence on ClpP in the absence of a highly-active PRCI (Fig. S41). Hence HrcA is a

pleiotropic regulator that can activate competence in healthy cells, but represses it in

response to stress.

Independent pathways limit the competent cell subpopulation

To test whether ManLMN and HrcA separately contributed to the difference between the

variants, the transformation efficiency of the RMV7$_{rare}$ *hrcA*::Janus *manLMN*::*cat* double

mutant was compared with that of the progenitor genotype, and the single mutants (Fig. 5D).

This demonstrated an approximately five-fold decrease in transformation for each single

mutant, and a ~25-fold reduction for the double mutant. This is consistent with HrcA and

ManLMN both regulating competence independently.

Experiments with two unlinked resistance markers were used to test whether the difference

in competence between variants reflected a uniform reduction in DNA import across cells, or

an alteration in the fraction of cells in which competence was induced. The excess of double

mutants, relative to the expected frequency inferred from the single mutants (Fig. S47),

demonstrated the latter explanation accounted for the distinct behaviours of the variants.

Most bacteria remained recalcitrant to CSP in both variants: with GlcNAc supplementation, it

was estimated that 1-2% of the RMV7$_{rare}$ population became competent for transformation,

whereas 0.5% of the RMV7$_{wt}$ population did under the same conditions (Fig. S47). Hence

HrcA and ManLMN independently changed the probability of an individual cell entering the

competent state.

Discussion

This analysis highlights important challenges in the functional genomic characterisation of

clinical pathogen isolates. The RMV7$_{domi}$ and RMV7$_{rare}$ phase variants differed by few

polymorphisms, and have the same gene content, with the key differences corresponding

reversible DNA excision-reintegration dynamics and alterations to methylation. Yet they

exhibited distinct phenotypes that confounded the interpretation of multiple genetic

experiments. These differences were attributed to the methylation patterns driven by the

*Spn*IV system, as the transformation efficiency of RMV7$_{wt}$ increased when the *tvr* locus was

removed, and fell when the *tvr*$_{domi}$::Janus locus was reinstated. Furthermore, the phenotype

could not be attributed to mutations outside of the the *tvr* locus. Yet the mechanism linking

the epigenetic cause with the phenotypic consequences was difficult to establish, as the

effects of DNA methylation were not primarily manifest at genes with modified bases in their

promoters. This is consistent with the effects of *Spn*III methylation at the capsule

polysaccharide synthesis locus, despite the lack of nearby modification sites (27, 30), and

genome-wide analyses of the effects of methylation in other species (48). Instead, it is likely

that afflicted genes have an intrinsic sensitivity to intracellular perturbations. For instance,

the contribution of the type 1 pilus to biofilm formation were only detectable in RMV7$_{rare}$,

while a previous study found the effects of the pilus on the same phenotype differed

between a wild-type bacterium and an unencapsulated derivative (83). Hence the substantial

impact of methylation variation on competence induction suggests a sensitivity to small

genetic, epigenetic and physiological changes that likely underlies its heterogeneity across

populations (9, 10, 84), and over the history of individual strains (8, 85).

This susceptibility to variation is likely symptomatic of the many pathways that regulate this

phenotype (Fig. 6). One of the important signals identified in this analysis was the availability

of GlcNAc, likely the most abundant non-glucose carbon source in the nasopharyngeal

mucosa, reaching concentrations similar to the supplements in this work (86). GlcNAc can

be liberated from host mucins by pneumococcal glycosylases (87), and used as a carbon

and nitrogen source either for growth or metabolism, making it a highly informative signal of

nutrient availability and cell physiology (88). Hence GlcNAc-6-phosphate is a regulatory

molecule recognised by proteins in some species (89), including *V. cholerae* (90), in which

GlcNAc also modulates the induction of competence by a quorum sensing signal (Fig. 6).

This analysis identified similarity with components of the *V. cholerae* GlcNAc-signalling

pathways in pneumococci, including TfoX, found in many bacterial phyla (Fig. S48), and

YjbK, which belongs to a recently-defined subset of CYTH proteins (91, 92) of unknown

function in gram-positive bacteria (74, 93). Despite their highly-conserved nature, the

corresponding genes were not substantially upregulated by CSP (Fig. S49). Hence these

proteins are likely to regulate multiple aspects of pneumococcal physiology, rather than

being specific regulators of transformation.

The effects of TfoX and YjbK were dependent on the primary glucose transporter and central

metabolic regulator ManLMN (94). As ManLMN is the only effective route by which GlcNAc

can be imported (64), signalling by this molecule is limited by the competence regulator

CiaRH, but not subject to carbon catabolite repression. Orthologues of ManLMN serve as

the main glucose transporter across many Firmicutes, including other streptococci,

Lactococcus lactis and *Listeria monocytogenes* (64), and the transporter has been

associated with regulation of biofilm formation and transformability in *Streptococcus mutans*

(95) suggesting its signalling role is likely to be common among Firmicutes. Hence the

heterogeneity of pneumococcal competence induction partly reflects highly-conserved

metabolic signalling networks intervening in competence-specific pathways.

HrcA is another widely-conserved key regulator of pneumococcal physiology that this

analysis found to regulate competence induction. This protein is sensitive to physiologically-

relevant concentrations of $CaCl_2$ (96, 97), the most common ionic compound in the

nasopharyngeal mucosa (86). Hence rather than Ca^{2+} aiding the translocation of DNA

molecules across the plasma membrane, as suggested previously (98), HrcA appears to

mediate a rare example of Ca^{2+} signalling in bacteria (79, 99). Another unusual aspect of

HrcA's activity is that its two conformations appear to have opposing effects on competence

activation, resulting in the regulator's phenotype depending on the epigenetic context of the

cell. Hence HrcA is able to modulate competence induction through integrating information

on intracellular stress and extracellular ion concentrations. The mechanism by which this

was achieved is not clear, but is more likely to involve protein-protein interactions than DNA

binding, given the slow transcriptional responses to Ca^{2+} administration, and the lack of

change in competence gene expression in an $hrcA^-$ genotype. This contrasts with the

independent effects of ManLMN, which limited the induction of late competence genes by

ComX, meaning general regulators of cell biology affect multiple steps of the competence

regulatory cascade (Fig. 6).

The structure of this regulatory network can help explain the paradox of two common, but

contrasting, aspects of competence regulation (100): quorum sensing, which drives

coordinated responses, and bet hedging, which underlies population-wide heterogeneity. In

isolated cells, bet hedging could result from intrinsic noise, the variation in gene activity

reflecting the inherent stochasticity of transcription and translation (101). Yet competent

pneumococci increase their production of CSP, propagating induction to neighbouring cells

(102), tending to homogenise the population-wide response. Maintaining heterogeneity

therefore requires cells within a clonally-related population, adapted to the stable niche of

the nasopharynx, stochastically differ in their susceptibility to the quorum sensing signal.

The examples of HrcA and ManLMN demonstrate how this is achieved by pneumococci.

Firstly, both leverage variation in cell clusters' microenvironments and intracellular

physiology (101) as sources of stochasticity, or extrinsic noise (103). Secondly, both

mechanisms act on steps of the regulatory cascade that do not affect CSP production,

meaning their effects occur independently within cells, and are not propagated intercellularly.

Thirdly, both act on independent steps of the cascade, rather than being integrated into a

common mechanism, enabling them to have uncorrelated, independent effects on

competence induction. Fourthly, although temperature, and the availability of Ca^{2+} and

GlcNAc, substantially altered the probability of competence induction, none alone had a

large enough effect to risk the entire population becoming competent, maintaining

heterogeneity in the population. Hence bet hedging can emerge as an apparently random

output of combining multiple noisy signals, with the phase-variable *Spn*III and *Spn*IV

restriction-modification systems further potentiating such variation (27, 57). The function of

CSP is therefore not to homogenise the population, but to coordinate the induction of this

transient state in a subset of pneumococci.

This complex regulation in *S. pneumoniae* is similar to that in *Bacillus subtilis* (104), as well

as *Vibrio cholerae* (105) and some other gram-negative bacteria (106). These distantly-

related species all have strongly-inducing quorum sensing signals that rapidly induce a

transient competence state in a subset of bacteria, with coordinated release of DNA from

conspecific cells through fratricide or cannibalism (107–109). In each case, the intercellular

signalling is modulated by multiple extracellular stimuli, although the signals themselves vary

between the bacteria, likely driven by their divergent ecologies. Hence *V. cholerae* responds

to chitin (66), while *B. subtilis* induces competence under starvation conditions (108),

whereas this work demonstrates that pneumococcal competence is favoured in healthy

bacteria, replete with host-derived nitrogen, carbon and ion sources. Instead, it is the

complex structure of the regulatory network that is shared, enabling environmental variation

and stochasticity of signalling to modulate responses to quorum sensing. Hence some

naturally transformable bacteria appear to have convergently evolved noisy regulatory

systems that stochastically segregate populations into donors and recipients, thereby

enabling the efficient transfer of DNA during a transient period of competence.

Other naturally transformable bacteria are either constitutively competent, in the case of

some *Neisseria* species (110), or do not employ quorum sensing, as appears to be the case

for *H. influenzae* (111). However, the import of DNA is constrained to molecules containing

DNA uptake sequences (DUSs) in these bacteria (110, 112). Therefore it has been

proposed that the purpose of the transient nature of competence by quorum sensing is to

synchronise the release and acquisition of DNA from conspecific bacteria, thereby serving

as an alternative mechanism to DUSs for ensuring imported genetic material comes from

close relatives (113). Hence it is highly unlikely the competence system primarily functions to

acquire nucleic acids as a source of nutrients (114–116), as both common types of

competence system limit their imported DNA to sequences that are sufficiently closely-

related to be integrated through homologous recombination. Rather, in species regulating

competence using quorum sensing, multiple signals are likely used to ensure both the

coordination of induction, and the emergence of population-level heterogeneity. Hence the

variable nature of species-wide transformation efficiency represents the delicate balance

between chaos and order necessary for the synchronised bet hedging that characterises

competence in many bacterial species.

Methods and materials

Cell culture

Genotypes used in this study are described in Table S5. Unless otherwise stated,

encapsulated *S. pneumoniae* were cultured statically at 35 °C in 5% CO_2. Culturing on solid

media used Todd-Hewitt broth supplemented with 0.5% yeast extract and 1.5% agar

(Sigma-Aldrich). Media were supplemented antibiotics for selection of mutated genotypes:

rifampicin (Fisher Scientific) at 4 μg mL^{-1}; kanamycin (Sigma-Aldrich) at 400 μg mL^{-1}, or

chloramphenicol (Sigma-Aldrich) at 4 μg mL^{-1}. Phase contrast microscopy of colonies used a

Leica DFC3000 G microscope.

Unless otherwise specified, culturing in liquid media used 10 mL of a mixture of a 2:3 ratio of

Todd-Hewitt broth (Sigma-Aldrich) with 0.5% yeast extract (Sigma-Aldrich), and Brain-Heart

Infusion media (Sigma-Aldrich); this referred to as "mixed" media. Transformation

experiments with *S. pneumoniae* R6 derivatives used a chemically-defined medium,

consisting of disodium β-glycerophosphate (20 g L^{-1}; Sigma-Aldrich), sodium pyruvate (0.1 g

L^{-1}; Fluorochem), choline (0.001 g L^{-1}; Alfa Aesar), cysteine (0.4 g L^{-1}; Tokyo Chemical

Industry UK), glucose (3.8 mM; Sigma-Aldrich) and galactose (12 mM; Sigma-Aldrich).

Carbon source supplements were added to liquid media at a final concentration of 30 mM,

unless otherwise specified.

Growth curves and surface adhesion assays

To measure growth curves, 2×10^4 cells from titrated frozen stocks were grown in mixed

liquid media in 96-well microtiter plates at 35°C with 5% CO_2 for 20 h. Measurements of

OD_{600} were taken at 30 min intervals over 16 hours using a FLUOstar Omega microplate

reader (BMG LABTECH). Three replicate wells were assayed for each tested genotype in

each experiment. The R package growthcurver was used for the inference of carrying

capacity, K, and replication rate, r (117). For measuring adhesion to abiotic surfaces, at the

end of the growth curve incubation, the microtitre plate was submerged in water and dried

for 10 min. Then 125 μl of a freshly-diluted 0.1% crystal violet solution (Scientific Laboratory

Supplies) was added to each well, followed by incubation for 30 minutes at room

temperature. Each well was then washed by repeatedly submerging the plate in water to

remove excess crystal violet. The plate was incubated at room temperature in an inverted

position for four hours. Subsequently, 125 μl of 30% acetic acid (Honeywell) was added to

each well. Adherence was quantified as OD_{550} across replicate wells, measured by a

FLUOstar Omega plate reader.

Mutagenesis

Disruption of genes for directed mutagenesis required the PCR amplification of 0.8-1 kb

regions flanking the gene of interest, using the oligonucleotide sequences listed in Table S6.

For each gene, the upstream region was amplified with the oligonucleotides labelled with the

gene name and the suffixes "Up_For" and "Up_Rev_ApaI", with the latter adding an *Apa*I

site to the 3' of the amplicon. The downstream region was amplified with the

oligonucleotides labelled with the gene name and the suffixes "Down_For_BamHI" and

"Down_Rev", with the former adding a *Bam*HI site to the 5' of the amplicon. The

corresponding antibiotic resistance markers were amplified with oligonucleotides that added

flanking *Bam*HI and *Apa*I restriction enzyme sites: Janus_For_ApaI and Janus_Rev_BamHI

for the Janus cassette, or Cat_For_ApaI and Cat_Rev_ApaI for the *cat* chloramphenicol

resistance marker (35). PCR products were digested with the appropriate restriction

enzymes (Promega) at 35 °C for 2-4 hours, and then ligated to the appropriate antibiotic

marker using T4 DNA ligase (Invitrogen). Ligation mixtures were used as templates to

amplify correctly-ligated constructs through PCR using the "Up_For" and "Down_Rev"

oligonucleotides, which generated the amplicons used for mutagenesis through

transformation of competent pneumococcal cells.

Restoration of genes disrupted using the Janus cassette depended on all genotypes in

which mutations were generated originally being resistant to streptomycin (Table S5), owing

to a mutation in *rpsL* (118). Mutants were isolated in which the Janus cassette restored

susceptibility to streptomycin. This enabled the Janus cassette to be removed through

transformation with PCR amplicons, encoding the intact gene, generated from the original

genotypes with the "Up_For" and "Down_Rev" oligonucleotides, using selection for

streptomycin-resistant cells. As intragenomic recombination generates false positive

streptomycin-resistant genotypes independently of transformation, identifying true positive

recombinants in which genes were restored through eliminating the Janus cassette was only

feasible in the more transformable RMV7$_{rare}$ variant. PCR amplification was used to ensure

the complete removal of the cassette.

The exception to this approach was the construction of RMV7 *tvr*$_{rare}$::Janus and RMV7

tvr$_{domi}$::Janus. Both genotypes derived from the RMV7$_{wt}$ *tvr*::*cat* mutant, which was

transformed with DNA from RMV7$_{domi}$ or RMV7 *tvr*$_{rare}$ *tvrR*::Janus, followed by selection on

plates supplemented with kanamycin. The RMV7 *tvr*$_{rare}$ *tvrR*::Janus genotype was the

original locked phase variant from which RMV7$_{rare}$ was derived (35), through removal of the

Janus cassette. Using RMV7 *tvr*$_{rare}$::Janus as the donor in this transformation allowed

kanamycin selection to be used to isolate transformed genotypes that had acquired the

modified *tvr* locus. Notably, the original isolation of different phase variants through

disruption of *tvrR* used different constructs, and therefore RMV7$_{domi}$ was locked through the

replacement of 681 bp from the 3' end of *tvrR*, whereas RMV7 *tvr*rare *tvrR*::Janus was locked

through the replacement of 267 bp from the 5' end of the gene (Fig. 1). Hence the

comparison of RNA-seq data across the two variants identified differential levels of

sequence read mapping to *tvrR* (Fig. 2).

Transformation assays

One milliliter of bacterial culture was collected at an optical density of 600 nm (OD_{600})

between 0.15 and 0.25. Cells were then incubated for 2 hours at 35 °C with 5 µl of 500 mM

$CaCl_2$ (Sigma-Aldrich), 250 ng of competence stimulating peptide 1 (CSP-1; Cambridge

Bioscience Ltd) and 100 ng of the purified *rpoB* gene, containing a base substitution that

conferred resistance to rifampicin (118). In experiments using carbon source supplements,

these were added at a final concentration of 33 mM. After two hours of incubation at 35 °C, a

volume of between 1 and 200 µl of the transformed culture was spread on an agar plate

supplemented with rifampicin. For precise quantification of transformation frequencies, 1 µl

of a 10^3-fold dilution of the same culture was spread on a non-selective plate in parallel.

Colonies were counted after 24 hours of incubation at 35 °C with 5% CO_2.

A logistic curve was used to analyse the relationship between transformation efficiency (in

transformants per 10^4 colony forming units), *t*, and the concentration of the $CaCl_2$, *x*. The

fitted function was:

21
$$t = \frac{a}{1 + e^{-b(x-c)}}$$

22   The values of the variables *a*, *b* and *c* were estimated using the Levenberg-Marquardt

23   nonlinear least-squares algorithm in the minpack R package (119), using the starting values

24   of 10 (increased to 500 for *S. pneumoniae* R6), 1 and 0.5, respectively. The confidence

25   intervals were calculated through refitting the function to 999 bootstrapped samples using

26   the car R package (120).

Estimation of the frequency of competence induction

The calculation of the proportion of cells in which competence was induced required the

transformation of cultures with two independent selectable markers. The frequency of

transformation with each single marker, f, can be expressed as the product of the proportion

of pneumococci that are competent for transformation, τ, and the probability of acquisition of

each marker: p_{rif} for rifampicin, and p_{kan} for kanamycin. Hence the product of these two

transformation efficiencies, P_{double}, is:

$$P_{double} = (\tau p_{rif})(\tau p_{kan}) = \tau^2 p_{rif} p_{kan}$$

The observed frequency of double mutants, O_{double}, can be expressed as the product of the

probability a cell is competent for transformation, p_{rif} and p_{kan}:

$$O_{double} = \tau p_{rif} p_{kan}$$

Therefore the proportion of bacteria that are competent for transformation within a culture

can be calculated as:

$$\frac{P_{double}}{O_{double}} = \frac{\tau^2 p_{rif} p_{kan}}{\tau p_{rif} p_{kan}} = \tau$$

Note that if all cells in the culture are transformable, then P_{double} = O_{double}, whereas an excess

of double mutants over the frequency expected from the single mutant frequencies is

evidence that only a subset of the population is competent for transformation.

Preparation of RNA samples and quantitative PCR

Three replicate cultures of RMV7 tvr_{domi}::Janus and RMV7 tvr_{rare}::Janus were grown in 25 mL

of mixed liquid media until they reached an OD_{600} of 0.15. A 5 mL sample of bacterial cells

was collected and 50 μL 100 ng mL^{-1} CSP1 was added to the remaining culture. Further 5

mL samples were taken from each culture 10 and 20 min post-CSP addition. Each sample

was immediately treated with 10 mL RNAprotect (Qiagen) and incubated at room

temperature for 5 min. Cell were then pelleted by centrifugation at 3,220 g for 10 min. RNA

was extracted from the washed pellets using the SV Total RNA Isolation System (Promega)

according to the manufacturer's instructions. The extracted RNA was used for RNA

sequencing or qRT-PCR.

All qRT-PCR experiments were conducted as described previously (35), using 0.2 μg of

RNA to generate cDNA with the First-Strand III cDNA synthesis kit (Invitrogen). Reactions

used the PowerUp™ SYBR™ Green Master Mix (Thermo Fisher) and the QuantStudio™ 7

Flex Real-Time PCR System (Applied Biosystems). All experiments use at least two

independent biological replicates, each of which was analysed with three technical

replicates.

Gene expression was quantified used standard curves. Genomic DNA was diluted to copy

numbers of $3x10^3$, $3x10^4$, $3x10^5$ and $3x10^6$, and cycle threshold (Ct) values were calculated

for *rpoA* and the gene of interest for each. The standard curve was used to convert Ct

values into absolute copy numbers. This allowed the quantification of expression as a

relative abundance of each gene of interest relative to *rpoA*.

To quantify the dynamics of *tvr* loci during culture, three independent passages of RMV7$_{wt}$ in

liquid media were each initiated through inoculation with a single colony. Each subsequent

24 h passage was inoculated with 10 µL of the previous culture. Genomic DNA was

extracted from the initial culture, and after four and eight days of passage, using the Wizard

Genomic DNA purification kit (Promega). Quantitative PCR was used to measure the

frequency of each distinguishable allele of the *tvr* locus, using triplicate technical replicates

on each sample, as described previously (121). The absolute copy numbers of each allele

were then estimated using standard curves, thereby enabling the *tvr*$_{domi}$:*tvr*$_{rare}$ ratio to be

calculated.

Generation and analysis of RNA-seq data

RNA samples were quantified using an Agilent Bioanalyser RNA Nano Chip (Agilent

Technologies), and treated with the RiboZero® rRNA Removal Kit for Bacteria (Illumina) to

deplete rRNA. The samples were then cleaned with Agencourt RNAClean Beads (Beckman

Coulter). Sequencing libraries were generated with the NEBNext® Ultra II Directional Library

Prep Kit for Illumina (New England BioLabs), modified to use oligonucleotide sequences

appropriate for the sequencing pipelines of the Wellcome Sanger Institute. The library was

amplified through nine PCR cycles using the Kapa HiFi HotStart Ready Mix (Roche) to

generate sufficient material for sequencing. All eighteen samples were sequenced as a

multiplexed library on a single lane of a HiSeq 4000 sequencing system (Illumina),

generating 200 nt paired end reads.

The set of genes used for expression analysis were the 2,088 protein coding sequences

annotated on the *S. pneumoniae* RMV7$_{domi}$ genome (accession code OV904788), and the

81 non-coding RNAs predicted by infernal version 1.1.2 (122) using the Rfam database

(123). RNA-seq reads were aligned to these sequences using kallisto version 0.46.2 (124),

using default settings and 100 bootstraps. Differential gene expression analysis used sleuth

version 0.30 (125). Wald tests were conducted to compare the pre-CSP samples for RMV7

tvr$_{domi}$::Janus and RMV7 *tvr*$_{rare}$::Janus, and to compare the 10 min and 20 min post-CSP

samples for each genotype to the corresponding pre-CSP samples. Visualisation and

plotting of data used the genoplotR (126), circlize (127), cowplot (128), ggpubr (129) and

tidyverse (130) packages.

Quantification of 3',5'-cAMP production

Quantification of 3',5'-cAMP used the Cyclic AMP XP® assay kit (Cell Signalling

Technology). Samples were harvested in the exponential (OD$_{600}$ of 0.2) and stationary

(OD$_{600nm}$ of 0.5) phases of statically-grown *S. pneumoniae* cultures. *E. coli* DH5α was grown

in 25 mL Luria-Bertani media (Sigma-Aldrich) at 37 °C, shaken at 250 revolutions per

minute, and samples were harvested in the exponential (OD$_{600}$ of 0.3) and stationary

(OD$_{600nm}$ of 1.0) phases. Cells were pelleted through centrifugation at 3,220 g for 10 min.

The cell pellets were re-suspended in 850 μL of the kit's lysis buffer and 150 μl of lysozyme

(10 mg mL^{-1}), followed by incubation at 35 °C for 20 minutes. Cells were pelleted by

centrifugation at 9,500 g for 5 minutes. The supernatants were collected, and the cell pellets

were resuspended in 400 μl phosphate-buffered saline. The protein concentrations of all the

collected samples were adjusted to 400 μg mL^{-1} using the Qubit protein broad range assay

kits (Qiagen). The Cyclic AMP XP® kit ELISA plates were prepared according to the

manufacturer's protocols, and a 50 μl sample of supernatant from each tested culture was

loaded in each well. The OD$_{450}$ was measured using the FLUOstar Omega plate reader.

Concentrations of 3',5'-cAMP were calculated using a standard curve, according to the

manufacturer's instructions.

Analysis of motif distribution and sequence diversity

The RMV7$_{domi}$ and RMV7$_{rare}$ sequences were aligned with nucmer (131). The overall

distribution of methylation motifs was calculated using DistAMo (132). The calculation of

distances between coding sequences and motifs used biopython (133).

The *S. pneumoniae* RMV7 TfoX and YjbK proteins were aligned to orthologues from *H.*

influenzae Rd and *V. cholerae* ATCC 39315 using Muscle (134). The predicted structures

were retrieved from the AlphaFoldDB (135).

Proteins containing the TfoX N-terminal domain were identified using EMBL SMART (136).

These amino acid sequences were aligned with MAFFT (137), and a phylogeny generated

with Fasttree2 (138) using default settings. Proteins were assigned to bacterial Families

using the NCBI Taxonomy (139).

Acknowledgements

MJK, AVI and NJC were supported by the UK Medical Research Council and Department for

International Development (grant nos MR/R015600/1 and MR/T016434/1), and a Sir Henry

Dale Fellowship, jointly funded by Wellcome and the Royal Society (grant no.

104169/Z/14/A). MJK and MRO were supported by the BBSRC (grant no. BB/N002903/1).

SDB was supported by Wellcome (grant no 206194). We thank the Bespoke team at the

Wellcome Sanger Institute for generating the sequencing libraries.

Financial disclosure

NJC has consulted for Antigen Discovery Inc and Pfizer. NJC has received an investigator-

initiated award from GlaxoSmithKline.

Data availability

The genome sequence and annotation of *S. pneumoniae* RMV7$_{domi}$ is available from the

European Nucleotide Archive (ENA) with the accession code OV904788. The RNA-seq data

are available from the ENA with the accession codes listed in Table S2. The expression

values and statistical tests calculated for all analysed genes in the RNA-seq analysis are

available from FigShare, alongside the raw gel images, micrographs and the results of qRT-

PCR and microbiological experiments

(https://figshare.com/projects/Diverse_regulatory_pathways_modulate_bet_hedging_of_com

petence_induction_in_epigenetically-

differentiated_phase_variants_of_Streptococcus_pneumoniae/171060), as detailed in Table

S7.

References

1. Griffith,F. (1928) The Significance of Pneumococcal Types. *J. Hyg. (Lond).*, **27**, 113–159.

2. Johnston,C., Campo,N., Bergé,M.J., Polard,P. and Claverys,J.P. (2014) *Streptococcus*

pneumoniae, le transformiste. *Trends Microbiol.*, **22**, 113–9.

3. Dowson,C.G., Hutchison,A., Brannigan,J.A., George,R.C., Hansman,D., Linares,J.,

Tomasz,A., Smith,J.M. and Spratt,B.G. (1989) Horizontal transfer of penicillin-binding

protein genes in penicillin-resistant clinical isolates of *Streptococcus pneumoniae*. *Proc.*

Natl. Acad. Sci., **86**, 8842–8846.

4. Sibold,C., Henrichsen,J., Konig,A., Martin,C., Chalkley,L. and Hakenbeck,R. (1994)

Mosaic *pbpX* genes of major clones of penicillin-resistant *Streptococcus pneumoniae*

have evolved from *pbpX* genes of a penicillin-sensitive *Streptococcus oralis*. *Mol*

Microbiol, **12**, 1013–1023.

5. Dowson,C.G., Coffey,T.J., Kell,C. and Whiley,R.A. (1993) Evolution of penicillin

resistance in *Streptococcus pneumoniae*; the role of *Streptococcus mitis* in the

formation of a low affinity PBP2B in *S. pneumoniae*. *Mol. Microbiol.*, **9**, 635–643.

6. D'Aeth,J.C., van der Linden,M.P.G., McGee,L., de Lencastre,H., Turner,P., Song,J.-H.,

Lo,S.W., Gladstone,R.A., Sá-Leão,R., Ko,K.S., *et al.* (2021) The role of interspecies

recombination in the evolution of antibiotic-resistant pneumococci. *Elife*, **10**, e67113.

7. Croucher,N.J., Chewapreecha,C., Hanage,W.P., Harris,S.R., McGee,L., van der

Linden,M., Song,J.-H.H., Ko,K.S., de Lencastre,H., Turner,C., *et al.* (2014) Evidence for

soft selective sweeps in the evolution of pneumococcal multidrug resistance and

vaccine escape. *Genome Biol. Evol.*, **6**, 1589–1602.

8. Golubchik,T., Brueggemann,A.B., Street,T., Gertz,R.E., Spencer,C.C.A., Ho,T.,

Giannoulatou,E., Link-Gelles,R., Harding,R.M., Beall,B., *et al.* (2012) Pneumococcal

genome sequencing tracks a vaccine escape variant formed through a multi-fragment

recombination event. *Nat. Genet.*, **44**, 352–355.

9. Croucher,N.J., Finkelstein,J.A., Pelton,S.I., Mitchell,P.K., Lee,G.M., Parkhill,J.,

Bentley,S.D., Hanage,W.P. and Lipsitch,M. (2013) Population genomics of post-vaccine

changes in pneumococcal epidemiology. *Nat. Genet.*, **45**, 656–663.

10. Croucher,N.J., Mitchell,A.M., Gould,K.A., Inverarity,D., Barquist,L., Feltwell,T.,

Fookes,M.C., Harris,S.R., Dordel,J., Salter,S.J., *et al.* (2013) Dominant role of

nucleotide substitution in the diversification of serotype 3 pneumococci over decades

and during a single infection. *PLoS Genet*, **9**, e1003868.

11. Evans,B.A. and Rozen,D.E. (2013) Significant variation in transformation frequency in

Streptococcus pneumoniae. ISME J., **7**, 791–799.

12. Hsieh,Y.C., Wang,J.T., Lee,W. Sen, Hsueh,P.R., Shao,P.L., Chang,L.Y., Lu,C.Y.,

Lee,C.Y., Huang,F.Y. and Huang,L.M. (2006) Serotype competence and penicillin

resistance in *Streptococcus pneumoniae. Emerg. Infect. Dis.*, **12**, 1709–1714.

13. Joloba,M.L., Kidenya,B.R., Kateete,D.P., Katabazi,F.A., Muwanguzi,J.K., Asiimwe,B.B.,

Alarakol,S.P., Nakavuma,J.L., Bajaksouzian,S., Windau,A., *et al.* (2010) Comparison of

transformation frequencies among selected *Streptococcus pneumoniae* serotypes. *Int.*

J. Antimicrob. Agents, **36**, 124–128.

14. Yother,J., McDaniel,L.S. and Briles,D.E. (1986) Transformation of encapsulated

Streptococcus pneumoniae. J. Bacteriol., **168**, 1463–1465.

15. Croucher,N.J., Harris,S.R., Fraser,C., Quail,M.A., Burton,J., van der Linden,M.,

McGee,L., von Gottberg,A., Song,J.H., Ko,K.S., *et al.* (2011) Rapid pneumococcal

evolution in response to clinical interventions. *Science (80-.).*, **331**, 430–434.

16. Croucher,N.J., Hanage,W.P., Harris,S.R., McGee,L., van der Linden,M., de

Lencastre,H., Sá-Leão,R., Song,J.-H., Ko,K.S., Beall,B., *et al.* (2014) Variable

recombination dynamics during the emergence, transmission and 'disarming' of a

multidrug-resistant pneumococcal clone. *BMC Biol.*, **12**, 49.

17. Croucher,N.J., Mostowy,R., Wymant,C., Turner,P., Bentley,S.D. and Fraser,C. (2016)

Horizontal DNA Transfer Mechanisms of Bacteria as Weapons of Intragenomic Conflict.

PLOS Biol., **14**, e1002394.

18. Johnston,C., Martin,B., Fichant,G., Polard,P. and Claverys,J.P. (2014) Bacterial

transformation: Distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.*, **12**, 181–196.

19. Håvarstein,L.S., Coomaraswamy,G. and Morrison,D.A. (1995) An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae. Proc. Natl. Acad. Sci.*, **92**, 11140 LP – 11144.

20. Alloing,G., Martin,B., Granadel,C. and Claverys,J.-P. (1998) Development of competence in *Streptococcus pneumoniae*: pheromone autoinduction and control of quorum sensing by the oligopeptide permease. *Mol. Microbiol.*, **29**, 75–83.

21. Lee,M.S. and Morrison,D.A. (1999) Identification of a New Regulator in *Streptococcus pneumoniae* Linking Quorum Sensing to Competence for Genetic Transformation. *J. Bacteriol.*, **181**, 5004–5016.

22. Campbell,E.A., Choi,S.Y. and Masure,H.R. (1998) A competence regulon in *Streptococcus pneumoniae* revealed by genomic analysis. *Mol. Microbiol.*, **27**, 929–939.

23. Pestova,E. V and Morrison,D.A. (1998) Isolation and characterization of three *Streptococcus pneumoniae* transformation-specific loci by use of a *lacZ* reporter insertion vector. *J. Bacteriol.*, **180**, 2701–2710.

24. Sung,C.K. and Morrison,D.A. (2005) Two Distinct Functions of ComW in Stabilization and Activation of the Alternative Sigma Factor ComX in *Streptococcus pneumoniae. J. Bacteriol.*, **187**, 3052–3061.

25. Weiser,J.N. and Kapoor,M. (1999) Effect of intrastrain variation in the amount of capsular polysaccharide on genetic transformation of *Streptococcus pneumoniae*: Implications for virulence studies of encapsulated strains. *Infect. Immun.*, **67**, 3690–3692.

26. Tettelin,H., Nelson,K.E., Paulsen,I.T., Eisen,J.A., Read,T.D., Peterson,S., Heidelberg,J., DeBoy,R.T., Haft,D.H., Dodson,R.J., *et al.* (2001) Complete genome sequence of a virulent isolate of *Streptococcus pneumoniae. Science (80-.).*, **293**, 498–506.

27. Manso,A.S., Chai,M.H., Atack,J.M., Furi,L., De Ste Croix,M., Haigh,R., Trappetti,C.,

Ogunniyi,A.D., Shewell,L.K., Boitano,M., *et al.* (2014) A random six-phase switch regulates pneumococcal virulence via global epigenetic changes. *Nat. Commun.*, **5**, 5055.

28. Croucher,N.J., Coupland,P.G., Stevenson,A.E., Callendrello,A., Bentley,S.D. and Hanage,W.P. (2014) Diversification of bacterial genome content through distinct mechanisms over different timescales. *Nat. Commun.*, **5**, 5471.

29. Li,J.W., Li,J., Wang,J., Li,C. and Zhang,J.R. (2019) Molecular mechanisms of *hsdS* inversions in the *cod* locus of *Streptococcus pneumoniae*. *J. Bacteriol.*, **201**, e00581-18.

30. Li,J., Li,J.W., Feng,Z., Wang,J., An,H., Liu,Y., Wang,Y., Wang,K., Zhang,X., Miao,Z., *et al.* (2016) Epigenetic Switch Driven by DNA Inversions Dictates Phase Variation in *Streptococcus pneumoniae*. *PLoS Pathog.*, **12**, e1005762.

31. Wang,J., Li,J.W., Li,J., Huang,Y., Wang,S. and Zhang,J.R. (2020) Regulation of pneumococcal epigenetic and colony phases by multiple two-component regulatory systems. *PLoS Pathog.*, **16**, e1008417.

32. De Ste Croix,M., Vacca,I., Kwun,M.J., Ralph,J.D., Bentley,S.D., Haigh,R., Croucher,N.J. and Oggioni,M.R. (2017) Phase-variable methylation and epigenetic regulation by type I restriction-modification systems. *FEMS Microbiol. Rev.*, **41**, S3–S15.

33. Veening,J.-W., Smits,W.K. and Kuipers,O.P. (2008) Bistability, epigenetics, and bet-hedging in bacteria. *Annu. Rev. Microbiol.*, **62**, 193–210.

34. Quillin,S.J. and Seifert,H.S. (2018) *Neisseria gonorrhoeae* host adaptation and pathogenesis. *Nat. Rev. Microbiol.*, **16**, 226–240.

35. Kwun,M.J., Oggioni,M.R., De Ste Croix,M., Bentley,S.D. and Croucher,N.J. (2018) Excision-reintegration at a pneumococcal phase-variable restriction-modification locus drives within- and between-strain epigenetic differentiation and inhibits gene acquisition. *Nucleic Acids Res.*, **46**, 11438–11453.

36. Slager,J., Aprianto,R. and Veening,J.W. (2019) Refining the pneumococcal competence regulon by RNA sequencing. *J. Bacteriol.*, **201**, e00780-18.

37. Sung,C.K., Li,H., Claverys,J.P. and Morrison,D.A. (2001) An *rpsL* cassette, Janus, for

gene replacement through negative selection in *Streptococcus pneumoniae. Appl.*

Environ. Microbiol., **67**, 5190–5196.

38. Gladstone,R.A., Lo,S.W., Lees,J.A., Croucher,N.J., van Tonder,A.J., Corander,J.,

Page,A.J., Marttinen,P., Bentley,L.J., Ochoa,T.J., *et al.* (2019) International genomic

definition of pneumococcal lineages, to contextualise disease, antibiotic resistance and

vaccine impact. *EBioMedicine*, **43**, 338–346.

39. Lees,J.A., Kremer,P.H.C., Manso,A.S., Croucher,N.J., Ferwerda,B., Serón,M.V.,

Oggioni,M.R., Parkhill,J., Brouwer,M.C., van der Ende,A., *et al.* (2017) Large scale

genomic analysis shows no evidence for pathogen adaptation between the blood and

cerebrospinal fluid niches during bacterial meningitis. *Microb. Genomics*,

10.1099/mgen.0.000103.

40. Trappetti,C., Kadioglu,A., Carter,M., Hayre,J., Iannelli,F., Pozzi,G., Andrew,P.W. and

Oggioni,M.R. (2009) Sialic Acid: A Preventable Signal for Pneumococcal Biofilm

Formation, Colonization, and Invasion of the Host. *J. Infect. Dis.*, **199**, 1497–1505.

41. Li,Y., Croucher,N.J., Thompson,C.M., Trzciński,K., Hanage,W.P. and Lipsitch,M. (2015)

Identification of pneumococcal colonization determinants in the stringent response

pathway facilitated by genomic diversity. *BMC Genomics*, **16**, 369.

42. Lamarche,M.G., Wanner,B.L., Crépin,S. and Harel,J. (2008) The phosphate regulon and

bacterial virulence: a regulatory network connecting phosphate homeostasis and

pathogenesis. *FEMS Microbiol. Rev.*, **32**, 461–473.

43. Aggarwal,S.D., Eutsey,R., West-Roberts,J., Domenech,A., Xu,W., Abdullah,I.T.,

Mitchell,A.P., Veening,J.-W., Yesilkaya,H. and Hiller,N.L. (2018) Function of BriC

peptide in the pneumococcal competence and virulence portfolio. *PLOS Pathog.*, **14**,

e1007328.

44. Peterson,S.N., Sung,C.K., Cline,R., Desai,B. V, Snesrud,E.C., Luo,P., Walling,J., Li,H.,

Mintz,M., Tsegaye,G., *et al.* (2004) Identification of competence pheromone responsive

genes in *Streptococcus pneumoniae* by use of DNA microarrays. *Mol. Microbiol.*, **51**,

1051–1070.

45. Majchrzykiewicz,J.A., Kuipers,O.P. and Bijlsma,J.J.E. (2010) Generic and specific adaptive responses of *Streptococcus pneumoniae* to challenge with three distinct antimicrobial peptides, bacitracin, LL-37, and nisin. *Antimicrob. Agents Chemother.*, **54**, 440–451.

46. Johnston,C.H.G., Soulet,A.-L., Bergé,M., Prudhomme,M., De Lemos,D. and Polard,P. (2020) The alternative sigma factor σ^X mediates competence shut-off at the cell pole in *Streptococcus pneumoniae. Elife*, **9**, e62907.

47. Weiser,J.N., Bae,D., Epino,H., Gordon,S.B., Kapoor,M., A.,Z.L. and Shchepetov,M. (2001) Changes in Availability of Oxygen Accentuate Differences in Capsular Polysaccharide Expression by Phenotypic Variants and Clinical Isolates of *Streptococcus pneumoniae. Infect. Immun.*, **69**, 5430–5439.

48. Bourgeois,J.S., Anderson,C.E., Wang,L., Modliszewski,J.L., Chen,W., Schott,B.H., Devos,N. and Ko,D.C. (2022) Integration of the *Salmonella* Typhimurium Methylome and Transcriptome Reveals That DNA Methylation and Transcriptional Regulation Are Largely Decoupled under Virulence-Related Conditions. *MBio*, **13**, e03464-21.

49. Buitrago,D., Labrador,M., Arcon,J.P., Lema,R., Flores,O., Esteve-Codina,A., Blanc,J., Villegas,N., Bellido,D., Gut,M., *et al.* (2021) Impact of DNA methylation on 3D genome structure. *Nat. Commun.*, **12**, 3243.

50. Solano-Collado,V., Espinosa,M. and Bravo,A. (2012) Activator role of the pneumococcal *mga*-like virulence transcriptional regulator. *J. Bacteriol.*, **194**, 197–4207.

51. Hemsley,C., Joyce,E., Hava,D.L., Kawale,A. and Camilli,A. (2003) MgrA, an Orthologue of Mga, Acts as a Transcriptional Repressor of the Genes within the *rlrA* Pathogenicity Islet in *Streptococcus pneumoniae. J. Bacteriol.*, **185**, 6640–6647.

52. Mascher,T., Heintz,M., Zähner,D., Merai,M. and Hakenbeck,R. (2006) The CiaRH system of *Streptococcus pneumoniae* prevents lysis during stress induced by treatment with cell wall inhibitors and by mutations in *pbp2x* involved in beta-lactam resistance. *J. Bacteriol.*, **188**, 1959–1968.

53. Guenzi,E., Gasc,A. -M, Sicard,M.A. and Hakenbeck,R. (1994) A two-component signal-

transducing system is involved in competence and penicillin susceptibility in laboratory

mutants of *Streptococcus pneumoniae*. *Mol. Microbiol.*, **12**, 505–515.

54. Martínez-Rubio,R., Quiles-Puchalt,N., Martí,M., Humphrey,S., Ram,G., Smyth,D.,

Chen,J., Novick,R.P. and Penadés,J.R. (2017) Phage-inducible islands in the Gram-

positive cocci. *ISME J.*, **11**, 1029–1042.

55. Halfmann,A., Kovács,M., Hakenbeck,R. and Brückner,R. (2007) Identification of the

genes directly controlled by the response regulator CiaR in *Streptococcus pneumoniae*:

Five out of 15 promoters drive expression of small non-coding RNAs. *Mol. Microbiol.*,

66, 110–126.

56. Hu,J.F., Yim,D., Ma,D., Huber,S.M., Davis,N., Bacusmo,J.M., Vermeulen,S., Zhou,J.,

Begley,T.J., DeMott,M.S., *et al.* (2021) Quantitative mapping of the cellular small RNA

landscape with AQRNA-seq. *Nat. Biotechnol.*, **39**, 978–988.

57. Kwun,M.J., Ion,A. V, Cheng,H.-C., D'Aeth,J.C., Dougan,S., Oggioni,M.R., Goulding,D.A.,

Bentley,S.D. and Croucher,N.J. (2022) Post-vaccine epidemiology of serotype 3

pneumococci identifies transformation inhibition through prophage-driven alteration of a

non-coding RNA. *Genome Med.*, **14**, 2022.09.21.508813.

58. Cassone,M., Gagne,A.L., Spruce,L.A., Seeholzer,S.H. and Sebert,M.E. (2012) The HtrA

protease from *Streptococcus pneumoniae* digests both denatured proteins and the

competence-stimulating peptide. *J. Biol. Chem.*, **287**, 38449–38459.

59. Sebert,M.E., Palmer,L.M., Rosenberg,M. and Weiser,J.N. (2002) Microarray-based

identification of *htrA*, a *Streptococcus pneumoniae* gene that is regulated by the CiaRH

two-component system and contributes to nasopharyngeal colonization. *Infect. Immun.*,

70, 4059–4067.

60. Stevens,K.E., Chang,D., Zwack,E.E. and Sebert,M.E. (2011) Competence in

Streptococcus pneumoniae is regulated by the rate of ribosomal decoding errors. *MBio*,

2, e00071-11.

61. Dagkessamanskaia,A., Moscoso,M., Hénard,V., Guiral,S., Overweg,K., Reuter,M.,

Martin,B., Wells,J. and Claverys,J.P. (2004) Interconnection of competence, stress and CiaR regulons in *Streptococcus pneumoniae*: Competence triggers stationary phase autolysis of *ciaR* mutant cells. *Mol. Microbiol.*, **51**, 1071–1086.

62. Mascher,T., Zähner,D., Merai,M., Balmelle,N., De Saizieu,A.B. and Hakenbeck,R. (2003) The *Streptococcus pneumoniae cia* regulon: CiaR target sites and transcription profile analysis. *J. Bacteriol.*, **185**, 60–70.

63. Gómez-Mejia,A., Gámez,G. and Hammerschmidt,S. (2018) *Streptococcus pneumoniae* two-component regulatory systems: The interplay of the pneumococcus with its environment. *Int. J. Med. Microbiol.*, **308**, 722–737.

64. Bidossi,A., Mulas,L., Decorosi,F., Colomba,L., Ricci,S., Pozzi,G., Deutscher,J., Viti,C. and Oggioni,M.R. (2012) A functional genomics approach to establish the complement of carbohydrate transporters in *Streptococcus pneumoniae*. *PLoS One*, **7**, e33320.

65. Yamamoto,S., Morita,M., Izumiya,H. and Watanabe,H. (2010) Chitin disaccharide (GlcNAc)$_2$ induces natural competence in *Vibrio cholerae* through transcriptional and translational activation of a positive regulatory gene *tfoX*$_{VC}$. *Gene*, **457**, 42–49.

66. Meibom,K.L., Blokesch,M., Dolganov,N.A., Wu,C.Y. and Schoolnik,G.K. (2005) Chitin induces natural competence in *Vibrio cholerae*. *Science (80-.).*, **310**, 1824–1827.

67. Lo Scrudato,M. and Blokesch,M. (2013) A transcriptional regulator linking quorum sensing and chitin induction to render *Vibrio cholerae* naturally transformable. *Nucleic Acids Res.*, **41**, 3644–3658.

68. Yamamoto,S., Izumiya,H., Mitobe,J., Morita,M., Arakawa,E., Ohnishi,M. and Watanabe,H. (2011) Identification of a chitin-induced small RNA that regulates translation of the *tfoX* gene, encoding a positive regulator of natural competence in *Vibrio cholerae*. *J. Bacteriol.*, **193**, 1953–1965.

69. Williams,P.M., Bannister,L.A. and Redfield,R.J. (1994) The *Haemophilus influenzae sxy-1* mutation is in a newly identified gene essential for competence. *J. Bacteriol.*, **176**, 6789–6794.

70. Chandler,M.S. (1992) The gene encoding cAMP receptor protein is required for

competence development in *Haemophilus influenzae* Rd. *Proc. Natl. Acad. Sci.*, **89**, 1626 LP – 1630.

71. Dorocicz,I.R., Williams,P.M. and Redfield,R.J. (1993) The *Haemophilus influenzae* adenylate cyclase gene: cloning, sequence, and essential role in competence. *J. Bacteriol.*, **175**, 7142–7149.

72. Wu,R., Zhao,M., Li,J., Gao,H., Kan,B. and Liang,W. (2015) Direct regulation of the natural competence regulator gene *tfoX* by cyclic AMP (cAMP) and cAMP receptor protein (CRP) in Vibrios. *Sci. Rep.*, **5**, 14921.

73. Croucher,N.J., Walker,D., Romero,P., Lennard,N., Paterson,G.K., Bason,N.C., Mitchell,A.M., Quail,M.A., Andrew,P.W., Parkhill,J., *et al.* (2009) Role of conjugative elements in the evolution of the multidrug-resistant pandemic clone *Streptococcus pneumoniae*[Spain23F] ST81. *J Bacteriol*, **191**, 1480–1489.

74. Zhang,Y., Agrebi,R., Bellows,L.E., Collet,J.-F., Kaever,V. and Gründling,A. (2017) Evolutionary Adaptation of the Essential tRNA Methyltransferase TrmD to the Signaling Molecule 3',5'-cAMP in Bacteria. *J. Biol. Chem.*, **292**, 313–327.

75. Bernheim,N.J. and Dobrogosz,W.J. (1970) Amino sugar sensitivity in *Escherichia coli* mutants unable to grow on *N*-acetylglucosamine. *J. Bacteriol.*, **101**, 384–391.

76. Uranga,L.A., Balise,V.D., Benally,C. V, Grey,A. and Lusetti,S.L. (2011) The *Escherichia coli* DinD protein modulates RecA activity by inhibiting postsynaptic RecA filaments. *J. Biol. Chem.*, **286**, 29480–29491.

77. Robertson,G.T., Ng,W.-L., Foley,J., Gilmour,R. and Winkler,M.E. (2002) Global Transcriptional Analysis of *clpP* Mutations of Type 2 *Streptococcus pneumoniae* and Their Effects on Physiology and Virulence. *J. Bacteriol.*, **184**, 3508–3520.

78. Roncarati,D., Danielli,A. and Scarlato,V. (2014) The HrcA repressor is the thermosensor of the heat-shock regulatory circuit in the human pathogen *Helicobacter pylori*. *Mol. Microbiol.*, **92**, 910–920.

79. Kwon,H.Y., Kim,S.N., Pyo,S.N. and Rhee,D.K. (2005) Ca^{2+}-dependent expression of the CIRCE regulon in *Streptococcus pneumoniae*. *Mol. Microbiol.*, **55**, 456 – 468.

80. Zuber,U. and Schumann,W. (1994) CIRCE, a novel heat shock element involved in

regulation of heat shock operon *dnaK* of *Bacillus subtilis*. *J. Bacteriol.*, **176**, 1359–1363.

81. Kim,S.N., Bae,Y.G. and Rhee,D.K. (2008) Dual regulation of *dnaK* and *groE* operons by

HrcA and Ca++ in *Streptococcus pneumoniae*. *Arch. Pharm. Res.*, **31**, 462–467.

82. Fox,M.S. and Hotchkiss,R.D. (1957) Initiation of bacterial transformation. *Nature*, **179**,

1322–1325.

83. Muñoz-Elías,E.J., Marcano,J. and Camilli,A. (2008) Isolation of *Streptococcus*

pneumoniae biofilm mutants and their characterization during nasopharyngeal

colonization. *Infect. Immun.*, **76**, 5049–5061.

84. Chewapreecha,C., Harris,S.R., Croucher,N.J., Turner,C., Marttinen,P., Cheng,L.,

Pessia,A., Aanensen,D.M., Mather,A.E., Page,A.J., *et al.* (2014) Dense genomic

sampling identifies highways of pneumococcal recombination. *Nat. Genet.*, **46**, 305–

309.

85. Mostowy,R., Croucher,N.J., Hanage,W.P., Harris,S.R., Bentley,S. and Fraser,C. (2014)

Heterogeneity in the Frequency and Characteristics of Homologous Recombination in

Pneumococcal Evolution. *PLoS Genet.*, **10**, e1004300.

86. Ruhluel,D., O'Brien,S., Fothergill,J.L. and Neill,D.R. (2022) Development of liquid culture

media mimicking the conditions of sinuses and lungs in cystic fibrosis and health.

F1000Research, **11**, 1007.

87. Weiser,J.N., Ferreira,D.M. and Paton,J.C. (2018) *Streptococcus pneumoniae*:

Transmission, colonization and invasion. *Nat. Rev. Microbiol.*, **16**, 355–367.

88. Komatsuzawa,H., Fujiwara,T., Nishi,H., Yamada,S., Ohara,M., McCallum,N., Berger-

Bächi,B. and Sugai,M. (2004) The gate controlling cell wall synthesis in *Staphylococcus*

aureus. *Mol. Microbiol.*, **53**, 1221–1231.

89. Uehara,T. and Park,J.T. (2004) The *N*-Acetyl-D-Glucosamine Kinase of *Escherichia coli*

and Its Role in Murein Recycling. *J. Bacteriol.*, **186**, 7273–7279.

90. Meibom,K.L., Li,X.B., Nielsen,A.T., Wu,C.-Y., Roseman,S. and Schoolnik,G.K. (2004)

The *Vibrio cholerae* chitin utilization program. *Proc. Natl. Acad. Sci. U. S. A.*, **101**, 2524

LP – 2529.

91. Iyer,L.M. and Aravind,L. (2002) The catalytic domains of thiamine triphosphatase and CyaB-like adenylyl cyclase define a novel superfamily of domains that bind organic phosphates. *BMC Genomics*, **3**, 33.

92. Vogt,M.S., Ngouoko Nguepbeu,R.R., Mohr,M.K.F., Albers,S.-V., Essen,L.-O. and Banerjee,A. (2021) The archaeal triphosphate tunnel metalloenzyme *Sa*TTM defines structural determinants for the diverse activities in the CYTH protein family. *J. Biol. Chem.*, **297**.

93. Mamou,G., Malli Mohan,G.B., Rouvinski,A., Rosenberg,A. and Ben-Yehuda,S. (2016) Early Developmental Program Shapes Colony Morphology in Bacteria. *Cell Rep.*, **14**, 1850–1857.

94. Fleming,E. and Camilli,A. (2016) ManLMN is a glucose transporter and central metabolic regulator in *Streptococcus pneumoniae*. *Mol. Microbiol.*, **102**, 467–487.

95. Abranches,J., Candella,M.M., Wen,Z.T., Baker,H. V. and Burne,R.A. (2006) Different Roles of EIIAB^{Man} and EII^{Glc} in Regulation of Energy Metabolism, Biofilm Development, and Competence in *Streptococcus mutans*. *J. Bacteriol.*, **188**, 3748–3756.

96. Rossol,M., Pierer,M., Raulien,N., Quandt,D., Meusch,U., Rothe,K., Schubert,K., Schöneberg,T., Schaefer,M., Krügel,U., *et al.* (2012) Extracellular Ca^{2+} is a danger signal activating the NLRP3 inflammasome through G protein-coupled calcium sensing receptors. *Nat. Commun.*, **3**, 1329.

97. Selvaraj,S., Liu,K., Robinson,A.M., Epstein,V.A., Conley,D.B., Kern,R.C. and Richter,C.-P. (2012) *In Vivo* Determination of Mouse Olfactory Mucus Cation Concentrations in Normal and Inflammatory States. *PLoS One*, **7**, e39600.

98. Seto,H. and Tomasz,A. (1976) Calcium-requiring step in the uptake of deoxyribonucleic acid molecules through the surface of competent pneumococci. *J. Bacteriol.*, **126**, 1113–1118.

99. Domínguez,D.C., Guragain,M. and Patrauchan,M. (2015) Calcium binding proteins and calcium signaling in prokaryotes. *Cell Calcium*, **57**, 151–165.

100. Grote,J., Krysciak,D. and Streit,W.R. (2015) Phenotypic Heterogeneity, a Phenomenon That May Explain Why Quorum Sensing Does Not Always Result in Truly Homogenous Cell Behavior. *Appl. Environ. Microbiol.*, **81**, 5280–5289.

101. Chalancon,G., Ravarani,C.N.J., Balaji,S., Martinez-Arias,A., Aravind,L., Jothi,R. and Babu,M.M. (2012) Interplay between gene expression noise and regulatory network architecture. *Trends Genet.*, **28**, 221–232.

102. Prudhomme,M., Berge,M., Martin,B. and Polard,P. (2016) Pneumococcal Competence Coordination Relies on a Cell-Contact Sensing Mechanism. *PLOS Genet.*, **12**, e1006113.

103. Raj,A. and van Oudenaarden,A. (2008) Nature, Nurture, or Chance: Stochastic Gene Expression and Its Consequences. *Cell*, **135**, 216–226.

104. Maier,B. (2020) Competence and Transformation in *Bacillus subtilis. Curr. Issues Mol. Biol.*, **37**.

105. Haycocks,J.R.J., Warren,G.Z.L., Walker,L.M., Chlebek,J.L., Dalia,T.N., Dalia,A.B. and Grainger,D.C. (2019) The quorum sensing transcription factor AphA directly regulates natural competence in *Vibrio cholerae. PLOS Genet.*, **15**, e1008362.

106. Seitz,P. and Blokesch,M. (2013) Cues and regulatory pathways involved in natural competence and transformation in pathogenic and environmental Gram-negative bacteria. *FEMS Microbiol. Rev.*, **37**, 336–363.

107. Claverys,J.-P. and Håvarstein,L.S. (2007) Cannibalism and fratricide: mechanisms and raisons d'etre. *Nat. Rev. Microbiol.*, **5**, 219–229.

108. González-Pastor,J.E. (2011) Cannibalism: a social behavior in sporulating Bacillus subtilis. *FEMS Microbiol. Rev.*, **35**, 415–424.

109. Borgeaud,S., Metzger,L.C., Scrignari,T. and Blokesch,M. (2015) The type VI secretion system of *Vibrio cholerae* fosters horizontal gene transfer. *Science (80-.).*, **347**, 63–67.

110. Goodman,S.D. and Scocca,J.J. (1988) Identification and arrangement of the DNA sequence recognized in specific transformation of *Neisseria gonorrhoeae. Proc. Natl. Acad. Sci.*, **85**, 6982–6986.

111. Cameron,A.D.S., Volar,M., Bannister,L.A. and Redfield,R.J. (2008) RNA secondary structure regulates the translation of *sxy* and competence development in *Haemophilus influenzae. Nucleic Acids Res.*, **36**, 10–20.

112. Sisco,K.L. and Smith,H.O. (1979) Sequence-specific DNA uptake in *Haemophilus transformation. Proc. Natl. Acad. Sci.*, **76**, 972–976.

113. Suckow,G., Seitz,P. and Blokesch,M. (2011) Quorum sensing contributes to natural transformation of *Vibrio cholerae* in a species-specific manner. *J. Bacteriol.*, **193**, 4914–4924.

114. Stewart,G.J. and Carlson,C.A. (1986) The Biology of Natural Transformation. *Annu. Rev. Microbiol.*, **40**, 211–231.

115. Redfield,R.J. (1993) Genes for breakfast: The have-your-cake and-eat-it-too of bacterial transformation. *J. Hered.*, **84**, 400–404.

116. Macfadyen,L.P., Chen,D., Vo,H.C., Liao,D., Sinotte,R. and Redfield,R.J. (2001) Competence development by *Haemophilus influenzae* is regulated by the availability of nucleic acid precursors. *Mol. Microbiol.*, **40**, 700–707.

117. Sprouffske,K. (2020) growthcurver: Simple Metrics to Summarize Growth Curves.

118. Apagyi,K.J., Fraser,C. and Croucher,N.J. (2018) Transformation asymmetry and the evolution of the bacterial accessory genome. *Mol. Biol. Evol.*, **35**, 575–581.

119. Elzhov,T. V, Mullen,K.M., Spiess,A.-N. and Bolker,B. (2016) minpack.lm: R Interface to the Levenberg-Marquardt nonlinear least-squares algorithm found in MINPACK, plus support for bounds.

120. Fox,J. and Weisberg,S. (2019) An {R} Companion to Applied Regression Third. Sage, Thousand Oaks {CA}.

121. Kwun,M.J., Oggioni,M.R., Bentley,S.D., Fraser,C. and Croucher,N.J. (2019) Synergistic Activity of Mobile Genetic Element Defences in *Streptococcus pneumoniae. Genes (Basel).*, **10**, 707.

122. Nawrocki,E.P. and Eddy,S.R. (2013) Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*, **29**, 2933–2935.

123. Kalvari,I., Argasinska,J., Quinones-Olvera,N., Nawrocki,E.P., Rivas,E., Eddy,S.R., Bateman,A., Finn,R.D. and Petrov,A.I. (2018) Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. *Nucleic Acids Res.*, **46**, D335–D342.

124. Bray,N.L., Pimentel,H., Melsted,P. and Pachter,L. (2016) Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.*, **34**, 525–527.

125. Pimentel,H., Bray,N.L., Puente,S., Melsted,P. and Pachter,L. (2017) Differential analysis of RNA-seq incorporating quantification uncertainty. *Nat. Methods*, **14**, 687–690.

126. Guy,L., Roat Kultima,J. and Andersson,S.G.E. (2010) genoPlotR: comparative gene and genome visualization in R. *Bioinformatics*, **26**, 2334–2335.

127. Gu,Z., Gu,L., Eils,R., Schlesner,M. and Brors,B. (2014) circlize implements and enhances circular visualization in R . *Bioinformatics*, **30**, 2811–2812.

128. Wilke,C. (2020) cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'.

129. Kassambara,A. (2020) ggpubr: 'ggplot2' Based Publication Ready Plots.

130. Wickham,H., Averick,M., Bryan,J., Chang,W., McGowan,L., François,R., Grolemund,G., Hayes,A., Henry,L., Hester,J., *et al.* (2019) Welcome to the Tidyverse. *J. Open Source Softw.*, **4**, 1686.

131. Delcher,A.L., Phillippy,A., Carlton,J. and Salzberg,S.L. (2002) Fast algorithms for large-scale genome alignment and comparison. *Nucleic Acids Res.*, **30**, 2478–2483.

132. Sobetzko,P., Jelonek,L., Strickert,M., Han,W., Goesmann,A. and Waldminghaus,T. (2016) DistAMo: A Web-Based Tool to Characterize DNA-Motif Distribution on Bacterial Chromosomes. *Front. Microbiol.*, **7**, 283.

133. Cock,P.J.A., Antao,T., Chang,J.T., Chapman,B.A., Cox,C.J., Dalke,A., Friedberg,I., Hamelryck,T., Kauff,F., Wilczynski,B., *et al.* (2009) Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics*, **25**, 1422–1423.

134. Edgar,R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.*, **32**, 1792–1797.

135. Varadi,M., Anyango,S., Deshpande,M., Nair,S., Natassia,C., Yordanova,G., Yuan,D., Stroe,O., Wood,G., Laydon,A., *et al.* (2022) AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res.*, **50**, D439–D444.

136. Letunic,I., Khedkar,S. and Bork,P. (2021) SMART: recent updates, new developments and status in 2020. *Nucleic Acids Res.*, **49**, D458–D460.

137. Katoh,K. and Standley,D.M. (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol. Biol. Evol.*, **30**, 772–780.

138. Price,M.N., Dehal,P.S. and Arkin,A.P. (2010) FastTree 2–approximately maximum-likelihood trees for large alignments. *PLoS One*, **5**, e9490.

139. Schoch,C.L., Ciufo,S., Domrachev,M., Hotton,C.L., Kannan,S., Khovanskaya,R., Leipe,D., Mcveigh,R., O'Neill,K., Robbertse,B., *et al.* (2020) NCBI Taxonomy: a comprehensive update on curation, resources and tools. *Database (Oxford).*, **2020**.

140. Stutzmann,S. and Blokesch,M. (2020) Comparison of chitin-induced natural transformation in pandemic *Vibrio cholerae* O1 El Tor strains. *Environ. Microbiol.*, **22**, 4149–4166.

141. Acebo,P., Martin-Galiano,A.J., Navarro,S., Zaballos,A. and Amblar,M. (2012) Identification of 88 regulatory small RNAs in the TIGR4 strain of the human pathogen *Streptococcus pneumoniae*. *RNA*, **18**, 530–546.

142. Piotrowski,A., Luo,P. and Morrison,D.A. (2009) Competence for genetic transformation in Streptococcus pneumoniae: termination of activity of the alternative sigma factor ComX is independent of proteolysis of ComX and ComW. *J. Bacteriol.*, **191**, 3359–3366.

Figures

Figure 1 Phenotypic differences between locked *tvr* variants. (A) Violin plot showing the

transformation efficiency of four pairs of *tvr* locus variants constructed from isolates RMV5,

RMV6, RMV7 and RMV8. Each individual point represents an independent transformation

experiment. The horizontal line within each violin shows the median for each genotype. The

brackets indicate the statistical significance of the comparison between variants from the

same isolate background, as calculated using a two-tailed Wilcoxon rank sum test. (B)

Schematic of the *tvr* loci from RMV7$_{wt}$, RMV7$_{domi}$ and RMV7$_{rare}$ to show the genes encoding

the methylase (*hsdM*), endonuclease (*hsdR*), regulatory system (*tvrAT*) and recombinase

(*tvrR*). The variants differ in their active *hsdS* genes, upstream of *tvrATR*. The RMV7$_{domi}$

HsdS protein comprises the TRD combination III-iii (recognising motif TGAN$_7$TCC), whereas

that of RMV7$_{rare}$ contains the TRDs III-I (recognising motif TGAN$_7$TATC). The black arrows

indicate the binding sites of a forward primer, in *hsdM*, and reverse primers, in *hsdS*

fragments. (C) Line graph showing the ratio of RMV7$_{domi}$ to RMV7$_{rare}$ loci in eight-day

passages of RMV7$_{wt}$. The RMV7$_{domi}$ variant was more common in each of the three

replicates at every measured timepoint. (D) Violin plot showing the higher transformation

efficiency of RMV$_{rare}$ relative to RMV7$_{domi}$ or RMV7$_{wt}$. Each individual point represents an

independent experiment in which the number of transformants, and number of overall

colony-forming units (cfu), was calculated. The horizontal line within each violin shows the

median for each genotype. Both mutants were compared with RMV7$_{wt}$; the horizontal

bracket shows a significant difference, as calculated from a two-tailed Wilcoxon rank sum

test. (E) Violin plot showing the adhesion of variants to an abiotic surface, as quantified by

the optical density at 550 nm after crystal violet staining of cells resisting washing from a

microplate well. The RMV7$_{rare}$ variant showed a significantly greater adhesion after 16 hours

incubation at 35 °C, as assessed by a two-tailed Wilcoxon rank sum test. (F) Violin plots

showing the transformation efficiency of knock-in mutants during a passage experiment. The

tvr loci of RMV7 *tvr*$_{domi}$::Janus and RMV7 *tvr*$_{rare}$::Janus (both containing a *tvrR* gene

interrupted by a Janus cassette) were each introduced into an RMV7 *tvr*::*cat* background

(Fig. S5). This pair were separately passaged in liquid cultures over six days in five

independent replicates. The number of transformants observed from three transformation

assays, conducted each day for both variants, is shown by the individual points' shapes and

colours. The violin plots summarise the median and distribution of these values. The

brackets indicate the statistical significance of the comparison between variants from the

same day of the passage, as calculated using a two-tailed Wilcoxon rank sum test. Across

all panels, significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p

values were subject to a Holm-Bonferroni correction within each panel.

Figure 2 Genes exhibiting significant differences in transcription between RNA-seq samples.

(A) Design of the RNA-seq experiment. (B)-(F) Volcano plots showing the variation in gene

transcription between RNA-seq samples. The horizontal axis shows the natural logarithm of

the fold difference in expression levels between the genotypes, β. The vertical axis shows

the negative base 10 logarithm of the q value, corresponding to a Benjamini-Hochberg

corrected p value. Points are coloured red where this value exceeds the threshold false

discovery rate threshold of 10^{-3}. The panels correspond to (B) the differences between the

pre-CSP samples of tvr_{domi}::Janus and tvr_{rare}::Janus; the differences between the pre-CSP

and 10 minute post-CSP samples for (C) tvr_{domi}::Janus and (D) tvr_{rare}::Janus; and the

differences between the pre-CSP and 20 minute post-CSP samples for (E) tvr_{domi}::Janus and

(F) tvr_{rare}::Janus. The significant difference in $tvrR$ expression between the tvr_{domi}::Janus and

tvr_{rare}::Janus mutants is an artefact of the different constructs used to disrupt $tvrR$ in these

two genotypes (see Methods). (G) Chromosomal distribution of differentially-expressed

genes. The outer ring shows the annotation of RMV7$_{domi}$ (accession code OV904788).

Protein coding sequences are represented as black boxes, with the vertical positioning

within the ring indicating the strand of the genome on which they are encoded. The next ring

inwards shows significant pre-CSP differences in transcription: green genes were more

highly expressed in RMV$_{domi}$, and blue genes were more highly expressed in RMV$_{rare}$. The

next ring inwards show significant changes in gene expression 10 minutes post-CSP in

RMV7$_{domi}$: pink genes were upregulated, and purple genes were downregulated. The third

ring inwards shows significant changes in gene expression 10 minutes post-CSP in

RMV7$_{rare}$: red genes were upregulated, and orange genes were downregulated. The two

inner rings repeat this representation for changes in gene expression 20 minutes post-CSP.

Figure 3 The dependence of transformation efficiency on import of carbon sources. (A)

Violin plots showing the transformation efficiency of RMV7$_{rare}$ relative to mutant derivatives.

The counter selectable Janus cassette was used to disrupt the *manLMN*, *tfoX* and *yjbK*

genes. This enabled the restoration of the *tfoX* and *yjbK* genes (similar data for *manL* are

shown in Fig. S28). Each genotype was transformed in unsupplemented media, and in the

presence of one of six carbon sources, as indicated by the plot colour (see key). Each point

represents an independent experiment, and the horizontal line within the violin plots show

the median for each combination of recipient cell genotype and carbon source. For each

genotype, Wilcoxon rank-sum tests were used to test for evidence of changes in

transformation efficiency caused by each carbon source, relative to the unsupplemented

media. Significant differences are indicated by the black brackets at the top of the panel. (B)

Violin plots showing the transformation efficiency of RMV7$_{wt}$ relative to mutant derivatives.

Data are displayed as in panel A. (C) Violin plots quantifying the effects of carbon source

supplementation, and disruption of *manLMN* and the *rlrA* pilus islet, on adhesion of bacteria

to an abiotic surface using OD$_{550}$. The density of the biofilm formed following growth in

GlcNAc-supplemented media was used as the comparator for Wilcoxon rank sum tests, as

all carbon source supplements increased biofilm formation relative to unsupplemented

media. (D) Violin plots showing the transformation efficiency of double mutants constructed

in the RMV7$_{rare}$ genotype. For comparison with panel A, the horizontal lines show the median

transformation efficiencies of the corresponding single mutants in unsupplemented and

GlcNAc-supplemented media: *manLMN*::Janus as solid lines; *tfoX*::Janus as dotted lines,

and *yjbK*::Janus as a dashed line. Data are shown on a logarithmic scale to resolve the

differences between these poorly-transformable genotypes. (E) Violin plots showing the

effect of GlcNAc supplementation on the transformation efficiency of the most common *tvr*

variants of the RMV5, RMV6 and RMV8 isolates. Across all panels, significance was coded

as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p values were subject to a Holm-

Bonferroni correction within each panel.

Figure 4. Effect of removing PRCI*dnaN* on the transformation efficiency of RMV7wt. (A) Violin

plots showing the number of transformants observed in assays of RMV7wt mutants in which

different parts of PRCI*dnaN* were replaced with a Janus cassette. The genotypes are arranged

left to right, and coloured black to purple, to represent the increasing proportion of the

element that was replaced by the cassette. Mutants removed different combinations of the

dinD-like gene (IONPJBJN_00496); the regulatory genes (*rep*); the replication genes (*rep*);

the integration genes (*int*), and the *att* site. The structure of RMV7 $PRCI_{dnaN}$ meant that

replacing the *int-rep* region also deleted the intervening *reg* genes. Each point represents an

independent transformation assay. The violin plot summarises the result for each mutant,

with a horizontal line indicating the median. Each mutant was compared against the parental

$RMV7_{wt}$ genotype using a Wilcoxon rank sum test. Black brackets at the top of the plot

indicate significant differences in the number of observed transformants. (B) Violin plot

quantifying the effect of $PRCI_{dnaN}$ and the chaperones HrcA and ClpP on transformation

efficiency in $RMV7_{wt}$. The comparison of $RMV7_{wt}$ with a mutant in which the PRCI and its *att*

site were removed was independent of the experiments presented in panel A, and more

accurately quantified transformation efficiency. This approach was also used to compare the

effects of single and double mutations that disrupted *hrcA*, *clpP* and $PRCI_{dnaN}$. (C) Violin

plots showing the effect of the disruption of $PRCI_{dnaN}$ on gene expression, quantified as

abundance relative to *rpoA* by qRT-PCR. IONPJBJN_00507 is a coding sequence within the

PRCI that is absent from $RMV7_{wt}$ $PRCI_{dnaN+att}$::Janus. The six points for each gene

correspond to three technical replicate assays on each of two biological replicates. The

horizontal line on the violin plot shows the median relative abundance for each gene in each

genotype. Across all panels, significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; p

$< 10^{-4}$, ****. All p values were subject to a Holm-Bonferroni correction within each panel.

Figure 5 The regulation of transformation by Ca²⁺ and heat shock in RMV7. (A) Violin plots

showing the transformation efficiency of RMV7$_{wt}$, RMV7$_{rare}$ and mutant derivatives in which

hrcA and *clpP* were disrupted during normal growth (35 °C) or during a 40 °C heat shock.

Each point corresponds to an independent transformation experiment, and the violin plots

have a horizontal line indicating the median transformation efficiency of each mutant at each

temperature. (B) Scatterplot showing the dose-dependent effect of $CaCl_2$ on transformation

efficiency in RMV7$_{rare}$ (cyan) and mutants in which *hrcA* had been disrupted (magenta), and

then restored (dark blue). Each point represents an independent transformation assay of

one genotype at the indicated $CaCl_2$ concentration. The best-fitting dose response logistic

models are shown, with the shaded areas corresponding to the 95% confidence intervals.

(C) Expression of *hrcA*, measured as transcript abundance relative to *rpoA* by qRT-PCR,

following the exposure of cells to CSP. CSP stimulated higher expression of *hrcA* unless co-

administered with 12.5 mM $CaCl_2$, demonstrating that $CaCl_2$ addition affects HrcA regulatory

activity. (D) Expression of the early competence gene *comX* and late competence gene

comEA following the addition of CSP in RMV7$_{rare}$ (cyan), RMV7$_{rare}$ *hrcA*::Janus (magenta)

and RMV7$_{rare}$ *manLMN*::Janus (peach), measured relative to the abundance of *rpoA* by qRT-

PCR. (E) Combined effects of chaperone and carbon source regulation on transformation

efficiency. Results are displayed as in panel A. Wilcoxon rank-sum tests were conducted

between all pairs of genotypes. These found both the single mutants, lacking *manLMN* or

hrcA, were significantly less transformable than the parental genotype. Furthermore, the

double mutant was less transformable than either single mutant. Across all panels,

significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p values

were subject to a Holm-Bonferroni correction within each panel.

Figure 6 Comparison of the regulation of competence in (A) *S. pneumoniae*, from this work,

and (B) *V. cholerae*, summarized from (140). In each, competence is regulated by a quorum-

sensing system: CSP in *S. pneumoniae*, and the cholera autoinducer 1 (CAI-1) in *V.*

cholerae. The production of CSP is known to be inhibited by the non-coding csRNA, and the

HtrA protease derades the signal. Similarly, the srn206 non-coding RNA represses the

ComD receptor of CSP (141). CAI-1 operates through inhibiting LuxO, thereby activating the

HapR protein, which indicates a high-cell density environment. HapR activates competence

through the Quorum-Sensing and TfoX-dependent Regulator (QstR). This regulator also

senses the activation of TfoX in response to GlcNAc being sensed by the transmembrane

regulator TfoS, via the TfoR small RNA. TfoX activity is also regulated by the Catabolite

Regulatory Protein (CRP), which is activated by 3',5'-cAMP, generated by the CyaA

adenylate cyclase under carbon source starvation conditions. Hence there are parallels with

the TfoX orthologue, and adenylate cyclase-like protein YjbK, responding to GlcNAc in *S.*

pneumoniae RMV7. GlcNAc also appears to promote competence through a TfoX/YjbK-

independent route, based on the behaviour of RMV7$_{wt}$. There are no known parallels in *V.*

cholerae for the regulation of competence by the chaperones ClpP and HrcA. ClpP

represses the induction of competence in response to stresses, such as MGE replication,

and has been previously shown to degrade ComX (142). One of HrcA's two conformations

appears to have a similar role, repressing competence in response to heat shock, albeit

through a different mechanism. The other active conformation of HrcA, denoted HrcA*,

appears to activate competence in response to Ca^{2+}, again through an unknown

mechanism.