Moonlighting proteins activate transformation in epigenetically-differentiated phase

variants of multidrug-resistant *Streptococcus pneumoniae*

Min Jung Kwun[1], Alexandru V. Ion[1], Marco R. Oggioni[2,3], Stephen D. Bentley[4], Nicholas J.

Croucher[1,*]

[1] MRC Centre for Global Infectious Disease Analysis, Sir Michael Uren Hub, White City

Campus, Imperial College London, London, W12 0BZ, UK

[2] Department of Genetics, University of Leicester, University Road, Leicester, LE1 7RH

[3] Dipartimento di Farmacia e Biotecnologie, Università di Bologna, Via Irnerio 42, 40126

Bologna, Italy

[4] Parasites & Microbes, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton,

Cambridge, CB10 1SA, UK

Abstract

Transformation of *Streptococcus pneumoniae* GPSC1 has enabled vaccine evasion and the

acquisition of antibiotic resistance. Epigenetic phase variants of GPSC1 isolate RMV7,

differentiated by rearrangements at the *tvr* restriction-modification locus, differed ~100-fold in

their transformation efficiency. This variation was recapitulated in knock-in mutants of the

relevant *tvr* alleles. RNA-seq showed the difference was due to blocking of the early

competence regulatory cascade. The more transformable variant upregulated expression of

manLMN, encoding a carbon source importer. This was shown to be necessary for efficient

competence induction, despite being dispensable for growth in rich media. Transformation

was promoted by import of *N*-acetylglucosamine, which activated competence through an

orthologue of the gram-negative competence regulator TfoX, and an enzyme likely involved

in nucleotide-mediated signalling. A mobile genetic element was more active in the less

transformable variant, which limited competence induction through inhibiting CIRCE motif

binding by the chaperone regulator HrcA. Correspondingly, both heat shock and decreased

Ca^{2+} concentrations reduced competence by limiting HrcA binding regulatory sequence

motifs. Hence both ManLMN and HrcA moonlighted as activators of competence. Such

proteins may be important in potentiating horizontal DNA exchange, as selection on their

primary function likely constrains them from mutating into alleles that selfishly downregulate

transformation.

Competence for natural transformation was first identified in *Streptococcus pneumoniae* (the

pneumococcus) in the early 20th century [1]. Cells can be "transformed" to express a new

phenotype through the acquisition of exogenous DNA, integrated into their genome through

homologous recombination following its import from the environment through the specialised

cell-encoded competence machinery [2]. Transformation has played a key role in the

emergence of antibiotic-resistant *S. pneumoniae*, both through generating "mosaic" alleles of

core loci [3–5] and the acquisition of specialised resistance genes [6]. It has also enabled

vaccine evasion through recombinations affecting the capsule polysaccharide synthesis

(*cps*) locus altering surface antigens [7,8].

Despite the ability of transformation to accelerate such adaptive evolution in *S. pneumoniae*,

considerable variation in the rate of diversification of strains through this mechanism persists

across the species [9]. Epidemiological studies have found the *r*/*m* ratio of base substitutions

introduced through homologous recombination, relative to those occurring through point

mutation, varies from well over 10 [7,9] to below 0.1 [9,10] across the species. Similarly, *in*

vitro assays have identified >100-fold differences in the transformation efficiency of *S.*

pneumoniae genotypes, with substantial variation even between isolates of the same

serotype or strain [11–14]. Many isolates are routinely found not to be transformable under

standard conditions [11]. This is often the consequence of integrative mobile genetic

elements (MGEs) disrupting genes necessary for transformation [6,15–17], selfishly

preventing themselves from being eliminated from the chromosome [17]. Yet in other non-

transformable isolates the highly-conserved competence machinery is intact [11,18]. This

suggests the variation in transformation rates also reflects differences in regulation of the

competence system.

The best-characterised signal inducing transformation in *S. pneumoniae* is the competence

stimulating peptide (CSP) pheromone, which acts as a quorum-sensing system that signals

through the ComDE two-component regulator [19]. This activates early competence genes

after about ten minutes [20]. These include *comX*, encoding an alternative sigma factor [21].

ComX enables the RNA polymerase to recognise late competence genes [20], which feature

a "combox" signal in their promoters [22,23]. This results in pneumococci entering a

transient competent state around 20 minutes post-CSP induction, after which the relevant

machinery is degraded [24], and the cells become temporarily refractory to induction [21].

Transformation efficiency is known to vary between isogenic pneumococci through phase

variation in capsule production. Transparent colony variants produce less capsule, and are

less virulent, but more transformable, than opaque colony variants [25]. This short-term

variation has been linked to rapid changes at the phase-variable inverting variable restriction

(*ivr*) locus, encoding the conserved Type I *Spn*III restriction-modification system (RMS) and

the IvrR recombinase that drives sequence inversions within the locus [26–31]. These

rearrangements alter the target recognition domains (TRDs) within the active HsdS

specificity protein, which determines the DNA motif that is targeted by both the methylase

and endonuclease activities of the system. Consequently, changes at this single locus affect

genome-wide methylation patterns, causing pleiotropy through its impact on the transcription

of multiple genes [27]. These phase-variable RMSs can thereby maintain phenotypic

heterogeneity within a genetically near-homogenous population [32], resulting in "bet

hedging" that can increase the chances of a species surviving a changing environment

[33,34].

The second pneumococcal phase-variable Type I restriction-modification system (named

*Spn*IV), encoded by the translocating variable restriction (*tvr*) locus, varies through TvrR

recombinase-mediated excision-reintegration [27,28,35,36]. This locus is generally

conserved across pneumococci, but the complement of TRDs varies between strains,

increasing the range of HsdS proteins, and possible methylation patterns, across the

species [28,35]. This locus is inactive in the R6x laboratory strain that is typically used to

study pneumococcal competence [28,35,37]. Here, we test pairs of otherwise-isogenic *tvr*

variants to identify the regulatory mechanisms underlying phenotypic heterogeneity in

genotypes commonly causing pneumococcal disease.

Results

Phase variants of *S. pneumoniae* RMV7 differ in their transformation efficiency

Four pairs of otherwise-isogenic "locked" phase variants, in which *tvrR* was knocked out or

disrupted by a selectable and counter-selectable Janus cassette marker [35,38], were

screened for differences in their transformation efficiency (Fig. 1A). In the RMV6 and RMV7

pairs, the variant in which the active *tvr* locus HsdS comprised the TRDs III-i (directing the

SpnIV system to target the bipartite motif $TGAN_7TATC$) was found to have significantly

higher transformation efficiency following induction by exogenous CSP. These closely-

related genotypes both originate from the multidrug-resistant GPSC1 strain [39]. Many

replicates were required to robustly establish the difference between the RMV6 variants,

whereas the RMV7 variants exhibited a clearer ~100-fold difference in their transformation

rate (Fig. 1A). This was because the RMV7 variant carrying the alternative form of the *tvr*

locus, with an active HsdS comprising the TRDs III-iii (which directs the SpnIV system to

target the motif $TGAN_7TCC$; Fig. 1B and S1) had an almost-undetectable transformation

efficiency. Culturing the wild-type isolate (RMV7$_{wt}$) over successive days demonstrated large changes in the relative frequency of these variants, although the less-transformable variant (RMV7 *tvrR*::Janus; henceforth, RMV7$_{domi}$, carrying *tvr*$_{domi}$) was typically more common than the more transformable variant (RMV7 Δ*tvrR*; henceforth, RMV7$_{rare}$, carrying *tvr*$_{rare}$; Fig. 1C). A comparison of RMV7$_{wt}$ and RMV7$_{domi}$ found them to have similarly low transformation efficiencies, whereas that of RMV7$_{rare}$ was confirmed to be ~50-fold higher, consistent with the relative proportions of the variants observed *in vitro* (Fig. 1D). Therefore, RMV7$_{domi}$ and RMV7$_{rare}$ exhibited distinctive phenotypes that correlated with their *tvr* locus arrangements.

RMV7$_{rare}$ was also significantly more adhesive to an abiotic surface (Fig. 1E), which can be considered a proxy for biofilm formation [40]. This could be consistent with RMV7$_{rare}$ being enriched for transparent phase variants. However, microscopy found no clear difference in colony morphology between the variants (Fig. S2). An alternative explanation for the phenotypic differences could be mutations that occurred during genetic manipulation of the isolates [41]. Alignment of the two variants' assemblies found them to be distinguished by seven non-synonymous single nucleotide polymorphisms and two premature stop codons outside the *tvr* locus, none of which were within genes known to directly affect the competence machinery (Table S1). Nevertheless, we tested the effect on transformation of mutations in RMV7$_{rare}$ that were absent from both the RMV7$_{domi}$ and RMV7$_{wt}$ sequences. A non-synonymous change in *phoB*, encoding a phosphate-sensitive response regulator [42], was not found to affect transformation efficiency (Fig. S3). Similarly, a premature stop codon in *pstS* was observed in RMV7$_{rare}$, but knocking out this gene also failed to explain any difference in transformability between the variants (Fig. S3). Hence the differences between RMV7 variants could not be explained by point mutations or alterations in encapsulation.

To test whether the phenotypic differences were causatively associated with variation in the SpnIV RMS, the *tvr* loci of RMV7$_{domi}$ and RMV7$_{rare}$ were introduced into RMV7$_{wt}$ *tvr*::*cat*. This generated the otherwise isogenic knock-in recombinants RMV7 *tvr*$_{domi}$::Janus and RMV7 *tvr*$_{rare}$::Janus, carrying two different locked *tvr* loci (Fig. S4). The transformability of both was assayed over five replicate six-day passages. This demonstrated a persistent distinction in the transformation efficiency of the two *tvr* genotypes, which replicated the difference between the original variants (Fig. 1F). Therefore, the RMV7 *tvr*$_{rare}$ and *tvr*$_{domi}$ alleles were associated with significant differences in transformation efficiency in otherwise isogenic isolates.

Induction of early and late competence genes is blocked in RMV7$_{domi}$

To understand how the *tvr* loci caused a difference in transformation, RNA-seq was used to

quantify patterns of transcription in the recombinants RMV7 *tvr*domi::Janus and RMV7

*tvr*rare::Janus. Samples were taken pre-CSP, 10 minutes post-CSP, and 20 minutes post-

CSP from each of three biological replicates (Fig. S4). The 18 RNA samples were

sequenced as 200 nt paired-end multiplexed libraries on a single Illumina HiSeq 4000 lane.

After alignment to the RMV7domi genome, analysis of the RNA-seq data found the fragment

length distributions (Fig. S5) and gene expression densities (Fig. S6) were consistent across

samples (Table S2). Q-Q plots suggested a Benjamini-Hochberg corrected *q* value of 10^{-3}

was an appropriate threshold for identifying significant transcriptional variation (Fig. S7-8).

This identified 154 genes that significantly differed in their expression between the two

genotypes prior to CSP exposure, or between the pre- and post-CSP samples from the

same genotype (Fig. 2 and S9-10; Table S3).

Clustering of the datasets found the biggest separation distinguished the post-CSP RMV7

*tvr*rare::Janus transcriptomes from the others (Fig. S11), suggesting a major difference in the

induction of the competence system between the variants. In RMV7 *tvr*rare::Janus, the early

competence genes showed elevated expression 10 minutes post-CSP (Fig. S12), with the

late competence genes exhibiting more variable patterns of transcription (Fig. S13). Also

upregulated was *briC*, a competence-induced biofilm formation signal [43], and some

nucleotide metabolism and transporter genes (*purA*, *tadA*, *dut*, *ribF*, *adeQ*; Fig. S14).

Another upregulated transporter, encoded by a three gene cluster, has reportedly been

induced in response to exposure to CSP [44] and antimicrobial peptides [45], and

consequently was named *pieABC* (peptide-induced exporter; CDSs IONPJBJN_01324-6 in

RMV7domi, corresponding to SP_0785-787 in TIGR4; Fig. S14, Table S3).

By contrast, CSP upregulated just five genes in RMV7 *tvr*domi::Janus: the quorum sensing

genes *comCDE* and *briC* (Fig. S12), and a gene encoding a CsbD stress response protein

that was not CSP-responsive in RMV7 *tvr*rare::Janus. However, the induction of *comCDE* was

much weaker, and there was no sign of late competence genes being activated by ComX.

However, expression of the *comX* gene itself can be difficult to determine through RNA-seq,

owing to the presence of two near-identical paralogues in pneumococcal genomes [46].

As an independent test of these transcriptional differences, qRT-PCR experiments were

undertaken on the original RMV7domi and RMV7rare variants, and the control genotypes

RMV7wt and RMV7wt *tvr*::*cat*, the latter of which lacked a *tvr* locus. Transformation assays

performed in parallel with these RNA extractions found RMV7wt *tvr*::*cat* was more efficiently

transformable than RMV7wt or RMV7 *tvr*domi::Janus, demonstrating the active, dominant form

of the locus was repressing transformation (Fig. S4). The qRT-PCR data showed genotypes

of both low (RMV7$_{wt}$ and RMV7$_{domi}$) and high (RMV7$_{rare}$ and RMV7$_{wt}$ *tvr*::Janus)

transformation efficiency up-regulated the early competence genes *comD* and *comX* in

response to CSP (Fig. S15)*.* However, the late competence genes *comEA* and *comYC* were

only significantly upregulated in the highly transformable strains. Hence the difference in

transformability between the RMV7 variants was a consequence of late competence gene

activation being blocked in RMV7$_{domi}$ through consequences of *tvr*$_{domi}$ expression.

Differences in pre-CSP gene expression

The two pre-CSP samples were compared to identify the cause of this difference in

competence induction. Pre-CSP, 33 genes exhibited higher expression in RMV7

tvr$_{domi}$::Janus, and 20 genes exhibited higher expression in RMV7 *tvr*$_{rare}$::Janus (Fig. 2; Table

S3). These did not include any *cps* locus genes, which were more highly expressed in RMV7

tvr$_{rare}$::Janus (Fig. S16), confirming that the elevated transformation efficiency of this

genotype did not reflect an enrichment of transparent phase variants [27,47]. To test

whether these differentially-expressed genes were affected by methylation within regulatory

or promoter sites, the distances from protein coding sequence start codons to the nearest

methylation site was plotted for both SpnIV motifs (Fig. S17). There was no general

relationship between differential expression and proximal methylation, although four cases of

variable methylation sites being within 100 bp of a differentially-expressed gene's start

codon were identified. In three cases, the coding sequences close to *tvr*$_{domi}$ motifs were

within a larger set of co-transcribed genes, and it was unlikely that transcription initiation

would be affected by methylation (Fig. S18). However, one *tvr*$_{rare}$ motif was within the

promoter of the *piuABCD* operon (Fig. S18), encoding an iron transporter, that was

upregulated in RMV7 *tvr*$_{rare}$::Janus (Fig. S14). However, knocking out *piuA* did not reduce

the transformation efficiency of RMV7$_{rare}$, suggesting this change was independent of those

affecting competence (Fig. S15).

As the induction of transformation has been suggested to represent a stress response [48],

the overall distribution of the SpnIV target motifs was analysed, to test whether differing

spatial arrangements across the chromosome might affect cell physiology. While the *tvr*$_{rare}$

motifs were uniformly distributed, the RMV7$_{domi}$ motifs were enriched in one segment of the

chromosome (Fig. S20). This suggested that the differences between the two variants was

not the result of strong effects at a small number of promoters, but instead a broader

response to a general change in the distribution of DNA modifications, as suggested for

other epigenetic phase variants [27].

Consistent with the methylation pattern of RMV7$_{domi}$ reducing transformation efficiency

through altering cell physiology, many of the differentially-expressed loci identified by RNA-

seq encoded regulators. The *mgrA* gene, encoding a regulator of cell adhesion, was

expressed more highly in RMV7$_{domi}$ (Fig. S21). Consistent with previous observations, this

correlated with repression of the *rlrA* pilus islet [49] and activation of the *accBC-yqhY-amaP-*

csbD cell wall stress operon (IONPJBJN_01032-6) [50] in RMV7$_{domi}$. The cell-surface

chaperone-encoding gene *prsA* was also upregulated, as was the chaperone regulator gene

hrcA. The *groES-groEL* and *grpE-dnaK*-IONPJBJN_02152-*dnaJ* operons of the *hrcA*

regulon were also more highly expression in RMV7$_{domi}$, although this difference only

achieved genome-wide significance for IONPJBJN_02152 (Fig. S22; Table S3).

Another indicator of cell stress in RMV7$_{domi}$ was the increased activity of a phage-related

chromosomal island (PRCI; also known as a phage-inducible chromosomal island, or PICI)

integrated adjacent to *dnaN* (PRCI$_{dnaN}$; Fig. S23; Table S3). This is one of two PRCIs

associated with GPSC1 [7,28], the other being integrated near *uvrA* (Fig. 2). The regulatory

mechanisms of these elements are not thoroughly characterised [51], and in the absence of

a helper prophage, it is unclear exactly what signal may have triggered this increased

activity. Given integrative MGEs are likely under selection to reduce host cell transformability

[17], the increased activity of PRCI$_{dnaN}$ could drive inhibition of the competence machinery. A

further indicator of cell stress was the increased expression of the *ciaRH* two-component

system, which is known to inhibit competence [52]. Hence PRCI$_{dnaN}$ and *ciaRH* were the

primary candidates for causing the observed difference in transformability between the

RMV7 variants.

ManLMN links competence induction to sugar metabolism

The higher expression of the *ciaRH* genes in the non-transformable genotypes RMV7$_{domi}$

and RMV7$_{wt}$, relative to the transformable genotype RMV7$_{rare}$, was confirmed by qRT-PCR

(Fig S15). CiaRH binds at least 15 promoter sequences, five of which drive the expression of

cia-dependent small RNAs (csRNA) that suppress the induction of competence for

transformation through inhibiting expression of the *comC* gene, encoding CSP [53]. The

remaining ten drive the expression of protein coding sequences (CDSs) [53]. Although

csRNA5 was the only non-coding sequence that appeared to respond to *ciaRH* activation,

many of the CiaRH regulon CDSs were more highly expressed in RMV7$_{domi}$ compared to

RMV7$_{rare}$ pre-CSP (Fig. S24). These included the extracytoplasmic chaperone and serine

protease HtrA [53–55], which can block competence induction at low CSP concentrations

through degrading extracellular CSP [54,56]. In agreement with some previous studies,

elimination of *htrA* further increased the transformability of RMV7$_{rare}$ [54], but the same

mutation had no significant effect in RMV7$_{wt}$ (Fig S25). This suggests HtrA inhibits the
induction of competence, but is unlikely to explain much of the difference between these
variants. Similarly, knock out of *ciaRH* itself had little effect in either genotype (Fig. S25),
suggesting an alternative locus was causing the difference in transformation efficiency
between the variants.
6
Among the CiaRH regulon, the most significant difference in expression between RMV7$_{rare}$
and RMV7$_{domi}$ was the ManLMN carbon source importer operon (Table S3). The *manLMN*
genes were more highly expressed in RMV7$_{rare}$, as CiaRH binds promoter sequences within
the operon [37,53,57,58] and acts as a repressor [59]. Disrupting the *manLMN* locus
reduced the transformation efficiency of RMV7$_{rare}$ by >5-fold, while having little effect in
RMV7$_{wt}$ (Fig. 3A-B). To test whether the observed transformation differences reflected a
growth defect, *manLMN*::Janus mutants of both RMV7$_{wt}$ and RMV7$_{rare}$ were cultured in the
same rich media (Fig. S26). However, removal of *manLMN* had little effect on growth,
suggesting the transporter's effect on transformation was through regulation rather than
proliferation. Hence the lower expression of *manLMN* in RMV7$_{domi}$ accounts for some of its
reduced transformation efficiency relative to RMV7$_{rare}$.
18
ManLMN is a phosphotransferase system (PTS) transporter that can facilitate the import of
glucose, mannose, galactose, fructose, aminoglucose and *N*-acetylglucosamine (GlcNAc)
[60]. To test whether any effects were dependent on the imported substrate, growth curves
and transformation experiments were undertaken in the presence of these carbon source
supplements. Few significant differences in growth rate were observed, except a small
positive effect of GlcNAc on the growth of RMV7$_{wt}$ (Fig. S26). However, GlcNAc increased
transformation efficiency ~10-fold in RMV7$_{wt}$ (Fig. 3A) and ~2-fold in RMV7$_{rare}$ (Fig. 3B). In
both variants, this effect was dependent upon *manLMN*. This was confirmed through
disrupting, and then restoring, *manL* alone (Fig. S27). Therefore, uptake of GlcNAc by
ManLMN increased the transformability of RMV7.
29
_N_-acetylglucosamine activates competence through TfoX and Yjbk
This result suggested two parallels with the regulation of competence in gram-negative
bacteria. Competence in *Vibrio cholerae* is induced GlcNAc disaccharides [61], thought to be
generated from degradation of chitin [62]. This is mediated through the Transformation
Factor X (TfoX) protein [63,64]. An orthologue of this protein (also called Sxy) is also central
to regulation of competence in *Haemophilus influenzae* [65], in which it facilitates induction
via 3',5'-cyclic AMP (cAMP) signalling [66,67]. Intracellular concentrations of 3',5'-cAMP rise
in many proteobacteria when the primary glucose PTS transporter is inactive, as the

accumulation of phosphorylated EIIA PTS subunits stimulates adenylate cyclase activity

[68]. We investigated whether analogues of either of these pathways existed in *S.*

pneumoniae.

An orthologue of the N terminal domain of TfoX was annotated, but not described, in *S.*

pneumoniae ATCC 700669 [69]. In RMV7, this gene ($tfoX_{Spn}$; IONPJBJN_02097) is

conserved in the same position, two genes upstream of the *comEA* competence operon

(Fig. S28). The amino acid sequence corresponded to only the N-terminal region of the

gram-negative orthologue, and was predicted to form a four-strand beta sheet flanked by

alpha helices (Fig. S29). The gene could be both disrupted, and restored, in $RMV7_{rare}$.

Transformation assays using a panel of sugars demonstrated the elevated induction of

competence by GlcNAc in this variant was dependent on $TfoX_{Spn}$ (Fig. 3C).

A gene encoding a candidate adenylate cyclase, *yjbK*, was also identified in RMV7

(IONPJBJN_01639). This protein is predicted to have a β-barrel structure, as observed for

orthologous enzymes synthesising 3',5'-cAMP (Fig. S30). The *yjbK* gene could be both

disrupted, and restored, in $RMV7_{rare}$. Transformation assays with these genotypes

demonstrated the effect of GlcNAc on competence induction in $RMV7_{rare}$ also depended on

YjbK (Fig. 3D). However, no 3',5'-cAMP signalling pathway is known in Firmicutes [70].

Therefore, an ELISA was used to compare the 3',5'-cAMP levels in $RMV7_{wt}$, $RMV7_{rare}$

yjbK::Janus and $RMV7_{rare}$ *yjbK* restored, relative to *Escherichia coli* DH5α. High levels of

3',5'-cAMP were found in fractions collected from both exponential and stationary phases of

E. coli. However, the concentrations of 3',5'-cAMP detected in *S. pneumoniae* were close to

the lower detection threshold, and unaffected by *yjbK* presence (Fig. 3E). Additionally,

exogenous 3',5'-cAMP had no effect on transformation efficiencies in any RMV7 genotypes

(Fig. S31). Therefore, it is unlikely that YjbK's regulatory role is mediated through 3',5'-cAMP

production.

To test whether TfoX and YjbK had a similar role in mediating the larger response to GlcNAc

in $RMV7_{wt}$, both were also knocked out in this background. However, the results showed a

different pattern (Fig. 3F). The disruption of *tfoX* had no effect on GlcNAc's ability to induce

competence, whereas $RMV7_{wt}$ *yjbK*::Janus failed to activate competence unless GlcNAc or

sialic acid were supplemented in the media. These results were consistent with ManLMN

affecting competence through at least two pathways, one of which is dependent upon TfoX

and YjbK. Disruption of these genes in the laboratory strain R6x was also found to reduce

transformation efficiency, although detecting these effects required culturing in a low-sugar

chemically-defined medium (see Methods; Fig. S32).

Variation in mobile element activity associated with cell stress

Although ManLMN had a role in regulating the competence machinery, it could not account

for the large difference in transformability between RMV7$_{domi}$ and RMV7$_{rare}$ alone. MGEs

have been found to strongly affect levels of competence in *S. pneumoniae* and other species

[16,71–73]. Therefore we tested the hypothesis that the increased activity of PRCI$_{dnaN}$ in

RMV7$_{domi}$ may inhibit the competence of the host cell, in order to prevent the MGE being

deleted through homologous recombination [17]. A candidate gene (IONPJBJN_00496) with

the potential to limit homologous recombination encoded a protein similar to DNA damage-

inducible protein D (DinD), which inhibits RecA activity in *E. coli* [74]. However, the RMV7$_{wt}$

IONPJBJN_00496::Janus mutant did not show any increase in transformability (Fig. 4A).

To test whether any other part of the PRCI might inhibit transformation, the entire element

was removed, either with (RMV7$_{wt}$ PRCI$_{dnaN+att}$::Janus) or without (RMV7$_{wt}$ PRCI$_{dnaN}$::Janus)

the flanking *att* sites. In both cases, a ~5-fold increase in transformation rates was observed

(Fig. 4A). This implied the PRCI encoded an activity that inhibited the activation of the

competence system. To identify where this was located within the MGE, four large mutations

were generated within the PRCI: one removing the regulatory genes; one removing the

regulatory genes and IONPJBJN_00496; one removing the replication genes; and one

removing the integration, regulatory and replication genes. However, none of these

mutations had such a large effect on transformation rates as the elimination of the entire

element (Fig. 4A). This suggested the inhibition of competence induction was not the

consequence of a single gene product, but instead the activity of the MGE itself.

A qRT-PCR assay was employed to test whether activation of PRCI$_{dnaN}$ could affect

transformation through upregulating other stress signals within the cell. This found neither

ciaR nor *manL* expression was altered when the PRCI was removed (Fig. 4C). However,

transcription of the chaperone regulator *hrcA* was halved in the absence of the MGE,

consistent with the higher *hrcA* expression detected when the MGE was more active in

RMV7$_{domi}$ pre-CSP (Table S3). This indicated that the repression of competence by PRCI$_{dnaN}$

could occur indirectly, through MGE-driven elevation of the protein misfolding responses in

the cell.

Repression of competence by conserved chaperone proteins

HrcA is encoded by a gene within the *dnaK* heat shock operon, and it autoregulates through

repressing this chaperone-encoding gene cluster through binding the CIRCE motif [75]. In *S.*

pneumoniae, HrcA binding of CIRCE is reduced at elevated temperatures, relieving its

repression of the *dnaK* and *groEL* operons, enabling a heat shock response [76]. By

contrast, Ca^{2+} ions facilitate HrcA-CIRCE motif binding, inhibiting chaperone expression [77].

Both low Ca^{2+} concentrations and extreme temperatures inhibit the induction of competence

[78]. To test whether these effects on transformation efficiency were mediated through their

reduction in HrcA-CIRCE interactions, the *hrcA* gene was disrupted with a Janus cassette,

then restored, in RMV7$_{rare}$.

Disruption of *hrcA* was confirmed to relieve repression of *dnaK* and *groL* expression through

qRT-PCR, with no changes in *ciaR* expression, demonstrating that any effects on

transformation were not mediated through CiaRH (Fig. 5A). In the absence of Ca^{2+}, the

hrcA::Janus mutant had a lower transformation efficiency than either RMV7$_{rare}$, or RMV7$_{rare}$

hrcA restored (Fig. 5B). This difference grew as the Ca^{2+} concentration increased, consistent

with HrcA-CIRCE binding increasing transformation efficiency. These effects could be

reproduced in *S. pneumoniae* R6x grown in a chemically-defined medium (Fig. S33).

Unfortunately, the essential nature of the chaperones regulated by HrcA prevented any

further identification of the specific mechanism by which competence was inhibited.

Replicating previous observations, a 40 °C heat shock reduced the transformation efficiency

of both RMV7$_{rare}$ and RMV7$_{wt}$ (Fig. 5C). However, the heat shock increased the

transformation efficiency of RMV7$_{rare}$ *hrcA*::Janus, demonstrating the competence machinery

itself to be resilient to misfolding at elevated temperatures. Hence the effect of elevated

temperature on transformation efficiency appears to be mediated through the decreased

CIRCE-HrcA interactions. This cannot explain the increased transformation efficiency of the

hrcA::Janus cells after heat shock, but the growth curves of these genotypes at 40 °C

suggests the loss of *hrcA* has pleiotropic effects on pneumococcal physiology (Fig. S34).

To check that this regulation was independent of the ManLMN-mediated effects on

competence, the transformation efficiency of the RMV7$_{rare}$ *hrcA*::Janus *manLMN*::*cat* double

mutant was compared to that of the progenitor genotype, and the single mutants (Fig. 5D).

This demonstrated an approximately five-fold decrease in transformation for each single

mutant, and a ~25-fold reduction for the double mutant. This is consistent with HrcA and

ManLMN both activating competence independently.

Discussion

The transformation efficiency of *S. pneumoniae* is heterogeneous across populations

[9,10,79] and over the history of individual strains [8,80]. While the competence machinery

itself is highly conserved across the species [2], studying the regulation of competence in

highly-transformable laboratory isolates will inevitably miss mechanisms that affect this process in clinical isolates. *S. pneumoniae* RMV7 is a representative of GPSC1 [39], which has diversified through homologous recombination exceptionally quickly, enabling vaccine escape and the acquisition of antibiotic resistance loci [7]. However, even within a single isolate of this strain, only a minority of the population were highly transformable *in vitro*. Phase variable systems enable such phenotypic heterogeneity to persist within a near-isogenic population [27,32], resulting in a "bet hedging" strategy in RMV7 that mimics the heterogeneity in transformation efficiency of *Bacillus subtilis* cultures [33,81,82]. Although the competence state is regarded as a stress response analogous to the SOS response in other bacteria [48], this work suggests its activation requires a nutrient-rich milieu and an absence of abiotic stresses, such as elevated temperature. An alternative stress state, observed in RMV7$_{domi}$, upregulates the chaperones and proteases needed to cope with protein misfolding, and proteins likely to ameliorate cell wall disruption, without the protein synthesis and energy expenditure of competence induction. A mixture of stress-response strategies may enable the pneumococcus to survive a greater diversity of environmental challenges.

This heterogeneity might be unexpected, given CSP's role in synchronising competence induction across pneumococcal populations. The competence inhibitor CiaRH inhibits pheromone signalling through induction of five csRNAs, inhibiting intracellular CSP production [53,83,84], and the upregulation of HtrA, which degrades extracellular CSP [54]. If the pheromone were freely diffusing throughout a population, such inhibition of competence would be homogenous across cells. However, the downregulation of *manLMN* by CiaRH reduces cells' ability to respond to a CSP stimulus, enabling heterogeneity within a single population.

Despite its major role in regulating the competence machinery, ManLMN was dispensable for efficient growth in rich media. This transporter is highly conserved across *S. pneumoniae* (Fig. S35; Table S4), and orthologues serve as the main glucose transporter across many Firmicutes, including other streptococci, *Lactococcus lactis* and *Listeria monocytogenes* [60]. In *S. pneumoniae*, it acts as the central metabolic regulator, with the ManL EIIAB component important in controlling the carbon catabolite preferences of *S. pneumoniae* cells [85]. Loss of ManLMN in *Streptococcus mutans* was associated with decreased biofilm formation and transformability [86], suggesting its regulatory role is likely to be shared across many streptococci.

While *S. pneumoniae* encodes a plethora of transporters [26], experiments in chemically-

defined media suggest ManLMN may be the only effective means of importing GlcNAc [60].

Crucially, ManLMN's role as the primary glucose importer means GlcNAc can act as an

intracellular signal without being subject to catabolite repression. On importation, GlcNAc is

converted to the cytotoxic compound GlcNAc-6-phosphate. This is typically used for cell wall

biosynthesis in proliferating cells, but any excess must be converted to fructose-6-phosphate

for glycolysis [87]. Hence GlcNAc is a uniquely informative signal of nutrient availability

relative to cell proliferation [88]. The increased availability of GlcNAc by *V. cholerae* has

been suggested to signal the colonisation of a copepod's chitinous surface [89].

Analogously, GlcNAc is a major component of mucins covering the upper respiratory tract

[90], although *S. pneumoniae* is an obligate commensal of the human nasopharynx, and

likely derives little ecological information from GlcNAc's availability from host structures. An

alternative interpretation is locally-elevated concentrations of GlcNAc may be indicative of

lysis of bacterial cell walls, as suggested in streptomycetes [91]. As *V. cholerae* and *S.*

pneumoniae both co-ordinate fratricide with their competence machinery [92], this would

represent a parsimonious explanation for both using GlcNAc as a signal inducing

competence.

These two pathogens also both regulate competence through quorum sensing (Fig. 6), and

when combined with elevated GlcNAc concentrations, the paired signals would imply high

local densities of conspecific bacteria undergoing cell wall lysis. This would indicate the

availability of genomic DNA suitable for integration through homologous recombination.

There are further similarities in these species' GlcNAc-signalling pathways. One shared

component is a TfoX protein, present in many bacterial phyla (Fig. S36). In gram-negative

bacteria, TfoX affects transcription [72] and responds to 3',5'-cAMP. It is unclear whether the

shorter orthologue found in gram-positive bacteria could fulfil either of these roles,

particularly given the apparent absence of 3,5'-cAMP in pneumococci, to whom it is likely

toxic [70]. The component of the pneumococcal regulatory system most similar to enzymes

generating 3',5'-cAMP was YjbK$_{Spn}$, which belongs to a recently-defined subset of CYTH

proteins [93,94]. However, like its orthologues in *B. subtilis* [95] and *Staphylococcus aureus*

[70], the function of YjbK$_{Spn}$ is unknown. Both TfoX and YjbK are conserved in *S.*

pneumoniae (Fig S35), although *tfoX* is absent from ~12% of isolates [96]. However, neither

is strongly upregulated by CSP, making it difficult to identify their regulatory functions

through gene expression studies alone (Fig. S37).

In vitro transformation of *S. pneumoniae* was also activated by the addition of CaCl$_2$, with

cells most sensitive to changes in Ca^{2+} concentration in the 0-5 mM range. This spans the

relevant concentrations of the ion in mucus [97] and interstitial fluid, which rise on cell

necrosis and inflammation [98]. Ca^{2+} was previously suggested to aid the translocation of

DNA molecules across the plasma membrane [99]. We found its effect was dependent upon

HrcA, the binding of which to CIRCE motifs is promoted by higher Ca^{2+} concentrations [77].

Both the deletion of *hrcA*, and the increased expression in RMV7$_{domi}$ associated with PRCI-

associated stress, were associated with reduced transformation efficiency. This suggests the

activation of competence is mediated by HrcA-CIRCE complexes. Any involvement of the

chaperones regulated by HrcA could not be tested in this genetic study. In other species,

orthologues of the chaperones regulated by HrcA-CIRCE have been found to block sigma

factor switching, as required for the activation of late competence genes [100]. However, the

change in the expression of chaperone genes in response to alterations in PRCI replication

was small compared to that of *hrcA*. Hence HrcA may represent a physiologically-relevant

competence regulator that acts through a chaperone-independent route.

The level of complexity of competence regulation in *S. pneumoniae* is similar to that in

Bacillus subtilis [101], as well as *Vibrio cholerae* [102] and other gram-negative bacteria

[103]. Such signalling networks may be the product of adaptive evolution, selecting for the

integration of many external cues to trigger competence only under optimal conditions.

However, this would require that *S. pneumoniae* benefits from transformation when GlcNAc

is abundant and there is no protein misfolding stress, whereas *H. influenzae* benefits during

starvation conditions [104–106], when both species are frequently isolated within the same

nasopharynx [107]. Perhaps tellingly, the regulatory roles of ManLMN and HrcA are

moonlighting functions. These are secondary physiological activities separate from their

primary physiological role [108]: the transporter activated transformation in media conditions

under which it did not affect growth (Fig. 3), and the chaperone regulator activated

transformation without being necessary for the correct folding of the proteins involved (Fig.

5).

Alternatively, the complexity of competence regulation may be a product of the conflict

between organism-level adaptation and gene-level selection. The conservation of the

competence machinery suggests individual cells must benefit from transformation. Yet any

gene in which variation can accumulate is expected to evolve into forms that selfishly reduce

transformation efficiency, as this limits the rate at which such loci can be replaced within a

recombination recipient by competing alleles from donors [17,105]. Previously documented

for MGEs [6,15–17], this study suggests non-mobile variable loci (e.g. the *tvr* locus) may

also evolve to suppress transformation. Counteracting this effect in pneumococci are the

moonlighting conserved regulators ManLMN and HrcA. Such proteins are well-suited to

synchronising and activating competence, because purifying selection on their primary

function impedes the emergence of selfish mutants that inhibit competence. Such conflicting

gene-level pressures may explain why so many loci regulate the competence machinery,

and why transformation efficiency is so variable across and within species [109].

Methods and materials

Cell culture and mutagenesis

Genotypes used in this study are described in Table S5. Unless otherwise stated,

encapsulated *S. pneumoniae* were cultured statically at 35 °C supplemented with 5% CO_2 in

10 mL of a mixed liquid media, consisting of a 2:3 ratio of Todd-Hewitt media with 0.5%

yeast extract (Sigma-Aldrich), and Brain-Heart Infusion media (Sigma-Aldrich).

Transformation experiments with *S. pneumoniae* R6x derivatives used a chemically-defined

medium, consisting of disodium β-glycerophosphate (20 g L^{-1}; Sigma-Aldrich), sodium

pyruvate (0.1 g L^{-1}; Fluorochem), choline (0.001 g L^{-1}; Alfa Aesar), cysteine (0.4 g L^{-1}; Tokyo

Chemical Industry UK), glucose (3.8 mM; Sigma-Aldrich) and galactose (12 mM; Sigma-

Aldrich). Carbohydrates were otherwise added as a supplement to liquid media at a final

concentration of 33 mM, unless otherwise specified.

Culturing on solid media used Todd-Hewitt media supplemented with 0.5% yeast extract.

For the generation of mutants, 0.8-1 kb PCR fragments of regions flanking the gene of

interest were amplified with an added *Bam*HI or *Eco*RV restriction enzyme site.

Oligonucleotide sequences are listed in Table S6. PCR products were digested with the

appropriate restriction enzymes (Promega) at 35 °C for 2-4 hours, and then ligated to the

chosen antibiotic markers using T4 DNA ligase (Invitrogen). Ligations were used as

templates for further amplifications using PCR, which generated the constructs used for

mutagenesis through transformation of competent pneumococcal cells.

Transformation assays

One millilitre of bacterial culture was collected at an OD_{600} between 0.15-0.25. Cells were

then incubated for 2 hours at 35 °C with 5 μl of 500 mM $CaCl_2$ (Sigma), 250 ng of

competence stimulating peptide 1 (CSP-1; Cambridge Bioscience Ltd) and 100 ng of the

purified *rpoB* gene, containing a base substitution that conferred resistance to rifampicin

[110]. After two hours of incubation at 35 °C, a volume of between 1 and 100 μl of the

transformed culture was spread on an agar plate supplemented with 4 μg mL^{-1} of rifampicin

(Fisher Scientific). For precise quantification of transformation frequencies, 1 μl of a 10^3-fold

dilution of the same culture was spread on a non-selective plate in parallel. Colonies were

counted after 24 hours of incubation at 35 °C with 5% CO_2. For other transformations, the

same conditions were used, and cells were incubated on agar plates supplemented with the

appropriate selective antibiotics: kanamycin (Sigma-Aldrich) at 600 μg mL^{-1}, or

chloramphenicol (Sigma-Aldrich) at 4 μg mL^{-1}.

A logistic curve was used to analyse the relationship between transformation efficiency (in

transformants per 10^4 cfu), t, and the volume of 25 mM $CaCl_2$ added to the 1 mL of cells, v.

The fitted function was:

10
$$t = \frac{a}{1 + e^{-b(v-c)}}$$

The values of the variables a, b and c were estimated using the Levenberg-Marquardt

nonlinear least-squares algorithm in the minpack R package [111], using the starting values

of 10 (increased to 500 for *S. pneumoniae* R6x), 1 and 0.5, respectively. The confidence

intervals were calculated through refitting the function to 999 bootstrapped samples using

the car R package [112].

Surface adhesion assays and growth curves

To measure growth curves, 2×10^4 cells from titrated frozen stocks were grown in mixed

liquid media in 96-well microtiter plates at 35°C with 5% CO_2 for 20 h. Measurements of

OD_{600} were taken at 30 minutes intervals over 16 hours using the FLUOstar Omega

microplate reader (BMG LABTECH). Three replicate wells were used for each tested

genotype in each experiment. For measuring adhesion to abiotic surfaces, at the end of the

growth curve incubation, the microtitre plate was submerged in water and dried for 10

minutes. Then 150 μl of a freshly-diluted 0.1% crystal violet solution (Scientific Laboratory

Supplies) was added to each well, followed by incubation for 30 minutes at room

temperature. Each well was then washed by repeatedly submerging the plate in water to

remove excess crystal violet. The plate was incubated at room temperature in an inverted

position for four hours. Subsequently, 150 μl of 30% acetic acid (Honeywell) was added to

each well. Adherence was quantified as OD_{550} across replicate wells, measured by a

FLUOstar Omega plate reader.

Preparations of RNA samples and quantitative PCR

Three replicate cultures of RMV7 tvr_{domi}::Janus and RMV7 tvr_{rare}::Janus were grown in 25 mL

of mixed liquid media until they reached an OD_{600} of 0.15. A 5 mL sample of bacterial cells

was collected and 50 μL 250 ng mL^{-1} CSP1 was added to the remaining culture. Further 5

mL samples were taken from each culture 10 and 20 min post-CSP addition. Each sample

was immediately treated with 10 mL RNAprotect (Qiagen) and incubated at room

temperature for 5 min. Cell were then pelleted by centrifugation at 3,220 *g* for 10 min. RNA

was extracted from the washed pellets using the SV Total RNA Isolation System (Promega)

according to the manufacturer's instructions. The extracted RNA was used for RNA

sequencing or qRT-PCR.

All qRT-PCR experiments were conducted as described previously [35], using 0.2 μg of RNA

to generate cDNA with the First-Strand III cDNA synthesis kit (Invitrogen). Reactions used

the PowerUp™ SYBR™ Green Master Mix (Thermo Fisher) and the QuantStudio™ 7 Flex

Real-Time PCR System (Applied Biosystems).

Generation and analysis of RNA-seq data

RNA samples were quantified using an Agilent Bioanalyser RNA Nano Chip (Agilent

Technologies), and treated with the RiboZero® rRNA Removal Kit for Bacteria (Illumina) to

deplete rRNA. The samples were then cleaned with Agencourt RNAClean Beads (Beckman

Coulter). Sequencing libraries were generated with the NEBNext® Ultra II Directional Library

Prep Kit for Illumina (New England BioLabs), modified to use oligonucleotide sequences

appropriate for the sequencing pipelines of the Wellcome Sanger Institute. The library was

amplified through nine PCR cycles using the Kapa HiFi HotStart Ready Mix (Roche) to

generate sufficient material for sequencing. All eighteen samples were sequenced as a

multiplexed library on a single lane of a HiSeq 4000 sequencing system (Illumina),

generating 200 nt paired end reads.

The set of genes used for expression analysis were the 2,088 protein coding sequences

annotated on the *S. pneumoniae* RMV7$_{domi}$ genome (accession code OV904788), and the

81 non-coding RNAs predicted by infernal version 1.1.2 [113] using the Rfam database

[114]. RNA-seq reads were aligned to these sequences using kallisto version 0.46.2 [115],

using default settings and 100 bootstraps. Differential gene expression analysis used sleuth

version 0.30 [116]. Wald tests were conducted to compare the pre-CSP samples for

RMV7$_{domi}$ and RMV7$_{rare}$, and to compare the 10 min and 20 min post-CSP samples for each

genotype to the corresponding pre-CSP samples. Visualisation and plotting of data used the

genoplotR [117], circlize [118], cowplot [119], ggpubr [120] and tidyverse [121] packages.

Quantification of 3',5'-cAMP production

Quantification of 3',5'-cAMP used the Cyclic AMP XP® assay kit (Cell Signalling

Technology). Samples were harvested in the exponential (OD$_{600}$ of 0.2) and stationary

(OD$_{600}$ of 0.5) phases of *S. pneumoniae* statically-grown cultures. *E. coli* DH5α was grown in

25 mL Luria-Bertani media (Sigma-Aldrich) at 37 °C, shaken at 250 revolutions per minute,

and samples were harvested in the exponential (OD$_{600}$ of 0.3) and stationary (OD$_{600}$ of 1.0)

phases. Cells were pelleted through centrifugation at 3,220 *g* for 10 min. The cell pellets

were re-suspended in 850 μL of the kit's lysis buffer and 150 μl of lysozyme (10 mg mL^{-1}),

followed by incubation at 35 °C for 20 minutes. Cells were pelleted by centrifugation at 9,500

g for 5 minutes. The supernatants were collected, and the cell pellets were resuspended in

400 μl phosphate-buffered saline. The protein concentrations of all the collected samples

were adjusted to 400 μg mL^{-1} using the Qubit protein broad range assay kits (Qiagen). The

Cyclic AMP XP® kit ELISA plates were prepared according to the manufacturer's protocols,

and a 50 μl sample of supernatant from each tested culture was loaded in each well. The

OD$_{450}$ was measured using the FLUOstar Omega plate reader. Concentrations of 3',5'-cAMP

were calculated using a standard curve, according to the manufacturer's instructions.

Analysis of motif distribution and sequence diversity

The overall distribution of methylation motifs was calculated using DistAMo [122]. The

calculation of distances between coding sequences and motifs used biopython.

The diversity of genes encoding regulatory proteins within *S. pneumoniae* used previously-

calculated pairwise genetic distances [123]. Only genes found in at least half the isolates in

the collection were used, to avoid including genes for which diversity might be inferred to be

low, due to a limited samples size for detecting polymorphisms.

Proteins containing the TfoX N-terminal domain were identified using EMBL SMART [124].

These amino acid sequences were aligned with MAFFT [125], and a phylogeny generated

with Fasttree2 [126] using default settings. Proteins were assigned to Families using the

NCBI Taxonomy [127].

References

1. Griffith F. The Significance of Pneumococcal Types. J Hyg (Lond). 1928;27: 113–159. doi:10.1017/S0022172400031879

2. Johnston C, Campo N, Bergé MJ, Polard P, Claverys JP. *Streptococcus pneumoniae*, le transformiste. Trends Microbiol. 2014;22: 113–9. doi:10.1016/j.tim.2014.01.002

3. Dowson CG, Hutchison A, Brannigan JA, George RC, Hansman D, Linares J, et al. Horizontal transfer of penicillin-binding protein genes in penicillin-resistant clinical isolates of *Streptococcus pneumoniae*. Proc Natl Acad Sci. 1989;86: 8842–8846. doi:10.1073/pnas.86.22.8842

4. Sibold C, Henrichsen J, Konig A, Martin C, Chalkley L, Hakenbeck R. Mosaic *pbpX* genes of major clones of penicillin-resistant *Streptococcus pneumoniae* have evolved from *pbpX* genes of a penicillin-sensitive *Streptococcus oralis*. Mol Microbiol. 1994/06/01. 1994;12: 1013–1023.

5. Dowson CG, Coffey TJ, Kell C, Whiley RA. Evolution of penicillin resistance in *Streptococcus pneumoniae*; the role of *Streptococcus mitis* in the formation of a low affinity PBP2B in *S. pneumoniae*. Mol Microbiol. 1993/08/01. 1993;9: 635–643. doi:10.1111/j.1365-2958.1993.tb01723.x

6. D'Aeth JC, van der Linden MPG, McGee L, de Lencastre H, Turner P, Song J-H, et al. The role of interspecies recombination in the evolution of antibiotic-resistant pneumococci. Elife. 2021;10: e67113. Available: https://doi.org/10.7554/eLife.67113

7. Croucher NJ, Chewapreecha C, Hanage WP, Harris SR, McGee L, van der Linden M, et al. Evidence for soft selective sweeps in the evolution of pneumococcal multidrug resistance and vaccine escape. Genome Biol Evol. 2014;6: 1589–1602. doi:10.1093/gbe/evu120

8. Golubchik T, Brueggemann AB, Street T, Gertz RE, Spencer CCA, Ho T, et al. Pneumococcal genome sequencing tracks a vaccine escape variant formed through a multi-fragment recombination event. Nat Genet. 2012;44: 352–355. doi:10.1038/ng.1072

9. Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, et al. Population genomics of post-vaccine changes in pneumococcal epidemiology. Nat Genet. 2013;45: 656–663. doi:10.1038/ng.2625

10. Croucher NJ, Mitchell AM, Gould KA, Inverarity D, Barquist L, Feltwell T, et al. Dominant role of nucleotide substitution in the diversification of serotype 3 pneumococci over decades and during a single infection. PLoS Genet. 2013;9: e1003868. doi:10.1371/journal.pgen.1003868

11. Evans BA, Rozen DE. Significant variation in transformation frequency in

Streptococcus pneumoniae. ISME J. 2013;7: 791–799. doi:10.1038/ismej.2012.170

12. Hsieh YC, Wang JT, Lee W Sen, Hsueh PR, Shao PL, Chang LY, et al. Serotype competence and penicillin resistance in *Streptococcus pneumoniae*. Emerg Infect Dis. 2006;12: 1709–1714. doi:10.3201/eid1211.060414

13. Joloba ML, Kidenya BR, Kateete DP, Katabazi FA, Muwanguzi JK, Asiimwe BB, et al. Comparison of transformation frequencies among selected *Streptococcus pneumoniae* serotypes. Int J Antimicrob Agents. 2010;36: 124–128. doi:10.1016/j.ijantimicag.2010.03.024

14. Yother J, McDaniel LS, Briles DE. Transformation of encapsulated *Streptococcus pneumoniae*. J Bacteriol. 1986;168: 1463–1465. doi:10.1128/jb.168.3.1463-1465.1986

15. Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, et al. Rapid pneumococcal evolution in response to clinical interventions. Science (80-). 2011/01/29. 2011;331: 430–434. doi:10.1126/science.1198545

16. Croucher NJ, Hanage WP, Harris SR, McGee L, van der Linden M, de Lencastre H, et al. Variable recombination dynamics during the emergence, transmission and "disarming" of a multidrug-resistant pneumococcal clone. BMC Biol. 2014;12: 49. doi:10.1186/1741-7007-12-49

17. Croucher NJ, Mostowy R, Wymant C, Turner P, Bentley SD, Fraser C. Horizontal DNA Transfer Mechanisms of Bacteria as Weapons of Intragenomic Conflict. Barton NH, editor. PLOS Biol. 2016;14: e1002394. doi:10.1371/journal.pbio.1002394

18. Johnston C, Martin B, Fichant G, Polard P, Claverys JP. Bacterial transformation: Distribution, shared mechanisms and divergent control. Nat Rev Microbiol. 2014;12: 181–196. doi:10.1038/nrmicro3199

19. Håvarstein LS, Coomaraswamy G, Morrison DA. An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. Proc Natl Acad Sci. 1995;92: 11140 LP – 11144. doi:10.1073/pnas.92.24.11140

20. Alloing G, Martin B, Granadel C, Claverys J-P. Development of competence in *Streptococcus pneumoniae*: pheromone autoinduction and control of quorum sensing by the oligopeptide permease. Mol Microbiol. 1998;29: 75–83. doi:https://doi.org/10.1046/j.1365-2958.1998.00904.x

21. Lee MS, Morrison DA. Identification of a New Regulator in *Streptococcus pneumoniae* Linking Quorum Sensing to Competence for Genetic Transformation. J Bacteriol. 1999;181: 5004–5016. doi:10.1128/JB.181.16.5004-5016.1999

22. Campbell EA, Choi SY, Masure HR. A competence regulon in *Streptococcus pneumoniae* revealed by genomic analysis. Mol Microbiol. 1998;27: 929–939.

doi:10.1046/j.1365-2958.1998.00737.x

23. Pestova E V, Morrison DA. Isolation and characterization of three *Streptococcus pneumoniae* transformation-specific loci by use of a *lacZ* reporter insertion vector. J Bacteriol. 1998;180: 2701–2710. doi:10.1128/JB.180.10.2701-2710.1998

24. Sung CK, Morrison DA. Two Distinct Functions of ComW in Stabilization and Activation of the Alternative Sigma Factor ComX in *Streptococcus pneumoniae*. J Bacteriol. 2005;187: 3052–3061. doi:10.1128/JB.187.9.3052-3061.2005

25. Weiser JN, Kapoor M. Effect of intrastrain variation in the amount of capsular polysaccharide on genetic transformation of *Streptococcus pneumoniae*: Implications for virulence studies of encapsulated strains. Infect Immun. 1999;67: 3690–3692.

26. Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, et al. Complete genome sequence of a virulent isolate of *Streptococcus pneumoniae*. Science (80-). 2001;293: 498–506. doi:10.1126/science.1061217

27. Manso AS, Chai MH, Atack JM, Furi L, De Ste Croix M, Haigh R, et al. A random six-phase switch regulates pneumococcal virulence via global epigenetic changes. Nat Commun. 2014;5: 5055. doi:10.1038/ncomms6055

28. Croucher NJ, Coupland PG, Stevenson AE, Callendrello A, Bentley SD, Hanage WP. Diversification of bacterial genome content through distinct mechanisms over different timescales. Nat Commun. 2014;5: 5471.

29. Li JW, Li J, Wang J, Li C, Zhang JR. Molecular mechanisms of *hsdS* inversions in the *cod* locus of *Streptococcus pneumoniae*. J Bacteriol. 2019;201: e00581-18. doi:10.1128/JB.00581-18

30. Li J, Li JW, Feng Z, Wang J, An H, Liu Y, et al. Epigenetic Switch Driven by DNA Inversions Dictates Phase Variation in *Streptococcus pneumoniae*. PLoS Pathog. 2016;12: e1005762. doi:10.1371/journal.ppat.1005762

31. Wang J, Li JW, Li J, Huang Y, Wang S, Zhang JR. Regulation of pneumococcal epigenetic and colony phases by multiple two-component regulatory systems. PLoS Pathog. 2020;16: e1008417. doi:10.1371/journal.ppat.1008417

32. De Ste Croix M, Vacca I, Kwun MJ, Ralph JD, Bentley SD, Haigh R, et al. Phase-variable methylation and epigenetic regulation by type I restriction-modification systems. FEMS Microbiol Rev. 2017;41: S3–S15. doi:10.1093/femsre/fux025

33. Veening J-W, Smits WK, Kuipers OP. Bistability, epigenetics, and bet-hedging in bacteria. Annu Rev Microbiol. 2008;62: 193–210. doi:10.1146/annurev.micro.62.081307.163002

34. Quillin SJ, Seifert HS. *Neisseria gonorrhoeae* host adaptation and pathogenesis. Nat Rev Microbiol. 2018;16: 226–240. doi:10.1038/nrmicro.2017.169

35. Kwun MJ, Oggioni MR, De Ste Croix M, Bentley SD, Croucher NJ. Excision-

reintegration at a pneumococcal phase-variable restriction-modification locus drives within- and between-strain epigenetic differentiation and inhibits gene acquisition. Nucleic Acids Res. 2018;46: 11438–11453. doi:10.1093/nar/gky906

36. Kwun MJ, Oggioni MR, Bentley SD, Fraser C, Croucher NJ. Synergistic Activity of Mobile Genetic Element Defences in *Streptococcus pneumoniae*. Genes (Basel). 2019;10: 707.

37. Slager J, Aprianto R, Veening JW. Refining the pneumococcal competence regulon by RNA sequencing. J Bacteriol. 2019;201: e00780-18. doi:10.1128/JB.00780-18

38. Sung CK, Li H, Claverys JP, Morrison DA. An *rpsL* cassette, Janus, for gene replacement through negative selection in *Streptococcus pneumoniae*. Appl Environ Microbiol. 2001;67: 5190–5196.

39. Gladstone RA, Lo SW, Lees JA, Croucher NJ, van Tonder AJ, Corander J, et al. International genomic definition of pneumococcal lineages, to contextualise disease, antibiotic resistance and vaccine impact. EBioMedicine. 2019;43: 338–346. doi:10.1016/j.ebiom.2019.04.021

40. Trappetti C, Kadioglu A, Carter M, Hayre J, Iannelli F, Pozzi G, et al. Sialic Acid: A Preventable Signal for Pneumococcal Biofilm Formation, Colonization, and Invasion of the Host. J Infect Dis. 2009;199: 1497–1505. doi:10.1086/598483

41. Li Y, Croucher NJ, Thompson CM, Trzciński K, Hanage WP, Lipsitch M. Identification of pneumococcal colonization determinants in the stringent response pathway facilitated by genomic diversity. BMC Genomics. 2015;16: 369. doi:10.1186/s12864-015-1573-6

42. Lamarche MG, Wanner BL, Crépin S, Harel J. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. FEMS Microbiol Rev. 2008;32: 461–473. doi:10.1111/j.1574-6976.2008.00101.x

43. Aggarwal SD, Eutsey R, West-Roberts J, Domenech A, Xu W, Abdullah IT, et al. Function of BriC peptide in the pneumococcal competence and virulence portfolio. PLOS Pathog. 2018;14: e1007328. Available: https://doi.org/10.1371/journal.ppat.1007328

44. Peterson SN, Sung CK, Cline R, Desai B V, Snesrud EC, Luo P, et al. Identification of competence pheromone responsive genes in *Streptococcus pneumoniae* by use of DNA microarrays. Mol Microbiol. 2004;51: 1051–1070. doi:https://doi.org/10.1046/j.1365-2958.2003.03907.x

45. Majchrzykiewicz JA, Kuipers OP, Bijlsma JJE. Generic and specific adaptive responses of *Streptococcus pneumoniae* to challenge with three distinct antimicrobial peptides, bacitracin, LL-37, and nisin. Antimicrob Agents Chemother. 2009/11/16.

2010;54: 440–451. doi:10.1128/AAC.00769-09

46.   Johnston CHG, Soulet A-L, Bergé M, Prudhomme M, De Lemos D, Polard P. The
alternative sigma factor σ^X mediates competence shut-off at the cell pole in
Streptococcus pneumoniae. Blokesch M, Storz G, Blokesch M, Dubnau D, editors.
Elife. 2020;9: e62907. doi:10.7554/eLife.62907

47.   Weiser JN, Bae D, Epino H, Gordon SB, Kapoor M, A. ZL, et al. Changes in
Availability of Oxygen Accentuate Differences in Capsular Polysaccharide Expression
by Phenotypic Variants and Clinical Isolates of <i>Streptococcus pneumoniae</i>.
Infect Immun. 2001;69: 5430–5439. doi:10.1128/IAI.69.9.5430-5439.2001

48.   Claverys J-P, Prudhomme M, Martin B. Induction of Competence Regulons as a
General Response to Stress in Gram-Positive Bacteria. Annu Rev Microbiol. 2006;60:
451–475. doi:10.1146/annurev.micro.60.080805.142139

49.   Hemsley C, Joyce E, Hava DL, Kawale A, Camilli A. MgrA, an Orthologue of Mga,
Acts as a Transcriptional Repressor of the Genes within the *rlrA* Pathogenicity Islet in
Streptococcus pneumoniae. J Bacteriol. 2003;185: 6640–6647.
doi:10.1128/JB.185.22.6640-6647.2003

50.   Solano-Collado V, Espinosa M, Bravo A. Activator role of the pneumococcal *mga*-like
virulence transcriptional regulator. J Bacteriol. 2012;194: 197–4207.
doi:10.1128/JB.00536-12

51.   Martínez-Rubio R, Quiles-Puchalt N, Martí M, Humphrey S, Ram G, Smyth D, et al.
Phage-inducible islands in the Gram-positive cocci. ISME J. 2017;11: 1029–1042.
doi:10.1038/ismej.2016.163

52.   Guenzi E, Gasc A -M, Sicard MA, Hakenbeck R. A two-component signal-transducing
system is involved in competence and penicillin susceptibility in laboratory mutants of
Streptococcus pneumoniae. Mol Microbiol. 1994;12: 505–515. doi:10.1111/j.1365-
2958.1994.tb01038.x

53.   Halfmann A, Kovács M, Hakenbeck R, Brückner R. Identification of the genes directly
controlled by the response regulator CiaR in *Streptococcus pneumoniae*: Five out of
15 promoters drive expression of small non-coding RNAs. Mol Microbiol. 2007;66:
110–126. doi:10.1111/j.1365-2958.2007.05900.x

54.   Cassone M, Gagne AL, Spruce LA, Seeholzer SH, Sebert ME. The HtrA protease
from *Streptococcus pneumoniae* digests both denatured proteins and the
competence-stimulating peptide. J Biol Chem. 2012;287: 38449–38459.
doi:10.1074/jbc.M112.391482

55.   Sebert ME, Palmer LM, Rosenberg M, Weiser JN. Microarray-based identification of
htrA, a *Streptococcus pneumoniae* gene that is regulated by the CiaRH two-
component system and contributes to nasopharyngeal colonization. Infect Immun.

2002;70: 4059–4067. doi:10.1128/IAI.70.8.4059-4067.2002

56.   Stevens KE, Chang D, Zwack EE, Sebert ME. Competence in *Streptococcus*

pneumoniae is regulated by the rate of ribosomal decoding errors. MBio. 2011;2:

e00071-11. doi:10.1128/mBio.00071-11

57.   Dagkessamanskaia A, Moscoso M, Hénard V, Guiral S, Overweg K, Reuter M, et al.

Interconnection of competence, stress and CiaR regulons in *Streptococcus*

pneumoniae: Competence triggers stationary phase autolysis of *ciaR* mutant cells.

Mol Microbiol. 2004;51: 1071–1086. doi:10.1111/j.1365-2958.2003.03892.x

58.   Mascher T, Zähner D, Merai M, Balmelle N, De Saizieu AB, Hakenbeck R. The

Streptococcus pneumoniae cia regulon: CiaR target sites and transcription profile

analysis. J Bacteriol. 2003;185: 60–70. doi:10.1128/JB.185.1.60-70.2003

59.   Gómez-Mejia A, Gámez G, Hammerschmidt S. *Streptococcus pneumoniae* two-

component regulatory systems: The interplay of the pneumococcus with its

environment. Int J Med Microbiol. 2018;308: 722–737.

doi:10.1016/j.ijmm.2017.11.012

60.   Bidossi A, Mulas L, Decorosi F, Colomba L, Ricci S, Pozzi G, et al. A functional

genomics approach to establish the complement of carbohydrate transporters in

Streptococcus pneumoniae. PLoS One. 2012;7: e33320.

doi:10.1371/journal.pone.0033320

61.   Yamamoto S, Morita M, Izumiya H, Watanabe H. Chitin disaccharide (GlcNAc)2

induces natural competence in *Vibrio cholerae* through transcriptional and

translational activation of a positive regulatory gene $tfoX_{VC}$. Gene. 2010;457: 42–49.

doi:10.1016/j.gene.2010.03.003

62.   Meibom KL, Blokesch M, Dolganov NA, Wu CY, Schoolnik GK. Chitin induces natural

competence in *Vibrio cholerae*. Science (80-). 2005;310: 1824–1827.

doi:10.1126/science.1120096

63.   Lo Scrudato M, Blokesch M. A transcriptional regulator linking quorum sensing and

chitin induction to render *Vibrio cholerae* naturally transformable. Nucleic Acids Res.

2013;41: 3644–3658. doi:10.1093/nar/gkt041

64.   Yamamoto S, Izumiya H, Mitobe J, Morita M, Arakawa E, Ohnishi M, et al.

Identification of a chitin-induced small RNA that regulates translation of the *tfoX* gene,

encoding a positive regulator of natural competence in *Vibrio cholerae*. J Bacteriol.

2011;193: 1953–1965. doi:10.1128/JB.01340-10

65.   Williams PM, Bannister LA, Redfield RJ. The *Haemophilus influenzae sxy*-1 mutation

is in a newly identified gene essential for competence. J Bacteriol. 1994;176: 6789–

6794. doi:10.1128/jb.176.22.6789-6794.1994

66.   Chandler MS. The gene encoding cAMP receptor protein is required for competence

development in *Haemophilus influenzae* Rd. Proc Natl Acad Sci. 1992;89: 1626 LP –
1630. doi:10.1073/pnas.89.5.1626

67.   Dorocicz IR, Williams PM, Redfield RJ. The *Haemophilus influenzae* adenylate
4   cyclase gene: cloning, sequence, and essential role in competence. J Bacteriol.
5   1993;175: 7142–7149. doi:10.1128/jb.175.22.7142-7149.1993

6   68.   Wu R, Zhao M, Li J, Gao H, Kan B, Liang W. Direct regulation of the natural
7   competence regulator gene *tfoX* by cyclic AMP (cAMP) and cAMP receptor protein
8   (CRP) in Vibrios. Sci Rep. 2015;5: 14921. doi:10.1038/srep14921

69.   Croucher NJ, Walker D, Romero P, Lennard N, Paterson GK, Bason NC, et al. Role
of conjugative elements in the evolution of the multidrug-resistant pandemic clone
Streptococcus pneumoniae[Spain23F] ST81. J Bacteriol. 2008/12/31. 2009;191: 1480–
1489. doi:10.1128/jb.01343-08

70.   Zhang Y, Agrebi R, Bellows LE, Collet J-F, Kaever V, Gründling A. Evolutionary
Adaptation of the Essential tRNA Methyltransferase TrmD to the Signaling Molecule
3',5'-cAMP in Bacteria. J Biol Chem. 2017;292: 313–327.
doi:10.1074/jbc.M116.758896

71.   Durieux I, Ginevra C, Attaiech L, Picq K, Juan P-A, Jarraud S, et al. Diverse
conjugative elements silence natural transformation in *Legionella* species. Proc Natl
Acad Sci. 2019;116: 18613 LP – 18618. doi:10.1073/pnas.1909374116

72.   Stutzmann S, Blokesch M. Comparison of chitin-induced natural transformation in
pandemic *Vibrio cholerae* O1 El Tor strains. Environ Microbiol. 2020;22: 4149–4166.
doi:https://doi.org/10.1111/1462-2920.15214

73.   Dalia AB, Seed KD, Calderwood SB, Camilli A. A globally distributed mobile genetic
element inhibits natural transformation of *Vibrio cholerae*. Proc Natl Acad Sci.
2015;112: 10485–10490. doi:10.1073/pnas.1509097112

74.   Uranga LA, Balise VD, Benally C V, Grey A, Lusetti SL. The *Escherichia coli* DinD
protein modulates RecA activity by inhibiting postsynaptic RecA filaments. J Biol
Chem. 2011;286: 29480–29491. doi:10.1074/jbc.M111.245373

75.   Zuber U, Schumann W. CIRCE, a novel heat shock element involved in regulation of
heat shock operon *dnaK* of *Bacillus subtilis*. J Bacteriol. 1994;176: 1359–1363.
doi:10.1128/jb.176.5.1359-1363.1994

76.   Kim SN, Bae YG, Rhee DK. Dual regulation of *dnaK* and *groE* operons by HrcA and
Ca^{++} in *Streptococcus pneumoniae*. Arch Pharm Res. 2008;31: 462–467.
doi:10.1007/s12272-001-1179-4

77.   Kwon HY, Kim SN, Pyo SN, Rhee DK. Ca^{2+}-dependent expression of the CIRCE
regulon in *Streptococcus pneumoniae*. Mol Microbiol. 2005;55: 456 – 468.
doi:10.1111/j.1365-2958.2004.04416.x

78. Fox MS, Hotchkiss RD. Initiation of bacterial transformation. Nature. 1957;179: 1322–1325. doi:10.1038/1791322a0

79. Chewapreecha C, Harris SR, Croucher NJ, Turner C, Marttinen P, Cheng L, et al. Dense genomic sampling identifies highways of pneumococcal recombination. Nat Genet. 2014;46: 305–309. doi:10.1038/ng.2895

80. Mostowy R, Croucher NJ, Hanage WP, Harris SR, Bentley S, Fraser C. Heterogeneity in the Frequency and Characteristics of Homologous Recombination in Pneumococcal Evolution. PLoS Genet. 2014;10: e1004300. doi:10.1371/journal.pgen.1004300

81. Süel GM, Garcia-Ojalvo J, Liberman LM, Elowitz MB. An excitable gene regulatory circuit induces transient cellular differentiation. Nature. 2006;440: 545–550. doi:10.1038/nature04588

82. Maamar H, Raj A, Dubnau D. Noise in gene expression determines cell fate in *Bacillus subtilis*. Science. 2007;317: 526–529. doi:10.1126/science.1140818

83. Laux A, Sexauer A, Sivaselvarajah D, Kaysen A, Brückner R. Control of competence by related non-coding csRNAs in *Streptococcus pneumoniae* R6. Front Genet. 2015;6: 246. doi:10.3389/fgene.2015.00246

84. Schnorpfeil A, Kranz M, Kovács M, Kirsch C, Gartmann J, Brunner I, et al. Target evaluation of the non-coding csRNAs reveals a link of the two-component regulatory system CiaRH to competence control in *Streptococcus pneumoniae* R6. Mol Microbiol. 2013;89: 334–349. doi:10.1111/mmi.12277

85. Fleming E, Camilli A. ManLMN is a glucose transporter and central metabolic regulator in *Streptococcus pneumoniae*. Mol Microbiol. 2016;102: 467–487. doi:https://doi.org/10.1111/mmi.13473

86. Abranches J, Candella MM, Wen ZT, Baker H V., Burne RA. Different Roles of EIIAB^{Man} and EII^{Glc} in Regulation of Energy Metabolism, Biofilm Development, and Competence in *Streptococcus mutans*. J Bacteriol. 2006;188: 3748–3756. doi:10.1128/JB.00169-06

87. Kawada-Matsuo M, Mazda Y, Oogai Y, Kajiya M, Kawai T, Yamada S, et al. GlmS and NagB Regulate Amino Sugar Metabolism in Opposing Directions and Affect *Streptococcus mutans* Virulence. PLoS One. 2012;7: e33382. Available: https://doi.org/10.1371/journal.pone.0033382

88. Komatsuzawa H, Fujiwara T, Nishi H, Yamada S, Ohara M, McCallum N, et al. The gate controlling cell wall synthesis in *Staphylococcus aureus*. Mol Microbiol. 2004;53: 1221–1231. doi:10.1111/j.1365-2958.2004.04200.x

89. Meibom KL, Li XB, Nielsen AT, Wu C-Y, Roseman S, Schoolnik GK. The *Vibrio cholerae* chitin utilization program. Proc Natl Acad Sci U S A. 2004;101: 2524 LP –

2529. doi:10.1073/pnas.0308707101

90. Weiser JN, Ferreira DM, Paton JC. *Streptococcus pneumoniae*: Transmission, colonization and invasion. Nat Rev Microbiol. 2018;16: 355–367. doi:10.1038/s41579-018-0001-8

91. Rigali S, Titgemeyer F, Barends S, Mulder S, Thomae AW, Hopwood DA, et al. Feast or famine: the global regulator DasR links nutrient stress to antibiotic production by *Streptomyces*. EMBO Rep. 2008;9: 670–675. doi:https://doi.org/10.1038/embor.2008.83

92. Veening J-W, Blokesch M. Interbacterial predation as a strategy for DNA acquisition in naturally competent bacteria. Nat Rev Microbiol. 2017;15: 621–629. doi:10.1038/nrmicro.2017.66

93. Iyer LM, Aravind L. The catalytic domains of thiamine triphosphatase and CyaB-like adenylyl cyclase define a novel superfamily of domains that bind organic phosphates. BMC Genomics. 2002;3: 33. doi:10.1186/1471-2164-3-33

94. Vogt MS, Ngouoko Nguepbeu RR, Mohr MKF, Albers S-V, Essen L-O, Banerjee A. The archaeal triphosphate tunnel metalloenzyme *Sa*TTM defines structural determinants for the diverse activities in the CYTH protein family. J Biol Chem. 2021;297. doi:10.1016/j.jbc.2021.100820

95. Mamou G, Malli Mohan GB, Rouvinski A, Rosenberg A, Ben-Yehuda S. Early Developmental Program Shapes Colony Morphology in Bacteria. Cell Rep. 2016;14: 1850–1857. doi:https://doi.org/10.1016/j.celrep.2016.01.071

96. Croucher NJ, Finkelstein JA, Pelton SI, Parkhill J, Bentley SD, Lipsitch M, et al. Population genomic datasets describing the post-vaccine evolutionary epidemiology of *Streptococcus pneumoniae*. Sci Data. 2015;2: 150058. doi:10.1038/sdata.2015.58

97. Selvaraj S, Liu K, Robinson AM, Epstein VA, Conley DB, Kern RC, et al. *In Vivo* Determination of Mouse Olfactory Mucus Cation Concentrations in Normal and Inflammatory States. PLoS One. 2012;7: e39600. Available: https://doi.org/10.1371/journal.pone.0039600

98. Rossol M, Pierer M, Raulien N, Quandt D, Meusch U, Rothe K, et al. Extracellular Ca^{2+} is a danger signal activating the NLRP3 inflammasome through G protein-coupled calcium sensing receptors. Nat Commun. 2012;3: 1329. doi:10.1038/ncomms2339

99. Seto H, Tomasz A. Calcium-requiring step in the uptake of deoxyribonucleic acid molecules through the surface of competent pneumococci. J Bacteriol. 1976;126: 1113–1118. doi:10.1128/jb.126.3.1113-1118.1976

100. Gamer J, Multhaup G, Tomoyasu T, McCarty JS, Rüdiger S, Schönfeld HJ, et al. A cycle of binding and release of the DnaK, DnaJ and GrpE chaperones regulates

activity of the *Escherichia coli* heat shock transcription factor sigma32. EMBO J. 1996;15: 607–617. doi:https://doi.org/10.1002/j.1460-2075.1996.tb00393.x

101. Maier B. Competence and Transformation in *Bacillus subtilis*. Current Issues in Molecular Biology. 2020. doi:10.21775/cimb.037.057

102. Haycocks JRJ, Warren GZL, Walker LM, Chlebek JL, Dalia TN, Dalia AB, et al. The quorum sensing transcription factor AphA directly regulates natural competence in *Vibrio cholerae*. PLOS Genet. 2019;15: e1008362. Available: https://doi.org/10.1371/journal.pgen.1008362

103. Seitz P, Blokesch M. Cues and regulatory pathways involved in natural competence and transformation in pathogenic and environmental Gram-negative bacteria. FEMS Microbiol Rev. 2013;37: 336–363. doi:10.1111/j.1574-6976.2012.00353.x

104. Stewart GJ, Carlson CA. The Biology of Natural Transformation. Annu Rev Microbiol. 1986;40: 211–231. doi:10.1146/annurev.mi.40.100186.001235

105. Redfield RJ. Genes for breakfast: The have-your-cake-and-eat-it-too of bacterial transformation. J Hered. 1993;84: 400–404. doi:10.1093/oxfordjournals.jhered.a111361

106. Macfadyen LP, Chen D, Vo HC, Liao D, Sinotte R, Redfield RJ. Competence development by *Haemophilus influenzae* is regulated by the availability of nucleic acid precursors. Mol Microbiol. 2001;40: 700–707. doi:https://doi.org/10.1046/j.1365-2958.2001.02419.x

107. Bosch AATM, Biesbroek G, Trzcinski K, Sanders EAM, Bogaert D. Viral and Bacterial Interactions in the Upper Respiratory Tract. PLOS Pathog. 2013;9: e1003057. Available: https://doi.org/10.1371/journal.ppat.1003057

108. Jeffery CJ. Protein moonlighting: what is it, and why is it important? Philos Trans R Soc B Biol Sci. 2018;373: 20160523. doi:10.1098/rstb.2016.0523

109. Maughan H, Redfield RJ. Extensive variation in natural competence in *Haemophilus influenzae*. Evolution (N Y). 2009;63: 1852–1866. doi:https://doi.org/10.1111/j.1558-5646.2009.00658.x

110. Apagyi KJ, Fraser C, Croucher NJ. Transformation asymmetry and the evolution of the bacterial accessory genome. Mol Biol Evol. 2018;35: 575–581. doi:10.1093/gbe/evs009

111. Elzhov T V, Mullen KM, Spiess A-N, Bolker B. minpack.lm: R Interface to the Levenberg-Marquardt nonlinear least-squares algorithm found in MINPACK, plus support for bounds. 2016. Available: https://cran.r-project.org/package=minpack.lm

112. Fox J, Weisberg S. An {R} Companion to Applied Regression. Third. Thousand Oaks {CA}: Sage; 2019. Available: https://socialsciences.mcmaster.ca/jfox/Books/Companion/

113. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. Bioinformatics. 2013;29: 2933–2935. doi:10.1093/bioinformatics/btt509

114. Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, et al. Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. Nucleic Acids Res. 2018;46: D335–D342. doi:10.1093/nar/gkx1038

115. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol. 2016;34: 525–527. doi:10.1038/nbt.3519

116. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of RNA-seq incorporating quantification uncertainty. Nat Methods. 2017;14: 687–690. doi:10.1038/nmeth.4324

117. Guy L, Roat Kultima J, Andersson SGE. genoPlotR: comparative gene and genome visualization in R. Bioinformatics. 2010;26: 2334–2335. doi:10.1093/bioinformatics/btq413

118. Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize implements and enhances circular visualization in R . Bioinformatics. 2014;30: 2811–2812. doi:10.1093/bioinformatics/btu393

119. Wilke C. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2." 2020.

120. Kassambara A. ggpubr: "ggplot2" Based Publication Ready Plots. 2020. Available: https://cran.r-project.org/package=ggpubr

121. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

122. Sobetzko P, Jelonek L, Strickert M, Han W, Goesmann A, Waldminghaus T. DistAMo: A Web-Based Tool to Characterize DNA-Motif Distribution on Bacterial Chromosomes. Front Microbiol. 2016;7: 283. doi:10.3389/fmicb.2016.00283

123. Croucher NJ, Campo JJ, Le TQ, Liang X, Bentley SD, Hanage WP, et al. Diverse evolutionary patterns of pneumococcal antigens identified by pangenome-wide immunological screening. Proc Natl Acad Sci U S A. 2017;114: E357–E366. doi:10.1073/pnas.1613937114

124. Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in 2020. Nucleic Acids Res. 2021;49: D458–D460. doi:10.1093/nar/gkaa937

125. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780. doi:10.1093/molbev/mst010

126. Price MN, Dehal PS, Arkin AP. FastTree 2–approximately maximum-likelihood trees for large alignments. PLoS One. 2010;5: e9490.

127. Schoch CL, Ciufo S, Domrachev M, Hotton CL, Kannan S, Khovanskaya R, et al. NCBI Taxonomy: a comprehensive update on curation, resources and tools.

1   Database (Oxford). 2020;2020. doi:10.1093/database/baaa062

1 Figures

3 **Figure 1** Variation in transformability between locked *tvr* variants. (A) Violin plot showing the

4 transformation efficiency of four pairs of *tvr* locus variants constructed from isolates RMV5,

5 RMV6, RMV7 and RMV8. Each individual point represents an independent transformation

experiment. The horizontal line within each violin shows the median for each genotype. The

brackets indicate the statistical significance of the comparison between variants from the

same isolate background, as calculated using a two-tailed Wilcoxon rank sum test. (B)

Schematic of the *tvr* loci from RMV7$_{wt}$, RMV7$_{domi}$ and RMV7$_{rare}$ to show the genes encoding

the methylase (*hsdM*), endonuclease (*hsdR*), regulatory system (*tvrAT*) and recombinase

(*tvrR*). The variants differ in their active *hsdS* genes, upstream of *tvrATR*. The RMV7$_{domi}$

HsdS protein comprises the TRD combination III-iii (recognising motif TGAN$_7$TCC), whereas

that of RMV7$_{rare}$ contains the TRDs III-I (recognising motif TGAN$_7$TATC). The black arrows

indicate the binding sites of a forward primer, in *hsdM*, and reverse primers, in *hsdS*

fragments. (C) Line graph showing the ratio of RMV7$_{domi}$ to RMV7$_{rare}$ loci in eight-day

passages of RMV7$_{wt}$. Although conversion to RMV7$_{rare}$ was detected, the RMV7$_{domi}$ variant

was more common in each of the three replicates at every measured timepoint. (D) Violin

plot showing the higher transformation efficiency of RMV$_{rare}$ relative to RMV7$_{domi}$ or RMV7$_{wt}$.

Each individual point represents an independent experiment in which the number of

transformants, and number of overall colony-forming units (cfu), was calculated. The

horizontal line within each violin shows the median for each genotype. Both mutants were

compared with RMV7$_{wt}$; the horizontal bracket shows a significant difference, as calculated

from a two-tailed Wilcoxon rank sum test. (E) Violin plot showing the adhesion of variants to

an abiotic surface, as quantified by OD$_{550}$ after crystal violet staining of cells resisting

washing from a microplate well. The RMV7$_{rare}$ variant showed a significantly greater

adhesion after 16 hours incubation at 35 °C, as assessed by a two-tailed Wilcoxon rank sum

test. (F) Violin plots showing the transformation efficiency of knock-in mutants during a

passage experiment. The *tvr* loci of RMV7 *tvr*$_{domi}$::Janus and RMV7 *tvr*$_{rare}$::Janus (both

containing a *tvrR* gene interrupted by a Janus cassette) were each introduced into an RMV7

tvr::*cat* background (Fig. S4). This isogenic pair were separately passaged in liquid cultures

over six days in five independent replicates. The number of transformants observed from

three transformation assays, conducted each day for both variants, is shown by the

individual points' shapes and colours. The violin plots summarise the median and distribution

of these values. The brackets indicate the statistical significance of the comparison between

variants from the same day of the passage, as calculated using a two-tailed Wilcoxon rank

sum test. Across all panels, significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; p

$< 10^{-4}$, ****. All p values were subject to a Holm-Bonferroni correction within each panel.

Figure 2 Chromosomal distribution of genes exhibiting significant differences in transcription

between RNA-seq samples. The outer ring shows the annotation of RMV7$_{domi}$ (accession

code OV904788). Protein coding sequences are represented as black boxes, with the

vertical positioning within the ring indicating the strand of the genome on which they are

encoded. The next ring inwards shows significant pre-CSP differences in transcription: green

genes were more highly expressed in RMV$_{domi}$, and blue genes were more highly expressed

in RMV$_{rare}$. The next ring inwards show significant changes in gene expression 10 minutes

post-CSP in RMV7$_{domi}$: pink genes were upregulated, and purple genes were

downregulated. The third ring inwards shows significant changes in gene expression 10

minutes post-CSP in RMV7$_{rare}$: red genes were upregulated, and orange genes were

downregulated. The two inner rings repeat this representation for changes in gene

expression 20 minutes post-CSP.

Figure 3 The dependence of transformation efficiency on import of carbon sources. (A)

Violin plots showing the transformation efficiency of RMV7$_{wt}$ relative to a mutant in which

manLMN had been disrupted by a Janus cassette. Each genotype was transformed in

unsupplemented media, and in the presence of one of six carbon sources, as indicated by

the plot colour (see key on bottom right). Each point represents an independent experiment,

and the horizontal line within the violin plots show the median for each combination of

recipient cell genotype and carbon source. For each genotype, Wilcoxon rank-sum tests

were used to test for evidence of changes in transformation efficiency caused by each

carbon source. Significant differences are indicated by the black brackets at the top of the

panel. (B) Violin plots showing the transformation efficiency of RMV7$_{rare}$ relative to a mutant

in which *manLMN* had been disrupted by a Janus cassette. Data are displayed as in panel

A. (C) Violin plots showing the transformation efficiency of RMV7$_{rare}$ relative to a mutant in

which *tfoX* had been disrupted by a Janus cassette, and a third genotype in which *tfoX* had

been restored. Data are displayed as in panel A; the data for RMV7$_{rare}$ are replicated from

panel B. (D) Violin plots showing the transformation efficiency of RMV7$_{rare}$ relative to a

mutant in which *yjbK* had been disrupted by a Janus cassette, and a third genotype in which

yjbK had been restored. Data are displayed as in panel A; the data for RMV7$_{rare}$ are

replicated from panel B. (E) Violin plots showing the concentration of 3',5'-cAMP in samples

taken from exponential and stationary phase cultures of *E. coli* DH5α, and *S. pneumoniae*

RMV7$_{wt}$ and RMV7$_{rare}$ genotypes differing in whether *yjbK* was intact or not. The genotypes

are indicated by the colours (see key on bottom left). (F) Violin plots showing the

transformation efficiency of RMV7$_{wt}$ relative to mutants in which either *tfoX* or *yjbK* had been

disrupted by a Janus cassette. Data are displayed as in panel A. Across all panels,

significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p values

were subject to a Holm-Bonferroni correction within each panel.

Figure 4. Effect of removing PRCI*dnaN* on RMV7wt. (A) Violin plots showing the number of transformants observed in assays of RMV7wt mutants in which different parts of PRCI*dnaN* were replaced with a Janus cassette. The genotypes are arranged left to right, and coloured black to purple, to represent the increasing proportion of the element that was replaced by the cassette. Mutants removed different combinations of the *dinD*-like gene (IONPJBJN_00496); the regulatory genes (*rep*); the replication genes (*rep*); the integration genes (*int*), and the *att* site. The structure of RMV7 PRCI*dnaN* meant that replacing the *int-rep* region also deleted the intervening *reg* genes. Each point is an independent transformation assay. The violin plot summarises the result for each mutant, with a horizontal line indicating the median. Each mutant was compared against the parental RMV7wt genotype using a Wilcoxon rank sum test. Black brackets at the top of the plot indicate significant differences in the number of observed transformants. (B) Violin plot quantifying the effect of PRCI*dnaN* on transformation efficiency. This comparison of RMV7wt with a mutant in which the PRCI and

its *att* site were removed was independent of the experiments presented in panel A, and

more accurately quantified transformation efficiency. The significant difference, calculated

from a two-tailed Wilcoxon rank sum test, is indicated by the bracket at the top of the panel.

(C) Violin plots showing gene expression, as quantified by qRT-PCR, in the genotypes

assayed in panel B. IONPJBJN_00507 is a coding sequence within the PRCI that is absent

from RMV7$_{wt}$ PRCI$_{dnaN+att}$::Janus. The six points for each gene correspond to three technical

replicate assays on each of two biological replicates. The horizontal line on the violin plot

shows the median relative abundance for each gene in each genotype. Across all panels,

significance is coded as: $p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p values

were subject to a Holm-Bonferroni correction within each panel.

Figure 5 The regulation of transformation by HrcA in RMV7. (A) Violin plots showing gene

expression, as quantified by qRT-PCR, in RMV7$_{rare}$ and a mutant derivative in which the

hrcA chaperone regulator gene was disrupted by a Janus cassette. The six points for each

gene correspond to three technical replicate assays of each of two biological replicates. The

horizontal line on the violin plot shows the median relative abundance for each gene in each

genotype. The absence of any change in *ciaR* expression demonstrated the effect on

transformation was independent of CiaRH. (B) Scatterplot showing the effect of CaCl$_2$ on

transformation efficiency in RMV7$_{rare}$ and mutants in which *hrcA* had been disrupted, and

then restored. Each point represents an independent transformation assay of one genotype

at the indicated CaCl$_2$ concentration. The best-fitting dose response logistic models are

shown, with the shaded areas corresponding to the 95% confidence intervals. (C) Violin

plots showing the transformation efficiency of RMV7$_{wt}$, RMV7$_{rare}$ and RMV7$_{rare}$ *hrcA*::Janus

during normal growth (35 °C) or following a 40 °C heat shock. Each point corresponds to an

independent transformation experiment, and the violin plots have a horizontal line indicating

the median transformation efficiency of each mutant at each temperature. (D) Combined

effects of chaperone and carbon source regulation on transformation efficiency. Results are

displayed as in panel C. Wilcoxon rank-sum tests were conducted between all pairs of

genotypes. These found both the single mutants, lacking *manLMN* or *hrcA*, were

significantly less transformable than the parental genotype. Furthermore, the double mutant

was less transformable than either single mutant. Across all panels, significance is coded as:

$p < 0.05$, *; $p < 0.01$, **; $p < 10^{-3}$, ***; $p < 10^{-4}$, ****. All p values were subject to a Holm-

Bonferroni correction within each panel.

1
Figure 6 Comparison of the regulation of competence in *S. pneumoniae* (left), from this

work, and *V. cholerae* (right), summarized from [72]. In each, competence is regulated by a

quorum-sensing system: CSP in *S. pneumoniae*, and the cholera autoinducer 1 (CAI-1) in *V.*

cholerae. CAI-1 operates through inhibiting LuxO, thereby activating the HapR protein,

which indicates a high-cell density environment. HapR activates competence through the

Quorum-Sensing and TfoX-dependent Regulator (QstR). This regulator also senses the

activation of TfoX in response to GlcNAc being sensed by the transmembrane regulator

TfoS, via the TfoR small RNA. TfoX activity is also regulated by the Catabolite Regulatory

Protein (CRP), which is activated by 3',5'-cAMP, generated by the CyaA adenylate cyclase

under carbon source starvation conditions. Hence there are parallels with the TfoX

orthologue, and adenylate cyclase-like protein YjbK, responding to GlcNAc in *S.*

pneumoniae RMV7. GlcNAc also appears to promote competence through a TfoX/YjbK-

independent route, based on the behaviour of RMV7$_{wt}$. There are no known parallels in *V.*

cholerae for the regulation of competence by HrcA, the binding of which to CIRCE

sequences is promoted by Ca^{2+}, thereby activating competence through an unknown

mechanism. Note in *S. pneumoniae*, these signals are shown to converge on the

competence-specific sigma factor ComX for simplicity, but the mechanisms and targets of

the signalling pathways are unknown.