1   *Mycobacterium tuberculosis* PhoP integrates stress response to intracellular survival by regulating

2   cAMP level

4   Hina Khan[1], Partha Paul[1,2], Harsh Goar [1,2,3], Bhanwar Bamniya[1,2,4], Navin Baid[1,5],

5   Dibyendu Sarkar[1,4,] *

7   [1]CSIR-Institute of Microbial Technology, Sector 39 A, Chandigarh 160036, India

8   [2]These authors contributed equally to this work.

10   [3]Present Address: Department of Medicine, Division of Hematology-Oncology

11   UT Southwestern Medical Center, Dallas, TX 75235

13   [4]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India

15   [5]Present address: Department of Biosciences and Bioengineering

16   Indian Institute of Technology Bombay

17   Powai, Mumbai-400076, India

18   Running title: Regulation of mycobacterial cAMP

Key Words: cAMP level; Mycobacterium; PhoP; stress response; virulence regulation

*Address correspondence to: Dibyendu Sarkar, CSIR-Institute of Microbial Technology,

Tel.: 091-172-6665291; Fax: 091-172-2690585; E-mail: dibyendu@imtech.res.in

Abstract

Survival of *M. tuberculosis* within the host macrophages requires the bacterial virulence regulator

PhoP, but the underlying reason remains unknown. cAMP is one of the most widely used second

messengers, which impacts on a wide range of cellular responses in microbial pathogens including *M.*

tuberculosis. Herein, we hypothesized that intra-bacterial cAMP level could be controlled by PhoP

since this major regulator plays a key role in bacterial responses against numerous stress conditions. A

transcriptomic analysis reveals that PhoP functions as a repressor of cAMP-specific

phosphodiesterase (PDE) Rv0805, which hydrolyses cAMP. In keeping with these results, we find

specific recruitment of the regulator within the promoter region of *rv0805* PDE, and absence of *phoP*

or ectopic expression of *rv0805* independently accounts for elevated PDE synthesis leading to

depletion of intra-bacterial cAMP level. Thus, genetic manipulation to inactivate PhoP-*rv0805*-cAMP

pathway decreases cAMP level, stress tolerance and intracellular survival of the bacillus.

Introduction

M. tuberculosis, the causative agent of pulmonary tuberculosis, encounters diverse environmental conditions during infection, persistence and transmission of the disease (Ehrt *et al*, 2018; Ernst, 2012; Russell, 2011). However, the pathogen is exceptionally capable to adjust and survive within diverse host environments. Mycobacterial adaptive response to different stages of infection is achieved via fine tuning of regulation of gene expression using an extensive repertoire of more than 100 transcriptional regulators, 11 two-component systems, 6-serine-threonine protein kinases and 13 alternative sigma factors suggesting that a very complex transcriptional program is important for *M. tuberculosis* pathogenesis. While regulation of gene expression as a consequence of interaction of tubercle bacilli with its immediate environment remains critically important (Rohde *et al*, 2007), defining these regulatory pathways represent a major challenge in the field.

3', 5'- cyclic adenosine monophosphate (cAMP), one of the most widely used second messengers, impacts on a wide range of cellular responses in microbial pathogens including *M. tuberculosis* (McDonough & Rodriguez, 2011). The bacterial genome encodes at least 15 adenylate cyclases, including one of the adenylate cyclases (Rv0386) which is required for virulence (Agarwal *et al*, 2009), and multiple cAMP regulated effector proteins(Banerjee *et al*, 2015; Johnson & McDonough, 2018; McDonough & Rodriguez, 2011; Shenoy & Visweswariah, 2006). cAMP levels are elevated upon infection of macrophages by pathogenic mycobacterium (Bai *et al*, 2009) and addition of exogenous cAMP was shown to influence mycobacterial protein expression (Gazdik & McDonough, 2005). While intra-bacterial cAMP responds to macrophage environment and carries out specific functions like gene expression and protein function under host-related conditions (Bai *et al*, 2011; Bai *et al.*, 2009; Johnson *et al*, 2017; Johnson & McDonough, 2018; Knapp & McDonough, 2014), secreted cAMP controls macrophage signalling (Agarwal *et al.*, 2009; Agarwal *et al*, 2006; Gazdik *et al*, 2009; Johnson *et al.*, 2017; Nambi *et al*, 2013; Ranganathan *et al*, 2016) (Rittershaus *et al*, 2018). cAMP signalling remains essential to *M. tuberculosis* pathogenesis. Agarwal and

colleagues had shown that a burst in synthesis of cAMP upon infection of macrophages, improved

bacterial survival by interfering with host signalling pathways (Agarwal *et al.*, 2009)**.** In keeping with

this, anti-tubercular compounds that interfere with mycobacterial cAMP levels also impact

intracellular growth of mycobacteria (Bai *et al.*, 2011; Johnson *et al.*, 2017; Knapp *et al*, 2015; Nambi

et al., 2013; Rittershaus *et al.*, 2018; VanderVen *et al*, 2015; Wilburn *et al*, 2022). Together, these

results underscore the significance of cAMP signalling in mycobacteria.

cAMP signalling is controlled at the transcriptional level. Previous *in silico* studies had

identified two out of 10 predicted nucleotide binding proteins of *M. tuberculosis* as members of the

CRP/FNR superfamily of transcriptional regulators (McCue *et al*, 2000). These are CRP (cyclic AMP

receptor protein), encoded by *rv3676* and CMR (cAMP macrophage regulator), encoded by *rv1675c*,

respectively. Of these two, CRP becomes activated upon cAMP binding, and functions as a global

regulator of ~100 genes (Agarwal *et al.*, 2006; Bai *et al*, 2005; Rickman *et al*, 2005). Consequently,

deletion of the *crp* locus significantly impairs mycobacterial growth and attenuates virulence of the

bacilli in a mouse model (Rickman *et al.*, 2005). In contrast, CMR is necessary for regulated

expression of genes involved in virulence and persistence including members of the dormancy

regulon (Gazdik *et al.*, 2009; Ranganathan *et al.*, 2016; Smith *et al*, 2017). These results strongly

suggest that cAMP homeostasis i.e. a balance between cAMP synthesis by adenylate cyclases and

cAMP degradation by phosphodiesterases contributes to rapid adaptive response of mycobacteria in a

hostile intracellular environment (Johnson & McDonough, 2018; McDonough & Rodriguez, 2011).

However, very little is known about the underlying mechanisms of regulation of mycobacterial cAMP

level.

M. tuberculosis encounters a hostile environment within the host. Among the hostile

conditions is the acidic pH stress and exposure to host immune effectors such as NO (Nathan &

Shiloh, 2000; Rustad *et al*, 2009; Wayne & Sohaskey, 2001). A growing body of evidence connects

virulence-associated mycobacterial *phoP* locus with varying environmental conditions, including acid

stress (Abramovitch *et al*, 2011; Bansal *et al*, 2017; Tan *et al*, 2013), heat-shock (Sevalkar *et al*, 2019;

Singh *et al*, 2014) and integration of acid stress response to redox homeostasis (Baker *et al*, 2019;

Baker *et al*, 2014; Goar *et al*, 2022). Disruption of *phoP,* the gene encoding the response regulator of

the PhoPR two-component signal transduction system (Gupta *et al*, 2006) significantly reduces *in vivo*

multiplication of the bacilli (Walters *et al*, 2006), suggesting that PhoPR remains essential for

virulence (Perez *et al*, 2001). Moreover, a mutant lacking this system shows a significantly lowered

synthesis of cell-wall components diacyltrehaloses, polyacyltrehaloses, and sulfolipids, specific to

pathogenic mycobacterial species (Gonzalo Asensio *et al*, 2006a; Goyal *et al*, 2011; Walters *et al.*,

2006). In fact, the significant attenuation of *phoPR* deletion strain forms the basis of the mutant being

considered in trials as a vaccine strain (Arbues *et al*, 2013). Recent transcriptomic analyses revealed

that approximately 2% of the H37Rv genome is regulated by PhoP (Solans *et al*, 2014; Walters *et al.*,

2006), and mycobacterial gene expression in response to acidic pH significantly overlaps with the

PhoP regulon (Rohde *et al.*, 2007). Consistent with this, a large subset of PhoPR-regulated low-pH-

inducible genes are induced immediately following *M. tuberculosis* phagocytosis and remain induced

during macrophage infection (Gonzalo Asensio *et al*, 2006b; Martin *et al*, 2006; Perez *et al.*, 2001;

Walters *et al.*, 2006). Additional evidences coupled with more recent results suggest that during onset

of macrophage infection, PhoPR activation is linked to acidic pH and the available carbon source,

suggesting a physiological link between pH, carbon source, and macrophage pathogenesis (Baker *et*

al., 2019).

112   In this study, we hypothesized that intra-mycobacterial cAMP level could be determined by

PhoP since the major regulator has been implicated in regulation of bacterial responses against

numerous stress conditions, many of which function as signals to activate cAMP synthesizing diverse

adenylate cyclases (Knapp & McDonough, 2014). Our results connect virulence regulator PhoP with

intra-mycobacterial cAMP level. We discovered that PhoP regulates expression of cAMP-specific

phosphodiesterase *rv0805*, which hydrolyses cAMP. To further probe the regulation, we demonstrate

that under the condition which activates PhoP-PhoR system, PhoP in a PhoR-dependent manner

represses transcription of *rv0805* through direct DNA binding at the upstream regulatory region.

These observations account for a consistently lower level of cAMP in a PhoPR-deleted *M.*

tuberculosis H37Rv (*phoPR*-KO) relative to the WT bacilli, and establishes the molecular mechanism

of regulation of cAMP level, absence of which strikingly impacts phagosome maturation, and reduces

mycobacterial survival within macrophages and mice. Together, the newly identified mechanism of

regulation of cAMP level allows intra-phagosomal survival and growth program of mycobacteria.

RESULTS

Intra-mycobacterial cAMP level is regulated by the *phoP* locus

We compared cAMP levels of WT and *phoPR*-KO mutant (lacking both the single copies of *phoP* and

phoR genes), grown under normal, NO stress and acid stress conditions (Fig. 1A). *phoPR*-KO showed

a significantly lower level of cAMP relative to the WT bacilli, both under normal and stress

conditions. Complementation of the mutant (Compl.) could restore cAMP to the WT level. A higher

cAMP level in the complemented strain under NO stress is possibly attributable to reproducibly

higher *phoP* expression in the complemented mutant under specific stress conditions (Khan et al.,

2022). Because bacterial growth often varies under stress conditions, and growth inhibition can

influence cAMP level, we compared viability of mycobacterial strains under normal and indicated

stress conditions (conditions of cAMP measurements) by determining the bacterial CFU (Figure 1 -

figure supplement 1). Note that for *in vitro* viability under specific stress conditions, indicated

mycobacterial strains were grown to mid log phase (OD_{600} 0.4-0.6) and exposed to acidic media (7H9

media, pH 4.5(Gouzy *et al*, 2021) for further two hours at 37˚C. Likewise, for NO stress, cells grown

to mid-log phase were exposed to 0.5 mM DetaNonoate for 40 minutes. Our results suggest that WT

and *phoPR*-KO under carefully controlled stress conditions display comparable viability, indicating

that variation in viable cell counts of the mutant under specific stress conditions do not account for

lower cAMP level. From these results, we conclude that PhoP plays a major role in maintaining intra-

mycobacterial cAMP level.

To investigate the possibility that *phoPR*-KO secretes out more cAMP and therefore,

shows a lower cytoplasmic level, we compared cAMP secretion of WT and the mutant (Fig. 1B). The

WT and mutant strains were grown as described in the Methods and culture filtrates (CF) as well as

cell lysates (CL) were collected as described previously (Anil Kumar *et al*, 2016). Our results

demonstrate that the mutant reproducibly secretes lower amount of cAMP relative to the WT bacilli,

and cAMP secretion is fully restored in the complemented mutant (Compl.). As the fold difference of

secretion (~ 1.25- fold) is not much different relative to the fold difference in intra-mycobacterial

cAMP level (~2- fold), we suggest that lower cAMP level of the mutant is not due to its higher

efficacy of cAMP secretion. Fig. 1C confirms absence of autolysis of mycobacterial cells as GroEL2,

a cytoplasmic protein, was undetectable in the culture filtrates (CF).

PhoP functions as a repressor of *rv0805*

We next investigated role of the *phoP* locus on expression of mycobacterial adenylate

cyclases (ACs) and phosphodiesterases (PDEs), which synthesize and degrade cAMP, respectively

(Fig. 2A). The selection of ACs and PDEs were based on two key points. First, we have chosen ACs

which are activated by known signals (Knapp & McDonough, 2014). Second, we reasoned that the

previously reported ACs were activated under environmental conditions, which are linked to

mycobacterial *phoP* locus (Bansal *et al.*, 2017; Goar *et al.*, 2022). Our RT-qPCR results using gene-

specific primer pairs (Supplementary file 1a) suggest that expression of adenylate cyclases including

rv0386, *rv1264*, *rv1647* and *rv2488c* (Agarwal *et al.*, 2009; Dass *et al*, 2008; Dittrich *et al*, 2006;

Knapp & McDonough, 2014) do not appear to be regulated by the *phoP* locus. However, a significant

activation of adenylate cyclase *rv0891c* (4±0.05-fold), and repression of phosphodiesterase *rv0805*

expression (6.5±0.7-fold), respectively, were dependent on the *phoP* locus. Although Rv0891c was

suggested as one of the *M. tuberculosis* H37Rv adenylate cyclases, the protein lacks most of the

important residues conserved for adenylate cyclase family of proteins (Zaveri *et al*, 2021). On the

other hand, *rv0805* encodes for a cAMP specific PDE, present only in slow growing pathogenic *M.*

tuberculosis (Matange *et al*, 2013; Shenoy *et al*, 2007; Shenoy *et al*, 2005). Although expression of

rv0805 was restored at the level of WT in the complemented mutant (Compl.), we observed poor

restoration of expression of *rv0891c* in the Compl. strain. Further, expression of PDE *rv1339* which

contributes to mycobacterial cAMP level (Thomson *et al*, 2022) remains unaffected by the *phoP*

locus. Therefore, we focussed our attention on the biological significance of PhoP-dependent

regulation of *rv0805*. It should be noted that expressions of *rv1357c* and *rv2837c*, encoding PDEs for

cyclic di-GMP (Flores-Valdez *et al*, 2015) and cyclic di-AMP (Valadares & Woo, 2017),

respectively, remained unchanged in *phoPR*-KO.

As we reproducibly observed activation of *rv0805* expression in *phoPR*-KO (relative to WT),

we investigated whether acidic pH conditions under which *phoPR* system is activated (Abramovitch

et al., 2011; Bansal *et al.*, 2017), impacts expression of *rv0805* in WT bacilli (Fig. 2B). Our results

show that repression of *rv0805* is significantly higher in WT grown under acidic conditions (pH 4.5)

relative to normal conditions (pH 7.0) of growth. This observation is consistent with RNA-seq data

displaying significant down-regulation of *rv0805* in WT bacilli grown under acidic pH conditions

relative to normal conditions of growth (GEO Accession number: GSE180161). As a control,

expression of PDE *rv1339* which also degrades cAMP, remains unaffected under acidic conditions of

growth. The finding that acidic pH (pH4.5) conditions of growth promoted PhoP-dependent

repression of *rv0805*, prompted us to investigate whether PhoP directly binds to *rv0805* promoter. To

this end, we next examined *in vivo* recruitment of PhoP within the *rv0805* promoter by chromatin

immunoprecipitation (ChIP) experiments (Fig. 2C). In this assay, formaldehyde-cross-linked DNA-

protein complexes of growing *M. tuberculosis* cells were sheared to generate fragments of average

size ≈500 bp. Next, ChIP experiments utilized anti-PhoP antibody, and IP DNA was analysed by real-

time qPCR relative to a mock sample (without antibody as a control) using FPrv0805up/RPrv0805up

as the primer pair (Supplementary file 1a). Our results show that under normal condition (light blue

bars) rv0805up showed an insignificant enrichment of PCR signal for PhoP relative to mock (no

antibody control) sample, suggesting low affinity DNA binding of PhoP under normal conditions.

However, IP samples from cells grown under acidic pH showed a significantly higher enrichment of

PhoP at the *rv0805* promoter (rv0805up; compare *light blue* bars with *dark blue* bars). As controls,

promoter of *msl3* (msl3up) which is controlled by PhoP, and non-specific 16S rRNAup showed

comparable enrichment, and no enrichment under normal and acidic conditions, respectively. Thus,

ChIP data showing PhoP recruitment under acidic pH conditions is in agreement with low pH-specific

impact of PhoP on *rv0805* expression (Fig. 2B). Note that PhoP binding to msl3up was used as a

positive control.

To examine DNA binding *in vitro*, we first probed for the PhoP binding site within the

upstream regulatory region of *rv0805* (rv0805up) by MEME Bioinformatic software using the

consensus PhoP binding sequence (He & Wang, 2014). Our results suggest that the likely PhoP

binding sequence spans from -127 to -110 (relative to the ORF start site) of rv0805up (Figure 2-figure

supplement 1A) (p=0.000726). Next, recombinant PhoP was phosphorylated by acetyl -phosphate

(AcP) and used in EMSA experiments as described earlier (Pathak *et al*, 2010). Consistent with the

presence of a PhoP binding site, EMSA results demonstrate that P~PhoP binds to radio-labelled

rv0805up to form a complex stable to gel electrophoresis (Figure 2-figure supplement 1B).

Probing PhoP-dependent regulation of *M. tuberculosis rv0805*

To examine whether the regulatory effect was attributable to PhoP activation via phosphorylation, we

next grew *phoPR*-KO complemented with either *phoP* (Fig. 3A) or the entire *phoPR* encoding ORFs

(Fig. 3B), both under normal (pH 7.0; empty bars) and acidic (pH 4.5; black bars) conditions and

compared relative expression of *rv0805*. Although both strains expressed *phoP*, the former strain

lacked a functional copy of *phoR*, the cognate sensor kinase which phosphorylates PhoP (Gupta *et al.*,

2006). *M. tuberculosis* H37Rv lacking a *phoR* gene (*phoPR*-KO*::phoP*) did not show a low pH-

dependent repression of *rv0805* expression. However, similar to WT bacilli, we observed a low pH-

dependent significant down-regulation of *rv0805* expression in *phoPR*-KO*::phoPR* (Compl.). Note

that a comparable expression of PDE *rv1339* was observed in both strains regardless of growth

conditions. These results indicate that acidic pH-dependent repression of *rv0805* expression *in vivo* is

attributable to P~PhoP requiring the presence of PhoR.

To examine the effect of Rv0805 on mycobacterial cAMP level, we next expressed a copy of

rv0805 in WT bacteria (referred to as WT-Rv0805) (Fig. 3C). *rv0805* ORF was cloned within the

multicloning site of pSTki (Parikh *et al*, 2013) between the EcoRI and HindIII sites under the control

of P$_{myc1}$*tetO* promoter, and expression of *rv0805* under non-inducing condition was verified by

determining the mRNA level (Figure 3 - figure supplement 1A). Although copy number of episomal

vectors with pAl5000 origin of replication (*oriM*) have been reported to be 3 by Southern

hybridization (Ranes *et al*, 1990), in this case wild-type and mutant Rv0805 proteins were expressed

from single-copy chromosomal integrants (Parikh *et al.*, 2013). We then assessed the impact of

Rv0805 on intra-mycobacterial cAMP level (Fig. 3C). Consistent with a previous study (Agarwal *et

al.*, 2009), WT-Rv0805 showed a significant depletion (2.1± 0.7-fold) of intra-mycobacterial cAMP

relative to WT bacteria. To confirm that the reduced level of mycobacterial cAMP is attributable to

rv0805 expression, we also expressed *rv0805M*, a mutant Rv0805 lacking phosphodiesterase activity.

As structural data coupled with biochemical evidences reveal that Asn-97 at the enzyme active site

plays a key role in phosphodiesterase activity of Rv0805 (Shenoy *et al.*, 2007; Shenoy *et al.*, 2005),

the mutant Rv0805M was constructed by changing the conserved Asn-97 to Ala. WT-Rv0805M

showed an insignificant variation of cAMP level relative to WT, suggesting that depletion of intra-

mycobacterial cAMP in WT-Rv0805 is indeed attributable to phosphodiesterase activity of Rv0805.

The corresponding mRNA levels of PDEs (wild-type and the mutant) are over-expressed

approximately 4.5-6 -fold relative to the genomic *rv0805* level of WT-H37Rv (Figure 3-figure

supplement 1A). In contrast, other PDE encoding genes (*rv1357* and *rv2387*), under identical

conditions, demonstrate comparable expression levels in WT-H37Rv and *rv0805* over-expressing

strains. Over-expression of these PDEs did not influence bacterial growth under normal conditions

(Figure 3 - figure supplement 1B).

To further probe regulation of Rv0805 expression and its control of intra-mycobacterial

cAMP level, we utilized a previously reported CRISPRi based approach (Singh *et al*, 2016) to

construct *rv0805* and *phoP* knockdown (*rv0805*-KD, and *phoP-KD*, respectively) mutants. Consistent

with *phoPR*-KO, *phoP*-KD shows a significantly higher *rv0805* expression in the presence of ATc

relative to its absence (Fig. 3D). However, despite a significant down regulation of *rv0805* expression

in presence of ATc, a comparable *phoP* expression was observed in *rv0805*-KD mutant both in the

absence or presence of ATc. As a control, we observed a comparable expression of 16S rRNA in both

knock-down mutants. Next, we determined intra-mycobacterial cAMP of the mutants as described in

Fig. 1 (Fig. 3E). cAMP level of *phoP*-KD (showing activation of Rv0805) was significantly lower

relative to WT bacteria. In contrast, *rv0805*-KD mutant demonstrated a significantly higher level of

cAMP relative to WT. We speculate that effective knocking down of *phoP* or *rv0805* is not truly

reflected in the extent of variation of cAMP levels possibly due to the presence of numerous other

mycobacterial PDEs. These data represent an integrated view of our results suggesting that PhoP-

dependant repression of *rv0805* regulates intra-mycobacterial cAMP level. In keeping with these

results, activated PhoP under acidic pH conditions significantly represses *rv0805*, and intracellular

mycobacteria most likely utilizes a higher level of cAMP to effectively mitigate stress for survival

under hostile environment including acidic pH of the phagosome.

PhoP contributes to mycobacterial stress tolerance by repressing the *rv0805* PDE expression

To investigate whether cAMP level influences mycobacterial susceptibility to stress, we

compared *in vitro* growth under acidic pH (pH 4.5) (Fig. 4A). As expected, *phoPR*-KO showed a

significant growth inhibition relative to WT under low pH (pH 4.5) (Bansal *et al.*, 2017). WT-

Rv0805, but not WT-Rv0805M displayed a comparable susceptibility to acidic pH as that of *phoPR*-

KO. However, all four mycobacterial strains showed comparable growth at pH 7.0. Next, to compare

growth of WT-Rv0805 and WT under oxidative stress, cells were grown in presence of increasing

concentrations of diamide, a thiol-specific oxidant, and examined by microplate-based Alamar Blue

assays (Figure 4 - figure supplement 1A). We have recently shown that *phoPR*-KO is significantly

more sensitive to diamide relative to WT (Goar *et al.*, 2022). Here, we uncovered that similar to

phoPR-KO, WT-Rv0805, but not WT-Rv0805M was significantly more susceptible to diamide stress

as compared to WT (Fig. 4B). A previous study had reported that *phoP*-deleted mutant strain was

more sensitive to Cumene Hydrogen Peroxide (CHP), suggesting a role of PhoP in regulating

mycobacterial stress response to oxidative stress (Walters *et al.*, 2006). To compare sensitivity to

CHP, we grew mycobacterial strains in presence of 50 µM CHP for 24 hours and determined their

survival by enumerating CFU values (Fig. 4C). In this case we were unable to perform Alamar Blue-

based survival assays requiring a longer time because of the bactericidal property of CHP. Our CFU

data highlight that WT-Rv0805, but not WT-Rv0805M, displayed a significantly higher growth

inhibition relative to WT in the presence of CHP. Together, these results reveal similar behaviour of

phoPR-KO, and WT-Rv0805 by demonstrating a comparably higher susceptibility of these strains to

acidic pH and oxidative stress relative to WT bacteria and indicate a link between intra-mycobacterial

cAMP level and bacterial stress response. It appears that at least one of the mechanisms by which

PhoP contributes to global stress response is attributable to maintenance of cAMP level.

A previous study showed that *rv0805* over-expression in *M. smegmatis* influences cell wall

permeability (Podobnik *et al*, 2009). Having shown a significantly higher sensitivity of WT-Rv0805

to low pH and oxidative stress (relative to WT), we sought to investigate whether altered cell wall

structure/properties of the mycobacterial strain contribute to elevated stress sensitivity. We compared

expression level of lipid biosynthetic genes, which encode part of cell-wall structure of the bacilli

(Gonzalo Asensio *et al.*, 2006a; Walters *et al.*, 2006). Our results suggest that in contrast to *phoPR*-

KO, both WT-Rv0805 and WT-Rv0805M share a comparable expression profile of complex lipid

biosynthesis genes as that of WT (Figure 4 - figure supplement 1B). These results suggest that both

strains expressing wild type or mutant PDEs share a largely similar cell-wall properties and are

consistent with (a) a recent study reporting no significant effect of cAMP dysregulation on

mycobacterial cell wall structure/permeability (Wong *et al*, 2023), and (b) role of PhoP in cell wall

composition and complex lipid biosynthesis (Gonzalo Asensio *et al.*, 2006a; Goyal *et al.*, 2011;

Walters *et al.*, 2006). These results support our view that higher susceptibility of WT-Rv0805 to stress

conditions, is attributable to its reduced cAMP level.

To investigate the impact of mycobacterial cAMP level *in vivo*, we studied infection of

murine macrophages using WT, WT-Rv0805, and WT-Rv0805M (Fig. 4D). In this assay, WT-H37Rv

inhibits phagosome maturation, whereas phagosomes with *phoPR*-KO mature into phagolysosomes

(Anil Kumar *et al.*, 2016). In our present experimental set up, although WT bacilli inhibited

phagosome maturation, infection of macrophages with WT-Rv0805 and *phoPR*-KO matured into

phagolysosomes, suggesting increased trafficking of the bacilli to lysosomes. Under identical

conditions, WT-Rv0805M could effectively inhibit phagosome maturation just as WT bacteria.

Results from co-localization experiments are plotted as Fig. 4E and as Pearson's correlation co-

efficient of the quantified co-localization signals as Fig. 4F. These data suggest reduced ability of

WT-Rv0805, but not WT-Rv0805M (relative to WT) to inhibit phagosome maturation. From these

results, we suggest that ectopic expression of *rv0805* impacts phagosome maturation arguing in

favour of a role of PhoP in influencing phagosome-lysosome fusion in macrophages.

Intra-bacterial cAMP level and its effect on *in vivo* survival of mycobacteria

To examine the effect of intra-bacterial cAMP level on *in vivo* survival, mice were infected

with mycobacterial strains via the aerosol route. Day 1 post-infection, CFU analyses revealed a

comparable count of four mycobacterial strains (~100 bacilli) in the mice lungs. However, for WT-

Rv0805, the CFU recovered from infected lungs 4 weeks post- infection declined by ~218-fold

relative to the lungs infected with WT bacteria (Fig. 5A). In contrast, the CFU recovered from

infected lungs after 4 weeks of infection by WT-Rv0805M marginally declined by ~7-fold relative to

the lungs infected with WT bacilli. These results suggest a significantly compromised ability of WT-

Rv0805 (relative to WT) to replicate in the mice lungs. Note that *phoPR*-KO, under the conditions

examined, showed a ~ 246-fold lower lung burden compared to WT. In keeping with these results,

while the WT bacilli disseminated to the spleens of infected mice, a significantly lower count of WT-

Rv0805 was recovered from the spleens after 4 weeks of infection (Fig. 5B). Thus, we suggest that

one of the reasons which accounts for an attenuated phenotype of *phoPR*-KO in both cellular and

animal models is attributable to PhoP-dependent repression of *rv0805* PDE activity, which controls

mycobacterial cAMP level.

M. tuberculosis H37Rv persists within granulomas where it is protected from the anti-

mycobacterial immune effectors of the host. Histopathological evaluations showed that WT bacilli -

infected lung sections displayed aggregation of granulocytes within alveolar spaces which degenerate

progressively to necrotic cellular debris. In contrast, *phoPR*-KO and WT-Rv0805 showed less severe

pathology as indicated by decreased tissue consolidation, smaller granulomas, and open alveolar

space (Fig. 5C). Together, these results suggest that ectopic expression of *rv0805* in WT bacilli is

phenotypically equivalent to deletion of *phoP*, suggesting that failure to maintain cAMP level most

likely accounts for attenuated phenotype of the bacilli and absence of immunopathology in the lungs

of infected mice.

Discussion

A number of studies suggest that conditions associated with host environment like low pH, and

macrophage interactions often influence mycobacterial cAMP levels (Bai *et al.*, 2009; Gazdik &

McDonough, 2005). Although many bacterial pathogens modulate host cell cAMP levels as a

common strategy, the mechanism of regulation of mycobacterial cAMP-level remains unknown. In

this study, we sought to investigate whether PhoP, a master regulator implicated in controlling diverse

mycobacterial stress response, regulates mycobacterial cAMP level. We find that under normal

conditions as well as under carefully controlled single stress conditions, *phoPR*-KO shows a

significantly lower level of cAMP relative to the WT bacilli (Fig. 1A), and complementation of the

mutant restored cAMP level. To investigate the mechanism, we next probed regulation of ACs and

PDEs (Fig. 2) and demonstrated that PhoP functions as a major repressor of *rv0805*, encoding cAMP

specific PDE. Indeed, this newly-identified *rv0805* regulation, coupled with a recent discovery that

phosphodiesterase activity of Rv0805 controls propionate detoxification (McDowell *et al*, 2023), fits

well with and explains, the previously puzzling *in vivo* observation by Abramovitch et al

(Abramovitch *et al.*, 2011) that PhoP -controlled *aprABC* locus is associated with the regulation of

genes of carbon and propionate metabolism.

Although a large number of ACs are present in *M. tuberculosis* genome, a class III metallo-

phosphoesterase Rv0805 was earlier considered the only PDE, specific for mycobacterial cAMP.

However, a recent study has identified an atypical class II PDE Rv1339, which upon overexpression

reduces cAMP level and contributes to antibiotic sensitivity (Thomson *et al.*, 2022). While, the

functional role of Rv1339 in *M. tuberculosis* is yet to be understood, crystal structure and biochemical

evidence suggest that dimeric Rv0805 is stabilized by the presence of a divalent cation, and remains

catalytically active on a broad range of linear and cyclic PDE substrates *in vitro* (Keppetipola &

Shuman, 2008; Shenoy *et al.*, 2007; Shenoy *et al.*, 2005). More recently, cyclic nucleotide hydrolytic

activity of mycobacterial Rv0805 has been implicated in propionate detoxification (McDowell *et al.*,

2023). However, the mechanism of regulation of Rv0805 and its effect on mycobacterial cAMP level

remained unknown before the present study.

To examine biological significance of PhoP-dependent Rv0805 expression, we studied

rv0805 expression under acidic conditions of growth as *phoPR* system is induced under acidic pH

both *in vitro* and in macrophages (Abramovitch *et al.*, 2011; Bansal *et al.*, 2017). A significantly

higher repression of *rv0805* expression under acidic pH relative to normal conditions is consistent

with activation of PhoP and subsequent repression of *rv0805* (Fig. 3). These results further suggest

that effective mitigation of stress by mycobacteria possibly requires a higher cAMP level for survival

under intra-phagosomal environment. In keeping with these results, we find that (a) P~PhoP binds to

rv0805 regulatory region (Fig. 2-figure supplement 1), and (b) PhoP-dependent *rv0805* expression

requires PhoR (Figs. 3A-B), the cognate kinase which activates PhoP in a signal-dependent manner

(Gupta *et al.*, 2006; Singh *et al*, 2023). These results account for a consistently lower level of cAMP

in *phoPR*-KO relative to the WT bacilli. Notably, except recently reported PDE Rv1339, Rv0805 has

been known as the only cAMP-specific PDE present in the slow growing pathogenic mycobacteria

and its closely related species (Matange, 2015; Shenoy *et al.*, 2007; Shenoy *et al.*, 2005), and Rv1339

expression does not appear to be regulated by the *phoP* locus (Fig. 2). Thus, the above results

showing PhoP-dependent repression of *rv0805* activity likely represent the most critical step of

regulation of mycobacterial cAMP level under stress. In this connection, our recent results that PhoP

interacts with cAMP receptor protein, CRP and a complex of two interacting regulators control

expression of virulence determinants (Khan *et al*, 2022), invite speculation of a complex regulatory

control of cAMP-responsive mycobacterial physiology.

399   As one might argue that PhoP deletion and *rv0805* over-expression could be unrelated and

independent events, we constructed *phoP* and *rv0805* knock-down mutants to further investigate the

PhoP-Rv0805-cAMP pathway. Our objective was to probe regulation of expression (Fig. 3D) and

examine the impact on mycobacterial cAMP (Fig. 3E). *phoP*-KD significantly elevated *rv0805*

expression; however, *phoP* expression remains unaffected in *rv0805*-KD (Fig. 3D). While elevated

rv0805 expression in *phoP*-KD reduces cAMP level, understandably cAMP level is elevated in

rv0805-KD mutant (Fig. 3E). These results integrate PhoP-dependent *rv0805* repression with

mycobacterial cAMP level, suggesting how *phoP*-deletion or knock-down, at least in part, mimics

Rv0805 over-expression. These considerations take on more significance given the fact that these two

events have similar consequences on relevant strains with respect to stress tolerance, and survival in

cellular and animal models. Thus, our results suggest that ectopic expression of *rv0805* is functionally

equivalent to deletion of the *phoP* locus. This observation is in apparent conflict with a previous work

by Matange and collaborators (Matange *et al.*, 2013), suggesting a cAMP-independent transcriptional

response in *rv0805* over-expressing *M. tuberculosis* H37Rv. Although both studies were performed

with *rv0805* over-expressing bacilli, the fact that important differences in the expression of PDEs in

this study (Matange *et al.*, 2013) and in our assays – yielding significantly different levels of *rv0805*

expression - most likely account for this discrepancy. While we cannot completely rule out the

possibility of cleavage of other cyclic nucleotide(s) by Rv0805 (Keppetipola & Shuman, 2008;

Shenoy *et al.*, 2007; Shenoy *et al.*, 2005) impacting our results, consistent with a previous study our

results correlate *rv0805* expression with intra-mycobacterial cAMP level (Agarwal *et al.*, 2009).

Further, our data on the effect of expression of cyclic nucleotide-specific PDE Rv0805 or its inactive

mutant (Rv0805M) correlate well with enzyme activities of the corresponding PDEs on mycobacterial

cAMP levels (Fig. 3C). Thus, we infer that PhoP-dependent regulation of *rv0805* is a critical regulator

of intra-mycobacterial cAMP level.

Our experiments to understand physiological significance of PhoP-dependent repression of

rv0805 expression uncovers a comparable stress tolerance of WT-Rv0805 and *phoPR*-KO

(significantly reduced relative to WT) (Fig. 4). These results are consistent with the notion that cAMP

level, at least in part, accounts for mycobacterial stress response. Along the line, WT-Rv0805

displayed a reduced ability to inhibit phagosome-lysosome fusion like *phoPR*-KO (Fig. 4). Further,

we show that WT-Rv0805, unlike the WT bacilli or WT-Rv0805M, shows a significantly reduced

intracellular growth in mice as that of *phoPR*-KO (Fig. 5). Thus, these results are of fundamental

significance to establish that PhoP contributes to maintenance of cAMP level and integrates it to

mechanisms of mycobacterial stress tolerance and intracellular survival. Together, we identify a novel

mycobacterial pathway as a therapeutic target and provide yet another example of an intimate link

between bacterial physiology and intracellular survival of the tubercle bacilli.

The results reported here are presented schematically in Fig. 6. In summary, upon sensing low

acidic pH as a signal PhoR activates PhoP, P~PhoP binds to *rv0805* upstream regulatory region and

functions as a specific repressor of Rv0805. Therefore, we observed (a) a reproducibly lower level of

cAMP in *phoPR*-KO relative to WT-H37Rv, (b) a significantly reduced expression of *rv0805* in WT-

H37Rv, grown under acidic pH relative to normal conditions, and (c) comparable cAMP levels in

phoPR-KO and WT-Rv0805. This is why the two strains remain ineffective to mount an appropriate

stress response, most likely due to their inability to coordinate regulation of gene expression because

of dysregulation of intra-mycobacterial cAMP level. However, without uncoupling regulatory control

of PhoPR and *rv0805* expression, we cannot confirm that dysregulation of cAMP level accounts for

virulence attenuation of *phoPR*-KO. Given the fact that *rv0805*-depleted *M. tuberculosis* is growth

attenuated *in vivo* (McDowell et al., 2023), paradoxically ectopic expression of *rv0805* leads to

dysregulated metabolic adaptation, thereby resulting in reduced stress tolerance and intracellular

survival.

Materials and Methods

Bacterial strains and culture conditions. *E. coli* DH5α was used for cloning experiments. *E. coli*

BL21(DE3), an *E. coli* B strain lysogenized with λDE3, a prophage expressing T7 RNA polymerase

from the IPTG (isopropyl-β-D-thiogalactopyranoside)-inducible *lac*UV5 promoter (Studier &

Moffatt, 1986) was used as the host for over-expression of recombinant proteins. *E. coli* strains were

grown in LB medium at 37°C with shaking, transformed according to standard procedures and the

transformants were selected on media containing appropriate antibiotics plates. WT- and mutant *M.*

tuberculosis and *M. smegmatis* mc^{2}155 were grown at 37°C in Middlebrook 7H9 broth (Difco)

containing 0.2% glycerol, 0.05% Tween-80 and 10% ADC (albumin-dextrose-catalase) or on 7H10-

agar medium (Difco) containing 0.5% glycerol and 10% OADC (oleic acid-albumin-dextrose-

catalase) enrichment. *phoPR* disruption mutant of *M. tuberculosis* H37Rv (*phoPR*-KO, a kind gift of

Dr. Issar Smith) was constructed as described (Walters *et al.*, 2006). To this end, a kanamycin-

resistant cassette from pUC-K4 was inserted into a unique EcoRV site within the coding region of

phoP gene, and disruption was confirmed by Southern blot analysis of chromosomal DNA isolated

from the mutant. Next, purified plasmid DNAs were electroporated into wild-type *M. tuberculosis*

strain by standard protocol (Jacobs *et al*, 1991). To complement *phoPR* expression, pSM607

containing a 3.6- kb DNA fragment of *M. tuberculosis phoPR* including 200-bp *phoP*

promoter region, a hygromycin resistance cassette, *attP* site and the gene encoding phage L5

integrase, as detailed earlier (Walters *et al.*, 2006) was used to transform *phoPR* mutant to

integrate at the L5 *attB* site. Growth, transformation of wild-type (WT), *phoPR*-KO, the

complemented mutant (Compl.) *M. tuberculosis* and selection of transformants on appropriate

antibiotics plates were performed as described (Goyal *et al.*, 2011). When appropriate, antibiotics

were used at the following concentrations: hygromycin (hyg), 250 µg/ml for *E. coli* or 50 µg/ml for

mycobacterial strains; streptomycin (str), 100 µg/ml for *E. coli* or 20 µg/ml for mycobacterial strains;

kanamycin (kan), 20 µg/ml for mycobacterial strains. For *in vitro* growth under specific stress

conditions, indicated mycobacterial strains were grown to mid log phase (OD_{600} 0.4-0.6) and exposed

to different stress conditions. For acid stress, cells were initially grown in 7H9 media, pH7.0 and on

attaining mid log phase it was transferred to acidic media (7H9 media, pH 4.5), and grown for further

two hours at 37˚C. For oxidative stress, cells were grown in presence of 50 µM CHP (Sigma) for 24

hours or indicated diamide concentration(s) for 7 days. For NO stress, cells grown to mid log phase

were exposed to 0.5 mM DataNonoate for 40 minutes (Voskuil *et al*, 2003).

cAMP measurement. Mycobacterial cell pellets were collected and washed with 1x PBS buffer, cells

were resuspended in IP buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA pH 8.0, 1 mM

PMSF, 5% glycerol and 1% TritonX 100) and cell lysates (CL) were prepared by lysing the cells in

presence of Lysing Matrix B (100 µm silica beads; MP Bio) using a FastPrep-24 bead beater (MP

Bio) at a speed setting of 6.0 for 30 seconds. The procedure was repeated for 10 cycles with

incubation on ice in between pulses. The supernatant was collected by centrifugation at 13,000 rpm

for 10 minutes and filtered through 0.22 µm filter (Millipore). cAMP levels in the cells were

determined in a plate reader by using fluorescence-based cAMP detection kit (Abcam) according to

the manufacturer's recommendations and normalized to the total protein present in the samples as

determined by a BCA protein estimation kit (Pierce). For secretion studies, each mycobacterial strain

was grown in Sauton's media as described (Anil Kumar *et al.*, 2016), comparable counts of bacterial

cells were pelleted, resuspended in 2 ml of Sauton's media in a 6-well plate format for 2 hours at

37°C, and the supernatants (culture filtrates, CF) were collected for cAMP measurements, as

described previously (Anil Kumar *et al.*, 2016).

Cloning. *M. tuberculosis* full-length ORFs of interest were cloned between EcoRI and HindIII sites of

the mycobacterial expression vector pSTKi (Parikh *et al.*, 2013) and expressed from the $P_{myc1}tetO$

promoter. Mutation in Rv0805 was introduced by two-stage overlap extension method using

mutagenic primers (Supplementary file 1b), and the construct was verified by DNA sequencing. For

over-expression of WT- or mutant PDEs, WT bacilli was transformed with pST-rv0805 or pST-

rv0805M to generate WT-Rv0805 or WT-Rv0805M, respectively.

EMSA. rv0805up DNA probe was PCR amplified, resolved on an agarose gel, recovered by gel

extraction, and end-labelled with [γ-^{32}P ATP] (1000 Ci nmol^{-1}) using T4 polynucleotide kinase. The

end-labelled DNA probe was purified from free label by Sephadex G-50 spin columns (GE

Healthcare), and incubated with increasing amounts of purified PhoP in a total volume of 10 µl

binding mix (50 mM Tris-HCl, pH 7.5, 50 mM NaCl, 0.2 mg/ml of bovine serum albumin, 10%

glycerol, 1 mM dithiothreitol, $\approx$ 50 ng of labelled DNA probe, and 0.2 µg of sheared herring sperm

DNA) at 20°C for 20 minutes. DNA-protein complexes were resolved by electrophoresis on a 6%

(w/v) polyacrylamide gel (non-denaturing) in 0.5X TBE (89 mM Tris-base, 89 mM boric acid and 2

mM EDTA) at 70 V and 4°C, and the position of the radioactive material was determined by exposure

to a phosphor storage screen.

Construction of *M. tuberculosis phoP* and *rv0805* knock-down mutants. In this study, we utilized

a previously reported CRISPRi system (Singh *et al.*, 2016) to construct knock-down mutants of *phoP*

and r*v0805* (*phoP*-KD, and *rv0805*-KD, respectively). This approach efficiently inhibits expression of

target genes via inducible expression of dCas9 along with gene specific guide RNAs (sgRNA). The

RNAs were 20 nt long and complementary to the non-template strand of the target gene. The sgRNAs

of *phoP* and *rv0805* were cloned in a vector pRH2521 using BbsI enzyme and the constructs were

confirmed by sequencing. The corresponding clones were used to transform *M. tuberculosis*

harbouring pRH2502, a vector expressing an inactive version of *Streptococcus pyogenes* cas9 (dcas9).

To express dcas9 and repress sgRNA-targeted genes (*phoP* or *rv0805*), the bacterial strains were

grown with or without 600 ng/ml of anhydro-tetracycline (ATc) every 48 hours, and cultures were

grown for 4 days. RNA isolation was carried out, and RT-qPCR experiments verified significant

repression of target genes. For the induced strains (in presence of ATc) expressing sgRNAs targeting

+155 to +175 (relative to *phoP* translational start site) and +224 to +244 sequences (relative to *rv0805*

translational start site), we obtained approximately 85% and 90% reduction of *phoP* and *rv0805* RNA

abundance, respectively, relative to corresponding un-induced strains. The oligonucleotides used to

generate gene-specific sgRNA constructs and the plasmids utilized in knock-down experiments are

listed in Supplementary file 1b.

RNA isolation. Total RNA was extracted from exponentially growing bacterial cultures grown with

or without specific stress as described above. Briefly, 25 ml of bacterial culture was grown to mid-log

phase (OD_{600}= 0.4 to 0.6) and combined with 40 ml of 5 M guanidinium thiocyanate solution

containing 1% β-mercaptoethanol and 0.5% Tween 80. Cells were pelleted by centrifugation, and

lysed by re-suspending in 1 ml Trizol (Ambion) in the presence of Lysing Matrix B (100 µm silica

beads; MP Bio) using a FastPrep-24 bead beater (MP Bio) at a speed setting of 6.0 for 30 seconds.

The procedure was repeated for 2-3 cycles with incubation on ice in between pulses. Next, cell lysates

were centrifuged at 13000 rpm for 10 minutes; supernatant was collected and processed for RNA

isolation using Direct-Zol^{TM} RNA isolation kit (ZYMO). Following extraction, RNA was treated with

DNAse I (Promega) to degrade contaminating DNA, and integrity was assessed using a Nanodrop

(ND-1000, Spectrophotometer). RNA samples were further checked for intactness of 23S and 16S

rRNA using formaldehyde-agarose gel electrophoresis, and Qubit fluorometer (Invitrogen).

Quantitative Real-Time PCR. cDNA synthesis and PCR reactions were carried out using total RNA

extracted from each bacterial culture, and Superscript III platinum-SYBR green one-step qRT-PCR

kit (Invitrogen) with appropriate primer pairs (2 µM) using an ABI real-time PCR detection system.

Oligonucleotide primer sequences used in RT-qPCR experiments are listed in Supplementary file 1a.

Control reactions with platinum Taq DNA polymerase (Invitrogen) confirmed absence of genomic

DNA in all our RNA preparations, and endogenously expressed *M. tuberculosis rpoB* was used as an

internal control. Fold difference in gene expression was calculated using $\Delta\Delta C_T$ method (Schmittgen &

Livak, 2008). Average fold differences in mRNA levels were determined from at least two biological

repeats each with two technical repeats. Non-significant difference is not indicated.

ChIP assays. ChIP experiments in actively growing cultures of *M. tuberculosis* were carried out

essentially as described previously (Fol *et al*, 2006). Immunoprecipitation (IP) was performed using

anti-PhoP antibody and protein A/G agarose beads (Pierce). qPCR reactions included PAGE purified

primer pairs (Supplementary file 1a) spanning specific promoter regions using suitable dilutions of

immunoprecipitated (IP) DNA in a reaction buffer containing SYBR green mix, and one unit of

Platinum Taq DNA polymerase (Invitrogen). An IP experiment without adding antibody (mock) was

used as the negative control, and data was analysed relative to PCR signal from the mock sample.

PCR amplifications were carried out for 40 cycles using serially diluted DNA samples (mock, IP

treated and total input) in a real-time PCR detection system (Applied Biosystems). In all cases melting

curve analysis confirmed amplification of a single product.

Immunoblotting. Cell lysates or culture filtrates were resolved by 12% SDS-PAGE and visualized by

Western blot analysis using appropriate antibodies. Briefly, resolved samples were electroblotted onto

polyvinyl difluoride (PVDF) membranes (Millipore, USA) and were detected by anti-GroEL2

antibody (Sigma), anti CFP-10 antibody (Abcam) or affinity-purified anti-PhoP antibody elicited in

rabbit (Alpha Omega Sciences, India). Goat anti-rabbit secondary antibody conjugated to horseradish

peroxidase was used, and blots were developed with Luminata Forte Chemiluminescence reagent

(Millipore, USA). RNA polymerase was used as a loading control and was detected with monoclonal

antibody against β-subunit of RNA polymerase, RpoB (Abcam).

Alamar Blue assay. In this assay, reduction of Alamar Blue correlates with the change of a non-

fluorescent blue to a fluorescent pink appearance, which is directly linked to bacterial growth. *M.*

tuberculosis H37Rv was grown in 7H9 media (Difco) with 10% ADS (albumin, dextrose and NaCl)

to an OD_{600} of 0.4, and freshly diluted to OD_{600} of 0.02. Next, increasing concentrations of diamide

was added to the wells of a 96 -well plate containing 0.05 ml 7H9 media followed by addition of 0.05

ml of *M. tuberculosis* H37Rv culture (0.02 OD_{600}). The plate was incubated at 37°C for 7 days.

Finally, 0.02 ml of 0.02% Resazurin (sodium salt, MP Bio), prepared in sterile 7H9 media was added

to each of the wells and the change in colour was examined after incubation at 37°C for 16 hours. The

fluorescence excitation was at 530 nm and emission was recorded at 590 nm. Efficiency of inhibition

was calculated relative to control wells which did not include diamide, and Rifampicin was included

as a positive control to confirm validity of the assay.

Macrophage Infections. Virulence of indicated H37Rv strains were assessed in murine macrophages

according to the previously published protocol (Solans *et al.*, 2014). Briefly, RAW264.7 macrophages

were grown in DMEM media containing 10% Fetal bovine serum at 37°C under 5% CO_2, and seeded

onto #1 thickness, 18 mm diameter glass coverslips in a 12-well plate at a 40% confluency (0.5

million cells). Cells were independently infected with titrated cultures of WT, WT-Rv0805, WT-

Rv0805M and *phoPR*-KO strains at a multiplicity of infection (MOI) of 1:5 for 3 hours at 37°C in 5%

CO_2, followed by 1X PBS washes thrice. The macrophages were further incubated for 3 hours at

37°C. After infection, extracellular bacteria were removed by washing thrice with PBS. To visualize

trafficking of the tubercle bacilli, mycobacterial strains were stained with phenolic auramine solution

(which selectively binds to mycolic acids) for 15 minutes followed by washing with acid alcohol

solution and finally with 1X PBS. The cells were stained with 150 nM LysoTracker Red DND-99

(Invitrogen) for 30 minutes in a CO_2 incubator. Next, the cells were fixed with 4% paraformaldehyde

for 15 minutes, washed thrice with PBS, the coverslips were mounted in Slow Fade-Anti-Fade

(Invitrogen) and analysed using laser scanning confocal microscope (Nikon) equipped with Argon

(488 nm excitation line; 510 nm emission detection) and LD (561 nm excitation line; 594 nm

emission detection) laser lines. Digital images were processed with IMARIS imaging software

(version 9.20). Details of the experimental methods and the laser/detector settings were optimized

using macrophage cells infected with WT-H37Rv as described previously (Anil Kumar *et al.*, 2016).

A standard set of intensity threshold was made applicable for all images, and percent bacterial co-

localization was determined by analyses of at least 50 infected cells originating from 10 different

fields of each of the three independent biological repeats.

Mouse infections. All experiments pertaining to mice were in accordance with Institutional

regulations after review of protocols and approval by the Institutional Animal Ethics Committee

(IAEC/17/05, and IAEC/19/02). Mice were maintained and bred in the animal house facility of CSIR-

IMTECH. Animal infection studies and subsequent experiments were carried out in the Institutional

BSL-3 facility as per institutional biosafety guidelines. Briefly, the experiments were conducted with

8-10 weeks old C57BL/6 mice, infected intranasally and euthanized post-infection for evaluation of

bacterial load in lungs and spleens. Infections were given through the respiratory route using an

inhalation exposure system (Glass-col) with passaged *M. tuberculosis* H37Rv cultures of mid log

phase. The actual bacterial load delivered to the animals was enumerated from 3-5 aerogenically

challenged mice, 1day post aerosol challenge. The animals were found to achieve a bacillary

deposition of 100 to 200 CFU in the lungs for each strain. Four weeks post infection, the animals were

sacrificed by cervical dislocation, lungs and spleens were isolated aseptically from the euthanized

animals, homogenized in sterile 1X PBS and plated after serially diluting the lysates on 7H11 agar

plates, supplemented with 10% OADC and antibiotics (50 μg/ml carbenicillin, 30 μg/ml polymyxin B,

10 μg/ml vancomycin, 20 μg/ml Trimethoprim, 20 μg/ml cycloheximide, and 20 μg/ml Amphotericin

B) to enumerate CFU. For histopathology, left lung lobes were fixed in 10% buffered formalin,

embedded in paraffin and stained with haematoxylin and eosin for visualization under the microscope.

The level of pathology was scored by analysing perivascular cuffing, leukocyte infiltration,

multinucleated giant cell formation and epithelial cell injury.

Statistical analysis. Data are presented as arithmetic means of the results obtained from multiple

replicate experiments ± standard deviations. Statistical significance was determined by Student's

paired t-test using Microsoft Excel or Graph Pad Prism. Statistical significance was considered at P

values of 0.05.

Acknowledgements

We are grateful to G. Marcela Rodriguez and Issar Smith (PHRI, New Jersey Medical School -

UMDNJ) for Δ*phoP*-H37Rv, and the complemented *M. tuberculosis* H37Rv strains, Adrie Steyn

(University of Alabama) for pUAB300/pUAB400 plasmids, Ashwani Kumar for very helpful

discussions, and Sanjeev Khosla for critical reading of the manuscript. We thank members of the

Institutional animal facility (iCARE) for their help with approval of our projects from the Institutional

Animal Ethics Committee. This study received financial support from intramural grants of CSIR-

IMTECH (OLP-0170), CSIR (MLP-0049) and by a research grant (to D.S) from SERB

(EMR/2016/004904), Department of Science and Technology (DST). H.K., P.P., H.G., B.B. and N.B.

were supported by CSIR pre-doctoral fellowships.

Supplemental Data

The supplemental data include three (3) supplemental figures, and two supplemental tables

(Supplemental files 1a, and 1b, respectively). All data generated or analysed during this study are

included in the manuscript and supporting file; source data files have been provided for all relevant

figures.

Competing interests

The authors declare that no competing interests exist.

REFERENCES

Abramovitch RB, Rohde KH, Hsu FF, Russell DG (2011) aprABC: a Mycobacterium tuberculosis
complex-specific locus that modulates pH-driven adaptation to the macrophage phagosome.
Molecular microbiology 80: 678-694
Agarwal N, Lamichhane G, Gupta R, Nolan S, Bishai WR (2009) Cyclic AMP intoxication of
macrophages by a Mycobacterium tuberculosis adenylate cyclase. *Nature* 460: 98-102
Agarwal N, Raghunand TR, Bishai WR (2006) Regulation of the expression of whiB1 in
Mycobacterium tuberculosis: role of cAMP receptor protein. *Microbiology* 152: 2749-2756
Anil Kumar V, Goyal R, Bansal R, Singh N, Sevalkar RR, Kumar A, Sarkar D (2016) EspR-dependent
ESAT-6 Protein Secretion of Mycobacterium tuberculosis Requires the Presence of Virulence
Regulator PhoP. *The Journal of biological chemistry* 291: 19018-19030
Arbues A, Aguilo JI, Gonzalo-Asensio J, Marinova D, Uranga S, Puentes E, Fernandez C, Parra A,
Cardona PJ, Vilaplana C *et al* (2013) Construction, characterization and preclinical evaluation of
MTBVAC, the first live-attenuated M. tuberculosis-based vaccine to enter clinical trials. *Vaccine* 31:
4867-4873
Bai G, Knapp GS, McDonough KA (2011) Cyclic AMP signalling in mycobacteria: redirecting the
conversation with a common currency. *Cellular microbiology* 13: 349-358
Bai G, McCue LA, McDonough KA (2005) Characterization of Mycobacterium tuberculosis Rv3676
(CRPMt), a cyclic AMP receptor protein-like DNA binding protein. *Journal of bacteriology* 187: 7795-
7804
Bai G, Schaak DD, McDonough KA (2009) cAMP levels within Mycobacterium tuberculosis and
Mycobacterium bovis BCG increase upon infection of macrophages. *FEMS immunology and medical*
microbiology 55: 68-73
Baker JJ, Dechow SJ, Abramovitch RB (2019) Acid Fasting: Modulation of Mycobacterium
tuberculosis Metabolism at Acidic pH. *Trends in microbiology* 27: 942-953
Baker JJ, Johnson BK, Abramovitch RB (2014) Slow growth of Mycobacterium tuberculosis at acidic
pH is regulated by phoPR and host-associated carbon sources. *Molecular microbiology* 94: 56-69
Banerjee A, Adolph RS, Gopalakrishnapai J, Kleinboelting S, Emmerich C, Steegborn C, Visweswariah
SS (2015) A universal stress protein (USP) in mycobacteria binds cAMP. *J Biol Chem* 290: 12731-
12743
Bansal R, Anil Kumar V, Sevalkar RR, Singh PR, Sarkar D (2017) Mycobacterium tuberculosis
virulence-regulator PhoP interacts with alternative sigma factor SigE during acid-stress response.
Molecular microbiology 104: 400-411
Dass BK, Sharma R, Shenoy AR, Mattoo R, Visweswariah SS (2008) Cyclic AMP in mycobacteria:
characterization and functional role of the Rv1647 ortholog in Mycobacterium smegmatis. *Journal of*
bacteriology 190: 3824-3834
Dittrich D, Keller C, Ehlers S, Schultz JE, Sander P (2006) Characterization of a Mycobacterium
tuberculosis mutant deficient in pH-sensing adenylate cyclase Rv1264. *International journal of*
medical microbiology : IJMM 296: 563-566
Ehrt S, Schnappinger D, Rhee KY (2018) Metabolic principles of persistence and pathogenicity in
Mycobacterium tuberculosis. *Nature reviews Microbiology* 16: 496-507
Ernst JD (2012) The immunological life cycle of tuberculosis. *Nature reviews Immunology* 12: 581-
591
Flores-Valdez MA, Aceves-Sanchez Mde J, Pedroza-Roldan C, Vega-Dominguez PJ, Prado-Montes de
Oca E, Bravo-Madrigal J, Laval F, Daffe M, Koestler B, Waters CM (2015) The Cyclic Di-GMP
Phosphodiesterase Gene Rv1357c/BCG1419c Affects BCG Pellicle Production and In Vivo
Maintenance. *IUBMB life* 67: 129-138

Fol M, Chauhan A, Nair NK, Maloney E, Moomey M, Jagannath C, Madiraju MV, Rajagopalan M
(2006) Modulation of Mycobacterium tuberculosis proliferation by MtrA, an essential two-
component response regulator. *Molecular microbiology* 60: 643-657
Gazdik MA, Bai G, Wu Y, McDonough KA (2009) Rv1675c (cmr) regulates intramacrophage and cyclic
AMP-induced gene expression in Mycobacterium tuberculosis-complex mycobacteria. *Molecular*
microbiology 71: 434-448
Gazdik MA, McDonough KA (2005) Identification of cyclic AMP-regulated genes in Mycobacterium
tuberculosis complex bacteria under low-oxygen conditions. *Journal of bacteriology* 187: 2681-2692
Goar H, Paul P, Khan H, Sarkar D (2022) Molecular Connectivity between Extracytoplasmic Sigma
Factors and PhoP Accounts for Coupled Mycobacterial Stress Response. *J Bacteriol* 204: e0011022
Gonzalo Asensio J, Maia C, Ferrer NL, Barilone N, Laval F, Soto CY, Winter N, Daffe M, Gicquel B,
Martin C *et al* (2006a) The virulence-associated two-component PhoP-PhoR system controls the
biosynthesis of polyketide-derived lipids in Mycobacterium tuberculosis. *J Biol Chem* 281: 1313-1316
Gonzalo Asensio J, Maia C, Ferrer NL, Barilone N, Laval F, Soto CY, Winter N, Daffé M, Gicquel B,
Martín C *et al* (2006b) The virulence-associated two-component PhoP-PhoR system controls the
biosynthesis of polyketide-derived lipids in Mycobacterium tuberculosis. *J Biol Chem* 281: 1313-1316
Gouzy A, Healy C, Black KA, Rhee KY, Ehrt S (2021) Growth of Mycobacterium tuberculosis at acidic
pH depends on lipid assimilation and is accompanied by reduced GAPDH activity. *Proc Natl Acad Sci*
U S A 118
Goyal R, Das AK, Singh R, Singh PK, Korpole S, Sarkar D (2011) Phosphorylation of PhoP protein plays
direct regulatory role in lipid biosynthesis of Mycobacterium tuberculosis. *J Biol Chem* 286: 45197-
45208
Gupta S, Sinha A, Sarkar D (2006) Transcriptional autoregulation by Mycobacterium tuberculosis
PhoP involves recognition of novel direct repeat sequences in the regulatory region of the promoter.
FEBS letters 580: 5328-5338
He X, Wang S (2014) DNA consensus sequence motif for binding response regulator PhoP, a
virulence regulator of Mycobacterium tuberculosis. *Biochemistry* 53: 8008-8020
Jacobs WR, Jr., Kalpana GV, Cirillo JD, Pascopella L, Snapper SB, Udani RA, Jones W, Barletta RG,
Bloom BR (1991) Genetic systems for mycobacteria. *Methods in enzymology* 204: 537-555
Johnson RM, Bai G, DeMott CM, Banavali NK, Montague CR, Moon C, Shekhtman A, VanderVen B,
McDonough KA (2017) Chemical activation of adenylyl cyclase Rv1625c inhibits growth of
Mycobacterium tuberculosis on cholesterol and modulates intramacrophage signaling. *Mol*
Microbiol 105: 294-308
Johnson RM, McDonough KA (2018) Cyclic nucleotide signaling in Mycobacterium tuberculosis: an
expanding repertoire. *Pathog Dis* 76
Keppetipola N, Shuman S (2008) A phosphate-binding histidine of binuclear
metallophosphodiesterase enzymes is a determinant of 2',3'-cyclic nucleotide phosphodiesterase
activity. *J Biol Chem* 283: 30942-30949
Khan H, Paul P, Sevalkar RR, Kachhap S, Singh B, Sarkar D (2022) Convergence of two global
regulators to coordinate expression of essential virulence determinants of Mycobacterium
tuberculosis. *Elife* 11
Knapp GS, Lyubetskaya A, Peterson MW, Gomes AL, Ma Z, Galagan JE, McDonough KA (2015) Role of
intragenic binding of cAMP responsive protein (CRP) in regulation of the succinate dehydrogenase
genes Rv0249c-Rv0247c in TB complex mycobacteria. *Nucleic Acids Res* 43: 5377-5393
Knapp GS, McDonough KA (2014) Cyclic AMP Signaling in Mycobacteria. *Microbiol Spectr* 2
Martin C, Williams A, Hernandez-Pando R, Cardona PJ, Gormley E, Bordat Y, Soto CY, Clark SO, Hatch
GJ, Aguilar D *et al* (2006) The live Mycobacterium tuberculosis phoP mutant strain is more
attenuated than BCG and confers protective immunity against tuberculosis in mice and guinea pigs.
Vaccine 24: 3408-3419
Matange N (2015) Revisiting bacterial cyclic nucleotide phosphodiesterases: cyclic AMP hydrolysis
and beyond. *FEMS Microbiol Lett* 362

Matange N, Hunt DM, Buxton RS, Visweswariah SS (2013) Overexpression of the Rv0805
phosphodiesterase elicits a cAMP-independent transcriptional response. *Tuberculosis (Edinb)* 93:
492-500
McCue LA, McDonough KA, Lawrence CE (2000) Functional classification of cNMP-binding proteins
and nucleotide cyclases with implications for novel regulatory pathways in Mycobacterium
tuberculosis. *Genome research* 10: 204-219
McDonough KA, Rodriguez A (2011) The myriad roles of cyclic AMP in microbial pathogens: from
signal to sword. *Nature reviews Microbiology* 10: 27-38
McDowell JR, Bai G, Lasek-Nesselquist E, Eisele LE, Wu Y, Hurteau G, Johnson R, Bai Y, Chen Y, Chan J
et al (2023) Mycobacterial phosphodiesterase Rv0805 is a virulence determinant and its cyclic
nucleotide hydrolytic activity is required for propionate detoxification. *Mol Microbiol* 119: 401-422
Nambi S, Gupta K, Bhattacharyya M, Ramakrishnan P, Ravikumar V, Siddiqui N, Thomas AT,
Visweswariah SS (2013) Cyclic AMP-dependent protein lysine acylation in mycobacteria regulates
fatty acid and propionate metabolism. *J Biol Chem* 288: 14114-14124
Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between
mammalian hosts and microbial pathogens. *Proceedings of the National Academy of Sciences of the
United States of America* 97: 8841-8848
Parikh A, Kumar D, Chawla Y, Kurthkoti K, Khan S, Varshney U, Nandicoori VK (2013) Development of
a new generation of vectors for gene expression, gene replacement, and protein-protein interaction
studies in mycobacteria. *Applied and environmental microbiology* 79: 1718-1729
Pathak A, Goyal R, Sinha A, Sarkar D (2010) Domain structure of virulence-associated response
regulator PhoP of Mycobacterium tuberculosis: role of the linker region in regulator-promoter
interaction(s). *J Biol Chem* 285: 34309-34318
Perez E, Samper S, Bordas Y, Guilhot C, Gicquel B, Martin C (2001) An essential role for phoP in
Mycobacterium tuberculosis virulence. *Molecular microbiology* 41: 179-187
Podobnik M, Tyagi R, Matange N, Dermol U, Gupta AK, Mattoo R, Seshadri K, Visweswariah SS (2009)
A mycobacterial cyclic AMP phosphodiesterase that moonlights as a modifier of cell wall
permeability. *J Biol Chem* 284: 32846-32857
Ranes MG, Rauzier J, Lagranderie M, Gheorghiu M, Gicquel B (1990) Functional analysis of pAL5000,
a plasmid from Mycobacterium fortuitum: construction of a "mini" mycobacterium-Escherichia coli
shuttle vector. *J Bacteriol* 172: 2793-2797
Ranganathan S, Bai G, Lyubetskaya A, Knapp GS, Peterson MW, Gazdik M, AL CG, Galagan JE,
McDonough KA (2016) Characterization of a cAMP responsive transcription factor, Cmr (Rv1675c), in
TB complex mycobacteria reveals overlap with the DosR (DevR) dormancy regulon. *Nucleic Acids Res*
44: 134-151
Rickman L, Scott C, Hunt DM, Hutchinson T, Menendez MC, Whalan R, Hinds J, Colston MJ, Green J,
Buxton RS (2005) A member of the cAMP receptor protein family of transcription regulators in
Mycobacterium tuberculosis is required for virulence in mice and controls transcription of the rpfA
gene coding for a resuscitation promoting factor. *Mol Microbiol* 56: 1274-1286
Rittershaus ESC, Baek SH, Krieger IV, Nelson SJ, Cheng YS, Nambi S, Baker RE, Leszyk JD, Shaffer SA,
Sacchettini JC *et al* (2018) A Lysine Acetyltransferase Contributes to the Metabolic Adaptation to
Hypoxia in Mycobacterium tuberculosis. *Cell Chem Biol* 25: 1495-1505.e1493
Rohde KH, Abramovitch RB, Russell DG (2007) Mycobacterium tuberculosis invasion of
macrophages: linking bacterial gene expression to environmental cues. *Cell Host Microbe* 2: 352-364
Russell DG (2011) Mycobacterium tuberculosis and the intimate discourse of a chronic infection.
Immunol Rev 240: 252-268
Rustad TR, Sherrid AM, Minch KJ, Sherman DR (2009) Hypoxia: a window into Mycobacterium
tuberculosis latency. *Cell Microbiol* 11: 1151-1159
Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nature
protocols* 3: 1101-1108

Sevalkar RR, Arora D, Singh PR, Singh R, Nandicoori VK, Karthikeyan S, Sarkar D (2019) Functioning of
Mycobacterial Heat Shock Repressors Requires the Master Virulence Regulator PhoP. *Journal of*
bacteriology 201
Shenoy AR, Capuder M, Draskovic P, Lamba D, Visweswariah SS, Podobnik M (2007) Structural and
biochemical analysis of the Rv0805 cyclic nucleotide phosphodiesterase from Mycobacterium
tuberculosis. *Journal of molecular biology* 365: 211-225
Shenoy AR, Sreenath N, Podobnik M, Kovacevic M, Visweswariah SS (2005) The Rv0805 gene from
Mycobacterium tuberculosis encodes a 3',5'-cyclic nucleotide phosphodiesterase: biochemical and
mutational analysis. *Biochemistry* 44: 15695-15704
Shenoy AR, Visweswariah SS (2006) New messages from old messengers: cAMP and mycobacteria.
Trends in microbiology 14: 543-550
Singh AK, Carette X, Potluri LP, Sharp JD, Xu R, Prisic S, Husson RN (2016) Investigating essential gene
function in Mycobacterium tuberculosis using an efficient CRISPR interference system. *Nucleic Acids*
Res 44: e143
Singh PR, Goar H, Paul P, Mehta K, Bamniya B, Vijjamarri AK, Bansal R, Khan H, Karthikeyan S, Sarkar
D (2023) Dual functioning by the PhoR sensor is a key determinant to Mycobacterium tuberculosis
virulence. *PLoS Genet* 19: e1011070
Singh R, Anil Kumar V, Das AK, Bansal R, Sarkar D (2014) A transcriptional co-repressor regulatory
circuit controlling the heat-shock response of Mycobacterium tuberculosis. *Mol Microbiol* 94: 450-
465
Smith LJ, Bochkareva A, Rolfe MD, Hunt DM, Kahramanoglou C, Braun Y, Rodgers A, Blockley A,
Coade S, Lougheed KEA *et al* (2017) Cmr is a redox-responsive regulator of DosR that contributes to
M. tuberculosis virulence. *Nucleic Acids Res* 45: 6600-6612
Solans L, Gonzalo-Asensio J, Sala C, Benjak A, Uplekar S, Rougemont J, Guilhot C, Malaga W, Martin
C, Cole ST (2014) The PhoP-dependent ncRNA Mcr7 modulates the TAT secretion system in
Mycobacterium tuberculosis. *PLoS Pathog* 10: e1004183
Studier FW, Moffatt BA (1986) Use of bacteriophage T7 RNA polymerase to direct selective high-
level expression of cloned genes. *Journal of molecular biology* 189: 113-130
Tan S, Sukumar N, Abramovitch RB, Parish T, Russell DG (2013) Mycobacterium tuberculosis
responds to chloride and pH as synergistic cues to the immune status of its host cell. *PLoS Pathog* 9:
e1003282
Thomson M, Liu Y, Nunta K, Cheyne A, Fernandes N, Williams R, Garza-Garcia A, Larrouy-Maumus G
(2022) Expression of a novel mycobacterial phosphodiesterase successfully lowers cAMP levels
resulting in reduced tolerance to cell wall-targeting antimicrobials. *J Biol Chem* 298: 102151
Valadares NF, Woo J (2017) Mechanism of Rv2837c from Mycobacterium tuberculosis remains
controversial. *J Biol Chem* 292: 13480
VanderVen BC, Fahey RJ, Lee W, Liu Y, Abramovitch RB, Memmott C, Crowe AM, Eltis LD, Perola E,
Deininger DD *et al* (2015) Novel inhibitors of cholesterol degradation in Mycobacterium tuberculosis
reveal how the bacterium's metabolism is constrained by the intracellular environment. *PLoS Pathog*
11: e1004679
Voskuil MI, Schnappinger D, Visconti KC, Harrell MI, Dolganov GM, Sherman DR, Schoolnik GK (2003)
Inhibition of respiration by nitric oxide induces a Mycobacterium tuberculosis dormancy program.
The Journal of experimental medicine 198: 705-713
Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I (2006) The Mycobacterium
tuberculosis PhoPR two-component system regulates genes essential for virulence and complex lipid
biosynthesis. *Mol Microbiol* 60: 312-330
Wayne LG, Sohaskey CD (2001) Nonreplicating persistence of mycobacterium tuberculosis. *Annual*
review of microbiology 55: 139-163
Wilburn KM, Montague CR, Qin B, Woods AK, Love MS, McNamara CW, Schultz PG, Southard TL,
Huang L, Petrassi HM *et al* (2022) Pharmacological and genetic activation of cAMP synthesis disrupts
cholesterol utilization in Mycobacterium tuberculosis. *PLoS Pathog* 18: e1009862

Wong AI, Beites T, Planck KA, Fieweger RA, Eckartt KA, Li S, Poulton NC, VanderVen BC, Rhee KY,
Schnappinger D *et al* (2023) Cyclic AMP is a critical mediator of intrinsic drug resistance and fatty
acid metabolism in M. tuberculosis. *Elife* 12
Zaveri A, Bose A, Sharma S, Rajendran A, Biswas P, Shenoy AR, Visweswariah SS (2021)
Mycobacterial STAND adenylyl cyclases: The HTH domain binds DNA to form biocrystallized
nucleoids. *Biophys J* 120: 1231-1246

Figure Legends

Figure 1: PhoP contributes to maintenance of mycobacterial cAMP level. (A) Intra-mycobacterial

cAMP levels were determined by a fluorescence-based assay as described in the Methods, and

compared for indicated mycobacterial strains, grown under normal or specific stress conditions. For

acid stress, mycobacterial strains were initially grown to mid-log phase (OD_{600} 0.4 to 0.6), and then it

was transferred to acidic pH (7H9 media, pH 4.5) for further 2 hours of growth at 37°C. For NO

stress, cells grown to mid-log phase were exposed to 0.5 mM DataNonoate for 40 minutes. The data

represent average values from three biological repeats (*$P \leq 0.05$, **$P \leq 0.01$). (B) To compare secretion

of cAMP by WT and *phoPR*-KO, cAMP levels were also determined in the corresponding culture

filtrates (CF). (C) Immunoblotting analysis of 10 µg of cell lysates (CL) and 20 µg of culture filtrates

(CF) of indicated *M. tuberculosis* strains. α-GroEL2 was used as a control to verify cytolysis of cells,

CFP-10 detected as a secreted mycobacterial protein in the culture filtrates, and RpoB used as the

loading control.

Figure 2: PhoP regulates expression of phosphodiesterase (PDE) *rv0805*. (A) To investigate

regulation of cAMP level, mRNA levels of well-characterized adenylate cyclases, and PDEs were

compared in indicated mycobacterial strains by RT-qPCR as described in the Methods. The results

show average values from biological triplicates, each with two technical repeats. Note that the

difference in expression levels of *rv0805* between WT and *phoPR*-KO was significant (p<0.01),

whereas the fold difference in mRNA level between WT and the complemented mutant (Compl.)

remains nonsignificant (not indicated). (B) To determine the effect of acidic pH conditions of growth,

mycobacterial *rv0805* expression was compared in WT grown under normal (pH 7.0) and acidic pH

(pH 4.5). Average fold difference in mRNA levels from biological duplicates (each with a technical

repeat) were measured as described in the Methods (**P≤0.01). As controls, expression of *rv1339* and

16S rDNA were also measured. Non-significant difference is not indicated. (C) *In vivo* PhoP binding

to *rv0805* promoter (rv0805up) was compared in WT grown under normal and acidic conditions of

growth using anti-PhoP antibody followed by ChIP-qPCR. Fold enrichment data represent mean

values of two independent experiments with a statistically significant fold difference (**P-value

<0.01; *P-value <0.05). The upstream regulatory regions of 16S rRNA (16S rRNAup) and msl3

(msl3up) were used as negative and positive controls, respectively. The assay conditions, sample

analyses, and detection are described in the Methods section.

Figure 3: PhoP dependent repression of *rv0805* to maintain mycobacterial cAMP level requires

the presence of PhoR. (A-B) To determine the impact of PhoR (the cognate sensor kinase of PhoP),

expression of *rv0805* was compared in indicated *M. tuberculosis* H37Rv strains: (A) *phoPR*-

KO*::phoP* (*phoPR* mutant complemented with *phoP*) and (B) *phoPR*-KO*::phoPR* (*phoPR* mutant

complemented with *phoP-phoR*) under normal and acidic conditions of growth. As expected, *phoPR*-

KO*::phoPR* (Compl.) shows a significant repression of *rv0805* (but not *rv1339*) under acidic pH

(***P<0.001). However, *rv0805* expression remains comparable in *phoPR*-KO*::phoP* under normal

as well as acidic conditions of growth. As a control, *rv1339* fails to show a variable expression in

indicated mycobacterial strains. (C) To determine the effect of ectopic expression of *rv0805* on intra-

mycobacterial cAMP level, WT and mutant Rv0805 proteins (Rv0805M, defective for

phosphodiesterase activity) were expressed in *M. tuberculosis* H37Rv (to construct WT-Rv0805, and

WT-Rv0805M, respectively) as described in the Methods section. Similar to *phoPR*-KO, WT-Rv0805

(but not WT-Rv0805M) showed a considerably lower level of cAMP relative to WT bacteria.

Significance in variation of cAMP levels were determined by paired student's t-test (**P<0.01). (D-E)

Relative expression of *phoP* and PDE in *phoP*-KD and *rv0805*-KD (*phoP* and *rv0805* knock-down

constructs, respectively). In keeping with elevated expression of *rv0805* in *phoPR*-KO, *phoP*-KD

shows an elevated expression of *rv0805* relative to WT bacilli. In contrast, *phoP* expression level

913 remains unaffected in *rv0805*-KD mutant. Panel E measured corresponding intra-bacterial cAMP

914 levels in the respective knock-down mutants, as described in the legend to Fig. 1A.

916 **Figure 4: Regulation of cAMP level and its effect on mycobacterial stress tolerance and survival**

917 **in macrophages**. (A) To compare susceptibility to low pH conditions, indicated mycobacterial strains

918 were grown at pH 4.5, and similar to *phoPR*-KO (grey circles), WT-Rv0805 (red circles) shows a

919 significant growth defect relative to WT (empty circles). However, WT-Rv0805M (green circles)

920 grows comparably well as that of the WT (empty circles). In contrast, at pH 7.0 all four mycobacterial

921 strains (WT, empty triangles; *phoPR*-KO, grey triangles; WT-Rv0805, red triangles; WT-Rv0805M,

922 green triangles) displayed comparable growth. (B) Microplate-based assay using Alamar Blue was

923 utilized to examine mycobacterial sensitivity to increasing concentrations of diamide. In this assay,

924 reduction of Alamar Blue correlates with the change of a non-fluorescent blue to a fluorescent pink

925 appearance, which is directly proportional to bacterial growth. Survival of indicated mycobacterial

926 strains, under normal conditions and in presence of 5 mM diamide, were determined by plotting

927 fluorescence intensity. The data is normalized relative to WT grown in presence of 5 mM diamide.

928 (C) To compare susceptibility to stress conditions, these mycobacterial strains were next grown in the

929 presence of 50 µM Cumene Hydrogen Peroxide (CHP). In presence of CHP, WT-Rv0805 (red

930 column) but not WT-Rv0805M (green column), shows a significant growth defect [relative to WT

931 (empty column)] in striking similarity to *phoPR*-KO (grey column). Note that similar to *phoPR*-KO,

932 WT-Rv0805 shows a comparably higher sensitivity to CHP relative to WT bacilli. However, WT-

933 Rv0805M expressing a mutant Rv0805, shows a significantly lower sensitivity to CHP relative to

934 WT-Rv0805, as measured by the corresponding CFU values. The growth experiments were

935 performed in biological duplicates, each with two technical replicates (**$P \leq 0.01$; ***$P \leq 0.001$). (D)

936 Murine macrophages were infected with indicated *M. tuberculosis* H37Rv strains. The cellular

937 organelle was made visible by LysoTracker; mycobacterial strains were stained with phenolic

938 auramine solution, and the confocal images display merge of two fluorescence signals (Lyso Tracker:

939 red; H37Rv: green; scale bar: 10 µm). The insets in the merge panels indicate bacteria which either

have inhibited or facilitated trafficking into lysosomes. White *arrowheads* in the merge panels

indicate non-colocalization which remains higher in WT-H37Rv and WT-Rv0805M relative to

phoPR-KO or WT-Rv0805. (E) Bacterial co-localization of *M. tuberculosis* H37Rv strains. The

percentage of auramine labelled strains co-localized with Lysotracker was determined by counting at

least 100 infected cells in 10 different fields. The results show the average values with standard

deviation determined from three independent experiments (***P$\leq$ 0.001). (F) Pearson's correlation

coefficient of images of internalized auramine-labelled mycobacteria and Lysotracker red marker in

RAW 264.7 macrophages. Data are representative of mean $\pm$ S.D., derived from three independent

experiments (*P<0.05; ***P<0.001).

Figure 5: Dysregulation of mycobacterial cAMP level impacts mycobacterial survival *in vivo*.

(A-B) Survival of mycobacterial strains in mice (A) lung, and (B) spleen after animals were given an

aerosol infection with ~100 CFU / lung. Mycobacterial load represents mean CFU values with

standard deviations obtained from at least five animals per strains used (**P< 0.01; ***P<0.001;

****P<0.0001). (C) Histopathology of lung sections after 4 weeks of infection with indicated

bacterial strains. Sections were stained with hematoxylin and eosin, observed under a light

microscope, and images of the pathology sections collected at x40 magnification display granulomas

(*filled arrows*) and alveolar space (*empty arrows*) (scale bar, 200 µm).

Figure 6: Increased cAMP level and effective stress response versus decreased cAMP level and

reduced stress tolerance of mycobacteria. In this model, upon activation by an appropriate signal

via the cognate sensor PhoR, P~PhoP binds to *rv0805* regulatory region and functions as a specific

repressor by preventing access for mycobacterial RNA polymerase (RNAP) to bind to the promoter

and initiate transcription. In keeping with PhoP-dependent *rv0805* repression, our results demonstrate

a reproducibly lower level of cAMP in *phoPR*-KO relative to WT bacilli. Thus, *phoPR*-KO (or WT-

Rv0805) remains ineffective to mount an appropriate stress response most likely due to its inability to

coordinate regulated gene expression because of dysregulation of cAMP level, accounts for their

reduced stress tolerance. Together, these molecular events suggest that failure to maintain cAMP level

accounts for attenuated phenotype of the bacilli.

Legends to supplementary figures

Figure 1-figure supplement 1: Defined and indicated stress conditions do not influence *in vitro*
growth of mycobacterial strains. To determine the impact of *in vitro* stress on bacterial growth
under conditions when intra-mycobacterial cAMP levels were measured, WT bacilli, *phoPR*-KO, and
the complemented mutant (Compl.) were exposed to carefully controlled stress conditions as detailed
in the Methods, and CFUs enumerated. The results clearly suggest that bacterial viability under
indicated conditions of stress do not impact intra-mycobacterial cAMP level.

Figure 2-figure supplement 1: Probing *in vitro* DNA binding of PhoP to the *rv0805* regulatory
region. (A) Core binding site of PhoP within the promoter region spanning -150 to +1-bp upstream
regulatory region of *rv0805* (rv0805up; relative to ORF start site) was identified by the MEME Suit
bioinformatic software using previously-identified consensus PhoP binding sequence (**He and Wang,**
2014). Note that the predicted binding site spans from -110 to -127 relative to the ORF start site
(p=0.000726). (B) To verify DNA binding *in vitro*, rv0805up was amplified, end-labelled, and EMSA
was carried out using radio-labelled rv0805up for binding of increasing concentrations of PhoP (lanes
2-4) and PhoP, pre-incubated in phosphorylation mix with acetyl-phosphate (AcP) (lanes 5-7),
respectively. Lane 1 shows the free probe. The binding mixtures in lanes 2-4, and lanes 5-7 contained
indicated proteins at 1, 2 and 4 μM, respectively. The position of the radioactive material was
determined by exposure to a phosphor storage screen. *Open* and *filled* arrows indicate free probe and
slower-moving complexes with band shifts produced in presence of P~PhoP, respectively.
991

Figure 3-figure supplement 1: Ectopic expression of wild-type and mutant *rv0805* (Rv0805M) in
WT bacilli. (A) Real-time RT-qPCR was carried out to compare expression of indicated
phosphodiesterases (PDEs) in WT (empty bar), WT-Rv0805 (red bar) and WT- Rv0805M (green bar),
respectively, as described in the Methods. Average fold difference in mRNA levels from two
biological repeats (each with a technical repeat) were determined as described in the Methods
(**P≤0.01; ***P≤0.001). Nonsignificant difference is not indicated. (B) To compare growth of WT
and the variant mycobacterial strains, CFU values were determined under normal conditions of
growth as described in the methods.

1000

Figure 4-figure supplement 1: PhoP-dependent *rv0805* expression contributes to mycobacterial
survival under oxidative stress. (A) In this experiment, we compared metabolic activity of WT,
phoPR-KO, WT-Rv0805, and WT-Rv0805M, grown in presence of increasing concentrations of
diamide by using Alamar Blue assay. Because reduction of Alamar Blue correlates with the change of
a non-fluorescent blue to a fluorescent pink appearance, mycobacterial metabolic activity could be
assessed by monitoring fluorescence. Two following controls were included in the Alamar Blue
assays: Rifampicin control, reflecting mycobacterial growth inhibition to confirm validity of the
assay, and no bacteria control indicated at the left top corner of the plate. (B) Next, effect of over

expression of *rv0805* on mycobacterial cell -wall structure was assessed by analysing expression of
lipid biosynthetic genes of indicated mycobacterial strains by RT-qPCR. Average fold difference in
mRNA levels from two biological repeats (each with a technical repeat) were determined as described
in the Methods (***$P \leq 0.001$). Nonsignificant difference is not indicated.
1013
1014 **Legends to supplementary Tables**

1015 Supplementary file 1a

1016 Sequences of oligonucleotide primers used in RT-qPCR and ChIP-qPCR measurements
1017 reported in this study
1018
1019 Supplementary file 1b
1020
1021 Sequences of oligonucleotide primers used for amplification and cloning, and plasmids used in this
1022 study
1023
1024

1025 **Source data files for supplementary figures**

1026 Figure 1-figure supplement 1

1027 Figure 2-figure supplement 1

1028 Figure 3-figure supplement 1

1029 Figure 4-figure supplement 1

1030

1031 **Author contributions**

1032

1033 H.K., P.P., H. G., B.B. and D.S. designed research; H.K., P. P., H.G., and B.B. performed research;

1034 N.B. contributed analytical tools; H.K., P. P., H. G., B.B., N.B., and D.S. analysed data; and D.S.

1035 wrote the manuscript.

1036

1037

1038 **Supplemental Data**

1039 The supplemental data include four supplemental figures and two supplemental tables (supplementary

1040 files 1a and 1b, respectively). All other data are part of the paper and its supplemental files.

1041   **Figure 1**

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052   **Figure 2**

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063 **Figure 3**

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077   **Figure 4**

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089   **Figure 5**

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100 **Figure 6**

1101

1102

1103

1104