Mycobacterium tuberculosis PhoP integrates stress response to intracellular survival by maintenance

of cAMP homeostasis

Hina Khan[1], Partha Paul[1,2], Harsh Goar [1,2,3], Bhanwar Bamniya[1,2,4], Navin Baid[1,5],

Dibyendu Sarkar[1,4, *]

[1]CSIR-Institute of Microbial Technology, Sector 39 A, Chandigarh 160036, India

[2]These authors contributed equally to this work.

[3]Present Address: Department of Medicine, Division of Hematology-Oncology

UT Southwestern Medical Center, Dallas, TX 75235

[4]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India

[5]Present address: Department of Biosciences and Bioengineering

Indian Institute of Technology Bombay

Powai, Mumbai-400076, India

Running title: mycobacterial cAMP homeostasis

Key Words: cAMP homeostasis; Mycobacterium; PhoP; stress response; virulence regulation

*Address correspondence to: Dibyendu Sarkar, CSIR-Institute of Microbial Technology,

Tel.: 091-172-6665291; Fax: 091-172-2690585; E-mail: dibyendu@imtech.res.in

Abstract

Survival of *M. tuberculosis* within the host macrophages requires the virulence regulator PhoP, but

the underlying mechanism remains unknown. Because growing evidence connects PhoP with varying

stress response, we hypothesized that the level of 3′,5′ cAMP, one of the most widely used second

messengers, was regulated by the *phoP* locus, linking numerous stress response with cAMP

production. A transcriptomic analysis discovers that PhoP functions as a repressor of cAMP-specific

phosphodiesterase (PDE) Rv0805, which hydrolytically degrades cAMP. The most fundamental

insight is derived from the PhoP-dependent regulation of *rv0805* expression by specific recruitment of

the regulator within the promoter region of the PDE. Consistent with these results, absence of PhoP

or ectopic expression of *rv0805* independently accounts for elevated PDE synthesis and depletion of

intra-mycobacterial cAMP level. Thus, genetic manipulation to inactivate PhoP-*rv0805*-cAMP

pathway leads to disruption of cAMP homeostasis, decreased stress tolerance and reduced survival of

the bacilli.

Introduction

M. tuberculosis, the causative agent of pulmonary tuberculosis, keeps encountering diverse environmental conditions during infection, persistence and transmission of the disease (Ehrt *et al*, 2018; Ernst, 2012; Russell, 2011). However, the intracellular pathogen is exceptionally capable to adjust and survive within diverse intracellular environments. Mycobacterial adaptive response to different stages of infection is achieved via fine tuning of regulation of gene expression using an extensive repertoire of more than 100 transcriptional regulators, 11 two-component systems, 6-serine-threonine protein kinases and 13 alternative sigma factors suggesting that a very complex transcriptional program is important for *M. tuberculosis* pathogenesis. While regulation of gene expression as a consequence of interaction of tubercle bacilli with its immediate environment remains critically important (Rohde *et al*, 2007), defining these regulatory pathways represent a major challenge in the field.

3', 5'- cyclic adenosine monophosphate (cAMP), one of the most widely used second messengers, impacts on a wide range of cellular responses in microbial pathogens including *M. tuberculosis* (McDonough & Rodriguez, 2011). The bacterial genome encodes at least 15 adenylate cyclases, including one of the adenylate cyclases (Rv0386) which is required for virulence (Agarwal *et al*, 2009), and multiple cAMP regulated effector proteins(Banerjee *et al*, 2015; Johnson & McDonough, 2018; McDonough & Rodriguez, 2011; Shenoy & Visweswariah, 2006). cAMP levels are elevated upon infection of macrophages by pathogenic mycobacterium (Bai *et al*, 2009) and addition of exogenous cAMP was shown to influence mycobacterial protein expression (Gazdik & McDonough, 2005). While intra-bacterial cAMP respond to macrophage environment and carry out specific functions like gene expression and protein function under host-related conditions (Bai *et al*, 2011; Bai *et al.*, 2009; Johnson *et al*, 2017; Johnson & McDonough, 2018; Knapp & McDonough, 2014), secreted cAMP controls macrophage signalling (Agarwal *et al.*, 2009; Agarwal *et al*, 2006; Gazdik *et al*, 2009; Johnson *et al.*, 2017; Nambi *et al*, 2013; Ranganathan *et al*, 2016) (Rittershaus *et*

al, 2018). Importantly, cAMP signalling remains essential to *M. tuberculosis* pathogenesis. Agarwal and colleagues had shown that bacterial cAMP burst, upon infection of macrophages, improved bacterial survival by interfering with host signalling pathways (Agarwal *et al.*, 2009)**.** In keeping with this, anti-tubercular compounds that interfere with mycobacterial cAMP levels also impact intracellular growth of mycobacteria (Bai *et al.*, 2011; Johnson *et al.*, 2017; Knapp *et al*, 2015; Nambi *et al.*, 2013; Rittershaus *et al.*, 2018; VanderVen *et al*, 2015; Wilburn *et al*, 2022). Together, these results underscore the significance of cAMP signalling in mycobacteria.

cAMP signalling is controlled at the transcriptional level. Previous *in silico* studies had identified two out of 10 predicted nucleotide binding proteins of *M. tuberculosis* as members of the CRP/FNR superfamily of transcriptional regulators (McCue *et al*, 2000). These are CRP (cyclic AMP Receptor protein), encoded by *rv3676* and CMR (cAMP macrophage regulator), encoded by *rv1675c*, respectively. Off these two, CRP becomes activated upon cAMP binding, and functions as a global regulator of ~100 genes (Agarwal *et al.*, 2006; Bai *et al*, 2005; Rickman *et al*, 2005). Consequently, deletion of the *crp* locus significantly impairs mycobacterial growth and attenuates virulence of the bacilli in a mouse model (Rickman *et al.*, 2005). In contrast, CMR is necessary for regulated expression of genes involved in virulence and persistence including members of the dormancy regulon (Gazdik *et al.*, 2009; Ranganathan *et al.*, 2016; Smith *et al*, 2017). Together, these results strongly suggest that balanced synthesis and degradation of cAMP contributes to rapid adaptive response of mycobacteria in a hostile intracellular environment (Johnson & McDonough, 2018; McDonough & Rodriguez, 2011). However, very little is known about the underlying mechanism of mycobacterial cAMP homeostasis.

M. tuberculosis encounters a hostile environment within the host. Among the hostile conditions is the acidic pH stress and exposure to host immune effectors such as NO (Nathan & Shiloh, 2000; Rustad *et al*, 2009; Wayne & Sohaskey, 2001). A growing body of evidence connects virulence-associated mycobacterial *phoP* locus with varying environmental conditions, including acid stress (Abramovitch *et al*, 2011; Bansal *et al*, 2017; Tan *et al*, 2013), heat-shock (Sevalkar *et al*, 2019;

Singh *et al*, 2014) and integration of acid stress response to redox homeostasis (Baker *et al*, 2019;

Baker *et al*, 2014; Goar *et al*, 2022). Disruption of *phoP,* the gene encoding the response regulator of

the PhoPR two-protein regulatory system (Gupta *et al*, 2006) significantly reduces *in vivo*

multiplication of the bacilli (Walters *et al*, 2006), suggesting that PhoPR remains essential for

virulence (Perez *et al*, 2001). Moreover, the mutant lacking this system shows a significantly lowered

synthesis of cell-wall components diacyltrehaloses, polyacyltrehaloses, and sulfolipids, specific to

pathogenic mycobacterial species (Gonzalo Asensio *et al*, 2006a; Goyal *et al*, 2011; Walters *et al.*,

2006). In fact, the significant attenuation of *phoPR* deletion strain forms the basis of the mutant being

considered in trials as a vaccine strain (Arbues *et al*, 2013). Recent transcriptomic analyses uncovered

that approximately 2% of the H37Rv genome is regulated by PhoP (Solans *et al*, 2014; Walters *et al.*,

2006), and mycobacterial gene expression in response to acidic pH significantly overlaps with the

PhoP regulon (Rohde *et al.*, 2007). Consistent with this, a large subset of PhoPR-regulated low-pH-

inducible genes are induced immediately following *M. tuberculosis* phagocytosis and remain induced

during macrophage infection(Gonzalo Asensio *et al*, 2006b; Martin *et al*, 2006; Perez *et al.*, 2001;

Walters *et al.*, 2006). Additional evidences coupled with more recent results suggest that during onset

of macrophage infection, PhoPR activation is linked to acidic pH and the available carbon source,

suggesting a physiological link between pH, carbon source, and macrophage pathogenesis (Baker *et*

al., 2019).

In this study, we hypothesized that intra-mycobacterial cAMP level could be determined by

the *phoP* locus since the major regulator has been implicated in regulation of bacterial responses

against numerous stress conditions, many of which function as signals to activate cAMP synthesizing

diverse adenylate cyclases (Knapp & McDonough, 2014). Interestingly, our results connect virulence

regulator PhoP with intra-mycobacterial cAMP level. We discovered that PhoP regulates expression

of cAMP-specific phosphodiesterase *rv0805*, which hydrolytically degrades cAMP. To further probe

the regulation, we demonstrate that under the condition which activates PhoP-PhoR system, PhoP in a

PhoR-dependent mechanism represses transcription of *rv0805* through direct DNA binding at the

upstream regulatory region. These observations account for a consistently lower level of cAMP in a

PhoPR-deleted *M. tuberculosis* H37Rv (*phoPR*-KO) relative to the WT bacilli, and establishes the

molecular mechanism of cAMP homeostasis, absence of which strikingly impacts phagosome

maturation, and reduces mycobacterial survival within macrophages and mice. Together, the newly

identified mechanism of cAMP homeostasis represents a molecular checkpoint during intra-

phagosomal survival and growth program of mycobacteria.

RESULTS

Intra-mycobacterial cAMP level is regulated by the *phoP* locus

As stress conditions are often linked to variation of cAMP levels, we compared cAMP levels of WT and *phoPR*-KO mutant (mutant lacking both copies of *phoP* and *phoR*), grown under normal and specific stress conditions, like NO stress and acid stress (Fig. 1A). Interestingly, *phoPR*-KO showed a significantly lower level of cAMP relative to the WT bacilli, both under normal and stress conditions. More importantly, complementation of the mutant (Compl.) could restore cAMP to the WT level. Because bacterial growth often varies under stress conditions, and growth inhibition can influence cAMP level, we compared viability of mycobacterial strains under normal and indicated stress conditions (conditions of cAMP measurements) by determining the bacterial CFU (Figure 1 - figure supplement 1). Note that for *in vitro* viability under specific stress conditions, indicated mycobacterial strains were grown to mid log phase (OD_{600} 0.4-0.6) and exposed to acidic media (7H9 media, pH 4.5(Gouzy *et al*, 2021) for further two hours at 37˚C. Likewise, for NO stress, cells grown to mid-log phase were exposed to 0.5 mM DetaNonoate for 40 minutes. Our results suggest that WT and *phoPR*-KO under carefully controlled stress conditions display comparable viability, indicating that variation in viable cell counts of the mutant under specific stress conditions do not account for lower cAMP level. From these results, we conclude that PhoP plays a major role in maintaining intra-mycobacterial cAMP level.

To investigate the possibility that *phoPR*-KO secretes out more cAMP and therefore, shows a lower cytoplasmic level, we compared cAMP secretion of WT and the mutant (Fig. 1B). The WT and mutant strains were grown as described in the Methods and culture filtrates (CF) as well as cell lysates (CL) were collected as described previously (Anil Kumar *et al*, 2016). Our results demonstrate that the mutant reproducibly secretes lower amount of cAMP relative to the WT bacilli, and cAMP secretion is fully restored in the complemented mutant (Compl.). As the fold difference of secretion (~ 1.25- fold) is not much different relative to the fold difference in intra-mycobacterial

cAMP level (~2- fold), we conclude that most likely lower cAMP level of the mutant is not due to its

higher efficacy of cAMP secretion. Fig. 1C confirms absence of autolysis of mycobacterial cells as

GroEL2 was undetectable in the culture filtrates (CF).

PhoP functions as a repressor of *rv0805*

We next investigated role of the *phoP* locus on expression of mycobacterial adenylate

cyclases and phosphodiesterases (PDEs), which synthesize and degrade cAMP, respectively (Fig.

2A). The selection of adenylate cyclases (ACs) and phosphodiesterases (PDEs) were based on two

key points. First, we have chosen ACs which are activated by known signals (Knapp & McDonough,

2014). Secondly, we reasoned that the previously reported ACs were activated under environmental

conditions, which are linked to mycobacterial *phoP* locus (Bansal *et al.*, 2017; Goar *et al.*, 2022). Our

RT-qPCR results using gene-specific primer pairs (Table S1) suggest that expression of adenylate

cyclases including *rv0386*, *rv1264*, *rv1647* and *rv2488c* (Agarwal *et al.*, 2009; Dass *et al*, 2008;

Dittrich *et al*, 2006; Knapp & McDonough, 2014) do not appear to be regulated by the *phoP* locus.

However, a significant activation of adenylate cyclase *rv0891c* (4±0.05-fold), and repression of

phosphodiesterase *rv0805* expression (6.5±0.7-fold), respectively, were dependent on the *phoP* locus.

Although Rv0891c was suggested as one of the *M. tuberculosis* H37Rv adenylate cyclases, the

protein lacks most of the important residues conserved for adenylate cyclase family of proteins

(Zaveri *et al*, 2021). On the other hand, *rv0805* encodes for a cAMP specific PDE, present only in

slow growing pathogenic *M. tuberculosis* (Matange *et al*, 2013; Shenoy *et al*, 2007; Shenoy *et al*,

2005). Although expression of *rv0805* was restored at the level of WT in the complemented mutant

(Compl.), we observed poor restoration of expression of *rv0891c* in the Compl. strain. Further,

expression of PDE *rv1339* which contributes to mycobacterial cAMP level (Thomson *et al*, 2022)

remains unaffected by the *phoP* locus. Therefore, we focussed our attention on the biological

significance of PhoP-dependent regulation of *rv0805*. It should be noted that expressions of *rv1357c*

and *rv2837c*, encoding PDEs for cyclic di-GMP (Flores-Valdez *et al*, 2015) and cyclic di-AMP

(Valadares & Woo, 2017), respectively, remained unchanged in *phoPR*-KO.

183   As we reproducibly observed activation of *rv0805* expression in *phoPR*-KO (relative to WT),

we investigated whether acidic pH conditions under which *phoPR* system is activated (Abramovitch

et al., 2011; Bansal *et al.*, 2017), impacts expression of *rv0805* in WT bacilli (Fig. 2B). Our results

show that repression of *rv0805* is significantly higher in WT grown under acidic conditions (pH 4.5)

relative to normal conditions (pH 7.0) of growth. This observation is consistent with RNA-seq data

displaying significant down-regulation of *rv0805* in WT bacilli grown under acidic pH conditions

relative to normal conditions of growth (GEO Accession number: GSE180161). As a control,

expression of PDE *rv1339* which also degrades cAMP, remains unaffected under acidic conditions of

growth. The finding that acidic pH (pH4.5) conditions of growth promoted PhoP-dependent

repression of *rv0805*, prompted us to investigate whether PhoP directly binds to *rv0805* promoter. To

this end, we next examined *in vivo* recruitment of PhoP within the *rv0805* promoter by chromatin

immunoprecipitation (ChIP) experiments (Fig. 2C). In this assay, formaldehyde-cross-linked DNA-

protein complexes of growing *M. tuberculosis* cells were sheared to generate fragments of average

size ≈500 bp. Next, ChIP experiments utilized anti-PhoP antibody, and IP DNA was analysed by real-

time qPCR relative to a mock sample (without antibody as a control) using FPrv0805up/RPrv0805up

as the primer pair (Table S1). Our results show that under normal condition (empty bars) rv0805up

showed an insignificant enrichment of PCR signal for PhoP relative to non-specific (16S rRNAup)

recruitment (compare empty bars with black bars). This is suggestive of low affinity rv0805up

binding of PhoP. However, IP samples from cells grown under acidic pH showed a significant

enrichment of PhoP at the *rv0805* promoter (compare empty bars with black bars). Notably, ChIP data

showing PhoP recruitment under acidic pH conditions is in agreement with low pH-specific impact of

PhoP on *rv0805* expression (Fig. 2B). Note that PhoP binding to *msl3* promoter region (msl3up) was

used as a positive control.

Probing PhoP-dependent regulation of *M. tuberculosis rv0805*

To examine whether the regulatory effect was attributable to PhoP activation via phosphorylation, we next grew *phoPR*-KO complemented with either *phoP* (Fig. 3A) or the entire *phoPR* encoding ORF (Fig. 3B), both under normal (pH 7.0; empty bars) and acidic (pH 4.5; black bars) conditions of growth and compared relative expression of *rv0805*. Although both strains expressed *phoP*, the former strain lacked a functional copy of *phoR*, the cognate sensor kinase which phosphorylates PhoP (Gupta *et al.*, 2006). Importantly, *M. tuberculosis* H37Rv lacking a *phoR* gene (*phoPR*-KO::*phoP*) failed to show a low pH-dependent repression of *rv0805* expression. However, similar to WT bacilli, we observed a low pH-dependent significant down-regulation of *rv0805* expression in *phoPR*-KO::*phoPR* (Compl.). Note that a comparable expression of PDE *rv1339* was observed in both strains regardless of growth conditions. These results confirm that acidic pH-dependent repression of *rv0805* expression *in vivo* is indeed attributable to P~PhoP requiring the presence of PhoR.

To examine the effect of Rv0805 on mycobacterial cAMP level, we next expressed a copy of *rv0805* in WT bacteria (referred to as WT-Rv0805) (Fig. 3C). *rv0805* ORF was cloned within the multicloning site of pSTki (Parikh *et al*, 2013) between the EcoRI and HindIII sites under the control of $P_{myc1}tetO$ promoter, and ectopic expression of *rv0805* was verified by determining the mRNA level (Figure 3 - figure supplement 1A). We then assessed the impact of Rv0805 on intra-mycobacterial cAMP. Consistent with a previous study (Agarwal *et al.*, 2009), WT-Rv0805 showed a significant depletion (2.1± 0.7-fold) of intra-mycobacterial cAMP relative to WT bacteria. To confirm that the reduced level of mycobacterial cAMP is attributable to *rv0805* expression, we also expressed *rv0805M*, a mutant Rv0805 lacking phosphodiesterase activity. As structural data coupled with biochemical evidences reveal that Asn-97 at the enzyme active site plays a key role in phosphodiesterase activity of Rv0805 (Shenoy *et al.*, 2007; Shenoy *et al.*, 2005), the mutant Rv0805M was constructed by changing the conserved Asn-97 to Ala. Most importantly, WT-Rv0805M showed an insignificant variation of cAMP level relative to WT, suggesting that depletion of intra-mycobacterial cAMP in WT-Rv0805 is indeed attributable to phosphodiesterase activity of

Rv0805. However, the corresponding mRNA levels of PDEs in WT suggest that both the WT and

mutant Rv0805 are over-expressed approximately 4.5-6 -fold relative to the genomic *rv0805* level of

WT (Figure 3 - figure supplement 1A). In contrast, other PDEs under identical conditions,

demonstrate comparable expression levels in *rv0805* over-expressing strains and WT. Thus, we

conclude that a significantly different protein levels is unlikely to account for variable cAMP levels in

the two PDEs (WT and mutant Rv0805) over-expressing strains utilizing identical expression

strategy. Notably, over-expression of these PDEs did not influence bacterial growth under normal

conditions (Figure 3 - figure supplement 1B).

To further probe regulation of Rv0805 expression and its control of intra-mycobacterial

cAMP homeostasis, we utilized a previously reported CRISPRi based approach (Singh *et al*, 2016) to

construct *rv0805* and *phoP* knockdown (*rv0805*-KD, and *phoP-KD*, respectively) mutants.

Importantly, consistent with *phoPR*-KO, *phoP*-KD shows a significantly higher *rv0805* expression in

the presence of ATc relative to its absence (Fig. 3D). However, despite a significant down regulation

of *rv0805* expression in presence of ATc, a comparable *phoP* expression was observed in *rv0805*-KD

mutant both in the absence or presence of ATc. As a control, we observed a comparable expression of

16S rRNA in both knock-down mutants. Next, we determined intra-mycobacterial cAMP of the

mutants as described in Fig. 1 (Fig. 3E). Importantly, cAMP level of *phoP*-KD (showing activation of

Rv0805) was significantly lower relative to WT bacteria. In contrast, *rv0805*-KD mutant

demonstrated a significantly higher level of cAMP relative to WT. Together, these data represent an

integrated view of our results suggesting that PhoP-dependant repression of *rv0805* regulates intra-

mycobacterial cAMP homeostasis. In keeping with these results, activated PhoP under acidic pH

conditions significantly represses *rv0805*, and intracellular mycobacteria most likely utilizes a higher

level of cAMP to effectively mitigate stress response for survival under stressful environment of the

phagosome.

PhoP contributes to mycobacterial stress tolerance and intracellular survival by maintenance of

cAMP homeostasis.

To investigate whether cAMP level influences mycobacterial susceptibility to stress, we

compared *in vitro* growth under acidic pH (pH 4.5) (Fig. 4A). As expected, *phoPR*-KO showed a

significant growth inhibition relative to WT under low pH (pH 4.5) (Bansal *et al.*, 2017). Remarkably,

WT-Rv0805, but not WT-Rv0805M displayed a comparable susceptibility to acidic pH as that of

phoPR-KO. However, all four mycobacterial strains showed comparable growth at pH 7.0. Next, to

compare growth of WT-Rv0805 and WT under oxidative stress, cells were grown in presence of

increasing concentrations of diamide, a thiol-specific oxidant, and examined by microplate-based

Alamar Blue assays (Figure 4 - figure supplement 1A). We have recently shown that *phoPR*-KO is

significantly more sensitive to diamide relative to WT (Goar *et al.*, 2022). Here, we uncovered that

similar to *phoPR*-KO, WT-Rv0805, but not WT-Rv0805M was significantly more susceptible to

diamide stress as compared to WT (Fig. 4B). Notably, a previous study had reported that *phoP*-deleted

mutant strain was more sensitive to Cumene Hydrogen Peroxide (CHP), suggesting role of PhoP in

regulating mycobacterial transcriptome and stress response (Walters *et al.*, 2006). To compare

sensitivity to CHP, we next grew mycobacterial strains in presence of 50 μM CHP for 24 hours and

determined their survival by enumerating CFU values (Fig. 4C). Note that in this case we were unable

to perform Alamar Blue-based survival assays requiring a longer time because of the bactericidal

property of CHP. Our CFU data highlight that WT-Rv0805, but not WT-Rv0805M, displayed a

significantly higher growth inhibition relative to WT in the presence of CHP. Together, these results

demonstrate a comparably higher susceptibility of *phoPR*-KO, and WT-Rv0805 to acidic pH and

oxidative stress relative to WT and establish a link between cAMP homeostasis and mycobacterial

stress response, suggesting that at least one of the mechanisms by which PhoP contributes to global

stress response is attributable to maintenance of mycobacterial cAMP homeostasis.

A previous study showed that *rv0805* over-expression in *M. smegmatis* influences cell wall

permeability (Podobnik *et al*, 2009). Having shown a significantly higher sensitivity of WT-Rv0805

to low pH and oxidative stress (relative to WT), we sought to investigate whether altered cell wall

structure/properties of the mycobacterial strain contribute to elevated stress sensitivity. Thus, we

compared expression level of lipid biosynthetic genes, which encode part of cell-wall structure of the bacilli (Gonzalo Asensio *et al.*, 2006a; Walters *et al.*, 2006). Our results clearly suggest that in contrast to *phoPR*-KO, both WT-Rv0805 and WT-Rv0805M share a comparable expression profile of complex lipid biosynthesis genes as that of WT (Figure 4 - figure supplement 1B). These results suggest that both strains expressing WT- or the mutant PDEs share a similar cell-wall properties and are consistent with a recent study reporting no significant effect of cAMP dysregulation on mycobacterial cell wall structure/ permeability (Wong *et al*, 2022). Together, our findings facilitate an integrated view of our results suggesting that higher susceptibility of WT-Rv0805 to stress conditions, is most likely controlled by mycobacterial cAMP level.

To investigate the impact of mycobacterial cAMP level *in vivo*, we studied infection of murine macrophages using WT, WT-Rv0805, and WT-Rv0805M (Fig. 4D). Although WT bacilli inhibited phagosome maturation, infection of macrophages with WT-Rv0805 and *phoPR*-KO matured into phagolysosomes, suggesting increased trafficking of the bacilli to lysosomes. Interestingly, under identical conditions, WT-Rv0805M could effectively inhibit phagosome maturation just as WT bacteria. Results from co-localization experiments are plotted as Fig. 4E and as Pearson's correlation co-efficient of the quantified co-localization signals as Fig. 4F. These data suggest reduced ability of WT-Rv0805, but not WT-Rv0805M (relative to WT) to inhibit phagosome maturation. From these results, we conclude that ectopic expression of *rv0805* impacts phagosome maturation arguing in favour of a role of PhoP in influencing phagosome-lysosome fusion in macrophages most likely by maintenance of mycobacterial cAMP homeostasis.

Intra-bacterial cAMP level and its effect on *in vivo* survival of mycobacteria

To examine the effect *in vivo*, mice were infected with mycobacterial strains via the aerosol route. Day 1 post-infection, CFU analyses revealed a comparable count of four mycobacterial strains (~100 bacilli) in the mice lungs. However, for WT-Rv0805, the CFU recovered from infected lungs 4 weeks post- infection declined by ≈218-fold relative to the lungs infected with WT bacteria (Fig. 5A).

In contrast, the CFU recovered from infected lungs after 4 weeks of infection by WT-Rv0805M

marginally declined by ≈7-fold relative to the lungs infected with WT bacilli. These results suggest a

significantly compromised ability of WT-Rv0805 (relative to WT) to replicate in the mice lungs. Note

that *phoPR*-KO, under the conditions examined, showed a ≈ 246-fold lower lung burden compared to

WT. In keeping with these results, while the WT bacilli disseminated to the spleens of infected mice,

a significantly lower count of WT-Rv0805 was recovered from the spleens after 4 weeks of infection

(Fig. 5B). Thus, we conclude that one of the reasons which accounts for an attenuated phenotype of

phoPR-KO in both cellular and animal models is likely attributable to PhoP-dependent repression of

rv0805 PDE activity, which controls mycobacterial cAMP homeostasis.

M. tuberculosis H37Rv persists within granulomas where it is protected from the anti-

mycobacterial immune effectors of the host. Histopathological evaluations showed that WT bacilli -

infected lung sections displayed aggregation of granulocytes within alveolar spaces which degenerate

progressively to necrotic cellular debris. In contrast, *phoPR*-KO and WT-Rv0805 showed less severe

pathology as indicated by decreased tissue consolidation, smaller granulomas, and open alveolar

space (Fig. 5C). Together, these results suggest that ectopic expression of *rv0805* in WT bacilli is

somewhat functionally equivalent to deletion of *phoP*, suggesting that failure to maintain cAMP

homeostasis most likely accounts for attenuated phenotype of the bacilli and absence of

immunopathology in the lungs of infected mice.

Discussion

A number of studies suggest that conditions associated with host environment like low pH, and

macrophage interactions often influence mycobacterial cAMP levels (Bai *et al.*, 2009; Gazdik &

McDonough, 2005). Although many bacterial pathogens modulate host cell cAMP levels as a

common strategy, the mechanism of mycobacterial cAMP-homeostasis remains unknown. In this

study, we sought to investigate whether PhoP, a master regulator implicated in controlling diverse

mycobacterial stress response, is connected to mycobacterial cAMP level. Intriguingly, we find that

under normal conditions as well as under carefully controlled single stress conditions, *phoPR*-KO

shows a significantly lower level of cAMP relative to the WT bacilli (Fig. 1A), and complementation

of the mutant restored cAMP level. To investigate the mechanism, we next probed regulation of

adenylate cyclases and PDEs (Fig. 2) and unambiguously demonstrated that PhoP functions as a

major repressor of *rv0805*, encoding cAMP specific PDE. Indeed, this newly-identified *rv0805*

regulation, coupled with a very recent discovery that phosphodiesterase activity of Rv0805 controls

propionate detoxification (McDowell *et al*, 2023), fits well with and explains, the previously puzzling

in vivo observation by Abramovitch et al (Abramovitch *et al.*, 2011) that PhoP -controlled *aprABC*

locus is associated with the regulation of genes of carbon and propionate metabolism.

Although a large number of ACs are present in *M. tuberculosis* genome, a class III metallo-

phosphoesterase Rv0805 was earlier considered the only PDE, specific for mycobacterial cAMP.

However, a recent study has identified an atypical class II PDE Rv1339, which upon overexpression

reduces cAMP level and contributes to antibiotic sensitivity (Thomson *et al.*, 2022). While, the

functional role of Rv1339 in *M. tuberculosis* is yet to be understood, crystal structure and biochemical

evidence suggest that dimeric Rv0805 is stabilized by the presence of a divalent cation, and remains

catalytically active on a broad range of linear and cyclic PDE substrates *in vitro* (Keppetipola &

Shuman, 2008; Shenoy *et al.*, 2007; Shenoy *et al.*, 2005). More recently, cyclic nucleotide hydrolytic

activity of mycobacterial Rv0805 has been implicated in propionate detoxification (McDowell *et al.*,

2023). However, the mechanisms of regulation of Rv0805 and maintenance of mycobacterial cAMP

homeostasis remain unknown.

To examine biological significance of PhoP-dependent Rv0805 expression, we studied

rv0805 expression under acidic conditions of growth as *phoPR* system is induced under acidic pH

both *in vitro* and in macrophages (Abramovitch *et al.*, 2011; Bansal *et al.*, 2017). Strikingly, a

significantly higher repression of *rv0805* expression under acidic pH relative to normal conditions is

consistent with activation of PhoP and subsequent repression of *rv0805* (Fig. 3). These results further

suggest that effective mitigation of stress response by mycobacteria possibly requires a higher cAMP

level for survival under intra-phagosomal environment. The most fundamental biological insight is

derived from the finding that PhoP-dependent *rv0805* repression requires PhoR (Figs. 3A-B), the

cognate kinase which activates PhoP in a signal-dependent manner (Gupta *et al.*, 2006). These results

account for a consistently lower level of cAMP in *phoPR*-KO relative to the WT bacilli. Notably,

except recently reported PDE Rv1339, Rv0805 has been known as the only cAMP-specific PDE

present in the slow growing pathogenic mycobacteria and its closely related species (Matange, 2015;

Shenoy *et al.*, 2007; Shenoy *et al.*, 2005), and Rv1339 expression does not appear to be regulated by

the *phoP* locus (Fig. 2). Thus, the above results showing PhoP-dependent repression of *rv0805*

activity likely represents the most critical step of mycobacterial cAMP homeostasis. In this

connection, our recent results that PhoP interacts with cAMP receptor protein, CRP and a complex of

two interacting regulators control expression of virulence determinants (Khan *et al*, 2022), invite

speculation of a complex regulatory control of cAMP-responsive mycobacterial physiology.

386   As one might argue that PhoP deletion and *rv0805* over-expression could be unrelated and

independent events, we constructed *phoP* and *rv0805* knock-down mutants to further investigate the

PhoP-Rv0805-cAMP pathway. Our objective was to probe regulation of expression (Fig. 3D) and

examine the impact on mycobacterial cAMP (Fig. 3E). Importantly, *phoP*-KD significantly elevated

rv0805 expression; however, *phoP* expression remains unaffected in *rv0805*-KD (Fig. 3D). More

importantly, elevated *rv0805* expression in *phoP*-KD reduces cAMP level, whereas cAMP level is

understandably elevated in *rv0805*-KD mutant (Fig. 3E). These results integrate PhoP-dependent

rv0805 repression with mycobacterial cAMP level, suggesting how *phoP*-deletion or knock-down, at

least in part, mimics Rv0805 over-expression. These considerations take on more significance given

the fact that these two events have similar consequences on correspondingly relevant strains with

respect to stress tolerance, and survival in cellular and animal models. Thus, our results appear to

suggest that ectopic expression of *rv0805* is somewhat functionally equivalent to deletion of the *phoP*

locus. This observation is in apparent conflict with a previous work by Matange and collaborators

(Matange *et al.*, 2013), suggesting a cAMP-independent transcriptional response in *rv0805* over-

expressing *M. tuberculosis* H37Rv. Although both studies were performed with *rv0805* over-

expressing bacilli, the fact that important differences in the expression of PDEs in this study (Matange

et al., 2013) and in our assays – yielding significantly different induction of *rv0805* expression - most

likely account for this discrepancy. While we cannot completely rule out the possibility of cleavage of

other cyclic nucleotide(s) by Rv0805 (Keppetipola & Shuman, 2008; Shenoy *et al.*, 2007; Shenoy *et*

al., 2005) impacting our results, consistent with a previous study our results correlate *rv0805*

expression with intra-mycobacterial cAMP level (Agarwal *et al.*, 2009). Further, our data on the effect

of expression of cyclic nucleotide-specific PDE Rv0805 or its inactive mutant (Rv0805M) correlate

well with enzyme activities of the corresponding PDEs on mycobacterial cAMP levels (Fig. 3C).

Thus, we conclude that PhoP-dependent regulation of *rv0805*, most likely, is a critical regulator of

intra-mycobacterial cAMP level.

Our experiments to understand physiological significance of PhoP-dependent repression of

rv0805 expression uncovers a strikingly similar stress tolerance of WT-Rv0805 and *phoPR*-KO

(significantly reduced relative to WT) (Fig. 4). These results are consistent with the notion that cAMP

homeostasis, at least in part, accounts for mycobacterial stress response. Along the line, WT-Rv0805

displayed a reduced ability to inhibit phagosome-lysosome fusion like *phoPR*-KO (Fig. 4). Further,

we show that WT-Rv0805, unlike the WT bacilli or WT-Rv0805M, shows a significantly reduced

intracellular growth in mice as that of *phoPR*-KO (Fig. 5). Thus, these results are of fundamental

significance to establish that PhoP contributes to maintenance of cAMP homeostasis and integrates it

to mechanisms of mycobacterial stress tolerance and intracellular survival. Together, we identify a

novel mycobacterial pathway as a therapeutic target and provide yet another example of an intimate

link between bacterial physiology and intracellular survival of the tubercle bacilli.

The results reported here are summarized schematically in Fig. 6. According to this model,

upon activation by an appropriate signal via the cognate sensor kinase PhoR, P~PhoP binds to *rv0805*

regulatory region and functions as a specific repressor of *rv0805*. Thus, we observed a reproducibly

lower level of cAMP in *phoPR*-KO relative to the WT bacilli. However, from these results we are

unable to claim that virulence attenuation of *phoPR*-KO is attributable to cAMP homeostasis or lack

thereof. While this would demand uncoupling of regulatory control of PhoPR and *rv0805* expression,

further support to PhoP-dependent *rv0805* repression was apparent from a significantly reduced

expression of *rv0805* in WT bacilli grown under acidic pH, a condition which activates mycobacterial

PhoPR system (Abramovitch *et al.*, 2011; Bansal *et al.*, 2017; Tan *et al.*, 2013). In keeping with these

results, we show comparable cAMP levels in *phoPR*-KO and WT-Rv0805. Thus, the above two

strains remain ineffective to mount an appropriate stress response most likely due to their inability to

coordinate regulation of gene expression because of dysregulation of cAMP homeostasis, facilitates

an integrated view of our results. Given the fact that *rv0805*-depleted *M. tuberculosis* is growth

attenuated *in vivo* (McDowell *et al.*, 2023), paradoxically ectopic expression of *rv0805* leads to

dysregulated metabolic adaptation, thereby resulting in reduced stress tolerance and intracellular

survival.

Materials and Methods

Bacterial strains and culture conditions. *E. coli* DH5α was used for cloning experiments. *E. coli* BL21(DE3), an *E. coli* B strain lysogenized with λDE3, a prophage expressing T7 RNA polymerase from the IPTG (isopropyl-β-D-thiogalactopyranoside)-inducible *lac*UV5 promoter (Studier & Moffatt, 1986) was used as the host for over-expression of recombinant proteins. *E. coli* strains were grown in LB medium at 37°C with shaking, transformed according to standard procedures and the transformants were selected on media containing appropriate antibiotics. WT- and mutant *M. tuberculosis* and *M. smegmatis* mc^{2}155 were grown at 37°C in Middlebrook 7H9 broth (Difco) containing 0.2% glycerol, 0.05% Tween-80 and 10% ADC (albumin-dextrose-catalase) or on 7H10-agar medium (Difco) containing 0.5% glycerol and 10% OADC (oleic acid-albumin-dextrose-catalase) enrichment. *phoPR* disruption mutant of *M. tuberculosis* H37Rv (*phoPR*-KO, a kind gift of Dr. Issar Smith) was constructed as described (Walters *et al.*, 2006). To this end, a kanamycin-resistant cassette from pUC-K4 was inserted into a unique EcoRV site within the coding region of *phoP* gene, and disruption was confirmed by Southern blot analysis of chromosomal DNA isolated from the mutant. Next, purified plasmid DNAs were electroporated into wild-type *M. tuberculosis* strain by standard protocol (Jacobs *et al*, 1991). To complement expression, pSM607 containing a 3.6-kb DNA fragment of *phoPR* including 200-bp *phoP* promoter region, a hygromycin resistance cassette, *attP* site and the gene encoding phage L5 Integrase, as described previously (Walters *et al.*, 2006) was transformed in the *phoPR* mutant strain to integrate at the L5 *attB* site. Growth, transformation of wild-type (WT), *phoPR*-KO, the complemented mutant (Compl.) *M. tuberculosis* and selection of transformants on appropriate antibiotics were performed as described (Goyal *et al.*, 2011). When appropriate, antibiotics were used at the following concentrations: hygromycin (hyg), 250 μg/ml for *E. coli* or 50 μg/ml for mycobacterial strains; streptomycin (str), 100 μg/ml for *E. coli* or 20 μg/ml for mycobacterial strains; kanamycin (kan), 20 μg/ml for mycobacterial strains. For *in vitro* growth under specific stress conditions, indicated mycobacterial strains were grown to mid log phase (OD$_{600}$ 0.4-0.6) and exposed to different stress conditions. For acid stress, cells were initially grown in 7H9 media, pH7.0 and on attaining mid log phase it was transferred to acidic media (7H9

media, pH 4.5) for further two hours at 37˚C. For oxidative stress, cells were grown in presence of 50

µM CHP (Sigma) for 24 hours or indicated diamide concentration(s) for 7 days. For NO stress, cells

grown to mid log phase were exposed to 0.5 mM DataNonoate for 40 minutes (Voskuil *et al*, 2003).

cAMP measurement. Mycobacterial cell pellets were collected and washed with 1x PBS buffer, cells

were resuspended in IP buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA pH 8.0, 1 mM

PMSF, 5% glycerol and 1% TritonX 100) and cell lysates (CL) were prepared by lysing the cells in

presence of Lysing Matrix B (100 µm silica beads; MP Bio) using a FastPrep-24 bead beater (MP

Bio) at a speed setting of 6.0 for 30 seconds. The procedure was repeated for 10 cycles with

incubation on ice in between pulses. The supernatant was collected by centrifugation at 13,000 rpm

for 10 minutes and filtered through 0.22 µm filter (Millipore). cAMP levels in the cells were

determined in a plate reader by using fluorescence-based cAMP detection kit (Abcam) according to

the manufacturer's recommendations and normalized to the total protein present in the samples as

determined by a BCA protein estimation kit (Pierce). For secretion studies, each mycobacterial strain

was grown in Sauton's media as described (Anil Kumar *et al.*, 2016), comparable counts of bacterial

cells were pelleted, resuspended in 2 ml of Sauton's media in a 6-well plate format for 2 hours at

37°C, and the supernatants (culture filtrates, CF) were collected for cAMP measurements, as

described previously (Anil Kumar *et al.*, 2016).

Cloning. *M. tuberculosis* full-length ORF was cloned between EcoRI and HindIII sites of the

mycobacterial expression vector pSTKi (Parikh *et al.*, 2013) and expressed from the $P_{myc1}tetO$

promoter. Mutation in Rv0805 was introduced by two-stage overlap extension method using

mutagenic primers (Table S2), and the construct was verified by DNA sequencing. For over-

expression of WT- or mutant PDEs, WT bacilli was transformed with pST-rv0805 or pST-rv0805M to

generate WT-Rv0805 or WT-Rv0805M, respectively.

Construction of *M. tuberculosis phoP* and *rv0805* knock-down mutants. In this study, we utilized

a previously reported CRISPRi system (Singh *et al.*, 2016) to construct knock-down mutants of *phoP*

and r*v0805* (*phoP*-KD, and *rv0805*-KD, respectively). This approach efficiently inhibits expression of

target genes via inducible expression of dCas9 along with gene specific guide RNAs (sgRNA). The

oligonucleotides were designed such that the expressed sgRNA comprises of a 20 bp sequence

complementary to the non-template strand of the target gene. The sgRNAs of *phoP* and *rv0805* were

cloned in a vector pRH2521 using BbsI enzyme and the constructs were confirmed by sequencing.

The corresponding clones were transformed into *M. tuberculosis* harbouring pRH2502, a vector

expressing an inactive version of *Streptococcus pyogenes* cas9 (dcas9). To express dcas9 and repress

sgRNA-targeted genes (*phoP* or *rv0805*), the bacterial strains were grown with or without 600 ng/ml

of anhydro-tetracycline (ATc) every 48 hours, and cultures were grown for 4 days. RNA isolation was

carried out, and RT-qPCR experiments verified significant repression of target genes. For the induced

strains (in presence of ATc) expressing sgRNAs targeting +155 to +175 (relative to *phoP* translational

start site) and +224 to +244 sequences (relative to *rv0805* translational start site), we obtained

approximately 85% and 90% reduction of *phoP* and *rv0805* RNA abundance, respectively, relative to

corresponding un-induced strains. The oligonucleotides used to generate gene-specific sgRNA

constructs and the plasmids utilized in knock-down experiments are listed in supplementary Table S2.

RNA isolation. Total RNA was extracted from exponentially growing bacterial cultures grown with

or without specific stress as described above. Briefly, 25 ml of bacterial culture was grown to mid-log

phase (OD_{600}= 0.4 to 0.6) and combined with 40 ml of 5 M guanidinium thiocyanate solution

containing 1% β-mercaptoethanol and 0.5% Tween 80. Cells were pelleted by centrifugation, and

lysed by re-suspending in 1 ml Trizol (Ambion) in the presence of Lysing Matrix B (100 μm silica

beads; MP Bio) using a FastPrep-24 bead beater (MP Bio) at a speed setting of 6.0 for 30 seconds.

The procedure was repeated for 2-3 cycles with incubation on ice in between pulses. Next, cell lysates

were centrifuged at 13000 rpm for 10 minutes; supernatant was collected and processed for RNA

isolation using Direct-Zol™ RNA isolation kit (ZYMO). Following extraction, RNA was treated with

DNAse I (Promega) to degrade contaminating DNA, and integrity was assessed using a Nanodrop

(ND-1000, Spectrophotometer). RNA samples were further checked for intactness of 23S and 16S

rRNA using formaldehyde-agarose gel electrophoresis, and Qubit fluorometer (Invitrogen).

Quantitative Real-Time PCR. cDNA synthesis and PCR reactions were carried out using total RNA

extracted from each bacterial culture, and Superscript III platinum-SYBR green one-step qRT-PCR

kit (Invitrogen) with appropriate primer pairs (2 μM) using an ABI real-time PCR detection system.

Oligonucleotide primer sequences used in RT-qPCR experiments are listed in Table S1. Control

reactions with platinum Taq DNA polymerase (Invitrogen) confirmed absence of genomic DNA in all

our RNA preparations, and endogenously expressed *M. tuberculosis rpoB* was used as an internal

control. Fold difference in gene expression was calculated using $\Delta\Delta C_T$ method (Schmittgen & Livak,

2008). Average fold differences in mRNA levels were determined from at least two biological repeats

each with two technical repeats. Non-significant difference is not indicated.

ChIP assays - ChIP experiments in actively growing cultures of *M. tuberculosis* were carried out

essentially as described previously (Fol *et al*, 2006). Immunoprecipitation (IP) was performed using

anti-PhoP antibody and protein A/G agarose beads (Pierce). qPCR reactions included PAGE purified

primer pairs (Table S1) spanning specific promoter regions using suitable dilutions of

immunoprecipitated (IP) DNA in a reaction buffer containing SYBR green mix, and one unit of

Platinum Taq DNA polymerase (Invitrogen). An IP experiment without adding antibody (mock) was

used as the negative control, and data was analysed relative to PCR signal from the mock sample.

PCR amplifications were carried out for 40 cycles using serially diluted DNA samples (mock, IP

treated and total input) in a real-time PCR detection system (Applied Biosystems). In all cases melting

curve analysis confirmed amplification of a single product.

Immunoblotting - Cell lysates or culture filtrates were resolved by 12% SDS-PAGE and visualized by Western blot analysis using appropriate antibodies. Briefly, resolved samples were electroblotted onto polyvinyl difluoride (PVDF) membranes (Millipore, USA) and were detected by anti-GroEL2 antibody (Sigma), anti CFP-10 antibody (Abcam) or affinity-purified anti-PhoP antibody elicited in rabbit (Alpha Omega Sciences, India). Goat anti-rabbit secondary antibody conjugated to horseradish peroxidase was used, and blots were developed with Luminata Forte Chemiluminescence reagent (Millipore, USA). RNA polymerase was used as a loading control and was detected with monoclonal antibody against β-subunit of RNA polymerase, RpoB (Abcam).

Alamar Blue assay. In this assay, reduction of Alamar Blue correlates with the change of a non-fluorescent blue to a fluorescent pink appearance, which is directly linked to bacterial growth. *M. tuberculosis* H37Rv was grown in 7H9 media (Difco) with 10% ADS (albumin, dextrose and NaCl) to an OD_{600} of 0.4, and freshly diluted to OD_{600} of 0.02. Next, increasing concentrations of diamide was added to the wells of a 96 -well plate containing 0.05 ml 7H9 media followed by addition of 0.05 ml of *M. tuberculosis* H37Rv culture (0.02 OD_{600}). The plate was incubated at 37°C for 7 days. Finally, 0.02 ml of 0.02% Resazurin (sodium salt, MP Bio), prepared in sterile 7H9 media was added to each of the wells and the change in colour was examined after incubation at 37°C for 16 hours. The fluorescence excitation was at 530 nm and emission was recorded at 590 nm. Efficiency of inhibition was calculated relative to control wells which did not include diamide, and Rifampicin was included as a positive control to confirm validity of the assay.

Macrophage Infections. Virulence of indicated H37Rv strains were assessed in murine macrophages according to the previously published protocol (Solans *et al.*, 2014). Briefly, RAW264.7 macrophages were grown in DMEM media containing 10% Fetal bovine serum at 37°C under 5% CO_2, and seeded onto #1 thickness, 18 mm diameter glass coverslips in a 12-well plate at a 40% confluency (0.5 million cells). Cells were independently infected with titrated cultures of WT, WT-Rv0805, WT-Rv0805M and *phoPR*-KO strains at a multiplicity of infection (MOI) of 1:5 for 3 hours at 37°C in 5%

CO_2, followed by 1X PBS washes thrice. The macrophages were further incubated for 3 hours at

37°C. After infection, extracellular bacteria were removed by washing thrice with PBS. To visualize

trafficking of the tubercle bacilli, mycobacterial strains were stained with phenolic auramine solution

(which selectively binds to mycolic acids) for 15 minutes followed by washing with acid alcohol

solution and finally with 1X PBS. The cells were stained with 150 nM acidotropic dye LysoTracker

Red DND-99 (Invitrogen) for 30 minutes in a CO_2 incubator. Next, the cells were fixed with 4%

paraformaldehyde for 15 minutes, washed thrice with PBS, the coverslips were mounted in Slow

Fade-Anti-Fade (Invitrogen) and analyzed using laser scanning confocal microscope (Nikon)

equipped with Argon (488 nm excitation line; 510 nm emission detection) and LD (561 nm excitation

line; 594 nm emission detection) laser lines. Digital images were processed with image-processing

software (Nikon). The settings of the laser and the detector were optimized using macrophage cells

infected with WT bacteria and the same standard set of intensity thresholds were applied to all

images.

Mouse infections.

All experiments pertaining to mice were in accordance with Institutional regulations after review of

protocols and approval by the Institutional Animal Ethics Committee (IAEC/17/05, and IAEC/19/02).

Mice were maintained and bred in the animal house facility of CSIR-IMTECH. Animal infection

studies and subsequent experiments were carried out in the Institutional BSL-3 facility as per

institutional biosafety guidelines. Briefly, the experiments were conducted with 8-10 weeks old

C57BL/6 mice, infected intranasally and euthanized post-infection for evaluation of bacterial load in

lungs and spleens. Infections were given through the respiratory route using an inhalation exposure

system (Glass-col) with passaged *M. tuberculosis* H37Rv cultures of mid log phase. The actual

bacterial load delivered to the animals was enumerated from 3-5 aerogenically challenged mice, 1day

post aerosol challenge. The animals were found to achieve a bacillary deposition of 100 to 200 CFU

in the lungs for each strain. Four weeks post infection, the animals were sacrificed by cervical

dislocation, lungs and spleens were isolated aseptically from the euthanized animals, homogenized in

sterile 1X PBS and plated after serially diluting the lysates on 7H11 agar plates, supplemented with

10% OADC and antibiotics (50 μg/ml carbenicillin, 30 μg/ml polymyxin B, 10 μg/ml vancomycin, 20

μg/ml Trimethoprim, 20 μg/ml cycloheximide, and 20 μg/ml Amphotericin B) to enumerate CFU. For

histopathology, left lung lobes were fixed in 10% buffered formalin, embedded in paraffin and stained

with haematoxylin and eosin for visualization under the microscope. The level of pathology was

scored by analysing perivascular cuffing, leukocyte infiltration, multinucleated giant cell formation

and epithelial cell injury.

Statistical analysis. Data are presented as arithmetic means of the results obtained from multiple

replicate experiments ± standard deviations. Statistical significance was determined by Student's

paired t-test using Microsoft Excel or Graph Pad Prism. Statistical significance was considered at P

values of 0.05.

Acknowledgements

We are grateful to G. Marcela Rodriguez and Issar Smith (PHRI, New Jersey Medical School -

UMDNJ) for Δ*phoP*-H37Rv, and the complemented *M. tuberculosis* H37Rv strains, Adrie Steyn

(University of Alabama) for pUAB300/pUAB400 plasmids, Ashwani Kumar for very helpful

discussions, and Sanjeev Khosla for critical reading of the manuscript. We thank members of the

Institutional animal facility (iCARE) for their help with approval of our projects from the Institutional

Animal Ethics Committee. This study received financial support from intramural grants of CSIR-

IMTECH (OLP-0170), CSIR (MLP-0049) and by a research grant (to D.S) from SERB

(EMR/2016/004904), Department of Science and Technology (DST). H.K., P.P., H.G., B.B. and N.B.

were supported by CSIR pre-doctoral fellowships.

Supplemental Data

The supplemental data include three (3) supplemental figures, and two supplemental tables

(Supplemental files 1a, and 1b, respectively). All data generated or analysed during this study are

included in the manuscript and supporting file; source data files have been provided for all relevant

figures.

Competing interests

The authors declare that no competing interests exist.

REFERENCES

Abramovitch RB, Rohde KH, Hsu FF, Russell DG (2011) aprABC: a Mycobacterium tuberculosis complex-specific locus that modulates pH-driven adaptation to the macrophage phagosome. *Molecular microbiology* 80: 678-694

Agarwal N, Lamichhane G, Gupta R, Nolan S, Bishai WR (2009) Cyclic AMP intoxication of macrophages by a Mycobacterium tuberculosis adenylate cyclase. *Nature* 460: 98-102

Agarwal N, Raghunand TR, Bishai WR (2006) Regulation of the expression of whiB1 in Mycobacterium tuberculosis: role of cAMP receptor protein. *Microbiology* 152: 2749-2756

Anil Kumar V, Goyal R, Bansal R, Singh N, Sevalkar RR, Kumar A, Sarkar D (2016) EspR-dependent ESAT-6 Protein Secretion of Mycobacterium tuberculosis Requires the Presence of Virulence Regulator PhoP. *The Journal of biological chemistry* 291: 19018-19030

Arbues A, Aguilo JI, Gonzalo-Asensio J, Marinova D, Uranga S, Puentes E, Fernandez C, Parra A, Cardona PJ, Vilaplana C *et al* (2013) Construction, characterization and preclinical evaluation of MTBVAC, the first live-attenuated M. tuberculosis-based vaccine to enter clinical trials. *Vaccine* 31: 4867-4873

Bai G, Knapp GS, McDonough KA (2011) Cyclic AMP signalling in mycobacteria: redirecting the conversation with a common currency. *Cellular microbiology* 13: 349-358

Bai G, McCue LA, McDonough KA (2005) Characterization of Mycobacterium tuberculosis Rv3676 (CRPMt), a cyclic AMP receptor protein-like DNA binding protein. *Journal of bacteriology* 187: 7795-7804

Bai G, Schaak DD, McDonough KA (2009) cAMP levels within Mycobacterium tuberculosis and Mycobacterium bovis BCG increase upon infection of macrophages. *FEMS immunology and medical microbiology* 55: 68-73

Baker JJ, Dechow SJ, Abramovitch RB (2019) Acid Fasting: Modulation of Mycobacterium tuberculosis Metabolism at Acidic pH. *Trends in microbiology* 27: 942-953

Baker JJ, Johnson BK, Abramovitch RB (2014) Slow growth of Mycobacterium tuberculosis at acidic pH is regulated by phoPR and host-associated carbon sources. *Molecular microbiology* 94: 56-69

Banerjee A, Adolph RS, Gopalakrishnapai J, Kleinboelting S, Emmerich C, Steegborn C, Visweswariah SS (2015) A universal stress protein (USP) in mycobacteria binds cAMP. *J Biol Chem* 290: 12731-12743

Bansal R, Anil Kumar V, Sevalkar RR, Singh PR, Sarkar D (2017) Mycobacterium tuberculosis virulence-regulator PhoP interacts with alternative sigma factor SigE during acid-stress response. *Molecular microbiology* 104: 400-411

Dass BK, Sharma R, Shenoy AR, Mattoo R, Visweswariah SS (2008) Cyclic AMP in mycobacteria: characterization and functional role of the Rv1647 ortholog in Mycobacterium smegmatis. *Journal of bacteriology* 190: 3824-3834

Dittrich D, Keller C, Ehlers S, Schultz JE, Sander P (2006) Characterization of a Mycobacterium tuberculosis mutant deficient in pH-sensing adenylate cyclase Rv1264. *International journal of medical microbiology : IJMM* 296: 563-566

Ehrt S, Schnappinger D, Rhee KY (2018) Metabolic principles of persistence and pathogenicity in Mycobacterium tuberculosis. *Nature reviews Microbiology* 16: 496-507

Ernst JD (2012) The immunological life cycle of tuberculosis. *Nature reviews Immunology* 12: 581-591

Flores-Valdez MA, Aceves-Sanchez Mde J, Pedroza-Roldan C, Vega-Dominguez PJ, Prado-Montes de Oca E, Bravo-Madrigal J, Laval F, Daffe M, Koestler B, Waters CM (2015) The Cyclic Di-GMP Phosphodiesterase Gene Rv1357c/BCG1419c Affects BCG Pellicle Production and In Vivo Maintenance. *IUBMB life* 67: 129-138

Fol M, Chauhan A, Nair NK, Maloney E, Moomey M, Jagannath C, Madiraju MV, Rajagopalan M
(2006) Modulation of Mycobacterium tuberculosis proliferation by MtrA, an essential two-
component response regulator. *Molecular microbiology* 60: 643-657
Gazdik MA, Bai G, Wu Y, McDonough KA (2009) Rv1675c (cmr) regulates intramacrophage and cyclic
AMP-induced gene expression in Mycobacterium tuberculosis-complex mycobacteria. *Molecular*
microbiology 71: 434-448
Gazdik MA, McDonough KA (2005) Identification of cyclic AMP-regulated genes in Mycobacterium
tuberculosis complex bacteria under low-oxygen conditions. *Journal of bacteriology* 187: 2681-2692
Goar H, Paul P, Khan H, Sarkar D (2022) Molecular Connectivity between Extracytoplasmic Sigma
Factors and PhoP Accounts for Coupled Mycobacterial Stress Response. *J Bacteriol* 204: e0011022
Gonzalo Asensio J, Maia C, Ferrer NL, Barilone N, Laval F, Soto CY, Winter N, Daffe M, Gicquel B,
Martin C *et al* (2006a) The virulence-associated two-component PhoP-PhoR system controls the
biosynthesis of polyketide-derived lipids in Mycobacterium tuberculosis. *J Biol Chem* 281: 1313-1316
Gonzalo Asensio J, Maia C, Ferrer NL, Barilone N, Laval F, Soto CY, Winter N, Daffé M, Gicquel B,
Martín C *et al* (2006b) The virulence-associated two-component PhoP-PhoR system controls the
biosynthesis of polyketide-derived lipids in Mycobacterium tuberculosis. *J Biol Chem* 281: 1313-1316
Gouzy A, Healy C, Black KA, Rhee KY, Ehrt S (2021) Growth of Mycobacterium tuberculosis at acidic
pH depends on lipid assimilation and is accompanied by reduced GAPDH activity. *Proc Natl Acad Sci*
U S A 118
Goyal R, Das AK, Singh R, Singh PK, Korpole S, Sarkar D (2011) Phosphorylation of PhoP protein plays
direct regulatory role in lipid biosynthesis of Mycobacterium tuberculosis. *J Biol Chem* 286: 45197-
45208
Gupta S, Sinha A, Sarkar D (2006) Transcriptional autoregulation by Mycobacterium tuberculosis
PhoP involves recognition of novel direct repeat sequences in the regulatory region of the promoter.
FEBS letters 580: 5328-5338
Jacobs WR, Jr., Kalpana GV, Cirillo JD, Pascopella L, Snapper SB, Udani RA, Jones W, Barletta RG,
Bloom BR (1991) Genetic systems for mycobacteria. *Methods in enzymology* 204: 537-555
Johnson RM, Bai G, DeMott CM, Banavali NK, Montague CR, Moon C, Shekhtman A, VanderVen B,
McDonough KA (2017) Chemical activation of adenylyl cyclase Rv1625c inhibits growth of
Mycobacterium tuberculosis on cholesterol and modulates intramacrophage signaling. *Mol*
Microbiol 105: 294-308
Johnson RM, McDonough KA (2018) Cyclic nucleotide signaling in Mycobacterium tuberculosis: an
expanding repertoire. *Pathog Dis* 76
Keppetipola N, Shuman S (2008) A phosphate-binding histidine of binuclear
metallophosphodiesterase enzymes is a determinant of 2',3'-cyclic nucleotide phosphodiesterase
activity. *J Biol Chem* 283: 30942-30949
Khan H, Paul P, Sevalkar RR, Kachhap S, Singh B, Sarkar D (2022) Convergence of two global
regulators to coordinate expression of essential virulence determinants of Mycobacterium
tuberculosis. *Elife* 11
Knapp GS, Lyubetskaya A, Peterson MW, Gomes AL, Ma Z, Galagan JE, McDonough KA (2015) Role of
intragenic binding of cAMP responsive protein (CRP) in regulation of the succinate dehydrogenase
genes Rv0249c-Rv0247c in TB complex mycobacteria. *Nucleic Acids Res* 43: 5377-5393
Knapp GS, McDonough KA (2014) Cyclic AMP Signaling in Mycobacteria. *Microbiol Spectr* 2
Martin C, Williams A, Hernandez-Pando R, Cardona PJ, Gormley E, Bordat Y, Soto CY, Clark SO, Hatch
GJ, Aguilar D *et al* (2006) The live Mycobacterium tuberculosis phoP mutant strain is more
attenuated than BCG and confers protective immunity against tuberculosis in mice and guinea pigs.
Vaccine 24: 3408-3419
Matange N (2015) Revisiting bacterial cyclic nucleotide phosphodiesterases: cyclic AMP hydrolysis
and beyond. *FEMS Microbiol Lett* 362

Matange N, Hunt DM, Buxton RS, Visweswariah SS (2013) Overexpression of the Rv0805
phosphodiesterase elicits a cAMP-independent transcriptional response. *Tuberculosis (Edinb)* 93:
492-500
McCue LA, McDonough KA, Lawrence CE (2000) Functional classification of cNMP-binding proteins
and nucleotide cyclases with implications for novel regulatory pathways in Mycobacterium
tuberculosis. *Genome research* 10: 204-219
McDonough KA, Rodriguez A (2011) The myriad roles of cyclic AMP in microbial pathogens: from
signal to sword. *Nature reviews Microbiology* 10: 27-38
McDowell JR, Bai G, Lasek-Nesselquist E, Eisele LE, Wu Y, Hurteau G, Johnson R, Bai Y, Chen Y, Chan J
et al (2023) Mycobacterial phosphodiesterase Rv0805 is a virulence determinant and its cyclic
nucleotide hydrolytic activity is required for propionate detoxification. *Mol Microbiol* 119: 401-422
Nambi S, Gupta K, Bhattacharyya M, Ramakrishnan P, Ravikumar V, Siddiqui N, Thomas AT,
Visweswariah SS (2013) Cyclic AMP-dependent protein lysine acylation in mycobacteria regulates
fatty acid and propionate metabolism. *J Biol Chem* 288: 14114-14124
Nathan C, Shiloh MU (2000) Reactive oxygen and nitrogen intermediates in the relationship between
mammalian hosts and microbial pathogens. *Proceedings of the National Academy of Sciences of the
United States of America* 97: 8841-8848
Parikh A, Kumar D, Chawla Y, Kurthkoti K, Khan S, Varshney U, Nandicoori VK (2013) Development of
a new generation of vectors for gene expression, gene replacement, and protein-protein interaction
studies in mycobacteria. *Applied and environmental microbiology* 79: 1718-1729
Perez E, Samper S, Bordas Y, Guilhot C, Gicquel B, Martin C (2001) An essential role for phoP in
Mycobacterium tuberculosis virulence. *Molecular microbiology* 41: 179-187
Podobnik M, Tyagi R, Matange N, Dermol U, Gupta AK, Mattoo R, Seshadri K, Visweswariah SS (2009)
A mycobacterial cyclic AMP phosphodiesterase that moonlights as a modifier of cell wall
permeability. *J Biol Chem* 284: 32846-32857
Ranganathan S, Bai G, Lyubetskaya A, Knapp GS, Peterson MW, Gazdik M, AL CG, Galagan JE,
McDonough KA (2016) Characterization of a cAMP responsive transcription factor, Cmr (Rv1675c), in
TB complex mycobacteria reveals overlap with the DosR (DevR) dormancy regulon. *Nucleic Acids Res*
44: 134-151
Rickman L, Scott C, Hunt DM, Hutchinson T, Menendez MC, Whalan R, Hinds J, Colston MJ, Green J,
Buxton RS (2005) A member of the cAMP receptor protein family of transcription regulators in
Mycobacterium tuberculosis is required for virulence in mice and controls transcription of the rpfA
gene coding for a resuscitation promoting factor. *Mol Microbiol* 56: 1274-1286
Rittershaus ESC, Baek SH, Krieger IV, Nelson SJ, Cheng YS, Nambi S, Baker RE, Leszyk JD, Shaffer SA,
Sacchettini JC *et al* (2018) A Lysine Acetyltransferase Contributes to the Metabolic Adaptation to
Hypoxia in Mycobacterium tuberculosis. *Cell Chem Biol* 25: 1495-1505.e1493
Rohde KH, Abramovitch RB, Russell DG (2007) Mycobacterium tuberculosis invasion of
macrophages: linking bacterial gene expression to environmental cues. *Cell Host Microbe* 2: 352-364
Russell DG (2011) Mycobacterium tuberculosis and the intimate discourse of a chronic infection.
Immunol Rev 240: 252-268
Rustad TR, Sherrid AM, Minch KJ, Sherman DR (2009) Hypoxia: a window into Mycobacterium
tuberculosis latency. *Cell Microbiol* 11: 1151-1159
Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nature
protocols* 3: 1101-1108
Sevalkar RR, Arora D, Singh PR, Singh R, Nandicoori VK, Karthikeyan S, Sarkar D (2019) Functioning of
Mycobacterial Heat Shock Repressors Requires the Master Virulence Regulator PhoP. *Journal of
bacteriology* 201
Shenoy AR, Capuder M, Draskovic P, Lamba D, Visweswariah SS, Podobnik M (2007) Structural and
biochemical analysis of the Rv0805 cyclic nucleotide phosphodiesterase from Mycobacterium
tuberculosis. *Journal of molecular biology* 365: 211-225

Shenoy AR, Sreenath N, Podobnik M, Kovacevic M, Visweswariah SS (2005) The Rv0805 gene from
Mycobacterium tuberculosis encodes a 3',5'-cyclic nucleotide phosphodiesterase: biochemical and
mutational analysis. *Biochemistry* 44: 15695-15704
Shenoy AR, Visweswariah SS (2006) New messages from old messengers: cAMP and mycobacteria.
Trends in microbiology 14: 543-550
Singh AK, Carette X, Potluri LP, Sharp JD, Xu R, Prisic S, Husson RN (2016) Investigating essential gene
function in Mycobacterium tuberculosis using an efficient CRISPR interference system. *Nucleic Acids*
Res 44: e143
Singh R, Anil Kumar V, Das AK, Bansal R, Sarkar D (2014) A transcriptional co-repressor regulatory
circuit controlling the heat-shock response of Mycobacterium tuberculosis. *Mol Microbiol* 94: 450-
465
Smith LJ, Bochkareva A, Rolfe MD, Hunt DM, Kahramanoglou C, Braun Y, Rodgers A, Blockley A,
Coade S, Lougheed KEA *et al* (2017) Cmr is a redox-responsive regulator of DosR that contributes to
M. tuberculosis virulence. *Nucleic Acids Res* 45: 6600-6612
Solans L, Gonzalo-Asensio J, Sala C, Benjak A, Uplekar S, Rougemont J, Guilhot C, Malaga W, Martin
C, Cole ST (2014) The PhoP-dependent ncRNA Mcr7 modulates the TAT secretion system in
Mycobacterium tuberculosis. *PLoS Pathog* 10: e1004183
Studier FW, Moffatt BA (1986) Use of bacteriophage T7 RNA polymerase to direct selective high-
level expression of cloned genes. *Journal of molecular biology* 189: 113-130
Tan S, Sukumar N, Abramovitch RB, Parish T, Russell DG (2013) Mycobacterium tuberculosis
responds to chloride and pH as synergistic cues to the immune status of its host cell. *PLoS Pathog* 9:
e1003282
Thomson M, Liu Y, Nunta K, Cheyne A, Fernandes N, Williams R, Garza-Garcia A, Larrouy-Maumus G
(2022) Expression of a novel mycobacterial phosphodiesterase successfully lowers cAMP levels
resulting in reduced tolerance to cell wall-targeting antimicrobials. *J Biol Chem* 298: 102151
Valadares NF, Woo J (2017) Mechanism of Rv2837c from Mycobacterium tuberculosis remains
controversial. *J Biol Chem* 292: 13480
VanderVen BC, Fahey RJ, Lee W, Liu Y, Abramovitch RB, Memmott C, Crowe AM, Eltis LD, Perola E,
Deininger DD *et al* (2015) Novel inhibitors of cholesterol degradation in Mycobacterium tuberculosis
reveal how the bacterium's metabolism is constrained by the intracellular environment. *PLoS Pathog*
11: e1004679
Voskuil MI, Schnappinger D, Visconti KC, Harrell MI, Dolganov GM, Sherman DR, Schoolnik GK (2003)
Inhibition of respiration by nitric oxide induces a Mycobacterium tuberculosis dormancy program.
The Journal of experimental medicine 198: 705-713
Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I (2006) The Mycobacterium
tuberculosis PhoPR two-component system regulates genes essential for virulence and complex lipid
biosynthesis. *Mol Microbiol* 60: 312-330
Wayne LG, Sohaskey CD (2001) Nonreplicating persistence of mycobacterium tuberculosis. *Annual*
review of microbiology 55: 139-163
Wilburn KM, Montague CR, Qin B, Woods AK, Love MS, McNamara CW, Schultz PG, Southard TL,
Huang L, Petrassi HM *et al* (2022) Pharmacological and genetic activation of cAMP synthesis disrupts
cholesterol utilization in Mycobacterium tuberculosis. *PLoS Pathog* 18: e1009862
Wong AI, Beites T, Planck KA, Li S, Poulton NC, Rhee K, Schnappinger D, Ehrt S, Rock J (2022) Cyclic
AMP is a critical mediator of intrinsic drug resistance and fatty acid metabolism in M.
tuberculosis. *bioRxiv*: 2022.2007.2007.499113
Zaveri A, Bose A, Sharma S, Rajendran A, Biswas P, Shenoy AR, Visweswariah SS (2021)
Mycobacterial STAND adenylyl cyclases: The HTH domain binds DNA to form biocrystallized
nucleoids. *Biophys J* 120: 1231-1246

Figure Legends

Figure 1: PhoP contributes to maintenance of mycobacterial cAMP level. (A) Intra-mycobacterial

cAMP levels were determined by a fluorescence-based assay as described in the Methods, and

compared for indicated mycobacterial strains, grown under normal or specific stress conditions. For

acid stress, mycobacterial strains were initially grown to mid-log phase (OD_{600} 0.4 to 0.6), and then it

was transferred to acidic pH (7H9 media, pH 4.5) for further 2 hours of growth at 37°C. For NO

stress, cells grown to mid-log phase were exposed to 0.5 mM DataNonoate for 40 minutes. The data

represent average values from three biological repeats (*$P \leq 0.05$, **$P \leq 0.01$). (B) To compare secretion

of cAMP by WT and *phoPR*-KO, cAMP levels were also determined in the corresponding culture

filtrates (CF). (C) Immunoblotting analysis of 10 μg of cell lysates (CL) and 20 μg of culture filtrates

(CF) of indicated *M. tuberculosis* strains. α-GroEL2 was used as a control to verify cytolysis of cells,

CFP-10 detected as a secreted mycobacterial protein in the culture filtrates, and RpoB used as the

loading control.

Figure 2: PhoP regulates expression of phosphodiesterase (PDE) *rv0805*. (A) To investigate

regulation of cAMP level, mRNA levels of well-characterized adenylate cyclases, and PDEs were

compared in indicated mycobacterial strains by RT-qPCR as described in the Methods. The results

show average values from biological triplicates, each with two technical repeats. Note that the

difference in expression levels of *rv0805* between WT and *phoPR*-KO was significant ($P < 0.01$),

whereas the mRNA level was insignificantly different in WT and the complemented mutant (Compl.).

Nonsignificant difference is not indicated. (B) To determine the effect of acidic pH conditions of

growth, mycobacterial *rv0805* expression was compared in WT grown under normal (pH 7.0) and

acidic pH (pH 4.5). Average fold difference in mRNA levels from biological duplicates (each with a

technical repeat) were measured as described in the Methods (**$P \leq 0.01$). As controls, expression of

rv1339 and *16S rDNA* were also measured. Non-significant difference is not indicated. (C) *In vivo*

PhoP binding to *rv0805* promoter (rv0805up) was compared in WT grown under normal and acidic

conditions of growth using anti-PhoP antibody followed by ChIP-qPCR. Fold enrichment data

represent mean values of two independent experiments with a statistically significant fold difference

(**P-value <0.001; *P-value <0.01). The upstream regulatory regions of 16S rRNA (16S rRNAup)

and msl3 (msl3up) were used as negative and positive controls, respectively. The assay conditions,

sample analyses, and detection are described in the Methods section.

Figure 3. PhoP dependent repression of *rv0805* to maintain mycobacterial cAMP homeostasis

requires the presence of PhoR. (A-B) To determine the impact of PhoR (the cognate sensor kinase

of PhoP), expression of *rv0805* was compared in indicated *M. tuberculosis* H37Rv strains: (A)

phoPR-KO*::phoP* (*phoPR* mutant complemented with *phoP*) and (B) *phoPR*-KO*::phoPR* (*phoPR*

mutant complemented with *phoP-phoR*) under normal and acidic conditions of growth. As expected,

phoPR-KO*::phoPR* (Compl.) shows a significant repression of *rv0805* (but not *rv1339*) under acidic

pH (***P<0.001). However, *rv0805* expression remains comparable in *phoPR*-KO*::phoP* under

normal as well as acidic conditions of growth. As a control, *rv1339* fails to show a variable expression

in indicated mycobacterial strains. (C) To determine the effect of ectopic expression of *rv0805* on

intra-mycobacterial cAMP level, WT and mutant Rv0805 proteins (Rv0805M, defective for

phosphodiesterase activity) were expressed in *M. tuberculosis* H37Rv (to construct WT-Rv0805, and

WT-Rv0805M, respectively) as described in the Methods section. Notably, similar to *phoPR*-KO,

WT-Rv0805 (but not WT-Rv0805M) showed a considerably lower level of cAMP relative to WT

bacteria. Significance in variation of cAMP levels were determined by paired student's t-test

(**P<0.01). (D-E) Relative expression of *phoP* and PDE in *phoP*-KD and *rv0805*-KD (*phoP* and

rv0805 knock-down constructs, respectively). In keeping with elevated expression of *rv0805* in

phoPR-KO, *phoP*-KD shows an elevated expression of *rv0805* relative to WT bacilli. In contrast,

phoP expression level remains unaffected in *rv0805*-KD mutant. Panel E measured corresponding

intra-bacterial cAMP levels in the respective knock-down mutants, as described in the legend to Fig.

1A.

Figure 4: cAMP homeostasis and its effect on mycobacterial stress tolerance and survival in

macrophages. (A) To compare susceptibility to low pH conditions, indicated mycobacterial strains

were grown at pH 4.5. Importantly, similar to *phoPR*-KO (grey circles), WT-Rv0805 (red circles)

shows a significant growth defect relative to WT (empty circles). However, WT-Rv0805M (green

circles) grows comparably well as that of the WT (empty circles). In contrast, at pH 7.0 all four

mycobacterial strains (WT, empty triangles; *phoPR*-KO, grey triangles; WT-Rv0805, red triangles;

WT-Rv0805M, green triangles) displayed comparable growth. (B) Microplate-based assay using

Alamar Blue was utilized to examine mycobacterial sensitivity to increasing concentrations of

diamide. In this assay, reduction of Alamar Blue correlates with the change of a non-fluorescent blue

to a fluorescent pink appearance, which is directly proportional to bacterial growth. Survival of

indicated mycobacterial strains, under normal conditions and in presence of 5 mM diamide, were

determined by plotting fluorescence intensity. The data is normalized relative to WT grown in

presence of 5 mM diamide. (C) To compare susceptibility to stress conditions, these mycobacterial

strains were next grown in the presence of 50 μM Cumene Hydrogen Peroxide (CHP). In presence of

CHP, WT-Rv0805 (red column) but not WT-Rv0805M (green column), shows a significant growth

defect [relative to WT (empty column)] in striking similarity to *phoPR*-KO (grey column). The

growth experiments were performed in biological duplicates, each with two technical replicates

(**$P \leq 0.01$; ***$P \leq 0.001$). (D) Murine macrophages were infected with indicated *M. tuberculosis*

H37Rv strains. The cells were made visible by LysoTracker; mycobacterial strains were stained with

phenolic auramine solution, and the confocal images display merge of two fluorescence signals (Lyso

Tracker: red; H37Rv: green; scale bar: 10 μm). (E) Bacterial co-localization of indicated *M.*

tuberculosis H37Rv strains. The percentage of auramine labelled strains co-localized with

Lysotracker was determined by counting at least 100 infected cells in 10 different fields. The results

show the average values with standard deviation determined from three independent experiments

(***$P \leq 0.001$). (F) Pearson's correlation coefficient of images of internalized auramine-labelled

mycobacteria and Lysotracker red marker in RAW 264.7 macrophages. Data are representative of

mean ± S.D., derived from three independent experiments (*$P < 0.05$; ***$P < 0.001$).

Figure 5: Dysregulation of mycobacterial cAMP homeostasis impacts mycobacterial survival *in*

***vivo*.** (A-B) Survival of mycobacterial strains in mice (A) lung, and (B) spleen after animals were

given an aerosol infection with ~100 CFU / lung. Mycobacterial load represents mean CFU values

with standard deviations obtained from at least five animals per strains used (**P< 0.01;

***P<0.001). (C) Histopathology of lung sections after 4 weeks of infection with indicated bacterial

strains. Sections were stained with hematoxylin and eosin, observed under a light microscope, and

images of the pathology sections collected at x40 magnification display granulomas (filled arrows)

and alveolar space (empty arrows) (scale bar, 200 μm).

Figure 6: Schematic summary of cAMP homeostasis and its impact on mycobacterial stress

tolerance. According to this model, upon activation by an appropriate signal via the cognate sensor

PhoR, P~PhoP binds to *rv0805* regulatory region and functions as a specific repressor by preventing

access for mycobacterial RNA polymerase (RNAP) to bind to the promoter and initiate transcription.

In keeping with the PhoP-dependent *rv0805* repression, our results demonstrate a reproducibly lower

level of cAMP in *phoPR*-KO relative to WT bacilli. Thus, *phoPR*-KO (or WT-Rv0805) remains

ineffective to mount an appropriate stress response most likely due to its inability to coordinate

regulated gene expression because of dysregulation of cAMP homeostasis, provides an explanation to

their reduced stress tolerance. Together, these molecular events suggest that failure to maintain cAMP

homeostasis most likely accounts for attenuated phenotype of the bacilli.

Legends to supplementary figures

Figure 1-figure supplement 1: Defined and indicated stress conditions do not influence *in vitro*
growth of mycobacterial strains. To determine the impact of *in vitro* stress on bacterial growth
under conditions when intra-mycobacterial cAMP levels were measured, WT bacilli, *phoPR*-KO, and
the complemented mutant (Compl.) were exposed to carefully controlled stress conditions as detailed
in the Methods, and CFUs enumerated. The results clearly suggest that bacterial viability under
indicated conditions of stress do not impact intra-mycobacterial cAMP level.

Figure 3-figure supplement 1: Ectopic expression of WT and mutant *rv0805* (Rv0805M) in WT
bacilli. (A) Real-time RT-qPCR was carried out to compare expression of indicated
phosphodiesterases (PDEs) in WT (empty bar), WT-Rv0805 (red bar) and WT- Rv0805M (green bar),
respectively, as described in the Methods. Average fold difference in mRNA levels from two
biological repeats (each with a technical repeat) were determined as described in the Methods
(**P≤0.01; ***P≤0.001). Nonsignificant difference is not indicated. **(B)** To compare growth of WT
and the variant mycobacterial strains, CFU values were determined under normal conditions of
growth as described in the methods.

Figure 4-figure supplement 1: PhoP-dependent *rv0805* expression contributes to mycobacterial
survival under oxidative stress. (A) In this experiment, we compared metabolic activity of WT,
phoPR-KO, WT-Rv0805, and WT-Rv0805M, grown in presence of increasing concentrations of
diamide by using Alamar Blue assay. Because reduction of Alamar Blue correlates with the change of
a non-fluorescent blue to a fluorescent pink appearance, mycobacterial metabolic activity could be
assessed by monitoring fluorescence. Two following controls were included in the Alamar Blue
assays: Rifampicin control, reflecting mycobacterial growth inhibition to confirm validity of the
assay, and no bacteria control indicated at the left top corner of the plate. **(B)** Next, effect of over
expression of *rv0805* on mycobacterial cell -wall structure was assessed by analysing expression of
lipid biosynthetic genes of indicated mycobacterial strains by RT-qPCR. Average fold difference in
mRNA levels from two biological repeats (each with a technical repeat) were determined as described
in the Methods (***P≤0.001). Nonsignificant difference is not indicated.
970

972 **Legends to supplementary Tables**

973 Supplementary file 1a

974 Sequences of oligonucleotide primers used in RT-qPCR and ChIP-qPCR measurements
975 reported in this study
976
977 Supplementary file 1b
978
979 Sequences of oligonucleotide primers for amplification and cloning, and plasmids used in this study
980
981
982

Source data files for supplementary figures

Figure 1-figure supplement 1

Figure 3-figure supplement 1

Figure 4-figure supplement 1

Author contributions

H.K., P.P., H. G., B.B. and D.S. designed research; H.K., P. P., H.G., and B.B. performed research;

N.B. contributed analytical tools; H.K., P. P., H. G., B.B., N.B., and D.S. analysed data; and D.S.

wrote the manuscript.

Supplemental Data

The supplemental data include three supplemental figures and two supplemental tables

(supplementary files 1a and 1b, respectively). All other data are part of the paper and its supplemental

files.

1001

1002

1003

1004

1005

1006

1007 **Figure 1**

1008

1009

1010

1011

1012

1013

1014

1015 **Figure 2**

1016

1017

1018

1019

1020

1021

1022

1023

1024 **Figure 3**

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038   **Figure 4**

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053   **Figure 5**

1054

1055

1056

1057   **Figure 6**

1058

1059

1060

1061