Excitatory and inhibitory D-serine binding to the NMDA receptor

Remy A. Yovanno[1], Tsung Han Chou[2], Sarah J. Brantley[3], Hiro Furukawa[2*], and Albert Y. Lau[1*]

Affiliations:

[1]Department of Biophysics and Biophysical Chemistry, Johns Hopkins University School of Medicine, 725 N. Wolfe Street, WBSB 706, Baltimore, MD 21205, USA.

[2] W.M. Keck Structural Biology Laboratory, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA.

[3]Department of Biology, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

* To whom correspondence should be addressed:

alau@jhmi.edu (A.Y.L.)

furukawa@cshl.edu (H.F.)

ABSTRACT

N-methyl-D-aspartate receptors (NMDARs) uniquely require binding of two different

neurotransmitter agonists for synaptic transmission. D-serine and glycine bind to one

subunit, GluN1, while glutamate binds to the other, GluN2. These agonists bind to the

receptor's bi-lobed ligand-binding domains (LBDs), which close around the agonist

during receptor activation. To better understand the unexplored mechanisms by which

D-serine contributes to receptor activation, we performed multi-microsecond molecular

dynamics simulations of the GluN1/GluN2A LBD dimer with free D-serine and glutamate

agonists. Surprisingly, we observed D-serine binding to both GluN1 and GluN2A LBDs,

suggesting that D-serine competes with glutamate for binding to GluN2A. This

mechanism is confirmed by our electrophysiology experiments, which show that D-

serine is indeed inhibitory at high concentrations. Although free energy calculations

indicate that D-serine stabilizes the closed GluN2A LBD, its inhibitory behavior suggests

that it either does not remain bound long enough or does not generate sufficient force

for ion channel gating. We developed a workflow using pathway similarity analysis to

identify groups of residues working together to promote binding. These conformation-

dependent pathways were not significantly impacted by the presence of N-linked

glycans, which act primarily by interacting with the LBD bottom lobe to stabilize the

closed LBD.

INTRODUCTION

The N-methyl-D-aspartate receptor (NMDAR) is an ionotropic glutamate receptor

(iGluR) that uniquely requires the binding of a co-agonist in addition to its primary

agonist for activation. This heterotetrameric ion channel comprises at least two different

subunits, GluN1 (isoforms 1-4a and 1-4b) and GluN2 (subtypes A-D), assembled as a

dimer of GluN1/GluN2 heterodimers. The GluN2 subunit binds the neurotransmitter

glutamate, while the GluN1 subunit can either bind the co-agonists glycine or D-serine.

Traditionally, glycine had been considered the dominant GluN1 agonist [1–3], but more

recent work has suggested that D-serine may in fact be the dominant co-agonist for

synaptic NMDARs in the brain [4]. D-serine is synthesized by the enzyme serine

racemase expressed in astroglia [5] and neurons [6] [7] and is released into the

postsynapse by the Asc-1 transporter [8] [9]. D-serine binding to these synaptic

NMDARs is responsible for inducing long-term potentiation (LTP), which is critical for

memory functions [10]. In addition, recent clinical efforts have indicated that D-serine

could be a promising therapeutic for the treatment of neuropsychiatric disorders

[11][12], most notably schizophrenia [13] and post-traumatic stress disorder (PTSD)

[14]. Unlike the more well-studied agonists glutamate and glycine, the role of D-serine is

less defined, causing it to be known as the "shape-shifting" agonist [9] that can adopt

different roles in neurotransmission.

60   Each NMDAR subunit consists of an amino-terminal domain (ATD), a ligand-

binding domain (LBD; also called an agonist-binding domain, ABD), a transmembrane

domain (TMD), and a disordered cytoplasmic C-terminal domain [15]. The LBDs adopt a

bi-lobed clamshell architecture that close upon agonist binding. Previous computational

studies of NMDAR LBDs have indicated that glycine binding to the GluN1 LBD and

glutamate binding to the GluN2A LBD drives the conformational equilibrium toward the

closed LBD [16]. While crystallographic studies have determined the binding pose of D-

serine bound to the closed GluN1 LBD [17], the molecular mechanisms by which D-

serine finds its way into and stabilizes NMDAR LBDs is not well understood.

Previous simulation studies have revealed the mechanisms by which glycine and

glutamate diffuse into the LBD binding site [18]. Specifically, they found that glycine

binds to the GluN1 subunit by freely diffusing into the binding pocket, where it is trapped

by energetically favorable interactions with key binding site residues. Glutamate, on the

other hand, was found to contact residues along the protein surface that helped guide

itself into its binding pocket, positioning it to interact stably with residues in the binding

site. These two binding mechanisms were referred to as "unguided" and "guided"

diffusion, respectively [19]. This paradigm established the two extremes by which

ligands enter their receptor sites: one in which stable ligand binding only depends upon

the identity of the binding site residues and another that also heavily relies on residues

outside the binding site to guide the ligand toward its bound pose.

Performing multi-microsecond molecular dynamics simulations of the

glycosylated GluN1/GluN2A LBD dimer, we identified binding mechanisms and residues

critical for promoting D-serine binding and stabilization by developing a new binding

pathway clustering workflow. Surprisingly, we observed D-serine binding to both GluN1

and GluN2A LBDs. We determined that D-serine binding to GluN2A partially stabilizes

the active LBD conformation. Inspired by these simulation results, we determined that

D-serine competes with glutamate for binding to GluN2A via a competitive inhibition

mechanism using electrophysiology measurements, where D-serine was found to be

inhibitory at high concentrations. Since NMDAR LBDs are glycosylated under

physiological conditions [20], including N-linked glycans in our simulations revealed that

glycans primarily regulate the binding process by stabilizing the active LBD. In total, we

investigated the molecular components contributing to D-serine binding and

stabilization, highlighting the complex components driving neurotransmission.

RESULTS

D-serine binding pathways for GluN2A and GluN1 LBDs

In simulating the GluN1/GluN2A LBD dimer, which is a physiological NMDAR

unit, we intended to focus our attention on the mechanisms by which D-serine binds to

the GluN1 LBD, the subunit to which D-serine is a potent agonist. However, in our

simulations, we also observed a significant number of D-serine binding events involving

the GluN2A LBD, an unexpected finding. These binding events are primarily made up of

guided-diffusion pathways in which D-serine contacts key residues on the LBD surface

to help guide it into the binding cleft. In our aggregate ~51 μs of sampling of the

glycosylated GluN1/GluN2A LBD dimer, we identified 99 guided-diffusion pathways for

GluN2A and 104 (plus 23 free diffusion events) for GluN1. Due to the stochastic nature

of these pathways, we needed to develop a reliable way to identify key features of

predominant binding pathways. To address this, we applied pathway similarity analysis

(PSA) [21] to quantify the spatial and geometric similarity between pairs of paths **(Fig.**

1A). Here, we extend this application to ligand binding pathways by monitoring the

change in ligand C_α position throughout each path. This allowed us to cluster paths

traversing similar regions of the LBD surface. To aid in describing the different faces of

the LBD, we use an order parameter (ξ_1, ξ_2) defined in previous work [16] to describe

whether D-serine primarily contacts residues on the ξ_1 or ξ_2 face of the LBD **(Fig. 1B,**

2A). For GluN2A, cluster analysis revealed four distinct regions of D-serine occupancy.

The clusters correspond to the following methods of binding: 1. D-serine approaches

the binding pocket from the ξ_2 face; 2. D-serine contacts the D1 residues on the ξ_1 face;

3. D-serine zigzags between D1 and D2 lobes on the ξ_1 face; 4. D-serine primarily

contacts residues on the D2 lobe of the ξ_1 face **(Fig. 1C-F)**. Similarly, for GluN1, cluster

analysis revealed four distinct clusters corresponding to similar pathways of binding: 1.

D-serine contacts the ξ_2 face; 2. D-serine zigzags between D1 and D2 lobes on the ξ_1

face; 3. D-serine contacts residues on the N-terminal (top) end of D1 of the ξ_1 face; 4.

D-serine contacts residues of D1 loop 2 that protrudes from the LBD into solution. We

then analyzed the resulting clusters to identify key residues that guide D-serine into the

binding site **(Fig. 2B-E)**. Interestingly, we observed that GluN1 pathways involve fewer

interactions between D-serine and D2 residues; most notably, there were fewer

contacts with Helix F (Helix E for GluN2A) compared to GluN2A pathways.

To quantify the extent to which these clusters involve similar residue contacts, we

used a pairwise similarity metric called the overlap coefficient (i.e., Szymkiewicz–

Simpson coefficient) that describes agreement between sets of residues [22]. Doing so

provides a way to determine whether these spatial clusters are mostly made up of

random contacts, or whether groups of residues tend to act together to promote binding,

allowing us to quantify the extent to which agonist diffusion is "guided" by contacts along

the LBD. For GluN2A, we computed the overlap coefficient for all path pairs in each

cluster for comparison with the global mean (global $\langle OC \rangle$ = 0.557) **(Fig. S1A)**. We found

that pathway pairs in three of the four clusters yielded an overlap coefficient greater

than the mean of all pairs of paths from all clusters, indicating that pathways in each

cluster are made up of specific residue contacts **(Fig. S1C)**. In contrast, for GluN1, a

significant cluster (26 paths) involving interactions with residues on the ξ_2 face of the

LBD has a cluster mean OC much less than the global mean (global $\langle OC \rangle$ = 0.671),

indicating that this cluster primarily comprises random contacts **(Fig. 1B, S1B,D)**. This

suggests that D-serine binding to GluN1 may be more diffusion-driven and less guided

than to GluN2A. Therefore, we propose that agonist binding mechanisms exist on a

spectrum ranging from unguided to guided diffusion. The difference in the specificity of

D-serine contacts along binding pathways for GluN2A and GluN1 suggests that the

extent to which agonists rely on pathways of guiding residues depends on LBD

architecture and not solely upon the identity of the agonist.

Mapping important pathway residues onto the intact GluN1/GluN2A NMDAR

(PDB ID: 6MMM [23]) further enriches our understanding of binding pathways by

allowing us to determine whether residues in particular pathways are accessible for

binding or obscured by other receptor domains and subunits. For GluN2A, access to

residues on the extreme of the ξ_2 face is slightly restricted by the presence of the GluN1

subunit of the adjacent LBD dimer **(Fig. S2A)**. However, this interface does not seem to

be near the specific residues identified as critical for binding. Even more restricted is

access to residues on the ξ_1 face of GluN1, which are obscured by GluN2A of the

adjacent LBD dimer, including residues identified as critical for binding pathways **(Fig.**

S2B). This might bias the pathways observed for the intact receptor by forcing the

agonist to favor residues on the ξ_2 face of the LBD. Since our overlap coefficient

analysis of the cluster that corresponds to the ξ_2 face of GluN1 identified more non-

specific interactions, it is possible that the D-serine mechanism would be biased to favor

unguided diffusion. It is also possible that access to residues near the N-terminal end of

D1 would be restricted by the R2 lobe of its own ATD.

We next investigated whether a specific LBD conformational state was favored

for successful D-serine binding pathways. We computed our (ξ_1, ξ_2) order parameter to

quantify the degree of closure of the LBDs for all trajectory frames identified as part of

binding (and unbinding) pathways and found that (ξ_1, ξ_2) = (16,14) for GluN2A **(Fig.**

S3A) and (ξ_1, ξ_2) = (11,13) for GluN1 **(Fig. S3B)**. These values correspond to a partially

open LBD. The LBD needs to be open enough for the ligand to diffuse into the pocket

but closed enough to form some stabilizing interactions with the ligand. However, we

notice that the ξ_1 is smaller for GluN1, indicating that agonist binding can occur at

slightly more closed LBD conformations. GluN1 pathways where (ξ_1, ξ_2) = (11,13) are

mostly in the cluster defined by D-serine interactions with Loop 2, highlighting the role of

Loop 2 residues in D-serine binding to GluN1. Overall, these results suggest that the

degree of LBD closure does influence the likelihood of successful binding.

Effects of D-serine binding on the LBD conformational free energy landscapes

Since we did not expect to see D-serine binding to the GluN2A LBD, we needed

to determine whether these GluN2A D-serine binding events are able to modulate the

GluN2A LBD conformation. Since full LBD closure occurs on multi-microsecond to

millisecond timescales [24][25][26], direct observation of such a conformational change

was not fully captured from our equilibrium binding trajectories. Instead, to ensure we

are sampling the full range of LBD conformations, we performed umbrella sampling free

energy molecular dynamics simulations to obtain the conformational free energy

landscape of GluN2A bound to D-serine **(Fig. 3A)**. We used the order parameter (ξ_1, ξ_2)

[16] that captures the opening and closing motion of the LBDs observed in crystal

structures of these domains. Since no crystal structure exists for D-serine bound to

GluN2A, we identified residues critical for stabilizing the agonist in the closed state by

analyzing contacts in lowest-energy ($\leq$1 kcal mol^{-1}) conformers **(Fig. S4A)**. For

reference, we compared the resulting energy landscape to those previously computed

for the apo- and glutamate-bound GluN2A monomers **(Fig. 3C,D)** [16]. We see that, like

glutamate, D-serine stabilizes the closed LBD conformation. The D-serine energy

landscape has a global minimum corresponding to (ξ_1, ξ_2) values of (11, 11.5 Å) and a

metastable minimum corresponding to (ξ_1, ξ_2) values of (15.5, 11.5 Å). The presence of

a metastable agonist-bound LBD partially open intermediate suggests that D-serine

may not stabilize the closed conformation to the same extent as glutamate and

generate sufficient force to control channel gating. We then compared different

conformers corresponding to these two states to determine residues critical for agonist

stabilization. The primary difference between the residue contacts in conformers of the

two states is the prevalence of interactions with Thr-690 **(Fig. S4B)**, which only contacts

D-serine in the more closed state centered at (ξ_1, ξ_2) = (11, 11.5 Å). This is supported by

our binding simulations; although we do not fully sample LBD closure, trajectory frames

with low (ξ_1, ξ_2) values involve contacts with Thr-690. This suggests that Thr-690 is

critically involved in promoting full GluN2A LBD closure upon agonist binding.

Experimental binding studies have indicated that D-serine may be a more potent

GluN1 agonist than glycine [27]. To better understand the molecular mechanism

responsible for this difference in agonist potency, we computed the conformational free

energy for the D-serine-bound GluN1 LBD **(Fig. 3F)**. Compared with the previously

computed glycine-bound and apo LBDs **(Fig. 3G,H)** [16], the presence of D-serine in

the binding cleft results in a greater population of conformers in the closed conformation

and fewer conformers adopting a more open conformation. Similar to GluN2A Thr-690,

GluN1 Asp-732 and (to a lesser extent) Ser-688 help stabilize D-serine in the closed

LBD conformation by interacting with the D-serine hydroxyl. For this reason, we propose

that D-serine's high potency is due, at least in part, to its ability to more strongly

stabilize a closed LBD through additional interactions with the D2 lobe.

D-serine and glutamate compete for binding to the GluN2A LBD

Since our simulations revealed that D-serine can enter the GluN2A LBD binding

pocket and partially stabilize the active conformation, we hypothesized that D-serine

might compete with glutamate for binding to GluN2A. In fact, we observed D-serine

binding to GluN2A, even in the presence of glutamate, although glutamate bound more

frequently than D-serine and with longer residence times in the binding site **(Datasets**

S2, S3). Since increasing the D-serine concentration would increase the frequency of D-

serine binding to GluN2A, it is possible that D-serine could function as an inhibitor

(competitive antagonist) at high concentrations. If true, this behavior may factor into

therapeutic strategies focused on increasing D-serine concentration in the synapse by

establishing an upper dosage limit after which a D-serine increase is no longer

potentiating.

To probe this behavior experimentally, we measured GluN1-2A NMDAR currents

using two-electrode voltage clamp (TEVC) electrophysiology. We observed that at high

(~1 mM) D-serine concentrations, NMDAR activity was inhibited **(Fig. 4A)**. The inhibition

was dependent on glutamate concentrations, implying that the inhibitory effect of D-

serine may be competitive **(Fig. 4B)**. Furthermore, dose-response curves of glutamate

activation were right-shifted in the presence of increasing concentrations of D-serine

(Fig. 4C). The calculated slope value of the Schild plot at 1.1 ± 0.1 implied that D-serine

and glutamate likely compete against each other **(Fig. 4C)**. Combined with our

simulation results, our electrophysiological data supports the hypothesis that D-serine at

high concentrations can bind to the GluN2A subunit and compete against glutamate.

Since a similar inhibitory effect was also observed at high glycine concentrations

by TEVC electrophysiology **(Fig. 4A)**, we repeated our umbrella sampling simulations

with glycine bound to the GluN2A LBD. We see that glycine also favors the closed LBD

(Fig. 3B). The lowest-energy conformers of GluN2A with glycine are fastened shut by

contacts between the N-terminal amine of glycine and Tyr-730. Although glutamate still

stabilizes the closed GluN2A LBD to the greatest extent, comparable thermodynamics

between different agonists suggests that kinetics of agonist binding and unbinding is a

critical driver of agonist-induced activation. The GluN2A LBD likely never closes around

glycine because glycine does not remain bound long enough to induce LBD closure.

Previous binding studies [18] have indicated that glutamate, the primary GluN2A

agonist, similarly relies on LBD surface residues to promote binding. To determine

whether D-serine and glutamate binding are guided by similar residue contacts, we

computed the overlap coefficient between residues in D-serine and glutamate pathways

to be 0.964 for the glycosylated GluN2A LBD, corresponding to a significant overlap in

agonist occupancy **(Fig. 5A)**. This high degree of overlap between glutamate and D-

serine pathway residues indicates that they bind through similar mechanisms.

Despite similar pathway residues, we identified key residues that distinguish

glutamate from D-serine binding pathways **(Fig. 5B and Dataset S7)**. Most of the

residues important for D-serine binding, but not for glutamate binding, are located on

the ξ_2 face of the LBD. Most notably, Glu-413, Tyr-730, Ser-511, and Asp-731 all occur

in D-serine binding pathways with a frequency of more than ten times their fractional

occurrence in glutamate binding pathways. It is important to note, however, that

glutamate does interact with residues on the ξ_2 face, but the specific nature of those

contacts differ between the two agonists. In contrast, we found that Lys-487 is

contacted with significantly greater frequency in glutamate binding pathways. Due to

these residues' close proximity to the binding cleft, it is likely that these residues are

responsible for facilitating proper positioning of the agonists in the binding site, based

on differences in agonist size and shape.

An important feature of glutamate binding to GluN2A is its ability to bind in an

inverted pose relative to the crystal structure, which we observed in previous

simulations [18] [19]. Since no experimental structure exists for glutamate bound in the

inverted pose, we performed umbrella sampling simulations to determine the free

energy landscape of the GluN2A LBD with glutamate bound in the inverted pose **(Fig.**

3E). We found that glutamate bound in the inverted pose prevents full LBD closure as

predicted in previous work [18]. Specifically, glutamate in the inverted pose stabilizes a

conformation centered around (ξ_1, ξ_2) values of (14, 13 Å). Comparing the low-energy

conformers of D-serine and inverted glutamate ($\leq$ 1 kcal mol^{-1}) with the glutamate-

bound crystal structure, we found that D-serine and glutamate are stabilized by the

same residues, although there are fewer interactions between Thr-690 and glutamate in

the inverted pose, further supporting the importance of this residue for stabilizing the

fully closed LBD.

Kinetic analysis of D-serine binding pathways

We computed the D-serine association rate constant (k_{on}) for GluN2A and GluN1

LBDs using a method described [28] and used in previous iGluR work [19] as

summarized in the equation below:

282
$$k_{on} = \frac{N_b}{\sum_i \frac{t_i [L_i]}{s_i}}.$$

Here, N_b is the number of association events, t_i is the time the agonist spends in bulk

solvent, s_i is the number of identical binding sites, and $[L_i]$ is the concentration of free

agonist. One advantage of this approach is the ability to combine simulations performed

at various concentrations of free agonist $[L_i]$. Here, k_{on} is a bulk property and relies on

fully sampling the LBD conformational landscape throughout the simulation. However,

our binding simulations fail to adequately sample the agonist-bound, closed LBD state.

This affects both the number of observed binding events N_b and the time the agonist

spends in bulk solvent (t_i). Since this value is most sensitive to the number of identified

binding events N_b, we computed the k_{on} for different N_b values based on the duration of

the resulting binding event. This minimizes contributions from extremely short binding

events that are unlikely to be functionally relevant. For GluN2A, this results in a D-serine

k_{on} with an upper bound of 7.8×10^7 M^{-1}s^{-1} (all events included) and a lower bound of

1.6×10^7 M^{-1}s^{-1} (only events with agonist residence times > 100 ns were included). For

GluN1, the upper bound for k_{on} is 9.0×10^7 M^{-1}s^{-1} and the lower bound is 7.0×10^6 M^{-}

^{1}s^{-1}. Based on these values, it is reasonable to expect that D-serine binds to GluN2A

and GluN1 at similar rates. For comparison, the association rate constants computed for

glutamate binding to GluN2A with this method range from 4.9×10^7 M^{-1}s^{-1} to 1.4×10^8

M^{-1}s^{-1}. Similar ranges of D-serine binding rate constants for GluN2A and GluN1 support

our data indicating a guided-diffusion mechanism. However, this definition of the

association rate constant does not capture the molecular details that produce this bulk

behavior.

For agonist binding mechanisms dominated by guided diffusion, we can monitor

how much time the agonist spends (1) in bulk solvent, (2) associated with the LBDs,

and (3) docked in the binding cleft (interacting with the conserved arginines Arg-523 for

GluN1 or Arg-518 for GluN2A). Transitions between these states can be represented by

the following three-step process:

$$P + L \rightleftharpoons PL_{\mathrm{assoc}} \rightleftharpoons PL_{\mathrm{docked}} \, .$$

Here, the PL_{assoc} state either results in successful binding (represented by pathways) or

nonspecific interactions resulting in dissociation. From the clusters of residues that we

identified in our pathway similarity analysis, we determined to what extent a particular

residue is critical for guiding the agonist into the binding site using a conditional

probability-based framework **(Datasets S9, S10)**. For GluN2A, given that a binding

event results in successful agonist docking, residues Asp-515, Glu-517, Arg-692, Asn-

687, Lys-487, Lys-484, and Ser-689, Lys-488, Ser-511, and Glu-413 are contacted

most frequently across all datasets. Given successful D-serine binding, contacts with

GluN1 residues Lys-496, Lys-495, Trp-498, Arg-489, and Glu-497 occur in the greatest

number of pathways. Slightly less agreement in crucial GluN1 binding residues across

datasets further supports a more diffusive/random binding mechanism for D-serine

binding to GluN1.

Role of N-linked glycans in D-serine binding pathways

In addition to identifying residues that are responsible for agonist specificity in

binding pathways, we also explored the effect of the N-linked $Man_5GlcNAc_2$ (Man5)

glycans **(Fig. S6A)** on the residues involved in agonist binding pathways. Previous

electrophysiological studies have indicated that glycans function as LBD potentiators

[29]. In our simulations, we observed that near-pocket glycans appear to "reach" into the

binding pocket. This reaching behavior was observed in previous simulations of the

glycosylated NMDAR LBDs in which the glycan forms a "cage" around the binding

pocket by forming interactions with the LBD D2 lobe and is believed to be associated

with NMDAR potentiation by glycans [29]. For GluN2A, there are two glycans that are

near the binding pocket: N443-Man5 and N444-Man5, both of which can interact with

the LBD D2 lobe **(Fig. 6A)**. For GluN1, there is a single glycan N491-Man5 that adopts

this caged conformation **(Fig. 6B)**. To quantify this behavior in our simulations, we

developed a general order parameter to describe the relationship between the glycan

and the LBD D2 lobe that measured the minimum distance between any glycan heavy

atom and any residue on the LBD D2 lobe. From this order parameter, we computed

glycan PMFs along the glycan-D2 order parameter for each near-pocket glycan **(Fig.**

6C-E).

We compared our glycosylated trajectories with an additional 30 μs of simulations

of the non-glycosylated GluN1/GluN2A LBD dimer to identify ways in which the

presence of glycans influence binding pathways. Our data indicate that residues on the

ξ_2 face are contacted more frequently in non-glycosylated simulations, although these

residues are important for D-serine binding with and without glycans **(Dataset S11)**.

GluN2A residues Asp-515 and Glu-517, are contacted more frequently in glycosylated

systems. The frequency with which D-serine interacts with GluN1 residue Arg-489 in

pathways is greater for glycosylated pathways than those without glycans. On average,

glycan-mediated D-serine interactions result in slightly longer pathways, suggesting that

the presence of glycans slows down the binding process, setting up small kinetic

"traps".

When we analyzed glycan behavior in our binding pathways, we found that very

few D-serine binding pathways (27% for both GluN2A and GluN1) involve contacts with

glycans. While glycan-agonist interactions make up a small percentage of time spent in

binding pathways (10% for GluN2A and GluN1 D-serine pathways), patterns in glycan

interactions with the agonist as it binds suggest that glycans contribute to binding

pathways in a consistent way. The most common glycan-mediated D-serine-LBD

interactions for GluN2A involve an interaction network formed by N443-Man5 with Glu-

412, Lys-438 **(Fig. S6B)**, Lys-738, Glu-413 **(Fig. S6C)**, Tyr-730, and Ser-511 **(Fig.**

S6D), as D-serine moves into the binding pocket. Another contact network formed by

N444-Man5 with Lys-487, Asn-687 **(Fig. S6E)**, Arg-692, Arg-695, **(Fig. S6F)**, and Glu-

413 (alongside the N443-Man5 glycan). For GluN1, the N491-Man5 glycan interacts

with D-serine, trapping it in a network of interactions dominated by Arg-489 **(Fig. S6G)**.

When formed, this contact network functions as a kinetic trap that results in longer

binding pathways. Additionally, the N440-Man5 glycan also contacts D-serine as it

interacts with Arg-489 and Glu-497 **(Fig S6H)**. It is interesting to note that, unlike the

glycan-mediated contacts identified for GluN2A, glycan-mediated agonist contacts for

GluN1 do not involve D2 lobe residues. These glycan-mediated interactions illustrate

how glycan conformation can play a functional role through involvement with agonist

binding and LBD conformational dynamics. However, since glycan-mediated

interactions are so infrequent, the potentiating effect of glycan-D2 interactions

dominates functionally.

We quantified the dependence of glycan conformation on agonist binding and

LBD conformation by comparing glycan PMFs for different LBD conformations. For

GluN2A, we found that glycan-D2 interactions occur more readily when the LBD is

closed (calculated using a 1-dimensional projection of our LBD order parameter ξ_{12}, see

Methods). This effect was more dramatic for N443-Man5 than for N444-Man5 **(Fig.**

S7A,B). A similar relationship was determined for the N491-Man5 glycan of GluN1 **(Fig.**

S7C); this is consistent with previous simulations [29] that suggest that N491-Man5 acts

as a latch that stabilizes LBD closure. No significant relationship between glycan-D2

distance and the presence of an agonist (D-serine, glutamate, or both) in the binding

site was observed.

DISCUSSION

Here, we characterized the guided-diffusion mechanism that drives D-serine

binding to NMDAR LBDs. Instead of binding solely to the GluN1 LBD, we observed

substantial D-serine binding to the GluN2A LBD, a subunit widely accepted to bind to

the neurotransmitter glutamate. We showed by electrophysiology that D-serine at high

concentration can compete against glutamate at GluN2A, which in turn inhibits the

channel activity. In the context of synaptic transmission, our finding implies that D-serine

could play a role in modulating the strength of synaptic transmission. The synaptic

concentration of glutamate ranges from nanomolar concentrations [30] to >1 mM

following an action potential [31]. The synaptic concentration of D-serine is unclear,

however; the extracellular concentration of D-serine ranges from 5 to 7 µM [32] [33].

Possible routes for D-serine to enter the synapse include vesicular release by astroglia

[34] and transport by Asc-1 [35].

Free energy landscapes computed for GluN2A bound to glutamate [16], D-

serine, and glycine all indicate stabilization of the closed LBD bi-lobe, which is the

conformational state required for receptor activation. Agonists that can interact

extensively with bottom-lobe residues stabilize this state. Since glutamate does this to

the greatest extent, it is likely that D-serine does not generate sufficient force to fully

gate the ion channel. Subtle differences in the thermodynamics of agonist stabilization

suggest that kinetics further distinguish individual agonists. While glutamate has a

slightly higher association rate than D-serine, differences between association rates

across agonists and subunits is not drastic. We hypothesize that, in order for agonist

binding to result in NMDAR activation, the agonist must remain in the binding site long

enough to induce closure – we found that this is largely dependent upon the number

and strength of stable contacts the agonist forms with both D1 and D2 lobe residues.

We determined the role of N-linked glycans in agonist binding and stabilization.

Glycans impact agonist binding kinetics less by direct glycan-agonist interactions and

more by stabilizing the closed LBD through glycan-D2 interactions. This bias toward

LBD closure would increase the agonist residence time and potentiate NMDAR activity.

Our adaptation of pathway similarity analysis allowed us to identify clusters of

residues critical for binding agonists. This also allowed us to determine that the

presence of pathways depends on the degree of LBD closure. We also observed that D-

serine binds to GluN2A using similar pathways and residues as glutamate, while the

locations of key D-serine cluster residues for GluN1 are different. Applied more broadly

to drug-binding simulations, this method of analyzing binding pathways provides a

useful framework for gleaning biological insight from noisy and diffusive binding data.

METHODS

Equilibrium Molecular Dynamics Simulations

A construct of the GluN1/GluN2A dimer based on crystal structure (PDB ID:

2A5T [36]) used in our previous study [18] was used as a starting model. The residue

numberings are based on the Uniprot numbering for GRIN1 and GRIN2a entries.

$Man_5GlcNAc_2$ (Man5) glycans were added using CHARMM-GUI *Glycan Reader &*

Modeler [37] [38] [39] [40] to asparagine residues 440, 471, 491, and 771 of GluN1 and

asparagine residues 443 and 444 of GluN2A in accordance with physiologically relevant

glycosylation sites [20]. GluN2A was chosen as the GluN2 subtype both to facilitate

comparison with previous simulation studies and because recent evidence has

suggested that the GluN2A subtype is the primary subtype at synapses, where D-serine

is the dominant co-agonist [4].

All systems were solvated in a 140 Å × 110 Å × 110 Å orthorhombic water box

with ~150 mM NaCl using CHARMM [41]. All systems were electrically neutral. All

simulations in this work were performed using the CHARMM36 forcefield [42] and

TIP3P water model [43]. The systems were pre-equilibrated using NAMD 2.13 [44] first

using NVT conditions and gradually relaxing backbone-sidechain restraints and then for

15 ns using NPT conditions at a pressure of 1 atm and a temperature of 310 K. The pre-

equilibrated systems were then simulated on Anton 2 provided by the Pittsburgh

Supercomputer Center [45]. A weak center-of-mass restraint of 0.5 kcal mol^{-1} Å^{-1} was

applied to GluN2A N, CA, and C atoms of residues 461-463. 507-509, and 523-525 to

prevent large protein translational motion. Simulations on Anton 2 were carried out at

310 K with the NPT ensemble and with the weak center-of-mass restraint of 0.3 kcal

mol^{-1} Å^{-1} in accordance with previous simulations [19]. Additional simulation details are

provided in **Dataset S1**.

Identification of binding pathways

Identifying frames in which the ligand is bound in the receptor's binding pocket

provides key information about the ligand's binding affinity and the bound ensemble;

however, it fails to account for the process by which the ligand enters and leaves the

binding pocket. In guided diffusion, the residues that guide the ligand into the binding

pocket are critical for promoting the bound state. While imposing a simple distance

cutoff is sufficient for identifying the fully bound state, identifying the pathways by which

the ligand binds is less trivial. Here, we introduce a "binding chains" paradigm for

defining the ligand's path along the protein. These binding chains are defined from

ligand association to dissociation. An association begins when any polar ligand heavy

atom comes within 6 Å of any protein polar heavy atom. The ligand is considered

associated until is diffuses beyond 10 Å from the protein. The resulting chains are then

filtered by contact with the selected "docking" residue(s). Here, we use the conserved

arginine residue for each subunit (Arg-523 for GluN1 and Arg-518 for GluN2A) as the

essential docking residue. These chains are filtered then split into their "binding" and

"unbinding" components by a more specific docking criterion. In our case, we require

that the NH1 and NH2 atoms of the conserved arginine be within 4 Å of the ligand

carboxyl in accordance with the following scheme:

Condition 1: Arg NH1 is within 4 Å of the ligand OT1 <u>AND</u> Arg NH2 is within 4 Å of the

ligand OT2

<u>OR</u>

Condition 2: Arg NH2 is within 4 Å of the ligand OT1 <u>AND</u> Arg NH1 is within 4 Å of the

ligand OT2

This scheme accounts for both the crystallographic binding pose (Condition 2) and a

"flipped" ligand orientation (Condition 1). Chains that fail to meet these criteria are

discarded. Since binding and unbinding pathways can be considered reversible, we

combine them in our analysis, reversing the order of the unbinding pathways so that all

pathways have the same directionality. This results in a series of binding pathways we

can characterize both geometrically and in terms of key residue interactions.

Pathway Similarity Analysis and Clustering

Pathway Similarity Analysis (PSA) was applied to each binding pathway by

monitoring the agonist position as it binds. PSA involves computing a pairwise distance

metric between paths that serves as a measure of geometric similarity [21]. The

weighted average Hausdorff distance was selected as the path metric because it gave

the most geospatially distinct clusters of agonist density around the protein. This

weighted average Hausdorff distance was computed for all pairs of paths using the

following formula as described in previous work [21] and implemented in the

MDAnalysis python package [46][47]. The weighted-average Hausdorff distance

between two paths A and B can be expressed as:

493
$$\delta_H^{w_{avg}}(A, B) = \frac{1}{2}\left[\frac{1}{|A|}\delta_H^{sum}(A|B) + \frac{1}{|B|}\delta_H^{sum}(B|A)\right],$$

495   where $|A|$ and $|B|$ are the number of frames in paths A and B, respectively, and δ_H^{sum} is

496   the one-sided summed Hausdorff distance from path A to path B,

498
$$\delta_H^{sum}(A|B) = \sum_{a \in A} \min_{b \in B} d(a, b) \ .$$

Here, $d(a, b)$ represents the distance between point a of path A and point b in path B.

For our system, each point a is the agonist C_α position for a single frame in path A, and

each point b is the agonist C_α position for a single frame in path B. Therefore, $d(a,b)$

represents the Euclidean distance between the agonist C_α's of points in paths A and B.

$\delta_H^{\text{sum}}(A|B)$ is then computed by summing the shortest distance from each point a in path

A to any point b of path B over all points in path A. Each of the normalized one-sided

sums $\delta_H^{\text{sum}}(A|B)$ and $\delta_H^{\text{sum}}(B|A)$ are then averaged with equal weights. This does not

give more weight to pathways with more frames, thus removing the temporal

component from the analysis. Temporal patterns in binding pathways are analyzed for

the spatial clusters separately.

These path pairs were then clustered using hierarchical clustering according to

their weighted-average Hausdorff distances with the Ward (minimum variance) linkage

criterion as described in previous work [21] and implemented in SciPy [48]. The

complete linkage criterion also gave reasonable clustering. This agglomerative metric

assigns clusters by successively combining clusters that minimize the sum of squared

errors between them. Hierarchical clustering presents an advantage here because it

does not assume the number of clusters *a priori*. Rather, final clusters were selected

using the Ward distances showed in the dendrograms (see supplemental) as a guide

and by overlaying the ligand occupancy density on the protein to ensure that each

cluster represents a distinct spatial region of the protein.

Quantifying residue similarity with the overlap coefficient (Szymkiewicz–Simpson

coefficient)

To quantify the similarity between two sets of residues A and B, the overlap

coefficient was computed by dividing the number of overlapping residues between A

and B by the size of the smaller set of residues and is illustrated in the equation below

[22]:

528
$$OC(A, B) = \frac{|A \cap B|}{\min (|A|, |B|)} \; .$$

Scaling the size of the intersection by the smallest set size normalizes the overlap and

accounts for the large range in pathway lengths. If A is a subset of B, then $OC(A, B) = 1$.

This scaling method is appropriate, since these pathways are stochastic and involve a

mixture of random residue contacts and "guiding" residue contacts critical for binding.

This would be problematic for the more common Jaccard similarity metric, which scales

the intersection by the total size of both sets, where many random contacts increase

pathway length and dilute the value of the similarity metric.

The overlap coefficient was used to quantify the residue overlap between pairs of

pathways in each cluster to validate the spatial clustering and determine whether

pathways within clusters involve similar residue contacts. In addition, this metric was

used to quantify the similarity between residues involved in D-serine and glutamate

binding.

Umbrella Sampling

All-atom models were constructed from monomeric GluN1 (PDB ID: 1PB8 [17])

and GluN2A (based on PDB ID: 2A5S [36]). Since no crystal structure of D-serine

bound GluN2A exists, LBDs were constructed using MODELLER [49] to fill in missing

residues, and sidechain remodeling was performed on those residues using SCWRL4

[50]. D-serine and glycine were modelled into the GluN2A LBD by superimposing the

conserved arginine of the 2A5S glutamate-bound crystal structure (Arg-518) with the

conserved arginine of the D-serine (1PB8) or glycine (1PB7) bound crystal structure,

since there exists no crystal structure of GluN2A bound to these agonists. Bound

crystallographic waters in the GluN2A (2A5S) and GluN1 (1PB8) structures were

retained in the simulations.

To generate windows for umbrella sampling, targeted molecular dynamics

simulations were performed by "opening" the closed LBD along the order parameter

(ξ_1, ξ_2) [16]. Specifically, ξ_1 and ξ_2 are defined as the center of mass distance between

the backbone atoms of the following residue selections: ξ_1 is defined by residues 484-

485 and 688-689 for GluN1 and residues 485-486 and 689-690 for GluN2A. ξ_2 is

defined by residues 405-407 and 714-715 for GluN1 and 413-414 and 713-714 for

GluN2A. 205 simulation windows were selected at 1 Å × 1 Å increments. Each window

was solvated with a solvent box with dimensions 94 Å × 72 Å × 68 Å and 150 mM

NaCl.

Umbrella sampling simulations were performed by applying a bias of 2 kcal mol^{-1}

to the (ξ_1, ξ_2) order parameter to each of the 205 simulation windows. Equilibration was

performed in an NVT ensemble by gradually relaxing backbone and sidechain

restraints, and production simulations were carried out in an NPT ensemble at 300 K

and 1 atm for best comparison with previously computed NMDAR LBD monomers [16].

To ensure that the agonist does not diffuse out of the binding site, a restraint of 2 kcal

mol^{-1} Å^{-1} between the carboxyl group of the agonist and the guanidinium group of the

conserved arginine (Arg-523 for GluN1 and Arg-518 for GluN2A) was applied if the

distance between these groups exceeded 3.2 Å. Previous work has indicated that these

restraints do not affect the results but ensures that only the bound population is

sampled [16]. A weak center-of-mass restraint of 0.5 kcal mol^{-1} Å^{-1} was used applied to

the N, CA, and C atoms of residues 461-463, 507-509, and 523-525 for GluN2A and

residues 460-462, 512-514, and 528-530 for GluN1 to prevent translational protein

motion. Biased trajectories were mathematically unbiased using the weighted histogram

analysis method (WHAM) [51] [52]. 5 ns of production sampling for each window were

used to compute the potential of mean force (PMF) for each simulation agonist.

Standard deviations of all PMFs were computed by block averaging with ten blocks of

trajectory for each window [53].

Computing energetics of glycan conformational dynamics

To quantify glycan conformational dynamics, a glycan-D2 order parameter was

defined as the minimum distance between the heavy atoms of the glycans near the

binding cleft (N491-Man5 for GluN1 and N443-Man5 and N444-Man5 for GluN2A) and

the bottom lobe C_α atoms (residues 537-544 and 663-754 for GluN1 and residues 533-

539 and 661-757 for GluN2A). One relative PMF was computed for each of the three

near-pocket glycans using a window size of 0.2 Å using all glycosylated datasets. Error

for each PMF was quantified using the standard deviation computed by block averaging

with five blocks **(Fig. S5A-D)**. Blocks for which the window is not sampled were omitted

from the error calculation; this was only necessary for high glycan distances >20 Å. A

1D projection of the (ξ_1, ξ_2) order parameter, ξ_{12}, which averages ξ_1 and ξ_2, was used as

a single measure of LBD closure for computing glycan PMFs [16][54][55].

Electrophysiology

cRNA encoding GluN1-4b and GluN2A was injected into defolliculated *Xenopus*

laevis oocytes (0.2–0.5 ng total cRNA per oocyte). The oocytes were incubated in

recovery medium (50% L-15 medium (Hyclone) buffered by 15mM Na-HEPES at a final

pH of 7.4), supplemented with 100 µg mL^{-1} streptomycin, and 100 U mL^{-1} penicillin at

18°C. Two electrode voltage clamp (TEVC; Axoclamp-2B) recording was performed

between 24 to 48 hours after injection using an extracellular solution containing 5 mM

HEPES, 100 mM NaCl, 0.3 mM $BaCl_2$, 10mM Tricine at final pH 7.4 (adjusted with

KOH). The current was measured using agarose-tipped microelectrode (0.4–0.9 MΩ) at

the holding potential of −60 mV. Maximal response currents were evoked by 50 µM of

D-serine and 100 µM of L-glutamate. Data was acquired by the program PatchMaster

(HEKA) and analyzed by Origin 8 (OriginLab Corp).

Acknowledgements

Anton 2 computer time (MCB130045P) was provided by the Pittsburgh Supercomputing

Center (PSC) through NIH grant R01GM116961 (to A.Y.L.); the Anton 2 machine at

PSC was generously made available by D.E. Shaw Research. We also used resources

provided by the Maryland Advanced Research Computing Center (MARCC) at Johns

Hopkins University. This work was funded by the Johns Hopkins Catalyst Award (to

A.Y.L.); NIH T32GM135131 (to R.A.Y. and S.J.B.); NIH NS111745 and MH085926 (to

H.F.); Robertson funds at CSHL, Doug Fox Alzheimer's fund, Austin's purpose,

Heartfelt Wing Alzheimer's fund, and the Gertrude and Louis Feil Family Trust (to H.F.).

Author Contributions

R.A.Y. conducted molecular dynamics simulations; R.A.Y., S.J.B., and A.Y.L. analyzed

the results; T.-H.C. and H.F. designed and conducted experiments related to

electrophysiology; R.A.Y., T.-H.C., H.F., and A.Y.L. wrote the manuscript.

Competing Interests

The authors declare no competing interests.

References

(1) Johnson, J. W.; Ascher, P. Glycine Potentiates the NMDA Response in Cultured

Mouse Brain Neurons. *Nature* **1987**, *325* (6104), 529–531.

https://doi.org/10.1038/325529a0.

(2) Forsythe, I. D.; Westbrook, G. L.; Mayer, M. L. Modulation of Excitatory Synaptic

Transmission by Glycine and Zinc in Cultures of Mouse Hippocampal Neurons. *J.*

Neurosci. **1988**, *8* (10), 3733–3741. https://doi.org/10.1523/jneurosci.08-10-

03733.

(3) Kleckner, N. W.; Dingledine, R. Requirement for Glycine in Activation of NMDA

Receptors Expressed in Xenopus Oocytes. *Science (80-.).* **1988**, *241* (4867),

835–837. https://doi.org/10.1126/science.2841759.

(4) Papouin, T.; Ladépêche, L.; Ruel, J.; Sacchi, S.; Labasque, M.; Hanini, M.; Groc,

L.; Pollegioni, L.; Mothet, J. P.; Oliet, S. H. R. Synaptic and Extrasynaptic NMDA

Receptors Are Gated by Different Endogenous Coagonists. *Cell* **2012**, *150* (3),

640   633–646. https://doi.org/10.1016/j.cell.2012.06.029.

641 (5) Wolosker, H.; Blackshaw, S.; Snyder, S. H. Serine Racemase: A Glial Enzyme

642   Synthesizing D-Serine to Regulate Glutamate-N-Methyl-D-Aspartate

643   Neurotransmission. *Proc. Natl. Acad. Sci. U. S. A.* **1999**, *96* (23), 13409–13414.

644   https://doi.org/10.1073/pnas.96.23.13409.

645 (6) Miya, K.; Inoue, R.; Takata, Y.; Abe, M.; Natsume, R.; Sakimura, K.; Hongou, K.;

646   Miyawaki, T.; Mori, H. Serine Racemase Is Predominantly Localized in Neurons in

647   Mouse Brain. *J. Comp. Neurol.* **2008**, *510* (6), 641–654.

648   https://doi.org/10.1002/cne.21822.

649 (7) Balu, D. T.; Takagi, S.; Puhl, M. D.; Benneyworth, M. A.; Coyle, J. T. D-Serine

650   and Serine Racemase Are Localized to Neurons in the Adult Mouse and Human

651   Forebrain. *Cell. Mol. Neurobiol.* **2014**, *34* (3), 419–435.

652   https://doi.org/10.1007/s10571-014-0027-z.

653 (8) Rutter, A. R.; Fradley, R. L.; Garrett, E. M.; Chapman, K. L.; Lawrence, J. M.;

654   Rosahl, T. W.; Patel, S. Evidence from Gene Knockout Studies Implicates Asc-1

655   as the Primary Transporter Mediating D-Serine Reuptake in the Mouse CNS. *Eur.*

656   *J. Neurosci.* **2007**, *25* (6), 1757–1766. https://doi.org/10.1111/j.1460-

657   9568.2007.05446.x.

658 (9) Coyle, J. T.; Balu, D.; Wolosker, H. D-Serine, the Shape-Shifting NMDA Receptor

659   Co-Agonist. *Neurochem. Res.* **2020**, 1–10. https://doi.org/10.1007/s11064-020-

660   03014-1.

661 (10) Henneberger, C.; Papouin, T.; Oliet, S. H. R.; Rusakov, D. A. Long-Term

662   Potentiation Depends on Release of d-Serine from Astrocytes. *Nature* **2010**, *463*

663   (7278), 232–236. https://doi.org/10.1038/nature08673.

664   (11)   Peyrovian, B.; Rosenblat, J. D.; Pan, Z.; Iacobucci, M.; Brietzke, E.; McIntyre, R.

665   S. The Glycine Site of NMDA Receptors: A Target for Cognitive Enhancement in

666   Psychiatric Disorders. *Progress in Neuro-Psychopharmacology and Biological*

667   *Psychiatry*. Elsevier Inc. June 8, 2019, pp 387–404.

668   https://doi.org/10.1016/j.pnpbp.2019.02.001.

669   (12)   MacKay, M. A. B.; Kravtsenyuk, M.; Thomas, R.; Mitchell, N. D.; Dursun, S. M.;

670   Baker, G. B. D-Serine: Potential Therapeutic Agent and/or Biomarker in

671   Schizophrenia and Depression? *Front. Psychiatry* **2019**, *10* (FEB), 18–23.

672   https://doi.org/10.3389/fpsyt.2019.00025.

673   (13)   Kantrowitz, J. T.; Malhotra, A. K.; Cornblatt, B.; Silipo, G.; Balla, A.; Suckow, R.

674   F.; D'Souza, C.; Saksa, J.; Woods, S. W.; Javitt, D. C. High Dose D-Serine in the

675   Treatment of Schizophrenia. *Schizophr. Res.* **2010**, *121* (1–3), 125–130.

676   https://doi.org/10.1016/j.schres.2010.05.012.

677   (14)   Heresco-Levy, U.; Javitt, D. C.; Ebstein, R.; Vass, A.; Lichtenberg, P.; Bar, G.;

678   Catinari, S.; Ermilov, M. D-Serine Efficacy as Add-on Pharmacotherapy to

679   Risperidone and Olanzapine for Treatment-Refractory Schizophrenia. *Biol.*

680   *Psychiatry* **2005**, *57* (6), 577–585. https://doi.org/10.1016/j.biopsych.2004.12.037.

681   (15)   Mayer, M. L. The Challenge of Interpreting Glutamate-Receptor Ion-Channel

682   Structures. *Biophysical Journal*. Biophysical Society November 21, 2017, pp

683   2143–2151. https://doi.org/10.1016/j.bpj.2017.07.028.

684   (16)   Yao, Y.; Belcher, J.; Berger, A. J.; Mayer, M. L.; Lau, A. Y. Conformational

685   Analysis of NMDA Receptor GluN1, GluN2, and GluN3 Ligand-Binding Domains

Reveals Subtype-Specific Characteristics. *Structure* **2013**, *21* (10), 1788–1799.

https://doi.org/10.1016/j.str.2013.07.011.

(17)   Furukawa, H.; Gouaux, E. Mechanisms of Activation, Inhibition and Specificity:

Crystal Structures of the NMDA Receptor NR1 Ligand-Binding Core. *EMBO J.*

2003, *22* (12), 2873–2885. https://doi.org/10.1093/emboj/cdg303.

(18)   Yu, A.; Lau, A. Y. Glutamate and Glycine Binding to the NMDA Receptor.

Structure **2018**, *26* (7), 1035-1043.e2. https://doi.org/10.1016/j.str.2018.05.004.

(19)   Yu, A.; Salazar, H.; Plested, A. J. R.; Lau, A. Y. Neurotransmitter Funneling

Optimizes Glutamate Receptor Kinetics. *Neuron* **2018**, *97* (1), 139-149.e4.

https://doi.org/10.1016/j.neuron.2017.11.024.

(20)   Kaniakova, M.; Lichnerova, K.; Skrenkova, K.; Vyklicky, L.; Horak, M. Biochemical

and Electrophysiological Characterization of *N*- Glycans on NMDA Receptor

Subunits. *J. Neurochem.* **2016**, *138* (4), 546–556.

https://doi.org/10.1111/jnc.13679.

(21)   Seyler, S. L.; Kumar, A.; Thorpe, M. F.; Beckstein, O. Path Similarity Analysis: A

Method for Quantifying Macromolecular Pathways. *PLoS Comput. Biol.* **2015**, *11*

(10), 1004568. https://doi.org/10.1371/journal.pcbi.1004568.

(22)   Vijaymeena, M. K.; Kavitha, K. A SURVEY ON SIMILARITY MEASURES IN

TEXT MINING. *Mach. Learn. Appl. An Int. J.* **2016**, *3* (1).

https://doi.org/10.5121/mlaij.2016.3103.

(23)   Jalali-Yazdi, F.; Chowdhury, S.; Yoshioka, C.; Gouaux, E. Mechanisms for Zinc

and Proton Inhibition of the GluN1/GluN2A NMDA Receptor. *Cell* **2018**, *175* (6),

1520-1532.e15. https://doi.org/10.1016/j.cell.2018.10.043.

(24) Sinitskiy, A. V.; Stanley, N. H.; Hackos, D. H.; Hanson, J. E.; Sellers, B. D.;

Pande, V. S. Computationally Discovered Potentiating Role of Glycans on NMDA

Receptors. *Sci. Rep.* **2017**, *7* (1), 1–10. https://doi.org/10.1038/srep44578.

(25) Dolino, D. M.; Adariani, S. R.; Shaikh, S. A.; Jayaraman, V.; Sanabria, H.

Conformational Selection and Submillisecond Dynamics of the Ligand-Binding

Domain of the N-Methyl-d-Aspartate Receptor. *J. Biol. Chem.* **2016**, *291* (31),

16175. https://doi.org/10.1074/JBC.M116.721274.

(26) Rajab, S.; Bismin, L.; Schwarze, S.; Pinggera, A.; Greger, I. H.; Neuweiler, H.

Allosteric Coupling of Sub-Millisecond Clamshell Motions in Ionotropic Glutamate

Receptor Ligand-Binding Domains. *Commun. Biol. 2021 41* **2021**, *4* (1), 1–11.

https://doi.org/10.1038/s42003-021-02605-0.

(27) Snyder, S. H.; Mustafa, A. K.; Kim, P. M. D-Serine as a Putative Glial

Neurotransmitter. *Neuron Glia Biol.* **2004**, *1* (3), 275–281.

https://doi.org/10.1017/S1740925X05000141.

(28) Dror, R. O.; Pan, A. C.; Arlow, D. H.; Borhani, D. W.; Maragakis, P.; Shan, Y.; Xu,

H.; Shaw, D. E. Pathway and Mechanism of Drug Binding to G-Protein-Coupled

Receptors. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108* (32), 13118–13123.

https://doi.org/10.1073/pnas.1104614108.

(29) Sinitskiy, A. V.; Pande, V. S. Simulated Dynamics of Glycans on Ligand-Binding

Domain of NMDA Receptors Reveals Strong Dynamic Coupling between Glycans

and Protein Core. *J. Chem. Theory Comput.* **2017**, *13* (11), 5496–5505.

https://doi.org/10.1021/acs.jctc.7b00817.

(30) Chiu, D. N.; Jahr, C. E. Extracellular Glutamate in the Nucleus Accumbens Is

Nanomolar in Both Synaptic and Non-Synaptic Compartments. *Cell Rep.* **2017**, *18* (11), 2576–2583. https://doi.org/10.1016/j.celrep.2017.02.047.

(31) Dzubay, J. A.; Jahr, C. E. The Concentration of Synaptically Released Glutamate Outside of the Climbing Fiber-Purkinje Cell Synaptic Cleft. *J. Neurosci.* **1999**, *19* (13), 5265–5274. https://doi.org/10.1523/jneurosci.19-13-05265.

(32) Matsui, T. -a; Sekiguchi, M.; Hashimoto, A.; Tomita, U.; Nishikawa, T.; Wada, K. Functional Comparison of D-Serine and Glycine in Rodents: The Effect on Cloned NMDA Receptors and the Extracellular Concentration. *J. Neurochem.* **1995**, *65* (1), 454–458. https://doi.org/10.1046/j.1471-4159.1995.65010454.x.

(33) Hashimoto, A.; Oka, T.; Nishikawa, T. Extracellular Concentration of Endogenous Free D-Serine in the Rat Brain as Revealed by in Vivo Microdialysis. *Neuroscience* **1995**, *66* (3), 635–643. https://doi.org/10.1016/0306-4522(94)00597-X.

(34) Mothet, J. P.; Pollegioni, L.; Ouanounou, G.; Martineau, M.; Fossier, P.; Baux, G. Glutamate Receptor Activation Triggers a Calcium-Dependent and SNARE Protein-Dependent Release of the Gliotransmitter D-Serine. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (15), 5606–5611. https://doi.org/10.1073/pnas.0408483102.

(35) Sason, H.; Billard, J. M.; Smith, G. P.; Safory, H.; Neame, S.; Kaplan, E.; Rosenberg, D.; Zubedat, S.; Foltyn, V. N.; Christoffersen, C. T.; Bundgaard, C.; Thomsen, C.; Avital, A.; Christensen, K. V.; Wolosker, H. Asc-1 Transporter Regulation of Synaptic Activity via the Tonic Release of -Serine in the Forebrain. *Cereb. Cortex* **2017**, *27* (2), 1573–1587. https://doi.org/10.1093/cercor/bhv350.

(36) Furukawa, H.; Singh, S. K.; Mancusso, R.; Gouaux, E. Subunit Arrangement and

Function in NMDA Receptors. *Nature* **2005**, *438* (7065), 185–192.

https://doi.org/10.1038/nature04089.

(37) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: A Web-Based Graphical User

Interface for CHARMM. *J. Comput. Chem.* **2008**, *29* (11), 1859–1865.

https://doi.org/10.1002/jcc.20945.

(38) Jo, S.; Song, K. C.; Desaire, H.; MacKerell, A. D.; Im, W. Glycan Reader:

Automated Sugar Identification and Simulation Preparation for Carbohydrates and

Glycoproteins. *J. Comput. Chem.* **2011**, *32* (14), 3135–3141.

https://doi.org/10.1002/jcc.21886.

(39) Park, S. J.; Lee, J.; Patel, D. S.; Ma, H.; Lee, H. S.; Jo, S.; Im, W. Glycan Reader

Is Improved to Recognize Most Sugar Types and Chemical Modifications in the

Protein Data Bank. *Bioinformatics* **2017**, *33* (19), 3051–3057.

https://doi.org/10.1093/bioinformatics/btx358.

(40) Park, S. J.; Lee, J.; Qi, Y.; Kern, N. R.; Lee, H. S.; Jo, S.; Joung, I.; Joo, K.; Lee,

J.; Im, W. CHARMM-GUI Glycan Modeler for Modeling and Simulation of

Carbohydrates and Glycoconjugates. *Glycobiology* **2019**, *29* (4), 320–331.

https://doi.org/10.1093/glycob/cwz003.

(41) Brooks, B. R.; Brooks, C. L.; Mackerell, A. D.; Nilsson, L.; Petrella, R. J.; Roux, B.;

Won, Y.; Archontis, G.; Bartels, C.; Boresch, S.; Caflisch, A.; Caves, L.; Cui, Q.;

Dinner, A. R.; Feig, M.; Fischer, S.; Gao, J.; Hodoscek, M.; Im, W.; Kuczera, K.;

Lazaridis, T.; Ma, J.; Ovchinnikov, V.; Paci, E.; Pastor, R. W.; Post, C. B.; Pu, J.

Z.; Schaefer, M.; Tidor, B.; Venable, R. M.; Woodcock, H. L.; Wu, X.; Yang, W.;

York, D. M.; Karplus, M. CHARMM: The Biomolecular Simulation Program. *J.*

Comput. Chem. **2009**, *30* (10), 1545–1614. https://doi.org/10.1002/jcc.21287.

(42)   MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.;

780   Field, M. J.; Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir,

781   L.; Kuczera, K.; Lau, F. T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.;

782   Prodhom, B.; Reiher, W. E.; Roux, B.; Schlenkrich, M.; Smith, J. C.; Stote, R.;

783   Straub, J.; Watanabe, M.; Wiórkiewicz-Kuczera, J.; Yin, D.; Karplus, M. All-Atom

784   Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J.*

785   *Phys. Chem. B* **1998**, *102* (18), 3586–3616. https://doi.org/10.1021/jp973084f.

786   (43)   Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L.

787   Comparison of Simple Potential Functions for Simulating Liquid Water. *J. Chem.*

788   *Phys.* **1983**, *79* (2), 926–935. https://doi.org/10.1063/1.445869.

789   (44)   Phillips, J. C.; Braun, R.; Wang, W.; Gumbart, J.; Tajkhorshid, E.; Villa, E.; Chipot,

790   C.; Skeel, R. D.; Kalé, L.; Schulten, K. Scalable Molecular Dynamics with NAMD.

791   *Journal of Computational Chemistry*. John Wiley and Sons Inc. December 1,

792   2005, pp 1781–1802. https://doi.org/10.1002/jcc.20289.

793   (45)   Shaw, D. E.; Grossman, J. P.; Bank, J. A.; Batson, B.; Butts, J. A.; Chao, J. C.;

794   Deneroff, M. M.; Dror, R. O.; Even, A.; Fenton, C. H.; Forte, A.; Gagliardo, J.; Gill,

795   G.; Greskamp, B.; Ho, C. R.; Ierardi, D. J.; Iserovich, L.; Kuskin, J. S.; Larson, R.

796   H.; Layman, T.; Lee, L. S.; Lerer, A. K.; Li, C.; Killebrew, D.; Mackenzie, K. M.;

797   Mok, S. Y. H.; Moraes, M. A.; Mueller, R.; Nociolo, L. J.; Peticolas, J. L.; Quan, T.;

798   Ramot, D.; Salmon, J. K.; Scarpazza, D. P.; Ben Schafer, U.; Siddique, N.;

799   Snyder, C. W.; Spengler, J.; Tang, P. T. P.; Theobald, M.; Toma, H.; Towles, B.;

800   Vitale, B.; Wang, S. C.; Young, C. Anton 2: Raising the Bar for Performance and

Programmability in a Special-Purpose Molecular Dynamics Supercomputer. In

International Conference for High Performance Computing, Networking, Storage

and Analysis, SC; IEEE Computer Society, 2014; Vol. 2015-January, pp 41–53.

https://doi.org/10.1109/SC.2014.9.

(46)   Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A

Toolkit for the Analysis of Molecular Dynamics Simulations. *J. Comput. Chem.*

2011, *32* (10), 2319–2327. https://doi.org/10.1002/jcc.21787.

(47)   Gowers, R.; Linke, M.; Barnoud, J.; Reddy, T.; Melo, M.; Seyler, S.; Domański, J.;

Dotson, D.; Buchoux, S.; Kenney, I.; Beckstein, O. MDAnalysis: A Python

Package for the Rapid Analysis of Molecular Dynamics Simulations. In

Proceedings of the 15th Python in Science Conference; SciPy, 2016; pp 98–105.

https://doi.org/10.25080/majora-629e541a-00e.

(48)   Virtanen, P.; Gommers, R.; Oliphant, T. E.; Haberland, M.; Reddy, T.;

Cournapeau, D.; Burovski, E.; Peterson, P.; Weckesser, W.; Bright, J.; van der

Walt, S. J.; Brett, M.; Wilson, J.; Millman, K. J.; Mayorov, N.; Nelson, A. R. J.;

Jones, E.; Kern, R.; Larson, E.; Carey, C. J.; Polat, İ.; Feng, Y.; Moore, E. W.;

VanderPlas, J.; Laxalde, D.; Perktold, J.; Cimrman, R.; Henriksen, I.; Quintero, E.

A.; Harris, C. R.; Archibald, A. M.; Ribeiro, A. H.; Pedregosa, F.; van Mulbregt, P.;

Vijaykumar, A.; Bardelli, A. Pietro; Rothberg, A.; Hilboll, A.; Kloeckner, A.;

Scopatz, A.; Lee, A.; Rokem, A.; Woods, C. N.; Fulton, C.; Masson, C.;

Häggström, C.; Fitzgerald, C.; Nicholson, D. A.; Hagen, D. R.; Pasechnik, D. V.;

Olivetti, E.; Martin, E.; Wieser, E.; Silva, F.; Lenders, F.; Wilhelm, F.; Young, G.;

Price, G. A.; Ingold, G. L.; Allen, G. E.; Lee, G. R.; Audren, H.; Probst, I.; Dietrich,

J. P.; Silterra, J.; Webber, J. T.; Slavič, J.; Nothman, J.; Buchner, J.; Kulick, J.;

Schönberger, J. L.; de Miranda Cardoso, J. V.; Reimer, J.; Harrington, J.;

Rodríguez, J. L. C.; Nunez-Iglesias, J.; Kuczynski, J.; Tritz, K.; Thoma, M.;

Newville, M.; Kümmerer, M.; Bolingbroke, M.; Tartre, M.; Pak, M.; Smith, N. J.;

Nowaczyk, N.; Shebanov, N.; Pavlyk, O.; Brodtkorb, P. A.; Lee, P.; McGibbon, R.

T.; Feldbauer, R.; Lewis, S.; Tygier, S.; Sievert, S.; Vigna, S.; Peterson, S.; More,

S.; Pudlik, T.; Oshima, T.; Pingel, T. J.; Robitaille, T. P.; Spura, T.; Jones, T. R.;

Cera, T.; Leslie, T.; Zito, T.; Krauss, T.; Upadhyay, U.; Halchenko, Y. O.;

Vázquez-Baeza, Y. SciPy 1.0: Fundamental Algorithms for Scientific Computing

in Python. *Nat. Methods* **2020**, *17* (3), 261–272. https://doi.org/10.1038/s41592-

019-0686-2.

(49)   Webb, B.; Sali, A. Comparative Protein Structure Modeling Using MODELLER.

Curr. Protoc. Bioinforma. **2016**, *2016*, 5.6.1-5.6.37. https://doi.org/10.1002/cpbi.3.

(50)   Krivov, G. G.; Shapovalov, M. V.; Dunbrack, R. L. Improved Prediction of Protein

Side-Chain Conformations with SCWRL4. *Proteins Struct. Funct. Bioinforma.*

2009, *77* (4), 778–795. https://doi.org/10.1002/prot.22488.

(51)   Kumar, S.; Rosenberg, J. M.; Bouzida, D.; Swendsen, R. H.; Kollman, P. A. THE

Weighted Histogram Analysis Method for Free-energy Calculations on

Biomolecules. I. The Method. *J. Comput. Chem.* **1992**, *13* (8), 1011–1021.

https://doi.org/10.1002/jcc.540130812.

(52)   Souaille, M.; Roux, B. Extension to the Weighted Histogram Analysis Method:

Combining Umbrella Sampling with Free Energy Calculations. *Comput. Phys.*

Commun. **2001**, *135* (1), 40–57. https://doi.org/10.1016/S0010-4655(00)00215-0.

847   (53)   Grossfield, A.; Zuckerman, D. M. Chapter 2 Quantifying Uncertainty and Sampling

848   Quality in Biomolecular Simulations. *Annual Reports in Computational Chemistry*.

849   NIH Public Access 2009, pp 23–48. https://doi.org/10.1016/S1574-

850   1400(09)00502-7.

851   (54)   Wied, T. J.; Chin, A. C.; Lau, A. Y. High Conformational Variability in the GluK2

852   Kainate Receptor Ligand-Binding Domain. *Structure* **2019**, *27* (1), 189-195.e2.

853   https://doi.org/10.1016/j.str.2018.09.008.

854   (55)   Chin, A. C.; Yovanno, R. A.; Wied, T. J.; Gershman, A.; Lau, A. Y. D-Serine

855   Potently Drives Ligand-Binding Domain Closure in the Ionotropic Glutamate

856   Receptor GluD2. *Structure* **2020**, *28* (10), 1168-1178.e2.

857   https://doi.org/10.1016/j.str.2020.07.005.

Fig. 1. Identifying D-serine binding pathways for GluN2A using pathway similarity analysis (PSA).
(A) Overview of the PSA workflow for quantifying similarity between D-serine binding pathways. **(B)** 2-
dimensional order parameter (ξ_1, ξ_2) that describes the degree of GluN2A LBD closure. For each of the
above **(C-F)**, the left image shows D-serine density, while the right image shows the residues most
frequently contacted by D-serine as it enters/leaves the binding site for each cluster. Labeled residues
demonstrate ≥ 0.2 fractional occurrence defined relative to the most contacted residue in each cluster,
but all residues with ≥ 0.1 fractional occurrence are shown in stick representation (see **Dataset S5**). **(C)**
Cluster 1 involves residues of the ξ_2 face of the LBD. **(D)** Cluster 2 involves residues of the ξ_1 face of the
D1 lobe. **(E)** In Cluster 3, D-serine zigzags between D1 and D2 lobe residues of the ξ_1 face. **(F)** Cluster 4
primarily involves D2 lobe residues on the ξ_1 face.

873
874
875
Fig. 2. Identifying D-serine binding pathways for GluN1 using pathway similarity analysis (PSA).
(A) 2-dimensional order parameter (ξ_1, ξ_2) that describes the degree of GluN1 LBD closure. For each of
the above **(B-E)**, the left image shows D-serine density, while the right image shows the residues most
frequently contacted by D-serine as it enters/leaves the binding site for each cluster. Labeled residues
demonstrate ≥ 0.2 fractional occurrence defined relative to the most contacted residue in each cluster,
but all residues with ≥ 0.1 fractional occurrence are shown in stick representation (see **Dataset S6**). **(B)**
In Cluster 1, D-Serine contacts residues on the ξ_2 face of the LBD. **(C)** Cluster 2 involves interactions with
both D1 and D2 residues of the ξ_1 face. **(D)** Cluster 3 involves contacts with residues at the top of the D1
lobe on the ξ_1 face. **(E)** Cluster 4 is defined by interactions with D1 loop 2 that reaches into solution.

Fig. 3. Conformational free energy landscapes for GluN2A and GluN1 LBDs. Umbrella sampling molecular dynamics simulations were used to compute the potential of mean force (PMF) along the (ξ_1, ξ_2) order parameter for **(A)** D-serine bound to GluN2A, **(B)** glycine bound to GluN2A, **(C)** apo GluN2A previously computed in [16], **(D)** glutamate bound to GluN2A in its crystallographic pose previously computed in [16], **(E)** glutamate bound to GluN2A in the inverted pose identified in [18], **(F)** D-serine bound to GluN1, **(G)** glycine bound to GluN1 previously computed in [16], **(H)** apo GluN1 previously computed in [16].

Fig. 4. D-serine completes glutamate binding as an antagonist at high concentration. (A)
Representative Two-electrode voltage clamp (TEVC) recording on GluN1/GluN2A NMDARs expressing
oocytes. The trace showed GluN1 agonist D-serine inhibited the NMDAR current at a high concentration.
6 μM of glutamate was present throughout the recording. (B) Glutamate-concentration dependent dose-
response curves of high-concentration D-serine inhibition. (C) D-serine-concentration dependent dose-
response curves of glutamate potentiation (left). Schild plot analysis of D-serine competition on glutamate
(right). The calculated slope of the Schild plot was 1.11±0.13 and the intercept was 2.38±0.26. DR stands
for dose ratio. All the dose-response experiments were repeated at least four times.

Fig. 5. Comparison of D-serine and glutamate binding to GluN2A. (A) Overlay of D-serine (teal) and glutamate (gray) density. **(B)** Residues that distinguish D-serine (teal) from glutamate (gray) binding pathways (see **Dataset S7**).

Fig. 6. Conformational dynamics of near-pocket glycans. N-linked $Man_5GlcNAc_2$ (Man5) glycans **(A)** N443-Man5 and N444-Man5 for GluN2A and **(B)** N491-Man5 for GluN1. Glycan conformational energy landscapes for **(C)** GluN2A N443-Man5, **(D)** GluN2A N444-Man5, and **(E)** GluN1 N491-Man5 were obtained by computing the minimum distance between all glycan heavy atoms and D2 lobe residues and binning the distribution from all glycosylated simulation systems. Shaded error regions were computed using a block-averaging scheme described in Methods.

Supplementary Information for

Excitatory and inhibitory D-serine binding to the NMDA receptor

Remy A. Yovanno[1], Tsung Han Chou[2], Sarah J. Brantley[3], Hiro Furukawa[2*], and Albert Y. Lau[1*]

Affiliations:

[1]Department of Biophysics and Biophysical Chemistry, Johns Hopkins University School of Medicine, 725

N. Wolfe Street, WBSB 706, Baltimore, MD 21205, USA.

[2] W.M. Keck Structural Biology Laboratory, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724,

USA.

[3]Department of Biology, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

[*] To whom correspondence should be addressed:

alau@jhmi.edu (A.Y.L.)

furukawa@cshl.edu (H.F.)

Figures 1, 2 – figure supplement 1. Overlap coefficient analysis for GluN2A and GluN1 binding pathways. The overlap coefficient was computed for each pair of paths within each cluster for **(A)** GluN2A and **(B)** GluN1. The dotted line in each plot indicates the global mean $\langle OC \rangle$ = 0.557 for GluN2A and $\langle OC \rangle$ = 0.671 for GluN1. The mean OC for each cluster was computed for **(C)** GluN2A and **(D)** GluN1. For reference, the black solid line indicates the global mean OC.

50
51
52
Figures 1, 2 – figure supplement 2. D-serine pathway residues mapped onto the intact GluN2A NMDAR
(PDB ID: 6MMM[1]). Sidechain atoms are shown as spheres for **(A)** GluN2A in teal and **(B)** GluN1 in
magenta and the adjacent domains labelled.

Figures 1, 2 – figure supplement 3. Degree of LBD closure during D-serine binding pathways to **(A)**
GluN2A and **(B)** GluN1. The two-dimensional (ξ_1, ξ_2) order parameter was computed for all frames in
binding/unbinding pathways and was plotted as a function of relative density as indicated by the colorbar.
62

Figure 3 – figure supplement 1. (A) Binding-site residues for D-serine bound to the GluN2A LBD computed from lowest-energy conformers ($\leq$ 1 kcal mol^{-1}) from umbrella sampling simulations. Residues with >50% contact frequency in the ensemble of lowest-energy conformers are labelled here. **(B)** D-serine interaction with Thr-690 present in lowest-energy conformers.

75
76
77 **Figure 3 – figure supplement 2.** Error of umbrella sampling PMFs computed by block averaging for **(A)**
78 D-serine bound to GluN2A, **(B)** glycine bound to GluN2A, **(C)** glutamate bound to GluN2A in the inverted
79 pose, and **(D)** D-serine bound to GluN1. The colors of the colorbar correspond to the standard deviation
80 in kcal mol[-1], which was computed for each window over 10 blocks.
81
82
83

Figure 6 – figure supplement 1. N-linked Man$_5$GlcNAc$_2$ (Man5) glycans interacts with D-serine as it binds. **(A)** Schematic of the Man5 glycan. **(B)** Contact network involving the GluN2A N443-Man5 glycan and residues Glu-412 and, Lys-438, **(C)** Lys-738 and Glu-413, **(D)** Glu-413,Tyr-730, and Ser-511. **(E)** Contact network involving the GluN2A N444-Man5 glycan and residues Lys-487 and Asn-687, **(F)** Lys-487, Arg-692, and Arg-695. **(G)** Contact network formed between D-serine, the GluN1 N491-Man5 glycan, and Arg-489. **(H)** Additional GluN1 contact network formed between both GluN1 N491-Man5 and N440-Man5 glycans, residues Arg-489 and Glu-497, and D-serine.

Figure 6 – figure supplement 2. Glycan-D2 distance dependence on LBD closure and agonist binding for the **(A)** GluN2A N443-Man5 glycan, **(B)** GluN2A N444-Man5 glycan, and **(C)** GluN1 N491-Man5 glycan.

93
94

96
97 **Figures 1, 2 – figure supplement 4.** Dendrograms for hierarchical clustering of weighted average
98 hausdorff distances for D-serine binding pathways for **(A)** GluN2A and **(B)** GluN1 according to the Ward
99 linkage criterion.

Legends for Datasets S1-S11. Data extracted from molecular dynamics simulations.

S1. 1_Simulation_Summary: overview of simulation systems
S2. 2_GluN2A: record of all successful binding pathways in each simulation system for D-serine binding to GluN2A.
S3. 3_Glu: record of all successful binding pathways in each simulation system for glutamate binding to GluN2A.
S4. 4_GluN1: record of all successful binding pathways in each simulation system for D-serine binding to GluN1.
S5. 5_contact_freq_N2_glyc: per residue contact frequency analysis for D-serine binding to GluN2A by cluster identified with PSA.
S6. 6_contact_freq_N1_glyc: per residue contact frequency analysis for D-serine binding to GluN1 by cluster identified with PSA.
S7. 7_Dser_vs_Glu: comparison of relative residue contact frequency for D-serine and glutamate.
S8. 8_bound_state_analysis: per residue contact frequency analysis of the bound state for each agonist computed from lowest-energy conformers extracted from umbrella sampling simulations.
S9. 9_conditional_prob_N2: GluN2A residues most frequently contacted by D-serine given that the pathway results in successful binding – listed for each simulation system.
S10. 10_conditional_prob_N1: GluN1 residues most frequently contacted by D-serine given that the pathway results in successful binding – listed for each simulation system.
S11. 11_glyc_vs_noglyc: comparison of relative residue contact frequency during GluN2A and GluN1 binding pathways for glycosylated and non-glycosylated simulations.

Legends for Movies S1-S2

Movie 1. Process of D-serine (spheres) binding to the GluN2A LBD (light orange cartoon). D-serine diffuses in bulk solvent until it contacts LBD residues (sticks) that guide it into the binding site. Shown also are the GluN1 LBD (light blue cartoon) and N-linked Man5 glycans (sticks).
Movie 2. Process of D-serine (spheres) binding to the GluN1 LBD (light blue cartoon). D-serine diffuses in bulk solvent until it contacts LBD residues (sticks) that guide it into the binding site. Shown also are the GluN2A LBD (light orange cartoon) and N-linked Man5 glycans (sticks).

SI References

1. Jalali-Yazdi, F.; Chowdhury, S.; Yoshioka, C.; Gouaux, E. Mechanisms for Zinc and Proton Inhibition of the GluN1/GluN2A NMDA Receptor. *Cell* **2018**, *175* (6), 1520-1532.e15. https://doi.org/10.1016/j.cell.2018.10.043.