1 # Differential DNA methylation in Pacific oyster reproductive

2 # tissue in response to ocean acidification

3
Running title: Low pH influence on oyster methylome

Yaamini R. Venkataraman[1,2,*], Samuel J. White[2], Steven B. Roberts[2]

[1]Biology Department, Woods Hole Oceanographic Institution, 266 Water Street, Woods Hole,

MA, 02543, USA

[2]School of Aquatic & Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle,

WA, USA

Abstract

Background: There is a need to investigate mechanisms of phenotypic plasticity in marine

invertebrates as negative effects of climate change, like ocean acidification, are experienced by

coastal ecosystems. Environmentally-induced changes to the methylome may regulate gene

expression, but methylome responses can be species- and tissue-specific. Tissue-specificity

has implications for gonad tissue, as gonad-specific methylation patterns may be inherited by

offspring. We used the Pacific oyster (*Crassostrea gigas)* — a model for understanding pH

impacts on bivalve molecular physiology due to its genomic resources and importance in global

aquaculture— to assess how low pH could impact the gonad methylome. Oysters were exposed

to either low pH (7.31 ± 0.02) or ambient pH (7.82 ± 0.02) conditions for seven weeks. Whole

genome bisulfite sequencing was used to identify methylated regions in female oyster gonad

samples. C->T single nucleotide polymorphisms were identified and removed to ensure

accurate methylation characterization.

Results: Analysis of gonad methylomes revealed a total of 1,284 differentially methylated loci

(DML) found primarily in genes, with several genes containing multiple DML. Gene ontologies

for genes containing DML were involved in development and stress response, suggesting

methylation may promote gonad growth homeostasis in low pH conditions. Additionally, several

of these genes were associated with cytoskeletal structure regulation, metabolism, and protein

ubiquitination — commonly-observed responses to ocean acidification. Comparison of these

DML with other *Crassostrea* spp. exposed to ocean acidification demonstrates that similar

pathways, but not identical genes, are impacted by methylation.

Conclusions: Our work suggests DNA methylation may have a regulatory role in gonad and

larval development, which would shape adult and offspring responses to low pH stress.

Combined with existing molluscan methylome research, our work further supports the need for

tissue- and species-specific studies to understand the potential regulatory role of DNA

methylation.

Keywords: Pacific oyster, bivalve, ocean acidification, DNA methylation, gonad development

Background

There is great interest in elucidating how changes in the environment can impact marine

invertebrate stress responses by examining DNA methylation [1, 2]. Environmental variation can

alter the positions of methyl groups on CpG dinucleotides, potentially regulating gene

expression by fine-tuning expression of housekeeping genes or providing additional

transcriptional opportunities for environmental response genes [3]. Methylation regulation of

gene expression could then influence plasticity in response to environmental stressors [1]. This

framework can be applied towards understanding rapid acclimatization of invasive species [4,

5], identification of target epialleles to aid assisted evolution [6], and rearing practices for

production of environmentally-resilient aquaculture species [7]. Ocean acidification [8–13], heat

stress [14], upwelling conditions [15], disease [16], salinity (Johnson et al. 2021), pesticide

exposure [17], and temperature and salinity [13, 18] have all elicited changes in marine

invertebrate methylomes, demonstrating a potential for methylation to regulate organismal

responses to a variety of environmental conditions.

Given the impact of ocean acidification on calcifying species like oysters [19–21], several

environmental epigenetic studies have examined the influence of ocean acidification on

molluscan methylomes (Table 1). For example, eastern oyster (*Crassostrea virginica*) mantle

[10], *C. virginica* reproductive tissue [12], Hong kong oyster (*Crassostrea hongkongensis*)

mantle [8], and *C. hongkongensis* larval methylomes [11] all display changes in DNA

methylation after experimental ocean acidification exposure. Between all four studies, only

Venkataraman et al. (2020) reports the percent of CpGs methylated: the stated 22% is

consistent with the baseline 15% methylation observed in initial Pacific oyster (*Crassostrea*

gigas) methylation studies [22], as well as examination of methylome responses to heat stress

[14] and diuron exposure [17]. In *C. virginica*, the mantle methylome was hypomethylated after

ocean acidification exposure [10], while there were no differences in global gonad methylation

patterns [12]. Experimental ocean acidification exposure yielded differential methylation

predominantly in gene bodies, but concentration in exons [11, 12] versus introns [8] varies.

Hyper- and hypomethylation occurs with equal frequency in response to ocean acidification.

This is consistent with diuron exposure [17], but not with heat stress [14]. Gene functions

enriched in differential methylation datasets were inconsistent between species and tissue types

(Table 1). While aminotransferase complex and biosynthetic processes were found to be

enriched in hypomethylated DML in *C. virginica* mantle tissue [10], there were no enriched

GOterms associated with genic DML in *C. virginica* gonad samples. In contrast, several

processes were enriched in methylation datasets for *C. hongkongensis*, including

acetoacetylco-A reductase activity, dehydrogenase activity, cellular response to pH, protein

xylosyltransferase activity, translation factor activity, RNA binding, and diacylglycerol kinase

activity in the mantle [8] and cytoskeletal and signal transduction, oxidative stress, metabolic

processes, and larval metamorphosis in larvae [11]. While considerable effort has been made to

understand molluscan methylation responses to ocean acidification, it is clear that there are

likely species- and tissue-specific responses.

In addition to being an important global aquaculture species, *C. gigas* has been used to further

our understanding of methylation in marine invertebrates [3, 7, 22–29]. Examination of diuron

exposure [17] and heat stress on different *C. gigas* phenotypes [14] reveal that environmental

conditions do influence the Pacific oyster methylome, and other oyster methylomes are

responsive to ocean acidification (Table 1). However, studies examining the influence of ocean

acidification on the *C. gigas* methylome are currently absent. Additionally, a recent study found

that female *C. gigas* exposure to low pH conditions, followed by four months in ambient pH

conditions prior to spawning, can still negatively impact larval survival 18 hours post-fertilization

[30]. Since maturation stage was not different between female *C. gigas* after low pH exposure,

this finding suggests environmental "memory" may be mediating the carryover effect [30].

Negative carryover effects were also found after parental exposure to low pH during

reproductive conditioning in the hard clam *Mercenaria mercenaria* and bay scallop *Argopectan*

irradians [31]. Although methylation differences associated with carryover effects have not been

reported in bivalves following low pH treatment, adult *C. gigas* exposure to the pesticide diuron

altered spat methylomes, demonstrating the potential for DNA methylation to influence

carryover effects in bivalves [17].

We sought to understand impacts of ocean acidification on the Pacific oyster gonad methylome

in order to not only understand DNA methylation as a potential mechanism for the observed

negative maternal carryover effect previously reported in Venkataraman et al. [30], but to also

examine the functional role of methylation in regulating biological responses to ocean

acidification. Female oyster gonad methylome sequencing revealed distinct methylation

changes associated with low pH exposure. These results provide the foundation for examining

phenotypic plasticity within a generation, as well as larval methylomes and potential

intergenerational effects.

Table 1. Studies exploring methylation responses to environmental stress in *Crassostrea* spp.

Results

Sequence Alignment

Whole genome bisulfite sequencing produced 1.38×10^9 total 150 bp paired-end reads for eight

total libraries. After quality trimming, 1.35×10^9 total paired reads remained (Supplementary

Table 1). Of the trimmed paired-end reads, 8.31×10^8 (61.7%) were uniquely aligned to the

bisulfite-converted _C. gigas_ genome with appended mitochondrial sequence (Supplementary

Table 1).

SNP Identification

C->T SNP variants were identified from WGBS data to remove loci that might have been

incorrectly characterized across individuals. A total of 13,234,183 unique SNPs were found in

individual or merged BAM files, including 300,278 C->T SNPs. These C->T SNPs were used to

exclude loci from downstream analyses (see DML Characterization and Enrichment Analysis).

Global Methylation Patterns

Figure 1. **Location of all CpGs with 5x coverage.** Percentage of highly methylated (≥ 50%),

moderately methylated (10-50%), and lowly methylated (≤ 10%) CpGs in various genome

features.

Combined data with 5x coverage threshold provided 11,238,223 CpG loci (84.8% of 13,246,547

total CpGs in *C. gigas* genome). After C->T SNP removal, 10,939,918 CpG loci (82.6% of total

C. gigas CpGs) were characterized. The majority of CpGs, 9,018,512 loci (82.4%), were lowly

methylated, followed by 966,655 highly methylated (8.8%) and 954,751 moderately methylated

(8.7%) CpGs. Highly methylated CpGs were found primarily in genic regions (903,933 CpGs,

93.5%), with 53,951 (6.0%) in exon UTR, 329,117 (36.4%) in CDS, and 523,386 (7.9%) in

introns (**Figure 1**). The distribution of highly methylated CpGs was significantly different than all

5x CpGs detected by WGBS (Supplementary Table 2). A Principal Component Analysis (PCA)

did not demonstrate any clear separation of samples by treatment (Supplementary Figure 1),

and consistently high Pairwise Pearson's correlation coefficients support the lack of global

percent methylation differences between pH treatments (Supplementary Figure 2).

DML Characterization

Initially, 1,599 CpG loci were identified as potential DML using a 50% methylation difference

threshold. Of these CpG loci, 315 (19.7%) overlapped with C->T SNPs, and thus were removed

from downstream analysis. A total of 1,284 DML were used in all remaining analyses, which

consisted of 654 (50.9%) hypomethylated and 630 (49.1%) hypermethylated DML (**Figure 2**).

The DML were distributed amongst the ten main chromosomes, as well as scaffolds not placed

in any chromosome (**Figure 3**, Supplementary Table 3).

Figure 2. **Percent methylation values for DML created using an euclidean distance matrix**.

Samples in low pH conditions are represented by black, and samples in ambient pH conditions

are represented by gray, with maturation stage along the bottom (0 = indeterminate, 3 = spawn-

ready mature female). Darker colors indicate higher percent methylation, and a density plot

depicts the distribution of percent methylation values for a panel. After excluding C->T SNPs,

1,284 DML were identified using a logistic regression, using a chi-squared test and 50%

methylation difference cut-off.

Figure 3. **Distribution of DML in main chromosomes**. Number of DML normalized by number

of CpG in each chromosome (bars) and number of genes (line) in each chromosome. Additional

DML were identified in scaffolds that were not mapped to any of the ten main linkage groups

(Supplementary Table 3).

The majority of DML (1,181; 92.0%) were located in genic regions (**Figure 4**), consisting of 859

unique genes. Most genes only contained one DML; however, several genes contained multiple

DML, ranging from 2-103 DML in one gene (Supplementary Table 4). The genic DML were

primarily located in introns (783; 61.0%), followed by CDS (442; 34.4%) and exon UTR (77;

6.0%). Putative promoter regions upstream of transcription start sites contained DML (10;

0.8%), but more were located in downstream flanking regions (61; 4.8%). The DML were also

found in transposable elements (434; 33.8%), lncRNA (9; 0.7%), and intergenic regions (38;

3.0%). The number of DML in all genome features differed significantly from all 5x CpGs

(Supplementary Table 5).

Figure 4. **Location of CpGs with 5x coverage and DML**. Percentage of 5x CpGs and DML

found in various genome features.

Enrichment Analysis

Ten biological processes and 4 cellular component GOterms were significantly enriched in

genes containing DML. A total of 339 genes with GOterm annotations containing 437 DML were

used for this analysis. The enriched biological process GOterms were involved in motility (ex.

cilium-dependent cell motility, cilium movement) and development (ex. insta larval or pupal

development, post-embryonic development) (Supplementary Table 6). Nine unique genes (37

associated transcripts) contained enriched biological process GOterms: *dynein heavy chain 5,*

axonemal, unconventional mysoin-VI, serine/threonine-protein kinase 36, helicase domino,

protein neuralized, cytoplasmic aconitate hydratase, serine-protein kinase ATM, cation-

independent mannose-6-phosphase receptor, uncharacterized LOC105324167. These genes

contained 16 DML, 13 of which were hypermethylated (Supplementary Table 6). Enriched

cellular component GOterms were involved in microtubule-associated complexes and various

acetyltransferase complexes (Supplementary Table 7). The six unique genes (25 associated

transcripts) that contained enriched cellular component GOterms were *dynein heavy chain 5,*

axonemal, cytoplasmic dynein 2 light intermediate chain-1-like, unconventional myosin-VI,

helicase domino, and kinesin-like protein KIF23 (Supplementary Table 7). Six of the seven DML

contained in these genes were hypermethylated (Supplementary Table 7).

Discussion

Our work examined environmentally-induced changes to the *C. gigas* reproductive tissue

methylome. We identified a total of 1,284 differentially methylated loci, showing that the

methylome is responsive to ocean acidification. We also found enriched biological processes

associated with genic DML, implying that low pH may impact regulation of distinct processes,

such as gonad development. Taken together, our findings suggest that environmentally-

responsive methylation may maintain gonad development in stressful conditions. Our findings

contribute to the growing body of work examining epigenetic responses in molluscan species,

and provide the foundation for future research examining intergenerational epigenetic

inheritance and its potential to mediate plastic responses to environmental stressors.

Functions of genes containing DML with enriched GOterms suggest role for

methylation regulation of gonad development

Enrichment of microtubule associated complex and cilium or flagellar motility GOterms imply

methylation may control microtubule movement in response to low pH. Motor proteins like

dynein heavy chain 5, axonemal, and _cytoplasmic dynein 2 light intermediate chain 1-like_ form

complexes to move flagella or cilia. Dyneins exhibit higher expression over the course of _C._

gigas spermatogenesis [32]. In female gonad tissue, hypermethylation may suppress

expression of dyneins to promote oogenesis. Indeed, lack of significant sex ratio and maturation

stage differences between low and ambient pH treatments suggests that oogenesis was

maintained in low pH conditions [30]. Studies examining molluscan somatic tissues have found

low pH impacts on dynein expression [33–35], with significantly lower dynein protein abundance

[33] or gene expression [34] in oysters exposed to ocean acidification. Similar to cytoplasmic

dyneins, _unconventional myosin-VI_ plays a role in organelle movement. Myosins generally

exhibit high expression in immature gonads, which reduces as the gonad matures [32, 36]. The

reduction in myosin expression may signify reorganization of female reproductive tissue to

accommodate growing oocytes. Hyper- and hypomethylation of unconventional myosin may

regulate cytoskeletal structure of cells to promote growth of reproductive over somatic tissue.

Dynein and myosin genes exhibit differential methylation in response to low pH in _C. virginica_

gonad as well [12]. Regulation of motility genes may be a conserved response to ocean

acidification, and methylation of these genes may promote homeostasis during gametogenesis.

Several genes with DML and enriched GOterms have links to stress response and gonad

development. In _C. gigas_, expression of cell cycle regulators _kinesin-like protein KIF23_ and

serine-protein kinase ATM changes in response to immune challenge or low pH and arsenic

exposure, respectively [37, 38]. Kinesin-like proteins have higher expression in mature _C. gigas_

gonads [32], and serine-protein kinase ATM is involved in razor clam *Sinonovacula constricta*

oogenesis [39]. Another kinase, *serine/threonine-protein kinase 36*, is involved in signaling

pathways. Several serine/threonine-protein kinases are highly expressed in mature *C. gigas*

female gonad, with certain homologues more expressed in females with mature gonad and

during sex determination and differentiation events [32, 36]. Serine/threonine-protein kinases

are also related to oocyte maturation in the king scallop *Pecten maximus* [40]. An investigation

of the *C. gigas* kinome found various serine/threonine-protein kinases present in eggs, including

serine/threonine-protein kinase 36, that were responsive to abiotic stressors [41].

Serine/threonine-protein kinase genes were also differentially methylated in *C. virginica* gonad

[12]. As a catalyst of the TCA cycle, *cytoplasmic aconitate hydratase* is upregulated during

spermatogenesis in *C. gigas* and downregulated during spermatogenesis in *P. maximus*, with

differing consequences for energy metabolism [42, 43]. Early examination of *C. virginia* found

cytoplasmic aconitate hydratase localized to eggs, with enzyme activity increasing during

embryogenesis [44]. Therefore, it's possible *cytoplasmic aconitate hydratase* may regulate

energy metabolism during oogenesis as well. Evidence of *cytoplasmic aconitate hydratase*

abundance changing in response to low pH has been documented in *C. gigas*, which implies

that low pH alters energy metabolism [38]. Energy metabolism would be important to regulate in

developing oocytes. *Helicase domino* regulates transcription through histone phosphorylation,

acetylation, and chromatin remodeling. Populations of the Olympia oyster (*Ostrea lurida*)

contain an outlier SNP loci in the *helicase domino* gene, which is implicated in immune or stress

response [45]. Upregulation of *helicase domino* was associated with oogenesis maintenance in

Acartia tonsa shrimp [46]. Taken together, hypermethylation of *kinesin-like protein 23*, *serine-*

protein kinase ATM, *cytoplasmic aconite hydratase*, and *helicase domino*, and hypomethylation

of *serine/threonine-protein kinase 36*, suggests that methylation may regulate gonad

development in stressful conditions, allowing the oyster to maintain homeostasis.

Similarities between differential methylation in gonads and larvae

Investigation of *C. hongkongensis* larvae exposed to ocean acidification revealed enrichment of

cytoskeletal and signal transduction processes in differentially methylated genes [11]. Genes

with DML in *C. gigas* gonad are enriched for similar processes, and have links to embryonic

development. Unconventional myosins, important cytoskeletal components, were found to be

important for fertilization, blastulation, and gastrulation in *S. purpuratus* [47]. Ocean acidification

drove decreased expression of another cytoskeletal component, kinesin-like protein, in *S.*

purpuratus larvae, which was related to measured reductions in calcification [48]. Expression of

signal transduction kinases found in embryos changed in response to various abiotic stressors

[41]. *Serine/threonine protein-kinase pim-3* gene was differentially methylated in *C.*

hongkongensis larvae [11], and this study found a DML in *serine/threonine-protein kinase 36* in

C. gigas gonad. Differential methylation in cytoskeleton and signal transduction genes in female

gonad may influence larval responses to low pH.

Enrichment of metabolic processes in *C. gigas* gonad and *C. hongkongensis* larval methylation

datasets [11] signify that methylation regulation of metabolism is important across life stages. In

S. purpuratus with a history of low pH exposure, day-old embryos exhibited higher expression of

cytoplasmic aconitate hydratase than embryos from populations without exposure history [49].

However, larvae from populations without exposure history had higher expression of

cytoplasmic aconitate hydratase after seven days [49]. Modulation of the TCA cycle through

cytoplasmic aconitate hydratase may be important for metabolic regulation in developing larvae,

and inheritance of hypermethylation within this gene could prime larval metabolism for low pH

exposure. *Cation-independent mannose-6-phosphate receptor* is a lysosomal enzyme

associated with post-embryonic development (GO:0009791). Expression of this gene was

upregulated during zebra mussel *Dreissena polymorpha* and shrimp *Marsupenaus japonicus*

immune challenges [50, 51]. Although no current research has explored this gene's role in larval

development or abiotic stress response, hypomethylation of *cation-independent mannose-6-*

phosphate receptor may impact lysosome metabolic activity and immune function during a

critical growth period. *Protein neuralized* is part of a group of ubiquitin ligases. Regulation of

ubiquitination is likely a conserved response to ocean acidification in oysters: elevated pCO_2

increased abundance of proteins involved in ubiquitination and decreased protein degradation in

C. gigas posterior gill lamellae [33], and upregulated ubiquitination gene expression in adult *C.*

virginica mantle [52]. Differential methylation of protein ubiquitination genes was also found in *C.*

virginica gonad [12] and *C. hongkongensis* larvae [11]. Hypermethylated DML within the *protein*

neuralized gene could regulate ubiquitination. Differential methylation in this gene may also

regulate cell division and larval growth, as upregulation of protein neuralized has been found to

suppress cell division and macromolecule synthesis in *Artemia sinica* during diapause [53].

Taken together, methylome similarities between *C. gigas* female gonad and *C. hongkongensis*

larvae suggest that differential methylation in these genes could also impact larval calcification,

cellular structure, and metabolism if patterns are inherited. Intergenerational methylation

inheritance has been demonstrated previously in *C. gigas*: adult exposure to the pesticide

diuron altered spat methylomes [17]. While we did not investigate *C. gigas* larval methylomes,

we did find that prior exposure of female oysters to low pH conditions negatively impacted larval

survival in ambient pH conditions 18 hours after fertilization [30]. It is possible that the

methylation patterns responsible for maintaining homeostasis in female gonad tissue were

maladaptive in larvae, especially when parent and larval environments were mismatched.

Considerations for marine invertebrate methylation research

The concentration of DML in genes is consistent with the literature examining molluscan

methylome responses to ocean acidification [8, 10–12], disease [16], pesticide exposure [17],

salinity (Johnson et al. 2021), and heat stress [14], as well as environmental responses in corals

[13, 18], urchins [15], and pteropods [9]. This differs from the vertebrate model of methylation, in

which CpG islands are found in promoter regions upstream of transcription start sites [1]. Similar

pathways exhibiting differential methylation — but not exact genes — between *C. gigas* gonad

(this study), *C. virginica* gonad [12] and mantle [10], and *C. hongkongensis* mantle [8] and

larvae [11] suggest tissue- and species-specific methylome responses to low pH. In *C. gigas*,

high levels of gene methylation are associated with increased gene expression and low

expression variability, implying that methylation leads to transcriptional stability [22, 25, 29].

Similarly, exposure to the pesticide diuron modified the *C. gigas* methylome such that DNA

methylation changes correlated with RNA abundance in a small group of highly methylated

genes [17], and increased methylation was also found to reduce transcriptional noise in

response to ocean acidification in corals [13]. While recent *Crassostrea* spp. studies have found

no connection between differentially methylated and differentially expressed genes after

exposure to low pH [8, 10], individual DML may play a role in shaping gene expression.

Comparison of a single gene with differing methylation levels between *C. gigas* mantle and male

gametes demonstrated that gene methylation was associated with alternative splicing and gene

expression [29]. Therefore, it is critical to examine how individual DML can influence gene

activity in response to environmental change.

The role of DML in shaping gene activity is likely dependent on their locations in genome

features. We found 442 and 941 DML in introns and CDS, respectively, with roughly half of the

genic DML being hypermethylated in treatment gonads. In *C. gigas*, exons included in the final

transcript and longer exons generally have higher methylation [29]. Genes with higher

methylation in exons or introns were highly expressed in *S. purpuratus*, and had a higher

likelihood of alternative splicing, alternative start site, or exon skipping in response to upwelling

conditions [54]. Differentially methylated positions in introns and CDS may have similar roles in

C. gigas response to low pH. Additionally, presence of multiple DML in a gene may have

conflicting effects on expression. Most genes examined in this study only contained one DML;

however, we found several genes with multiple DML, ranging from 2-103 in one gene. High

methylation in both exons and introns did not produce high expression levels in *S. purpuratus*,

suggesting increased methylation in both of these genome features act antagonistically to

impact gene expression [54]. Multiple DML in CDS and introns may influence *C. gigas*

expression similarly. Future work should examine how environmentally-influenced methylation

impacts gene expression by considering individual DML, not averaged methylation for a gene.

Confounding elements may have influenced gonad methylome responses to ocean acidification,

and our interpretation of those responses. Maturation stage is known to produce distinct

baseline methylation patterns in *C. gigas* (Zhang et al., 2018). While the overall maturation

stage was used as a covariate when identifying DML, it is possible that each sample methylome

contains information from cells with various maturation stages or cell types. Single-cell

sequencing efforts or investigations within a single maturation stage would clarify how these

sources of variation influenced our findings. We also identified several C->T SNPs that

overlapped with DML even though specimens were sibling oysters with similar genomic

backgrounds. Recent examination of *C. virginica* methylomes suggest methylation is largely

linked to genotype, emphasizing the need to account for genotypic differences in methylation

calls [55]. Our work demonstrates how SNP identification from WGBS can be used to exclude

loci with incorrectly characterized methylation due to genomic variation. While the lack of

annotated genes used in the enrichment analysis limits our interpretation of which processes

are most impacted by pH-sensitive methylation, removing SNP-DML overlaps prior to

enrichment analysis provides a more biologically accurate understanding of how methylation

may impact gonad growth.

Conclusion

Our work — which is the first to explore DNA methylation responses to ocean acidification in

Pacific oysters — found that low pH treatment altered the *C. gigas* female gonad methylome

after a seven-week exposure. We found that differential methylation occurs primarily in genic

regions, which is consistent with other studies examining oyster methylome responses to ocean

acidification. In conjunction with histology data from Venkataraman et al. [30], enriched

biological processes and cellular components in genes containing DML suggest that methylation

may be used to maintain gonad growth in low pH conditions. Our work expands on previous

molluscan methylation research by not only demonstrating a potential role for methylation in

response to ocean acidification, but also showing the species- and tissue-specific nature of

these responses.

As inheritance of environmentally-induced methylation — and any resulting phenotypic plasticity

— is contingent on epigenetic marks being present in the germline, withstanding chromatin

reorganization, and being packaged into gametes [1], the next step in this work is correlating

germline methylation with offspring methylation, and examining gene expression, alternative

splicing, chromatin regulation, and phenotype. Comparing parental and offspring phenotype will

allow researchers to assess whether parental experience and resulting epigenetic responses

are accurate, reliable cues for adaptive plasticity in offspring.

Methods

DNA Extraction and Library Preparation

Adult hatchery-raised *C. gigas* (average shell length = 117.46 ± 19.16 mm) were exposed to

either low pH (7.31 ± 0.02) or ambient pH (7.82 ± 0.02) conditions from February 15, 2017 to

April 8, 2017 at the Kenneth K. Chew Center for Shellfish Research and Restoration at the

National Oceanic and Atmospheric Administration Manchester Field Station (47° 34' 09.1" N

122° 33' 19.0" W, Manchester, WA). The experimental design, seawater chemistry analysis with

carbon chemistry parameters, and histological analysis are described in Venkataraman et al.

[30]. To determine how pH exposure altered maternal gonad methylation, four female (or

presumed female) oysters sampled at the end of the seven-week pH exposure were selected

for each treatment. DNA was extracted from histology blocks for these gonad tissue samples

using the PAXgene Tissue DNA Kit (Qiagen) with modifications specified below. For each

sample, up to 0.02 g of histology tissue embedded in paraffin was cut from the histology block

and placed in a 2 mL round-bottomed tube. The tissue was then further homogenized within the

tube. After adding 1 mL of xylene, each sample was vortexed for 20 seconds, incubated at room

temperature for three minutes, then centrifuged at maximum speed for three minutes. To

evaporate ethanol from samples after ethanol addition and vortexing steps, samples were

placed on a heat block at 37°C for 10-20 minutes.

The resuspended pellet was lysed using a TissueTearor at setting 1. Prior to lysis and between

samples, the TissueTearor was run in a 10% bleach solution for 20 seconds, followed by two

consecutive 20 second runs in DI water to clean the instrument. Lysates were transferred to

clean, labeled 1.5 mL microcentrifuge-safelock tubes. Proteinase K (20 µL) was added to each

sample, and pulse-vortexed for 15 seconds to mix. The sample-Proteinase K solution was

incubated 56°C for 60 minutes. Every ten minutes, the samples were briefly removed from the

heat block to vortex at maximum speed for one minute. After 60 minutes, RNase A (4 µL, 100

mg/mL) was added to each sample. Samples were vortexed at maximum speed for 20 seconds

to mix and incubated for two minutes at room temperature to obtain RNA-free genomic DNA

and reduce possible interference for downstream enzyme reactions. Samples were then kept at

80°C for 60 minutes, and vortexed at maximum speed for one minute in ten-minute intervals.

After the incubation, Buffer TD2 and ethanol (200 µL) were added to each sample, vortexing

thoroughly to mix.

DNA was isolated for each sample after lysis and incubation following manufacturer's

instructions. To elute DNA, samples were loaded onto the spin column with Buffer TD5 (50 µL)

and incubated at room temperature for five minutes to increase yield, then centrifuged at

maximum speed for one minute. A Qubit™ 3.0 and dsDNA BR Assay (Invitrogen) was used to

quantify sample yield using 1 µL of DNA for each sample. For two samples, DNA was further

concentrated using an ethanol precipitation.

Bisulfite conversion was performed with the Zymo-Seq WGBS Library Kit (Cat. #D5465) using

100 ng of genomic DNA following manufacturer's instructions. Samples were spiked with a

mixture of six unique double-stranded synthetic amplicons (180-200 bp) from the *Escherichia*

coli K12 genome. Since each amplicon represents a different percent methylation, the spike-in

was used to determine bisulfite conversion efficiency. Libraries were barcoded and pooled into a

single lane to generate 150 bp paired-end reads in a single Illumina NovaSeq flowcell.

Genome Information and Feature Tracks

Nuclear genome [56] (NCBI GenBank: GCA_902806645.1, RefSeq: GCF_902806645.1) and

mitochondrial genome sequences (NCBI Reference Sequence: NC_001276.1) were used in

downstream analysis. These sequences were combined for downstream alignment. The

`fuzznuc` function within EMBOSS was used to identify CG motifs in the combined genomes.

The RefSeq annotation was used to obtain genome feature tracks, and create additional tracks.

The annotation included a combination of Gnomon, RefSeq, cmsearch, and tRNAscan-SE

predictions. Gene, coding sequence, exon, and lncRNA tracks were extracted from the RefSeq

annotation. These tracks were then used to obtain additional genome feature information. To

create an intron track, the complement of the exon track was generated with BEDtools

`complementBed` v.2.26.0 to create a non-coding sequence track [57]. The overlap between the

genes and the non-coding sequences was found with `intersectBed` and defined as introns.

Similarly, untranslated regions of exons were obtained by subtracting coding sequences from

exon information with `subtractBed`. Flanking regions were defined as the 1000 bp upstream

or downstream of genes. Upstream flanks were generated by adding 1000 bp upstream of

genes, taking into account strand, with `flankBed`. Existing genes were removed from flanking

sequences using `subtractBed`. A similar process was performed to generate downstream

flanking regions. Intergenic regions were isolated by finding the complement of genes with

`complementBed`, then using `subtractBed` to remove any flanks. Transposable element

locations were obtained from RepeatMasker using the NCBI RefSeq annotation (RefSeq:

GCF_902806645.1) [58, 59]. The number of CG motifs in a given feature track was obtained

using `intersectBed`. All genome feature tracks are available in the associated Open Science

Framework repository (see *Availability of data and materials*).

Sequence Alignment

Reads were trimmed three times prior to alignment using TrimGalore! v.0.6.6 [60]. Known

Illumina adapter sequences were removed (`--illumina`), along with 10 bp from both the 5' (`-`

`-clip_R1 10` and `--clip_R2 10`) and 3' ends (`--three_prime_clip_R1 10` and `--`

`three_prime_clip_R2 10`) of 150 bp paired-end reads. Any remaining adapters were

removed in a second round of trimming. Finally, poly-G tails were trimmed out of samples by

manually specifying adapter sequences (`--adapter`

`GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG` and `--adapter2`

`GGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG`). Sequence quality was

assessed with FastQC v.0.11.9 [61] and MultiQC [62] after each round of trimming.

Trimmed samples were then aligned to the combined nuclear and mitochondrial genomes for *C.*

gigas [56]. The genome was bisulfite converted with Bowtie 2-2.3.4 (Linux x84_64 version [63])

using the `bismark_genome_preparation` function within Bismark v.0.22.3 [64]. Reads were

aligned to the bisulfite-converted genome. Non-directional input was specified (`--`

`non_directional`), and alignment scores could be no lower than -90 (`-score_min L,0,-`

`0.6`). Resultant BAM files were deduplicated (`--deduplicate_bismark`), and methylation

calls were extracted from deduplicated BAM files (`bismark_methylation_extractor`).

Deduplicated BAM files were also sorted and indexed for downstream applications using

SAMtools v.1.10 [65]. Using coverage files generated from methylation extraction, CpG

information was consolidated between strands to report 1-based chromosomal coordinates

(`coverage2cystosine`). Using `coverage2cytosine` coverage file output, genomic

coordinates for CpGs with at least 5x coverage across all samples were written to BEDgraphs.

Summary reports with alignment information for each sample were generated

(`bismark2report` and `bismark2summary`) and concatenated with MultiQC. Code for all

sequence alignment steps can be found in the associated Open Science Framework repository

(see *Availability of data and materials*).

SNP Identification

During bisulfite conversion, cytosines methylated in a CpG context are retained as cytosines,

while unmethylated cytosines are converted to uracils. These uracils are then converted to

thymines during PCR amplification. As C->T single nucleotide polymorphisms (SNPs) would be

interpreted by Bismark as an unmethylated cytosine, removing these loci is an important step to

ensure accurate methylation characterization. SNP variants across all samples and in

individuals were identified with `BS-Snper` [66]. This program takes into consideration nucleotide

identity on the opposite strand: a thymine paired with an adenine indicates a C->T SNP, while a

thymine paired with a guanine is a bisulfite-converted unmethylated cytosine. Sorted

deduplicated BAM files generated with Bismark v.0.22.3 and SAMtools v.1.10 [64, 65] were

merged (`samtools merge`) to identify SNPs. Variants were also identified in individual sorted

deduplicated BAM files. Default `BS-Snper` settings were used, except for a minimum coverage

of 5x to be consistent with methylation analyses. The C->T SNPs were used to exclude loci of

interest.

Global Methylation Characterization

Global methylation patterns were characterized by averaging DNA methylation across all

samples using loci with at least 5x coverage that did not overlap with C->T SNPs. A CpG locus

was considered highly methylated if average percent methylation was greater than or equal to

50%, moderately methylated if percent methylation was between 10-50%, and lowly methylated

if percent methylation was less than or equal to 10%. The genomic location of highly,

moderately, and lowly methylated CpGs in relation to UTR, CDS, introns, up- and downstream

flanking regions, transposable elements, and intergenic regions were determined by using

`intersectBed`. We tested the null hypothesis that there was no association between the

genomic location of CpG loci and methylation status (all CpGs versus highly methylated CpGs)

with a chi-squared contingency test (`chisq.test` in R Version 3.5.3 [67]).

Identification of Differentially Methylated Loci

Identification of differentially methylated loci, or DML, was conducted with `methylKit`

`v.1.17.4` [68] in R Version 3.5.3 [67]. Prior to DML identification, data were imported and

processed. Data with at least 2x coverage were imported using `methRead` from merged CpG

coverage files produced by `coverage2cytosine`. Imported data were filtered again

(`filterByCoverage`) to require 5x coverage for each CpG locus (`lo.count = 5`) in each

sample. Potential PCR duplicates in each sample were also removed by excluding CpG loci in

the 99.9th percentile of coverage (`high.perc = 99.9`). Once filtered, data were normalized

across samples (`normalizeCoverage`) to avoid over-sampling reads from one sample during

downstream statistical analysis. Histograms of percent methylation (`getMethylationStats`)

and CpG coverage (`getCoverageStats`) were used to confirm normalization. After filtering

and normalization, data at each CpG locus with 5x were combined (`unite`) to create a CpG

background, ensuring that each locus had at least 5x coverage in each sample.

Methylation differences were then identified for *C. gigas* gonad samples. A logistic regression

(`calculateDiffMeth`) modeled the log odds ratio based on the proportion of methylation at

each CpG locus. A covariate matrix with gonad stage information (`covariates`) was applied to

the model for the *C. gigas* data to account for varying maturation stages in samples.

Additionally, an overdispersion correction with a chi-squared test (`overdispersion = "MN"`,

`test = "Chisq"`) was applied.

Differentially methylated loci, or DML, were defined as CpG dinucleotides that did not overlap

with C->T SNPs with at least a 50% methylation difference between pH treatments, and a q-

value < 0.01 based on correction for false discovery rate with the SLIM method [69].

Hypermethylated DML were defined as those with significantly higher percent methylation in low

pH samples, and hypomethylated DML with significantly lower percent methylation in low pH

samples. BEDfiles with DML locations were created and viewed with the Integrative Genomics

Viewer (IGV) version 2.9.2 [70].

DML Characterization

The location of DML was characterized in relation to various genome feature tracks. Presence

of DML in UTR, CDS, introns, up- and downstream flanking regions, transposable elements,

and intergenic regions was discerned using `intersectBed`. A chi-squared contingency test

was used to test the null hypothesis of no association between genomic location and differential

methylation by comparing the genomic location of all CpGs with 5x data and DML. Additionally,

the number of DML present in each chromosome was quantified to see if DML were distributed

uniformly across the genome. Overlaps between DML and SNPs were identified to determine if

DML were solely attributed to treatment differences, or if genetic differences contributed to

differential methylation.

Enrichment Analysis

To determine if genes containing DML were associated with overrepresented biological

processes, cellular component, or molecular function gene ontologies (GO), functional

enrichment analysis was performed. Prior to this analysis, the *C. gigas* genome was annotated.

A `blastx` alignment was performed against the Uniprot-SwissProt database (accessed June

01, 2021) to get Uniprot Accession information for each transcript [71]. The transcript nucleotide

sequences were extracted from the genome

(GCF_902806645.1_cgigas_uk_roslin_v1_rna_from_genomic.fna available in the [NCBI

Annotation Release 102](#)). Transcript IDs from the `blastx` output were matched with GO terms

from the Uniprot-Swissprot database using Uniprot Accession codes. These transcript IDs and

corresponding GO terms were then used to create a gene ID-to-GO term database

(`geneID2GO`) for manual GO term annotation. The transcript IDs were filtered for genes that

contained CpGs with at least 5x coverage in all samples, as the same parameters were used to

generate the CpG background for differential methylation analysis. Each line of the database

contained transcript ID in one column, and all corresponding GO terms in another column.

Gene enrichment analysis was conducted with `topGO` v.2.34.0 in R [72]. The `geneID2GO`

database was used as the gene universe for enrichment, while transcript IDs for genes with

DML were used as the genes of interest. These transcript IDs were filtered such that they did

not include any transcripts associated with DML-SNP overlaps. A `topGO` object was generated

for each DML list and GO category (biological process, cellular component, or molecular

function), with GO term annotation performed using the `geneID2GO` database. A Fisher's exact

test was used to identify GO terms in each DML list significantly enriched with respect to the

gene background (P-value < 0.01). Keeping with `topGO` default settings, we did not correct for

multiple comparisons [72]. Enriched GOterms were clustered by semantic similarity using

`simplifyEnrichment` v.1.2.0 default settings [73]. A semantic similarity matrix was created

from enriched GO IDs using the "Relevance" similarity measure and GO tree topology

(`GO_similarity`). The semantic similarity matrix was clustered using the binary cut method,

and visualized as a word cloud alongside a heatmap of semantic similarity (`simplifyGO`).

Finally, the enriched GOterms and cluster information were appended to a list of genic DML and

annotations to understand which genes with DML also had enriched GOterms.

List of Abbreviations

WGBS: whole genome bisulfite sequencing

DML: differentially methylated locus/loci

SNP: single nucleotide polymorphism

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data, genome feature tracks, scripts, and a supplementary materials list are available in the

Oyster Gonad Methylation repository, doi.org/10.17605/OSF.IO/YGCTB. All raw data can be

accessed at the NCBI Sequence Read Archive under BioProject accession number

PRJNA806944 (https://www.ncbi.nlm.nih.gov/bioproject/806944).

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by National Science Foundation award 1634167 to SBR. The Hall

Conservation Genetics Research Fund (YRV) supported sequencing for this project.

Authors' contributions

YRV conceived the experimental design and conducted the pH exposure experiment with

assistance from SBR. YRV extracted DNA for sequencing and analyzed sequence data with

SJW and SBR. YRV wrote the initial manuscript with input from SJW and SBR. All authors

reviewed and approved the final manuscript.

Acknowledgements

References

1. Eirin-Lopez JM, Putnam HM. Marine Environmental Epigenetics. Ann Rev Mar Sci. 2018.
https://doi.org/10.1146/annurev-marine-010318-095114.

2. Hofmann GE. Ecological Epigenetics in Marine Metazoans. Frontiers in Marine Science.
2017;4.

3. Gavery MR, Roberts SB. A context dependent role for DNA methylation in bivalves. Brief
Funct Genomics. 2014;13:217–22.

4. Huang X, Li S, Ni P, Gao Y, Jiang B, Zhou Z, et al. Rapid response to changing environments
during biological invasions: DNA methylation perspectives. Mol Ecol. 2017;26:6621–33.

5. Ni P, Murphy KJ, Wyeth RC, Bishop CD, Li S, Zhan A. Significant population methylation
divergence and local environmental influence in an invasive ascidian Ciona intestinalis at fine
geographical scales. Mar Biol. 2019;166:143.

6. van Oppen MJH, Oliver JK, Putnam HM, Gates RD. Building coral reef resilience through
assisted evolution. Proc Natl Acad Sci U S A. 2015;112:2307–13.

7. Gavery MR, Roberts SB. Epigenetic considerations in aquaculture. PeerJ. 2017;5:e4147.

8. Chandra Rajan K, Yuan M, Yu Z, Roberts SB, Thiyagarajan V. Oyster biomineralisation under
ocean acidification: from genes to shell. Glob Chang Biol. 2021.
https://doi.org/10.1111/gcb.15675.

9. Bogan SN, Johnson KM, Hofmann GE. Changes in Genome-Wide Methylation and Gene
Expression in Response to Future pCO2 Extremes in the Antarctic Pteropod Limacina helicina
antarctica. Frontiers in Marine Science. 2020;6:788.

10. Downey-Wall AM, Cameron LP, Ford BM, McNally EM, Venkataraman YR, Roberts SB, et
al. Ocean Acidification Induces Subtle Shifts in Gene Expression and DNA Methylation in
Mantle Tissue of the Eastern Oyster (Crassostrea virginica). Frontiers in Marine Science.
2020;7:828.

11. Lim Y-K, Cheung K, Dang X, Roberts SB, Wang X, Thiyagarajan V. DNA methylation
changes in response to ocean acidification at the time of larval metamorphosis in the edible
oyster, Crassostrea hongkongensis. Mar Environ Res. 2020;163:105214.

12. Venkataraman YR, Downey-Wall AM, Ries J, Westfield I, White SJ, Roberts SB, et al.
General DNA methylation patterns and environmentally-induced differential methylation in the
eastern oyster (Crassostrea virginica). Frontiers in Marine Science. 2020.
https://doi.org/10.3389/fmars.2020.00225.

13. Liew YJ, Zoccola D, Li Y, Tambutté E, Venn AA, Michell CT. Epigenome-associated
phenotypic acclimatization to ocean acidification in a reef-building coral. Science Advances.
2018.

14. Arredondo-Espinoza R, Ibarra AM, Roberts SB, Sicard-Gonzalez MT, Escobedo-Fregoso C.
Differentially methylated gene regions between resistant and susceptible heat-phenotypes of
the Pacific oyster Crassostrea gigas. Aquaculture. 2021;543:736923.

15. Strader ME, Wong JM, Kozal LC, Leach TS, Hofmann GE. Parental environments alter DNA
methylation in offspring of the purple sea urchin, Strongylocentrotus purpuratus. J Exp Mar Bio
Ecol. 2019;517:54–64.

16. Johnson KM, Sirovy KA, Casas SM, La Peyre JF, Kelly MW. Characterizing the Epigenetic
and Transcriptomic Responses to Perkinsus marinus Infection in the Eastern Oyster
Crassostrea virginica. Frontiers in Marine Science. 2020;7.

17. Rondon R, Grunau C, Fallet M, Charlemagne N, Sussarellu R, Chaparro C, et al. Effects of
a parental exposure to diuron on Pacific oyster spat methylome. Environ Epigenet. 2017;3.

18. Liew YJ, Howells EJ, Wang X, Michell CT, Burt JA, Idaghdour Y, et al. Intergenerational
epigenetic inheritance in reef-building corals. bioRxiv. 2018;:269076.

19. Melzner F, Mark FC, Seibel BA, Tomanek L. Ocean Acidification and Coastal Marine
Invertebrates: Tracking CO2 Effects from Seawater to the Cell. Ann Rev Mar Sci. 2019.
https://doi.org/10.1146/annurev-marine-010419-010658.

20. Parker LM, Ross PM, O'Connor WA, Pörtner HO, Scanes E, Wright JM. Predicting the
response of molluscs to the impact of ocean acidification. Biology . 2013;2:651–92.

21. Kroeker KJ, Kordas RL, Crim RN, Singh GG. Meta-analysis reveals negative yet variable
effects of ocean acidification on marine organisms. Ecol Lett. 2010;13:1419–34.

22. Gavery MR, Roberts SB. Predominant intragenic methylation is associated with gene
expression characteristics in a bivalve mollusc. PeerJ. 2013;1:e215.

23. Roberts SB, Gavery MR. Is there a relationship between DNA methylation and phenotypic
plasticity in invertebrates? Front Physiol. 2012;2.

24. Riviere G, Wu G-C, Fellous A, Goux D, Sourdaine P, Favrel P. DNA methylation is crucial
for the early development in the Oyster C. gigas. Mar Biotechnol . 2013;15:739–53.

25. Olson CE, Roberts SB. Genome-wide profiling of DNA methylation and gene expression in
Crassostrea gigas male gametes. Front Physiol. 2014;5:224.

26. Olson CE, Roberts SB. Indication of family-specific DNA methylation patterns in developing
oysters. bioRxiv. 2015;:012831.

27. Saint-Carlier E, Riviere G. Regulation ofHoxorthologues in the oysterCrassostrea
gigasevidences a functional role for promoter DNA methylation in an invertebrate. FEBS Letters.
2015;589:1459–66.

28. Riviere G, He Y, Tecchio S, Crowell E, Gras M, Sourdaine P, et al. Dynamics of DNA
methylomes underlie oyster development. PLoS Genet. 2017;13:e1006807.

29. Song K, Li L, Zhang G. The association between DNA methylation and exon expression in
the Pacific oyster Crassostrea gigas. PLoS One. 2017;12:e0185224.

30. Venkataraman YR, Spencer LH, Roberts SB. Larval Response to Parental Low pH
Exposure in the Pacific Oyster Crassostrea gigas. Journal of Shellfish Research. 2019;38:743.

31. Griffith AW, Gobler CJ. Transgenerational exposure of North Atlantic bivalves to ocean
acidification renders offspring more vulnerable to low pH and additional stressors. Sci Rep.
2017;7:11394.

32. Dheilly NM, Lelong C, Huvet A, Kellner K, Dubos M-P, Riviere G, et al. Gametogenesis in
the Pacific oyster Crassostrea gigas: a microarrays-based analysis identifies sex and stage
specific genes. PLoS One. 2012;7:e36353.

33. Timmins-Schiffman E, Coffey WD, Hua W, Nunn BL, Dickinson GH, Roberts SB. Shotgun
proteomics reveals physiological response to ocean acidification in Crassostrea gigas. BMC
Genomics. 2014;15:951.

34. Goncalves P, Jones DB, Thompson EL, Parker LM, Ross PM, Raftos DA. Transcriptomic
profiling of adaptive responses to ocean acidification. Mol Ecol. 2017;26:5974–88.

35. De Wit P, Durland E, Ventura A, Langdon CJ. Gene expression correlated with delay in shell
formation in larval Pacific oysters (Crassostrea gigas) exposed to experimental ocean
acidification provides insights into shell formation mechanisms. BMC Genomics. 2018;19:160.

36. Broquard C, Saowaros S-A, Lepoittevin M, Degremont L, Lamy J-B, Morga B, et al. Gonadal
transcriptomes associated with sex phenotypes provide potential male and female candidate
genes of sex determination or early differentiation in Crassostrea gigas, a sequential
hermaphrodite mollusc. BMC Genomics. 2021;22:609.

37. de Lorgeril J, Zenagui R, Rosa RD, Piquemal D, Bachère E. Whole transcriptome profiling of
successful immune response to Vibrio infections in the oyster Crassostrea gigas by digital gene
expression analysis. PLoS One. 2011;6:e23142.

38. Moreira A, Figueira E, Mestre NC, Schrama D, Soares AMVM, Freitas R, et al. Impacts of
the combined exposure to seawater acidification and arsenic on the proteome of Crassostrea
angulata and Crassostrea gigas. Aquat Toxicol. 2018;203:117–29.

39. Yao H, Lin Z, Dong Y, Kong X, He L, Xue L. Gonad Transcriptome Analysis of the Razor
Clam (Sinonovacula constricta) Revealed Potential Sex-Related Genes. Frontiers in Marine
Science. 2021;8:1246.

40. Pauletto M, Milan M, Huvet A, Corporeau C, Suquet M, Planas JV, et al. Transcriptomic
features of Pecten maximus oocyte quality and maturation. PLoS One. 2017;12:e0172805.

41. Epelboin Y, Quintric L, Guévélou E, Boudry P, Pichereau V, Corporeau C. The Kinome of
Pacific Oyster Crassostrea gigas, Its Expression during Development and in Response to
Environmental Factors. PLoS One. 2016;11:e0155435.

42. Kingtong S, Kellner K, Bernay B, Goux D, Sourdaine P, Berthelin CH. Proteomic
identification of protein associated to mature spermatozoa in the Pacific oyster Crassostrea
gigas. J Proteomics. 2013;82:81–91.

43. Boonmee A, Heude Berthelin C, Kingtong S, Pauletto M, Bernay B, Adeline B, et al.
Differential protein expression during sperm maturation and capacitation in an hermaphroditic
bivalve, Pecten maximus (Linnaeus, 1758). J Molluscan Stud. 2016;82:575–84.

44. Black RE. THE CONCENTRATIONS OF SOME ENZYMES OF THE CITRIC ACID CYCLE
AND ELECTRON TRANSPORT SYSTEM IN THE LARGE GRANULE FRACTION OF EGGS
AND TROCHOPHORES OF THE OYSTER, CRASSOSTREA VIRGINICA. Biol Bull.
1962;123:71–9.

45. Silliman K. Population structure, genetic connectivity, and adaptation in the Olympia oyster
(Ostrea lurida) along the west coast of North America. Evol Appl. 2019;12:923–39.

46. Nilsson B, Jepsen PM, Bucklin A, Hansen BW. Environmental stress responses and
experimental handling artifacts of a model organism, the copepod Acartia tonsa (Dana). Front
Mar Sci. 2018;5.

47. Sirotkin V, Seipel S, Krendel M, Bonder EM. Characterization of sea urchin unconventional
myosins and analysis of their patterns of expression during early embryogenesis. Mol Reprod
Dev. 2000;57:111–26.

48. Padilla-Gamiño JL, Kelly MW, Evans TG, Hofmann GE. Temperature and CO2 additively
regulate physiology, morphology and genomic responses of larval sea urchins,
Strongylocentrotus purpuratus. Proceedings of the Royal Society B: Biological Sciences.
2013;280:20130155.

49. Evans TG, Pespeni MH, Hofmann GE, Palumbi SR, Sanford E. Transcriptomic responses to
seawater acidification among sea urchin populations inhabiting a natural pH mosaic. Mol Ecol.
2017;26:2257–75.

50. Zhang K-Y, Yuan W-J, Xu J-D, Wang J-X. Cation-dependent mannose-6-phosphate
receptor functions as a pattern recognition receptor in anti-bacterial immunity of Marsupenaeus
japonicus. Dev Comp Immunol. 2018;89:122–30.

51. Leprêtre M, Almunia C, Armengaud J, Salvador A, Geffard A, Palos-Ladeiro M. The immune
system of the freshwater zebra mussel, Dreissena polymorpha, decrypted by proteogenomics of
hemocytes and plasma compartments. J Proteomics. 2019;202:103366.

52. Tomanek L, Zuzow MJ, Ivanina AV, Beniash E, Sokolova IM. Proteomic response to
elevated PCO2 level in eastern oysters, Crassostrea virginica: evidence for oxidative stress. J
Exp Biol. 2011;214 Pt 11:1836–44.

53. Jiang G, Xu X, Jing Y, Wang R, Fan T. Comparative studies on sorting cells from Artemia
sinica at different developmental stages for in vitro cell culture. In Vitro Cell Dev Biol Anim.
2011;47:341–5.

54. Bogan SN, Strader ME, Hofmann GE. Gene regulation by DNA methylation is contingent on
chromatin accessibility during transgenerational plasticity in the purple sea urchin. bioRxiv.
2021;:2021.09.23.461091.

55. Johnson KM, Sirovy KA, Kelly MW. Differential DNA methylation across environments has
no effect on gene expression in the eastern oyster. J Anim Ecol. 2021.
https://doi.org/10.1111/1365-2656.13645.

56. Peñaloza C, Gutierrez AP, Eöry L, Wang S, Guo X, Archibald AL, et al. A chromosome-level
genome assembly for the Pacific oyster Crassostrea gigas. Gigascience. 2021;10.

57. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26:841–2.

58. Tarailo-Graovac M, Chen N. Using RepeatMasker to Identify Repetitive Elements in
Genomic Sequences. Current Protocols in Bioinformatics. 2009.

59. Smit AFA, Hubley R, Green P. 2013--2015. RepeatMasker Open-4.0. 2013.

60. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011;17:10–2.

61. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010.

62. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics. 2016;32:3047–8.

63. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9:357–9.

64. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics. 2011;27:1571–2.

65. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

66. Gao S, Zou D, Mao L, Liu H, Song P, Chen Y, et al. BS-SNPer: SNP calling in bisulfite-seq
data. Bioinformatics. 2015;31:4006–8.

67. R Core Team. R: A language and environment for statistical computing. 2019.

68. Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, et al.
methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation
profiles. Genome Biol. 2012;13:R87.

69. Wang H-Q, Tuominen LK, Tsai C-J. SLIM: a sliding linear model for estimating the
proportion of true null hypotheses in datasets with dependence structures. Bioinformatics.
2011;27:225–31.

70. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. Brief Bioinform. 2013;14:178–92.

71. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.
2019;47:D506–15.

72. Alexa A, Rahnenfuhrer J, Others. topGO: enrichment analysis for gene ontology. R package
version. 2010;2:2010.

73. Gu Z, Hübschmann D. simplifyEnrichment: an R/Bioconductor package for Clustering and
Visualizing Functional Enrichment Results. bioRxiv. 2021;:2020.10.27.312116.