Title: Composition and functionality of bacterioplankton communities in marine coastal

zones adjacent to finfish aquaculture

Author names and affiliations:

Da Silva, R.R.P.[1], White, C.A.[1], Bowman, J.P.[2] & Ross, D.J.[1]

1 Institute for Marine and Antarctic Studies (IMAS), Nubeena Crescent, Taroona, Tasmania

7053, Australia

2 Tasmanian Institute of Agriculture (TIA), University of Tasmania, Hobart, Tasmania 7001,

Australia

Corresponding author:

E-mail address: ricardo.silva@utas.edu.au (Ricardo R P Silva)

Abstract:

Finfish aquaculture is one of the fastest-growing primary industries globally and is increasingly common

in coastal ecosystems. Bacterioplankton is ubiquitous in marine environment and respond rapidly to

environmental changes. However, little is known about the effect of the aquaculture in the

bacterioplankton community. This study aims to examine aquaculture effects in the composition

and functional profiles of the bacterioplankton community using amplicon sequencing along a distance

gradient from two finfish leases in a marine embayment. Our results revealed natural stratification in

bacterioplankton strongly associated to NOx, conductivity, salinity, temperature and PO_4. Among the

differentially abundant bacteria in leases, we found members associated with nutrient enrichment and

aquaculture activities. Abundant predicted functions near leases were assigned to organic matter

degradation, fermentation, and antibiotic resistance. This study provides a first effort to describe

changes in the bacterioplankton community composition and function due to finfish aquaculture in a

semi-enclosed and highly stratified embayment.

1. Introduction

Aquaculture is one of the fastest-growing primary industries worldwide and is increasingly

common in coastal ecosystems. Aquaculture of carnivorous fish species, such as Atlantic

salmon (*Salmo salar*) is feed-additive, with a significant portion of feed flowing to the

environment as metabolic waste products, faeces and waste feed (Navarro et al., 2008, Burridge

et al., 2010, Zhang et al., 2015). As industries expand there is increasing concern regarding

potential effects that this waste can cause to surrounding environments. Key to sustainable

aquaculture operations in marine coastal systems is a better understanding around how pelagic

environments assimilate and mitigate additional nutrient loadings.(Asami et al., 2005, á Norði

et al., 2011, Belias et al., 2003, Elizondo-Patrone et al., 2015). There are two main pathways

of enrichment in pelagic environments, being the solid and dissolved waste streams (Bouwman

et al., 2013, Tsukamoto et al., 2008, Wang et al., 2012, Amirkolaie, 2011). The solid waste

stream consists of waste feed and faecal material and is available to pelagic primary consumers

as the solid debris settles out of the water column through sedimentation (Macleod et al. 2004).

In contrast, metabolic waste products, in particular bioactive inorganic nitrogen compounds

such as ammonia and nitrate, are released into the water column as dissolved nutrients and

available for assimilation by planktonic organisms (Wang et al. 2012).

Bacterial communities are known to be ubiquitous in the marine environment making up a

large fraction of the planktonic ecosystem. Bacterioplankton communities (i.e., bacterial

communities of the water column) play an important role in nutrient cycling and

biogeochemical processes (Azam et al., 1983, Pomeroy et al., 2007, Fuhrman and Steele, 2008).

Bacterioplankton are highly dynamic, responding to natural or anthropogenic changes (Newell

et al., 2013, Bowman and Ducklow, 2015, Yu et al., 2014, Reji et al., 2020). Previous studies

on sediments beneath aquaculture operations have identified changes in the bacterial

community due to nutrient enrichment (McCaig et al., 1999, Bissett et al., 2006, Dowle et al.,

2015, Rubio-Portillo et al., 2019). However, bacterioplankton community change across

enrichment gradients caused by aquaculture activities have yet to be fully defined. Olsen et al.

(2017) studying the response of the bacterial community to nutrient input in surface waters

adjacent to fish cages in southern Chile found an impact on community structure, although

changes in diversity were not observed. While this suggests stability and functional resilience

in bacterioplankton communities, there are very few other studies that examined response,

particularly across different receiving environments (Yoshikawa and Eguchi, 2013, Elizondo-

Patrone et al., 2015, Girvan et al., 2005). Aquatic ecosystems are complex, with cumulative or

synergetic effects frequently acting in a multifactorial way. Natural variation of the system and

management of aquaculture inputs are key factors to consider when assessing the impact of

waste on the pelagic environment (Wu, 1995, Buschmann et al., 2006). Therefore, it is

important to study the responses of bacterial community to environmental variations to enhance

our ability to understand changes caused by both natural and anthropogenic environmental

changes.

The advancement of molecular-based techniques has greatly advanced the field of microbial

ecology (Pace, 1997, Gilbert et al., 2010, Caporaso et al., 2011, Yarza et al., 2014, Martínez-

Porchas and Vargas-Albores, 2017, DeLong et al., 1993). The high-throughput sequencing of

16S rRNA has been used as a powerful tool to study the composition of bacterial community

from a variety of habitats (Caporaso et al., 2011, Logares et al., 2014, Morelan, 2019, Han et

al., 2017). Additionally, the availability of genomic data now allows inference of functional

capabilities based on taxon identification and relative abundances (Langille et al., 2013, Sevigny et al., 2019, Asshauer et al., 2015, Nagpal et al., 2016, Ortiz-Estrada et al., 2019). Previous studies using of 16S rRNA gene amplicon sequencing have showed a response of the microbial community and function to aquaculture activities in various aquatic systems (Keeley et al., 2018, Kolda et al., 2020, Ortiz-Estrada et al., 2019, Martins et al., 2018, Moncada et al., 2019, Frühe et al., 2021). Cordier (2020) could detect bacterial compositional and functional changes in the sediments close to aquaculture, where molecular-based techniques performed similarly to traditional microscopy-based techniques in predicting organic enrichment. However, natural local characteristics such as water column stratification and natural variation in environmental factors have major effects on the bacterial community structure and function (Labbate et al., 2016, Martiny et al., 2006, Zorz et al., 2019, Da Silva et al., 2021). The strong influence of the environment can make comparisons between studies undertaken in different locations difficult. However, the more we understand the natural variability across different marine systems, the better placed we are to understand common responses in bacteria community different perturbations, both natural and anthropogenic.

Macquarie Harbour is highly stratified fjord-like system with restricted water exchange with the ocean. The water column is characterized by a brackish tannin-rich water layer on top of more saline marine waters. The major freshwater source flows from the southern end carrying high amounts of organic matter with mean monthly flows varying according to the season ($<$ 100 m^3 sec^{-1} in summer and early autumn, and 500 m^3 sec^{-1} late autumn, winter, and spring) (Hartstein et al., 2019, Carpenter et al., 1991, Teasdale et al., 2003). Marine water enters through a narrow and shallow inlet in the north, allowing the renewal of the dissolved oxygen (DO) in the deeper waters. Marine incursions into the Harbour are likely driven by forces such as tide, wind, atmospheric pressure and freshwater inputs (Cresswell et al., 1989, Hartstein et

al., 2019). The Harbour has a history of anthropogenic impacts including mining, damming

and aquaculture (King and Tyler, 1982, Crawford et al., 2003, Koehnken, 2005, Kirkpatrick et

al., 2019). Pelagic DO concentrations naturally drop with increasing depth reaching around 4

to 6 mg O_2 L^{-1} at bottom layers. However, a recent report observed a considerable decline, with

levels now 2 to 4 mg O_2 L^{-1} or lower below 25 m in depth (Ross et al., 2021). This decline is

thought to be related to a reduction in the frequency and scalability of DO recharge along with

a high biological oxygen demand due to input of organic matter from fish farms. Additionally,

previous studies have suggested that pelagic microbial oxygen is likely to be an important

determinant of the decline in pelagic oxygen within the Harbour (Revill et al., 2016, Ross et

al., 2021, Da Silva et al., 2021). However, the impact of fish farm inputs on the

bacterioplankton community remain unclear.

For this reason, the main goal of this study is to better characterize the relationship between

the bacterioplankton community and proximity to aquaculture within Macquarie Harbour. This

will provide a starting point for further studies to examine the effects of aquaculture in the

bacterioplankton community, especially in a region with restricted water exchange. The study

aims (i) to examine the composition and distribution of bacterioplankton community and (ii)

to assess the influence of aquaculture activity on microbial community structure and

distribution. Based on previous studies and taking into account the hydrodynamics of the

Harbour, we hypothesize that the effects of aquaculture in the bacterioplankton community will

be localized.

2. Material and methods

Location and sample collection

In December of 2019, sampling was conducted at two fish farm leases ("Lease 1" and "Lease

2") and control stations ("Control") in Macquarie Harbour (42°18'13.8"S; 145°22'22.7"E),

Tasmania, Australia (Fig. 1). At each of the leases, samples were collected at sites on replicate

transects radiating from a fully stocked Atlantic salmon cage (*Salmo salar*) fed a pellet diet

using 10 L Niskin bottles (Tables S1 and S2). Seawater was collected from the surface (0-4 m),

middle waters (10 – 15 m) and bottom waters (20 – 37 m) at each site except for the site inside

the cage. Each transect (Lease 1: n = 2; Lease 2: n=3) included 4 distances (inside the cage +

3 distances) at the surface, and 3 distances at middle and bottom layers. Sampling over distance

was carried out by letting the boat drift along the direction of the current. Three control stations

(>1500 m away from cages) were sampled at the three depths mentioned above. Genetic

material was collected on 0.22 µM polyethersulfone Sterivex™ filter cartridges (Millipore,

Darmstadt, Germany) using a multichannel peristaltic pump. Each sample was stored at -80°C

until DNA extraction. Haversine's formula was used to calculate the distance between sample

points by use of latitude and longitude data (Sinnott, 1984).

Fig. 1 – Map on the left displays the locations of sites sampled within this study. Green points are control sites
and squares are leases located in the Macquarie Harbour. Sampled leases are the blue (Lease 1) and orange (Lease
2) ones. The two maps on the right show the sampled leases where the red points are the cages and the grey points
are the samples collected along a distance gradient.
145

Environmental variables

Physiochemical parameters including temperature ($^\circ$C), dissolved oxygen (DO, mg L^{-1}),

turbidity (NTU) and salinity were measured *in situ* at each site using a YSI 6600 V2 Multi

Parameter Water Quality Sonde with a YSI 650 MDS logger. Nutrient analyses were assayed

using a Bran + Luebbe AA3 HR segmented flow analyzer following standard

spectrophotometric methods (Grasshoff et al., 2009). Detection limits for NOx (NO$_3^-$ + NO$_2^-$),

nitrite (NO$_2^-$) were 0.015 µmol/L, for phosphate (PO$_4^{-3}$) to 0.01 µmol/L, and for ammonium

(NH$_4^+$) to 0.004 µmol/L.

DNA extraction and sequencing

DNA was extracted using a modified phenol:chloroform:isoamyl based DNA extraction

protocol of the DNeasy PowerWater SterivexTM Kit (MoBio-Qiagen,vHilden, Germany) as

described in Appleyard et al. (2013). Amplicon sequencing targeted the V1 to V3 region of the

bacterial 16S rRNA gene (27F–519R) (Lane, 1991, Lane et al., 1985). Libraries were

separately generated and sequenced for each sample at the Ramaciotti Centre for Genomics

(University of New South Wales, Sydney) using 300 bp paired-end sequencing on the MiSeq

platform (Illumina Inc., USA). Raw sequence reads and metadata are publicly available at the

Sequence Read Archive under the BioProject accession PRJNA803985

(https://www.ncbi.nlm.nih.gov/sra/PRJNA803985).

Bioinformatic Pipeline

Paired end sequences of the 16S rRNA gene amplicons were analyzed through the DADA2

package pipeline, version 1.20.0 (Callahan et al., 2016) in R 4.1.0 environment (R Core Team,

2021). Reads were filtered with the following parameters: truncLen = c(290, 225), truncQ = 2,

maxEE = c(2, 2). Merging of the forward and reverse reads was done with the mergePairs

function using the default parameters (minOverlap = 12, maxMismatch = 0). Chimeras were

removed using *removeBimeraDenovo* with default parameters. Taxonomic classification was

performed using the *assignTaxonomy* function using the parameters (tryRC = FALSE, minBoot

= 50) with the SILVA v138 database (Yilmaz et al., 2014) using the naive Bayesian classifier

method described in Wang et al. (2007). Mitochondria-derived sequences were removed from

the original dataset, obtaining 24575 ASVs. ASV abundance tables, taxonomy and

environmental data were imported into the phyloseq R package version 1.36.0 (McMurdie and

Holmes, 2013). To reduce the data complexity (Cao et al., 2021), bacterial ASVs were filtered

to keep taxa observed at least 2 times in at least 2 samples resulting in 3260 ASVs for further

analyses (2.7 % of total reads removed).

Functional Annotations

Functional composition were predicted using PICRUSt2 version 2.4.1 (Douglas et al., 2020).

First, the nearest sequenced taxon index (NSTI) values are obtained as a metric to determine

the degree to which ASVs in a sample are related to available reference genomes and

environmentally sampled organisms. Then, gene content inference per ASV was performed

using hidden-state prediction (HSP) approaches. Before prediction, data was normalized by

their 16S rRNA gene copy number abundance. Finally, metagenome inference was used to

generate MetaCyc pathway predictions (Caspi et al., 2018). MetaCyc pathways were linked to

their respective secondary superclass levels using the tools available at https://metacyc.org.

Data Analysis and Statistics

All data analysis and statistical tests were conducted in R 4.1.0 environment (R Core Team,

2021). Environmental variables were explored using the tidyverse package v.1.3.1 (Wickham

et al., 2019). Bacterial and functional communities were centered log-ratio (CLR) transformed

to address the negative correlation bias intrinsic to compositional data (Gloor et al., 2017) using

CoDaSeq v.0.99.6 (Gloor, 2016) and zCompositions v.1.3.4 (Palarea-Albaladejo and Martín-

Fernández, 2015) packages. To visualize differences among samples based on environmental

and community (ASVs composition and functional annotations) data, Principal Composition

Analysis (PCA) followed by K-means clustering and distance-based Redundancy Analysis (db-

RDA) on Aitchison distance (Aitchison, 1986) were performed, respectively. PCA were

implemented using tidymodels v.0.1.3 (Kuhn and Wickham, 2020), K-mean clustering and

validation methods using factoextra v.1.0.7 (Kassambara and Mundt, 2020), and db-RDA using

vegan v.2.5.7 (Oksanen et al., 2018) packages. Environmental variables which could explain

variation found on db-RDA ordinations were fitted using both the *envfit*, and *ordistep* functions,

the latter with forward and backward stepwise selections. Significant correlation between

environmental variables and ordination (p-value < 0.01 in both analyses) were plotted as

vectors (arrows) whose direction and length indicate the environmental variable gradient and

the correlation, respectively. Prior to both ordination methods, autocorrelated environmental

variables (i.e., r > 0.7, p-value ≤ 0.05) were combined into one for downstream analyses.

To verify the magnitude of variation along a gradient of environmental changes between

taxonomic and functional profiles, a correlation between bacterial community and functional

annotation distance matrices was calculated using Mantel tests (Spearman rank correlation and

9999 permutations). To evaluate the difference in community composition between leases and

control sites, ASVs and functional annotations were subjected to two widely used statistical

methods: ALDEx2 (Fernandes et al., 2013) and DESeq2 (Love et al., 2014) packages. The

combination of these two methods aims to overcome the drawbacks found in differential

abundance methods, such as low power and high False Discovery Rate (FDR) (Hawinkel et al.,

2019). For ALDEx2, clr-transformed abundance table was used as the input for the analyses,

whereas DESeq2 used variance-stabilized data to perform the evaluation. ASVs and pathways

were considered differentially abundant if they satisfied the following assumptions: an effect

size > |1.5| in ALDEx2 and an adjusted P value (FDR) < 0.05 and a log2foldchange > |2| in

DESeq2 analysis. The package ggtern v.3.3.5 (Hamilton and Ferry, 2018) was used for data

visualization. Then, the differentially abundant ASVs and pathways were used as model input

data (clr-transformed count table) to verify the relationship with the environmental variables

used in the previous ordination analyses (i.e., NH_4, NOx, turbidity, and oxygen) as described

below.

Random forest regression modelling was implemented through tidymodels v.0.1.3 (Kuhn and

Wickham, 2020) and randomForest v.4.6.14 (Liaw and Wiener, 2002) packages to test the

ability of the differentially abundant ASVs to predict estimates of the variability of the

aforementioned environmental variables. For this process normalized clr-transformed data was

split into training and testing sets in a proportion of ¾. To tune the hyperparameters (mtry and

min_n), a k-fold cross validation scheme (k = 10 with 3 repeats) with 501 trees of the training

set was used to estimate the best model based on root-mean-square error (RMSE) metric. The

model was finalized using the *finalize_workflow* and *select_best* functions. After that, the

model performance was tested in the training set and the metrics RMSE and R^2 were collected

using the *collect_metrics* function. Variable importance was measured by the residual sum of

squares. Model   predictions   and   variable   importance   were   extracted   with   the

collect_predictions function and vip v.0.3.2 package (Greenwell and Boehmke, 2020),

respectively and visualized using tidyverse package v.1.3.1 (Wickham et al., 2019). Code used

for data analyses in this study can be found in the associated GitHub repository

(https://github.com/ricrocha82/MH_2019_16s).

3. Results

Environmental Variables

The samples from leases and control sites showed a uniform trend in most environmental

variables when distances from cages were considered, however for NH_4, NO_2 and NOx,

concentrations were highly variable at both leases (Fig. 2; Tables S2, S3 and S4). Despite this

variation, NH_4 level was observed to decrease as distance from cage increased at the surface of

Lease 2 (2.84 ± 0.47 μM to 0.1 ± 0.03 μM) and NOx decreased in concentration with distance

in the middle waters of Lease 1 (6.38 ± 1.22 μM to 1.29 ± 0.46 μM). Also, at Lease 1, NO_2

showed an increase in concentration as distance increased after ~18 m (0.04 ± 0.05 μM to 0.28

± 0.04 μM).

Fig. 2 – Environmental variables along a distance gradient in each lease area. Each line in the graph displays environmental variable measurements in each depth. Red dashed line shows the farthest distance from the cage. (i.e., end of cage influenced zone). After this line cage site distances are displayed. Distance 0 m indicates sample from inside the cages (2 cages in Lease 1 and 3 cages in Lease 2). Dissolved nutrients (μmol l^{-1}), conductivity, fluorescence, oxygen (mg/l), temperature (°C), turbidity (NTU)

When depth is considered, stratification is clear for all environmental variables at all leases and

control sites (Fig. 3 – A; Tables S1-S4). Despite the variation observed in nutrient

concentrations, NH_4 (1.53 ± 0.85 µM to 0.3 ± 0.48 µM) and NO_2 (0.28 ± 0.08 µM to 0.14 ±

0.07 µM) decreased with depth whereas NOx (1.29 ± 0.51 µM to 7.57 ± 0.63 µM) and PO_4

(0.03 ± 0.02 µM to 0.3 ± 0.07 µM) displayed the opposite trend. Similarly to NOx and PO_4,

conductivity (6914.73 ± 2781.28 to 36573.31 ± 548.15), salinity (5 ± 2.26 to 29.7 ± 0.5) and

temperature (13.28 ± 0.41 °C to 14.45 ± 0.07 °C) were greater at depth, whereas fluorescence

(2.68 ± 0.2 to 1.26 ± 0.022 °C), oxygen (8.10 ± 1.14 mg/l to 0.86 ± 0.4 mg/l), PAR (0.62 ± 1

to 0.003 ± 0.001) and turbidity (2 ± 0.4 NTU to 0.72 ± 0.6 NTU) were all lower at depth as

seen for NH_4 and NO_2 concentrations. NOx was highly correlated to conductivity, salinity,

temperature and PO_4, whereas NH_4 was correlated to NO_2 and oxygen to PAR and fluorescence

(Fig.S2). Therefore, NOx, NH_4 and oxygen were used as proxies of their highly correlated

peers.

Fig. 3 – Ordination plots of the samples and vectors (arrows). (**A**) Principal Component Analysis (PCA) of
environmental data. Dashed lines highlight the results of K-mean clustering analyses. (**B-C**) Distance-based
Redundancy Analysis (db-RDA) of the composition differences of bacterial and functional communities. To avoid

multicollinearity, NOx included conductivity, salinity, temperature and PO_4, and NH_4 included NO_2. Vectors
indicate a significant correlation between environmental variables and ordination ($p < 0.01$) using *envfit* function
of vegan package. (**D**) Differences in ASV bacterial and functional compositions. Equal changes are shown by
the red line. Mantel test shows a significant correlation between the two distances matrices (Spearman rho =
0.6154).
285

Bacterial and Functional Communities Structure

At phylum and class levels, bacterial community at all sites was dominated by *Proteobacteria*

(Control: 58.4 %; Lease 1: 55.3 %; Lease 2: 57.9 %) and *Gammaproteobacteria* (Control:

30.4 %; Lease 1: 30.7 %; Lease 2: 31.7 %), respectively. *Proteobacteria* was followed by

Bacteroidota (Control: 15.6 %; Lease 1: 17 %; Lease 2: 18.1 %) at all sites, *Actinobacteriota*

(5.6 %) at Control sites and *Campylobacterota* (Lease 1: 7.63 %; Lease 2: 5.45 %) at lease

sites. *Gammaproteobacteria* was followed by *Alphaproteobacteria* (Control: 28.1 %; Lease 1:

24.6 %; Lease 2: 21.6 %) and *Bacteroidia* (Control: 15.2 %; Lease 1: 16.7 %; Lease 2: 17.9 %)

at all sites. ASVs encompassing the class *Actinobacteria* (Control: 4.3 %; Lease 1: 1.93 %;

Lease 2: 1.90 %) were more abundant at control sites whereas *Campylobacteria* ASVs were

more abundant at lease sites (Control: 3.3 %; Lease 1: 7.63 %; Lease 2: 5.45 %) (Fig. S3).

Functional annotations were uniformly distributed among leases and control sites. Amino-acid

biosynthesis was the most abundant pathway (Control: 17 %; Lease 1: 16.8 %; Lease 2: 16.9 %),

followed by nucleotide-biosynthesis (Control: 16.2 %; Lease 1: 16.3 %; Lease 2: 16.3 %) and

vitamin-biosynthesis (Control: 9.93 %; Lease 1: 10.2 %; Lease 2: 10.1 %) (Fig. S4). Despite

little variation in both communities among sites, depth stratification was evident when

ordination analyses were performed (Fig. 3 - B and C). db-RDA explained 49.47% and 31.18%

of the variation in bacterial and functional communities, respectively. The ordination of

bacterial community distribution was significantly correlated to NOx ($r^2 = 0.76$; $p < 0.01$), NH_4

($r^2 = 0.49$; $p < 0.01$) and turbidity ($r^2 = 0.36$; $p < 0.01$) whereas functional profiles were

significantly related to NOx ($r^2 = 0.74$; $p < 0.01$) and turbidity ($r^2 = 0.41$; $p < 0.01$) (Table 1).

The results indicate that environmental parameters and both bacterial and functional

communities are more affected by depth stratification than by aquaculture activity. Moreover,

the correlation between the distance matrices of bacterial ASVs and functional compositions

suggested a significant relationship (rho: 0.62; $p < 0.05$). In other words, the differences in the

taxonomic profile between samples co-vary with the differences in the functional profile

between samples. However, bacterial community showed a more prominent overall shift than

the associated functional data (Fig. 3 - D).

Table 1 - Result of redundancy analysis (*ordistep* and *envif* functions) and random forest (rf) of bacterial and
functional communities in the Macquarie Harbour. Regarding the db-RDA, we considered significant, if the
environmental variable had a p-value < 0.01 in both analyses. To avoid multicollinearity, NOx included
conductivity, salinity, temperature and PO_4. NH_4 included NO_2.

Community	env variable	p-value (ordistep)	p-value (envfit)	r2 (envfit)	RMSE (rf)	R2 (rf)
Bacteria	NOx*	0.005	0.001	0.76	1.14	0.87
	turbidity	0.005	0.001	0.36	0.66	0.44
	NH_4**	0.005	0.001	0.49	0.70	0.53
Functional annotation	NOx*	0.005	0.001	0.74	1.23	0.82
	turbidity	0.005	0.001	0.41	0.53	0.59

319 *NOx included conductivity, salinity, temperature and PO_4.
320 **NH_4 included NO_2.
321

Differential Abundance Among Sites by Depth

To detect key members in bacterial and functional communities at lease and control sites,

differential abundance analyses were performed considering the effect of depth stratification

observed in the community structure analyses. Therefore, the analyses were conducted by

executing pairwise comparisons between sites nested by depth (i.e., surface, middle and bottom

water layers). ALDEx2 and DESeq2 found a total of 90 and 167 bacterial ASVs and 82 and 61

functional annotations differentially abundant, respectively. At all depths, ASVs and functional

annotations selected by both methods were more abundant in both lease influential zones (i.e.,

relative abundance $\geq 60\%$) compared to control site. The difference in bacterial community

was pronounced at the bottom and surface layers (Fig. 4). Most of the enriched ASVs found in

Lease 1 area inhabited the bottom layers (Fig. S5 and Table S5). The taxon with the highest

number of differentially abundant ASVs were those belonging to the genus *Psychromonas*

(family *Psychromonadaceae*) followed by *Spirochaeta* 2 (family *Spirochaetaceae*), unknown

Arcobacteraceae (order *Campylobacterales*), unknown *Bacteroidales*, *Hypnocyclicus* and an

unknown genus (both assigned to the family *Leptotrichiaceae*), *Draconibacterium* and three

unknown genera related to the order *Bacteroidales*, *Sulfurimonas* (family *Sulfurimonadaceae*),

Desulfovibrio (family *Desulfovibrionaceae*), *Pseudomonas* (family *Pseudomonadaceae*), and

Woeseia (family *Steroidobacterales*). In contrast, at the surface of Lease 2, *Aliivibrio* (family:

Vibrionaceae) had the highest number of differentially abundant ASVs (Fig. S6 and Table S6).

In the control, taxa related to *Nitrospina* and *Gammaproteobacteria* were more abundant in the

bottom layers (Fig. S7 and Table S7).

Fig. 4 – Ternary plots of per-depth relative abundances of differentially abundant ASVs. Each point represents an
ASV, and its position indicates the proportion of its relative abundance at each site. Points closer to the ternary
plot corners indicate that a greater proportion of the total relative abundance of this ASV was found in this site.
Point colors indicate the Phylum and Class of the ASV. The size of each point indicates the mean relative
abundance of the ASV in all samples. The red lines inside the ternary plot indicate the 60% level of relative
abundance of each of the sites.

Regarding the functional community, the number of enriched functional annotation was the

highest at Lease 1, followed by Lease 2 and control sites (Fig. 5, S8 and S9; Tables S8 and S9).

In the bottom layer at Lease 1, the analysis detected pathways related with organic matter

degradation such as fermentation, carbohydrates, amino-acids, aromatic compounds, and

alcohol degradations, as well as those associated with trace metal and macro-nutrient

limitations (secondary metabolites, cofactor, vitamins, and amino acids biosynthesis),

methanogenesis and cell metabolism. Pathways associated with vitamin degradation and cell

metabolism (cell-structure/cell-wall biosynthesis) and those related to alcohol degradation and

vitamin biosynthesis were more abundant at the surface and middle layers, respectively. At the

control site, lipid biosynthesis and secondary metabolite biosynthesis were differentially

expressed at the middle and bottom layers, respectively. These results suggest that the

differences observed in bacterioplankton community (16S rRNA gene sequencing) and

functional profile (functional annotation: PICRUSt2) at each depth between the lease and

control stations are likely to reflect changes in the local environment due to aquaculture

activities.

Fig. 5 – Ternary plots of per-depth relative abundances of differentially abundant functions. Point colors indicate the pathway of the function according to https://metacyc.org. The size of each point indicates the mean relative abundance of the function in all samples. The red lines inside the ternary plot indicate the 60% level of relative abundance of each of the sites.

Predictive Performance of the Selected ASVs

To measure the importance of these differentially expressed features in predicting the environmental drivers found in the db-RDA analysis (Fig. 3 and Table 1), supervised machine learning was used to estimate the relationship between feature distribution and environmental variable measurements. Then, RF regression models were trained with 257 ASVs and 143 functional annotations. Only NOx (included conductivity, salinity, temperature and PO_4) concentration prediction models yielded desirable performance for both communities (Table 1). The most important ASV and functional annotation in reducing variance and uncertainty to NOx values were assigned to *Oligoflexaceae* (Fig. 6 and S10; Table S8) and to vitamin biosynthesis, respectively (Fig. 7 and S11; Table S9). Additionally, the most important features in both communities were uniformly distributed among sites, except for some ASVs and functional annotations. For instance, ASVs that could predict NOx concentration, OM27 clade

385 (*Bdellovibrionaceae*) and *Planctomycetes*, were found more abundant (> 60% relative

386 abundance) at the control and Lease 1 samples, respectively. And two functional annotations

387 assigned to methanogenesis and vitamin biosynthesis, both important in NOx prediction, were

388 more enriched in Lease 1 samples.

390
391 Fig. 6 – Ternary plot showing the distribution of the 10 most important ASVs based on the residuals sum of square
392 in regression-based prediction model of the environmental drivers of the bacterial community (NH4, NOx and
393 turbidity). The size of each point indicates the mean relative abundance of the ASV in all samples. The red lines
394 inside the ternary plot indicate the 60% level of relative abundance of each of the sites. To avoid multicollinearity,
395 NOx included conductivity, salinity, temperature and PO4; and NH4 included NO2.
396

Fig. 7 – Ternary plot showing the distribution of the 10 most important pathways based on the residuals sum of square in regression-based prediction model of the environmental drivers of the functional community (NOx and turbidity). The size of each point indicates the mean relative abundance of the function in all samples. To avoid multicollinearity, NOx included conductivity, salinity, temperature and PO_4.

4. Discussion

In this study, we used taxonomic and functional profiles to characterize the bacterioplankton community of a coastal embayment containing finfish aquaculture operations in Tasmania (Australia). Using unsupervised and supervised analysis techniques of 16S rRNA bacterial community composition, we observed that the effects of depth stratification on bacterioplankton dominates over aquaculture effects. Despite this natural influence, we also detected differentially abundant ASVs and functional annotations between both marine farm leases (< 85 m) and control sites (> 1500 m), that could be used as good bioindicators for aquaculture activity. According to the random forest regression model, these features could only predict NOx concentrations along with its highly correlated peers (i.e., conductivity, salinity, temperature and PO_4). Given the environmental heterogeneity and complexity of Macquarie Harbour, it is not possible to draw comprehensive conclusions regarding the impact of salmon aquaculture in the Harbour. Rather, our study provides initial insight into the

interaction between salmon aquaculture and the bacterioplankton community, as discussed

further below.

4.1. Community variation and depth stratification

The main driver of variation in the environmental variables, bacterial community and

functional annotations was water column stratification. Macquarie Harbour is considered a

highly stratified marine embayment with distinct communities inhabiting each layer in the

water column (Da Silva et al., 2021). The higher turbidity and NH_4 concentration observed at

the surface could be related to the tannin-rich waters from Gordon River affecting light

penetration which can favor nitrification and organic matter degradation within the water

column (Terry, 2001, Hartstein et al., 2019, Bartl et al., 2018, Hou et al., 2018). This could

explain the increasing NOx and decreasing DO and NH_4 concentrations with increasing depth.

Additionally, the stratified distribution of both communities (ASVs and functional annotations)

could be explained by the variation in the environmental variables, especially those which

increased with depth. Most of the differentially abundant ASVs and functional annotation used

in the regression analysis to predict NOx concentration were enriched at the bottom layers.

Thus, it is unsurprising that the vertical differences in both community distributions observed

in this study were strongly correlated to NOx concentration along with conductivity, salinity,

temperature and PO_4 that followed the same vertical profile.

4.2. Effects of aquaculture on bacterial community composition

In this study, differential abundance analysis revealed an effect on the taxonomic and

functional profiles at both the surface and the near-bottom water layers in lease areas. Most

previous aquaculture-environmental impact studies focus on sediments and associated effects

to the benthic microbial community (Borja et al., 2009, Fodelianakis et al., 2014, Dowle et al.,

2015, Aylagas et al., 2017, Stoeck et al., 2018, Cordier et al., 2018, Keeley et al., 2021). The

pelagic environment is less studied and as such the information regarding bacterioplankton

response to aquaculture operation is limited, despite their known sensitivity to environmental

changes (Wei et al., 2016, Lindh and Pinhassi, 2018, Kolda et al., 2020). Aquaculture activities

are known to increase nutrient loading and change trophic status near the fish cages from waste,

by-products, fish feces or feed products inputs (Belias et al., 2003, Buschmann et al., 2006,

Amirkolaie, 2011). Various microbial groups have been associated with high level of nutrient

inputs in the water column (Haro-Moreno et al., 2020, Yang et al., 2019), sediments

(Verhoeven et al., 2018, Fruhe et al., 2021, Zhang et al., 2020, Asami et al., 2005, Dully et al.,

2021) and land-based aquaculture operations (Mahmoud and Magdy, 2021, Kandel et al., 2014,

Aalto et al., 2021, Blancheton et al., 2013) and some have been reported as potential

aquaculture impact biomarkers. (Haro-Moreno et al., 2020, Mahmoud and Magdy, 2021,

Kandel et al., 2014, Verhoeven et al., 2018, Frühe et al., 2021, Zhang et al., 2020, Aalto et al.,

2021, Blancheton et al., 2013, Yang et al., 2019, Rasigraf et al., 2017, Dully et al., 2021). In

accordance with these studies, we found differentially abundant ASVs in lease areas that could

be related to higher levels of organic matter and the presence of farmed fish. These include

members of the taxa *Psychromonadaceae* (*Psychromonas), Fusobacteriales*

(Fusobacteriaceae), Vibrionaceae, Desulfovibrionaceae (Desulfovibrio), Sulfurimonadaceae

(Sulfurimonas), Sulfurovaceae (Sulfurovum), Fusibacteraceae (Fusibacter), Rhizobiales,

Sphingomonadaceae, Methylotenera, Hyphomonadaceae, *Spirosomaceae, Flavobacteriaceae*

(NS3a marine group) and *Bacteriovoracaceae*. The abundant ASVs detected around leases in

this study correlate well with previous studies conducted in different environments. It suggests

that bacterioplankton community from the Macquarie Harbour had similar response to

aquaculture activities. Therefore, the techniques applied here could be potential candidates for

biomonitoring in coastal salmon aquaculture.

The abundance of members linked to sulphur cycling (i.e., *Desulfovibrionaceae* and *Fusibacter*)

have been found in a range of impacted aquatic systems (Blancheton et al., 2013, Aranda et al.,

2015, Frühe et al., 2021, Verhoeven et al., 2018, Won et al., 2017). They contribute to organic

matter and sulphur-cycling in environments with high levels of organic matter. These findings

support previous studies linking nutrient enrichment and sulphur cycle leading to oxygen

depletion and the increased production of sulphide (Barton and Hamilton, 2007, Kwon et al.,

2016). Moreover, other ASVs detected in higher relative abundance in proximity to aquaculture

were those often related to nitrogen and methane cycling. The presence of *Thioglobaceae*

(SUP05 cluster), *Beijerinckiaceae* (*Methylocystis*) and *Methylophilaceae* (*Methylotenera*) have

been previously reported in both pelagic (Haro-Moreno et al., 2020) and benthic (Fan et al.,

2019) aquaculture sites as well as in Atlantic salmon hatcheries (Lokesh et al., 2019, Minich et

al., 2020) and could be linked to oxygen concentration and feed inputs (Fan et al., 2019,

Hernandez et al., 2015). Interestingly, the predatory *Bacteriovoracaceae* were also found

differentially abundant in around leases and could be related to the increase in the number of

prey items such as *Vibrio* (Chen et al., 2011, Pineiro et al., 2007). This supports a previous

study that detected *Bacteriovoracaceae* as a potential indicator for aquaculture activity (Frühe

et al., 2021). It is plausible that these enriched bacterial taxa are most likely related to organic

enrichment caused by fish faeces or feed inputs. These distinctive groups of microorganisms

represent potential biomarkers and could be tested in further studies to trace and monitor

changes caused by aquaculture.

Psychromonadaceae (*Psychromonas*), *Fusobacteriales (Fusobacteriaceae), Spirochaetaceae,*

and *Leptotrichiaceae*, *Enterobacterales*, *Pseudomonadales* (*Pseudomonas*) and *Vibrionaceae*

(*Vibrio* and *Aliivibrio*) were also found in higher differential abundance in lease influenced

zones compared with control sites. Those ASVs have been reported as members of the marine

and freshwater fish gut microbiome (Algammal et al., 2020, Fogarty et al., 2019) often found

in bacterial profiling studies using molecular techniques in sediments (Verhoeven et al., 2016,

Fruhe et al., 2021, Verhoeven et al., 2018, Kolda et al., 2020), water column (Kopprio et al.,

2021, Haro-Moreno et al., 2020), and other aquaculture operation (Martins et al., 2013). Their

presence in leases and their absence or low abundance in the control sites suggests an influence

of fish faeces on the bacterioplankton community at the aquaculture sites. Additionally, some

of these ASVs have been reported as potential pathogens such as *Aliivibrio, Pseudomonas,* and

Arcobacteraceae (Zarkasi et al., 2016, Bozzi et al., 2021, Wakabayashi, 1980, Loch et al., 2012,

Ho et al., 2008) and should be considered in further studies to analyse the role of fish farms as

potential reservoirs for pathogens.

4.3. The influence of aquaculture activity on functional structure of bacterioplankton

Taxonomic composition of bacterial communities through the use of 16S rRNA gene profiling

is useful in showing differences in community composition between sites, as well as providing

understanding regarding taxonomic response to the presence of fish cages in the Harbour.

However, evidence on potential functional composition can provide important insights into the

interaction between metabolic pathways and environmental stress (Mukherjee et al., 2017, Wu

et al., 2019, Laroche et al., 2021, Raes et al., 2021). PICRUSt2 is a computational approach to

predict the functional composition of a metagenome from the 16S rRNA sequence (Douglas et

al., 2020), and has been used widely in previous studies on aquaculture environmental impact

(Qu et al., 2021, Vargas-Albores et al., 2019, Ortiz-Estrada et al., 2019, Laroche et al., 2021,

Hornick and Buschmann, 2017). By applying PICRUSt2, we are able to explore how the

functional community changes with the presence of aquaculture activities.

In this study, PICRUSt2 inferred a methanogenesis pathway from bacterial 16S rRNA

amplicon sequencing data. It is well known that methanogenesis is only seen in the archaeal

kingdom. Although only bacterial sequences were used as input, this inference could be related

to an incorrect clustering with reference archaeal sequences. For these reasons, results from

functional prediction tools are useful and insightful but should be interpreted with caution

(Douglas et al., 2020). The differences were more pronounced in the near-bottom water layers

of leases areas where enriched pathways were associated with organic compounds and vitamin

degradation, fermentation, methanogenesis. These pathways are likely related to environment

change caused by the presence of fish, including organic enrichment and oxygen depletion, and

is consistent with studies on functional alteration along pollution gradients (Galand et al., 2016,

Cordier et al., 2019). Interestingly, antibiotic resistance pathway was also found enriched

closed to the finfish farms supporting previous studies on aquaculture impact (Tamminen et al.,

2011, Hornick and Buschmann, 2017, Chen et al., 2017). The application of antimicrobials can

lead to selective pressures in which resistant taxa thrive to the detriment of others (Miranda

and Zemelman, 2002). While outputs from functional prediction tools need to be interpreted

within their limitations, results from this study suggest that an accumulative effect of organic

enrichment, oxygen depletion and antibiotic usage is likely to affect the functional profile of

the bacterial community in Macquarie Harbour. Further studies are necessary to test this

hypothesis.

Overall, we found that the relative stability of the functional community was greater than the

taxonomic community between lease and control sites. The stability detected in this study may

be related to the concept of functional redundancy among prokaryotes (Hornick and

Buschmann, 2017, Cordier, 2020, Louca et al., 2016, Louca et al., 2018, Bissett et al., 2006,

Bissett et al., 2007). This same relationship was observed in previous studies that assessed

bacterial community and functional prediction as indicators of anthropogenic pressures such

as aquaculture and oil and gas activities (Cordier, 2020, Laroche et al., 2018). Both authors

suggested that functional redundancy along with 16S rRNA sequencing could be a useful proxy

to assess ecosystem resilience, since deep metagenomic sequencing across many samples is

economically prohibitive for many studies. (Wemheuer et al., 2020, Sun et al., 2021, Teichert

et al., 2018). Therefore, this functional trait-based approach should be considered as a potential

tool for assessing the impacts of aquaculture. That said, metagenome prediction tools can lead

to incorrect predictions as a result of gene transfer rate or phylogenetic variability. Furthermore,

the performance of these tools for inference on genes associated with "housekeeping" drops

significantly in environmental samples (Sun et al., 2020, Langille et al., 2013). Therefore, it is

important to highlight that the improvement of currently available reference databases used by

these inference tools is crucial for environmental impact assessment, since most of the genomes

available are associated to human health and biotechnology use (Sun et al., 2020).

4.4. Implications for environmental variability

It is important to note a greater number of differentially abundant ASVs associated with the

Lease 1 local area, particularly in the near-bottom waters, compared to Lease 2 and control

locations. The difference could be related to the production (e.g., stock density, faeces),

management practices (e.g., feed input rate), hydrodynamics and location (e.g., shallow or

deeper waters) of that specific lease (Belias et al., 2003, Sarà et al., 2007, Tsagaraki et al., 2011,

Jansen et al., 2016, Sanderson et al., 2008). Hydrodynamics and location of both leases could

be important factors explaining this difference. While Lease 1 is located near the southern end

of Macquarie Harbour, Lease 2 is further north, near the deepest basin with a depth of

approximately 52 m. Macquarie Harbour is a fjord-like system with periods of deep-water

renewal entering by a narrow and shallow entrance located at the Harbour's northern end. Due

to its geological formation, and freshwater, seasonal or weather-related influences, the

distribution of the marine water intrusions are heterogeneously distributed leading to a less

frequent and less pronounced deep water renewal events in the southern Harbour (Hartstein et

al., 2019, Ross et al., 2021). Also, Teasdale et al. (2003) revealed increasing organic carbon

and sulphide concentrations from north to south which was influenced by riverine inputs. This

north-south environmental gradient may affect the oxygen and organic matter concentration in

the water column in the Lease 1 area, causing changes in the environmental conditions as we

observed in our study. Although our findings suggest bacterioplankton as a potential indicator

for aquaculture activities, more work is needed to confirm its reliability over time in the context

of natural variation. Understanding seasonal change in the bacterial community across the

Harbour will be particularly important to improve our capacity to detect disturbances caused

by both natural and anthropogenic environmental changes (Allison and Martiny, 2008,

Magurran et al., 2010, Lindh et al., 2015).

5. Conclusions

In summary, investigation of bacterioplankton communities and predicted functions in water

column revealed localised changes in the presence of salmon cages. Local enriched taxa were

related to biochemical processes (i.e., sulphur, nitrogen, methane, and carbon cycling),

potential pathogens or members of the fish gut microbiome, while the relative abundance of

function annotations was associated with organic compounds and vitamin degradation,

fermentation, methanogenesis and antibiotic resistance. Our study demonstrates that

molecular-based methods could be a valuable tool in monitoring aquaculture effects. Trends in

nutrient concentrations associated with aquaculture were likely be masked by the natural

environmental variation in the water column, strongly influencing the distribution of the

bacterial taxonomic and functional communities. The robustness of the aquaculture-associated

indicators suggested by this study could be further validated through sampling across both

season and farm production cycle to examine temporal variation in taxonomic and functional

bacterioplankton communities. Overall, the study showed that the 16S rRNA sequencing and

functional annotation of the bacterioplankton communities have the potential to become

effective tools in understanding the interaction between finfish aquaculture and the pelagic

environment.

6. Author contributions

RRPS, CW and JR: Conceptualization. **RRPS**: DNA extraction, Formal analysis, Writing-

Original draft preparation. **CW, JR and JB**: Supervision, Writing- Reviewing and Editing**.**

7. Declaration of competing interest

We declare that the submitted work was conducted in the absence of any commercial or

financial relationships that could potentially be construed as a conflict of interest.

8. Acknowledgments

This study was supported by the Fisheries Research & Development Council grant 2016-067

and the Sustainable Marine Research Collaborative Agreement between the Institute for

Marine and Antarctic Studies and the State Government of Tasmania. The authors would like

to thank Samuel Kruimink, Patricia Peinado Fuentes, Ruth Eriksen, Andy Revill, Kirsten Karsh,

Lev Bodrossy and Eric Raes for their assistance in obtaining field samples and laboratory data.

We appreciate the help and comments of the anonymous reviewers,

that improved our manuscript notably.

9. References

á Norði, G., Glud, R. N., Gaard, E. and Simonsen, K. (2011) 'Environmental impacts of coastal fish farming: carbon and nitrogen budgets for trout farming in Kaldbaksfjør7%%FONT_ERR%%ur (Faroe Islands)', *Marine Ecology Progress Series,* 431, pp. 223-241. doi: 10.3354/meps09113

Aalto, S. L., Suurnakki, S., von Ahnen, M., Tiirola, M. and Pedersen, P. B. (2021) 'Microbial communities in full-scale woodchip bioreactors treating aquaculture effluents', *J Environ Manage,* 301, pp. 113852. doi: 10.1016/j.jenvman.2021.113852

Aitchison, J. (1986) *The Statistical Analysis of Compositional Data.* London; New York: Chapman and Hall.

Algammal, A. M., Mabrok, M., Sivaramasamy, E., Youssef, F. M., Atwa, M. H., El-Kholy, A. W., Hetta, H. F. and Hozzein, W. N. (2020) 'Emerging MDR-Pseudomonas aeruginosa in fish commonly harbor oprL and toxA virulence genes and blaTEM, blaCTX-M, and tetA antibiotic-resistance genes', *Sci Rep,* 10(1), pp. 15961. doi: 10.1038/s41598-020-72264-4

Allison, S. D. and Martiny, J. B. (2008) 'Resistance, resilience, and redundancy in microbial communities', *Proceedings of the National Academy of Sciences,* 105(Supplement 1), pp. 11512-11519. doi:

Amirkolaie, A. K. (2011) 'Reduction in the environmental impact of waste discharged by fish farms through feed and feeding', *Reviews in Aquaculture,* 3(1), pp. 19-26. doi: 10.1111/j.1753-5131.2010.01040.x

Appleyard, S., Abell, G. and Watson, R. (2013) *Tackling microbial related issues in cultured shellfish via integrated molecular and water chemistry approaches. Seafood CRC Final Report (2011/729) (pp. 89).* Available at: http://frdc.com.au/Archived-Reports/FRDC%20Projects/2011-729-DLD.pdf (Accessed: 20 August 2019).

Aranda, C. P., Valenzuela, C., Matamala, Y., Godoy, F. A. and Aranda, N. (2015) 'Sulphur-cycling bacteria and ciliated protozoans in a Beggiatoaceae mat covering organically enriched sediments beneath a salmon farm in a southern Chilean fjord', *Mar Pollut Bull,* 100(1), pp. 270-278. doi: 10.1016/j.marpolbul.2015.08.040

Asami, H., Aida, M. and Watanabe, K. (2005) 'Accelerated sulfur cycle in coastal marine sediment beneath areas of intensive shellfish aquaculture', *Appl Environ Microbiol,* 71(6), pp. 2925-33. doi: 10.1128/AEM.71.6.2925-2933.2005

Asshauer, K. P., Wemheuer, B., Daniel, R. and Meinicke, P. (2015) 'Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data', *Bioinformatics,* 31(17), pp. 2882-4. doi: 10.1093/bioinformatics/btv287

Aylagas, E., Borja, A., Tangherlini, M., Dell'Anno, A., Corinaldesi, C., Michell, C. T., Irigoien, X., Danovaro, R. and Rodriguez-Ezpeleta, N. (2017) 'A bacterial community-based index to assess the ecological status of estuarine and coastal environments', *Mar Pollut Bull,* 114(2), pp. 679-688. doi: 10.1016/j.marpolbul.2016.10.050

Azam, F., Fenchel, T., Field, J., Grey, J., Meyer-Reil, L. and Thingstad, F. (1983) 'The ecological role of water-column microbes', *Mar. ecol. Prog. ser,* 10, pp. 257-263. doi:

Bartl, I., Liskow, I., Schulz, K., Umlauf, L. and Voss, M. (2018) 'River plume and bottom boundary layer–Hotspots for nitrification in a coastal bay?', *Estuarine, Coastal and Shelf Science,* 208, pp. 70-82. doi:

Barton, L. L. and Hamilton, W. A. (2007) *Sulphate-reducing bacteria: environmental and engineered systems.* Cambridge University Press.

Belias, C. V., Bikas, V. G., Dassenakis, M. J. and Scoullos, M. J. (2003) 'Environmental impacts of coastal aquaculture in eastern mediterranean bays the case of astakos gulf, Greece', *Environmental Science and Pollution Research,* 10(5), pp. 287. doi: 10.1065/espr2003.06.159

Bissett, A., Bowman, J. and Burke, C. (2006) 'Bacterial diversity in organically-enriched fish farm sediments', *FEMS Microbiol Ecol,* 55(1), pp. 48-56. doi: 10.1111/j.1574-6941.2005.00012.x

Bissett, A., Burke, C., Cook, P. L. and Bowman, J. P. (2007) 'Bacterial community shifts in organically
perturbed sediments', *Environ Microbiol,* 9(1), pp. 46-60. doi: 10.1111/j.1462-
2920.2006.01110.x

Blancheton, J. P., Attramadal, K. J. K., Michaud, L., d'Orbcastel, E. R. and Vadstein, O. (2013) 'Insight
into bacterial population in aquaculture systems and its implication', *Aquacultural Engineering,*
53, pp. 30-39. doi: 10.1016/j.aquaeng.2012.11.009

Borja, Á., Rodríguez, J. G., Black, K., Bodoy, A., Emblow, C., Fernandes, T. F., Forte, J., Karakassis, I.,
Muxika, I., Nickell, T. D., Papageorgiou, N., Pranovi, F., Sevastou, K., Tomassetti, P. and Angel,
D. (2009) 'Assessing the suitability of a range of benthic indices in the evaluation of
environmental impact of fin and shellfish aquaculture located in sites across Europe',
Aquaculture, 293(3-4), pp. 231-240. doi: 10.1016/j.aquaculture.2009.04.037

Bouwman, L., Beusen, A., Glibert, P. M., Overbeek, C., Pawlowski, M., Herrera, J., Mulsow, S., Yu, R.
and Zhou, M. (2013) 'Mariculture: significant and expanding cause of coastal nutrient
enrichment', *Environmental Research Letters,* 8(4). doi: 10.1088/1748-9326/8/4/044026

Bowman, J. S. and Ducklow, H. W. (2015) 'Microbial Communities Can Be Described by Metabolic
Structure: A General Framework and Application to a Seasonally Variable, Depth-Stratified
Microbial Community from the Coastal West Antarctic Peninsula', *PLoS One,* 10(8), pp.
e0135868. doi: 10.1371/journal.pone.0135868

Bozzi, D., Rasmussen, J. A., Carøe, C., Sveier, H., Nordøy, K., Gilbert, M. T. P. and Limborg, M. T. (2021)
'Salmon gut microbiota correlates with disease infection status: potential for monitoring
health in farmed animals', *Animal microbiome,* 3(1), pp. 1-17. doi:

Burridge, L., Weis, J. S., Cabello, F., Pizarro, J. and Bostick, K. (2010) 'Chemical use in salmon
aquaculture: A review of current practices and possible environmental effects', *Aquaculture,*
306(1), pp. 7-23. doi: https://doi.org/10.1016/j.aquaculture.2010.05.020

Buschmann, A., Riquelme, V., Hernandezgonzalez, M., Varela, D., Jimenez, J., Henriquez, L., Vergara,
P., Guinez, R. and Filun, L. (2006) 'A review of the impacts of salmonid farming on marine
coastal ecosystems in the southeast Pacific', *ICES Journal of Marine Science,* 63(7), pp. 1338-
1345. doi: 10.1016/j.icesjms.2006.04.021

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. and Holmes, S. P. (2016) 'DADA2:
High-resolution sample inference from Illumina amplicon data', *Nat Methods,* 13(7), pp. 581-
3. doi: 10.1038/nmeth.3869

Cao, Q., Sun, X., Rajesh, K., Chalasani, N., Gelow, K., Katz, B., Shah, V. H., Sanyal, A. J. and Smirnova, E.
(2021) 'Effects of Rare Microbiome Taxa Filtering on Statistical Analysis', *Frontiers in
Microbiology,* 11. doi: 10.3389/fmicb.2020.607325

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., Fierer,
N. and Knight, R. (2011) 'Global patterns of 16S rRNA diversity at a depth of millions of
sequences per sample', *Proc Natl Acad Sci U S A,* 108 Suppl 1, pp. 4516-22. doi:
10.1073/pnas.1000080107

Carpenter, P. D., Butler, E. C. V., Higgins, H. W., Mackey, D. J. and Nichols, P. D. (1991) 'Chemistry of
trace elements, humic substances and sedimentary organic matter in Macquarie Harbour,
Tasmania', *Marine and Freshwater Research,* 42(6). doi: 10.1071/mf9910625

Caspi, R., Billington, R., Fulcher, C. A., Keseler, I. M., Kothari, A., Krummenacker, M., Latendresse, M.,
Midford, P. E., Ong, Q. and Ong, W. K. (2018) 'The MetaCyc database of metabolic pathways
and enzymes', *Nucleic acids research,* 46(D1), pp. D633-D639. doi:

Chen, C., Zheng, L., Zhou, J. and Zhao, H. (2017) 'Persistence and risk of antibiotic residues and
antibiotic resistance genes in major mariculture sites in Southeast China', *Science of the Total
Environment,* 580, pp. 1175-1184. doi:

Chen, H., Athar, R., Zheng, G. and Williams, H. N. (2011) 'Prey bacteria shape the community structure
of their predators', *ISME J,* 5(8), pp. 1314-22. doi: 10.1038/ismej.2011.4

Cordier, T. (2020) 'Bacterial communities' taxonomic and functional turnovers both accurately predict marine benthic ecological quality status', *Environmental DNA,* 2(2), pp. 175-183. doi: 10.1002/edn3.55

Cordier, T., Forster, D., Dufresne, Y., Martins, C. I. M., Stoeck, T. and Pawlowski, J. (2018) 'Supervised machine learning outperforms taxonomy-based environmental DNA metabarcoding applied to biomonitoring', *Molecular Ecology Resources,* 18(6), pp. 1381-1391. doi: 10.1111/1755-0998.12926

Cordier, T., Frontalini, F., Cermakova, K., Apotheloz-Perret-Gentil, L., Treglia, M., Scantamburlo, E., Bonamin, V. and Pawlowski, J. (2019) 'Multi-marker eDNA metabarcoding survey to assess the environmental impact of three offshore gas platforms in the North Adriatic Sea (Italy)', *Mar Environ Res*. doi: 10.1016/j.marenvres.2018.12.009

Crawford, C. M., Macleod, C. K. A. and Mitchell, I. M. (2003) 'Effects of shellfish farming on the benthic environment', *Aquaculture,* 224(1-4), pp. 117-140. doi: 10.1016/s0044-8486(03)00210-2

Cresswell, G., Edwards, R. and Barker, B. 'Macquarie Harbour, Tasmania-seasonal oceanographic surveys in 1985'. *Papers and Proceedings of the Royal Society of Tasmania*, 63-66.

Da Silva, R. R. P., White, C. A., Bowman, J. P., Raes, E., Bisset, A., Chapman, C., Bodrossy, L. and Ross, D. J. (2021) 'Environmental influences shaping microbial communities in a low oxygen, highly stratified marine embayment', *Aquatic Microbial Ecology,* 87, pp. 185-203. doi: 10.3354/ame01978

DeLong, E. F., Franks, D. G. and Alldredge, A. L. (1993) 'Phylogenetic diversity of aggregate‐attached vs. free‐living marine bacterial assemblages', *Limnology and oceanography,* 38(5), pp. 924-934. doi:

Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M., Huttenhower, C. and Langille, M. G. (2020) 'PICRUSt2 for prediction of metagenome functions', *Nature biotechnology,* 38(6), pp. 685-688. doi:

Dowle, E., Pochon, X., Keeley, N. and Wood, S. A. (2015) 'Assessing the effects of salmon farming seabed enrichment using bacterial community diversity and high-throughput sequencing', *FEMS microbiology ecology,* 91(8), pp. fiv089. doi. Available at: https://watermark.silverchair.com/fiv089.pdf?token=AQECAHi208BE49Ooan9kkhW_Ercy7D m3ZL_9Cf3qfKAc485ysgAAAkYwggJCBgkqhkiG9w0BBwagggIzMIICLwIBADCCAigGCSqGSIb3D QEHATAeBglghkgBZQMEAS4wEQQMQWvb8ISYQTRAZ8wwAgEQgIIB-fmA6b-VKmxRIKm_00Vh6ojIDonSb5oQGpfAQhoAK7igi1-yzZ-2H5wf9AwmnCPN9mmtNwSHu1U_sKBJ9DUlyT_9xnAYifzcC4yMxRzWZgbKMhZ8s4RrZFNl496 VoXcuokjiJPTNitieX4ec1Ek9-BwSn-pHuX2QmqAW1LHrtagPHJeWv55y3148eIz_fZTo9J2wpimBscNkCnSi0r1irXrpljF2GlBgJmPFpSh Ls3b3kg6y3nXICUHXucM-1TxqhYkasn0KE4-3M8q0tewG8pM823-9bjLCkff61kwjFjR0GHs15jfmg6cGq85Y-3suRr9Q_ec0GGZtVzTyOP-yx9w_49Yj-N147JH4X72e0EilBqgzws1h6KdSTv1T8BGWECVCZSKGy7dZK5RD_v1kDwLJZprNL-hlNQbmPUDomKoSxRXaIZV_Rb5qP6scjFz_K1y2fjove3wKvuO_sAJ5NTLu8Yvx3PtvY5fsaYVPgr anqdas5IacMVyOPydOSb48ZMbNXzXjkDvDboUi2SfJ5NljVB1IoAO64B9LRlKuJ-tlecpjpcd74CpzZAsUKZyCi3SlpZC2MqKa7tqkFSG1jK9K_B_hbmVrhOFrKWZKY06E_eEtfjlgiHAw egrbj9Z4y4bIWGdoaXm0BK9kpJHeJLOHwrHN3gvu_3Y

Dully, V., Balliet, H., Frühe, L., Däumer, M., Thielen, A., Gallie, S., Berrill, I. and Stoeck, T. (2021) 'Robustness, sensitivity and reproducibility of eDNA metabarcoding as an environmental biomonitoring tool in coastal salmon aquaculture – An inter-laboratory study', *Ecological Indicators,* 121. doi: 10.1016/j.ecolind.2020.107049

Elizondo-Patrone, C., Hernández, K., Yannicelli, B., Olsen, L. M. and Molina, V. (2015) 'The response of nitrifying microbial assemblages to ammonium (NH4+) enrichment from salmon farm activities in a northern Chilean Fjord', *Estuarine, Coastal and Shelf Science,* 166, pp. 131-142. doi: 10.1016/j.ecss.2015.03.021

Fan, L., Qiu, L., Song, C., Meng, S., Zheng, Y., Li, D., Hu, G. and Chen, J. (2019) 'Effects of feed input and
planting of submerged aquatic vegetation on methanotrophic communities in the surface
sediments of aquaculture ponds', *Applied soil ecology,* 143, pp. 10-16. doi:
Fernandes, A. D., Macklaim, J. M., Linn, T. G., Reid, G. and Gloor, G. B. (2013) 'ANOVA-like differential
expression (ALDEx) analysis for mixed population RNA-Seq', *PLoS One,* 8(7), pp. e67019. doi:
10.1371/journal.pone.0067019
Fodelianakis, S., Papageorgiou, N., Karakassis, I. and Ladoukakis, E. D. (2014) 'Community structure
changes in sediment bacterial communities along an organic enrichment gradient associated
with fish farming', *Annals of Microbiology,* 65(1), pp. 331-338. doi: 10.1007/s13213-014-0865-
4
Fogarty, C., Burgess, C. M., Cotter, P. D., Cabrera-Rubio, R., Whyte, P., Smyth, C. and Bolton, D. J. (2019)
'Diversity and composition of the gut microbiota of Atlantic salmon (Salmo salar) farmed in
Irish waters', *J Appl Microbiol,* 127(3), pp. 648-657. doi: 10.1111/jam.14291
Frühe, L., Dully, V., Forster, D., Keeley, N. B., Laroche, O., Pochon, X., Robinson, S., Wilding, T. and
Stoeck, T. (2021) 'Global trends of benthic bacterial diversity and community composition
along organic enrichment gradients of salmon farms', *Frontiers in Microbiology,* 12, pp. 853.
doi:
Fruhe, L., Dully, V., Forster, D., Keeley, N. B., Laroche, O., Pochon, X., Robinson, S., Wilding, T. A. and
Stoeck, T. (2021) 'Global Trends of Benthic Bacterial Diversity and Community Composition
Along Organic Enrichment Gradients of Salmon Farms', *Front Microbiol,* 12, pp. 637811. doi:
10.3389/fmicb.2021.637811
Fuhrman, J. A. and Steele, J. A. (2008) 'Community structure of marine bacterioplankton: patterns,
networks, and relationships to function', *Aquatic Microbial Ecology,* 53, pp. 69-81. doi:
10.3354/ame01222
Galand, P. E., Lucas, S., Fagervold, S. K., Peru, E., Pruski, A. M., Vetion, G., Dupuy, C. and Guizien, K.
(2016) 'Disturbance Increases Microbial Community Diversity and Production in Marine
Sediments', *Front Microbiol,* 7, pp. 1950. doi: 10.3389/fmicb.2016.01950
Gilbert, J. A., Field, D., Swift, P., Thomas, S., Cummings, D., Temperton, B., Weynberg, K., Huse, S.,
Hughes, M. and Joint, I. (2010) 'The taxonomic and functional diversity of microbes at a
temperate coastal site: a 'multi-omic'study of seasonal and diel temporal variation', *PLoS one,*
5(11),   pp.   e15545.   doi.   Available   at:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2993967/pdf/pone.0015545.pdf
Girvan, M., Campbell, C., Killham, K., Prosser, J. I. and Glover, L. A. (2005) 'Bacterial diversity promotes
community stability and functional resilience after perturbation', *Environmental microbiology,*
7(3), pp. 301-313. doi:
Gloor, G. (2016) 'CoDaSeq: Analyzing HTS using compositional data analysis', *F1000Research,* 5. doi:
Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. and Egozcue, J. J. (2017) 'Microbiome Datasets Are
Compositional: And This Is Not Optional', *Front Microbiol,* 8, pp. 2224. doi:
10.3389/fmicb.2017.02224
Grasshoff, K., Kremling, K. and Ehrhardt, M. (2009) *Methods of seawater analysis.* John Wiley & Sons.
Greenwell, B. M. and Boehmke, B. C. (2020) ' Variable Importance Plots—An Introduction to the vip
Package. ', *The R Journal,* 12(1), pp. 343--366. doi: https://doi.org/10.32614/RJ-2020-013.
Hamilton, N. E. and Ferry, M. (2018) 'ggtern: Ternary diagrams using ggplot2', *Journal of Statistical
Software,* 87(1), pp. 1-17. doi:
Han, P., Klumper, U., Wong, A., Li, M., Lin, J. G., Quan, Z., Denecke, M. and Gu, J. D. (2017) 'Assessment
of molecular detection of anaerobic ammonium-oxidizing (anammox) bacteria in different
environmental samples using PCR primers based on 16S rRNA and functional genes', *Appl
Microbiol Biotechnol,* 101(20), pp. 7689-7702. doi: 10.1007/s00253-017-8502-3
Haro-Moreno, J. M., Coutinho, F. H., Zaragoza-Solas, A., Picazo, A., Almagro-Moreno, S. and Lopez-
Perez, M. (2020) 'Dysbiosis in marine aquaculture revealed through microbiome analysis:

reverse ecology for environmental sustainability', *FEMS Microbiol Ecol,* 96(12). doi:
10.1093/femsec/fiaa218

Hartstein, N. D., Maxey, J. D., Loo, J. C. H. and Then, A. Y.-H. (2019) 'Drivers of deep water renewal in
Macquarie Harbour, Tasmania', *Journal of Marine Systems,* 199(103226). doi:
10.1016/j.jmarsys.2019.103226

Hawinkel, S., Mattiello, F., Bijnens, L. and Thas, O. (2019) 'A broken promise: microbiome differential
abundance methods do not control the false discovery rate', *Brief Bioinform,* 20(1), pp. 210-
221. doi: 10.1093/bib/bbx104

Hernandez, M. E., Beck, D. A., Lidstrom, M. E. and Chistoserdova, L. (2015) 'Oxygen availability is a
major factor in determining the composition of microbial communities involved in methane
oxidation', *PeerJ,* 3, pp. e801. doi:

Ho, H., Lipman, L. and Gaastra, W. (2008) 'Is Arcobacter a Food Related Pathogen Causing an Emerging
Disease?', *International Journal of Infectious Diseases,* 12, pp. e325-e326. doi:

Hornick, K. M. and Buschmann, A. H. (2017) 'Insights into the diversity and metabolic function of
bacterial communities in sediments from Chilean salmon aquaculture sites', *Annals of
Microbiology,* 68(2), pp. 63-77. doi: 10.1007/s13213-017-1317-8

Hou, L., Xie, X., Wan, X., Kao, S.-J., Jiao, N. and Zhang, Y. (2018) 'Niche differentiation of ammonia and
nitrite oxidizers along a salinity gradient from the Pearl River estuary to the South China Sea',
Biogeosciences, 15(16), pp. 5169-5187. doi: 10.5194/bg-15-5169-2018

Jansen, H. M., Reid, G. K., Bannister, R. J., Husa, V., Robinson, S. M. C., Cooper, J. A., Quinton, C. and
Strand, Ø. (2016) 'Discrete water quality sampling at open-water aquaculture sites: limitations
and strategies', *Aquaculture Environment Interactions,* 8, pp. 463-480. doi: 10.3354/aei00192

Kandel, P. P., Pasternak, Z., van Rijn, J., Nahum, O. and Jurkevitch, E. (2014) 'Abundance, diversity and
seasonal dynamics of predatory bacteria in aquaculture zero discharge systems', *FEMS
Microbiol Ecol,* 89(1), pp. 149-61. doi: 10.1111/1574-6941.12342

Author (2020) *factoextra: Extract and Visualize the Results of Multivariate Data
Analyses. R package version 1.0.7. https://CRAN.R-project.org/package=factoextra*.

Keeley, N., Laroche, O., Birch, M. and Pochon, X. (2021) 'A Substrate-Independent Benthic Sampler
(SIBS) for Hard and Mixed-Bottom Marine Habitats: A Proof-of-Concept Study', *Frontiers in
Marine Science,* 8. doi: 10.3389/fmars.2021.627687

Keeley, N., Wood, S. A. and Pochon, X. (2018) 'Development and preliminary validation of a multi-
trophic metabarcoding biotic index for monitoring benthic organic enrichment', *Ecological
Indicators,* 85, pp. 1044-1057. doi: 10.1016/j.ecolind.2017.11.014

King, R. D. and Tyler, P. A. (1982) 'Downstream effects of the Gorden River Power Development, south-
west Tasmania', *Marine and Freshwater Research,* 33(3), pp. 431-442. doi:

Kirkpatrick, J. B., Kriwoken, L. K. and Styger, J. (2019) 'The reverse precautionary principle: science, the
environment and the salmon aquaculture industry in Macquarie Harbour, Tasmania, Australia',
Pacific Conservation Biology, 25(1). doi: 10.1071/pc17014

Koehnken, L. (2005) 'Overview of Water Quality in Macquarie Harbour and Assessment of Risks due
to Copper Levels'. doi:

Kolda, A., Gavrilović, A., Jug-Dujaković, J., Ljubešić, Z., El-Matbouli, M., Lillehaug, A., Lončarević, S.,
Perić, L., Knežević, D., Vukić Lušić, D. and Kapetanović, D. (2020) 'Profiling of bacterial
assemblages in the marine cage farm environment, with implications on fish, human and
ecosystem health', *Ecological Indicators,* 118. doi: 10.1016/j.ecolind.2020.106785

Kopprio, G. A., Cuong, L. H., Luyen, N. D., Duc, T. M., Ha, T. H., Huong, L. M. and Gärdes, A. (2021)
'Carrageenophyte-attached and planktonic bacterial communities in two distinct bays of
Vietnam: Eutrophication indicators and insights on ice-ice disease', *Ecological Indicators,* 121.
doi: 10.1016/j.ecolind.2020.107067

Author (2020) *Tidymodels: a collection of packages for modeling and machine learning using tidyverse
principles. https://www.tidymodels.org*.

Kwon, M. J., O'Loughlin, E. J., Boyanov, M. I., Brulc, J. M., Johnston, E. R., Kemner, K. M. and
Antonopoulos, D. A. (2016) 'Impact of organic carbon electron donors on microbial community
development under iron-and sulfate-reducing conditions', *PloS one,* 11(1), pp. e0146689. doi:
Labbate, M., Seymour, J. R., Lauro, F. and Brown, M. V. (2016) 'Editorial: Anthropogenic Impacts on
the Microbial Ecology and Function of Aquatic Environments', *Front Microbiol,* 7, pp. 1044.
doi: 10.3389/fmicb.2016.01044
Lane, D. (1991) '16S/23S rRNA sequencing', *Nucleic acid techniques in bacterial systematics*, pp. 115-
175. doi:
Lane, D. J., Pace, B., Olsen, G. J., Stahl, D. A., Sogin, M. L. and Pace, N. R. (1985) 'Rapid determination
of 16S ribosomal RNA sequences for phylogenetic analyses', *Proceedings of the National*
Academy of Sciences, 82(20), pp. 6955-6959. doi:
Langille, M. G., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A., Clemente, J. C.,
Burkepile, D. E., Vega Thurber, R. L., Knight, R., Beiko, R. G. and Huttenhower, C. (2013)
'Predictive functional profiling of microbial communities using 16S rRNA marker gene
sequences', *Nat Biotechnol,* 31(9), pp. 814-21. doi: 10.1038/nbt.2676
Laroche, O., Meier, S., Mjøs, S. A. and Keeley, N. (2021) 'Effects of fish farm activities on the sponge
Weberella bursa, and its associated microbiota', *Ecological Indicators,* 129. doi:
10.1016/j.ecolind.2021.107879
Laroche, O., Pochon, X., Tremblay, L. A., Ellis, J. I., Lear, G. and Wood, S. A. (2018) 'Incorporating
molecular-based functional and co-occurrence network properties into benthic marine impact
assessments', *FEMS Microbiol Ecol,* 94(11). doi: 10.1093/femsec/fiy167
Liaw, A. and Wiener, M. (2002) 'Classification and regression by randomForest', *R news,* 2(3), pp. 18-
22. doi:
Lindh, M. V., Lefebure, R., Degerman, R., Lundin, D., Andersson, A. and Pinhassi, J. (2015)
'Consequences of increased terrestrial dissolved organic matter and temperature on
bacterioplankton community composition during a Baltic Sea mesocosm experiment', *Ambio,*
44 Suppl 3, pp. 402-12. doi: 10.1007/s13280-015-0659-3
Lindh, M. V. and Pinhassi, J. (2018) 'Sensitivity of Bacterioplankton to Environmental Disturbance: A
Review of Baltic Sea Field Studies and Experiments', *Frontiers in Marine Science,* 5. doi:
10.3389/fmars.2018.00361
Loch, T., Scribner, K., Tempelman, R., Whelan, G. and Faisal, M. (2012) 'Bacterial infections of Chinook
salmon, Oncorhynchus tshawytscha (Walbaum), returning to gamete collecting weirs in
Michigan', *Journal of Fish Diseases,* 35(1), pp. 39-50. doi:
Logares, R., Sunagawa, S., Salazar, G., Cornejo-Castillo, F. M., Ferrera, I., Sarmento, H., Hingamp, P.,
Ogata, H., de Vargas, C., Lima-Mendez, G., Raes, J., Poulain, J., Jaillon, O., Wincker, P., Kandels-
Lewis, S., Karsenti, E., Bork, P. and Acinas, S. G. (2014) 'Metagenomic 16S rDNA Illumina tags
are a powerful alternative to amplicon sequencing to explore diversity and structure of
microbial communities', *Environ Microbiol,* 16(9), pp. 2659-71. doi: 10.1111/1462-2920.12250
Lokesh, J., Kiron, V., Sipkema, D., Fernandes, J. M. and Moum, T. (2019) 'Succession of embryonic and
the intestinal bacterial communities of Atlantic salmon (Salmo salar) reveals stage‐specific
microbial signatures', *MicrobiologyOpen,* 8(4), pp. e00672. doi:
Louca, S., Parfrey, L. W. and Doebeli, M. (2016) 'Decoupling function and taxonomy in the global ocean
microbiome', *Science,* 353(6305), pp. 1272-1277. doi. Available at:
http://science.sciencemag.org/content/353/6305/1272.long
Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I., Ackermann, M., Hahn,
A. S., Srivastava, D. S., Crowe, S. A., Doebeli, M. and Parfrey, L. W. (2018) 'Function and
functional redundancy in microbial systems', *Nat Ecol Evol,* 2(6), pp. 936-943. doi:
10.1038/s41559-018-0519-1
Love, M. I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2', *Genome Biol,* 15(12), pp. 550. doi: 10.1186/s13059-014-0550-8

Magurran, A. E., Baillie, S. R., Buckland, S. T., Dick, J. M., Elston, D. A., Scott, E. M., Smith, R. I., Somerfield, P. J. and Watt, A. D. (2010) 'Long-term datasets in biodiversity research and monitoring: assessing change in ecological communities through time', *Trends Ecol Evol,* 25(10), pp. 574-82. doi: 10.1016/j.tree.2010.06.016

Mahmoud, M. A. A. and Magdy, M. (2021) 'Metabarcoding profiling of microbial diversity associated with trout fish farming', *Sci Rep,* 11(1), pp. 421. doi: 10.1038/s41598-020-80236-x

Martínez-Porchas, M. and Vargas-Albores, F. (2017) 'Microbial metagenomics in aquaculture: a potential tool for a deeper insight into the activity', *Reviews in Aquaculture,* 9(1), pp. 42-56. doi: 10.1111/raq.12102

Martins, P., Cleary, D. F., Pires, A. C., Rodrigues, A. M., Quintino, V., Calado, R. and Gomes, N. C. (2013) 'Molecular analysis of bacterial communities and detection of potential pathogens in a recirculating aquaculture system for Scophthalmus maximus and Solea senegalensis', *PLoS One,* 8(11), pp. e80847. doi: 10.1371/journal.pone.0080847

Martins, P., Coelho, F. J. R. C., Cleary, D. F. R., Pires, A. C. C., Marques, B., Rodrigues, A. M., Quintino, V. and Gomes, N. C. M. (2018) 'Seasonal patterns of bacterioplankton composition in a semi-intensive European seabass (Dicentrarchus labrax) aquaculture system', *Aquaculture,* 490, pp. 240-250. doi: 10.1016/j.aquaculture.2018.02.038

Martiny, J. B., Bohannan, B. J., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L., Horner-Devine, M. C., Kane, M., Krumins, J. A., Kuske, C. R., Morin, P. J., Naeem, S., Ovreas, L., Reysenbach, A. L., Smith, V. H. and Staley, J. T. (2006) 'Microbial biogeography: putting microorganisms on the map', *Nat Rev Microbiol,* 4(2), pp. 102-12. doi: 10.1038/nrmicro1341

McCaig, A. E., Phillips, C. J., Stephen, J. R., Kowalchuk, G. A., Harvey, S. M., Herbert, R. A., Embley, T. M. and Prosser, J. I. (1999) 'Nitrogen cycling and community structure of proteobacterial β-subgroup ammonia-oxidizing bacteria within polluted marine fish farm sediments', *Applied and Environmental Microbiology,* 65(1), pp. 213-220. doi. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC91005/pdf/am000213.pdf

McMurdie, P. J. and Holmes, S. (2013) 'phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data', *PloS one,* 8(4), pp. e61217. doi:

Minich, J. J., Poore, G. D., Jantawongsri, K., Johnston, C., Bowie, K., Bowman, J., Knight, R., Nowak, B. and Allen, E. E. (2020) 'Microbial ecology of Atlantic salmon (Salmo salar) hatcheries: impacts of the built environment on fish mucosal microbiota', *Applied and environmental microbiology,* 86(12), pp. e00411-20. doi:

Miranda, C. D. and Zemelman, R. (2002) 'Bacterial resistance to oxytetracycline in Chilean salmon farming', *Aquaculture,* 212(1-4), pp. 31-47. doi:

Moncada, C., Hassenruck, C., Gardes, A. and Conaco, C. (2019) 'Microbial community composition of sediments influenced by intensive mariculture activity', *FEMS Microbiol Ecol,* 95(2). doi: 10.1093/femsec/fiz006

Morelan, I. A. (2019) '16S rRNA Gene Amplicon Sequencing Reveals Trends in Marine Bacterial Diversity and Taxonomic Composition in Natural and Human-built Systems'. doi:

Mukherjee, A., Chettri, B., Langpoklakpam, J. S., Basak, P., Prasad, A., Mukherjee, A. K., Bhattacharyya, M., Singh, A. K. and Chattopadhyay, D. (2017) 'Bioinformatic approaches including predictive metagenomic profiling reveal characteristics of bacterial response to petroleum hydrocarbon contamination in diverse environments', *Scientific reports,* 7(1), pp. 1-22. doi:

Nagpal, S., Haque, M. M. and Mande, S. S. (2016) 'Vikodak--A Modular Framework for Inferring Functional Potential of Microbial Communities from 16S Metagenomic Datasets', *PLoS One,* 11(2), pp. e0148347. doi: 10.1371/journal.pone.0148347

Navarro, N., Leakey, R. J. G. and Black, K. D. (2008) 'Effect of salmon cage aquaculture on the pelagic environment of temperate coastal waters: seasonal changes in nutrients and microbial community', *Marine Ecology Progress Series,* 361, pp. 47-58. doi. Available at: https://www.int-res.com/abstracts/meps/v361/p47-58/

963   Newell, S. E., Fawcett, S. E. and Ward, B. B. (2013) 'Depth distribution of ammonia oxidation rates and
964   ammonia-oxidizer community composition in the Sargasso Sea', *Limnology and Oceanography,*
965   58(4), pp. 1491-1500. doi: 10.4319/lo.2013.58.4.1491

966   Oksanen, J. F., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Peter, R. and Minchin, R.
967   (2018) 'O'Hara; Gavin, L.; et al', *The Vegan Community Ecology Package. Available online:*
968   *https://cran. r-project. org/web/packages/vegan/(accessed on 1 March 2019)*. doi:

969   Olsen, L. M., Hernández, K. L., Ardelan, M. V., Iriarte, J. L., Bizsel, K. C. and Olsen, Y. (2017) 'Responses
970   in bacterial community structure to waste nutrients from aquaculture: an in situ microcosm
971   experiment in a Chilean fjord', *Aquaculture Environment Interactions,* 9, pp. 21-32. doi.
972   Available at: https://www.int-res.com/abstracts/aei/v9/p21-32/

973   Ortiz-Estrada, Á. M., Gollas-Galván, T., Martínez-Córdova, L. R. and Martínez-Porchas, M. (2019)
974   'Predictive functional profiles using metagenomic 16S rRNA data: a novel approach to
975   understanding the microbial ecology of aquaculture systems', *Reviews in Aquaculture,* 11(1),
976   pp. 234-245. doi: 10.1111/raq.12237

977   Pace, N. R. (1997) 'A Molecular View of Microbial Diversity and the Biosphere', *Science,* 276(5313), pp.
978   734-740. doi: 10.1126/science.276.5313.734

979   Palarea-Albaladejo, J. and Martín-Fernández, J. A. (2015) 'zCompositions—R package for multivariate
980   imputation of left-censored data under a compositional approach', *Chemometrics and*
981   *Intelligent Laboratory Systems,* 143, pp. 85-96. doi:

982   Pineiro, S. A., Stine, O. C., Chauhan, A., Steyert, S. R., Smith, R. and Williams, H. N. (2007) 'Global survey
983   of diversity among environmental saltwater Bacteriovoracaceae', *Environ Microbiol,* 9(10), pp.
984   2441-50. doi: 10.1111/j.1462-2920.2007.01362.x

985   Pomeroy, L. R., leB. WILLIAMS, P. J., Azam, F. and Hobbie, J. E. (2007) 'The microbial loop',
986   *Oceanography,* 20(2), pp. 28-33. doi:

987   Qu, J., Yang, H., Liu, Y., Qi, H., Wang, Y. and Zhang, Q. (2021) 'The study of natural biofilm formation
988   and microbial community structure for recirculating aquaculture system', *IOP Conference*
989   *Series: Earth and Environmental Science,* 742(1). doi: 10.1088/1755-1315/742/1/012018

990   R Core Team 2021. R: A language and environment for statistical computing. R Foundation for
991   Statistical Computing, Vienna,
992   Austria. URL https://www.R-project.org/.

993   Raes, E. J., Karsh, K., Sow, S. L. S., Ostrowski, M., Brown, M. V., van de Kamp, J., Franco-Santos, R. M.,
994   Bodrossy, L. and Waite, A. M. (2021) 'Metabolic pathways inferred from a bacterial marker
995   gene illuminate ecological changes across South Pacific frontal boundaries', *Nat Commun,*
996   12(1), pp. 2213. doi: 10.1038/s41467-021-22409-4

997   Rasigraf, O., Schmitt, J., Jetten, M. S. M. and Luke, C. (2017) 'Metagenomic potential for and diversity
998   of N-cycle driving microorganisms in the Bothnian Sea sediment', *Microbiologyopen,* 6(4). doi:
999   10.1002/mbo3.475

1000   Reji, L., Tolar, B. B., Chavez, F. P. and Francis, C. A. (2020) 'Depth-Differentiation and Seasonality of
1001   Planktonic Microbial Assemblages in the Monterey Bay Upwelling System', *Front Microbiol,*
1002   11, pp. 1075. doi: 10.3389/fmicb.2020.01075

1003   Revill, A. T., Ross, J. a. and Thompson, P. A. (2016) *Investigating Dissolved Oxygen Drawdown in*
1004   *Macquarie Harbour. CSIRO, Australia. Report to Huon Aquaculture*: Institute for Marine and
1005   Antarctic Studies (IMAS), Hobart. Available at: https://www.huonaqua.com.au/wp-
1006   content/uploads/2015/08/Appendix-4-part-4.pdf (Accessed: 16 September 2019).

1007   Ross, J., Beard, J., Wild-Allen, K., Andrewartha, Moreno, D., Semmens, J., Davey, A., Hortle, J., Pender,
1008   Quigley B., Andrewartha, J., Stehfest, K., Durand, A. and Macleod, C. a. (2021) *Understanding*
1009   *oxygen dynamics and the importance for benthic recovery in Macquarie Harbour. Hobart,*
1010   *Australia CC BY 3.0*: Institute for Marine and Antarctic Studies, University of Tasmania.

1011   Rubio-Portillo, E., Villamor, A., Fernandez-Gonzalez, V., Antón, J. and Sanchez-Jerez, P. (2019)
1012   'Exploring changes in bacterial communities to assess the influence of fish farming on marine
1013   sediments', *Aquaculture,* 506, pp. 459-464. doi: 10.1016/j.aquaculture.2019.03.051

Sanderson, J. C., Cromey, C. J., Dring, M. J. and Kelly, M. S. (2008) 'Distribution of nutrients for seaweed cultivation around salmon cages at farm sites in north–west Scotland', *Aquaculture,* 278(1-4), pp. 60-68. doi: 10.1016/j.aquaculture.2008.03.027

Sarà, G., Lo Martire, M., Buffa, G., Mannino, A. M. and Badalamenti, F. (2007) 'The fouling community as an indicator of fish farming impact in Mediterranean', *Aquaculture Research,* 38(1), pp. 66-75. doi: 10.1111/j.1365-2109.2006.01632.x

Sevigny, J. L., Rothenheber, D., Diaz, K. S., Zhang, Y., Agustsson, K., Bergeron, R. D. and Thomas, W. K. (2019) 'Marker genes as predictors of shared genomic function', *BMC Genomics,* 20(1), pp. 268. doi: 10.1186/s12864-019-5641-1

Sinnott, R. W. (1984) 'Virtues of the Haversine', *S&T,* 68(2), pp. 158. doi:

Stoeck, T., Fruhe, L., Forster, D., Cordier, T., Martins, C. I. M. and Pawlowski, J. (2018) 'Environmental DNA metabarcoding of benthic bacterial communities indicates the benthic footprint of salmon aquaculture', *Mar Pollut Bull,* 127, pp. 139-149. doi: 10.1016/j.marpolbul.2017.11.065

Sun, S., Jones, R. B. and Fodor, A. A. (2020) 'Inference-based accuracy of metagenome prediction tools varies across sample types and functional categories', *Microbiome,* 8(1), pp. 46. doi: 10.1186/s40168-020-00815-y

Sun, X., Dong, J., Hu, C., Zhang, Y., Chen, Y. and Zhang, X. (2021) 'Use of macrofaunal assemblage indices and biological trait analysis to assess the ecological impacts of coastal bivalve aquaculture', *Ecological Indicators,* 127, pp. 107713. doi:

Tamminen, M., Karkman, A., Corander, J., Paulin, L. and Virta, M. (2011) 'Differences in bacterial community composition in Baltic Sea sediment in response to fish farming', *Aquaculture,* 313(1-4), pp. 15-23. doi: 10.1016/j.aquaculture.2011.01.020

Teasdale, P. R., Apte, S. C., Ford, P. W., Batley, G. E. and Koehnken, L. (2003) 'Geochemical cycling and speciation of copper in waters and sediments of Macquarie Harbour, Western Tasmania', *Estuarine, Coastal and Shelf Science,* 57(3), pp. 475-487. doi: 10.1016/s0272-7714(02)00381-5

Teichert, N., Lepage, M. and Lobry, J. (2018) 'Beyond classic ecological assessment: the use of functional indices to indicate fish assemblages sensitivity to human disturbance in estuaries', *Science of the Total Environment,* 639, pp. 465-475. doi:

Terry, C. (2001) 'Numerical Modelling of Macquarie Harbour, Tasmania', *In: Conference on Hydraulics in Civil Engineering (6th : 2001 : Hobart, Tas.). 6th Conference on Hydraulics in Civil Engineering: The State of Hydraulics; Proceedings. Barton, A.C.T.: Institution of Engineers, Australia, 2001: 345-354.*

Tsagaraki, T. M., Petihakis, G., Tsiaras, K., Triantafyllou, G., Tsapakis, M., Korres, G., Kakagiannis, G., Frangoulis, C. and Karakassis, I. (2011) 'Beyond the cage: ecosystem modelling for impact evaluation in aquaculture', *Ecological modelling,* 222(14), pp. 2512-2523. doi:

Tsukamoto, K., Kawamura, T., Takeuchi, T., Beard Jr, T. and Kaiser, M. (2008) 'Environmental Impact of Aquaculture on Coastal Planktonic Ecosystems', *Fisheries for Global Welfare and Environment*, pp. 181. doi:

Vargas-Albores, F., Martínez-Córdova, L. R., Gollas-Galván, T., Garibay-Valdez, E., Emerenciano, M. G. C., Lago-Leston, A., Mazorra-Manzano, M. and Martínez-Porchas, M. (2019) 'Inferring the functional properties of bacterial communities in shrimp-culture bioflocs produced with amaranth and wheat seeds as fouler promoters', *Aquaculture,* 500, pp. 107-117. doi: 10.1016/j.aquaculture.2018.10.005

Verhoeven, J. T. P., Salvo, F., Hamoutene, D. and Dufour, S. C. (2016) 'Bacterial community composition of flocculent matter under a salmonid aquaculture site in Newfoundland, Canada', *Aquaculture Environment Interactions,* 8, pp. 637-646. doi: 10.3354/aei00204

Verhoeven, J. T. P., Salvo, F., Knight, R., Hamoutene, D. and Dufour, S. C. (2018) 'Temporal Bacterial Surveillance of Salmon Aquaculture Sites Indicates a Long Lasting Benthic Impact With Minimal Recovery', *Front Microbiol,* 9, pp. 3054. doi: 10.3389/fmicb.2018.03054

Wakabayashi, H. (1980) 'Bacterial gill disease of salmonid fish', *Fish pathology,* 14(4), pp. 185-189. doi:

Wang, Q., Garrity, G. M., Tiedje, J. M. and Cole, J. R. (2007) 'Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy', *Appl Environ Microbiol,*
73(16), pp. 5261-7. doi: 10.1128/AEM.00062-07

Wang, X., Olsen, L. M., Reitan, K. I. and Olsen, Y. (2012) 'Discharge of nutrient wastes from salmon
farms: environmental effects, and potential for integrated multi-trophic aquaculture',
Aquaculture Environment Interactions, 2(3), pp. 267-283. doi: 10.3354/aei00044

Wei, G., Li, M., Li, F., Li, H. and Gao, Z. (2016) 'Distinct distribution patterns of prokaryotes between
sediment and water in the Yellow River estuary', *Appl Microbiol Biotechnol,* 100(22), pp. 9683-
9697. doi: 10.1007/s00253-016-7802-3

Wemheuer, F., Taylor, J. A., Daniel, R., Johnston, E., Meinicke, P., Thomas, T. and Wemheuer, B. (2020)
'Tax4Fun2: prediction of habitat-specific functional profiles and functional redundancy based
on 16S rRNA gene sequences', *Environ Microbiome,* 15(1), pp. 11. doi: 10.1186/s40793-020-
00358-7

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A.,
Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson,
D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K. and Yutani, H. (2019)
'Welcome to the Tidyverse', *Journal of Open Source Software,* 4(43), pp. 1686. doi:
10.21105/joss.01686

Won, N. I., Kim, K. H., Kang, J. H., Park, S. R. and Lee, H. J. (2017) 'Exploring the Impacts of
Anthropogenic Disturbance on Seawater and Sediment Microbial Communities in Korean
Coastal Waters Using Metagenomics Analysis', *Int J Environ Res Public Health,* 14(2), pp. 130.
doi: 10.3390/ijerph14020130

Wu, D. M., Dai, Q. P., Liu, X. Z., Fan, Y. P. and Wang, J. X. (2019) 'Comparison of bacterial community
structure and potential functions in hypoxic and non-hypoxic zones of the Changjiang Estuary',
PLoS One, 14(6), pp. e0217431. doi: 10.1371/journal.pone.0217431

Wu, R. S. S. (1995) 'The environmental impact of marine fish culture: Towards a sustainable future',
Marine Pollution Bulletin, 31(4), pp. 159-166. doi: https://doi.org/10.1016/0025-
326X(95)00100-2

Yang, Y., Gao, Y., Huang, X., Ni, P., Wu, Y., Deng, Y. and Zhan, A. (2019) 'Adaptive shifts of
bacterioplankton communities in response to nitrogen enrichment in a highly polluted river',
Environ Pollut, 245, pp. 290-299. doi: 10.1016/j.envpol.2018.11.002

Yarza, P., Yilmaz, P., Pruesse, E., Glockner, F. O., Ludwig, W., Schleifer, K. H., Whitman, W. B., Euzeby,
J., Amann, R. and Rossello-Mora, R. (2014) 'Uniting the classification of cultured and
uncultured bacteria and archaea using 16S rRNA gene sequences', *Nat Rev Microbiol,* 12(9),
pp. 635-45. doi: 10.1038/nrmicro3330

Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig,
W. and Glöckner, F. O. (2014) 'The SILVA and "all-species living tree project (LTP)" taxonomic
frameworks', *Nucleic acids research,* 42(D1), pp. D643-D648. doi:

Yoshikawa, T. and Eguchi, M. (2013) 'Planktonic processes contribute significantly to the organic
carbon budget of a coastal fish-culturing area', *Aquaculture Environment Interactions,* 4(3), pp.
239-250. doi:

Yu, Z., Yang, J., Amalfitano, S., Yu, X. and Liu, L. (2014) 'Effects of water stratification and mixing on
microbial community structure in a subtropical deep reservoir', *Sci Rep,* 4, pp. 5821. doi:
10.1038/srep05821

Zarkasi, K. Z., Taylor, R. S., Abell, G. C., Tamplin, M. L., Glencross, B. D. and Bowman, J. P. (2016)
'Atlantic salmon (Salmo salar L.) gastrointestinal microbial community dynamics in relation to
digesta properties and diet', *Microbial ecology,* 71(3), pp. 589-603. doi:

Zhang, K., Zheng, X., He, Z., Yang, T., Shu, L., Xiao, F., Wu, Y., Wang, B., Li, Z., Chen, P. and Yan, Q. (2020)
'Fish growth enhances microbial sulfur cycling in aquaculture pond sediments', *Microb
Biotechnol,* 13(5), pp. 1597-1610. doi: 10.1111/1751-7915.13622

Zhang, Q., Tang, F., Zhou, Y., Xu, J., Chen, H., Wang, M. and Laanbroek, H. J. (2015) 'Shifts in the pelagic
1116   ammonia-oxidizing microbial communities along the eutrophic estuary of Yong River in
1117   Ningbo City, China', *Front Microbiol,* 6, pp. 1180. doi: 10.3389/fmicb.2015.01180
1118   Zorz, J., Willis, C., Comeau, A. M., Langille, M. G. I., Johnson, C. L., Li, W. K. W. and LaRoche, J. (2019)
1119   'Drivers of Regional Bacterial Community Structure and Diversity in the Northwest Atlantic
1120   Ocean', *Front Microbiol,* 10, pp. 281. doi: 10.3389/fmicb.2019.00281
1121