The impact of SNP density on quantitative genetic analyses of body size traits in a wild population of Soay sheep

Caelinn James[1*], Josephine M. Pemberton[1], Pau Navarro[2], Sara Knott[1]

[1] Institute of Evolutionary Biology, School of Biological Sciences, The University of Edinburgh, Edinburgh

[2] MRC Institute of Genetics and Cancer, The University of Edinburgh, Edinburgh

[*] Corresponding author

Corresponding author mailing address: Caelinn James, Office 133, Ashworth Laboratories, Kings Buildings, Charlotte Auerbach Road, Edinburgh EH9 3FL

Corresponding author email address: c.james-5@sms.ed.ac.uk

Running head: Impact of SNP density on quantitative analyses

Abstract

Understanding the genetic architecture underpinning quantitative traits in wild populations is pivotal to understanding the processes behind trait evolution. The 'animal model' is a popular method for estimating quantitative genetic parameters such as heritability and genetic correlation and involves fitting an estimate of relatedness between individuals in the study population. Genotypes at genome-wide markers can be used to estimate relatedness; however, relatedness estimates vary with marker density, potentially affecting results. Increasing density of markers is also expected to increase the power to detect quantitative trait loci (QTL). We estimated heritability and performed genome-wide association studies (GWAS) on five body size traits in an unmanaged population of Soay sheep using two different SNP densities: a dataset of 37,037 genotyped SNPs, and an imputed dataset of 417,373 SNPs. Heritability estimates did not differ between the two SNP densities, but the high-density imputed SNP dataset revealed five new SNP-trait associations that were not found with the lower density dataset. Conditional GWAS analyses after fitting the most significant SNPs revealed two more novel SNP-trait associations.

Introduction

Investigating the genetic architecture behind heritable traits is key to understanding the biological diversity of wild populations. If we know the number of loci influencing a trait and their effect size, we can better understand the evolutionary processes that underpin traits, improve inferences about trait evolution (Barton and Keightley 2002), and understand micro-evolutionary dynamics that occur due to environmental change. Most quantitative genetic research in animals is carried out in artificial populations; either domestic, agricultural or laboratory. Such populations experience controlled environmental conditions which make it easier to account for environmental factors when studying the effect of genetic variants on phenotypic variation. However, given that environmental factors can influence the phenotype of a quantitative trait (Charmantier et al. 2014), and that the presence of genotype-by-environment interactions can cause additive genetic variance to differ between environmental conditions, studies on artificial populations arguably cannot be fully extrapolated to wild populations (Kruuk et al. 2008). Therefore, it is important to also study quantitative traits in wild populations in their natural habitats. There is a wealth of quantitative genetics research in human populations (for examples, see (Kang et al. 2010; Locke et al. 2015; Manolio et al. 2009; Xia et al. 2016; Xia et al. 2021; Yang et al. 2010; Zaitlen et al. 2013)), but humans also arguably experience a more buffered environment than wild populations.

A popular method to decompose phenotypic variation in wild populations into genetic variance and environmental variance is the 'animal model', originally developed by animal breeders (Henderson 1984; Kruuk 2004; Wilson et al. 2010). As part of the model, genetic relatedness is fitted, which is often derived from a pedigree. Pedigrees can be constructed using field observations, assigning parentage using genetic markers, or a mixture of both (Pemberton 2008). However, wild pedigrees are often short, incomplete, and contain errors: observational data may be inaccurate due to incorrect parent-offspring assumptions, and if the genetic markers chosen are not sufficiently discriminatory they may result in misassigned parentage. Erroneous pedigree links can bias results of analyses using animal models; for example, misidentification of sires in cattle resulted in decreased heritability estimates for milk yield, fat yield, and milk-fat ratio (Van Vleck 1970).

In place of a pedigree, genotypes at multiple polymorphic loci can be used to estimate relatedness for use in an animal model. This has the advantage of not relying on recovering a pedigree and thus is not affected by incomplete or incorrect familial links (though knowledge of pedigree relationships is still valuable, for example so that maternal effects can be fitted). Relatedness estimated using genotype data is also potentially more precise than that from pedigrees – for example, with a pedigree it is presumed that full-sibs have a relatedness of 0.5, however the exact relationship varies depending on which DNA segments each sib has inherited (Visscher et al. 2006). Despite this greater accuracy, genotype-based relatedness estimates can still vary depending on which variants in the population have been genotyped, and the density of the genotyped variants. Increasing the density of genotyped polymorphisms means they are more likely to be in linkage disequilibrium (LD) with causal variants for the trait of interest, either by being physically closer to the causal variants or by matching the allele frequency of the causal variants more accurately. Thus, in species such as humans where genotyping is commonly of unrelated individuals and LD is generally low, the estimated heritability of a trait increases with SNP density due to an increase in the number of causal variants being in LD with genotyped SNPs (for instance, heritability of human height was estimated to be 0.45 when using 294,831 SNPs (Yang et

al. 2010) and 0.56 when using ~17 million imputed SNPs (Yang et al. 2015)). However, increasing the
number of genotyped markers means larger, denser genotyping arrays with prices increasing with
density. For commonly studied species, high-density arrays are more affordable due to high demand,
but for more niche species, including wild populations, large genotyping arrays are often unaffordable.
Genotyping-by-sequencing, e.g. ddRAD (Peterson et al. 2012) is a potentially useful alternative for
upscaling SNP density, though the combination of bioinformatics and samples sizes required in
quantitative genetic research means that this approach is not yet in widespread use.

As an alternative to expensive high-density genotyping, genotype imputation can be used to increase
the number of variants analysed (Burdick et al. 2006). Imputation involves predicting genotypes at
untyped SNPs in a 'target' population using a subset of the study population – or more generally a
"reference" population – genotyped at a higher density, either through a high density SNP array or by
genotyping-by sequencing. The genotypes at these untyped SNPs for individuals in the target population
are inferred using their genotypes at typed markers and taking advantage of existing linkage
disequilibrium (LD) between SNPs. Pedigree information can also be used to increase the accuracy of the
imputation by identifying haplotype blocks that are identical by descent (Burdick et al. 2006).

The Soay sheep *(Ovis aries)* of St Kilda are a primitive, unmanaged breed of sheep that have been the
focus of a longitudinal, individual-based study since 1985 (Soay sheep: Dynamics and selection in an
island population 2003). As part of the study, life history and environmental data is collected, DNA
samples are collected, and a pedigree has been constructed using observation and genetic parentage
inference. 7630 sheep have been genotyped on the Ovine SNP50 Illumina Beadchip, on which 37,037
SNPs are autosomal and polymorphic in this population.

To date, quantitative genetic analyses of the Soays have been performed using either the pedigree or
the 50K SNP data. Bérénos et al. (2014) (Bérénos et al. 2014) investigated the difference in quantitative
genetic parameter estimates when using the pedigree or a genomic relationship matrix (GRM)
constructed from the 50K SNP data. The authors estimated heritability, maternal genetic effects and
genetic correlations for body size traits (weight, foreleg length, hindleg length, metacarpal length and
jaw length) across four age groups. The additive genetic variance and the heritability estimates using the
GRM were lower than when using the pedigree to estimate relatedness, with the SNPs explaining 84% of
the additive genetic variance of the pedigree on average, though for the majority of the traits the
standard errors of the pedigree-based and SNP-based heritability estimates were within one standard
error of each other. Genetic correlations were found to differ little between analyses using the different
relatedness estimates. A SNP rarefaction analysis of the heritability estimates was conducted on the
adult traits, and it was found that heritability estimates asymptoted when 50% of the SNPs were used to
estimate relatedness.

Linkage disequilibrium is high in Soay sheep (Bérénos et al. 2014), which may explain the results of the
rarefaction analysis: the same causal variations can be represented by multiple genotyped SNPs, and so
many genotyped SNPs are not providing unique information towards the additive genetic variance.
However, given that heritability estimates when using SNP-based estimates of relatedness were lower
than when using pedigree-based estimates, it is also possible that there are variants that contribute to
variation in these traits that aren't in perfect LD with the genotyped SNPs. This may be depressing
estimates of additive genetic variance in comparison to pedigree-based estimates. Increasing the

density of genotyped SNPs may therefore increase the SNP-based heritability estimates of these traits to
closer match the pedigree-based estimates.

Bérénos et al. (2015) (Bérénos et al. 2015) focused on the same traits as (Bérénos et al. 2014) in adults
only, and partitioned the genetic variance for each trait, first by chromosome and then by 150 SNP
windows, and performed GWAS. For the three leg length phenotypes, a disproportionately high amount
of variance was explained by two SNPs on chromosome 16 (s23172.1) and chromosome 19 (s74894.1).
For the remaining traits, the proportion of additive genetic variance explained by a chromosome was
proportional to its length, suggesting that variation in these traits is influenced by many genetic variants
of small effect, and no SNP-trait associations were discovered. The authors performed a two-step GWAS
analysis; first they modelled the traits by fitting fixed and random effects, and then they extracted the
residuals from the models and tested for association between SNPs and the residuals. Three traits
(weight, foreleg length and hindleg length) are measured in the same individual across multiple years –
for these traits, the authors analysed the mean residual values.

Since the studies from 2014 and 2015, 1895 more individuals have been both phenotyped and
genotyped on the Ovine SNP50 Illumina Beadchip. In addition, 188 individuals were genotyped on the
Ovine Infinium High Density chip with 600K attempted SNPs, which has enabled the genotypes of the
remaining individuals to be accurately imputed to that higher density using LD and pedigree data (Stoffel
et al. 2021).

In this study we performed a direct comparison of heritability estimates and GWAS associations
between the lower density SNP data and the imputed high density SNP data in the Soay population
using an increased sample size and compared the results to (Bérénos et al. 2015; Bérénos et al. 2014).
We focused on the same five traits as the previous studies in neonates, lambs and adults. Unlike the
2015 study (Bérénos et al. 2015), we performed GWAS by fitting fixed and random effects in the same
step as testing for SNP-trait associations. This has the advantage of correctly propagating error
throughout the analysis, reducing the chance of false positive results. We also carried out a two-step
GWAS approach similar to that of (Bérénos et al. 2015), focusing on the adult traits using the 50K SNP
data, to investigate whether any SNP-trait associations identified using our approach were due to the
increased population size or due to the different methodology (single-step vs. two-step GWAS).

Our aims were as follows:

1) To determine whether the increased density of SNPs changes the heritability estimates of the
traits using the same individuals for both the low SNP density and the high SNP density analysis.
Given the previous rarefaction analysis (Bérénos et al. 2014), our prediction was that there
would be no change.
146
147 2) To determine whether the imputed SNP data enables the identification of new SNP-trait
148   associations via GWAS. We predicted we would find more associations, either due to increased
149   power to detect small effect size SNPs through increased LD with the imputed SNPs or due to
150   tagging of new causal variants.
151
3) To examine how a single-step GWAS methodology compares the two-step approach previously
used on the Soay population, to investigate whether any novel SNP-trait associations identified

with the 50K SNP data were due to increased population size or due to the difference in
methods.

METHODS

Phenotypic data

We focused on five body size traits in three age groups: neonates, lambs, and adults. Of the five traits,
three (weight, foreleg length and hindleg length) are live measures, recorded in April for neonates and in
August for lambs and adults. The remaining two traits (metacarpal length and jaw length) are *post*
mortem measures taken from skeletal material. The sheep are ear-tagged when they are first captured
which allows for reidentification for life. Both birth and August weight are measured to the nearest
0.1kg, whilst the remaining traits are all measured to the nearest mm. A detailed description of trait
measurements can be found in (Beraldi et al. 2007).

We defined neonates as individuals who were caught and weighed between two and ten days after birth
– birth weight was the only trait recorded for this age group. Lambs were classed as individuals who had
phenotypic data recorded in the August of their birth year for the live traits, and as individuals who died
before 14 months of age for the *post mortem* measures. Individuals were classed as adults if they had
August phenotypic data recorded at least two years after birth, or if they died after 26 months of age for
post mortem measures. Unlike (Bérénos et al. 2014), we chose not to analyse yearling data due to the
small sample sizes in comparison to the other age classes, which is due to high first winter mortality.

Genetic data

Most of the sheep in our study population have been genotyped using the Ovine SNP50 Illumina
BeadChip, which targets 54,241 SNPs across the sheep genome. After removing SNPs which failed
quality control standards (minor allele frequency (MAF) > 0.001, call rate > 0.99, deviation from Hardy-
Weinberg Equilibrium P > 1e-05), 39,368 polymorphic variants remained for 7630 individuals (3643
female, 3987 male). See Bérénos et al. 2014 (Bérénos et al. 2014) for information on genetic sampling
protocol and marker characteristics).

Of these 7630 individuals, 188 have also been genotyped using the Ovine Infinium HD SNP BeadChip
which types 606,066 SNPs, of which 417,373 were autosomal and polymorphic in the Soays. This has
allowed for the low density genotypes to be imputed to the higher density using AlphaImpute, which
combines shared haplotype and pedigree information for phasing and genotype imputation (Hickey et
al. 2012) (see Stoffel et al. (2021)(Stoffel et al. 2021) for information on imputation). We used imputed
genotype "hard" calls (rather than genotype probabilities) in downstream analyses. After filtering SNPs
that failed quality control standards (minor allele frequency (MAF) > 0.001, call rate > 0.99, deviation
from Hardy-Weinberg Equilibrium P > 1e-05) and imputation QC, 343,889 autosomal SNPs remained for
7362 individuals (3529 females, 3736 males).

Both the 50K SNP data and the imputed SNP data are mapped to the OAR_v3.1 genome assembly.

Narrow sense heritability estimation

We used animal models to partition the phenotypic variance for each trait in each age class into genetic
and non-genetic variance components. Fixed and random effects were fitted for all models, with the

effects differing between traits and age classes (Table 1). We implemented these analyses in DISSECT
(Canela-Xandri et al. 2015) using the following model:

$$y = X\beta + \sum_r Z_r u_r + Wg + \varepsilon$$

where y is the vector of phenotypic values; X is a design matrix linking individual records with the vector
of fixed effects β, Z_r is an incidence matrix that relates a random effect to the individual records; u_r is the
associated vector of non-genetic random effects; g is the vector of additive genetic random effects with
W the incidence matrix; and ε is the vector of residuals. It is assumed that g ~ MVN(0, $M\sigma_g^2$), where σ_g^2 is
the additive genetic variance and M is the genomic relationship matrix (GRM). For each trait in each age
class, we ran this model twice: first with M being a GRM calculated from the 50K genotype data, and
second with M being a GRM calculated from the imputed SNP genotypes. The GRMs were computed
using DISSECT(Canela-Xandri et al. 2015), and the genetic relationship between individuals i and j is
computed as:

$$A_{ij} = \frac{1}{N} \sum_{k=1}^{N} \frac{(s_{ik} - 2p_k)(s_{jk} - 2p_k)}{2p_k(1 - p_k)}$$

where s_{ik} is the number of copies of the reference allele for SNP k of the individual i, p_k is the frequency
of the reference allele for the SNP k, and N is the number of SNPs.

The narrow sense heritability was estimated by dividing the additive genetic variance (the variance
explained by the GRM) by the total estimated phenotypic variance (the sum of the variance explained by
the GRM and other fitted random effects after fitting fixed effects).

In adults, there are multiple records for August weight, foreleg length and hindleg length for the same
individual due to individuals being caught across multiple years. For these traits we used a repeatability
model in order that uncertainty was correctly propagated through all estimations (Mrode 2014). To
implement a repeatability model in DISSECT, we edited the input files so that each measurement had its
own row in the genotype and covariate files. Individual ID was replaced with a unique capture reference
number, and individual permanent environment was fitted as a random effect (see Supplementary
Methods for a more detailed explanation).

Sample sizes and total number of phenotypic measurements for all traits are shown in Table 1, with
effects fitted in all models.

Genome wide association analysis

We also conducted genome-wide association analyses using DISSECT (Canela-Xandri et al. 2015). We
fitted the same fixed and random effects for each trait and age class as for the heritability estimation
(Table 1). To account for population structure, when testing SNPs on a given chromosome for

association with the phenotype, a GRM calculated from the remaining autosomes (referred to as Leave
One Chromosome Out GRM (Yang et al. 2014)) was fitted. Input files for repeated-measure traits were
reformatted as above. Our significance threshold was corrected for multiple testing using the SimpleM
(Gao et al. 2008) method, which accounts for linkage disequilibrium between markers in order to
calculate the effective number of independent tests.

We estimated the variance explained by SNPs that passed the significance threshold using the equation

$$V(SNP) = 2pq\alpha^2$$

where p and q are the major and minor allele frequencies of the SNP, and α is the estimated SNP effect.
We then calculated the proportion of additive genetic variance explained by each SNP by dividing by the
total additive genetic variance estimated for that trait.

For any trait for which several SNPs in the same region were associated with variation in the trait and
thus had strong support for at least one QTL in the region, we carried out conditional analysis to
understand if the region could harbour potentially several independent QTL, or if further QTL could be
uncovered elsewhere in the genome. To that aim, the genotypes of the SNP with the smallest
association p value from each associated region (hereafter called the "top SNP") were added to the
GWAS model as a fixed covariate and removed from the GRMs and genotype data. The GWAS analysis
was re-run accounting for those associations to try and reveal novel peaks either in the same regions or
elsewhere in the genome.

Two-step GWAS analysis

To investigate whether any novel SNP associations identified (since (Bérénos et al. 2015)) by performing
GWAS on the adult traits using the 50K SNP data were due to the increased population sample or due to
the change in methodology, we also performed a two-step GWAS, focusing on adults only and using the
50K SNP data. We performed mixed model analyses using ASReml-R (Butler et al. 2017) for each trait
fitting the same fixed and random effects as in our single-step analyses, including whole-genome
relatedness (in the form of a GRM) and, for repeated-measure traits, permanent environment. We then
extracted the residuals from the mixed models and performed GWAS with the residuals as the trait
phenotypes using DISSECT (Canela-Xandri et al. 2015). For repeated measure traits, we used the mean
residual value for each individual. We used the Bonferroni correction calculated in (Bérénos et al. 2015)
to determine the significance threshold for our two-step GWAS in order to best compare with previous
GWAS performed on the Soay sheep.

RESULTS

Heritability estimation

Neonates

In neonates, the heritability of birth weight was 0.058 (S.E. 0.021) when using the 50K SNPs to calculate
relatedness, and 0.049 (S.E. 0.020) when using the imputed SNPs (Figure 1, Supplementary Table 1). The
two estimates were within one standard error of each other, meaning the difference between the two
estimates was not significant.

Lambs

In lambs, the heritability estimates for the live August measures were lower than those for the *post*
mortem measures (Figure 1, Supplementary Table 1). Across all the traits, heritability estimates were
similar when using the 50K SNP data and the imputed SNP data, with the biggest difference being 0.016
for jaw length. For all traits, estimates were within one standard error of each other, meaning that the
small differences in heritability estimates between the two SNP densities were not significant.

Adults

As observed in lambs, heritability estimates for live measures in adults were lower than those of the
post mortem measures. Across all traits, heritability estimates were higher in adults than in lambs.
Estimates obtained using the 50K SNPs and using the imputed SNPs were similar and were within one
standard error of each other (Figure 1, Supplementary Table 1), meaning that the imputed SNPs
provided no additional information to partition the variation into genetic and environmental variance.

Estimates for all variance components are listed in Supplementary Table 1.

GWAS

50K SNP data

To correct for multiple testing, we calculated the effective number of tests to be 20082 using the
SimpleM method (Gao et al. 2008), giving a genome-wide significance threshold of $2.49e^{-06}$ for the 50K
SNP data.

For weight in neonates (birth weight), lambs and adults (August weight), no SNPs were found to have an
association p value smaller than this threshold, suggesting that any variants that influence weight
variation are either of small effect or were not tagged by SNPs in the 50K SNP data (Figure 2A,
Supplementary Figure 1B and 1G).

For all three leg length measures in lambs, we found associations with the same region on chromosome
16. SNP s23172.1 was the SNP with the lowest p value for lamb foreleg and hindleg, explaining 0.45%
and 0.54% of the genetic variance for each trait respectively (Supplementary Table 2, Supplementary
Figure 1C and 1D). For lamb metacarpal, SNP 22142.1 in the same chromosome 16 region had the
lowest p value and explained 0.98% of the genetic variance. There was also a single SNP on chromosome
3 (OAR3_100483326.1) and a cluster of SNPs on chromosome 19 that had p values smaller than the
genome-wide significance threshold and were associated with variation in lamb metacarpal length,

explaining 1.95% and 2.32% of the genetic variance respectively (Supplementary Table 2, Supplementary
Figure 1E).

The two regions on chromosomes 16 and 19 that were associated with lamb metacarpal length variation
were also significantly associated with all three leg length measures in adults, with SNP s22142.1 on
chromosome 16 and SNP s74894.1 on chromosome 19 respectively explaining 0.74% and 1.96% of the
genetic variation in adult foreleg, 0.83% and 1.44% of the genetic variation in adult hindleg, and 1.08%
and 1.96% of the genetic variation in adult metacarpal length. There were other regions of the genome
also associated with variation in the adult leg length traits; a region on chromosome 11 was significant
across all three adult leg length traits, with the most significant SNP explaining 2.84%, 2.35% and 2.72%
of the genetic variance in adult foreleg, hindleg and metacarpal respectively (Figure 2B, Supplementary
Table 2, Supplementary Figure 1H and 1J). For adult foreleg, a SNP on chromosome 7 and two on
chromosome 9 were also associated, with the most significant SNPs in each region explaining 1.17% and
2.99% of the genetic variance respectively for this trait (Supplementary Table 2, Supplementary Figure
1H).

In lambs, there were no associations with jaw length found (Supplementary Figure 1F). In adults, a
region on chromosome 20 was associated with jaw length variation, with 2.69% of the genetic variance
for this trait being explained by SNP s07047.1 (Supplementary Table 2, Supplementary Figure 1K).

Imputed data

Using the SimpleM method (Gao et al. 2008), we calculated the number of effective tests to be 48635,
giving a genome-wide significance threshold of 1.03e-06.

When performing GWAS using the imputed SNP data, we were able to recover significant SNPs in the
same locations for all traits as those we found using the 50K SNP data, with the exception of the
association between s48811.1 on chromosome 7 and adult foreleg length (Supplementary Table 2). Of
the 84 SNP-trait associations that we identified with the 50K SNP data, only 78 were significant using the
imputed SNP data – the remaining 6 SNPs were no longer significant due to the increased multiple
testing burden (which leads to a more stringent significance threshold) between the 50K SNP data and
the imputed SNP data (2.49e^{-06} and 1.03e^{-06} respectively).

We also identified 845 new SNP-trait associations using the imputed SNP data (Supplementary Table 2).
The majority of new associations were in the same regions as the SNPs identified using the 50K SNP
data, but we also found new associations: a SNP on chromosome 7 was associated with birth weight
(Figure 2A, Supplementary Table 2), and two SNPs on chromosome 13 were associated with adult
hindleg length (Figure 2B, Supplementary Table 2).

Manhattan and QQ plots for all traits can be found in Supplementary Figure 1.

Conditional analysis

For any trait that had at least two SNPs on the same region associated with variation in that trait, we
fitted the genotype of the SNP with the lowest p value in each region in the GWAS model and removed
the SNP from the genotype file. For traits that had multiple SNP associations on more than one
chromosome, we fitted the genotypes of the SNP with the lowest p value from each associated
chromosome simultaneously. We performed conditional analysis on all three leg length traits in both

lambs and adults, as well as on adult jaw length (See Supplementary Table 2 for all SNPs that were fitted
for each trait).

Of the traits we performed conditional analysis on, only two traits had further significant SNPs after
fitting the SNPs with the lowest p value; lamb metacarpal length and adult foreleg length. After fitting
the genotypes for s22142.1 on chromosome 16 and s74894.1 on chromosome 19, three SNPs were
associated with lamb metacarpal length, both when using the 50K SNP data and when using the imputed
SNP data (Figure 3A, Supplementary Table 3). All three SNPs were on chromosome 3 – when we
performed GWAS without fitting SNP genotypes, we found an association between this same region on
chromosome 3 and lamb metacarpal length, however only one SNP had previously reached the
significance threshold and thus had not been fitted in the conditional analysis.

After fitting the genotypes for the SNPs with the smallest p value on chromosomes 9, 11, 16 and 19 for
adult foreleg length, two further SNPs were found to be significant when performing conditional analysis
with the 50K SNP data; one SNP on chromosome 7 (s48811.1) and one SNP on chromosome 11
(s72261.1) (Figure 3B, Supplementary Table 3). The SNP on chromosome 7 was found to be associated
with adult hindleg length when using the 50K SNP data, but was not fitted in the conditional analysis due
to being the only SNP on this chromosome to pass the significance threshold. The SNP on chromosome
11 is located 1.2Mb upstream of the SNP from chromosome 11 that was fitted during the conditional
analysis. When performing the conditional analysis with the imputed SNP data, three new SNPs were
found to be associated with adult foreleg length, all in the same region of chromosome 7 previously
identified when we performed GWAS (Figure 3B, Supplementary Table 3.)

Two-step GWAS

To compare our results using the 50K SNP data to previous GWAS of adult traits in the Soays (Yang et al.
2010), we also performed a two-step GWAS. For our two-step analysis, we used the significance
threshold previously calculated in (Bérénos et al. 2015) ($1.35e^{-6}$), higher than the significance threshold
we calculated using the SimpleM method for the same data.

Across all 5 traits, we recovered the SNP-trait associations identified by (Bérénos et al. 2015). However,
we were unable to recover any of the novel SNP-trait associations we had found when performing our
single-step GWAS on the 50K SNP data, with the exception of the association between chromosome 16
and adult foreleg (however Bérénos et al. noted that SNPs in this region approached significance in their
analysis). Our QQ plots using the two-step method also matched the QQ plots of (Bérénos et al. 2015).
In both, the observed p values were higher than the expected p values, causing the majority of points in
the plots to fall below the x=y line (Supplementary Figure 3).

DISCUSSION

Heritability

Our results corroborate previous findings that all five body size traits we studied in Soay sheep are
influenced by genetic variation in the population (Bérénos et al. 2014), that *post mortem* measures
(metacarpal length and jaw length) have higher heritability estimates than live measures (weight,
foreleg length and hindleg length), that leg measures have higher heritability than weight (Beraldi et al.
2007; Bérénos et al. 2014; Wilson et al. 2006), and that heritability estimates increase with age (Bérénos
et al. 2014; Wilson et al. 2006).

The heritability estimates for the 50K data were very similar to those estimated using a GRM based on
the 50K data in a smaller sample of the same population of sheep by Bérénos et al. (2014) (Bérénos et
al. 2014), with estimates for the same trait falling within one standard error of each other. The biggest
difference was in adult metacarpal length with a heritability difference of 0.049 (estimates were 0.643
(0.049) and 0.594 (0.047) for our and Bérénos et al.'s results respectively). Given that we used the same
models as Bérénos et al., it is likely that the small differences between heritability estimates for each
trait may be due to our increased sample sizes.

Comparing the heritabilities estimated using the imputed SNP data against the estimates using the 50K
SNP data, we found little difference between the two SNP densities in any traits in any age class. The
additional genotypes at the imputed SNPs do not give any additional information on additive genetic
variation for these traits. This result is not surprising given the previous rarefaction analysis showing that
the heritability of these body size traits in adults asymptoted when about half the 50K SNP data was
used (Bérénos et al. 2014). There is high LD between nearby SNPs in the Soay sheep genome, which
suggests that most, if not all, of the causal variants tagged by the imputed SNP data may have already
been tagged by the 50K SNPs. The high LD was reflected when calculating GWAS significance thresholds
– whilst the number of SNPs between the 50K SNP data and the imputed SNP data increased by a factor
of ten, the number of effective tests only doubled (39K SNPs, 20082 effective tests and 401K SNPs and
48635 effective tests respectively).

For some of the traits we have analysed there is still a difference in heritability estimated using SNP data
versus heritability estimated using pedigree – for example, the highest SNP-based heritability estimate
for lamb metacarpal length (the estimate using the imputed SNP data) gave an estimate 59% of Bérénos
et al.'s pedigree-based estimate (Bérénos et al. 2014). Given that our SNP-based heritability estimates
were similar when using the 50K SNP data as when using the imputed SNP data, and the results of
Bérénos et al.'s rarefaction analysis (Bérénos et al. 2014), we believe it is unlikely that increasing the
density of genotyped SNPs that are common in the population will increase heritability estimates of
these traits. It is possible instead that the difference in heritability estimates obtained from pedigree
and genomic data is due to rare familial variants that do not segregate widely in the population, as well
as due to dominance and epistasis.

GWAS

Body size traits have been the focus of many kinds of analyses in Soay sheep (Ashraf et al. 2021; Bérénos
et al. 2015; Bérénos et al. 2014; Ozgul et al. 2009; Pemberton et al. 2017; Regan et al. 2017), and several

SNPs have already been identified as being associated with variation in these traits. A 2014 study aiming
to find SNP-trait associations for these body size traits in adults identified QTL for leg length measures
on chromosomes 16 and 19 (s23172.1 and s74894.1 respectively) (Bérénos et al. 2015). A more recent
study comparing genomic prediction methods in Soays using the 50K SNP data identified s48811.1 on
chromosome 7 and s50107.1 on chromosome 9 as having a probability higher than 0.9 of having a non-
zero effect on adult foreleg length (Ashraf et al. 2021). We were able to identify all four of these
associations in our GWAS, alongside associations that have not previously been identified in this
population. Use of the imputed SNP data allowed us to discover two more associations with loci that
were not genotyped in the 50K SNP data, suggesting that future identification of polymorphisms
influencing trait variation in the Soay sheep may benefit from using the imputed data.

Performing a two-step analysis confirmed that the novel SNP-trait associations we were able to identify
using the 50K SNP data were due to being able to fit the fixed and random effects for each trait whilst
performing GWAS all in a single step, rather than the increased population sample. Given the increase in
SNP-trait associations when using the single-step methodology, we suggest that a two-step GWAS is
redundant with the availability of software like DISSECT which is able to fit fixed and random effects
whilst performing GWAS. As we have shown, although DISSECT does not currently have the option to
automatically run a repeated measures GWAS, it is possible to modify input files to allow for repeated
measures.

The imputed SNP data revealed SNP-trait associations in two regions of the genome that were not
discovered using the 50K SNP data; one between a region on chromosome 7 and birth weight, and one
between a region on chromosome 13 and adult hindleg length. (Supplementary Table 2). When
examining the Manhattan plot for the 50K data for each trait (Figure 2A and 2B, Supplementary Table 2)
it is clear that there was a small cluster of SNPs just under the significance threshold in the 50K analyses.
The additional (imputed) SNPs may have matched the allele frequency of the underlying causal variants
more accurately, resulting in a smaller association p value.

We performed conditional analysis on all three leg length traits in both lambs and adults, as well as on
adult jaw length. For each trait, we simultaneously fitted the genotype for the SNP with the lowest p
value for any chromosome that had at least two SNPs found to be associated with the trait (see
Supplementary Table 2 for a list of SNPs fitted for each trait). We found that, with the exception of
s72261.1 on chromosome 11 associated with adult foreleg length, SNPs that were significant in the
GWAS analysis were no longer significant in the conditional analysis when a significant SNP on the same
chromosome was fitted (Figure 3, Supplementary Table 3). S72261.1 is located more than 1Mb away
from the SNP on chromosome 11 that was fitted during the conditional analysis of adult foreleg, and we
feel confident in stating that the effect that this SNP has on the trait is independent of the effects of the
SNP fitted as a covariate. We suggest that any future work looking to pinpoint the exact location of the
genetic variants affecting body size traits in Soay sheep primarily focus on the regions around the SNPs
listed in Supplementary Table 2, but keep in mind the results of our conditional analyses when
examining chromosome 11 and adult foreleg due to the potential for multiple causal variants on this
chromosome.

Across all traits for all age classes, the QQ plots showed deviation from the expected distribution of test
statistics under the null hypothesis (x=y line) for a wide range of test statistics, including low values,
indicative of underlying population structure not accounted for by the GRMs. Preliminary analysis

suggests that there is a low level of population stratification associated with sire ID, with the first 20
principal components of the GRM accounting for just under 10% of the observed genetic variation.
Preliminary work using LMM-OPS (Conomos et al. 2018), a method that splits genetic sample structure
into familial relatedness (alleles inherited that are identical by descent from recent common ancestors)
and population structure (alleles shared due to distant common ancestors) appears to reduce the
genomic inflation for the lower range of test statistics, while increasing the significance of more robust
associations. However further investigation of the application of that methodology to our population is
still required.

In order to have sufficient power to detect associations between markers and a trait of interest, GWAS
primarily requires two factors: i) a very high density of genotyped SNPs, and ii) a large number of
individuals that have been genotyped and phenotyped (Santure and Garant 2018). For intensively
studied organisms, both are achievable; such populations tend to have more individuals accessible to
collect data from, high density genotyping can be done at a lower cost due to higher demand, and, as in
humans, data from different populations can be combined to create larger sample sizes. GWA studies of
humans are the most obvious example of this; studies often have study populations made up of
hundreds of thousands of individuals and human SNP chips commonly genotype hundreds of thousands
of variants (for example, see (Ishigaki et al. 2020; Wood et al. 2014; Wu et al. 2021)). In comparison, wild
study population samples are much smaller – often struggling to reach one thousand individuals – and
the number of SNPs genotyped is much lower (for example, see (Malenfant et al. 2018; Perrier et al.
2018; Silva et al. 2017)). Analyses of wild populations therefore generally lack the power of more
intensively studied study organisms. Here, we have increased power by increasing the number of
genotyped markers via imputation. Despite high LD in the Soay sheep population, use of imputed data
has allowed us to identify two new SNP-trait associations, including an association with birth weight,
which had yet to be associated with any QTL in the Soay population. We have therefore shown that for a
given sample size, more information can be obtained by increasing the density of markers for those
individuals have been phenotyped. We suggest that, where possible, analyses of wild populations
impute SNP data in order to increase power and obtain results that may otherwise remain
undiscovered.

Although we have discovered new SNP-trait associations, it is likely that there are still causative variants
that remain undetected. GWAS lacks power to detect rare causative variants and variants with very
small effect sizes (Yang et al. 2010). Also, GWAS power drops when the same amount of phenotypic
variation is a consequence of multiple variants in the same region as opposed to a single variant
(Nagamine et al. 2012). Regional mapping methods have been developed that partition trait variance
into regions by simultaneously fitting a whole genome and a regional GRM, with the regions either being
defined as fixed SNP windows (Nagamine et al. 2012) or haplotype blocks (Shirali et al. 2018). Such
methodologies have the potential to identify regions of the genome that contain variants associated
with a trait that are unable to be identified by GWAS either due to being rare, or individually having
small effects on trait variation. Genomic prediction, which simultaneously estimates all marker effects
drawn from multiple distributions, can also be used to study the genetic architecture of traits by
estimating the posterior inclusion probability of a SNP having a non-zero effect on a trait. Genomic
prediction has already been used on adult body size traits in Soays, and has identified several of the
SNPs we identified through our GWAS approach (Ashraf et al. 2021). Ultimately, we believe that it is

important to use a variety of methodologies when studying the genetic architecture of complex traits, as

different analyses have different strengths and may be able to identify different QTL.

References

Ashraf B, Hunter DC, Bérénos C, Ellis PA, Johnston SE, Pilkington JG, Pemberton JM, Slate J. 2021.
495   Genomic prediction in the wild: A case study in soay sheep. Molecular Ecology. 1– 15.
496   Barton NH, Keightley PD. 2002. Understanding quantitative genetic variation. Nature reviews Genetics.
497   3(1):11-21.
498   Beraldi D, McRae AF, Gratten J, Slate J, Visscher PM, Pemberton JM. 2007. Mapping quantitative trait
499   loci underlying fitness-related traits in a free-living sheep population. 61(6):1403-1416.
500   Bérénos C, Ellis PA, Pilkington JG, Lee SH, Gratten J, Pemberton JM. 2015. Heterogeneity of genetic
501   architecture of body size traits in a free-living population. 24(8):1810-1830.
502   Bérénos C, Ellis PA, Pilkington JG, Pemberton JM. 2014. Estimating quantitative genetic parameters in
503   wild populations: A comparison of pedigree and genomic approaches. 23(14):3434-3451.
504   Burdick JT, Chen W-M, Abecasis GR, Cheung VG. 2006. In silico method for inferring genotypes in
505   pedigrees. Nature genetics. 38(9):1002-1004.
506   Butler DG, Cullis BR, Gilmour AR, G. GB, R. T. 2017. Asreml-r reference manual version 4. Hemel
507   Hempstead, HP1 1ES, UK.: VSN International Ltd.
508   Canela-Xandri O, Law A, Gray A, Woolliams JA, Tenesa A. 2015. A new tool called dissect for analysing
509   large genomic data sets using a big data approach. Nature Communications. 6(1):10162.
510   Charmantier A, Garant D, Kruuk LEB. 2014. Quantitative genetics in the wild. Oxford University Press.
511   Conomos MP, Reiner AP, McPeek MS, Thornton TA. 2018. Genome-wide control of population structure
512   and relatedness in genetic association studies via linear mixed models with orthogonally
513   partitioned structure. bioRxiv.409953.
514   Gao X, Starmer J, Martin ER. 2008. A multiple testing correction method for genetic association studies
515   using correlated single nucleotide polymorphisms. 32(4):361-369.
516   . Applications of linear models in animal breeding. 1984.
517   Hickey JM, Kinghorn BP, Tier B, van der Werf JH, Cleveland MA. 2012. A phasing and imputation method
518   for pedigreed populations that results in a single-stage genomic evaluation. Genetics, selection,
519   evolution : GSE. 44(1):9.
520   Ishigaki K, Akiyama M, Kanai M, Takahashi A, Kawakami E, Sugishita H, Sakaue S, Matoba N, Low S-K,
521   Okada Y et al. 2020. Large-scale genome-wide association study in a japanese population
522   identifies novel susceptibility loci across different diseases. Nature genetics. 52(7):669-679.
523   Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S-y, Freimer NB, Sabatti C, Eskin E. 2010. Variance
524   component model to account for sample structure in genome-wide association studies. Nature
525   genetics. 42(4):348-354.
526   Kruuk LEB. 2004. Estimating genetic parameters in natural populations using the "animal model".
527   Philosophical transactions of the Royal Society of London Series B, Biological sciences.
528   359(1446):873-890.
529   Kruuk LEB, Slate J, Wilson AJ. 2008. New answers for old questions: The evolutionary quantitative
530   genetics of wild animal populations. 39(1):525-548.
531   Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J
532   et al. 2015. Genetic studies of body mass index yield new insights for obesity biology. Nature.
533   518(7538):197-206.
534   Malenfant RM, Davis CS, Richardson ES, Lunn NJ, Coltman DW. 2018. Heritability of body size in the
535   polar bears of western hudson bay. Molecular Ecology Resources. 18(4):854-866.

Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon
LR, Chakravarti A et al. 2009. Finding the missing heritability of complex diseases. Nature.
461(7265):747-753.
Mrode RA. 2014. Linear models for the prediction of animal breeding values.
Nagamine Y, Pong-Wong R, Navarro P, Vitart V, Hayward C, Rudan I, Campbell H, Wilson J, Wild S, Hicks
AA et al. 2012. Localising loci underlying complex trait variation using regional genomic
relationship mapping. PloS one. 7(10):e46501.
Ozgul A, Tuljapurkar S, Benton TG, Pemberton JM, Clutton-Brock TH, Coulson T. 2009. The dynamics of
phenotypic change and the shrinking sheep of st. Kilda. Science (New York, NY). 325(5939):464-
467.
Pemberton JM. 2008. Wild pedigrees: The way forward. Proceedings Biological sciences. 275(1635):613-
621.
Pemberton JM, Ellis PE, Pilkington JG, Bérénos C. 2017. Inbreeding depression by environment
interactions in a free-living mammal population. Heredity (Edinb). 118(1):64-77.
Perrier C, Delahaie B, Charmantier A. 2018. Heritability estimates from genomewide relatedness
matrices in wild populations: Application to a passerine, using a small sample size. Molecular
Ecology Resources. 18(4):838-853.
Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012. Double digest radseq: An inexpensive
method for de novo snp discovery and genotyping in model and non-model species. PloS one.
7(5):e37135.
Regan CE, Pilkington JG, Bérénos C, Pemberton JM, Smiseth PT, Wilson AJ. 2017. Accounting for female
space sharing in st. Kilda soay sheep (ovis aries) results in little change in heritability estimates.
Journal of Evolutionary Biology. 30(1):96-111.
Santure AW, Garant D. 2018. Wild gwas—association mapping in natural populations. Molecular Ecology
Resources. 18(4):729-738.
Shirali M, Knott SA, Pong-Wong R, Navarro P, Haley CS. 2018. Haplotype heritability mapping method
uncovers missing heritability of complex traits. Scientific Reports. 8(1):4982.
Silva CNS, McFarlane SE, Hagen IJ, Rönnegård L, Billing AM, Kvalnes T, Kemppainen P, Rønning B, Ringsby
TH, Sæther BE et al. 2017. Insights into the genetic architecture of morphological traits in two
passerine bird species. Heredity (Edinb). 119(3):197-205.
Soay sheep: Dynamics and selection in an island population. 2003. Clutton-Brock TH, Pemberton JM,
editors. Cambridge: Cambridge University Press.
Stoffel MA, Johnston SE, Pilkington JG, Pemberton JM. 2021. Genetic architecture and lifetime dynamics
of inbreeding depression in a wild mammal. Nature communications. 12(1):2972.
Van Vleck LD. 1970. Misidentification in estimating the paternal sib correlation. Journal of Dairy Science.
53(10):1469-1474.
Visscher PM, Medland SE, Ferreira MAR, Morley KI, Zhu G, Cornes BK, Montgomery GW, Martin NG.
2006. Assumption-free estimation of heritability from genome-wide identity-by-descent sharing
between full siblings. PLOS Genetics. 2(3):e41.
Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB. 2006. Quantitative
genetics of growth and cryptic evolution of body size in an island population. Evolutionary
Ecology. 21(3):337.
Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, Kruuk LE, Nussey DH. 2010. An
ecologist's guide to the animal model. The Journal of animal ecology. 79(1):13-26.
Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, Chu AY, Estrada K, Luan Ja, Kutalik Z et al.
2014. Defining the role of common variation in the genomic and biological architecture of adult
human height. Nature genetics. 46(11):1173-1186.

583 Wu Y, Murray GK, Byrne EM, Sidorenko J, Visscher PM, Wray NR. 2021. Gwas of peptic ulcer disease
584   implicates helicobacter pylori infection, other gastrointestinal disorders and depression. Nature
585   communications. 12(1):1146-1146.
586 Xia C, Amador C, Huffman J, Trochet H, Campbell A, Porteous D, Hastie ND, Hayward C, Vitart V, Navarro
587   P et al. 2016. Pedigree- and snp-associated genetics and recent environment are the major
588   contributors to anthropometric and cardiometabolic trait variation. PLoS Genet.
589   12(2):e1005804.
590 Xia C, Canela-Xandri O, Rawlik K, Tenesa A. 2021. Evidence of horizontal indirect genetic effects in
591   humans. Nature human behaviour. 5(3):399-406.
592 Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AA, Lee SH, Robinson MR, Perry JR, Nolte IM, van Vliet-
593   Ostaptchouk JV et al. 2015. Genetic variance estimation with imputed variants finds negligible
594   missing heritability for human height and body mass index. Nature genetics. 47(10):1114-1120.
595 Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath AC, Martin NG,
596   Montgomery GW et al. 2010. Common snps explain a large proportion of the heritability for
597   human height. Nature genetics. 42(7):565-569.
598 Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. 2014. Advantages and pitfalls in the application
599   of mixed-model association methods. Nature genetics. 46(2):100-106.
600 Zaitlen N, Kraft P, Patterson N, Pasaniuc B, Bhatia G, Pollack S, Price AL. 2013. Using extended genealogy
601   to estimate components of heritability for 23 quantitative and dichotomous traits. PLOS
602   Genetics. 9(5):e1003520.

Age	Trait	No. individuals	No. records	Fixed effects	Random effects
Neonate	Birth weight	2678	2678	Sex	Year of birth
				Litter size	Mother ID
				Population size year before birth	
				Age of mother (quadratic)	
				Ordinal date of birth	
				Age (days)	
Lamb	Weight	2228	2228	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Foreleg	2284	2284	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Hindleg	2349	2349	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Metacarpal	2059	2059	Sex	Year of birth
				Litter size	Mother ID
				Age at death (months)	
	Jaw	2113	2113	Sex	Year of birth
				Litter size	Mother ID
				Age at death (months)	
Adult	Weight	1152	3553	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Foreleg	1121	3331	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Hindleg	1135	3444	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Metacarpal	945	945	Sex	Year of birth
				Age at death (years)	
	Jaw	991	991	Sex	Year of birth
				Age at death (years)	

Table 1 Number of individuals and records, fixed and random effects fitted in each trait x age class
model in addition to the GRM. The same individuals and records were used for both heritability
estimates and for GWAS.

Figure 1 Estimates of VA/VP for body size traits in neonates, lambs, and adult Soay sheep when using a
GRM calculated from the 50K SNP data (blue) compared with using a GRM calculated from the imputed
SNP data (yellow). Error bars represent standard error estimates.

Figure 2 Manhattan plots for A) birth weight GWAS using 50K SNP data (left) and imputed SNP data
(right); and B) adult hindleg GWAS using 50K SNP data (left) and imputed SNP data (right). The red line
represents the significance threshold (2.49e-06 for the 50K SNP data and 1.03e-06 for the imputed SNP
data) – any SNPs above this threshold are considered to be significantly associated with variation in
their respective traits.

Figure 3 Miami plots for A) lamb metacarpal GWAS using 50K SNP data (top left) and imputed SNP data
(top right), lamb metacarpal conditional analysis using 50K SNP data (bottom left) and imputed SNP data
(bottom right); and B) adult foreleg GWAS using 50K SNP data (top left) and imputed SNP data (top
right), lamb metacarpal conditional analysis using 50K SNP data (bottom left) and imputed SNP data
(bottom right). The red line represents the significance threshold (2.49e-06 for the 50K SNP data and
1.03e-06 for the imputed SNP data) – any SNPs above this threshold are considered to be significantly
associated with variation in their respective traits.

Data Availability Statement

All data and scripts can be found at
https://github.com/CaelinnJames/Impact_of_SNPDensity_on_Soay_Sheep

Acknowledgments

We thank the National Trust for Scotland and Scottish Natural Heritage for permission to work on St
Kilda and QinetiQ and Eurest for logistics and other support on the island. We also thank all those who
have been involved in the long-term project, including those who helped with field work on the island.
We thank the Wellcome Trust Clinical Research Facility Genetics Core in Edinburgh for SNP genotyping.

Funding

This work was supported by a NERC Doctoral Training Partnership grant (NE/S007407/1). The long-term
field project on St Kilda has been largely funded by the UK Natural Environment Research Council. The
SNP genotyping was funded by a European Research Council Advanced Grant.

Lamb Metacarpal Conditional GWAS Analysis

A

Original GWAS, 50K SNP data

Original GWAS, imputed SNP data

Conditional Analysis, 50K SNP data

Conditional Analysis, imputed SNP data

Adult Foreleg Conditional GWAS Analysis

B

Original GWAS, 50K SNP data

Original GWAS, imputed SNP data

Conditional Analysis, 50K SNP data

Conditional Analysis, imputed SNP data