The impact of SNP density on quantitative genetic analyses of body size traits in a wild population of Soay sheep

Caelinn James[1*], Josephine M. Pemberton[1], Pau Navarro[2], Sara Knott[1]

[1] Institute of Ecology and Evolution, School of Biological Sciences, The University of Edinburgh, Edinburgh

[2] MRC Human Genetics Unit, Institute of Genetics and Cancer, The University of Edinburgh, Edinburgh

[*] Corresponding author

Corresponding author mailing address: Caelinn James, Ashworth Laboratories, Kings Buildings, Charlotte Auerbach Road, Edinburgh EH9 3FL

Corresponding author email address: c.james-5@sms.ed.ac.uk

Key words: GWAS, heritability, wild population, SNP density, animal model

Abstract

Understanding the genetic architecture underpinning quantitative traits in wild populations is pivotal to understanding the processes behind trait evolution. The 'animal model' is a popular method for estimating quantitative genetic parameters such as heritability and genetic correlation and involves fitting an estimate of relatedness between individuals in the study population. Genotypes at genome-wide markers can be used to estimate relatedness; however, relatedness estimates vary with marker density, potentially affecting results. Increasing density of markers is also expected to increase the power to detect quantitative trait loci (QTL). In order to understand how the density of genetic markers affects the results of quantitative genetic analyses, we estimated heritability and performed genome-wide association studies (GWAS) on five body size traits in an unmanaged population of Soay sheep using two different SNP densities: a dataset of 37,037 genotyped SNPs, and an imputed dataset of 417,373 SNPs. Heritability estimates did not differ between the two SNP densities, but the high-density imputed SNP dataset revealed four new SNP-trait associations that were not found with the lower density dataset, as well as confirming all previously-found QTL. We also demonstrated that fitting fixed and random effects in the same step as performing GWAS is a more powerful approach than pre-correcting for covariates in a separate model.

Introduction

Investigating the genetic architecture behind heritable traits is key to understanding the biological diversity of wild populations. If we know the number of loci influencing a trait and their effect size, we can better understand the evolutionary processes that underpin traits, improve inferences about trait evolution, and understand micro-evolutionary dynamics that occur due to environmental change (Barton and Keightley 2002). Most quantitative genetic research in animals is carried out in artificial populations; either domestic, agricultural or laboratory. Such populations experience controlled environmental conditions which make it easier to account for environmental factors when studying the effect of genetic variants on phenotypic variation. However, given that environmental factors can influence the phenotype of a quantitative trait (Charmantier et al. 2014), and that the presence of genotype-by-environment interactions can cause additive genetic variance to differ between environmental conditions, studies on artificial populations arguably cannot be fully extrapolated to wild populations (Kruuk et al. 2008). Therefore, it is important to also study quantitative traits in wild populations in their natural habitats. There is a wealth of quantitative genetics research in human populations (for examples, see Manolio et al. 2009; Kang et al. 2010; Yang et al. 2010; Zaitlen et al. 2013; Locke et al. 2015; Xia et al. 2016; Xia et al. 2021), but humans also arguably experience a more buffered environment than wild populations and inferences in wild populations are lacking in comparison.

When using molecular markers to inform quantitative genetic analyses, the results are dependent on the genetic polymorphisms used in the analysis: low numbers of polymorphisms can result in downwardly biased heritability estimates and regions containing causal variants may not appear as significant if there are no genotyped polymorphisms in linkage disequilibrium (LD) with the causal variant. Increasing the density of genotyped polymorphisms means they are more likely to be in LD with causal variants for the trait of interest, either by being physically closer to the causal variants or by matching the allele frequency of the causal variants more accurately. However, increasing the number of genotyped markers means larger, denser genotyping arrays with costs increasing with density. For commonly studied species, high-density arrays are more affordable due to high demand, but for more niche species, including wild populations, large genotyping arrays are often unaffordable. Genotyping-by-sequencing, e.g. ddRAD (Peterson et al. 2012) is a potentially useful alternative for upscaling SNP density, though the combination of bioinformatics and samples sizes required in quantitative genetic research means that this approach is not yet in widespread use.

As an alternative to expensive high-density genotyping, genotype imputation can be used to increase the number of variants analysed (Burdick et al. 2006). Imputation involves predicting genotypes at untyped SNPs in a 'target' population using a subset of the study population – or more generally a "reference" population – genotyped at a higher density, either through a high density SNP array or by genotyping-by sequencing. The genotypes at these untyped SNPs for individuals in the target population are inferred using their genotypes at typed markers and taking advantage of existing linkage disequilibrium (LD) between SNPs. Pedigree information can also be used to increase the accuracy of the imputation by identifying haplotype blocks that are identical by descent (Burdick et al. 2006).

The Soay sheep *(Ovis aries)* of St Kilda are a primitive, unmanaged breed of sheep that have been the
focus of a longitudinal, individual-based study since 1985 (Clutton-Brock and Pemberton 2003). As part
of the study, morphometric, life history and environmental data are collected, DNA samples are
collected, and a pedigree has been constructed using observation and genetic parentage inference.
7630 sheep have been genotyped on the Ovine SNP50 Illumina Beadchip, on which 37,037 SNPs are
autosomal and polymorphic in this population. In addition, 188 individuals have been genotyped on the
Ovine Infinum HD Beadchip, which contains 419,281 autosomal SNPs that are polymorphic in the
population – this has allowed for imputation of the remaining sheep to this higher density (Stoffel et al.
2021).

In this study we performed a direct comparison of heritability estimates and GWAS associations
between the lower density SNP data and the imputed high density SNP data in the Soay population,
focusing on five body size traits in neonates, lambs and adults. We performed GWAS by fitting fixed and
random effects in the same step as testing for SNP-trait associations, which has the advantage of
correctly propagating error throughout the analysis, reducing the chance of false positive results and
increasing power by disentangling potential correlations. We also carried out a two-step GWAS
approach previously used on a smaller sample size of the Soays (Bérénos et al. 2015) to investigate
whether any SNP-trait associations identified using our approach were due to the increased sample size
or due to the different methodology (single-step vs. two-step GWAS).

Our aims were as follows:

1) To determine whether the increased density of SNPs changes the heritability estimates of the
traits.
93

94   2) To determine whether the imputed SNP data enables the identification of new SNP-trait
95   associations via GWAS.
96

97   3) To compare a single-step GWAS methodology with the two-step approach previously used on
98   the study population.
99

Whilst we have used the Soay sheep as our study population, we believe that our objectives are also
relevant for other wild populations. Keeping costs down is important for all research groups, and we aim
to show that imputation is a way to do so whilst improving the power results of quantitative genetic
analyses. We also intend to highlight the benefit of fitting fixed and random effects whilst performing
GWAS instead of pre-correcting.

METHODS

Phenotypic data

The sheep are ear-tagged when they are first captured which allows for reidentification for life. We
focused on five body size traits in three age groups: neonates, lambs, and adults. Of the five traits, three
(weight, foreleg length and hindleg length) are live measures, recorded in April for neonates and in
August for lambs and adults. The remaining two traits (metacarpal length and jaw length) are *post*
mortem measures taken from skeletal material. Both birth and August weight are measured to the
nearest 0.1kg, whilst the remaining traits are all measured to the nearest mm. A detailed description of
trait measurements can be found in Beraldi et al. (2007).

We defined neonates as individuals who were caught and weighed between two and ten days after birth
– birth weight was the only trait recorded for this age group. Lambs were classed as individuals who had
phenotypic data recorded in the August of their birth year for the live traits, and as individuals who died
before 14 months of age for the *post mortem* measures. Individuals were classed as adults if they had
August phenotypic data recorded at least two years after birth, or if they died after 26 months of age for
post mortem measures. We chose not to analyse yearling data due to the small sample sizes in
comparison to the other age classes, which is due to high first winter mortality.

Genetic data

Most of the sheep in our study population have been genotyped using the Ovine SNP50 Illumina
BeadChip, which targets 54,241 SNPs across the sheep genome. After removing SNPs which failed
quality control standards (minor allele frequency (MAF) > 0.001, call rate > 0.99, deviation from Hardy-
Weinberg Equilibrium P > 1e-05) and individuals with a call rate < 0.95, 39,368 polymorphic variants
remained for 7630 individuals (3643 female, 3987 male). See Bérénos et al. (2014) for information on
genetic sampling protocol and marker characteristics).

Of these 7630 individuals, 188 have also been genotyped using the Ovine Infinium HD SNP BeadChip
which targets 606,066 SNPs. This has allowed for the low density genotypes to be imputed to the higher
density using AlphaImpute, which combines shared haplotype and pedigree information for phasing and
genotype imputation (Hickey et al. 2012) (see Stoffel et al. (2021) for information on imputation and
quality control). We used imputed genotype "hard" calls (rather than genotype probabilities) in
downstream analyses. After filtering SNPs that failed quality control standards, 419,281 autosomal SNPs
remained for 7621 individuals (3639 females, 3982 males).

Both the 50K SNP data and the imputed SNP data are mapped to the OAR_v3.1 genome assembly.

Narrow sense heritability estimation

We used animal models to partition the phenotypic variance for each trait in each age class into genetic
and non-genetic variance components. Fixed and random effects were fitted for all models, with the
effects differing between traits and age classes (Table 1). We implemented these analyses in DISSECT
(Canela-Xandri et al. 2015) using the following model:

$$y = X\beta + \sum_{r} Z_r u_r + Wg + \varepsilon$$

where y is the vector of phenotypic values; X is a design matrix linking individual records with the vector
of fixed effects β, Z_r is an incidence matrix that relates the random effect r to the individual records; u_r is
the associated vector of non-genetic random effects; g is the vector of additive genetic random effects
with W the incidence matrix; and ε is the vector of residuals. It is assumed that g ~ *MVN*(0, Mσ_g^2), where
σ_g^2 is the additive genetic variance and M is the genomic relationship matrix (GRM). For each trait in
each age class, we ran this model twice: first with M being a GRM calculated from the 50K genotype
data, and second with M being a GRM calculated from the imputed SNP genotypes. The GRMs were
computed using DISSECT (Canela-Xandri et al. 2015) using VanRaden's Method 2 GRM calculation, for
which the genetic relationship between individuals i and j is computed as:

$$A_{ij} = \frac{1}{N} \sum_{k=1}^{N} \frac{(s_{ik} - 2p_k)(s_{jk} - 2p_k)}{2p_k(1 - p_k)}$$

where s_{ik} is the number of copies of the reference allele for SNP k of the individual i, p_k is the frequency
of the reference allele for the SNP k, and N is the number of SNPs (VanRaden 2008).

The narrow sense heritability was estimated by dividing the additive genetic variance (the variance
explained by the GRM) by the total estimated phenotypic variance (the sum of the variance explained by
the GRM and other fitted random effects after fitting fixed effects).

In adults, there are multiple records for August weight, foreleg length and hindleg length for the same
individual due to individuals being caught across multiple years. For these traits we used a repeatability
model in order that uncertainty was correctly propagated through all estimations (Mrode 2014). To
implement a repeatability model in DISSECT, we edited the input files so that each measurement had its
own row in the genotype and covariate files. Individual ID was replaced with a unique capture reference
number, and individual permanent environment was fitted as a random effect (see Supplementary
Methods for a more detailed explanation).

Sample sizes and total number of phenotypic measurements for all traits are shown in Table 1, with
effects fitted in all models.

Genome wide association analysis

Principal component analysis (PCA) using the GRM was performed prior to the genome-wide association
analyses (GWAS) using DISSECT (Canela-Xandri et al. 2015) in order to examine the underlying
population structure.

GWAS was also conducted using DISSECT (Canela-Xandri et al. 2015) using the following model:

$$y = \text{X}\beta + \text{S}\alpha + \sum_r \text{Z}_r \text{u}_r + \text{W}g_{LOCO} + \varepsilon$$

where y, X, β, Z_r, u_r, W and ε are the same as in the model for heritability estimation, s is the effect of
the focal SNP, S is the design matrix linking individual records to the number of effect alleles for α, the
estimated SNP effect (coded as 0, 1 or 2), and gLOCO is the vector of additive genetic random effects
excluding the chromosome on which the focal SNP resides.

We fitted the same fixed and random effects for each trait and age class as for the heritability
estimation (Table 1). To account for population structure, when testing SNPs on a given chromosome for
association with the phenotype, a GRM calculated from the remaining autosomes (referred to as Leave
One Chromosome Out GRM (gLOCO) (Yang et al. 2014)) was fitted. Input files for repeated-measure
traits were reformatted as above. Our significance threshold was corrected for multiple testing using the
SimpleM method (Gao et al. 2008), which accounts for linkage disequilibrium between markers in order
to calculate the effective number of independent tests.

We estimated the variance explained by SNPs that passed the significance threshold using the equation

$$V(SNP) = 2pq\alpha^2$$

where p and q are the major and minor allele frequencies of the SNP, and α is the estimated SNP effect.
We then calculated the proportion of additive genetic variance explained by each SNP by dividing by the
total additive genetic variance estimated for that trait.

For any trait for which several SNPs in the same region were associated with variation in the trait and
thus had strong support for at least one QTL in the region, we carried out conditional analysis to
understand if the region could potentially harbour several independent QTL, or if further QTL could be
uncovered elsewhere in the genome. To that aim, the genotypes of the SNP with the smallest
association p value from each associated region (hereafter called the "top SNP") were added to the
GWAS model as a fixed covariate and removed from the GRMs and genotype data. The GWAS analysis
was re-run accounting for those associations to try and reveal novel peaks either in the same regions or
elsewhere in the genome.

Genes in QTL regions

For each trait x SNP association, we investigated the genes within a 0.5Mb window either side of the top
SNP to identify any genes which could be contributing to trait variation. We extracted a list of genes for
each trait using the biomaRt package in R (Durinck et al. 2005; Durinck et al. 2009) from the OAR_v3.1
genome assembly and reviewed each gene against the NCBI Gene (Bethesda (MD): National Library of
Medicine (US) 2004 - 2022) (including information from the Sheep Tissue Atlas (Jiang et al. 2014)),
Animal QTLdb (Hu et al. 2022), and Ensembl (Howe et al. 2020) databases to examine function and
expression annotations. When possible, we also compared with human and mouse orthologues due to
the high level of annotation data available for these two species.

Two-step GWAS analysis

GWAS has previously been performed on the adult traits in a smaller sample of Soay sheep using the
50K SNP data (Bérénos et al. 2015). The authors performed GWAS by first running a mixed model
analysis, fitting fixed and random effects including whole-genome relatedness in the form of a GRM and,
for repeated-measure traits, permanent environment. The residuals were then extracted and used as
the phenotypic values for GWAS. For repeated measure traits, the mean residual value was used for
each individual.

To investigate whether any novel SNP associations identified in this study were due to the increased
sample size or due to the change in methodology, we also performed a two-step GWAS, focusing on
adults only and using the 50K SNP data. Like Bérénos et al. (2015), we performed mixed model analyses
using ASReml-R (Butler et al. 2017) and performed GWAS with the residuals as the trait phenotypes
using DISSECT (Canela-Xandri et al. 2015). We used the Bonferroni correction calculated in Bérénos et al.
(2015) to determine the significance threshold of 1.35e-6 for our two-step GWAS in order to compare
with the previously published analysis.

RESULTS

Heritability estimation

Neonates

In neonates, the heritability of birth weight was 0.051 (S.E. 0.020) both when using the 50K SNPs to
calculate relatedness, and when using the imputed SNPs (Figure 1, Supplementary Table 1). Given that
both estimates are identical to 3 decimal places, there is no difference between the estimates.

Lambs

In lambs, the heritability estimates for the live August measures were lower than those for the *post*
mortem measures (Figure 1, Supplementary Table 1). Across all the traits, heritability estimates were
similar when using the 50K SNP data and the imputed SNP data, with the biggest difference being 0.024
for metacarpal length. For all traits, estimates were within one standard error of each other, indicating
that the small differences in heritability estimates between the two SNP densities were not significant.

Adults

As observed in lambs, heritability estimates for live measures in adults were lower than those of the
post mortem measures. Across all traits, heritability estimates were higher in adults than in lambs.
Estimates obtained using the 50K SNPs and using the imputed SNPs were similar and were within one
standard error of each other (Figure 1, Supplementary Table 1), meaning that the imputed SNPs
provided no additional information to partition the variation into genetic and environmental variance.

Estimates for all variance components are listed in Supplementary Table 1.

GWAS

PCA

The top 20 principal components together explained 10.68% of the variance in the genetic data, with the
top two principal components explaining 0.98% and 0.85% of the variance. We concluded that any
population structure that was likely to affect the GWAS results would be corrected for by fitting a LOCO
GRM in our GWAS model, as this will account for any structure caused by relationships.

50K SNP data

To correct for multiple testing, we calculated the effective number of tests to be 20082 using the
SimpleM method (Gao et al. 2008), giving a genome-wide significance threshold of $2.49e^{-06}$ for the 50K
SNP data.

For weight in neonates (birth weight), and lambs (August weight), no SNPs were found to have an
association p value smaller than this threshold, suggesting that any variants that influence weight
variation are either of small effect or were not tagged by SNPs in the 50K SNP data (Figure 2A,
Supplementary Figure 1B and 1G). For adult August weight, three SNPs had a p value lower than the
genome-wide significance threshold; one SNP on chromosome 6 and two SNPs on chromosome 9.

For all three leg length measures in lambs, we found associations with the same region on chromosome
16. SNP s23172.1 was the SNP with the lowest p value for lamb foreleg and hindleg, explaining 0.52%
and 0.69% of the genetic variance for each trait respectively (Supplementary Table 2, Supplementary
Figure 1C and 1D). For lamb metacarpal, SNP 22142.1 in the same chromosome 16 region had the
lowest p value and explained 0.97% of the genetic variance. There was also a single SNP on chromosome
3 (OAR3_100483326.1) and a cluster of SNPs on chromosome 19 that had p values smaller than the
genome-wide significance threshold and were associated with variation in lamb metacarpal length, with
the SNP with the lowest p value from each region explaining 2.08% and 2.40% of the genetic variance
respectively (Supplementary Table 2, Supplementary Figure 1E).

The two regions on chromosomes 16 and 19 that were associated with lamb metacarpal length variation
were also significantly associated with all three leg length measures in adults, with SNP s22142.1 on
chromosome 16 and SNP s74894.1 on chromosome 19 respectively explaining 0.80% and 2.04% of the
genetic variation in adult foreleg, 0.88% and 1.32% of the genetic variation in adult hindleg, and 0.55%
and 2.02% of the genetic variation in adult metacarpal length. There were other regions of the genome
also associated with variation in the adult leg length traits; a region on chromosome 11 was significant
across all three adult leg length traits, with the most significant SNP explaining 2.35%, 2.25% and 1.13%
of the genetic variance in adult foreleg, hindleg and metacarpal respectively (Figure 2B, Supplementary
Table 2, Supplementary Figure 1H and 1J). For adult foreleg, a SNP on chromosome 7 and two on
chromosome 9 were also associated, with the most significant SNPs in each region explaining 1.31% and
2.99% of the genetic variance respectively for this trait (Supplementary Table 2, Supplementary Figure
1H).

In lambs, there were no associations with jaw length found (Supplementary Figure 1F). In adults, a SNP
on chromosome 20 was associated with jaw length variation, explaining 2.05% of the genetic variance
for this trait (Supplementary Table 2, Supplementary Figure 1K).

In total, we identified 85 SNP-trait associations with 39 unique SNPs.

Imputed data

Using the SimpleM method (Gao et al. 2008), we calculated the number of effective tests to be 48635,
giving a genome-wide significance threshold of $1.03e^{-06}$.

When performing GWAS using the imputed SNP data, we were able to recover significant SNPs in the
same locations for all traits as those we found using the 50K SNP data. Of the 85 SNP-trait associations
that we identified with the 50K SNP data, 81 were significant using the imputed SNP data – the
remaining four SNPs were no longer significant due to the increased multiple testing burden (which
leads to a more stringent significance threshold) between the 50K SNP data and the imputed SNP data
($2.49e^{-06}$ and $1.03e^{-06}$ respectively).

We also identified 795 new SNP-trait associations using the imputed SNP data with 425 unique SNPs
(Supplementary Table 2). The majority of new associations were in the same regions as the SNPs
identified using the 50K SNP data, but we also found new associations: four SNPs on chromosome 1 and
three SNP on chromosome 7 was associated with birth weight (Figure 2A, Supplementary Table 2), one
SNP on chromosome 3 was associated with adult August weight (Figure 2B, Supplementary Table 2), and

one SNP on chromosome 17 was associated with adult metacarpal length (Figure 2C, Supplementary
Table 2).

Manhattan and QQ plots for all traits can be found in Supplementary Figure 1.

Conditional analysis

We performed conditional analysis on all three leg length traits in both lambs and adults, as well as on
birth weight, adult August weight adult jaw length (See Supplementary Table 2 for all SNPs that were
fitted for each trait). For all of these traits we performed the conditional analysis using both the 50K SNP
data and the imputed SNP data, with the exception of birth weight, which did not have any significant
SNP associations using the 50K data.

Six of the nine traits we performed conditional analysis on had significant SNPs after fitting the SNPs
with the lowest p value, however for four of these traits (lamb metacarpal length, adult August weight,
foreleg length and hindleg length), these were SNPs that were also significant in our original GWAS
analysis but were not fitted in the conditional analysis due to being the only SNP that was significantly
associated with the trait in that region (Supplementary Table 3). The remaining two traits (birth weight
and adult jaw length) both had a new association, both of which were on chromosome 2. For birth
weight, nine SNPs had p values lower than the genome-wide significance threshold, all around ~81Mb
(Figure 3A, Supplementary Table 3). For adult jaw length, only one SNP had a lower p value than the
genome-wide significance threshold, at position 137,162,126 (Figure 3B, Supplementary Table 3).

Genes in QTL regions

Given that all of the region-trait associations that were found to be significant with the 50K SNP data
were also significant with the imputed SNP data, we chose to focus on top SNPs in the imputed dataset,
including the two regions that were only significantly associated when performing the conditional
analysis (See Supplementary Table 2 and 3 for the list of SNPs, and Supplementary Table 4 for the list of
genes).

We found a total of 179 genes in the regions around the SNPs associated with our traits. 56 of these
genes were unannotated in the current sheep genome build, and of those that were annotated, three
did not have a listed mouse homologue and a further six had neither a mouse nor a human homologue.
Of the genes that did have annotation and homologue data, we found nine that are associated with
similar traits to our focal traits in humans and mice, suggesting that they may be contributing to the
genetic variation of our traits (Table 2). However, without intimate knowledge of the genes surrounding
the focal SNPs, it is likely that there are other genes that are also contributing. It is also worth noting
that variation in coding regions of genes in proximity to SNPs we identified as being associated with our
traits may not be responsible for variation in the phenotypes, but instead the causal variants may lie in
regulatory sequences that modulate expression of either these or other genes.

We also compared our GWAS results with QTL from Animal QTLdb (Hu et al. 2022). We found that the
region on chromosome 6 that we found to be associated with adult August weight overlaps with a
region previously found to be associated with carcass weight and final body weight in an (Awassi x
Merino) x Merino backcross population (Cavanagh et al. 2010) and is ~0.5Mb upstream of a 2.5Mb

region that has also previously been associated with body weight in a population of Australian Merino
sheep (Al-Mamun et al. 2015). In addition, the region on chromosome 9 we found to be associated with
adult August weight is 1Mb upstream of a region previously found to be associated with yearling live
weight in a population of Chinese Merino sheep.

Chromosome 6 has previously been associated with adult body weight using linkage analysis in a smaller
sample of Soays (Beraldi et al. 2007), however the microsatellite markers flanking the associated region
are not located close to the region we identified.

Two-step GWAS

Across all 5 traits when using the two-step approach, we recovered the SNP-trait associations identified
by Bérénos et al. (2015). However, we were unable to recover any of the novel SNP-trait associations we
found when performing our single-step GWAS on the 50K SNP data, with the exception of the
association between chromosome 16 and adult foreleg (though Bérénos et al. noted that SNPs in this
region approached significance in their analysis). Despite the genome-wide significance threshold used
by Bérénos et al. (2015) being more stringent than the significance threshold we calculated using the
SimpleM method for the 50K SNP data, no additional associations are recovered when using our less
stringent threshold

Our QQ plots using the two-step method also matched the QQ plots of Bérénos et al. (2015). In both,
the observed p values were higher than the expected p values, causing the majority of points in the
plots to fall below the x=y line (Supplementary Figure 3).

DISCUSSION

Heritability

Our results corroborate findings from previous studies in the Soay sheep. For instance, we showed that
all five body size traits we studied in Soay sheep are influenced by genetic variation in the population
(Bérénos et al. 2014), and that *post mortem* measures have higher heritability estimates than live
measures . We also found that leg measures have higher heritability than weight (Wilson et al. 2006;
Beraldi et al. 2007; Bérénos et al. 2014), and that heritability estimates increase with age (Wilson et al.
2006; Bérénos et al. 2014).

The heritability estimates for the 50K data were very similar to those estimated using a GRM based on
the 50K data in a smaller sample of the same population of sheep by Bérénos et al. (2014), with
estimates for the same trait falling within one standard error of each other. The biggest difference was
in adult metacarpal length with a heritability difference of 0.05 (estimates were 0.644 (0.047) and 0.594
(0.047) for our and Bérénos et al.'s results respectively). Given that we used the same models as
Bérénos et al., it is likely that the small differences between heritability estimates for each trait is due to
our increased sample sizes.

Comparing the heritabilities estimated using the imputed SNP data against the estimates using the 50K
SNP data, we found little difference between the two SNP densities in any traits in any age class. The
additional genotypes at the imputed SNPs do not give any additional information on additive genetic
variation for these traits. This result is not surprising given the previous rarefaction analysis showing that
the heritability of these body size traits in adults asymptoted when about half the 50K SNP data was
used (Bérénos et al. 2014). There is high LD between nearby SNPs in the Soay sheep genome, which
suggests that most, if not all, of the causal variants tagged by the imputed SNP data may have already
been tagged by the 50K SNPs. The high LD was reflected when calculating GWAS significance thresholds
– whilst the number of SNPs between the 50K SNP data and the imputed SNP data increased by a factor
of ten, the number of effective tests only doubled (39K SNPs, 20082 effective tests and 401K SNPs and
48635 effective tests respectively).

For some of the traits we have analysed there is still a difference in heritability estimated using SNP data
versus heritability estimated using pedigree – for example, the highest SNP-based heritability estimate
for lamb metacarpal length (the estimate using the imputed SNP data) gave an estimate 59% of Bérénos
et al.'s pedigree-based estimate (Bérénos et al. 2014). Given that our SNP-based heritability estimates
were similar when using the 50K SNP data as when using the imputed SNP data, and the results of
Bérénos et al.'s rarefaction analysis (Bérénos et al. 2014), we believe it is unlikely that increasing the
density of genotyped SNPs that are common in the population will increase heritability estimates of
these traits. It is possible instead that the difference in heritability estimates obtained from pedigree
and genomic data is due to rare familial variants that do not segregate widely in the population, as well
as due to dominance and epistasis.

GWAS

Body size traits have been the focus of many kinds of analyses in Soay sheep (Beraldi et al. 2007; Ozgul
et al. 2009; Bérénos et al. 2014; Bérénos et al. 2015; Pemberton et al. 2017; Regan et al. 2017; Ashraf et

al. 2021), and several SNP associations have already been identified for these traits. A 2015 study aiming
to find SNP-trait associations identified QTL for adult leg length measures on chromosomes 16 and 19
(s23172.1 and s74894.1 respectively) (Bérénos et al. 2015). A more recent study comparing genomic
prediction methods in Soays using the 50K SNP data identified s48811.1 on chromosome 7 and s50107.1
on chromosome 9 as having a probability higher than 0.9 of having a non-zero effect on adult foreleg
length in addition to the previously discovered regions on chromosomes 16 and 19 (Ashraf et al. 2021).
We were able to identify all four of these associations in our GWAS, alongside associations that have not
previously been identified in this population. Use of the imputed SNP data allowed us to discover four
more associations with loci that were not genotyped in the 50K SNP data, indicating that future
identification of polymorphisms influencing trait variation in the Soay sheep will benefit from using the
imputed data.

Performing a two-step analysis confirmed that the novel SNP-trait associations we were able to identify
using the 50K SNP data were due to fitting the fixed and random effects for each trait simultaneously,
performing GWAS in a single step, rather than due to having a larger sample size. Given the increase in
SNP-trait associations when using the single-step methodology, and with the availability of software like
DISSECT which is able to fit fixed and random effects whilst performing GWAS, two-step GWAS should
be avoided. As we have shown, although DISSECT does not currently have the option to automatically
run a repeated measures GWAS, it is possible to modify input files to allow for repeated measures to be
appropriately modelled.

The imputed SNP data revealed SNP-trait associations in four regions of the genome that were not
discovered using the 50K SNP data; a region on chromosome 1 and a region on chromosome 7 and birth
weight, a region on chromosome 3 and adult August weight, and a region on chromosome 17 and adult
metacarpal length. (Supplementary Table 2). When examining the Manhattan plot for the 50K data for
each trait (Figure 2A, 2B and 2C, Supplementary Table 2) it is clear that, with the exception of the region
on chromosome 1 associated with birth weight, there was a small cluster of SNPs just under the
significance threshold in the 50K analyses. The additional (imputed) SNPs may have matched the allele
frequency of the underlying causal variants more accurately, resulting in a smaller association p value.

We performed conditional analysis on all three leg length traits in both lambs and adults, as well as on
birth weight (only using the imputed SNP data), adult August weight and adult jaw length. For each trait,
we simultaneously fitted the genotype for the SNP with the lowest p value for any chromosome that had
at least two SNPs found to be associated with the trait (see Supplementary Table 2 for a list of SNPs
fitted for each trait). We found that all of the SNPs that were significant in the GWAS analysis were no
longer significant in the conditional analysis when a significant SNP on the same chromosome was fitted
(Figure 3, Supplementary Table 3). We suggest that any future work looking to pinpoint the exact
location of the genetic variants affecting body size traits in Soay sheep primarily focus on the regions
around the SNPs listed in Supplementary Table 2.

We identified 179 genes within 0.5Mb of the top SNPs for each trait (Supplementary Table 4), and of
these genes, we found nine that are potential candidate genes for further analyses due to their
association with similar traits in other species. Two of these genes (*CYP26B1* and *EXOC6B*) are associated
with the same trait (lamb metacarpal length) and are located within 0.5Mb of the same top SNP on
chromosome 3. *CYP26B1* is associated with skeletal abnormalities in humans and zebrafish (Laue et al.
2011) and *CYP26B1* knockouts produced reduced limbs in mice (Yashiro et al. 2004), whilst *EXOC6B* is

associated with spondyloepimetaphyseal dysplasia in humans which symptoms include skeletal
malformations affecting the long bones of the limbs and short stature (Campos-Xavier et al. 2018).

PTH1R is located in the region on chromosome 19 associated with lamb metacarpal length and all three
adult leg traits, and has been found to be involved in osteoblast development in mice (Qiu et al. 2015),
and is associated with skeletal disorders such as Eiken's syndrome (Duchatelet et al. 2005), Jansen's
metaphyseal chondrodysplasia and Blomstrand's lethal chondrodysplasia (Schipani and Provot 2003). As
the top SNPs on chromosome 19 for both lamb and adult metacarpal are in a slightly different position
to the top SNP for the live measures, *LTF* is in the region associated with both lamb and adult
metacarpal length on chromosome 19, and has been found to increase bone growth when injected into
piglets (Li et al. 2018), stimulate osteoblast proliferation in humans (Cornish and Naot 2010), and has
high expression levels in human bone marrow (Fagerberg et al. 2014).

BMP6, located in the region on chromosome 20 associated with adult jaw length, has also been found to
be involved in bone development and expression in mice jaw bone (Oralová et al. 2014).

We found three potential candidate genes associated with weight. *FAM13A* in the region on
chromosome 6 associated with adult August weight has been found to modulate body fat distribution
and adipocyte function in humans and mice (Fathzadeh et al. 2020), affects adipose insulin signalling in
mice (Wardhana et al. 2018), and is associated with obesity also in mice (Tang et al. 2019). *PENK*, located
in the region on chromosome 9 associated with adult August weight has been found to cause diminished
food motivation, lower baseline body weight and attenuated weight gain when knocked out in mice
(Mendez et al. 2015). *ONECUT1*, located in the region on chromosome 7 associated with birth weight
has been found to be associated with birth weight in humans (Warrington et al. 2019). Unlike the other
potential candidate gene-trait associations we found, this association was only found when performing
GWAS with the imputed SNP data.

Although we have discovered new SNP-trait associations, it is likely that there are still causative variants
that remain undetected. GWAS lacks power to detect rare causative variants and variants with very
small effect sizes (Yang et al. 2010). Also, GWAS power drops when the same amount of phenotypic
variation is a consequence of multiple variants in the same region as opposed to a single variant
(Nagamine et al. 2012). Regional mapping methods have been developed that partition trait variance
into regions by simultaneously fitting a whole genome and a regional GRM, with the regions either being
defined as fixed SNP windows (Nagamine et al. 2012) or haplotype blocks (Shirali et al. 2018). Such
methodologies have the potential to identify regions of the genome that contain variants associated
with a trait that are unable to be identified by GWAS either due to being rare, or individually having
small effects on trait variation. Genomic prediction, which simultaneously estimates all marker effects
drawn from multiple distributions, can also be used to study the genetic architecture of traits by
estimating the posterior inclusion probability of a SNP having a non-zero effect on a trait. Genomic
prediction has already been used on adult body size traits in Soays, and has identified several of the
SNPs we identified through our GWAS approach (Ashraf et al. 2021). Ultimately, we believe that it is
important to use a variety of methodologies when studying the genetic architecture of complex traits, as
different analyses have different strengths and may be able to identify different QTL.Across all traits for
all age classes, the QQ plots showed deviation from the expected distribution of test statistics under the
null hypothesis (x=y line) for a wide range of test statistics, including low values, which may be either
due to underlying population structure not accounted for by the GRMs or due to trait architecture. The

first 20 genomic principal components accounted for only 10.68% of the variance in the genetic data,
and repeating the GWAS analysis fitting these first 20 genomic principal components in addition to the
GRM did not change the p values of the SNPs nor the QQ plots (data not shown). This shows that the
principal components in this case were not useful in adjusting for population structure in the presence
of the GRM, and that population structure is not likely to cause any issues in our analyses, especially
after fitting a LOCO GRM.

In order to have sufficient power to detect associations between markers and a trait of interest, GWAS
primarily requires two factors: i) a very high density of genotyped SNPs, and ii) a large number of
individuals that have been genotyped and phenotyped (Santure and Garant 2018). For intensively
studied organisms, both are achievable; such populations tend to have more individuals accessible to
collect data from, high density genotyping can be done at a lower cost due to higher demand, and, as in
humans, data from different populations can be combined to create larger sample sizes. GWA studies of
humans are the most obvious example of this; studies often have study populations made up of
hundreds of thousands of individuals and human SNP chips commonly genotype hundreds of thousands
of variants (for example, see Wood et al. 2014; Ishigaki et al. 2020; Wu et al. 2021). In comparison, wild
study population samples are much smaller – often struggling to reach one thousand individuals – and
the number of SNPs genotyped is much lower (for example, see Silva et al. 2017; Malenfant et al. 2018;
Perrier et al. 2018). Analyses of wild populations therefore generally lack the power of more intensively
studied study organisms. Here, we have increased power by increasing the number of genotyped
markers via imputation. Despite high LD in the Soay sheep population, use of imputed data has allowed
us to identify four new SNP-trait associations, including an association with birth weight, which had yet
to be associated with any QTL in the Soay population. We have therefore shown that for a given sample
size, more information can be obtained by increasing the density of markers for those individuals have
been phenotyped. We suggest that, where possible, analyses of wild populations impute SNP data in
order to increase power and obtain results that may otherwise remain undiscovered.

In our population, we were able to identify new SNP-trait associations when performing GWAS with the
higher density imputed dataset that were not found with the lower density non-imputed data – one of
which allowed for the identification of a potential candidate gene for that trait. We have shown that
analyses performed with imputed data, such as heritability estimation and GWAS, can still pick up
additive genetic variation that would have been identified with the non-imputed data. We suggest that,
where possible, analyses of wild populations impute SNP data in order to increase power and obtain
associations that may otherwise remain undiscovered. In addition, we demonstrated the benefits of
fitting fixed and random effects during GWAS instead of pre-correcting and performing GWAS on the
residuals. We therefore recommend that researchers use this "one-step" method when performing
GWAS on wild populations.

References

Al-Mamun HA, Kwan P, Clark SA, Ferdosi MH, Tellam R, Gondro C. 2015. Genome-wide association study
of body weight in australian merino sheep reveals an orthologous region on oar6 to human and
bovine genomic regions affecting height and weight. Genetics, selection, evolution : GSE.
47(1):66.
Arts HH, Bongers EM, Mans DA, van Beersum SE, Oud MM, Bolat E, Spruijt L, Cornelissen EA, Schuurs-
Hoeijmakers JH, de Leeuw N et al. 2011. C14orf179 encoding ift43 is mutated in sensenbrenner
syndrome. Journal of medical genetics. 48(6):390-395.
Ashraf B, Hunter DC, Bérénos C, Ellis PA, Johnston SE, Pilkington JG, Pemberton JM, Slate J. 2021.
Genomic prediction in the wild: A case study in soay sheep. Molecular Ecology. 1– 15.
Barton NH, Keightley PD. 2002. Understanding quantitative genetic variation. Nature reviews Genetics.
3(1):11-21.
Beraldi D, McRae AF, Gratten J, Slate J, Visscher PM, Pemberton JM. 2007. Mapping quantitative trait
loci underlying fitness-related traits in a free-living sheep population. 61(6):1403-1416.
Bérénos C, Ellis PA, Pilkington JG, Lee SH, Gratten J, Pemberton JM. 2015. Heterogeneity of genetic
architecture of body size traits in a free-living population. 24(8):1810-1830.
Bérénos C, Ellis PA, Pilkington JG, Pemberton JM. 2014. Estimating quantitative genetic parameters in
wild populations: A comparison of pedigree and genomic approaches. 23(14):3434-3451.
Ncbi gene. 2004 - 2022. [accessed 11/02/2022]. https://www.ncbi.nlm.nih.gov/gene/.
Burdick JT, Chen W-M, Abecasis GR, Cheung VG. 2006. In silico method for inferring genotypes in
pedigrees. Nature genetics. 38(9):1002-1004.
Campos-Xavier B, Rogers RC, Niel-Bütschi F, Ferreira C, Unger S, Spranger J, Superti-Furga A. 2018.
Confirmation of spondylo-epi-metaphyseal dysplasia with joint laxity, exoc6b type.
176(12):2934-2935.
Canela-Xandri O, Law A, Gray A, Woolliams JA, Tenesa A. 2015. A new tool called dissect for analysing
large genomic data sets using a big data approach. Nature Communications. 6(1):10162.
Cavanagh CR, Jonas E, Hobbs M, Thomson PC, Tammen I, Raadsma HW. 2010. Mapping quantitative trait
loci (qtl) in sheep. Iii. Qtl for carcass composition traits derived from ct scans and aligned with a
meta-assembly for sheep and cattle carcass qtl. Genetics, selection, evolution : GSE. 42(1):36.
Charmantier A, Garant D, Kruuk LEB. 2014. Quantitative genetics in the wild. Oxford University Press.
Clutton-Brock TH, Pemberton JM. 2003. Soay sheep: Dynamics and selection in an island population.
Clutton-Brock TH, Pemberton JM, editors. Cambridge: Cambridge University Press.
Cornish J, Naot D. 2010. Lactoferrin as an effector molecule in the skeleton. BioMetals. 23(3):425-430.
Duchatelet S, Ostergaard E, Cortes D, Lemainque A, Julier C. 2005. Recessive mutations in pthr1 cause
contrasting skeletal dysplasias in eiken and blomstrand syndromes. Human molecular genetics.
14(1):1-5.
Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W. 2005. Biomart and
bioconductor: A powerful link between biological databases and microarray data analysis.
Bioinformatics (Oxford, England). 21(16):3439-3440.
Durinck S, Spellman PT, Birney E, Huber W. 2009. Mapping identifiers for the integration of genomic
datasets with the r/bioconductor package biomart. Nature protocols. 4(8):1184-1191.
Fagerberg L, Hallström BM, Oksvold P, Kampf C, Djureinovic D, Odeberg J, Habuka M, Tahmasebpoor S,
Danielsson A, Edlund K et al. 2014. Analysis of the human tissue-specific expression by genome-
wide integration of transcriptomics and antibody-based proteomics. Molecular & cellular
proteomics : MCP. 13(2):397-406.

Fathzadeh M, Li J, Rao A, Cook N, Chennamsetty I, Seldin M, Zhou X, Sangwung P, Gloudemans MJ, Keller
M et al. 2020. Fam13a affects body fat distribution and adipocyte function. Nature
communications. 11(1):1465.
Gao X, Starmer J, Martin ER. 2008. A multiple testing correction method for genetic association studies
using correlated single nucleotide polymorphisms. 32(4):361-369.
Hickey JM, Kinghorn BP, Tier B, van der Werf JH, Cleveland MA. 2012. A phasing and imputation method
for pedigreed populations that results in a single-stage genomic evaluation. Genetics, selection,
evolution : GSE. 44(1):9.
Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov AG, Bennett R,
Bhai J et al. 2020. Ensembl 2021. Nucleic Acids Research. 49(D1):D884-D891.
Hu Z-L, Park CA, Reecy JM. 2022. Bringing the animal qtldb and corrdb into the future: Meeting new
challenges and providing updated services. Nucleic Acids Research. 50(D1):D956-D961.
Ishigaki K, Akiyama M, Kanai M, Takahashi A, Kawakami E, Sugishita H, Sakaue S, Matoba N, Low S-K,
Okada Y et al. 2020. Large-scale genome-wide association study in a japanese population
identifies novel susceptibility loci across different diseases. Nature genetics. 52(7):669-679.
Jiang Y, Xie M, Chen W, Talbot R, Maddox JF, Faraut T, Wu C, Muzny DM, Li Y, Zhang W et al. 2014. The
sheep genome illuminates biology of the rumen and lipid metabolism. Science (New York, NY).
344(6188):1168-1173.
Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S-y, Freimer NB, Sabatti C, Eskin E. 2010. Variance
component model to account for sample structure in genome-wide association studies. Nature
genetics. 42(4):348-354.
Kruuk LEB, Slate J, Wilson AJ. 2008. New answers for old questions: The evolutionary quantitative
genetics of wild animal populations. 39(1):525-548.
Laue K, Pogoda H-M, Daniel PB, van Haeringen A, Alanay Y, von Ameln S, Rachwalski M, Morgan T, Gray
MJ, Breuning MH et al. 2011. Craniosynostosis and multiple skeletal anomalies in humans and
zebrafish result from a defect in the localized degradation of retinoic acid. Am J Hum Genet.
89(5):595-606.
Li Q, Zhao J, Hu W, Wang J, Yu T, Dai Y, Li N. 2018. Effects of recombinant human lactoferrin on
osteoblast growth and bone status in piglets. Animal Biotechnology. 29(2):90-99.
Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J
et al. 2015. Genetic studies of body mass index yield new insights for obesity biology. Nature.
518(7538):197-206.
Malenfant RM, Davis CS, Richardson ES, Lunn NJ, Coltman DW. 2018. Heritability of body size in the
polar bears of western hudson bay. Molecular Ecology Resources. 18(4):854-866.
Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon
LR, Chakravarti A et al. 2009. Finding the missing heritability of complex diseases. Nature.
461(7265):747-753.
Mendez IA, Ostlund SB, Maidment NT, Murphy NP. 2015. Involvement of endogenous enkephalins and
β-endorphin in feeding and diet-induced obesity. Neuropsychopharmacology : official
publication of the American College of Neuropsychopharmacology. 40(9):2103-2112.
Mrode RA. 2014. Linear models for the prediction of animal breeding values.
Nagamine Y, Pong-Wong R, Navarro P, Vitart V, Hayward C, Rudan I, Campbell H, Wilson J, Wild S, Hicks
AA et al. 2012. Localising loci underlying complex trait variation using regional genomic
relationship mapping. PloS one. 7(10):e46501.
Oralová V, Chlastáková I, Radlanski RJ, Matalová E. 2014. Distribution of bmp6 in the alveolar bone
during mouse mandibular molar eruption. Connective tissue research. 55(5-6):357-366.

Ozgul A, Tuljapurkar S, Benton TG, Pemberton JM, Clutton-Brock TH, Coulson T. 2009. The dynamics of phenotypic change and the shrinking sheep of st. Kilda. Science (New York, NY). 325(5939):464-467.

Pemberton JM, Ellis PE, Pilkington JG, Bérénos C. 2017. Inbreeding depression by environment interactions in a free-living mammal population. Heredity (Edinb). 118(1):64-77.

Perrier C, Delahaie B, Charmantier A. 2018. Heritability estimates from genomewide relatedness matrices in wild populations: Application to a passerine, using a small sample size. Molecular Ecology Resources. 18(4):838-853.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012. Double digest radseq: An inexpensive method for de novo snp discovery and genotyping in model and non-model species. PloS one. 7(5):e37135.

Qiu T, Xian L, Crane J, Wen C, Hilton M, Lu W, Newman P, Cao X. 2015. Pth receptor signaling in osteoblasts regulates endochondral vascularization in maintenance of postnatal growth plate. J Bone Miner Res. 30(2):309-317.

Regan CE, Pilkington JG, Bérénos C, Pemberton JM, Smiseth PT, Wilson AJ. 2017. Accounting for female space sharing in st. Kilda soay sheep (ovis aries) results in little change in heritability estimates. Journal of Evolutionary Biology. 30(1):96-111.

Santure AW, Garant D. 2018. Wild gwas—association mapping in natural populations. Molecular Ecology Resources. 18(4):729-738.

Schipani E, Provot S. 2003. Pthrp, pth, and the pth/pthrp receptor in endochondral bone development. Birth defects research Part C, Embryo today : reviews. 69(4):352-362.

Shirali M, Knott SA, Pong-Wong R, Navarro P, Haley CS. 2018. Haplotype heritability mapping method uncovers missing heritability of complex traits. Scientific Reports. 8(1):4982.

Silva CNS, McFarlane SE, Hagen IJ, Rönnegård L, Billing AM, Kvalnes T, Kemppainen P, Rønning B, Ringsby TH, Sæther BE et al. 2017. Insights into the genetic architecture of morphological traits in two passerine bird species. Heredity (Edinb). 119(3):197-205.

Stoffel MA, Johnston SE, Pilkington JG, Pemberton JM. 2021. Genetic architecture and lifetime dynamics of inbreeding depression in a wild mammal. Nature communications. 12(1):2972.

Tang J, Zhou H, Sahay K, Xu W, Yang J, Zhang W, Chen W. 2019. Obesity-associated family with sequence similarity 13, member a (fam13a) is dispensable for adipose development and insulin sensitivity. International Journal of Obesity. 43(6):1269-1280.

VanRaden PM. 2008. Efficient methods to compute genomic predictions. J Dairy Sci. 91(11):4414-4423.

Wardhana DA, Ikeda K, Barinda AJ, Nugroho DB, Qurania KR, Yagi K, Miyata K, Oike Y, Hirata KI, Emoto N. 2018. Family with sequence similarity 13, member a modulates adipocyte insulin signaling and preserves systemic metabolic homeostasis. Proceedings of the National Academy of Sciences of the United States of America. 115(7):1529-1534.

Warrington NM, Beaumont RN, Horikoshi M, Day FR, Helgeland Ø, Laurin C, Bacelis J, Peng S, Hao K, Feenstra B et al. 2019. Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk factors. Nature genetics. 51(5):804-814.

Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB. 2006. Quantitative genetics of growth and cryptic evolution of body size in an island population. Evolutionary Ecology. 21(3):337.

Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, Chu AY, Estrada K, Luan Ja, Kutalik Z et al. 2014. Defining the role of common variation in the genomic and biological architecture of adult human height. Nature genetics. 46(11):1173-1186.

Wu Y, Murray GK, Byrne EM, Sidorenko J, Visscher PM, Wray NR. 2021. Gwas of peptic ulcer disease implicates helicobacter pylori infection, other gastrointestinal disorders and depression. Nature communications. 12(1):1146-1146.

Xia C, Amador C, Huffman J, Trochet H, Campbell A, Porteous D, Hastie ND, Hayward C, Vitart V, Navarro
P et al. 2016. Pedigree- and snp-associated genetics and recent environment are the major
contributors to anthropometric and cardiometabolic trait variation. PLoS Genet.
12(2):e1005804.
Xia C, Canela-Xandri O, Rawlik K, Tenesa A. 2021. Evidence of horizontal indirect genetic effects in
humans. Nature human behaviour. 5(3):399-406.
Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath AC, Martin NG,
Montgomery GW et al. 2010. Common snps explain a large proportion of the heritability for
human height. Nature genetics. 42(7):565-569.
Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. 2014. Advantages and pitfalls in the application
of mixed-model association methods. Nature genetics. 46(2):100-106.
Yashiro K, Zhao X, Uehara M, Yamashita K, Nishijima M, Nishino J, Saijoh Y, Sakai Y, Hamada H. 2004.
Regulation of retinoic acid distribution is required for proximodistal patterning and outgrowth of
the developing mouse limb. Developmental cell. 6(3):411-422.
Zaitlen N, Kraft P, Patterson N, Pasaniuc B, Bhatia G, Pollack S, Price AL. 2013. Using extended genealogy
to estimate components of heritability for 23 quantitative and dichotomous traits. PLOS
Genetics. 9(5):e1003520.

Age	Trait	No. individuals	No. records	Fixed effects	Random effects
Neonate	Birth weight	2678	2678	Sex	Year of birth
				Litter size	Mother ID
				Population size year before birth	
				Age of mother (quadratic)	
				Ordinal date of birth	
				Age (days)	
Lamb	Weight	2228	2228	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Foreleg	2284	2284	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Hindleg	2349	2349	Sex	Year of birth
				Litter size	Mother ID
				Population size	Permanent environment
				Age (days)	
	Metacarpal	2059	2059	Sex	Year of birth
				Litter size	Mother ID
				Age at death (months)	
	Jaw	2113	2113	Sex	Year of birth
				Litter size	Mother ID
				Age at death (months)	
Adult	Weight	1152	3553	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Foreleg	1121	3331	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Hindleg	1135	3444	Sex	Year of capture
				Population size	Permanent environment
				Age (years)	
	Metacarpal	945	945	Sex	Year of birth
				Age at death (years)	
	Jaw	991	991	Sex	Year of birth
				Age at death (years)	

676 **Table 1** Number of individuals and records, fixed and random effects fitted in each trait x age class
677 model in addition to the GRM. The same individuals and records were used for both heritability
678 estimates and for GWAS.

Gene Name	Ensembl Gene ID	Chr	Associated trait	Effects in other species
Cytochrome P450 26B1	ENSOARG00000011582	3	Lamb metacarpal length	Associated with skeletal abnormalities in humans and zebrafish (Laue et al. 2011), knockouts produce reduced limbs in mice (Yashiro et al. 2004).
EXOC6B	ENSOARG00000011607	3	Lamb metacarpal length	Associated with spondyloepimetaphyseal dysplasia (resulting in short stature) in humans (Campos-Xavier et al. 2018).
FAM13A	ENSOARG00000018727	6	Adult August weight	Modulates body fat distribution and adipocyte function in humans and mice (Fathzadeh et al. 2020) as well as adipose insulin signalling in mice (Wardhana et al. 2018), also linked with obesity in mice (Tang et al. 2019)
ONECUT1	ENSOARG00000020928	7	Birth weight	Associated with birth weight in humans (Warrington et al. 2019).
IFT43	ENSOARG00000002065	7	Adult foreleg length	Associated with Sensenbrenner syndrome (resulting in growth retardation and dwarfism due to femoral and humeral limb shortening) in humans (Arts et al. 2011).
PENK	ENSOARG00000020184	9	Adult August weight	PENK knock-out mice found to have diminished food motivation, lower baseline body weight and attenuated weight gain (Mendez et al. 2015)
PTH1R	ENSOARG00000006638	19	Lamb metacarpal length, adult foreleg length, adult hindleg length, adult metacarpal length	Involved in osteoblast development in mice (Qiu et al. 2015), associated with skeletal disorders such as EKNS (Duchatelet et al. 2005), JMC and BLC (Schipani and Provot 2003) in humans.
LTF	ENSOARG00000008620	19	Lamb metacarpal length, adult metacarpal length	Human LTF associated with increased bone growth when injected into piglets (Li et al. 2018), found to stimulate osteoblast

				proliferation (Cornish and Naot 2010). High expression levels in human bone marrow (Fagerberg et al. 2014).
BMP6	ENSOARG00000017264	20	Adult jaw length	Involved in bone development and expressed in the jaw bone in mice (Oralová et al. 2014).

Table 2 Potential candidate genes for future analyses. From left to right: gene name, Ensembl gene ID,
chromosome, associated trait, and evidence for association in other species.

Figure 1 Estimates of heritability for body size traits in neonates, lambs, and adult Soay sheep when
using a GRM calculated from the 50K SNP data (blue) compared with using a GRM calculated from the
imputed SNP data (yellow). Error bars represent standard error estimates.

Figure 2 Manhattan plots for A) birth weight GWAS using 50K SNP data (left) and imputed SNP data
(right); B) adult August weight GWAS using 50K SNP data (left) and imputed SNP data (right); and C)
adult metacarpal length GWAS using 50K SNP data (left) and imputed SNP data (right). The red line
represents the significance threshold ($2.49e^{-06}$ for the 50K SNP data and $1.03e^{-06}$ for the imputed SNP
data) – any SNPs above this threshold are considered to be significantly associated with variation in their
respective traits.

Figure 3 Miami plots for A) birth weight using imputed SNP data (top) and birth weight conditional
analysis using imputed SNP data (bottom); and B) adult jaw length GWAS using 50K SNP data (top left)
and imputed SNP data (top right), adult jaw length conditional analysis using 50K SNP data (bottom left)
and imputed SNP data (bottom right). The red line represents the significance threshold ($2.49e^{-06}$ for the
50K SNP data and $1.03e^{-06}$ for the imputed SNP data) – any SNPs above this threshold are considered to
be significantly associated with variation in their respective traits.

Data Availability Statement

All scripts and data can be found at
https://github.com/CaelinnJames/Impact_of_SNPDensity_on_Soay_Sheep

Acknowledgments

We thank the National Trust for Scotland and Scottish Natural Heritage for permission to work on St
Kilda and QinetiQ and Eurest for logistics and other support on the island. We also thank all those who
have been involved in the long-term project, including those who helped with field work on the island.
We thank the Wellcome Trust Clinical Research Facility Genetics Core in Edinburgh for SNP genotyping.

Funding

This work was supported by a NERC Doctoral Training Partnership grant (NE/S007407/1). The long-term
field project on St Kilda has been largely funded by the UK Natural Environment Research Council. The
SNP genotyping was funded by a European Research Council Advanced Grant. P. Navarro was supported
by the Medical Research Council (United Kingdom, grant number MC_UU_00007/10).

Heritability estimates for body size traits in neonates, lambs and adults

A

Birth weight GWAS

DSR SNP data

Imputed SNP data

B

Adult August weight GWAS

DSR SNP data

Imputed SNP data

C

Adult metacarpal length GWAS

DSR SNP data

Imputed SNP data

A
Birth weight conditional GWAS analysis

Conditioned in MDD, imputed SNP data

Conditioned in MDD, imputed SNP data

B
Adult jaw length conditional GWAS analysis

Original GWAS, with SNP data

Original GWAS, imputed SNP data

Conditional analysis, SNP data

Conditional analysis, imputed SNP data