G matrix stability in clinally diverging populations of an annual weed.

Georgia A. Henry[1*], John R. Stinchcombe[12]

[1] Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada

[2] Koffler Scientific Reserve at Joker's Hill, University of Toronto, King City, Ontario, Canada

***Correspondence:** georgia.henry@mail.utoronto.ca

Abstract

How phenotypic and genetic divergence among populations is influenced by the genetic

architecture of those traits, and how microevolutionary changes in turn affect the within-

population patterns of genetic variation, are of major interest to evolutionary biology. Work on

Ipomoea hederacea, an annual vine, has found genetic clines in the means of a suite of

ecologically important traits, including flowering time, growth rate, seed mass, and corolla

width. Here we investigate the genetic (co)variances of these clinally varying traits in two

northern range-edge and two central populations of *Ipomoea hederacea* to evaluate the influence

of the genetic architecture on divergence across the range. We find 1) limited evidence for clear

differentiation between northern and southern populations in the structure of **G**, suggesting

overall stability of **G** across the range despite mean trait divergence and 2) that the axes of

greatest variation (g_{max}) were unaligned with the axis of greatest multivariate divergence.

Together these results indicate the role of constraint on the divergence among populations across

the range.

Keywords: G-matrix, quantitative genetics, multivariate evolution, clinal divergence, *Ipomoea*
hederacea

Running title: Stable **G** across diverging populations

Introduction

A major goal in evolutionary biology is to understand the relationship between genetic variation within populations and phenotypic divergence between populations (Antonovics 1976, Endler 1977, Walsh & Blows 2009). Range edge populations provide an interesting setting in which to investigate this relationship. Populations existing on the margin of species' ranges are expected to have smaller population sizes (Brown et al. 1995), which are more susceptible to drift, and limited gene flow compared to more central populations (Sexton et al. 2009). Environmental gradients will often additionally create different selective pressures favouring phenotypic and genetic divergence. Here we examine the divergence and evolutionary potential of four populations of an annual plant, *Ipomoea hederacea*, sampled from the center and northern edge of the species' range with respect to five ecologically relevant traits.

The evolutionary response of a population undergoing selection is dependent not only on local selection, but on the underlying genetic architecture of the traits undergoing selection. The ability to adapt to the local environment depends on the amount of standing genetic variation as well as the genetic variance-covariance structure, represented as the **G** matrix (Antonovics 1976, Lande 1979, Lande & Arnold 1983, Falconer & Mackay 1996, Agrawal & Stinchcombe 2009). **G** is expected to shape evolutionary responses to selection as the relationships between traits lead to correlated responses selection (Lande 1979, Lande & Arnold 1983). Divergence among populations in response to local selection may thus be constrained or facilitated by the existing trait (co)variances. The multivariate trait combination with the greatest amount of genetic variation, dubbed the "genetic line of least resistance" by Schluter (1996) has been shown to influence macroevolutionary responses by deflecting the response to selection toward itself (Schluter 1996).

Whether **G** matrices themselves change along with trait means is not known. While **G**

matrices exhibit some stability over evolutionary timescales (Schluter 1996, Arnold et al. 2008)

we also expect population-level **G** matrices to undergo changes due to: bottlenecks (Roff 2000),

inbreeding (Phillips et al. 2001), drift (Lande 1979, Jones et al. 2004), strong selection (with

some genetic architectures) (Roff 2000), gene flow (Guillaume & Whitlock 2007), and genotype-

by-environment interactions (Wood & Brodie 2015). The **G** matrices of populations located in

ephemeral habitats or otherwise prone to extinction/colonization dynamics may reflect genetic

drift because of small population sizes prior to extirpation (Lande 1992) and/or the effects of

migration from limited or varied source populations due to colonization history (Whitlock &

McCauley 1990), and may not be at equilibrium. Many of the biological forces that can affect **G**,

in principle, could have contrasting effects between range-edge and central populations, making

strong *a priori* predictions difficult. For instance, if range-edge populations are smaller and more

subject to genetic drift, or under very strong selection because of novel conditions, we might

predict less overall genetic variance, or different patterns of genetic covariance, in these

populations. In contrast, if range-edge populations frequently experience swamping gene flow

from the center of the range (Mayr 1954, Mayr 1963, Kirkpatrick & Barton 1997; Garcia-Ramos

& Kirkpatrick 1997), their patterns of genetic covariance might show few differences compared

to core populations. Consequently, whether and how range edge populations differ in **G** remains

an empirical question.

Ipomoea hederacea, an annual plant which grows across the eastern U.S.A. from

Pennsylvania to Florida, has latitudinal genetic clines in a variety of quantitative traits (Stock et

al. 2014) and leaf shape (Bright 1998, Campitelli & Stinchcombe 2013), a Mendelian trait. These

clines do not appear to be due to drift, but rather a response to selection, as no latitudinal patterns

in neutral loci have been found (Campitelli & Stinchcombe 2013; Campitelli & Stinchcombe

2014). Previous work by Stock and colleagues (2014) focused on among-population patterns and

broad sampling found evidence of significant clinal divergence in three of the five traits

(flowering time, anther-stigma distance and corolla width) while the remaining traits (seed mass,

growth rate) had marginally significant divergence. These traits did not, however, exhibit a

reduction of genetic variation in northern range edge populations compared to the central

populations as was expected, suggesting there may still be sufficient variation for continued

adaptation. Stock et al. (2014) estimated the axis of greatest multivariate divergence, that is the

mean trait combination that is most different between populations of *I. hederacea* across

latitudes, which we use here alongside more deeply sampled populations to compare within-

population trait organization and among-population trait divergence.

Our research aims were: 1) to determine whether northern and southern populations

differed in the overall heritability and genetic variance of the focal traits, 2) evaluate how genetic

variation within populations was aligned relative to the axis of multivariate genetic divergence,

and 3) assess population level divergence of the **G** matrices. To do so, we estimated trait

heritability and the genetic variance-covariance matrices of four populations of *Ipomoea*

hederacea; collected from the northern range edge (Pennsylvania and Maryland) and the core of

the range (Hoffman and Ellerby, North Carolina). We expected the axis of trait divergence

between populations to be biased toward the axis of greatest variation within populations, as this

is the genetic line of least resistance (Schluter 1996).

Methods

Study species

Ipomoea hederacea is an annual weedy vine with an eastern North American distribution

that spans Florida to Pennsylvania, USA (Bright 1998). While *I. hederacea* has existed in the

present range for at least 150 years, its provenance is disputed, with some sources claiming it is

native to its current range while others suggest it is introduced from the American tropics (Bright

1998, Campitelli & Stinchcombe 2014). *Ipomoea hederacea* is capable of mixed mating, but

typically has a high selfing rate, with overall estimates ranging from 66-94% (Ennos 1981, Hull-

Sanders et al. 2005, Campitelli & Stinchcombe 2014). Despite significant population structure,

work using putatively neutral loci have found weak isolation-by-distance and no evidence of

clines in neutral diversity (Campitelli & Stinchcombe 2013). There are, however, latitudinal

clines in leaf shape (Bright & Rausher 2008, Campitelli & Stinchcombe 2013) and a suite of

quantitative traits (Klingaman and Oliver 1996; Stock et al. 2014)

Propagation and trait measurements

We harvested seed from populations from the edge of agricultural fields in Pennsylvania

(40.116611°, 76.398889°), Maryland (39.581861°, 77.816861°), Hoffman, North Carolina

(35.074139°, 79.556694°) and Ellerby, North Carolina (35.091722°, 79.742722°). We sampled

seed pods haphazardly from vines 1-2 m from one another, to reduce the chance of sampling

twice from the same maternal plants, which were the founders of the matrilines we used in our

experiments. We grew one seed from 50 randomly chosen matrilines per population in a

common greenhouse environment and allowed these individuals to self-fertilize. One death prior

to seed set resulted in 49 matrilines in Ellerby, NC. We harvested selfed seeds from each

maternal plant and grew approximately 11 focal individuals (range: 2-11, mean = 10.7) from

each line for this study, resulting in a total sample of 2,133 plants. While our use of selfed

progeny precludes estimating additive genetic variance, selection acts on the broad sense

variation rather than just the additive components of variation in mainly selfing species

(Roughgarden 1979), making this design appropriate for *I. hederacea*.

We grew plants in cone-tainers (Stuewe & Sons, Oregon, USA) filled with Pro-Mix soil

in a glasshouse at the University of Toronto from April-October 2018. All plants were fertilized

with a weak 10-52-10 fertilizer solution every 2 weeks. Day length was initially 16 hours at

28°C, changed to 12 hours at 25°C on day 32, and 8 hours at 22°C on day 38 to promote

flowering.

Trait measurements followed Stock et al. (2014). Prior to planting, we weighed each seed

and recorded individual mass. We counted leaves at 17 days after planting and again on the 26th

or 27th day of planting; we estimated growth rate as the difference in leaf counts divided by the

number of days, and as such it is in units of new leaves/day. We measured corolla width and

anther-stigma distance on the first flower of each individual using calipers. We measured the

distance from the lowest and highest anthers to the stigma and calculated the mean of the

absolute distance to characterize anther-stigma distance.

Data preparation

Four of the five focal traits, (seed mass, early growth rate, corolla width and anther-

stigma distance) were standardized to mean = 0 and standard deviation = 1 using the grand mean

and standard deviation across populations to allow for meaningful comparison of differences

(Hine et al. 2009, Hansen & Houle 2008). We elected to use this standardization to eliminate

differences in the scale and units of our traits (e.g., anther stigma distance in mm, growth rate in

leaves per day, and flowering time in days) and because mean standardization can be difficult to

apply to traits like flowering time, which have an arbitrary rather than natural zero (Hansen &

Houle 2008, Houle et al. 2011). As the glasshouse was exposed to ambient light, the change in

artificial light used to induce flowering was largely overwhelmed by the natural photoperiod. A

change from lengthening to shortening days interrupted the flowering schedule. For this reason

we transformed the number of days until the first flower using ordered quantile normalization

(ORQ), which preserves the order of flowering but results in a gaussian distribution (Peterson

and Cavanaugh 2019). The ORQ normalization results in a distribution centered on zero with a

standard deviation of one, which is the same scale as all other traits.

Univariate analysis

To determine broad sense heritabilities of the traits we fit Bayesian generalized linear

mixed-effect models for each trait in each population separately. We estimated the models using

MCMCglmm in R (Hadfield 2010) using the mixed model,

$$y = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}_1\mathbf{u}_1 + \mathbf{Z}_2\mathbf{u}_2 + \mathbf{e} \qquad\qquad (1)$$

For each trait, y, we ran a separate model. In (1), $\mathbf{X}$, $\mathbf{Z}_1$ and $\mathbf{Z}_2$ represent design matrices for the

fixed effect of population, and random effects of matriline and greenhouse block, respectively. $\boldsymbol{\beta}$

and $\mathbf{u}$ are vectors of the related parameters. We included the environmental block to control for

the effect of microenvironmental differences within the glasshouse. For seed mass, we used the

greenhouse block of the maternal plant. The residual error is represented by $\mathbf{e}.$

We tested six priors and summed the Deviance Information Criteria for each of the five

univariate models. We used the prior with the lowest summed DIC score, although due to the

amount of data in the model the prior choice was not very influential. The effective size of the

samples was greater than 85% of the total number of samples and the autocorrelation was below

0.05 for all parameters (Hadfield 2010). To evaluate whether there was meaningful broad sense

heritability, we ran models within each population with and without the matriline variable and

compared DIC values, per Puentes et al. (2016). We considered models where the matriline

reduced the DIC score >2 to have a "significant" genetic component and thus H2 was significant.

We constructed comparable models for validation, using REML, which are presented in the

Supplemental Material. Heritability estimates were similar, with the greatest difference in

Heritability of 5.2%, and an average absolute difference of 1.7%.The 95% Highest Posterior

Density (HPD) intervals from Bayesian models overlapped REML heritability estimates in all

cases, and as such Bayesian estimates are presented in the main text.

Multivariate Analyses

We estimated the total genetic variance-covariance matrices separately for each

population using Bayesian generalized linear mixed-effect models through MCMCglmm in R

(Hadfield 2010). We again used the mixed model,

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}_1\mathbf{u}_1 + \mathbf{Z}_2\mathbf{u}_2 + \mathbf{e} \tag{3}$$

where $\mathbf{X}$, $\mathbf{Z}_1$ and $\mathbf{Z}_2$ represent design matrices for the vectors of trait means ($\boldsymbol{\beta}$)**,** the total genetic

effects ($\mathbf{u}_1$) and the Greenhouse Block effects ($\mathbf{u}_2$). The residual error is represented by the term

$\mathbf{e}$.

We used weakly informative inverse-Wishart priors for estimating the variance and

covariances where the distribution was described by the variance (V) and the belief parameter

(nu) and the default prior for the fixed effect estimates, $N(0,10^8)$. We tested a variety of priors to

determine the robustness of estimates and selected the model with the best fitting prior,

determined by the lowest average Deviation Information Criteria score to use for our analyses

(see Supplementary Materials). The final prior for the residual variance was given by a diagonal

matrix of 1's with the degree of belief slightly above *n*-1, for *n* traits (V=diag(5), nu = 4.001).

The best prior for the random effects was a matrix of ⅓ the phenotypic variance-covariance

matrix and a belief parameter slightly above *n*-1 (V = (**P**/3)*10, nu=4.001, where **P** is the

phenotypic covariance matrix). All the response variables in the model were multiplied by a

factor of 10, which improves the estimation of small variances (and is subsequently factored out

of the genetic variance-covariance matrix). To account for this factor of 10 in the response

variables, the Phenotypic variance-covariance matrix (**P**) was also multiplied by a factor of 10 in

the priors.

We ran full models for 5,005,000 iterations, with a burn in of 5,000 and a thinning

interval of 1,000, resulting in 5000 iterations sampled from the posterior distribution. The

effective size of the samples was greater than 85% of the total number of samples and the

autocorrelation was below 0.05 for all parameters (Hadfield 2010).

Null Models

For some analyses, a null model was necessary, as MCMC methods can only estimate

variance greater than zero (matrices must be positive definite). We constructed randomized **G**

matrices where the values only reflected sampling error as a point of comparison to our actual

estimates. Following Walter et al. (2018) and McGoey and Stinchcombe (2021), we randomly

assigned individuals to matrilines within populations, without replacement, resulting in **G**

matrices that are only due to sampling error, as any phenotypic similarity among individuals is

only due to random sampling, rather than true genetic ancestry. Our design includes significant

fixed effects, which Morrissey et al. (2019) note will make these null distributions larger than

they should be, which should make our approach conservative. We also note that because each

randomized **G** matrix only reflects sampling variation, the expectation of a large number of **G**

matrix comparisons should be near 0 (as each population only reflects sampling variation), but

the distribution of these comparisons reflects the range of potential outcomes one can obtain if

each **G** only reflects sampling, making it a sensible null model for comparisons. The within-

population randomization approach also allows us to compare whether each estimated **G** differs

from expectations due to sampling (Sztepanacz and Blows 2017).

We fed each randomized "population" into the same models as the observed data, run for

20,000 iterations with a burn in of 5,000 and a thinning interval of 100. We iterated the

randomization and model estimates 1000 times. We combined the final posterior sample of each

of the randomized models within populations to be used as a composite posterior. We assessed

chains visually for stability and with Gelman and Rubin convergence diagnostic criteria.

G matrix comparisons

We used several metrics to compare G matrices, each of which has different strengths

and weaknesses (Krzanowski 1979, Kirkpatrick 2009, Hine et al 2009, Aguirre et al. 2014,

Puentes et al. 2016, Walter et al. 2018, Teplitsky et al. 2014). In general, our goal was to use a

variety of metrics to compare G matrix size, shape, concordance between patterns of genetic

variance and clinal divergence, and similarity in the responses to selection that each G matrix

would produce.

G matrix dimensionality

We calculated the trace of the **G** matrices to determine the total amount of genetic

variance in each population. We then evaluated the overall strength of the covariance by

calculating the effective number of dimensions, n_D (Kirkpatrick 2009), which is the sum of the

eigenvalues divided by the leading eigenvalue. The effective number of dimensions ranges

between 1, which would indicate that all of the genetic variance was in a single direction in

multivariate space, and the number of traits, which would indicate that the traits are equally

genetically variable and uncorrelated.

230   Genetic variation in the direction of maximum clinal divergence

To evaluate the alignment between the axis of greatest genetic variation (g_{max}) of each

population, and between g_{max} and the vector of greatest multivariate divergence (Cline$_{max}$ from

Stock et al. 2014) we calculated the correlation coefficient between these vectors. We used

random unit vectors as a null comparison. We note that Stock et al. (2014) used and presented

results from mean-standardized data; we re-analyzed their data to estimate the vector of greatest

clinal divergence using standard deviation standardized data, to match our use of standard

deviation-standardized data from the greenhouse experiment.

We projected the vector of maximum clinal divergence through each **G** matrix (using

b^T**G**b, where b is the vector of clinal divergence; see Lin and Allaire 1977) to calculate the

genetic variance in direction of clinal divergence. To estimate the proportion of genetic variance

along the direction of maximum clinal divergence, we estimate b^T**G**b/$\boldsymbol{\lambda}_1$, where $\boldsymbol{\lambda}_1$ is the first

eigenvalue of **G**, representing the vector of maximum genetic variance in each population. Given

that the **G** matrices had different shapes, the maximum amount of genetic variation in any

direction differed between the populations. To allow for fair comparison, we standardized by the

proportion of variance associated with g_{max}, the first eigenvector of **G**.

246   Random skewers

We performed a random skewers analysis (Cheverud & Marriog 2007, Aguirre et al.

2014) following Aguirre et al. 2014. Briefly, 1000 randomly generated normal vectors were

projected through each population's **G**. We compared the response from each projection between

populations to determine which random vectors which produced response differences. We then

calculate the variance-covariance matrix of the skewers which result in response differences

between populations and perform eigenanalysis to generate the **R** matrix. The **R** matrix describes

the trait space where differences in the response to skewers exist between populations, with the

first eigenvector describing the axis with the greatest differences between populations.

Krzanowski's subspace analysis

256   Krzanowski's subspace analysis provides a method of evaluating the similarity in

257 geometry of multiple variance-covariance matrices. The approach works by examining whether

258 the subspaces containing most of the genetic variation (e.g., the leading PCs) in each population

259 are shared, or in common. We first determine the shared space (**H**) of all populations, from a

260 subset of the leading PCs (eigenvectors) of each population, using:

261    $\mathbf{H} = \sum_{t=1}^{p} \boldsymbol{A}_t \boldsymbol{A}_t^T$         **(4)**

where superscript T indicates transposition, and a subset t of PCs of each **G** matrix are included

in **A**, and summation is to p populations ($p = 4$ in our case; Krzanowski 1979; Aguirre et al.

2014). As per Aguirre et al. 2014, the number of eigenvectors used for each population were

those which explained >90% of the total variation to account for differences in the shape of the

population **G** matrices. The eigenvalues of **H** can range from 0 to p, which indicates common

sub-spaces to the matrices being compared (i.e., the leading PCs describe similar directions of

multivariate space). If the eigenvalues of H are less than p (4 in our case), this indicates that

genetic variation described by that eigenvector differs among the populations. To evaluate which

populations are most similar we determine the smallest angle between the mean posterior

subspace of each population and the mean shared subspace **H**, as per Aguirre et al. 2014.

Tensor analysis

Differences between matrices can be evaluated by calculating the 4th order covariance

tensor, describing the variances and covariances between matrices (Hine et al. 2009). We used

the 4th Order Genetic Covariance Tensor analysis developed by Hine et al. (2009), and outlined

by Aguirre et al. (2014), and Walter et al. (2018). We refer interested readers to these papers for

full details (especially Fig. 2 of Walter et al. 2018), and only provide a brief overview here. The

fourth order tensor of the matrices describes the variance and covariances among the population

G matrices. The tensor can be represented as a symmetric (n(n+1)/2) matrix (**S**) (Basser &

Pajevic 2007), with elements describing the variances and covariances of genetic variances and

covariances in the four populations. The matrix **S** can then be subject to eigenanalysis, where

eigenvalues and eigentensors describe how the **G** matrices vary. Eigentensors with larger

eigenvalues (which explain a larger percentage of variance) describe dimensions of greater

variation in the **G** matrices; the eigentensors can themselves be subjected eigenanalysis, to

determine the linear combination of traits (eigenvectors) that lead to the difference in **G** matrices

captured by the eigentensors. The covariance tensor analysis can thus be used to evaluate axes of

differences among the genetic variance-covariance structure of the populations.

Results

Univariate Analyses

We estimated broad-sense heritability of each trait within populations using a Bayesian

model (Table S1, Figure S1), and confirmed results with REML (Table S2) as described above.

All traits had heritability greater than zero in at least 3 of the 4 populations (Table S1). The

Pennsylvania population lacked significant heritability for anther-stigma distance (Table S1).

The mean of all the heritability estimates was 0.213 (ranging from 0.011 to 0.535). There was no

significant difference between northern and southern populations, although northern populations

tended to have lower heritability estimates than southern populations (Table S1).

G matrix dimensionality

We present population **G** matrices, 95% HPD intervals, and null expectations due to

sampling variation in the Supplemental Material (Table S3, Table S4), and in the main text focus

on comparisons of the matrices. Our **G** matrices all contain variances with HPD intervals that do

not overlap the 95% HPD intervals of the null **G** matrices, although some traits in some

populations did have overlapping 95% HPD intervals indicating that the variation may be very

small (Table S3). Two of the traits, corolla width and anther-stigma distance, have variation

indistinguishable from sampling variation in all populations except Hoffman (Table S3); we

nonetheless retained these traits in our analyses to allow comparisons of **G** properties to

divergence in trait means, which requires these traits. All populations have an effective number

of dimensions considerably less than the number of traits measured (Table 1), indicating that the

genetic covariance is important in all cases. There were no population level differences, as the

95% HPD intervals of all populations overlapped one another. The northern populations

(Pennsylvania and Maryland) had slightly smaller effective dimensions indicating overall greater

correlation among traits, but the 95% HPD intervals overlapped.

We found that Hoffman, NC had significantly greater total genetic variation than all other

populations, which we estimated as the trace of the **G** matrices. Hoffman had approximately

twice the amount of genetic variation than populations from Ellerby, NC and Maryland, and

three times that of Pennsylvania. The 95% HPD intervals of Hoffman did not overlap with those

of any other population.

Table 1: Effective number of dimensions (N dimensions) for each population **G** matrix and the total
amount of genetic variation in each population, estimated as the trace of **G**. Low and high values
demarcate the 95% highest posterior density interval (HPD).

Population	N dimensions	Lower HPD	Upper HPD	Trace of G	Lower HPD	Upper HPD
Pennsylvania	1.887	1.440	2.371	0.518	0.367	0.685
Maryland	1.887	1.454	2.378	0.764	0.545	1.006
Hoffman, NC	2.068	1.579	2.608	1.615	1.208	2.054
Ellerby, NC	2.007	1.535	2.509	0.886	0.644	1.176

Genetic variation in the direction of maximum clinal divergence

Overall the proportion of genetic variation in the direction of maximum clinal divergence was

small, roughly on par with the third eigenvalues of each population, and there were no significant

differences between any populations (Table 2). To fairly compare between populations, which

differ in their total amount of genetic variation (Table 1) and the variation along g_{max} (Table S5),

we standardized the genetic variance in the direction of clinal divergence by the genetic variance

associated with g_{max}, so the values are relative to the linear combination of traits with the

maximum possible genetic variance in each population. The standardized projections of the

northern populations were slightly larger than the southern populations, but there were no

significant differences between them. In general, genetic variance in the direction of clinal

variation was ~25-50% of the genetic variance associated with g_{max}.

Table 2: Genetic variance in the direction of maximum clinal divergence for each population. In the
standardized column (Variance (std)), the variances are standardized by the eigenvalue of the leading
eigenvector of each population, to express this relative to the direction of greatest genetic variation.
Lower and higher HPD values demarcate the 95% highest posterior density interval.

Population	Variance	Lower HPD	Higher HPD	Variance (std)	Lower HPD	Higher HPD
Pennsylvania	0.102	0.047	0.176	0.371	0.174	0.602
Maryland	0.157	0.076	0.252	0.383	0.203	0.570
Hoffman, NC	0.204	0.092	0.325	0.266	0.104	0.456
Ellerby, NC	0.227	0.106	0.361	0.512	0.266	0.782

The correlation between gmax from each of the G matrices and direction of maximum clinal

divergence is low and not significantly different from the expected correlation between gmax

and a random vector (Figure 1). Although we might expect the direction of maximum clinal

divergence to be aligned with directions of genetic variation, this does not appear to be the case

for any population. The correlation between PC2 through PC5 of the population G matrices and

direction of maximum clinal divergence followed the same pattern (Figure S2).

Figure 1: The correlation between the g_{max} of each population **G** matrices and the vector of greatest
multivariate clinal divergence (Observed, in blue) compared to the correlation between the g_{max} of each
population **G** matrices and randomized vectors (Randomized, in, in gray). Error bars represent the 95%
HPD intervals. "Penn" is Pennsylvania, "Mary" is Maryland, "Hoff" is Hoffman, NC and "Ellr" is
Ellerby, NC.

Random skewers

The majority of random vectors produced some difference in response between

populations, with 588 of 1000 resulting in responses where at least one pairwise comparison had

95% HPD intervals which did not overlap. Following Aguirre et al. (2014), we took this subset

of vectors which resulted in response differences and created a variance-covariance matrix of the

vectors. We then performed eigenanalysis to evaluate the trait space which describes differences

between populations. The highest trait loading of the first eigenvector is corolla width, followed

by seed mass and flowering time (table 3). These differences in response are largely attributable

to Hoffman, NC. Projecting the eigenvectors of the **R** matrix through each population's **G**

matrix, we found that Hoffman had significantly more genetic variation along this eigenvector

than the other three populations (Figure 2). In other words, the response difference we detected

with random skewers appears to be due to the Hoffman population having more genetic variation

for a linear combination of traits made up primarily of seed mass and corolla width, with lesser

but nearly equal contributions of growth rate and anther stigma distance, than the other three

populations.

Table 3: Eigenanalysis of the **R** matrix from the Random Skewers analysis which resulted in difference
in response when projected through populations. The **R** matrix summarized the trait combinations which
resulted in differences to the projected response. Each eigenvector (RSe1 through RSe5) describes an
independent combination of traits which capture differences among populations. The eigenvalues describe
the proportion of variation of each eigenvectors within **R**, larger values indicate more of the variation in
the skewers are explained.

Trait	RSe1	RSe2	RSe3	RSe4	RSe5
Seed mass	-0.579	-0.147	0.211	-0.404	0.660
Growth	-0.262	-0.131	0.776	-0.075	-0.554
Flow time	-0.337	-0.587	-0.106	0.727	0.052
Corolla	-0.622	0.124	-0.556	-0.234	-0.483
AS dist.	-0.309	0.776	0.180	0.498	0.149
Eigenvalues	0.303	0.253	0.232	0.115	0.095

The second and third eigenvalues of **R** are of similar magnitude, and when projected

through the **G** matrices, reveal similar patterns. The second axis of **R** captures trait combinations

with opposing and larger values for anther-stigma distance and flowering time (Table 2); we

observe that Hoffman had more genetic variation for this combination of traits than the Maryland

and Pennsylvania populations. The third axis of **R** reflects trait combinations with opposing and

larger values for corolla width and growth rate, and Hoffman had more variation for that

combination of traits than the Maryland population. Collectively, these data indicate that the

differences in the **G** matrices detected by random skewers were primarily driven by the Hoffman

population, and it having more variation linear combinations of traits involving all of the traits

measured (*RSe1*: corolla width and seed mass in concert; *RSe2*: flowering time and anther stigma

distance in opposition; *RSe3*: corolla width and growth rate in opposition). The Hoffman

population also had the highest heritability for all of the measured traits (Table S1), the largest

effective number of dimensions (Table 1), and the largest trace of **G** (Table 1).

Figure 2: Genetic variance (Vg) in the direction of each eigenvector of the **R** matrix, where the **R** matrix is composed of vectors which resulted in response differences between populations.

Krzanowski's subspace analysis

To construct **H**, the shared subspace, we used the number of eigenvectors which

explained at least 90% of the variation in each population. The first three eigenvectors of each

population satisfied this condition. Lambda, which ranges from 1 to 4, approaches 4 in the first

two eigenvectors of **H** (Table 4, Figure 4). A lambda value nearing 4 means the populations all

contain the variation sufficient to recreate the shared trait space. Thus, the populations contain

considerable shared trait space in the first two dimensions of **H**. The third eigenvalue is lower,

but is similar to the randomized comparison (Figure 4). Overall there is no evidence of

divergence among populations in the shared subspace.

Table 4: Eigenvalues (lambda) for each eigenvector of the shared subspace **H** with the 95% HPD
intervals. A value of 4 would indicate that all the populations have the variation in their subspaces
required to recreate the respective eigenvalue of **H**, lower values indicate that populations do not and thus
lack similarity.

Eigenvector	Lambda	Lower HPD	Higher HPD
h1	3.840866	3.655673	3.978588
h2	3.04281	2.477581	3.798831
h3	2.284866	1.538301	3.009639

To evaluate any population level divergence from the shared subspace we can calculate

the angle between the axes of **H** and the subspaces of the populations ($\mathbf{A}_t\mathbf{A}_t^T$), which capture at

least 90% of the genetic variation in each population. Larger angles indicate that population

differs in some manner along that axis of shared subspace. The first three eigenvectors of

Pennsylvania, Hoffman and Ellerby, and Maryland were used to generate the shared subspace.

Overall the angles between **H** and the population subspaces are similar and small with increasing

uncertainty for higher eigenvectors (Figure S3), meaning differences between the populations

and the shared trait subspace are minimal.

Figure 3: Eigenvalues (lambda) for each eigenvector of the shared subspace **H**, (where h > min k).
Higher values of lambda indicate greater similarity of population subspaces.

Covariance tensor analysis

We calculated the fourth order covariance tensor for the population **G** matrices to assess

areas of divergence between the populations. We used the randomized **G** matrices, which

represent sampling variation within populations, as a null contrast to assess significance. For the

observed **G** matrices, we estimated three non-zero eigentensors for tensor **S**. Of these three

eigentensors, the first two have eigenvalues significantly different from a null expectation based

on sampling variation creating each **G** matrix independently (Table S6, Figure 6).

Figure 4: A) Eigenvalues (α) for each non-zero eigentensor from the observed and randomized **G**
matrices. Note that the upper and lower bounds of the randomized data, reflecting only sampling
variation, are small and hard to distinguish on the figure. B) Coordinates of each population in the first
non-zero eigentensor of **S**, E1. C) Coordinates of each population in the second non-zero eigentensor of **S,**
E2.

To determine the contributions of each population to the eigentensor, we estimated the

coordinates of the populations within each eigentensor. The greater the absolute value of the

coordinate the greater the contribution a population has. In the first eigentensor (E1, which

explains 81.5% of the variation across all non-zero eigentensors), Hoffman again stands out from

the other populations and has the greatest contribution. Thus the first eigentensor, which

describes the greatest amount of variation among the population **G** matrices, is largely driven by

differences between Hoffman, NC and all the other populations. In contrast, Ellerby, NC has the

largest coordinate value in the second eigentensor (E2), which explains 12.8% of the variation

among **G** matrices.

Eigenanalysis of the eigentensors reveals that the first eigenvector of E1, (*e11*) which

explains 47.6% of overall variation (58.4% of variation within E1), is most heavily weighted by

seed mass and corolla width (similar to *e1* of the **R** matrix, as analyzed by Random Skewers),

with all trait loadings being of the same sign (Table 5). The first eigenvector of E2 *(e21)*, which

explains 5.9% of overall variation among populations (45.9% of variation in E2), describes a

dimension most heavily weighted by the seed mass and growth rate of opposing signs (Table 5).

Thus these two eigentensors describe population differences in the relationship between seed

mass and other traits, especially with respect to the southern populations.

Table 5: Eigenvectors of each non-zero eigentensor of observed **S**. The variance explained by each
eigentensor of **S** is given, along with the eigenvalues of each eigentensor in the direction of each
eigenvector. These values describe the contribution of the independent trait combinations described by the
eigentensors. The proportion of variance is the proportion of variance of the eigentensors explained by
each eigenvector. The trait loadings for eigenvalues of each of the three eigentensor are given, along with
the eigenvalue of the eigentensors.

Eigenvector	σ of E_x	Eigenvalue	Prop. of σ	Seed mass	Growth	Flow time	Corolla	AS dist.
e1.1	0.815	-0.912	0.584	-0.611	-0.152	-0.323	-0.612	-0.354
e1.2		-0.317	0.203	0.044	0.379	-0.674	-0.140	0.617
e1.3		-0.254	0.163	0.568	0.519	-0.088	-0.342	-0.532
e1.4		0.049	0.031	0.542	-0.751	-0.312	-0.210	0.034
e1.5		-0.031	0.020	0.089	0.031	0.581	-0.667	0.457
e2.1	0.128	0.830	0.459	0.696	-0.639	-0.284	-0.097	0.134
e2.2		0.406	0.225	0.631	0.621	0.247	-0.387	-0.077
e2.3		-0.329	0.182	0.070	-0.288	0.896	0.167	0.286
e2.4		0.189	0.105	-0.253	0.019	-0.104	-0.596	0.754
e2.5		-0.054	0.030	-0.222	-0.352	0.212	-0.676	-0.571
e3.1	0.057	0.651	0.347	-0.005	0.811	0.441	0.315	0.220
e3.2		-0.640	0.341	0.946	-0.089	0.000	0.300	-0.081
e3.3		0.362	0.193	-0.228	-0.349	0.567	0.473	-0.530
e3.4		0.184	0.098	-0.116	-0.415	0.047	0.455	0.778
e3.5		-0.041	0.022	-0.197	0.202	-0.695	0.617	-0.240

We next projected these eigenvectors through the population **G** matrices, as we did for

the random skewers, to assess how much variation each population has along these axes. Not

unexpectedly, Hoffman, NC has significantly greater genetic variation along the first and second

eigenvector of E1 (Figure 5). Hoffman again has greater variation than the other population

along the first eigenvector of E2, although the HPD intervals overlap for Ellerby, NC, the other

southern population. The second eigenvector of E2 similarly has overlapping 95% HPD intervals

for all populations but the southern populations have slightly more genetic variation in this

direction than the northern populations.

Figure 5: Genetic variation in the direction of eigenvectors of the fourth-order covariance tensor **S**. e11 is
the first eigenvector of the first eigentensor, e12 is the second eigenvector of the first eigentensor, e21 is
the first eigenvector of the second eigentensor, e22 is the second eigenvector of the second eigentensor.
The vectors were projected through each of the **G** matrices to determine the amount of genetic variation
that exists in that dimension.

Similarity between skewers and tensors

Given the similarities between the first eigenvector of E1 of **S** and the results from the

random skewers **R** matrix projection we calculated the correlation between the two vectors and

478 found that they are almost perfectly correlated (corr = 0.992). This congruence between two

479 vectors of differentiation calculated with entirely different methods provides additional support

480 for this axis describing considerable differences among populations.

Discussion

Understanding patterns of genetic variation within populations, and how they relate to

phenotypic differentiation among them, is a central goal in evolutionary biology. Our analyses of

genetic (co)variance matrices and divergence of four natural populations of *Ipomoea hederacea*

point to three main results. First, we found that overall correlations between traits are important,

reducing the dimensionality of **G** in all four populations, and thus the potential for independent

phenotypic trait evolution. While northern populations tended to have greater dimensionality

reduction, reduced overall genetic variation, and lower trait heritabilities, the posterior

distributions of these parameters overlapped for northern and southern populations. Second,

contrary to our expectations, the axis of greatest multivariate clinal divergence was not aligned

with the axis of greatest variation in any population, suggesting a role for constraint and selection

in the divergence of these populations. Third, we found that the populations tended to have

similar patterns of **G** matrix structure and inhabit the same trait space, with the notable exception

of one population. Below, we discuss these results in light of how genetic constraints and

reduced dimensionality of **G** influence phenotypic divergence, and divergence in **G** these

populations.

Evolutionary potential and divergence of G

Our study of **G** within these four populations allows both an evaluation of the

evolutionary potential within each population, and the divergence (or lack thereof) of **G** itself

between range-edge adjacent populations and the central core. Using a variety of approaches,

including estimating the effective number of dimensions, the trace of **G**, and a handful of

methods for comparing **G** among populations (Krzanowski's subspace, covariance tensors,

random skewers), we found a consistent pattern of results. These populations all appear to be of

reduced rank, have similar amounts of total genetic variance, and there do not appear to be

appreciable differences between range-edge populations and core populations in the structure,

size, or shape of **G**.

We first assessed the general structure of the populations and found that all four have an

effective number of dimensions around two, appreciably less than the number of traits evaluated.

Our results indicate that while we might observe and measure these traits separately (and can

detect significant genetic variation in them, individually), the patterns of genetic covariances in

them mean that there are effectively fewer independent traits. Reduced effective dimensionality

suggests the importance of these trait correlations in all of the populations. Much of the

multidimensional trait space lacks detectable genetic variation (Kirkpatrick 2009), and

populations are likely to be evolutionarily constrained if selection were favoring trait

combinations completely lacking variation (see Kingsolver et al. 2015). While there were no

differences in the effective number of dimensions, there were differences in the amount of total

genetic variation among populations. Hoffman, NC, one of the southern populations, possessed

significantly greater variation than all other populations. While the second southern population,

Ellerby, NC did have slightly higher genetic variation, the HPD intervals overlap those of the

other populations. Similarly, trait heritabilities tended to be higher in southern populations (Table

S1), albeit again with overlapping HPD intervals. As such, it appears that the Hoffman

population is distinct from the rest, rather than southern populations on the whole differing from

northern populations. Two of the traits, corolla width and anther-stigma distance, had very small

variances indistinguishable from sampling variance in three of the populations (Hoffman again

being the exception). The lack of variation in in these traits in **G** make observing differences

among populations impossible with respect to corolla width and anther-stigma distance. We

retained these traits in our **G** matrices so as to allow for comparison with the axis of greatest

multivariate divergence, Cline$_{max}$ (from Stock et al. 2014). In Cline$_{max}$ both corolla width and

anther-stigma distance showed significant among-population divergence of the means, which

suggests that the paucity of within-population variation for these traits is due to selection.

Our analysis of the geometric similarity of **G** through Krzanowski's subspace analysis

showed that the majority of within-population genetic (co)variation of our populations share

overlapping trait space. Our results are suggestive of the overall stability of the **G** matrices across

populations, with no evidence of differences between range-edge and core populations in the trait

space where most (co)variation lies. In this regard, our results are similar to McGoey and

Stinchcombe (2021), who found little difference in **G** matrices between introduced and native

populations of ragweed (*Ambrosia artemisiifolia*) despite expectations to the contrary. More

generally, our results add to other findings of **G** being relatively stable across geography (Arnold

et al. 2008; Delahaie et al. 2017). Work on the cuticular hydrocarbons of *Drosophila serrata*

population by Hine and colleagues (Hine et al 2009, Chenoweth et al. 2008) did find patterns in

mean trait divergence, reduction in associated variation in g_{max}, and divergence among

populations along g_{max}. Similarly, Aguirre et al (2014) found that the populations share

considerable trait space, although there were axes of divergence among the populations.

The independently estimated sexual selection gradient is nearly orthogonal with respect to g_{max},

however, and thus the mostly likely explanation for the majority of the divergence among these

populations was drift (Hine et al 2009).

Both the covariance tensor and random skewers analysis suggest that the Hoffman, NC

population differed from the others (this population also had the most genetic variation overall,

the highest effective number of dimensions, and the most genetic variance in the direction of

clinal divergence). The covariance tensor and skewers approaches are complementary, in that the

tensor reflects patterns of variation and covariation among G, while the skewers analysis is not

sensitive to differences in the total amount of genetic variance and magnitude of the response (as

it uses the correlation between response vectors, but not their magnitude; Hansen and Houle

2008) and so these differences are not only due to Hoffman's greater overall genetic variation.

The first eigenvector of the first eigentensor (*e11*), the first eigenvector of the random skewers

analysis (*RSe1*), and g_{max} within the Hoffman population were all highly correlated, with heavy

loading from flowering time, corolla width, and seed mass. While the greater overall genetic

variation in Hoffman may not have directly led to these differences between matrices, it may

have allowed Hoffman greater evolvability, allowing for an increase in observable differences

when compared to the other populations. The Hoffman, NC population site did not differ

substantially from the others: all were collected from roadsides adjacent to agricultural fields,

and were reasonably large to allow for sampling from 70-100 individuals. Hoffman was,

however, the only population where we observed both leaf shape genotypes (leaf shape is a

simple Mendelian polymorphism in this species; Bright 1998). Some speculative possibilities are

that it was founded by heterozygous individuals, multiple founders of opposite leaf shape

genotypes, or is subject to balancing selection on the leaf shape locus (as has been shown by

Bright and Rausher 2008), and thus has more variation for these reasons.

Collectively, our diverse set of analyses points to a single population being qualitatively

different from the others, with divergence driven by a handful of traits in that population, rather

than range-edge populations having any inherent differences in the structure of **G**. At face value,

these findings might suggest that range-edge populations are not small enough to experience

substantial genetic drift that could reduce genetic variance compared to central populations;

alternatively, it may be that all populations experience frequent extinction and colonization

dynamics, across the range, and that this does not have differential effects on genetic variance.

We suspect the latter possibility is more likely.

Clinal divergence and G

In general, phenotypic divergence is expected to be aligned with **G** under three overall

conditions. First, under genetic drift, we expect that phenotypic divergence in a suite of traits

should be related to the overall amount of standing quantitative genetic variance in those traits

(Lande 1979, Phillips et al. 2001). Second, when **G** matrices are ill-conditioned, or of reduced

rank, we expect evolutionary responses to be biased towards the directions of multivariate trait

space with genetic variance (Chenoweth et al. 2010). In other words, once a **G** matrix differs

from spherical, evolutionary responses will be dominated by the directions containing the most

variance. Finally, when $\beta = g_{max}$, we expect the maximal evolutionary response, and a

concordance between Δz, g_{max}, and β (Gaydos et al. 2013). Hangartner et al. (2019) found that

the **G** matrices of range edge populations of *Drosophila melanogaster* were in fact aligned with

the axis of clinal divergence and that the trait covariances improved the adaptive potential of

peripheral populations. In contrast, phenotypic divergence does not have to be aligned with **G** if

selection has been strong and persistently favoring alternative combinations of traits. For

example, McGuigan et al. (2005), found that the direction of hydrodynamic adaptation in

rainbow fish was only weakly associated with g_{max}, and was primarily aligned with the trailing

eigenvectors of **G**; they interpreted this as a result of long-term selection towards a new

phenotypic optimum.

With this context in mind, many of our results suggest that we should have observed a

relationship between clinal divergence and **G**: the traits are genetically variable, both

individually (Table S1) and as a linear combination (Table 2), and the **G** matrices appear to be

ill-conditioned and of low effective number of dimensions (Table 1). It also seems unlikely that

the phenotypic divergence detected is a result of long-term selection, similar to McGuigan et al.

(2005), as *I. hederacea* currently inhabits (and was collected from) ephemeral, disturbance-prone

habitats such as roadsides, agricultural fields, and cleared areas. Consequently, many of the

conditions that can produce a relationship between **G** and divergence appear to be met.

Two potential explanations for why we do not see such a relationship between **G** and

divergence occur to us. The first is that our previous work (Simonsen and Stinchcombe 2010;

Campitelli and Stinchcombe 2013; Stock et al. 2014) and this study inadvertently omitted an

ecologically important, correlated trait. For this scenario to explain the lack of relationship

between divergence and **G**, there would have to be an omitted trait that is both highly variable

within populations, and highly diverged between populations, such that its inclusion would

predominate both g_{max} and any estimate of a divergence vector. The problem of missing traits is

inherent to studies of phenotypic evolution; while the traits we measured capture aspects of size,

growth, phenology, and floral morphology, it is possible that other traits related to

ecophysiology, seed bank dynamics, or other aspects of the life cycle are more important.

Second, it may be the case that natural selection is acting in a direction that is nearly

orthogonal to g_{max}, thus leading to the divergence that is at a substantial angle from g_{max}. If this

were the case, it would be possible for the **G** matrices we estimated to produce evolutionary

responses that could lead to the observed divergence. One clue in support of this hypothesis is

that flowering time appears to load relatively weakly on g_{max} in the Pennsylvania population, but

in the past has been detected to be under very strong directional selection in this species

(Simonsen and Stinchcombe 2010; Campitelli and Stinchcombe 2013). Similarly, Campitelli and

Stinchcombe (2013) found selection favoring decreased flowering time and increased early and

mid-season growth rates (i.e., β elements of opposite signs), while we found that growth rate and

flowering time load in the same direction (i.e., are positively correlated) of g_{max} in three of four

populations. As such, there are past observations of selection acting strongly on traits with

relatively little genetic variance, or in opposite directions on positively correlated traits, both of

which could lead to β being orthogonal to g_{max}. Indeed, Simonsen and Stinchcombe (2010) found

strong natural selection almost orthogonal to g_{max} in a field study of size traits and flowering

time, due to size traits being highly variable and under weak selection, and flowering time being

less variable but under very strong selection. The divergence observed may thus be the product

of selection, but without requiring recourse to long-term selection towards a single optimum.

While there are field studies of natural selection on some of the traits we studied, an overall

study of all of them is necessary to distinguish between the missing trait explanation and β being

orthogonal to g_{max}.

Conclusions and Future Directions

Determining the relationship between **G** and phenotypic divergence, or divergence of **G**

matrices themselves, is a challenging empirical and statistical endeavor. Practical constraints

make constructing well estimated **G** matrices for dozens of populations nearly impossible

(Arnold et al. 2008), and is certainly the case for *I. hederacea*. The most tractable and testable

hypothesis to emerge from our work is that natural selection on these five traits is poorly aligned

with g_{max}. Future field studies will be required to further elucidate the action of natural selection

on this species, and the ecological mechanisms behind selection for this suite of traits and others.

Acknowledgments

We thank NSERC Canada for funding and a Koffler Scientific Reserve graduate fellowship

(GH) for support. We thank Tia Harrison for field assistance, Brechann McGoey for sharing

relevant R scripts and advice, Jacqueline Sztepanacz for comments on the manuscript, and Aneil

Agrawal, Asher Cutter, Matt Osmond, and Stephen Wright for discussion. Greenhouse support

from William Cole, Tom Gludovacz, Colin Bonner, and Andrew Petrie was invaluable.

References

Agrawal, A. F., and J. R. Stinchcombe. 2009. How much do genetic covariances alter the

rate of adaptation? Proc. Biol. Sci. 276:1183–1191.

Aguirre, J. D., E. Hine, K. McGuigan, and M. W. Blows. 2014. Comparing G:

multivariate analysis of genetic variation in multiple populations. Heredity 112:21–29.

Antonovics, J. 1976. The Nature of Limits to Natural Selection. Ann. Mo. Bot. Gard.

63:224–247. Missouri Botanical Garden Press.

Arnold, S. J., R. Bürger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008.

Understanding the evolution and stability of the G-matrix. Evolution 62:2451–2461.

Basser, P. J., and S. Pajevic. 2007. Spectral decomposition of a 4th-order covariance

tensor: Applications to diffusion tensor MRI. Signal Processing 87:220–236.

Bright-Emlen, K. L. 1998. Geographic variation and natural selection on a leaf shape

polymorphism in the ivyleaf morning glory (*Ipomoea hederacea*). Duke University, Ann Arbor,

United States.

Bright, K. L., and M. D. Rausher. 2008. Natural selection on a leaf-shape polymorphism

in the ivyleaf morning glory (*Ipomoea hederacea*). Evolution 62:1978–1990.

Brown, J. H., D. W. Mehlman, and G. C. Stevens. 1995. Spatial variation in abundance. Ecology

76:2028–2043.

Campitelli, B. E., and J. R. Stinchcombe. 2013. Natural selection maintains a single-locus

leaf shape cline in Ivyleaf morning glory, *Ipomoea hederacea*. Mol. Ecol. 22:552–564.

Campitelli, B. E., and J. R. Stinchcombe. 2014. Population dynamics and evolutionary

history of the weedy vine *Ipomoea hederacea* in North America. G3 4:1407–1416.

678 Chenoweth, S. F., H. D. Rundle, and M. W. Blows. 2008. Genetic constraints and the

679 evolution of display trait sexual dimorphism by natural and sexual selection. Am. Nat. 171:22–

680 34.

681 Chenoweth, S. F., H. D. Rundle, and M. W. Blows. 2010. The contribution of selection

682 and genetic constraints to phenotypic divergence. Am. Nat. 175:186–196.

683 Cheverud, J. M., and G. Marroig. 2007. Comparing covariance matrices: Random

684 skewers method compared to the common principal components model. Genetics and Molecular

685 Biology 30:461–469.

686 Delahaie, B., A. Charmantier, S. Chantepie, D. Garant, M. Porlier, and C. Teplitsky.

687 2017. Conserved G-matrices of morphological and life-history traits among continental and

688 island blue tit populations. Heredity 119:76–87.

689 Endler, J. A. 1977. Geographic Variation, Speciation and Clines. Princeton University

690 Press.

691 Ennos, R. A. 1981. Quantitative studies of the mating system in two sympatric species of

692 *Ipomoea* (Convolvulaceae). Genetica 57:93–98.

693 Falconer, D. S., and T. F. C. Mackay. 1996. Introduction to quantitative genetics. Fourth.

694 Essex, UK: Longman Group.

695 Garcia-Ramos, G., and M. Kirkpatrick. 1997. Genetic models of adaptation and gene

696 flow in peripheral populations.

697 Gaydos, T. L., N. E. Heckman, M. Kirkpatrick, J. R. Stinchcombe, J. Schmitt, J.

698 Kingsolver, and J. S. Marron. 2013. Visualizing genetic constraints. Ann. Appl. Stat. 7:860–882.

699 Guillaume, F., and M. C. Whitlock. 2007. Effects of migration on the genetic covariance

700 matrix. Evolution 61:2398–2409.

Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. J. Stat. Softw. 33:1–22.

Hangartner, S., Lasne, C., Sgrò C.M., Connallon T., and K. Monro. 2019. Genetic

covariances promote climatic adaptation in Australian *Drosophila*. Evolution 74: 326-337.

Hansen, T. F., and D. Houle. 2008. Measuring and comparing evolvability and constraint

in multivariate characters. J. Evol. Biol. 21:1201–1219.

Hine, E., S. F. Chenoweth, H. D. Rundle, and M. W. Blows. 2009. Characterizing the

evolution of genetic variance using genetic covariance tensors. Philos. Trans. R. Soc. Lond. B

Biol. Sci. 364:1567–1578.

Hull-Sanders, H. M., M. D. Eubanks, and D. E. Carr. 2005. Inbreeding depression and

selfing rate of *Ipomoea hederacea* var. *integriuscula* (Convolvulaceae). Am. J. Bot. 92:1871–

1877.

Jones, A. G., S. J. Arnold, and R. Bürger. 2004. Evolution and stability of the G-matrix

on a landscape with a moving optimum. Evolution 58:1639–1654.

Kingsolver, J. G., N. Heckman, J. Zhang, P. A. Carter, J. L. Knies, J. R. Stinchcombe,

and K. Meyer. 2015. Genetic variation, simplicity, and evolutionary constraints for function-

valued traits. Am. Nat. 185:E166–81.

Kirkpatrick, M. 2009. Patterns of quantitative genetic variation in multiple dimensions.

Genetica 136:271–284.

Kirkpatrick, M., and N. H. Barton. 1997. Evolution of a species' range. Am. Nat. 150:1–

23.

Klingaman, T. E., and L. R. Oliver. 1996. Existence of ecotypes among populations of

entireleaf morningglory (*Ipomoea hederacea* var. *integriuscula*). Weed Sci. 44:540–544.

Krzanowski, W. J. 1979. Between-groups comparison of principal components. J. Am.

Stat. Assoc. 74:703–707.

Lande, R. 1992. Neutral theory of quantitative genetic variance in an island model with

local extinction and colonization. Evolution 46:381–389.

Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain:

body size allometry. Evolution 33:402–416.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.

Evolution 37:1210–1226.

Lin, C. Y., and F. R. Allaire. 1977. Heritability of a linear combination of traits. Theor.

Appl. Genet. 51:1–3.

Mayr, E. 1963. Animal Species and Evolution. Harvard University Press.

Mayr, E. 1954. Change of genetic environment and evolution. George Allen & Unwin

Limited.

McGoey, B. V., and J. R. Stinchcombe. 2021. Introduced populations of ragweed show

as much evolutionary potential as native populations. Evol. Appl. 14:1436–1449.

McGuigan, K., S. F. Chenoweth, and M. W. Blows. 2005. Phenotypic divergence along

lines of genetic variance. Am. Nat. 165:32–43.

Morrissey, M. B., S. Hangartner, and K. Monro. 2019. A note on simulating null

distributions for G matrix comparisons. Evolution 73:2512–2517.

Peterson, R. A., and J. E. Cavanaugh. 2020. Ordered quantile normalization: a

semiparametric transformation built for the cross-validation era. J. Appl. Stat. 47:2312–2327.

Phillips, P. C., M. C. Whitlock, and K. Fowler. 2001. Inbreeding changes the shape of the

genetic covariance matrix in *Drosophila melanogaste*r. Genetics 158:1137–1145.

Puentes, A., G. Granath, and J. Ågren. 2016. Similarity in G matrix structure among

natural populations of *Arabidopsis lyrata*. Evolution 70:2370–2386.

Roff, D. 2000. The evolution of the G matrix: selection or drift? Heredity 84:135–142.

Roughgarden, J. 1979. Theory of population genetics and evolutionary ecology : an

introduction. Macmillan, New York.

Sexton, J. P., P. J. McIntyre, A. L. Angert, and K. J. Rice. 2009. Evolution and ecology

of species range limits. Annu. Rev. Ecol. Evol. Syst. 40:415–436. Annual Reviews.

Simonsen, A. K., and J. R. Stinchcombe. 2010. Quantifying evolutionary genetic

constraints in the ivyleaf morning glory, *ipomoea hederacea*. Int. J. Plant Sci. 171:972–986.

Stock, A. J., B. E. Campitelli, and J. R. Stinchcombe. 2014. Quantitative genetic variance

and multivariate clines in the Ivyleaf morning glory, *Ipomoea hederacea*. Philos. Trans. R. Soc.

Lond. B Biol. Sci. 369:20130259.

Sztepanacz, J. L., and M. W. Blows. 2017. Accounting for Sampling Error in Genetic

Eigenvalues Using Random Matrix Theory. Genetics 206:1271–1284.

Teplitsky, C., M. R. Robinson, and J. Merilä. 2014. Evolutionary potential and

constraints in wild populations. Quantitative genetics in the wild 190–208. Oxford University

Press Oxford, UK.

Walsh, B., and M. W. Blows. 2009. Abundant genetic variation + strong selection =

multivariate genetic constraints: a geometric view of adaptation. Annu. Rev. Ecol. Evol. Syst.

40:41–59.

Walter, G. M., J. David Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018. Evolution

of genetic variance during adaptive radiation. Am. Nat. 191:E108–E128.

Whitlock, M. C., and D. E. McCauley. 1990. Some population genetic consequences of

colony formation and extinction: genetic correlations within founding groups. Evolution

44:1717–1724.

Wood, C. W., and E. D. Brodie 3rd. 2015. Environmental effects on the structure of the

G-matrix. Evolution 69:2927–2940.