Research article

Flexible estimation of biodiversity with short-range multispectral imaging

in a temperate grassland

Jackson, J.[1], Lawson, C. S.[2], Adelmant, C.[1], Huhtala, E.[1], Fernandes, P.[1], Hodgson, R.[1], King, H.[1],

Williamson, L., Maseyk, K.[2], Hawes, N.[3], Hector, A.[1], & Salguero-Gómez, R[1,4]

1. Department of Biology, University of Oxford, UK, OX1 3SZ

2. School of Environment, Earth & Ecosystem Sciences, The Open University, UK, MK7 6AA

3. Oxford Robotics Institute, Department of Engineering Science, University of Oxford, UK, OX1 3PJ

4. Max Planck Institute for Demographic Research, Rostock, Germany, DE 18057

Corresponding author: John Jackson, john.jackson@zoo.ox.ac.uk, 0000-0002-4563-2840

Declaration of Competing Interest

We declare that there are no competing interests.

Data Availability Statement

We intend to submit all data to the dryad repository on acceptance, and archive code used in analyses

using Zenodo.

Abstract

1. Image sensing technologies are rapidly increasing the cost-effectiveness of biodiversity monitoring efforts. Species differences in the reflectance of electromagnetic radiation have recently been highlighted as a promising target to estimate plant biodiversity using multispectral image data.

2. However, these efforts are currently hampered by logistical difficulties in broad-scale implementation and their use in characterizing biodiversity at different spatial scales.

3. Here, we investigate the utility of multispectral imaging technology from commercially available unmanned aerial vehicles (UAVs, or drones) in estimating biodiversity metrics at short-range (<10 m image recording height) in a temperate calcareous grassland ecosystem in Oxfordshire, UK. We calculate a suite of moments (coefficient of variation, standard deviation, skew, kurtosis) for the distribution of radiance from multispectral images at five wavelength bands (Blue 450±16 nm; Green 560±16 nm; Red 650±16 nm; Red Edge 730±16 nm; Near Infrared 840±16 nm) and test their effectiveness at estimating ground-truthed biodiversity metrics from *in-situ* botanical surveys for 37 - 1 m × 1 m quadrats.

4. We find positive associations between the average coefficient of variation in spectral radiance and both the Shannon-Weiner and Simpsons biodiversity indices. Furthermore, we find that the average coefficient of variation in spectral radiance is consistent and highly repeatable, across sampling days and recording heights. Positive associations with biodiversity indices hold irrespective of the image recording height (2-8 m), but we report reductions in estimates of spectral diversity with increases to UAV recording height. UAV imaging reduced sampling time by 16-fold relative to *in-situ* botanical surveys.

5. *Synthesis* - We demonstrate the utility of multispectral radiance moments as an indicator of grassland biodiversity metrics at high spatial resolution using a widely available UAV monitoring system at a coarse spectral resolution. The use of UAV technology with multispectral sensors has far-reaching potential to provide cost-effective and high-resolution monitoring of biodiversity in complex environments.

Keywords: Autonomous monitoring, Biodiversity Drone, Remote sensing, Unmanned Aerial Vehicle (UAV)

1. Introduction

With over 1 million species expected to go extinct by 2100, cost-effectively monitoring biodiversity is a critical task in the Anthropocene (Díaz et al., 2019). Image sensing technologies, which can be used to monitor biological systems through the measurement of reflected and emitted radiation, have emerged as a critical tool that can increase this cost-effectiveness (Turner, 2014). The characterisation of floral biodiversity with remote sensing, particularly with satellite imagery, is well-established in biodiversity research (Pettorelli et al., 2005). Multiple efforts have been made towards using remote sensing data, particularly at large spatial scales and in forest ecosystems, to estimate plant diversity (e.g., Frye et al., 2021; Jetz et al., 2016; Tuanmu & Jetz, 2015; Turner et al., 2003). However, there are limitations in the use of long-range remote sensing, including low spatial resolution that does not necessarily highlight biodiversity at small spatial scales (Mairota et al., 2015), high sensor costs (*e.g.,* $98,700, Headwall Photonics, 2022) and monitoring costs (*e.g.,* $60,000, Jet Propulsion Laboratory, 2022), and reliance on publicly available satellite data (e.g., The European Space Agency, 2022). Flexible application of remote sensing concepts and technology at a wide range of spatial scales, in complex changing environments, and with increased cost-effectiveness, will provide vital resources for monitoring biodiversity (Turner, 2014).

Reflectance of electromagnetic (EM) radiation beyond the visible range (380-700 nm) has recently been shown accurate proxy for biodiversity (Cavender-Bares et al., 2020; Wang & Gamon, 2019). The general concept of 'spectral diversity' is founded on the principle that, due to differences in functional form (both growth form and pigmentation), plant species have differential reflectance signals across the electromagnetic (EM) spectrum (Gamon et al., 1997). Thus, for a multispectral image, the diversity of spectral reflectance can be a proxy for the number of different plant species, or species diversity (Gholizadeh et al., 2019; Laliberté et al., 2020). The spectral diversity concept was recently applied in the hyper-diverse Cape Floristic Region, where destructively sampled leaf reflectance spectra were used to obtain a robust proxy ($R^2 > 0.9$) of species diversity across 1,267 - 10 m $\times$ 5 m quadrats

(Frye et al., 2021). Therefore, integrating sensing data at a range of spatial scales (Laliberté et al., 2020;

Turner, 2014) and the use of spectral surrogates for biodiversity (Frye et al., 2021) can rapidly improve

biodiversity monitoring.

Recently, there have been several applications of spectral diversity from close range imaging

data (Gholizadeh et al., 2019; Lopatin et al., 2017). In prairie grassland ecosystems, close associations

have been found between species diversity and spectral diversity, captured using aircraft-mounted

hyperspectral sensors and images at a spatial resolution of 1 m × 1 m (pixel resolution) (Gholizadeh et

al., 2018, 2019, 2020). Gholizadeh et al. (2019) primarily use the average coefficient of variation across

pixels and spectral bands as the metric of spectral diversity, which we also adopt here. Furthermore, at

a fine resolution of < 0.5 cm × 0.5 cm, static monitoring of grassland plots has been used to estimate

not only biodiversity metrics (Imran et al., 2021; Villoslada et al., 2020), but to reconstruct species

percentage cover and extract detailed features of community dynamics (Lopatin et al., 2017). However,

a key limitation of these close-range imaging approaches is their reliance on expensive hyperspectral

sensors (> $50,000 sensors; Gholizadeh et al., 2019; Imran et al., 2021; Lopatin et al., 2017) and

monitoring ($1,200 per hour using the CALMIT aerial sensor from Gholizadeh et al., 2019).

Overcoming these cost limitations will facilitate further use of spectral imaging in biodiversity research.

Despite advances in image sensing, there is a need for monitoring systems that i) are deployable

in a diverse range of habitats, ii) can span a range of spatial resolutions, iii) are cost-effective, and iv)

user friendly. One potential solution is the use of commercial unmanned aerial vehicles (UAVs), or

drones, which have rapidly increased in popularity over the last decade (Colomina & Molina, 2014).

Here, we investigate the efficacy of multispectral imaging UAV technology in the estimation of

biodiversity in a temperate calcareous grassland. We used a commercially available UAV system with

a five-band multispectral band sensor (Blue 450±16 nm; Green 560±16 nm; Red 650±16 nm; Red Edge

730±16 nm; Near Infrared 840±16 nm) to image 37 - 1 m × 1 m quadrats that were also characterised

using *in situ* biodiversity assessments from botanical surveys. Then, by extracting moments of spectral

diversity from multispectral image data, we estimate spectral diversity and explored associations

between spectral diversity and biodiversity. Our goal is to provide a framework for cost-effective

biodiversity monitoring in complex environments using readily and commercially available technologies.

2. Materials and Methods

2.1 Study site and in-situ biodiversity data

Data collection took place at the five-acre section of the Upper Seeds field site (51°46'16.8"N 1°19'59.1"W) in Wytham woods, Oxfordshire, UK between 16th June-14th July 2021, the peak of the growing season. The Upper Seeds site is a recovering and managed calcareous grassland, which was used for agriculture in the 1950s, before encroaching scrub vegetation was removed and the site was managed as a grassland beginning in 1978 (Gibson, 1986). Management on upper seeds is implemented with mowing of the site in mid-July at the peak of the growing season, and again in early October, coinciding with the end of the growing season. The site has a low average soil depth (300-500 mm), generally alkaline soils (Gibson & Brown, 1991), a daily average temperature range of -5 °C to 26 °C (2016-2020), a daily total precipitation range of 0-40 mm (2016-2020), and high general biodiversity, in which graminoids are the dominant functional group (59.1% by biomass). A total of 37 1 m × 1 m experimental quadrats were used in the current study, which display a large degree of variation in species composition and biomass. There were between 16 and 33 vascular plant species per m^2, with a mean richness of 25.77 species and a median richness of 26 species. Total above-ground dry biomass across quadrats varied between 166.8 g/m^2 and 931.5 g/m^2, with a mean of 397.9 g/m^2 and median of 327.2 g/m^2.

We explored biodiversity and spectral diversity associations in the context of two long-term experiments that aim to explore the response of grasslands to environmental change (full site map in Fig. S1). These experiments are the Disturbance and Resources Across Global Grasslands (DRAGNet, n = 20 plots) coordinated research network (https://nutnet.org/dragnet) and the global drought network (DroughtNet, n = 17 plots) coordinated research network (https://drought-net.colostate.edu/). All DRAGNet plots (5 m × 5 m plots) were ambient controls, with no experimental treatments applied prior to the collection of the data reported here. Each 5 m × 5 m plot from DroughtNet was one of four

experimental treatments: ambient control plots (n = 5), -50% rainfall shelters to simulate drought (n =

5), +50% irrigated plots to simulate increased rainfall (n = 5), and procedural controls (rainfall shelter

with no change to rainfall) (n = 2; three plots were inaccessible for the UAV as the rainfall shelters were

fixed). For analyses, ambient control treatments (n = 25) across the two research networks were pooled

as we did not observe substantial differences between biodiversity metrics (Fig. S2). To account for

replicated observations of the same quadrats and estimate the consistency of spectral diversity measures,

we explored quadrat-level variance using intercept-only random effects for the quadrat ID.

To estimate the efficacy of multispectral sensors in predicting biodiversity, we collected data

from two sources, *in-situ* biodiversity assessments and UAV derived multi-spectral image data (Fig. 1).

For the *in-situ* assessments, we quantified biodiversity metrics using species-level percentage cover and

dry above-ground biomass data. We estimated percentage cover data for all vascular plant species in a

plot using a 1 m × 1 m gridded quadrat (10 cm grid), focusing on four broad functional groups:

graminoids, legumes, forbs, and woody species (Fig. 1). Because species overlapped spatially,

percentage cover estimates could exceed 100%. Using relative proportions, p, calculated from

percentage cover estimates, we calculated three biodiversity metrics: i) vascular plant species richness,

ii) the Shannon-Weiner diversity index, H (equation 1; Shannon & Weaver, 1963), and iii) the

Simpson's diversity index, D (equation 2; Simpson, 1949):

$$H = -\sum p \ln p, \qquad \text{Eqn. 1}$$

$$D = \sum p^2 \qquad \text{Eqn. 2}$$

We estimated above ground biomass after UAV image/percentage cover data collection, using

a clip strip of all vascular plant material in 1 m × 0.2 m (DRAGNet; collected from standardised

locations in the plot) or 1 m × 0.25 m (DroughtNet; collected from the centre of each quadrat). Clip

strips were gathered using hand trimmers at a height of 1-2cm above the soil surface. Within one day

of collection, we sorted clip strips in to five functional groups: graminoids, legumes, forbs, woody

species, and bryophytes (not included in species-level percentage cover estimates) and dried them at

70°C for 48hr, before weighing the dry biomass at an accuracy of ±0.1g. The estimates of biomass were

scaled to g/m^2 for analyses.

2.2 UAV image data collection

To obtain spectral diversity data, we collected image data using manual flights of the DJI Phantom 4 multispectral UAV (https://www.dji.com/p4-multispectral). The sensor payload of the DJI Phantom 4 multispectral consists of six 4.96 mm × 3.72 mm complementary metal–oxide–semiconductor (CMOS) sensors: one RGB sensor for visible range colour images, and five monochrome sensors for multispectral imaging. The five multispectral sensors are sensitive at the following electromagnetic wavelengths: Blue - 450 nm ± 16 nm, Green - 560 nm ± 16 nm, Red - 650 nm ± 16 nm, Red edge - 730 nm ± 16 nm and Near-infrared - 840 nm ± 26 nm (Fig. 1). Each sensor has an effective resolution of 2.08 MP. All six image sensors are triggered simultaneously when capturing data, with negligible (but non-0) positional differences between sensors. A dorsal spectral sunlight sensor on the P4 multispectral sensor provides image exposure compensation of multispectral image data.

Fig. 1. Schematic for assessing the efficacy of spectral distribution moments for capturing biodiversity in a temperate grassland. For each $1m \times 1m$

observation quadrat, we collected both UAV image data (top) and in-situ biodiversity data (bottom). In-situ biodiversity data were collected by botanical surveys

for vascular plant percentage cover across the quadrat (from which richness, Shannon-Weiner and Simpson's indices were calculated) and using dry above-

ground biomass for clip strips (area determined by the coordinated research networks DRAGNet and DroughtNet; see 2.1), after UAV images were taken. UAV

images were collected for each plot at four recording heights (2 m, 4 m, 6 m, and 8 m) across five multispectral bands for which at-sensor radiance Digital

Number (DN) value distributions were summarised using four moments. Finally, *in situ* biodiversity data were combined with spectral distribution moments to

examine their potential relationships using a Bayesian linear regression framework.

To obtain spectral diversity metrics, we collected multispectral images for each quadrat over

several flights across the sampling period, capturing quadrat-level variability with weather/light

conditions. However, to minimise visual interference (from rain or low sun), all images were taken

during dry weather and between 10:30-15:30 (BST). The corners of each DRAGNet quadrat were

marked with flags (Fig. S1b). For DroughtNet, the quadrat was approximated using the outer edges of

the 5 m × 5 m plot. All images were collected facing the western edge of each plot. We collected images

at increasing approximate image recording heights of 2 m, 4 m, 6 m, and 8 m above the ground to

capture changes in image resolution and consequences for estimating biodiversity (Fig. 1). Flying height

is recorded relative to the UAV's take-off location, and although the topographical variation at the site

is < 5 m, image record heights were approximated using structures of known height (*i.e.*, rainfall

shelters, see Fig. S5). A total of 1,878 individual images were collected for the 37 quadrats over seven

sampling days.

2.3 Image processing

To extract spectral reflectance metrics from the raw image data, we standardised raw images across

quadrats for each sample. The raw images encompassed the full field of view of the sensors, and we

first batch-cropped images with Adobe Lightroom v. 5 (Adobe, 2021) to include only data for the

desired 1 m^2 quadrats. Images were exported as .tif files maintaining at-sensor radiance values with

minimal post-processing. We did not perform post-processing to account for interference from other

sources solar radiation (e.g. Gholizadeh et al., 2019) due to the low recording height of UAV flights in

the current study. Instead we estimated quadrat-level variability in spectral signals across the study

period. The data stored in each .tif image were at-sensor radiance Digital Number (DN) values, which

were used in subsequent analyses. The use of raw DN values was appropriate for the purpose of this

study because we aimed to capture moments of the distribution of DN values in each image. Scaling

DN values to normalised reflectance values was not necessary to explore the radiance distributions

because relative differences between DN values, and thus distributional moments remain unaltered.

Furthermore, the omission of calibration to reflectance values, which is typically performed with

standardised reflectance plates, was more appropriate for the application of the current protocol and

aims *i.e.* deployment in complex environments. Multispectral .tif images were treated as rasters for

further image processing, and all subsequent analysis was carried out using R version 4.0.5 (R Core

Team, 2021).

We calculated moments of spectral radiance for each image using the *raster* package (Hijmans,

2020). Following Gholizadeh et al. (2019), we calculated the coefficient of variation, standard

deviation, skewness, and kurtosis across raster pixels to capture the shape of the at-sensor radiance DN

distribution (Fig. 1). We averaged moment values of radiance DNs across all multispectral bands for a

single observation (a given quadrat at a given recording height in each sampling event) to calculate

overall distributional moments. Thus, here we define the spectral coefficient of variation as the mean

coefficient of variation of the spectral radiance across raster pixels and multispectral bands for a single

image. Observations were discarded if the image recording height was >8 m and replicate images were

not obtained for all quadrats at all image recording heights. Therefore, the final sample size for the

averaged spectral moment data was 193. In addition, to identify the spectral bands that were most

sensitive to biodiversity metrics, we also tested band-level associations, where radiance distributions

were not averaged across spectral bands for the same raw data, and in this case spectral radiance

distributions of each band were related to biodiversity indices.

2.4 Statistical analyses

We explored the efficacy of spectral radiance distribution moments in describing *in-situ* biodiversity

indices using a Bayesian hierarchical linear regression model selection framework using the *brms*

package (Bürkner, 2017; Fig. 1). All variables were z-scored (mean and variance centered on 0) for

analysis to meet the distributional assumptions of linear regressions. The key response variable was the

spectral coefficient of variation (Gholizadeh et al., 2019), and the key predictor variables were the *in-*

situ biodiversity indices. However, we also tested other spectral moment-biodiversity associations,

namely, the skewness of spectral radiance and biomass.

We then estimated the out-of-sample predictive performance of models including biodiversity

indices relative to base models. For each explored pair-wise combination of spectral distribution

moment and biodiversity indices, we performed leave-one-out cross validation with the *loo* criterion

and the expected log-wise predictive density (*elpd*, where $\Delta elpd$ gives the change in *elpd* relative to

another explanatory model) (Vehtari et al., 2017). Base models did not include any predictor variables,

including only an intercept-only random effect for quadrat. We also investigated the performance of

models including image recording height, and two-way interaction terms between biodiversity indices

and height to explore how image resolution change influenced the efficacy of spectral diversity indices.

In addition to models on averaged spectral moments, we also used band-level moments to

investigate the relationship between biodiversity indices and the spectral coefficient of variation for

each individual EM band. We included univariate and two-way interaction terms between the EM band

and biodiversity indices variables. Finally, because a small number of quadrats used in the current study

were also exposed to long-term drought/irrigation/control treatments, we explored whether there were

differences in average spectral moments between ambient (n = 25 plots), control (n = 2), irrigated (n =

5) and drought (n = 5) using a categorical predictor for treatment.

To account for repeated observations from the same quadrat at different heights or across

sampling events, all models included an intercept-only random effect for the quadrat $\sigma_{quadrat}$. From

this random effect, we also estimated the intraclass correlation (ICC) or repeatability (R). This term

indicates the proportion of quadrat-level variance $\sigma_{quadrat}$ with respect to the population-level variance

σ (Nakagawa & Schielzeth, 2010). We used this estimate of repeatability enabled us to assess the

consistency of spectral radiance distributions across observations of the same quadrat.

In analyses with average spectral distribution moments, we used weakly informed normal priors

for the population-level intercept and coefficient terms of $N(0,1)$. The $\sigma_{quadrat}$ term was fit using an

exponential prior with a rate of two. For band-level analyses (with a greater number of parameters),

models were fit using $N(0,0.7)$ intercept/coefficient priors and exponential $\sigma_{quadrat}$ priors with a rate

of four. Models were run across four serial chains for 2,000 iterations with 1,000 warmup iterations,

and the model's convergence across chains was assessed by inspecting $\hat{R}$ values (Bürkner, 2017).

3. Results

We found consistent positive associations between the average coefficient of variation in spectral radiance and biodiversity, namely, the Shannon-Weiner and Simpson's indices (Fig. 2). The model including the Shannon-Weiner index and image recording height as univariate terms outperformed the base model, with $\Delta elpd = 123.6$ (Table S1). Increases in the Shannon-Weiner index were associated with increases in the average spectral coefficient of variation ($\beta_{shannon} = 0.19$ [-0.04, 0.43], 95% credible intervals; Fig. 2a). Furthermore, as expected, there was a strong negative association between image recording height and the average spectral coefficient of variation ($\beta_{height} = -0.28$ [-0.31, -0.26]; Fig. 2a). This negative association suggests that the resolution of spectral diversity decreases rapidly with recording height at this spatial resolution (~ 30 % decrease in scaled spectral variation per 1 m height increase). The positive association with the spectral coefficient of variation was stronger for the Simpson's biodiversity index (Fig. 2b). Although the full model including a two-way interaction between recording height and Simpson's index was the best predictive model, we selected the model including only univariate effects ($\Delta elpd = 125.0$), because of a lack of a clear interaction effect (Table S2). Here, a similar patten with image recording height was accompanied by a stronger positive association between Simpson's index and spectral coefficient of variation ($\beta_{simpsons} = 0.33$ [0.12, 0.54]; Fig. 2b).

Generally, we did not observe associations between the skewness and kurtosis in spectral radiance distributions and biodiversity indices (Fig. S3). Furthermore, there was no clear evidence for a relationship between total above-ground biomass and any of the spectral distribution moments (Fig. S3). Specifically, although there was increased model predictive performance from models including biomass, there was no clear relationship between the skewness of spectral radiance distribution and biomass ($\beta_{biomass} = 0.08$ [-0.19, 0.35]; Fig. S3; Table S5)

In addition to overall effects, in band-level analyses where raw data were not averaged across bands, there was evidence for an interaction effect between the spectral band and both the Shannon-Weiner ($\Delta elpd = 772.0$) and Simpson's indices ($\Delta elpd = 771.5$) (Table S3 & S4, respectively). Generally, the green (560±16 nm) and red (650±16 nm) spectral bands displayed higher variability in

the coefficient of variation across quadrats, and stronger associations with the Shannon-Weiner and Simpson's indices (Fig. 3). The Red Edge (730±16 nm) and Near Infrared (840±16 nm) bands exhibited weaker associations with biodiversity indices (Fig. 3).

When assessing the influence of treatment on spectral radiance, we also observed reductions in the average spectral coefficient of variation in both drought and procedural control quadrats in comparison to ambient or irrigated treatments (Fig. S4). However, given the congruence of procedural control and drought treatments in DroughtNet, both of which are characterised by metal rainfall shelters, we conclude that the reduction in spectral coefficient of variation in drought and procedural treatments is likely a result of structural interference from the rain shelter structures (Fig. S5).

Finally, we tested the consistency of the spectral coefficient of variation across observation days and heights for each quadrat in the best predictive Shannon-Weiner and Simpson's index models (Table S1 & S2, respectively). The average coefficient of variation was highly consistent for each quadrat when images at different heights or across sampling events were compared (Fig. S6). Both the Shannon-Weiner and Simpson's models with the average coefficient of variation exhibited quadrat-level variance that exceeded the population-level variance and a high degree of repeatability (Fig. S5; 0.76 [0.65, 0.85] and 0.72 [0.60, 0.82], respectively).

Fig. 2. Consistent positive associations between biodiversity indices and the average spectral coefficient of variation. The positive association between (a) Shannon-Weiner biodiversity index and (b) Simpsons index, and the average spectral coefficient of variation (averaged across five spectral bands) for different image recording heights: 2 m, 4 m, 6 m, and 8 m (panels). Points are observations from a single quadrat at a given height. Both biodiversity indices and spectral coefficient of variation are z-scored. Lines represented the posterior prediction mean over 4,000 simulations averaged over all quadrats, with the 90% credible intervals. Insets showcase density distributions of the posterior estimates for the image recording height (β_{height}) and biodiversity indices ($\beta_{shannon}$ and β_{height}) .

Fig. 3. Green and Red spectral bands are the most sensitive to biodiversity indices. Posterior

predictions for the band-level spectral coefficient of variation with the Shannon-Weiner (a) and

Simpson's (b) biodiversity indices at an image recording height of 2 m. Here, raw spectral moments

were not averaged across spectral bands (as in Fig. 2). Points are raw observations from a single plot,

and the colour denotes the spectral band (B = Blue, G = Green, R = Red, RE = Red Edge, NIR = Near

Infrared). Both biodiversity indices and spectral coefficient of variation are z-scored. Lines are the

posterior prediction mean over 4,000 simulations averaged over plots, with the 90% credible intervals.

4. Discussion

Despite rapid technological advancements in image sensing over the last four decades, biodiversity monitoring is not currently able to track the full extent of human impacts on the biosphere (Wang & Gamon, 2019; Wilson, 2017). We urgently need more cost-effective and widely available systems to monitor detailed changes in biodiversity (Turner, 2014; Turner et al., 2003). Here, using a commercially available short-range Unmanned Aerial Vehicle (UAV, drone), we find a consistent, repeatable association between variation in spectral radiance and species diversity in a biodiverse temperate grassland. The coefficient of variation in spectral radiance was positively associated with the Shannon-Weiner and Simpsons indices, and in particular the green and red bands of the electromagnetic (EM) spectrum were most indicative of grassland biodiversity. Our results build on extensive work in grassland ecosystems exploring the use of spectral diversity as a surrogate for biodiversity (Frye et al., 2021; Gholizadeh et al., 2019; Villoslada et al., 2020) and species composition (Lopatin et al., 2017). However, our research in a diverse temperate grassland community contrast with previous findings that highlighted limitations to the characterisation of biodiversity using spectral imaging in species-rich environments (Imran et al., 2021). We highlight the importance of close-range remote sensing for biodiversity monitoring (Turner, 2014; Turner et al., 2003; Wang & Gamon, 2019). Crucially, we demonstrate the feasibility of the spectral diversity concept using a commercially available UAV with a low spectral resolution sensor, which has far-reaching potential as a tool to explore biodiversity change at high spatio-temporal resolution and in complex environments.

The key advantage of using commercially available UAV technology is its cost-effectiveness relative to reliance on *in-situ* monitoring or the use of long-range or high spectral resolution sensors. In the current study, where we collected data from 37 quadrats at four different image recording heights, the total flight time was 134 min. If we reasonably allocate one researcher 60 min to do a full botanical survey (species percentage cover and biomass clip – ignoring biomass processing of 30 min per sample), the full *in-situ* sampling time is 37 hr. Thus, remote monitoring would have saved on sampling time by a factor of 16. In the current study, the reduction in sampling time is of course limited to proxies of broad biodiversity metrics. However, with increases to sensor resolution and decreasing costs, reconstructing species-level biodiversity data with flexible remote monitoring may also be possible

(Lopatin et al., 2017). Increases in cost-effectiveness may also be increased by automated flight paths over survey locations, for which unsupervised spectral reflectance data could be collected. Furthermore, while rapid advancements have been made on spectral diversity, previous studies have utilised high-resolution multi/hyperspectral sensors that are either immobile, high-cost, or both (Frye et al., 2021; Gholizadeh et al., 2019; Imran et al., 2021; Lopatin et al., 2017). The current complete monitoring system is available for purchase for < 10,000 USD, relative to >50,000 USD for many hyperspectral sensors. Moreover, because the most sensitive spectral radiance moments for biodiversity indices were in the green and red bands, which lie within the visible range, we provide evidence that spectral radiance from visible-only imaging may be sufficient to act as a proxy for biodiversity indices. The present method provides a low-cost and mobile solution to rapidly characterise biodiversity in complex environments.

The rapid increase in the public use of drone technology provides an opportunity for a rapid expansion of detailed biodiversity monitoring at flexible spatio-temporal scales (Colomina & Molina, 2014). For example, drone technology with multispectral imaging sensors has recently been applied to characterise fungal disease in lemon myrtle trees (Heim et al., 2019) and macroalgal community structure in intertidal habitats (Tait et al., 2019), when combined with classification algorithms to disentangle biological communities. These recent findings suggest that drone technology may be applicable in many ecosystems, when combined with ground-truthed biodiversity data. However, biodiversity metrics are not the only ecological indicators, and functional traits are also widely used as ecological indicators of environmental change (e.g., Bjorkman et al., 2018). Because spectral reflectance is related to functional form (Gamon et al., 1997), the applicability of sensing technology is not limited to biodiversity, and can also act as a proxy for functional diversity (Frye et al., 2021). In the current study site, incorporating UAV monitoring to long-term experimental manipulations will enable the investigation of how environmental disturbances such as drought and nutrient addition influence biodiversity, functional diversity and community structure.

Another advantage of the current study is our use of raw digital radiance values and moments of spectral radiance distributions as opposed to ground-calibrated reflectance values that are typically applied to long-range multispectral imaging (Schläpfer et al., 2020). We therefore call for the

application of the current approach across ecosystems. The public interest in drone technology may also

assist in the development of crowd sourcing or citizen science programs to provide high-resolution

biodiversity monitoring globally. Furthermore, the utility of this approach to gather large amounts of

image data will increase when combined with machine learning algorithms to identify single species or

high-resolution community dynamics. Currently, machine learning has been applied to agricultural

imaging challenges (Heim et al., 2019) and in static species-cover assessments (Lopatin et al., 2017).

Therefore, there is now an opportunity to combine this technology with large UAV-derived datasets.

Ultimately, capitalising on drone technology has the potential to revolutionise biodiversity research.

Two key limitations to the current approach that may influence its application in other

environments is the spatial and temporal resolution of data. Integrating spatial scales has long been a

central issue in remote sensing applications (Turner, 2014). We found a negative association between

spectral diversity and recording height. With increases in recording height there are decreases in pixel

resolution, which are then unable to discern micro-structures from plant species that contribute to

spectral reflectance variation. While this negative relationship was insufficient to influence species-

diversity relationships in the current study below 10 m in recording height, this finding suggests that at

greater recording heights associations with high-resolution measures of species diversity may

breakdown. Previous studies have either focussed on short-range/destructive sampling (Frye et al.,

2021; Lopatin et al., 2017) or long-range multispectral imaging (Gholizadeh et al., 2019), and therefore

there is a need to explore species-spectral diversity associations across spatial scales. Recently, novel

dissimilarity approaches have been applied to satellite imaging data at a range of spatial scales (Rossi

et al., 2021). Integrating these approaches using UAV technology in novel habitats with alternate plant

communities will be crucial for future research.

Temporal resolution is also a key factor in need of consideration when assessing spectral-

species diversity associations. The data used in the current study represent a 'static' measure of

biodiversity at the peak of the growing season in a temperate grassland. However, grassland

communities exhibit a high degree of temporal variability, particularly in response to environmental

drivers (Harrison et al., 2015; Thorhallsdottir, 1990). When implementing unmanned biodiversity

monitoring, this temporal component may hinder the accuracy of spectral diversity measures. A solution

developed by Rossi et al., (2021) applies a novel dissimilarity index between pairs of spectral images

over the same region to disentangle temporal components of community change linked to management

and phenology. However, this approach has yet to be applied to images collected using drone

technology in varying habitats.

5. Conclusions

Taking advantage of technological advancements in unmanned sensing will greatly improve the cost-

effectiveness of biodiversity monitoring. UAVs have the potential to span spatial scales, access

challenging environments, and provide high resolution data on the impact of environmental change on

ecosystems. Our study adds to a growing body of literature highlighting links between spectral and

species diversity. Integrating these patterns at varying spatio-temporal scales and in novel habitats will

provide vital insights to aid in documenting changes in the biosphere.

Credit author statement

The study was designed by JJ, RS-G, AH and NH. Core DroughtNet biodiversity data were collected

and experimentation designed by CSL, AH, KM. DRAGNet biodiversity data and image data were

collected by JJ, CA, EH, RH, LW, PF, and RS-G, with support from CSL. JJ analysed the data and

wrote the manuscript, with initial feedback from RS-G. Further manuscript feedback was provided by

CSL, KM, CA, LW and AH. All authors approved the manuscript for publication.

Acknowledgements

We thank the many researchers making use of the RainDrop Wytham Woods site in summer 2021, who

contributed vital discussions on botany and field logistics. We thank the Environmental Change

Network, and in particular S. Schäfer and D. Pallett from the Centre for Ecology & Hydrology for

access to the local weather data. Thanks to S. Middleton and D. Encarnation for support in botanical

surveys. Special thanks to N. Fisher, N. Havercroft, and K. Crawford for field logistic support. JJ was

funded by the Amazon Web Service Test Bed Funding scheme "Monitoring and Predicting Biodiversity

Resilience through AI & Robotics" to RS-G and NH. RS-G was funded by a NERC IRF

NE/M018458/1. NH was funded by EPSRC Programme Grant "From Sensing to Collaboration"

(EP/V000748/1). AH was supported by the John Fell Fund. AH, KM and the Raindrop project were

supported by the John Fell Fund, the Ecological Continuity Trust, the Patsy Wood Trust and the British

Ecological Society.

References

Adobe Lightroom (5.0). (2021). Adobe.

Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Rüger, N., Beck, P. S. A.,
Blach-Overgaard, A., Blok, D., Cornelissen, J. H. C., Forbes, B. C., Georges, D., Goetz, S. J.,
Guay, K. C., Henry, G. H. R., HilleRisLambers, J., Hollister, R. D., Karger, D. N., Kattge, J.,
Manning, P., … Weiher, E. (2018). Plant functional trait change across a warming tundra biome.
Nature, *562*(7725), 57–62. https://doi.org/10.1038/s41586-018-0563-7

Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of*
Statistical Software, *80*. https://doi.org/10.18637/jss.v080.i01

Cavender-Bares, J., Gamon, J. A., & Townsend, P. A. (2020). *Remote Sensing of Plant Biodiversity*.
Springer Open.

Colomina, I., & Molina, P. (2014). Unmanned aerial systems for photogrammetry and remote sensing:
A review. In *ISPRS Journal of Photogrammetry and Remote Sensing* (Vol. 92, pp. 79–97).
Elsevier B.V. https://doi.org/10.1016/j.isprsjprs.2014.02.013

Díaz, S., J. Settele, E. S. Brondízio E.S., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K.
A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M.
Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, … C. N. Zayas. (2019).
IPBES: Summary for policymakers of the global assessment report on biodiversity and
ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and
Ecosystem Services. www.ipbes.net

Frye, H. A., Aiello-Lammens, M. E., Euston-Brown, D., Jones, C. S., Kilroy Mollmann, H., Merow,
C., Slingsby, J. A., van der Merwe, H., Wilson, A. M., & Silander, J. A. (2021). Plant spectral
diversity as a surrogate for species, functional and phylogenetic diversity across a hyper-diverse
biogeographic region. *Global Ecology and Biogeography*, *30*(7), 1403–1417.
https://doi.org/10.1111/geb.13306

Gamon, J. A., Serrano, L., & Surfus, J. S. (1997). The photochemical reflectance index: an optical
indicator of photosynthetic radiation use efficiency across species, functional types, and nutrient
levels. *Oecologia*, *112*, 492–501.

Gholizadeh, H., Gamon, J. A., Helzer, C. J., & Cavender-Bares, J. (2020). Multi-temporal assessment
of grassland α- and β-diversity using hyperspectral imaging. *Ecological Applications*, *30*(7).
https://doi.org/10.1002/eap.2145

Gholizadeh, H., Gamon, J. A., Townsend, P. A., Zygielbaum, A. I., Helzer, C. J., Hmimina, G. Y.,
Yu, R., Moore, R. M., Schweiger, A. K., & Cavender-Bares, J. (2019). Detecting prairie
biodiversity with airborne remote sensing. *Remote Sensing of Environment*, *221*, 34–49.

Gholizadeh, H., Gamon, J. A., Zygielbaum, A. I., Wang, R., Schweiger, A. K., & Cavender-Bares, J.
(2018). Remote sensing of biodiversity: Soil correction and data dimension reduction methods
improve assessment of α-diversity (species richness) in prairie ecosystems. *Remote Sensing of*
Environment, *206*, 240–253. https://doi.org/10.1016/j.rse.2017.12.014

Gibson, C. W. D. (1986). Management History in Relation to Changes in the Flora of Different
Habitats on an Oxfordshire Estate, England. In *Biological Conservation* (Vol. 38).

Gibson, C. W. D., & Brown, V. K. (1991). The Nature and Rate of Development of Calcareous
Grassland in Southern Britain. In *Biological Conservation* (Vol. 51).

Harrison, S. P., Gornish, E. S., & Copeland, S. (2015). Climate-driven diversity loss in a grassland
community. *Proceedings of the National Academy of Sciences of the United States of America*,
112(28), 8672–8677. https://doi.org/10.1073/pnas.1502074112

Headwall Photonics. (2022). *Headwall VNIR 400-1000nm sensor quote*.

Heim, R. H. J., Wright, I. J., Scarth, P., Carnegie, A. J., Taylor, D., & Oldeland, J. (2019).
Multispectral, Aerial Disease Detection for Myrtle Rust (Austropuccinia psidii) on a Lemon
Myrtle Plantation. *Drones*.

Hijmans, R. J. (2020). *raster: Geographic Data Analysis and Modeling*. (3.4-5). CRAN.

Imran, H. A., Gianelle, D., Scotton, M., Rocchini, D., Dalponte, M., Macolino, S., Sakowska, K.,
Pornaro, C., & Vescovo, L. (2021). Potential and limitations of grasslands α-diversity prediction
using fine-scale hyperspectral imagery. *Remote Sensing*, *13*(14).
https://doi.org/10.3390/rs13142649

Jet Propulsion Labratory. (2022, January 26). *AVIRIS: Airborne Visible/Infrared Imaging*
Spectrometer.

Jetz, W., Cavender-Bares, J., Pavlick, R., Schimel, D., Davis, F. W., Asner, G. P., Guralnick, R.,
Kattge, J., Latimer, A. M., Moorcroft, P., Schaepman, M. E., Schildhauer, M. P., Schneider, F.
D., Schrodt, F., Stahl, U., & Ustin, S. L. (2016). Monitoring plant functional diversity from
space. In *Nature Plants* (Vol. 2, Issue 3). Palgrave Macmillan Ltd.
https://doi.org/10.1038/NPLANTS.2016.24

Laliberté, E., Schweiger, A. K., & Legendre, P. (2020). Partitioning plant spectral diversity into alpha
and beta components. *Ecology Letters*, *23*(2), 370–380. https://doi.org/10.1111/ele.13429

Lopatin, J., Fassnacht, F. E., Kattenborn, T., & Schmidtlein, S. (2017). Mapping plant species in
mixed grassland communities using close range imaging spectroscopy. *Remote Sensing of*
Environment, *201*, 12–23. https://doi.org/10.1016/j.rse.2017.08.031

Mairota, P., Cafarelli, B., Didham, R. K., Lovergine, F. P., Lucas, R. M., Nagendra, H., Rocchini, D.,
& Tarantino, C. (2015). Challenges and opportunities in harnessing satellite remote-sensing for
biodiversity monitoring. *Ecological Informatics*, *30*, 207–214.
https://doi.org/10.1016/j.ecoinf.2015.08.006

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A
practical guide for biologists. In *Biological Reviews* (Vol. 85, Issue 4, pp. 935–956).
https://doi.org/10.1111/j.1469-185X.2010.00141.x

Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J. M., Tucker, C. J., & Stenseth, N. C. (2005). Using
the satellite-derived NDVI to assess ecological responses to environmental change. In *Trends in*

Ecology and Evolution (Vol. 20, Issue 9, pp. 503–510).
https://doi.org/10.1016/j.tree.2005.05.011

R Core Team. (2021). *R: A language and environment for statistical computing.* (4.0.4). R Foundation
for Statistical Computing. http://www.r-project.org/

Rossi, C., Kneubühler, M., Schütz, M., Schaepman, M. E., Haller, R. M., & Risch, A. C. (2021).
Remote sensing of spectral diversity: A new methodological approach to account for spatio-
temporal dissimilarities between plant communities. *Ecological Indicators*, *130*.
https://doi.org/10.1016/j.ecolind.2021.108106

Schläpfer, D., Popp, C., & Richter, R. (2020). Drone data atmospheric correction concept for multi-
and hyperspectral imagery-The droacor model. *International Archives of the Photogrammetry,
Remote Sensing and Spatial Information Sciences - ISPRS Archives*, *43*(B3), 473–478.
https://doi.org/10.5194/isprs-archives-XLIII-B3-2020-473-2020

Shannon, C. B., & Weaver, W. (1963). *The Mathematical Theory of Communication*. Urbana.

Simpson, E. H. (1949). Measurement of Diversity. *Nature*, *163*, 688.

Tait, L., Bind, J., Charan-Dixon, H., Hawes, I., Pirker, J., & Schiel, D. (2019). Unmanned aerial
vehicles (UAVs) for monitoring macroalgal biodiversity: Comparison of RGB and multispectral
imaging sensors for biodiversity assessments. *Remote Sensing*, *11*(19).
https://doi.org/10.3390/rs11192332

The European Space Agency. (2022, January 26). *Sentinel Online*.

Thorhallsdottir, T. E. (1990). The Dynamics of a Grassland Community: A Simultaneous
Investigation of Spatial and Temporal Heterogeneity at Various Scales. In *Source: Journal of
Ecology* (Vol. 78, Issue 4).

Tuanmu, M. N., & Jetz, W. (2015). A global, remote sensing-based characterization of terrestrial
habitat heterogeneity for biodiversity and ecosystem modelling. *Global Ecology and
Biogeography*, *24*(11), 1329–1339. https://doi.org/10.1111/geb.12365

Turner, W. (2014). Sensing biodiversity: Sophisticated networks are required to make the best use of
biodiversity data from satellites and in situ sensors. *Science*, *346*(6207), 301–302.
https://doi.org/10.1126/science.1256014

Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., & Steininger, M. (2003). Remote
sensing for biodiversity science and conservation. In *Trends in Ecology and Evolution* (Vol. 18,
Issue 6, pp. 306–314). Elsevier Ltd. https://doi.org/10.1016/S0169-5347(03)00070-3

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out
cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.
https://doi.org/10.1007/s11222-016-9696-4

Villoslada, M., Bergamo, T. F., Ward, R. D., Burnside, N. G., Joyce, C. B., Bunce, R. G. H., & Sepp,
K. (2020). Fine scale plant community assessment in coastal meadows using UAV based
multispectral data. *Ecological Indicators*, *111*. https://doi.org/10.1016/j.ecolind.2019.105979

Wang, R., & Gamon, J. A. (2019). Remote sensing of terrestrial plant biodiversity. *Remote Sensing of
Environment*, *231*. https://doi.org/10.1016/j.rse.2019.111218

Wilson, E. O. (2017). Biodiversity research requires more boots on the ground: Comment. In *Nature
Ecology and Evolution* (Vol. 1, Issue 11, pp. 1590–1591). Nature Publishing Group.
https://doi.org/10.1038/s41559-017-0360-y