Multiple carotenoid-based signals are enhanced despite poor body condition

in urban male and female Northern Cardinals (*Cardinalis cardinalis*)

Daniel T. Baldassarre[1,*], Haley S. Bedell[1], Kristie M. Drzewiecki[1], Brooke D. Goodman[1],

Mya L. Mills[1], and Denis A. Ramos[1]

[1]Department of Biological Sciences, SUNY Oswego, Oswego, NY, USA

*Corresponding author: daniel.baldassarre@oswego.edu

ABSTRACT—If humans aim to sustainably coexist with wildlife, we must understand how our

activity impacts the communication systems of urban animal populations. We know much about

the effects of anthropogenic noise on bird song, but relatively little about how avian visual

signals are affected by urbanization. One way such an effect may occur if urbanization alters the

food available to species with color based on carotenoids, which they must obtain from their diet.

Over three years, we compared a comprehensive suite of visual signals in male and female

Northern Cardinals (*Cardinalis cardinals*) in a rural and an urban population. We predicted that

urban birds would have enhanced carotenoid-based signals as they likely have access to more

carotenoids from invasive plants, especially honeysuckle (*Lonicera* spp.), that thrive in cities.

We used reflectance spectrometry, digital image analyses, and avian visual models to quantify

hue, saturation, and brightness of chest (male), underwing (female), and bill (male and female)

signals. Compared to rural males, urban males had redder chest feathers in one year and redder

bills in every year. Urban females had more saturated underwing color than rural females in

every year. These color differences were sufficient to be distinguished by the avian visual

system. Urbanization did not affect female bill color. Interestingly, urban birds had significantly

reduced mass-related body condition compared to rural birds. These results show that both male

and female urban birds can display enhanced carotenoid-based signals despite being in relatively

poor condition. The consequences of this color enhancement are unknown, but it could affect the

information content of the signals and the dynamics of the social and mating systems. These

results stand in stark contrast to the predominant trend in birds of decreased color in urban areas

and highlight the complex and varied potential effects of urbanization on animal communication.

Key words: avian visual model, bill color, *Lonicera*, plumage color, urbanization.

RESUMEN (Spanish)—*Please translate, thank you*

Humans have drastically altered the natural landscape and it is imperative to understand

how our activity affects animal populations if we aim to promote sustainable coexistence with

wildlife (Seddon et al. 2016, Elmqvist et al. 2019). Beyond the negative impacts of human

activities like direct exploitation (e.g., poaching, pet trade) and habitat destruction on wildlife

population viability, humans can also more subtly alter evolutionary dynamics (Szulkin et al.

2020). For instance, considerable research demonstrates that humans can disrupt animal

communication systems (Shannon et al. 2016, Slabbekoorn et al. 2018). In particular,

urbanization has a myriad of impacts on signal production, transmission, and perception (Sepp et

al. 2018, Sepp et al. 2020, Heinen-Kay et al. 2021). Urban environments present several

challenges that may affect communication systems, including anthropogenic noise and light,

heavy metal pollution, habitat fragmentation, and changes in resource availability (Heinen-Kay

et al. 2021).

In birds, much of this work has focused on anthropogenic disruption of vocal

communication, especially in urban areas (Slabbekoorn and den Boer-Visser 2006, Slabbekoorn

2013, Derryberry and Luther 2021). Less well understood is how visual signals may also be

impacted by human development. Carotenoid-based color patches may be especially affected by

urbanization as carotenoids cannot be synthesized de novo by birds and must be obtained from

the diet (Hill 1991). One way urbanization can directly impact these signals, and hence the

communication system, is if rural and urban areas differ in the access to, amount, or types of

resources containing carotenoids. Moreover, carotenoids can undergo extensive physiological

modification before being deposited in tissue (Goodwin 1984), leaving room for other more

indirect effects of urbanization on color (e.g., via stress or parasites; Møller 2009, Hutton and

McGraw 2016). Additionally, rural and urban habitats likely differ in lighting regime, which can

also affect carotenoid-based color via selection on conspicuousness (Leveau 2021). Importantly,

these mechanisms are not mutually exclusive and might either reduce or enhance color in urban

populations. To date, research comparing carotenoid-based signals between rural and urban

populations has found reduced ornamentation in urban populations of House Finches

(*Haemorhous mexicanus*; Hasegawa et al. 2014, Giraudeau et al. 2015, Giraudeau et al. 2018)

and Great Tits (*Parus major*; Grunst et al. 2020). However, urban birds could have more heavily-

pigmented, enhanced carotenoid-based color if, for example, they have access to carotenoid-rich

invasive plants that thrive in disturbed landscapes (Hobbs 2000, Vilà and Ibáñez 2011). In the

midwestern and northeastern United States, invasive honeysuckle (*Lonicera* spp.) has been

implicated in the enhancement of carotenoid-based signals in a variety of bird species (e.g.,

Northern Cardinal, *Cardinalis cardinalis*; Cedar Waxwing, *Bombycilla cedrorum*; Baltimore

Oriole, *Icterus galbula*; Northern Flicker, *Colaptes auratus*; Mulvihill et al. 1992, Hudon et al.

2013, Hudon et al. 2017, Hudon and Mulvihill 2017). *Lonicera* thrives in urban areas (Borgmann

and Rodewald 2005, Rodewald 2005, Leston and Rodewald 2006), providing a potential link

between urbanization and enhanced color. However, to our knowledge, no study has found

enhanced carotenoid-based signals in an urban population of any species. Quantifying the color

of rural and urban populations of a species that occurs in both habitats provides a powerful

framework to test this idea.

The Northern Cardinal (hereafter "cardinal") is well suited to such a study because it

readily occurs in both rural and urban areas, and both males and females exhibit carotenoid-

based color patches that are displayed during various territorial, courtship, and pre-copulatory

interactions and likely function as signals (Linville et al. 1998, Wolfenbarger 1999a,

Wolfenbarger 1999b, Jawor et al. 2003, Jawor et al. 2004, Jawor and Breitwisch 2004). Males

have bright red plumage overall and orange-red bills, while females have pink-red underwing

patches (hereafter "wing") and orange-red bills. The putative functions of these colors as signals

are complex and variable across studies. For example, males with redder chest color had higher

quality territories (Wolfenbarger 1999a, Rodewald et al. 2011) and reproductive success

(Wolfenbarger 1999a), but females did not prefer redder males (Wolfenbarger 1999b). Males and

females mated assortatively based on chest, wing, and bill color (Jawor et al. 2003); and female

wing and bill color co-varied with body condition (Jawor et al. 2004). The color of carotenoid-

based plumage color in males is related to carotenoid availability in fruits, which they favor

during molt in late summer and fall (Linville and Breitwisch 1997); and captive studies confirm

that carotenoid intake affects plumage color (McGraw and Hill 2001). Cardinals have been

studied in an urban context where males, but not females, had reduced brightness as urbanization

increased across a gradient in Ohio (Jones et al. 2010), which was attributed to the generally

negative effects of urban development. In Florida, rural and urban males did not differ in

plumage color (Leigh 2012). Unlike plumage, the effects of urbanization on bill color have not

been studied in this or any species of which we are aware.

Our goal was to compare multiple carotenoid-based signals (chest, bill, and wing color)

in rural and urban males and females while accounting for the sensitivity of the avian visual

system, which has not been implemented in this species. We hypothesized that urbanization

would affect the color of these signals, specifically that urban birds would have enhanced signals

(e.g., redder hue, increased saturation) compared to rural birds, potentially due to increased

access to carotenoids. We also explored relationships between mass-related body condition,

urbanization, and color.

107   **Methods**

Field sites and data collection

We studied cardinals in 2 populations in Central New York that we chose because they differed

drastically in the extent of human development (Fig. 1). The rural site, Rice Creek Field Station

(43.43°N, 76.55°W; Fig. 1a), was located outside of Oswego (population 17,470), while the

urban field site, Barry Park (43.03°N, 76.12°W; Fig. 1b), was located within the city limits of

Syracuse (population 142,874). The 2 sites were separated by 57 km. At each site, we captured

wild individuals using a territory-based targeted mist net approach between May and July in

2019, 2020, and 2021. We did not re-capture any individuals across years. Although we did not

know the precise age of most individuals, we limited our analyses to those we could age as

"after-hatching-year" (i.e., at least one-year-old) based on plumage. No captured individuals had

yet initiated their prebasic molt.

To quantify body condition, we measured mass to the nearest 0.5 g with a spring scale,

and tarsus length to the nearest 0.01 mm with digital calipers. Both populations were in the midst

of breeding, but we did not have precise information on nest stage for most individuals. For

males and females separately, we regressed mass on tarsus length. Males exhibited a significant

positive relationship between tarsus length and mass ($F_{1,63} = 18.0$, $P < 0.001$), so we extracted

model residuals and used them as an index of body condition. This index described whether an

individual was heavier (positive values) or lighter (negative values) than would be predicted by

body size. There was no significant relationship between female mass and tarsus length ($F_{1,23} =$

0.7, $P = 0.41$), so we used mass as an index of body condition.

We collected 10 feathers from the same central region of the chest of each captured male

for analysis via reflectance spectrometry (rural $n = 43$, urban $n = 22$). In the lab, we mounted the

feathers in an overlapping pattern on black construction paper (Strathmore Artagain Coal Black)

and measured ultraviolet (UV) and visible light reflectance from 300-700 nm using an Ocean

Optics Jaz reflectance spectrometer and a PX2 lamp. The probe was mounted at a 90° angle in a

block that held the probe tip 5 mm from the feather surface and excluded ambient light. Between

each sample, we calibrated the spectrometer using a WS-1 white standard. We took 3

measurements across different regions of the feather sample and averaged the reflectance curves

before analysis (Supplemental Fig. S1). To quantify male bill (rural $n = 40$, urban $n = 21$), female

bill (rural $n = 16$, urban $n = 9$), and female wing (rural $n = 15$, urban $n = 9$) color, we took

photographs in the field using a Canon EOS Rebel T6 digital camera (Supplemental Fig. S2).

Variation in sample size across signals was due to some photographs being unusable. We used

automatic settings because there were no systematic differences in ISO (t = -0.7, df = 47.3, P =

0.5) or aperture (t = -0.8, df = 42.9, P = 0.41) between sites. Each photograph contained color

standards and a scale bar (X-Rite ColorChecker Passport) and was taken under diffuse natural

sunlight without shadows. The inclusion of color standards in each photo allowed us to compare

among images controlling for variation in ambient light. We took 3 photos of each color patch

that varied in exposure (one stop under-exposed, normal, and one stop over-exposed) and then

selected the best-exposed, sharpest image for analysis.

Color analyses

We analyzed reflectance curves of male chest feathers using the program *pavo 2* (Maia et al.

2019) in R 3.6.0 (R Core Team 2019). We obtained quantum catch values for each of the 4 avian

single cones using the Blue Tit (*Cyanistes caeruleus*) visual model (plus double-cone sensitivity

for luminance, hereafter "brightness") and idealized illuminant D65. We then used the quantum

catch values to represent each color as a point in tetrahedral color space where the vertices

represented stimulation of the u, s, m, and l cones. From the color space, we extracted the angle

of the color vector relative to the x-y (theta) and z (phi) plane, which together describe hue.

Theta and phi were highly correlated (R = -0.94, P < 0.001), so we only used theta for

subsequent analyses. To quantify saturation, we used the distance between the point and the

achromatic center relative to the maximum possible distance. Together these measurements

described hue, saturation, and brightness of the chest feathers accounting for UV reflectance and

the sensitivity of the avian visual system.

We analyzed digital photographs of male and female bills and female wing patches using

the Quantitative Color Pattern Analysis (QCPA; van den Berg et al. 2020) framework in the

ImageJ 1.53a (Schneider et al. 2012) plugin micaToolbox (Troscianko and Stevens 2015). We

first generated a cone catch model accounting for the camera's photosensitivity, the Blue Tit

visual model, and idealized illuminant D65. This calibration was achieved by measuring an

image of the color standards, which have known reflectance values. The camera was not UV-

sensitive, so these analyses were restricted to the visible spectrum (400-700 nm). However, these

orange-red color patches reflect predominantly long, visible wavelengths, where most of the

variation occurs (Supplemental Fig. S1). Once a cone catch model was built, we converted each

image into a normalized multispectral stack standardized by the 3% and 92% reflectance

standards and selected regions of interest to measure (Supplemental Fig. S2a–b). We then ran

each image through the cone catch model, followed by the QCPA framework that applies spatial

acuity and viewing distance correction and then uses the Receptor Noise Limited (RNL) model

(Vorobyev and Osorio 1998) to reduce the image into discernable color clusters that can be

measured (Supplemental Fig. S2c–d). We used Gaussian acuity correction with 6 cycles per

degree and a simulated viewing distance of 0.5 m. We used RNL ranked filtering with the

following settings: iterations: 5, radius: 5, falloff: 3. We used RNL clustering with the following

settings: visual system Weber fractions: Blue Tit 0.05, luminance Weber fraction: 0.1, color JND

threshold: 2, luminance JND threshold: 3, loops: 20, radius multiplier: 2, minimum cluster size:

10. For each color cluster produced by QCPA, we extracted the quantum catch values for the 3

avian single cones (excluding the UV-sensitive cone) and the double cone (i.e., brightness).

When an image was split into more than one cluster by QCPA, we computed a weighted average

of each quantum catch value accounting for the area of each cluster. Using *pavo 2*, we converted

these values into a color point in triangular color space where the vertices represented

stimulation of the s, m, and l cones. From the color space, we extracted the single angle of the

color vector (theta), which describes hue. To quantify saturation, we used the distance between

the point and the achromatic center. Together these measurements described hue, saturation, and

brightness of the male and female bill and female wing patches considering the visible spectrum

and the sensitivity of the avian visual system. Note that although both the tetrahedral and

triangular color spaces produce angle theta to describe hue, the difference in geometry is such

that higher values for male chest correspond to more orange hue, while they correspond to redder

hue for male and female bill and female wing.

If we discovered a significant difference in color variables between factors, we further

explored whether these differences would be recognizable by the signal receiver. We input the

quantum catch values into the noise-weighted color distance (dS) calculator in *pavo 2* using the

following settings: Weber fraction: 0.05, noise: neural, photoreceptor densities: u = 1, s = 2, m =

2, l = 4. We compared the resulting values to dS = 1, or the perceptual threshold indicating colors

that can be distinguished by the avian visual system (Vorobyev and Osorio 1998).

Statistical analyses

Because hue, saturation, and brightness represent different and potentially informative aspects of

a color, we analyzed them as separate response variables for each signal. We considered site

(rural vs. urban) and year (2019, 2020, 2021) as factors because color could vary depending on

habitat characteristics or could change systematically due to environmental differences over

time. When checking for collinearity among predictors, we found a significant correlation

between site and body condition using a linear model (see Results), and thus did not include

body condition as a factor in these models. We used a model selection approach, with all

possible combinations of factors as predictors in a linear model, as well as an intercept-only

model. We ranked models by Akaike's Information Criterion corrected for small sample sizes

(AICc) and considered the model with the lowest AICc score to be best supported by the data.

We considered models with ΔAICc < 2 to have similar fits compared to the top model.

Additionally, we used log likelihoods and normalized Akaike weights to compare models. We

conducted post-hoc comparisons using Tukey's HSD corrected for multiple comparisons. To

determine whether dS values between factors were larger than the avian perceptual threshold, we

tested whether the average pairwise dS value of all cross-factor comparisons (e.g., between every

rural and urban color) was significantly greater than one using a one-sample *t*-test. Analyses

were performed using R 3.6.0 (R Core Team 2019) and the *MuMIn* package (Bartón 2020).

Results

The top model for male chest hue included an interaction between site and year (Supplemental

Table S1). There was a significant effect of year ($F_{2,59} = 18.7$, $P < 0.001$) and the interaction

between site and year ($F_{2,59} = 5.0$, $P = 0.01$). Male chest hue in 2020 was higher (more orange)

than 2019 and 2021 (both $P < 0.001$), and urban hue was lower (redder) than rural hue in 2020

($P = 0.03$, Fig. 2a). The top model for male chest saturation included year only (Supplemental

Table S1). There was a significant effect of year ($F_{2,62} = 8.7$, $P < 0.001$) such that saturation in

2019 was higher than 2020 ($P = 0.003$) and 2021 ($P = 0.01$, Fig. 2b). Similarly, the top model for

male chest brightness also included year only (Supplemental Table S1). There was a significant

effect of year ($F_{2,62} = 13.4$, $P < 0.001$) such that brightness in 2021 was lower than 2019 and

2020 (both $P < 0.001$, Fig. 2c). In 2020, the significant difference in male chest hue between the

rural and urban site resulted in an average pairwise dS value that was significantly greater than

the perceptual threshold ($t = 6.0$, df $= 35$, $P < 0.001$, Fig. 4a).

The top model for male bill hue included site only (Supplemental Table S2). There was a

significant effect of site ($F_{1,59} = 12.4$, $P = 0.001$) such that urban hue was higher (redder) than

rural hue (Fig. 3a). There were no significant effects of any factors on male bill saturation or

brightness, as the top models were indistinguishable from an intercept-only model (Supplemental

Table S2). The significant difference in male bill hue between the rural and urban site resulted in

an average pairwise dS value that was significantly greater than the perceptual threshold ($t =$

35.5, df $= 839$, $P < 0.001$, Fig. 4a). There were no significant effects of any factors on female bill

hue, saturation, or brightness (Supplemental Table S3). For female wing saturation, the top

model included site only (Supplemental Table S4). There was a significant effect of site ($F_{1,22} =$

4.9, $P = 0.04$) such that urban saturation was higher than rural saturation (Fig. 3b). There were no

significant effects of any factors on female wing hue or brightness (Supplemental Table S4). The

significant difference in female wing saturation between the rural and urban site resulted in an

average pairwise dS value that was significantly greater than the perceptual threshold ($t = 11.1$,

df $= 134$, $P < 0.001$, Fig. 4a).

Mass-related body condition of both males and females was affected by site. Male body

condition score was higher in rural compared to urban birds ($F_{1,63} = 14.9$, $P < 0.001$, Figure 5a),

and female mass was higher in rural compared to urban birds ($F_{1,23} = 4.5$, $P = 0.05$, Figure 5b).

250   **Discussion**

The color of multiple carotenoid-based signals was enhanced in males and females compared to

rural birds. Specifically, these colors were redder or more saturated in urban birds (Fig. 2–3),

despite these individuals being in relatively poor mass-related body condition compared to rural

birds (Fig. 5). Furthermore, although subtle to the human eye, these color differences were large

enough to be distinguished by the avian visual system (Fig. 4). These findings suggest that urban

living may reduce mass-related body condition, but that urban birds have enough access to

carotenoids that they are nonetheless able to produce what are typically thought of as costly

honest signals. In rural habitats, carotenoids may be a much more limiting resource, such that

rural birds have reduced ornamentation overall, but color may be a more honest signal of

condition. Preliminary analyses have found weak or no relationships between body condition and

ornament color within either population, but we are hesitant to over-interpret these results due to

low sample sizes. Our findings are in line with previous work on cardinals that found a

dampened relationship between body condition, plumage color, and reproductive success as

urbanization increased (Jones et al. 2010, Rodewald et al. 2011). Whether cardinals attend to the

color variation documented here, whether that variation is linked to reproductive success, and

whether the urban color enhancement disrupts the typical signaling dynamics are unknown in our

populations. However, our study stands apart in that we found a distinct enhancement of multiple

carotenoid-based signals in urban birds of both sexes, apparently not associated with increased

mass-related body condition. This is the first such result we are aware of, as the overwhelming

trend in the literature is for reduced ornamentation in urban areas, even among melanin-based or

structural colors (Sepp et al. 2018, Sepp et al. 2020, Heinen-Kay et al. 2021).

A closer inspection of the chest color results supports the idea that increased and/or

consistent access to carotenoids in urban areas could be what promoted the color enhancement.

In 2020 when we found a significant difference in chest hue between rural and urban males,

overall hue was more orange than in other years, but this difference was greater in the rural

habitat, which drove the rural-urban difference (Fig. 2A). This pattern suggests that carotenoid

availability may fluctuate more widely over time in the rural habitat, but that urban birds have a

more stable carotenoid supply that allows them to maintain redder plumage. Leigh (2012) made

a similar interpretation based on urban male cardinals in Florida exhibiting less among-

individual variation in chest brightness than rural males. We suggest that the relative abundance

of exotic fruiting shrubs, especially *Lonicera* spp., in urban areas (Borgmann and Rodewald

2005, Borgmann and Rodewald 2005, Leston and Rodewald 2006) is responsible for the pattern

we observed. Although we have not quantified plant diversity at our sites, anecdotally, *Lonicera*

is much more established at the urban site compared to a more diverse vegetation community at

the rural site (DTB, 2019, pers. observ.). We did not choose these sites a priori based on this

difference. Furthermore, an increase in sampling sites would help determine if the observed color

variation is due to *Lonicera* abundance specifically or other potential differences between rural

and urban sites. Additionally, work is ongoing to quantify carotenoids in blood plasma and

feathers to determine more conclusively if urban birds ingest and/or deposit more carotenoids

than rural birds.

Whether or not carotenoids are more available to urban birds, it is important to note that

many other unmeasured factors may influence whether they are acquired and deposited in tissue.

For example, even if carotenoid-based color is a condition-dependent trait in cardinals, our mass-

related measures of body condition offer only a small glimpse into what might be physiologically

relevant to true condition. Rural and urban birds may differ in other more meaningful measures

of condition such as immune function or stress. Wright and Fokidis (2016) found that urban male

cardinals had lower circulating corticosterone and a lower stress response than rural males.

Urban habitats may be less stressful for cardinals and/or they may modulate their stress response

such that they can more efficiently metabolize and deposit carotenoids in their tissues regardless

of availability (Giraudeau et al. 2015). Related to this idea, in a suite of Western Palearctic

species, Møller (2009) found that urban birds had a stronger immune response than rural birds,

which might free up carotenoids to be deposited in tissue. Additionally, if the lighting conditions

and/or visual background differs between sites, urban and rural birds may experience disruptive

selection to maximize the conspicuousness of their visual signals (Dalrymple et al. 2018, Leveau

2019). In summary, it is difficult with the present data to determine the mechanism of the

observed color variation, and there are several, non-mutually exclusive possibilities.

It is unclear why urban birds were in worse mass-related body condition than rural birds.

Potential explanations include limited access to protein-rich insect prey, benefits of reduced mass

for predator evasion, and adaptive changes in nestling development. In cardinals, previous

studies have found little to no effect of urbanization on similar body condition metrics

(Rodewald and Shustack 2008, Jones et al. 2010) or even that urban birds were in better

condition than rural birds (Wright and Fokidis 2016). However, the potential impacts of

urbanization on body condition may be site-specific, and there is evidence in other species of a

negative effect (Jiménez-Peñuela et al. 2019). Additionally, urban birds may not need to

accumulate energy reserves in the form of body mass in urban areas where food availability is

more predictable and nutritional stress is not as severe as in rural areas (Salleh Hudin et al.

2016). Regardless of the cause, we found that despite this reduction in mass-related body

condition, urban birds were still able to produce enhanced color signals compared to rural birds.

We found an enhancement of both feather and bill color in urban birds. The timing of

production of these two signal types has different implications for the relationship with

carotenoid availability. Cardinals molt once per year after the breeding season, so chest and wing

color should mainly be influenced by carotenoid deposition at that time only. However, it is

possible that the color variation is due to differences between rural and urban birds in the extent

to which feather color wears over time (e.g., due to differences in sun exposure). In contrast to

plumage, carotenoid-based bill color can vary much more dynamically and over shorter time

periods (Rosen and Tarvin 2006, Ardia et al. 2010). If the rural-urban color difference in bill

color is driven by carotenoid availability, these results may suggest that urban birds have greater

access to carotenoids not only during molt, but potentially throughout the breeding season (when

these birds were captured and photographed). Since *Lonicera* fruits are mainly available in late

summer and fall, carotenoids producing bill color may come from different sources (e.g., insect

prey), although fruits can ripen as early as June (DTB, 2020, pers. observ.). However, the extent

to which cardinal bill color changes over time is unknown, and color measured during the

summer breeding season may well have resulted from carotenoids ingested and sequestered

previously. Importantly, the factors influencing color (e.g., carotenoid availability, body

condition, stress) may vary across the signals measured in this study.

Conclusion

Urbanization typically dulls or reduces the extent of color in birds (Sepp et al. 2018, Leveau

2019, Heinen-Kay et al. 2021). Here we show instead that both male and female urban cardinals

had significantly enhanced color of multiple carotenoid-based signals compared to rural birds.

We suggest this enhancement is potentially due to the widespread and stable availability of

carotenoids in exotic plants that thrive in disturbed areas, although there are other possible

explanations, and these mechanisms deserve additional investigation. If carotenoids are a

limiting resource in rural but not urban habitats, the reliability of these signals as honest

indicators of condition may deteriorate (Jones et al. 2010, Rodewald et al. 2011). In support of

this idea, we also found that urban birds were in worse mass-related body condition compared to

rural birds, suggesting a decoupling of condition and color. Together, these results underscore

the complex and subtle impacts of urbanization on wildlife. Not only can urbanization negatively

affect populations by destroying and fragmenting habitat, it may also disrupt communication

systems and potentially alter the evolutionary dynamics of signals.

352   **Acknowledgments**

Susan DeVries, Sylvia Halkin, Jodie Jawor, Gavin Leighton, and Daniel Shustack provided

helpful comments on early drafts of this manuscript. Comments from anonymous reviewers

greatly improved the manuscript. Shyla Luna helped collect data in the field. We are grateful to

Syracuse University and the Onondaga County Department of Water Environment Protection for

access to Barry Park, and to Alan Harris for logistical support at Rice Creek Field Station.

Lauren Silvernail provided childcare while fieldwork was being conducted. This research was

supported by Rice Creek Associates and the Scholarly and Creative Activities Committee at

SUNY Oswego. This research was approved by the SUNY Oswego Institutional Animal Care

and Use Committee (protocol 2018.05) and conducted under appropriate federal (banding permit

24047), state (NYS DEC License 229), and local permits.

364   **Literature cited**

365   Ardia DR, Broughton DR, Gleicher MJ. 2010. Short-term exposure to testosterone propionate

366   leads to rapid bill color and dominance changes in Zebra Finches. Hormones and

367   Behavior. 58:526–532.

Bartón K. 2020. MuMIn: multi-model inference. https://CRAN.R-project.org/package=MuMIn.

van den Berg CP, Troscianko J, Endler JA, Marshall NJ, Cheney KL. 2020. Quantitative Colour
Pattern Analysis (QCPA): a comprehensive framework for the analysis of colour patterns
in nature. Methods in Ecology and Evolution. 11:316–332.

Borgmann KL, Rodewald AD. 2005. Forest restoration in urbanizing landscapes: interactions
between land uses and exotic shrubs. Restoration Ecology. 13:334–340.

Dalrymple RL, Flores-Moreno H, Kemp DJ, White TE, Laffan SW, Hemmings FA, Hitchcock
TD, Moles AT. 2018. Abiotic and biotic predictors of macroecological patterns in bird
and butterfly coloration. Ecological Monographs. 88:204–224.

Derryberry EP, Luther D. 2021. What is known—and not known—about acoustic
communication in an urban soundscape. Integrative and Comparative Biology. 61:1783–
1794.

Elmqvist T, Andersson E, Frantzeskaki N, McPhearson T, Olsson P, Gaffney O, Takeuchi K,
Folke C. 2019. Sustainability and resilience for transformation in the urban century.
Nature Sustainability. 2:267–273.

Giraudeau M, Chavez A, Toomey MB, McGraw KJ. 2015. Effects of carotenoid
supplementation and oxidative challenges on physiological parameters and carotenoid-
based coloration in an urbanization context. Behavioral Ecology and Sociobiology.
69:957–970.

Giraudeau M, Toomey MB, Hutton P, McGraw KJ. 2018. Expression of and choice for

condition-dependent carotenoid-based color in an urbanizing context. Behavioral

Ecology. 29: 1307–1315.

Goodwin TW. 1984. The biochemistry of the carotenoids. Second. New York (NY): Chapman

and Hall.

Grunst ML, Grunst AS, Pinxten R, Bervoets L, Eens M. 2020. Carotenoid- but not melanin-

based plumage coloration is negatively related to metal exposure and proximity to the

road in an urban songbird. Environmental Pollution. 256:113473.

Hasegawa M, Ligon RA, Giraudeau M, Watanabe M, McGraw KJ. 2014. Urban and colorful

male House Finches are less aggressive. Behavioral Ecology. 25:641–649.

Heinen-Kay JL, Kay AD, Zuk M. 2021. How urbanization affects sexual communication.

Ecology and Evolution. 11:17625–17650.

Hill GE. 1991. Plumage coloration is a sexually selected indicator of male quality. Nature.

350:337–339.

Hobbs RJ. 2000. Land-use changes an invasions. In: Mooney, HA, Hobbs, RJ, editors. Invasive

species in a changing world. Washington (DC): Island Press;p. 55–64.

Hudon J, Derbyshire D, Leckie S, Flinn T. 2013. Diet-induced plumage erythrism in Baltimore

Orioles as a result of the spread of introduced shrubs. Wilson Journal of Ornithology.

125:88–96.

Hudon J, Driver RJ, Rice NH, Lloyd-Evans TL, Craves JA, Shustack DP. 2017. Diet explains

red flight feathers in Yellow-shafted Flickers in eastern North America. The Auk.

134:22–33.

Hudon J, Mulvihill R. 2017. Diet-induced plumage erythrism as a result of the spread of alien

shrubs in North America. North American Bird Bander. 42:95–103.

Hutton P, McGraw KJ. 2016. Urban impacts on oxidative balance and animal signals. Frontiers

in Ecology and Evolution. 4:54.

Jawor JM, Breitwisch R. 2004. Multiple ornaments in male Northern Cardinals, *Cardinalis*

cardinalis, as indicators of condition. Ethology. 110:113–126.

Jawor JM, Gray N, Beall SM, Breitwisch R. 2004. Multiple ornaments correlate with aspects of

condition and behaviour in female Northern Cardinals, *Cardinalis cardinalis*. Animal

Behavior. 67:875–882.

Jawor JM, Linville SU, Beall SM, Breitwisch R. 2003. Assortative mating by multiple ornaments

in Northern Cardinals (*Cardinalis cardinalis*). Behavioral Ecology. 14:515–520.

Jiménez-Peñuela J, Ferraguti M, Martínez-de la Puente J, Soriguer R, Figuerola J. 2019.

Urbanization and blood parasite infections affect the body condition of wild birds.

Science of The Total Environmen. 651:3015–3022.

Jones TM, Rodewald AD, Shustack DP. 2010. Variation in plumage coloration of Northern

Cardinals in urbanizing landscapes. Wilson Journal of Ornithology. 122:326–333.

Leigh BA. 2012. Does urbanization influence coloration of Northern Cardinals (*Cardinalis*
cardinalis)? [honors thesis]. Tampa (FL): University of South Florida.

Leston LFV, Rodewald AD. 2006. Are urban forests ecological traps for understory birds? An
examination using Northern Cardinals. Biological Conservation. 131:566–574.

Leveau L. 2021. United colours of the city: a review about urbanisation impact on animal
colours. Austral Ecology. 46:670–679.

Leveau LM. 2019. Urbanization induces bird color homogenization. Landscape and Urban
Planning. 192:103645.

Linville SU, Breitwisch R. 1997. Carotenoid availability and plumage coloration in a wild
population of Northern Cardinals. The Auk. 114:796–800.

Linville SU, Breitwisch R, Schilling AJ. 1998. Plumage brightness as an indicator of parental
care in Northern Cardinals. Animal Behavior. 55:119–127.

Maia R, Gruson H, Endler JA, White TE. 2019. pavo 2: New tools for the spectral and spatial
analysis of colour in R. Methods in Ecology and Evolution. 10:1097–1107.

McGraw KJ, Hill GE. 2001. Carotenoid access and intraspecific variation in plumage
pigmentation in male American Goldfinches (*Carduelis tristis*) and Northern Cardinals
(*Cardinalis cardinalis*). Functional Ecology. 15:732–739.

Møller AP. 2009. Successful city dwellers: a comparative study of the ecological characteristics
of urban birds in the Western Palearctic. Oecologia. 159:849–858.

Mulvihill RS, Parkes KC, Leberman RC, Wood DS. 1992. Evidence supporting a dietary basis

for orange-tipped rectrices in the Cedar Waxwing. Journal of Field Ornithology. 63:212–

216.

R Core Team. 2019. R: a language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing. http://www.R-project.org.

Rodewald AD. 2005. Interaction between exotic shrubs and breeding birds in riparian forests. In:

Cardina, J, editor. Ohio invasive plant research conference: bridging the gap between

land management and research. Special Circular 196: Aug 2005. Ohio Agricultural and

Development Center. Wooster (OH): The Ohio State University; p. 43–48.

Rodewald AD, Shustack DP. 2008. Consumer resource matching in urbanizing landscapes: are

synanthropic species over-matching? Ecology. 89:515–521.

Rodewald AD, Shustack DP, Jones TM. 2011. Dynamic selective environments and evolutionary

traps in human-dominated landscapes. Ecology. 92:1781–1788.

Rosen RF, Tarvin KA. 2006. Sexual signals of the male American Goldfinch. Ethology.

112:1008–1019.

Salleh Hudin N, Strubbe D, Teyssier A, De Neve L, White J, Janssens GPJ, Lens L. 2016.

Predictable food supplies induce plastic shifts in avian scaled body mass. Behavioral

Ecology. 27:1833–1840.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nature Methods. 9:671–675.

Seddon N, Mace GM, Naeem S, Tobias JA, Pigot AL, Cavanagh R, Mouillot D, Vause J,

Walpole M. 2016. Biodiversity in the Anthropocene: prospects and policy. Proceedings

of The Royal Society of London B: Biological Sciences. 283:20162094.

Sepp T, McGraw KJ, Giraudeau M. 2020. Urban sexual selection. In: Szulkin M, Munshi-South

J, Charmantier A, editors. Urban evolutionary biology. Oxford (UK): Oxford University

Press; p. 234–252.

Sepp T, McGraw KJ, Kaasik A, Giraudeau M. 2018. A review of urban impacts on avian life-

history evolution: does city living lead to slower pace of life? Global Change Biology.

24:1452–1469.

Shannon G, McKenna MF, Angeloni LM, Crooks KR, Fristrup KM, Brown E, Warner KA,

Nelson MD, White C, Briggs J, et al. 2016. A synthesis of two decades of research

documenting the effects of noise on wildlife: effects of anthropogenic noise on wildlife.

Biological Reviews. 91:982–1005.

Slabbekoorn H. 2013. Songs of the city: noise-dependent spectral plasticity in the acoustic

phenotype of urban birds. Animal Behavior. 85:1089–1099.

Slabbekoorn H, den Boer-Visser A. 2006. Cities change the songs of birds. Current Biology.

16:2326–2331.

Slabbekoorn H, Dooling R, Popper AN, Fay RR. 2018. Effects of anthropogenic noise on

animals. New York (NY): Springer.

Szulkin M, Munshi-South J, Charmantier A. 2020. Urban evolutionary biology. Oxford (UK):

Oxford University Press.

Troscianko J, Stevens M. 2015. Image calibration and analysis toolbox – a free software suite for

objectively measuring reflectance, colour and pattern. Methods in Ecology and

Evolution. 6:1320–1331.

Vilà M, Ibáñez I. 2011. Plant invasions in the landscape. Landscape Ecology. 26:461–472.

Vorobyev M, Osorio D. 1998. Receptor noise as a determinant of colour thresholds. Proceedings

of The Royal Society of London B: Biological Sciences. 265:351–358.

Wolfenbarger LL. 1999a. Red coloration of male Northern Cardinals correlates with mate quality

and territory quality. Behavioral Ecology. 10:80–90.

Wolfenbarger LL. 1999b. Female mate choice in Northern Cardinals (*Cardinalis cardinalis*): is

there a preference for redder males? Wilson Bulletin. 111:76–83.

Wright S, Fokidis HB. 2016. Sources of variation in plasma corticosterone and

dehydroepiandrosterone in the male Northern Cardinal (*Cardinalis cardinalis*): II. Effects

of urbanization, food supplementation and social stress. General and Comparative

Endocrinology. 235:201–209.

503

504   **Figure 1.** A map of the rural (Rice Creek Field Station, a) and urban (Barry Park, b) field sites.

Figure 2. Variation in male chest hue (a), saturation (b), and brightness (c) as a function of
urbanization and/or year. There was no significant effect of site on saturation or brightness, so
only the effect of year is shown. Lowercase letters indicate which factors are (different letter) or
are not (same letter) significantly different from each other.

525 **Figure 3.** Variation in male bill hue (a) and female wing saturation (b) as a function of

526 urbanization. Asterisks separate factors that are significantly different.

Figure 4: Pairwise chromatic contrast between rural and urban male chest (2020 only), male bill,

and female wing (a). The dashed line indicates dS = 1, or the perceptual threshold above which

colors can be distinguished by the avian visual system. Photos exemplifying the average rural

and urban male chest (b, c), male bill (d, e), and female wing (f, g) signals.

552 **Figure 5.** Variation in male body condition score (a) and female mass (b) as a function of

553 urbanization.