Enhanced signal color in urban Northern Cardinals

RESEARCH ARTICLE

Urbanization enhances multiple carotenoid-based signals despite poor body condition in

male and female Northern Cardinals

Daniel T. Baldassarre[1,*], Haley S. Bedell[1], Kristie M. Drzewiecki[1], Brooke D. Goodman[1], Mya

L. Mills[1], and Denis A. Ramos[1]

[1]Department of Biological Sciences, SUNY Oswego, Oswego, NY, USA

*corresponding author: daniel.baldassarre@oswego.edu

ACKNOWLEDGEMENTS

Susan DeVries, Sylvia Halkin, Jodie Jawor, Gavin Leighton, and Daniel Shustack provided

helpful comments on early drafts of this manuscript. Comments from anonymous reviewers

greatly improved the manuscript. Shyla Luna helped collect data in the field. We are grateful to

Syracuse University and the Onondaga County Department of Water Environment Protection for

access to Barry Park, and to Alan Harris for logistical support at Rice Creek Field Station.

Lauren Silvernail provided childcare while fieldwork was being conducted.

Funding statement: This research was supported by Rice Creek Associates and the Scholarly

and Creative Activities Committee at SUNY Oswego.

Ethics statement: This research was approved by the SUNY Oswego Institutional Animal Care

and Use Committee. D.B. had appropriate federal, state, and local permits.

Author contributions: D.B. conceived the idea, developed methods, collected data, analyzed

data, and wrote the paper. H.B., K.D., B.G., M.M., and D.R. collected data and substantially

edited the paper.

Data availability: Analyses in this article can be reproduced using the data provided by

Baldassarre et al. (2022).

ABSTRACT

If humans aim to sustainably coexist with wildlife, we must understand how our activity impacts

the communication systems of urban animal populations. We know much about the effects of

anthropogenic noise on bird song, but relatively little about how avian visual signals are affected

by urbanization. Such an effect may occur if urbanization alters the food available to species

with color based on carotenoids, which they must obtain from their diet. Over three years, we

compared a comprehensive suite of visual signals in male and female Northern Cardinals

(*Cardinalis cardinals*) in a rural and an urban population. We predicted that urban birds would

have enhanced carotenoid-based signals as they likely have access to more carotenoids from

invasive plants, especially honeysuckle (*Lonicera* spp.), that thrive in cities. We used reflectance

spectrometry, digital image analyses, and avian visual models to quantify hue, saturation, and

brightness of chest (male), underwing (female), and bill (male and female) signals. Compared to

rural males, urban males had redder chest feathers in one year and redder bills in every year.

Urban females had more saturated underwing color than rural females in every year. These color

differences were sufficient to be distinguished by the avian visual system. Urbanization did not

affect female bill color. Interestingly, urban birds had significantly reduced mass-related body

condition compared to rural birds. These results show that both male and female urban birds can

display enhanced carotenoid-based signals despite being in relatively poor condition. The

consequences of this color enhancement are unknown, but it could affect the information content

of the signals and the dynamics of the social and mating systems. These results stand in stark

contrast to the predominant trend in birds of decreased color in urban areas and highlight the

complex and varied potential effects of urbanization on animal communication.

Keywords: avian visual model, bill color, carotenoids, *Lonicera*, Northern Cardinal, plumage

color, signals, urbanization

LAY SUMMARY

• Urbanization typically dulls bird color, but carotenoid-based signals may be enhanced if

urban birds have increased access to carotenoids from invasive plants.

• Urban Northern Cardinals of both sexes had enhanced signal color compared to rural

birds, but were in worse body condition.

• Urban birds likely had greater access to carotenoids from invasive plants, allowing them

to produce enhanced color despite poor condition.

• Urbanization can disrupt animal communication and possibly alter evolutionary

dynamics.

INTRODUCTION

Humans have drastically altered the natural landscape and it is imperative to understand how our

activity affects animal populations if we aim to promote sustainable coexistence with wildlife

(Seddon et al. 2016, Elmqvist et al. 2019). Beyond the negative impacts of human activities like

direct exploitation (e.g., poaching, pet trade) and habitat destruction on wildlife population

viability, humans can also more subtly alter evolutionary dynamics (Szulkin et al. 2020). For

instance, considerable research demonstrates that humans can disrupt animal communication

systems (Shannon et al. 2016, Slabbekoorn et al. 2018). In particular, urbanization has a myriad

of impacts on signal production, transmission, and perception (Sepp et al. 2018, 2020; Heinen-

Kay et al. 2021). Urban environments present several challenges that may affect communication

systems, including anthropogenic noise and light, heavy metal pollution, habitat fragmentation,

and changes in resource availability.

In birds, much of this work has focused on anthropogenic disruption of vocal

communication, especially in urban areas (Slabbekoorn and den Boer-Visser 2006, Slabbekoorn

2013, Derryberry and Luther 2021). Less well understood is how visual signals may also be

impacted by human development. Carotenoid-based color patches may be especially affected by

urbanization as carotenoids cannot be synthesized de novo by birds and must be obtained from

the diet (Hill 1991). One way urbanization can directly impact these signals, and hence the

communication system, is if rural and urban areas differ in the access to, amount, or types of

resources containing carotenoids. Moreover, carotenoids can undergo extensive physiological

modification before being deposited in tissue (Goodwin 1984), leaving room for other more

indirect effects of urbanization on color (e.g., via stress or parasites). To date, research

comparing carotenoid-based signals between rural and urban populations has found reduced

ornamentation in urban populations of House Finches (*Haemorhous mexicanus*; Hasegawa et al.

2014, Giraudeau et al. 2015, 2018) and Great Tits (*Parus major*; Grunst et al. 2020). However,

urban birds could have enhanced carotenoid-based color if, for example, they have access to

carotenoid-rich invasive plants that thrive in disturbed landscapes (Hobbs 2000, Vilà and Ibáñez

2011). In the midwestern and northeastern United States, invasive honeysuckle (*Lonicera* spp.)

has been implicated in the enhancement of carotenoid-based signals in a variety of bird species

(e.g., Northern Cardinal, *Cardinalis cardinalis*; Cedar Waxwing, *Bombycilla cedrorum*;

Baltimore Oriole, *Icterus galbula*; Northern Flicker, *Colaptes auratus*; Mulvihill et al. 1992,

Hudon et al. 2013, 2017; Hudon and Mulvihill 2017). *Lonicera* thrives in urban areas (Rodewald

2005, Borgmann and Rodewald 2005, Leston and Rodewald 2006), providing a potential link

between urbanization and enhanced color. However, to our knowledge, no study has found

enhanced carotenoid-based signals in any species because of urbanization. Quantifying the color

of rural and urban populations of the same species provides a powerful framework to test this

idea.

The Northern Cardinal (hereafter "cardinal") is well suited to such a study because it

readily occurs in both rural and urban areas, and both males and females exhibit carotenoid-

based color patches that are displayed during various territorial, courtship, and pre-copulatory

interactions and likely function as signals (Linville et al. 1998, Wolfenbarger 1999a, b; Jawor et

al. 2003, 2004; Jawor and Breitwisch 2004). Males have bright red plumage overall and orange-

red bills, while females have pink-red underwing patches (hereafter "wing") and orange-red bills.

The putative functions of these colors as signals are complex and variable across studies. For

example, males with redder chest color had higher quality territories (Wolfenbarger 1999b,

Rodewald et al. 2011) and reproductive success (Wolfenbarger 1999b), but females did not

prefer redder males (Wolfenbarger 1999a). Males and females mated assortatively based on

chest, wing, and bill color (Jawor et al. 2003); and female wing and bill color co-varied with

body condition (Jawor et al. 2004). The color of carotenoid-based plumage color in males is

related to carotenoid availability in fruits, which they favor during molt in late summer and fall

(Linville and Breitwisch 1997); and captive studies confirm that carotenoid intake affects

plumage color (McGraw and Hill 2001). Cardinals have been studied in an urban context where

males, but not females, had reduced brightness as urbanization increased across a gradient in

Ohio (Jones et al. 2010), which was attributed to the generally negative effects of urban

development. It is not obvious how carotenoid access would affect brightness per se, whereas the

link between increased pigmentation and more saturated, redder hue is clear. Unlike plumage,

the effects of urbanization on bill color have not been studied in any species of which we are

aware.

Our goal was to compare multiple carotenoid-based signals (chest, bill, and wing color)

in rural and urban males and females while accounting for the sensitivity of the avian visual

system. We hypothesized that urbanization would affect the color of these signals, specifically

that urban birds would have enhanced signals (e.g., redder hue, increased saturation) compared

to rural birds due to increased access to carotenoids. We also explored relationships between

mass-related body condition, urbanization, and color.

METHODS

Field Sites and Data Collection

We studied cardinals in two populations in Central New York that differed drastically in the

extent of human development (Figure 1). The rural site, Rice Creek Field Station (43.43°N,

76.55°W; Figure 1A), was located outside of Oswego (population 17,470), while the urban field

site, Barry Park (43.03°N, 76.12°W; Figure 1B), was located within the city limits of Syracuse

(population 142,874). The two sites were separated by 57 km. At each site, we captured wild

individuals using a territory-based targeted mist net approach between May and August in 2019,

2020, and 2021. We did not re-capture any individuals across years. Although we did not know

the precise age of most individuals, we limited our analyses to those we could age as "after-

hatching-year" (i.e. 1+ years old) based on plumage. To avoid comparing color among

individuals with worn and new feathers, we excluded individuals that had initiated their prebasic

molt.

To quantify body condition, we measured mass to the nearest 0.5 g with a spring scale,

and tarsus length to the nearest 0.01 mm with digital calipers. There were no obvious differences

between populations in breeding status (i.e. both were in the midst of breeding), but we did not

have precise information on nest stage for most individuals. For males and females separately,

we regressed mass on tarsus length. Males exhibited a significant positive relationship between

tarsus length and mass ($F = 18.0$, df $= 1$ and 63, $P < 0.001$), so we extracted model residuals and

used them as an index of body condition. This index described whether an individual was heavier

(positive values) or lighter (negative values) than would be predicted by body size. There was no

significant relationship between female mass and tarsus length ($F = 0.7$, df $= 1$ and 23, $P = 0.41$),

so we used mass as an index of body condition.

We collected 10 feathers from the chest of each captured male for analysis via reflectance

spectrometry (rural $n = 43$, urban $n = 22$). In the lab, we mounted the feathers in an overlapping

pattern on black construction paper (Strathmore Artagain Coal Black) and measured ultraviolet

(UV) and visible light reflectance from 300-700 nm using an Ocean Optics Jaz reflectance

spectrometer and a PX2 lamp. The probe was mounted at a 90° angle in a block that held the

probe tip 5 mm from the feather surface and excluded ambient light. Between each sample, we

calibrated the spectrometer using a WS-1 white standard. We took three measurements across

different regions of the feather sample and averaged the reflectance curves before analysis

(Supplementary Material Figure S1). To quantify male bill (rural $n = 40$, urban $n = 21$), female

bill (rural $n = 16$, urban $n = 9$), and female wing (rural $n = 15$, urban $n = 9$) color, we took

photographs in the field using a Canon EOS Rebel T6 digital camera (Supplementary Material

Figure S2). Variation in sample size across signals was due to some photographs being unusable.

We used automatic settings because there were no systematic differences in ISO ($t = -0.7$, df $=$

47.3, $P = 0.5$) or aperture ($t = -0.8$, df $= 42.9$, $P = 0.41$) between sites. Each photograph

contained color standards and a scale bar (X-Rite ColorChecker Passport) and was taken under

diffuse natural sunlight without shadows. The inclusion of color standards in each photo allowed

us to compare among images controlling for variation in ambient light. We took three photos of

each color patch that varied in exposure (one stop under-exposed, normal, and one stop over-

exposed) and then selected the best-exposed, sharpest image for analysis.

Color Analyses

We analyzed reflectance curves of male chest feathers using the program *pavo 2* (Maia et al.

2019) in R 3.6.0 (R Core Team 2019). We obtained quantum catch values for each of the four

avian single cones using the Blue Tit visual model (plus double-cone sensitivity for luminance,

hereafter "brightness") and idealized illuminant D65. We then used the quantum catch values to

represent each color as a point in tetrahedral color space where the vertices represented

stimulation of the u, s, m, and l cones. From the color space, we extracted the angle of the color

vector relative to the x-y (theta) and z (phi) plane, which together describe hue. Theta and phi

were highly correlated (chest $r = -0.94$, $P < 0.001$), so we only used theta for subsequent

analyses. To quantify saturation, we used the distance between the point and the achromatic

center relative to the maximum possible distance. Together these measurements described hue,

saturation, and brightness of the chest feathers accounting for UV reflectance and the sensitivity

of the avian visual system.

We analyzed digital photographs of male and female bills and female wing patches using

the Quantitative Color Pattern Analysis (QCPA; van den Berg et al. 2020) framework in the

ImageJ 1.53a (Schneider et al. 2012) plugin micaToolbox (Troscianko and Stevens 2015). We

first generated a cone catch model accounting for the camera's photosensitivity, the Blue Tit

visual model, and idealized illuminant D65. This calibration was achieved by measuring an

image of the color standards, which have known reflectance values. The camera was not UV-

sensitive, so these analyses were restricted to the visible spectrum (400-700 nm). However, these

orange-red color patches reflect predominantly long, visible wavelengths, where most of the

variation occurs (Supplementary Material Figure S1). Once a cone catch model was built, we

converted each image into a normalized multispectral stack standardized by the 3% and 92%

reflectance standards and selected regions of interest to measure (Supplementary Material Figure

S2A, B). We then ran each image through the cone catch model, followed by the QCPA

framework that applies spatial acuity and viewing distance correction and then uses the Receptor

Noise Limited (RNL) model (Vorobyev and Osorio 1998) to reduce the image into discernable

color clusters that can be measured (Supplementary Material Figure S2C, D). We used Gaussian

acuity correction with six cycles per degree and a simulated viewing distance of 0.5 m. We used

RNL ranked filtering with the following settings: iterations: 5, radius: 5, falloff: 3. We used RNL

clustering with the following settings: visual system Weber fractions: Blue Tit 0.05, luminance

Weber fraction: 0.1, color JND threshold: 2, luminance JND threshold: 3, loops: 20, radius

multiplier: 2, minimum cluster size: 10. For each color cluster produced by QCPA, we extracted

the quantum catch values for the three avian single cones (excluding the UV-sensitive cone) and

the double cone (i.e. brightness). When an image was split into more than one cluster by QCPA,

we computed a weighted average of each quantum catch value accounting for the area of each

cluster. Using *pavo 2*, we converted these values into a color point in triangular color space

where the vertices represented stimulation of the s, m, and l cones. From the color space, we

extracted the single angle of the color vector (theta), which describes hue. To quantify saturation,

we used the distance between the point and the achromatic center. Together these measurements

described hue, saturation, and brightness of the male and female bill and female wing patches

considering the visible spectrum and the sensitivity of the avian visual system. Note that

although both the tetrahedral and triangular color spaces produce angle theta to describe hue, the

difference in geometry is such that higher values for male chest correspond to more orange hue,

while they correspond to redder hue for male and female bill and female wing.

If we discovered a significant difference in color variables between factors, we further

explored whether these differences would be recognizable by the signal receiver. We input the

quantum catch values into the noise-weighted color distance (dS) calculator in *pavo 2* using the

following settings: Weber fraction: 0.05, noise: neural, photoreceptor densities: u = 1, s = 2, m =

2, l = 4. We compared the resulting values to dS = 1, or the perceptual threshold indicating colors

that can be distinguished by the avian visual system (Vorobyev and Osorio 1998).

Statistical Analyses

Because hue, saturation, and brightness represent different and potentially informative aspects of

a color, we analyzed them as separate response variables for each signal. We considered site

(rural vs. urban) and year (2019, 2020, 2021) as factors because color could vary depending on

habitat characteristics or could change systematically due to environmental differences over

time. When checking for collinearity among predictors, we found a significant correlation

between site and body condition using a linear model (see Results), and thus did not include

body condition as a factor in these models. We used a model selection approach, with all

possible combinations of factors as predictors in a linear model, as well as an intercept-only

model. We ranked models by Akaike's Information Criterion corrected for small sample sizes

(AICc) and considered the model with the lowest AICc score to be best supported by the data.

We considered models with ΔAICc < 2 to have similar fits compared to the top model.

Additionally, we used log likelihoods and normalized Akaike weights to compare models. We

conducted post-hoc comparisons using Tukey's HSD corrected for multiple comparisons. To

determine whether dS values between factors were larger than the avian perceptual threshold, we

tested whether the average pairwise dS value of all cross-factor comparisons (e.g., between every

rural and urban color) was significantly greater than one using a one-sample t-test. Analyses

were performed using R 3.6.0 (R Core Team 2019) and the *MuMIn* package (Bartón 2020).

RESULTS

The top model for male chest hue included an interaction between site and year (Supplementary

Material Table S1). There was a significant effect of year ($F = 18.7$, df = 2 and 59, $P < 0.001$)

and the interaction between site and year ($F = 5.0$, df = 2 and 59, $P = 0.01$). Male chest hue in

2020 was higher (more orange) than 2019 and 2021 (both $P < 0.001$), and urban hue was lower

(redder) than rural hue in 2020 ($P = 0.03$, Figure 2A). The top model for male chest saturation

included year only (Supplementary Material Table S1). There was a significant effect of year (F

$= 8.7$, df = 2 and 62, $P < 0.001$) such that saturation in 2019 was higher than 2020 ($P = 0.003$)

and 2021 ($P = 0.01$, Figure 2B). Similarly, the top model for male chest brightness also included

year only (Supplementary Material Table S1). There was a significant effect of year ($F = 13.4$, df

$= 2$ and 62, $P < 0.001$) such that brightness in 2021 was lower than 2019 and 2020 (both $P <$

0.001, Figure 2C). In 2020, the significant difference in male chest hue between the rural and

urban site resulted in an average pairwise dS value that was significantly greater than the

perceptual threshold ($t = 6.0$, df = 35, $P < 0.001$, Figure 4A).

The top model for male bill hue included site only (Supplementary Material Table S2).

There was a significant effect of site ($F = 12.4$, df = 1 and 59, $P = 0.001$) such that urban hue was

higher (redder) than rural hue (Figure 3A). There were no significant effects of any factors on

male bill saturation or brightness, as the top models were indistinguishable from an intercept-

only model (Supplementary Material Table S2). The significant difference in male bill hue

between the rural and urban site resulted in an average pairwise dS value that was significantly

greater than the perceptual threshold ($t = 35.5$, df = 839, $P < 0.001$, Figure 4A). There were no

significant effects of any factors on female bill hue, saturation, or brightness (Supplementary

Material Table S3). For female wing saturation, the top model included site only (Supplementary

Material Table S4). There was a significant effect of site ($F = 4.9$, df = 1 and 22, $P = 0.04$) such

that urban saturation was higher than rural saturation (Figure 3B). There were no significant

effects of any factors on female wing hue or brightness (Supplementary Material Table S4). The

significant difference in female wing saturation between the rural and urban site resulted in an

average pairwise dS value that was significantly greater than the perceptual threshold ($t = 11.1$,

df = 134, $P < 0.001$, Figure 4A).

Mass-related body condition of both males and females was affected by site. Male body

condition score was higher in rural compared to urban birds ($F = 14.9$, df = 1 and 63, $P < 0.001$,

Figure 5A), and female mass was higher in rural compared to urban birds ($F = 4.5$, df = 1 and 23,

$P = 0.05$, Figure 5B).

DISCUSSION

Urbanization enhanced the color of multiple carotenoid-based signals in both males and females.

Specifically, these colors were redder or more saturated in urban birds (Figures 2, 3), despite

these individuals being in relatively poor mass-related body condition compared to rural birds

(Figure 5). Furthermore, these color differences – although subtle to the human eye – were large

enough to be distinguished by the avian visual system (Figure 4). These findings suggest that

urban living may reduce mass-related body condition, but that urban birds have enough access to

carotenoids that they are nonetheless able to produce what are typically thought of as costly

honest signals. In rural habitats, carotenoids may be a much more limiting resource, such that

rural birds have reduced ornamentation overall, but color may be a more honest signal of

condition. Preliminary analyses have found weak or no relationships between body condition and

ornament color within either population, but we are hesitant to over-interpret these results due to

low sample sizes. Our findings are in line with previous work on cardinals that found a

dampened relationship between body condition, plumage color, and reproductive success as

urbanization increased (Jones et al. 2010, Rodewald et al. 2011). Whether cardinals attend to the

color variation documented here, whether that variation is linked to reproductive success, and

whether the urban color enhancement disrupts the typical signaling dynamics are unknown in our

populations. However, our study stands apart in that we found a distinct enhancement of multiple

carotenoid-based signals in urban birds of both sexes, apparently not associated with increased

mass-related body condition. This is the first such result we are aware of, as the overwhelming

trend in the literature is for reduced ornamentation in urban areas, even among melanin-based or

structural colors (Sepp et al. 2018, 2020; Heinen-Kay et al. 2021).

A closer inspection of the chest color results supports the idea that increased and/or

consistent access to carotenoids in urban areas could be what promoted the color enhancement.

In 2020 when we found a significant difference in chest hue between rural and urban males,

overall hue was more orange than in other years, but this difference was greater in the rural

habitat, which drove the rural-urban difference (Figure 2A). This pattern suggests that carotenoid

availability may fluctuate more widely over time in the rural habitat, but that urban birds have a

more stable carotenoid supply that allows them to maintain redder plumage. We suggest that the

relative abundance of exotic fruiting shrubs, especially *Lonicera* spp., in urban areas (Borgmann

and Rodewald 2005; Leston and Rodewald 2006) is responsible for this pattern. Although we

have not quantified plant diversity at our sites, anecdotally, *Lonicera* is much more established at

the urban site compared to a more diverse vegetation community at the rural site. Furthermore,

work is ongoing to quantify carotenoids in blood plasma and feathers to determine more

conclusively if urban birds ingest and/or deposit more carotenoids than rural birds.

Even if carotenoids are more available to urban birds, it is important to note that many

other unmeasured factors may influence whether they are acquired and deposited in tissue. For

example, even if carotenoid-based color is a condition-dependent trait in cardinals, our mass-

related measures of body condition offer only a small glimpse into what might be physiologically

relevant to true condition. Rural and urban birds may differ in other more meaningful measures

of condition such as immune function or stress. Wright and Fokidis (2016) found that urban male

cardinals had lower circulating corticosterone and a lower stress response than rural males.

Urban habitats may be less stressful for cardinals and/or they may modulate their stress response

such that they can more efficiently metabolize and deposit carotenoids in their tissues regardless

of availability (Giraudeau et al. 2015). Additionally, if the lighting conditions and/or visual

background differs between sites, urban and rural birds may experience disruptive selection to

maximize the conspicuousness of their visual signals (Dalrymple et al. 2018, Leveau 2019).

It is unclear why urban birds were in worse mass-related body condition than rural birds.

Potential explanations include limited access to protein-rich insect prey, benefits of reduced mass

for predator evasion, and adaptive changes in nestling development. In cardinals, previous

studies have found little to no effect of urbanization on similar body condition metrics

(Rodewald and Shustack 2008, Jones et al. 2010) or even that urban birds were in better

condition than rural birds (Wright and Fokidis 2016). However, the potential impacts of

urbanization on body condition may be site-specific, and there is evidence in other species of a

negative effect (Jiménez-Peñuela et al. 2019). Additionally, urban birds may not need to

accumulate energy reserves in the form of body mass in urban areas where food availability is

more predictable and nutritional stress is not as severe as in rural areas (Salleh Hudin et al.

2016). Regardless of the cause, we found that despite this reduction in mass-related body

condition, urban birds were still able to produce enhanced color signals compared to rural birds.

We found an enhancement of both feather and bill color in urban birds. The timing of

production of these two signal types has different implications for the relationship with

carotenoid availability. Cardinals molt once per year after the breeding season, so chest and wing

color would be influenced by carotenoid intake at that time only. In contrast, carotenoid-based

bill color can vary much more dynamically and over shorter time periods (Rosen and Tarvin

2006, Ardia et al. 2010). If the rural-urban color difference in bill color is driven by carotenoid

availability, these results suggest that urban birds have greater access to carotenoids not only

during molt, but potentially throughout the breeding season (when these birds were captured and

photographed). Since *Lonicera* fruits are mainly available in late summer and fall, carotenoids

producing bill color may come from different sources (e.g., insect prey), although fruits can ripen

as early as June (personal observation). However, the extent to which cardinal bill color changes

over time is unknown, and color measured during the summer breeding season may well have

resulted from carotenoids ingested and sequestered previously. Importantly, the factors

influencing color (e.g., carotenoid availability, body condition, stress) may vary across the

signals measured in this study.

CONCLUSION

Urbanization typically dulls or reduces the extent of color in birds (Sepp et al. 2018, Leveau

2019, Heinen-Kay et al. 2021). Here we show instead that both male and female urban cardinals

had significantly enhanced color of multiple carotenoid-based signals compared to rural birds.

We suggest this enhancement is potentially due to the widespread and stable availability of

carotenoids in exotic plants that thrive in disturbed areas, although there are other possible

explanations, and this relationship deserves additional investigation. If carotenoids are a limiting

resource in rural but not urban habitats, the reliability of these signals as honest indicators of

condition may deteriorate (Jones et al. 2010, Rodewald et al. 2011). In support of this idea, we

also found that urban birds were in worse mass-related body condition compared to rural birds,

suggesting a decoupling of condition and color. Together, these results underscore the complex

and subtle impacts of urbanization on wildlife. Not only can urbanization negatively affect

populations by destroying and fragmenting habitat, it may also disrupt communication systems

and potentially alter the evolutionary dynamics of signals.

LITERATURE CITED

Ardia, D. R., D. R. Broughton, and M. J. Gleicher (2010). Short-term exposure to testosterone propionate leads to rapid bill color and dominance changes in Zebra Finches. Hormones and Behavior 58:526–532.

Bartón, K. (2020). MuMIn: multi-model inference. R package version 1.43.17. https://CRAN.R-project.org/package=MuMIn

van den Berg, C. P., J. Troscianko, J. A. Endler, N. J. Marshall, and K. L. Cheney (2020). Quantitative Colour Pattern Analysis (QCPA): A comprehensive framework for the analysis of colour patterns in nature. Methods in Ecology and Evolution 11:316–332.

Borgmann, K. L., and A. D. Rodewald (2005). Forest restoration in urbanizing landscapes: interactions between land uses and exotic shrubs. Restoration Ecology 13:334–340.

Dalrymple, R. L., H. Flores-Moreno, D. J. Kemp, T. E. White, S. W. Laffan, F. A. Hemmings, T. D. Hitchcock, and A. T. Moles (2018). Abiotic and biotic predictors of macroecological patterns in bird and butterfly coloration. Ecological Monographs 88:204–224.

Derryberry, E. P., and D. Luther (2021). What is known—and not known—about acoustic communication in an urban soundscape. Integrative and Comparative Biology 61:1783–1794.

Elmqvist, T., E. Andersson, N. Frantzeskaki, T. McPhearson, P. Olsson, O. Gaffney, K. Takeuchi, and C. Folke (2019). Sustainability and resilience for transformation in the urban century. Nature Sustainability 2:267–273.

Giraudeau, M., A. Chavez, M. B. Toomey, and K. J. McGraw (2015). Effects of carotenoid supplementation and oxidative challenges on physiological parameters and carotenoid-based coloration in an urbanization context. Behavioral Ecology and Sociobiology 69:957–970.

Giraudeau, M., M. B. Toomey, P. Hutton, and K. J. McGraw (2018). Expression of and choice for condition-dependent carotenoid-based color in an urbanizing context. Behavioral Ecology 29.

Goodwin, T. W. (1984). The biochemistry of the carotenoids. Second. Chapman and Hall, New York.

Grunst, M. L., A. S. Grunst, R. Pinxten, L. Bervoets, and M. Eens (2020). Carotenoid- but not melanin-based plumage coloration is negatively related to metal exposure and proximity to the road in an urban songbird. Environmental Pollution 256:113473.

Hasegawa, M., R. A. Ligon, M. Giraudeau, M. Watanabe, and K. J. McGraw (2014). Urban and colorful male House Finches are less aggressive. Behavioral Ecology 25:641–649.

Heinen-Kay, J. L., A. D. Kay, and M. Zuk (2021). How urbanization affects sexual
communication. Ecology and Evolution 11:17625–17650.

Hill, G. E. (1991). Plumage coloration is a sexually selected indicator of male quality. Nature
350:337–339.

Hobbs, R. J. (2000). Land-use changes an invasions. In Invasive species in a changing world.
Island Press, Washington, DC, pp. 55–64.

Hudon, J., D. Derbyshire, S. Leckie, and T. Flinn (2013). Diet-induced plumage erythrism in
Baltimore Orioles as a result of the spread of introduced shrubs. The Wilson Journal of
Ornithology 125:88–96.

Hudon, J., R. J. Driver, N. H. Rice, T. L. Lloyd-Evans, J. A. Craves, and D. P. Shustack (2017).
Diet explains red flight feathers in Yellow-shafted Flickers in eastern North America.
The Auk 134:22–33.

Hudon, J., and R. Mulvihill (2017). Diet-induced plumage erythrism as a result of the spread of
alien shrubs in North America. North American Bird Bander 42:95–103.

Jawor, J. M., and R. Breitwisch (2004). Multiple ornaments in male Northern Cardinals,
Cardinalis cardinalis, as indicators of condition. Ethology 110:113–126.

Jawor, J. M., N. Gray, S. M. Beall, and R. Breitwisch (2004). Multiple ornaments correlate with
aspects of condition and behaviour in female Northern Cardinals, *Cardinalis cardinalis*.
Animal Behaviour 67:875–882.

Jawor, J. M., S. U. Linville, S. M. Beall, and R. Breitwisch (2003). Assortative mating by
multiple ornaments in Northern Cardinals (*Cardinalis cardinalis*). Behavioral Ecology
14:515–520.

Jiménez-Peñuela, J., M. Ferraguti, J. Martínez-de la Puente, R. Soriguer, and J. Figuerola (2019).
Urbanization and blood parasite infections affect the body condition of wild birds.
Science of The Total Environment 651:3015–3022.

Jones, T. M., A. D. Rodewald, and D. P. Shustack (2010). Variation in plumage coloration of
Northern Cardinals in urbanizing landscapes. The Wilson Journal of Ornithology
122:326–333.

Leston, L. F. V., and A. D. Rodewald (2006). Are urban forests ecological traps for understory
birds? An examination using Northern cardinals. Biological Conservation 131:566–574.

Leveau, L. M. (2019). Urbanization induces bird color homogenization. Landscape and Urban
Planning 192:103645.

Linville, S. U., and R. Breitwisch (1997). Carotenoid availability and plumage coloration in a
wild population of Northern Cardinals. The Auk 114:796–800.

Linville, S. U., R. Breitwisch, and A. J. Schilling (1998). Plumage brightness as an indicator of
460    parental care in Northern Cardinals. Animal Behaviour 55:119–127.

461 Maia, R., H. Gruson, J. A. Endler, and T. E. White (2019). pavo 2: New tools for the spectral and
462    spatial analysis of colour in R. Methods in Ecology and Evolution 10:1097–1107.

463 McGraw, K. J., and G. E. Hill (2001). Carotenoid access and intraspecific variation in plumage
464    pigmentation in male American Goldfinches (*Carduelis tristis*) and Northern Cardinals
465    (*Cardinalis cardinalis*). Functional Ecology 15:732–739.

466 Mulvihill, R. S., K. C. Parkes, R. C. Leberman, and D. S. Wood (1992). Evidence supporting a
467    dietary basis for orange-tipped rectrices in the Cedar Waxwing. Journal of Field
468    Ornithology 63:212–216.

469 R Core Team (2019). R: a language and environment for statistical computing. R Foundation for
470    Statistical Computing, Vienna, Austria.

471 Rodewald, A. D. (2005). Interaction between exotic shrubs and breeding birds in riparian forests.
472    Proceedings of the Ohio invasive plan research conference: bridging the gap between
473    land management and research 196:43–48.

474 Rodewald, A. D., and D. P. Shustack (2008). Consumer resource matching in urbanizing
475    landscapes: are synanthropic species over-matching? Ecology 89:515–521.

476 Rodewald, A. D., D. P. Shustack, and T. M. Jones (2011). Dynamic selective environments and
477    evolutionary traps in human-dominated landscapes. Ecology 92:1781–1788.

478 Rosen, R. F., and K. A. Tarvin (2006). Sexual signals of the male American Goldfinch. Ethology
479    112:1008–1019.

480 Salleh Hudin, N., D. Strubbe, A. Teyssier, L. De Neve, J. White, G. P. J. Janssens, and L. Lens
481    (2016). Predictable food supplies induce plastic shifts in avian scaled body mass.
482    Behavioral Ecology 27:1833–1840.

483 Schneider, C. A., W. S. Rasband, and K. W. Eliceiri (2012). NIH Image to ImageJ: 25 years of
484    image analysis. Nature Methods 9:671–675.

485 Seddon, N., G. M. Mace, S. Naeem, J. A. Tobias, A. L. Pigot, R. Cavanagh, D. Mouillot, J.
486    Vause, and M. Walpole (2016). Biodiversity in the Anthropocene: prospects and policy.
487    Proceedings of the Royal Society B: Biological Sciences 283:20162094.

488 Sepp, T., K. J. McGraw, and M. Giraudeau (2020). Urban sexual selection. In Urban
489    evolutionary biology. Oxford University Press.

490 Sepp, T., K. J. McGraw, A. Kaasik, and M. Giraudeau (2018). A review of urban impacts on
491    avian life-history evolution: Does city living lead to slower pace of life? Global Change
492    Biology 24:1452–1469.

Shannon, G., M. F. McKenna, L. M. Angeloni, K. R. Crooks, K. M. Fristrup, E. Brown, K. A.
Warner, M. D. Nelson, C. White, J. Briggs, S. McFarland, and G. Wittemyer (2016). A
synthesis of two decades of research documenting the effects of noise on wildlife: effects
of anthropogenic noise on wildlife. Biological Reviews 91:982–1005.

Slabbekoorn, H. (2013). Songs of the city: noise-dependent spectral plasticity in the acoustic
phenotype of urban birds. Animal Behaviour 85:1089–1099.

Slabbekoorn, H., and A. den Boer-Visser (2006). Cities change the songs of birds. Current
Biology 16:2326–2331.

Slabbekoorn, H., R. Dooling, A. N. Popper, and R. R. Fay (2018). Effects of anthropogenic noise
on animals. Springer, New York.

Szulkin, M., J. Munshi-South, and A. Charmantier (Editors) (2020). Urban evolutionary biology.
Oxford University Press, Oxford, UK.

Troscianko, J., and M. Stevens (2015). Image calibration and analysis toolbox – a free software
suite for objectively measuring reflectance, colour and pattern. Methods in Ecology and
Evolution 6:1320–1331.

Vilà, M., and I. Ibáñez (2011). Plant invasions in the landscape. Landscape Ecology 26:461–472.

Vorobyev, M., and D. Osorio (1998). Receptor noise as a determinant of colour thresholds.
Proceedings of the Royal Society of London. Series B: Biological Sciences 265:351–358.

Wolfenbarger, L. L. (1999a). Female mate choice in Northern Cardinals (*Cardinalis cardinalis*):
is there a preference for redder males? Wilson Bulletin 111:76–83.

Wolfenbarger, L. L. (1999b). Red coloration of male Northern Cardinals correlates with mate
quality and territory quality. Behavioral Ecology 10:80–90.

Wright, S., and H. B. Fokidis (2016). Sources of variation in plasma corticosterone and
dehydroepiandrosterone in the male Northern Cardinal (*Cardinalis cardinalis*): II. Effects
of urbanization, food supplementation and social stress. General and Comparative
Endocrinology 235:201–209.

Figure 1. A map of the rural (Rice Creek Field Station, **A**) and urban (Barry Park, **B**) field sites.

Figure 2. Variation in male chest hue (**A**), saturation (**B**), and brightness (**C**) as a function of

urbanization and/or year. There was no significant effect of site on saturation or brightness, so

only the effect of year is shown. Lowercase letters indicate which factors are (different letter) or

are not (same letter) significantly different from each other.

Figure 3. Variation in male bill hue (**A**) and female wing saturation (**B**) as a function of

urbanization. Asterisks separate factors that are significantly different.

Figure 4: Pairwise chromatic contrast between rural and urban male chest (2020 only), male bill,

and female wing (**A**). The dashed line indicates dS = 1, or the perceptual threshold above which

colors can be distinguished by the avian visual system. Photos exemplifying the average rural

and urban male chest (**B**, **C**), male bill (**D**, **E**), and female wing (**F**, **G**) signals.

576   **Figure 5.** Variation in male body condition score (**A**) and female mass (**B**) as a function of

577   urbanization.

586
587

SUPPLEMENTARY MATERIAL

591　**Table S1.** Summary of model selection results examining male chest hue, saturation, and
592　brightness ($n = 65$). Only models with ΔAICc < 5 are shown.

Response	Factors	k	logLik	ΔAICc	Weight
hue	site*year	7	170.22	0	0.86
hue	year	4	164.76	3.63	0.14
saturation	year	4	173.58	0	0.66
saturation	site+year	5	173.58	2.35	0.21
saturation	site*year	7	175.62	3.23	0.13
brightness	year	4	175.87	0	0.58
brightness	site*year	7	178.50	2.05	0.21
brightness	site+year	5	176.01	2.07	0.21

596　**Table S2.** Summary of model selection results examining male bill hue, saturation, and
597　brightness ($n = 61$). Only models with ΔAICc < 5 are shown.

Response	Factors	k	logLik	ΔAICc	Weight
hue	site	3	120.41	0	0.50
hue	site+year	5	122.53	0.43	0.40
hue	site*year	7	123.61	3.29	0.10
saturation	site	3	115.65	0	0.29
saturation	intercept	2	114.51	0.07	0.28
saturation	site+year	5	117.69	0.58	0.22
saturation	year	4	116.48	0.63	0.21
brightness	intercept	2	116.29	0	0.39
brightness	year	4	118.24	0.62	0.29
brightness	site	3	116.68	1.44	0.19
brightness	site+year	5	118.62	2.24	0.13

611 **Table S3.** Summary of model selection results examining female bill hue, saturation, and
612 brightness ($n = 25$). Only models with ΔAICc < 5 are shown.
613

Response	Factors	k	logLik	ΔAICc	Weight
hue	intercept	2	47.09	0	0.50
hue	year	4	49.10	1.43	0.25
hue	site	3	47.35	2.07	0.18
hue	site+year	5	49.41	3.98	0.07
saturation	year	4	50.13	0	0.43
saturation	intercept	2	47.29	0.23	0.38
saturation	site	3	47.30	2.81	0.10
saturation	site+year	5	50.15	3.12	0.09
brightness	year	4	46.56	0	0.45
brightness	intercept	2	43.60	0.47	0.36
brightness	site	3	43.60	3.07	0.10
brightness	site+year	5	46.56	3.15	0.09

614
615
616 **Table S4.** Summary of model selection results examining female wing hue, saturation, and
617 brightness ($n = 24$). Only models with ΔAICc < 5 are shown.
618

Response	Factors	k	logLik	ΔAICc	Weight
hue	intercept	2	42.30	0	0.51
hue	site	3	42.93	1.37	0.26
hue	year	4	43.97	2.21	0.17
hue	site+year	5	44.57	4.22	0.06
saturation	site	3	52.73	0	0.75
saturation	intercept	2	50.34	2.15	0.25
brightness	year	4	27.03	0	0.33
brightness	intercept	2	24.21	0.11	0.31
brightness	site+year	5	28.10	1.09	0.19
brightness	site	3	24.94	1.27	0.17

Figure S1. Examples of reflectance curves for the signals measured in the study. The dark line represents average reflectance and the shaded area represents one standard deviation. The male chest curve was generated from ten random males measured in the study. The male bill, female bill, and female wing curves were generated from ten measurements of museum skins. All signals reflect predominantly long wavelengths and have minimal ultraviolet (UV) reflectance.

Figure S2. Examples representing the analysis of male and female bill (**A**) and female wing (**B**) photos. After selecting regions of interest, we used the QCPA framework to break the color patch up into discernable clusters (**C**, **D**) that could be measured for hue, saturation, and brightness.