Community diversity is associated with intra-species genetic diversity and gene loss in the human gut microbiome

Naïma Madi[1], Daisy Chen[2,3,^], Richard Wolff[4,^], B. Jesse Shapiro[1,5,6,7,*], and Nandita Garud[4,8,*]

1. Département de sciences biologiques, Université de Montréal, Canada;
2. Computational and Systems Biology, University of California, Los Angeles
3. Bioinformatics and Systems Biology Program, University of California, San Diego▢
4. Department of Ecology and Evolutionary Biology, University of California, Los Angeles
5. Department of Microbiology and Immunology, McGill University, Canada;
6. McGill Genome Centre, McGill University, Canada
7. Quebec Centre for Biodiversity Science, Canada
8. Department of Human Genetics, University of California, Los Angeles

* Correspondence to jesse.shapiro@mcgill.ca and ngarud@ucla.edu. These authors contributed equally.

^ These authors contributed equally.

Abstract

The human gut microbiome contains a diversity of microbial species that varies in composition over time and across individuals. These species (and strains within species) can migrate across hosts and evolve by mutation and recombination within hosts. How the ecological process of community assembly interacts with intra-species diversity and evolutionary change is a longstanding question. Two contrasting hypotheses have been proposed based on ecological observations and theory: Diversity Begets Diversity (DBD), in which taxa tend to become more diverse in already diverse communities, and Ecological Controls (EC), in which higher community diversity impedes diversification within taxa. Previously, using 16S rRNA gene amplicon data across a range of environments, we showed a generally positive relationship between taxa diversity and community diversity at higher taxonomic levels, consistent with the predictions of DBD (Madi et al., 2020). However, this positive 'diversity slope' reaches a plateau at high levels of community diversity. Here we show that this general pattern holds at much finer genetic resolution, by analyzing intra-species strain and nucleotide variation in static and temporally sampled shotgun-sequenced fecal metagenomes from cohorts of healthy human hosts. We find that both intra-species polymorphism and strain number are positively correlated with community Shannon diversity. This trend is consistent with DBD, although we cannot exclude abiotic drivers of diversity. Shannon diversity is also predictive of increases in polymorphism over time scales up to ~4-6 months, after which the diversity slope flattens and then becomes negative—consistent with DBD eventually giving way to EC. Also supporting a complex mixture of DBD and EC, the number of strains per focal species is positively associated with Shannon diversity but negatively associated with richness. Finally, we show that higher community diversity predicts gene loss in a focal species at a future time point. This observation is broadly consistent with the Black Queen Hypothesis, which posits that genes with functions provided by the community are less likely to be retained in a focal species' genome. Together, our results show that a mixture of DBD, EC, and Black Queen may operate simultaneously in the human gut microbiome, adding to a growing body of evidence that these eco-evolutionary processes are key drivers of biodiversity and ecosystem function.

Introduction

Our understanding of microbial evolution and diversification has been enriched by experimental studies of bacterial isolates in the laboratory, but it remains a challenge to study evolution in the context of more complex communities (Lenski, 2017). Ongoing advances in culture-independent technologies have allowed us to study bacteria in the complex and dense communities in which they naturally occur (Garud and Pollard, 2020). Within a community, individual players engage in many negative and positive ecological interactions. Negative interactions can originate from competition for resources and biomolecular warfare (Hibbing et al., 2010; Mitri and Foster, 2013), while positive interactions can stem from secreted metabolites that are used by other members of the community (cross-feeding) (Venturelli et al., 2018). These ecological interactions can create new niches and selective pressures, leading to eco-evolutionary feedbacks whose nature are yet to be fully understood.

Ecological interactions can yield positive or negative effects on the diversification of a focal species. Under the "Diversity Begets Diversity" (DBD) hypothesis, higher levels of community diversity increase the rate of speciation (or diversification, more generally) due to positive feedback mechanisms such as niche construction (Calcagno et al., 2017; Schluter and Pennell, 2017). Competition for limited niche space could also drive DBD if species diversify into new niches to avoid competition (Meyer and Kassen, 2007; Mitri and Foster, 2013; Schluter, 2000). By contrast, the "Ecological Controls" (EC) hypothesis posits that competition for a limited number of niches at high levels of community diversity results in a negative effect on further diversification. Metabolic models predict that DBD may initially spur diversification due to cross-feeding, but the diversification rate eventually slows and reaches a plateau as metabolic niches are filled (San Roman and Wagner, 2021). These theoretical predictions are largely supported by our previous study involving 16S rRNA gene amplicon sequencing data from the Earth Microbiome Project, in which we observed a generally positive relationship (which we call the diversity slope; **Figure 1**) between community diversity and focal-taxon diversity at most taxonomic levels, reaching a plateau at the highest levels of diversity (Madi et al., 2020).

In this previous study, we found stronger support for DBD in the animal gut relative to more diverse microbiomes such as soils and sediments, which were closer to a plateau of diversity

(Madi et al., 2020). While diversity slopes were generally positive at taxonomic levels as fine as

amplicon sequence variants (akin to species or strains) within a genus, they were most positive at

higher levels such as classes or phyla. A recent experiment on soil bacteria also found evidence

of DBD at the family level, likely driven by niche construction and metabolic cross-feeding

(Estrela et al., 2022). It therefore remains unclear if the predictions of DBD hold primarily at

these higher taxonomic levels, involving the ecological process of community assembly, or if

they also apply at the finer intra-species level. Within-host intra-species diversity can arise by

co-colonization of a host by genetically distinct strains belonging to the same species or

evolutionary diversification of a lineage via *de novo* mutation and gene gain/loss events within a

host.

Such fine-scale strain-level variation has important functional and ecological consequences;

among other things, strains are known to engage in interactions that cannot be predicted from

their species identity alone (Goyal et al., 2022). Although closely-related bacteria are expected to

have broadly similar niche preferences, finer-scale niches may differ below the species level

(Martiny et al., 2015). For example, the acquisition of a carbohydrate-active enzyme by

Bacteroides plebeius allows it to exploit a new dietary niche in the guts of people consuming

nori (seaweed) (Hehemann et al., 2010), and single nucleotide adaptations permit *Enterococcus*

gallinarum translocation across the intestinal barrier resulting in inflammation (Yang et al.,

2022). Despite their potential phenotypic effects, it is unknown if such fine-scale genetic changes

are favored by higher community diversity (due for example to niche construction, as predicted

by DBD) or suppressed (due to competition for limited niche space, as predicted by EC).

Competition could also lead to DBD if focal species evolve new niche preferences to avoid

extinction (Mitri and Foster, 2013; Schluter, 2000) – an idea with some support in experimental

microcosms (Meyer and Kassen, 2007) but largely unexplored in natural communities.

Here, we investigate the relationship between intra-species genetic diversity and community

diversity in the human gut microbiome, a well-studied system in which we previously found

support for DBD at higher taxonomic levels. We use static and temporal shotgun metagenomic

data from a large panel of healthy adult hosts from the Human Microbiome Project (Lloyd-Price

et al., 2017; The Human Microbiome Project Consortium, 2012) as well as from four healthy

individuals sampled almost daily over the course of one year (Poyet et al., 2019). Using
metagenomic data allows us to track change in single nucleotide variation, strain diversity, and
gene gain or loss events within relatively abundant species in the microbiome, and study how
these measures of intra-species diversity are associated with community diversity. Although such
analyses of natural diversity cannot fully control for unmeasured confounding environmental
factors, they are an important complement to controlled experimental and theoretical studies
which lack real-world complexity.

Figure 1. Diversity Begets Diversity (DBD) and Ecological Controls (EC) hypotheses illustrated.
The top panel shows patterns predicted by alternative hypotheses and the bottom panel illustrates possible
underlying mechanisms, including strain invasion and *de novo* mutation within a focal species. Under
DBD, high community diversity is associated with high focal species polymorphism (red line) yielding a
positive diversity slope. In the bottom panels, the bacterial community within a host (large circle) is
represented as small rectangles with different bacterial species in different colors. The focal species is
shown in light green outlined in black and the invading strain (striped colors) is a different strain of the
resident focal species.

Results

We investigated the relationship between community diversity and within-species genetic diversity in human gut microbiota using two shotgun metagenomic datasets. First, we analyzed data from a panel of 249 healthy hosts (Lloyd-Price et al., 2017; The Human Microbiome Project Consortium, 2012), in which stool samples were collected 1-3 times from each host at approximately 6-month intervals. Second, we analyzed data from four individuals sampled more densely over the course of ~18 months (Poyet et al., 2019). In both cases, we only consider intra-species diversity of relatively abundant species that are well sampled in these metagenomic datasets (Methods).

We examined several metrics of community diversity and intra-species diversity and calculated the slope of their relationship, defined as the diversity slope (**Figure 1**). We note that intra-species diversity can arise within hosts via *de novo* point mutation, gene gain or loss, or the coexistence of genetically distinct strains that diverged before colonizing the host. To quantify community diversity, we calculated Shannon diversity and richness at the species level. Shannon diversity is relatively insensitive to sampling effort (Madi et al., 2020; Walters and Martiny, 2020) but richness can be underestimated in low sample sizes. We therefore computed richness on data rarefied to an equal number of reads per sample, yielding generally similar results to unrarefied data (described below). In all cases, we included the number of reads per sample (coverage) as a covariate in our models, as this could affect estimates of both community diversity and intra-species diversity. To quantify intra-species diversity, we used a reference genome-based approach to call single nucleotide variants (SNVs) and gene copy number variants (CNVs) within each focal species and computed polymorphism rates, measured as the fraction of synonymous nucleotide sites in a species' core genome with intermediate allele frequencies (between 0.2 and 0.8) within a host (Methods). We also repeated the analysis on nonsynonymous sites, as these are subject to stronger selective constraints. As an additional metric of intra-species diversity, we inferred the number of strains within each species using StrainFinder applied to all polymorphic sites (including those outside the 0.2-0.8 frequency range) (Smillie et al., 2018).

Community diversity is positively associated with intra-species polymorphism in the human gut microbiome

As an exploratory visualization, we began by plotting the relationship between community diversity and intra-species polymorphism rate calculated at synonymous sites in cross-sectional HMP metagenomes for the nine most prevalent species (**Figure 2A,B**). The slope of this relationship (the diversity slope; **Figure 1**) provides an indicator of the evidence for DBD (positive slope) or EC (flat or negative slope). The relationship between polymorphism rate and community diversity was mostly positive in the top nine most prevalent species in HMP hosts (**Figure 2A,B**). These nine species are used as a simple illustration of the diversity slope, not as a formal hypothesis-testing framework.

Figure 2. Positive association between community diversity and within-species polymorphism in cross-sectional Human Microbiome Project samples. (A) Scatter plots showing the relationship between community Shannon diversity and within-species polymorphism rate (estimated at synonymous sites) in the nine most prevalent species in HMP. (B) Scatter plots showing the relationship between species richness and within-species polymorphism rate in the nine most prevalent species in HMP. These are simple correlations to show the relationships in the raw data. Significant correlations are shown with red trendlines (Spearman correlation, $P<0.05$); non-significant trendlines are in gray. **Results of generalized additive models (GAMs)** predicting polymorphism rate in a focal species as a function of (C) Shannon diversity, (D) species richness estimated on all sequence data, and (E) species richness estimated on rarefied sequence data. GAMs are based on data from 69 bacterial species across 249 HMP stool donors. Adjusted R^2 and Chi-square P-values corresponding to the predictor effect are displayed in each panel. Shaded areas show the 95% confidence interval of each model prediction. See Supplementary File 1a and supplementary file 2 section 1 for detailed model outputs.

To generalize across species and to formally test the predictions of DBD, we fit generalized additive models (GAMs) to the HMP data. Using GAMs, we are able to model non-linear relationships evident in the data (**Figure 2A,B**), account for random variation in the strength of the diversity slope across bacterial species, and account for the uneven number of samples per host and the non-independence of samples from the same host (Methods; see **Supplementary File 1a** and **Supplementary File 2 section 1** for additional model details). These GAMs included 69 focal species with sufficient coverage to quantify within-species polymorphism (Methods); the results therefore apply to relatively abundant species in the human gut microbiome. GAMs showed an overall positive association between within-species polymorphism and Shannon diversity (**Fig 2C**, GAM, $P=0.031$, Chi-square test) as well as between within-species polymorphism and community richness after controlling for coverage as a covariate (**Fig 2D**, GAM, $P=0.017$, Chi-square test) or rarefying samples to an equal number of reads (**Fig 2E**, GAM, $P=2.63e-04$, Chi-square test). The random effect of species identity is highly significant in all models, indicating that each bacterial species has its own characteristic diversity slope (**Supplementary File 1a**). It appears that synonymous polymorphism reaches a plateau at high levels of community richness, which is particularly evident when using rarefied data (**Fig 2E**). Using the same GAMs applied to nonsynonymous polymorphism, we found no significant associations between diversity and within-species polymorphism rate (GAM, $P>0.05$, Chi-square test) (**Supplementary File 1b, Supplementary File 2 section 4**). This could be due to lower statistical power, since there are fewer nonsynonymous than synonymous sites, or could reflect a true difference in the diversity slope between these site categories.

These generally positive correlations between focal species polymorphism and species-level
measures of community diversity also hold when community diversity is measured at higher
taxonomic levels; specifically, synonymous polymorphism rate was significantly positively
associated with Shannon diversity calculated at the genus and family levels (GAMs, $P<0.05$,
Chi-square test) (**Figure 2-figure supplement 1, Supplementary File 1c**). However,
synonymous polymorphism rate was not significantly associated with Shannon diversity
calculated at the highest taxonomic levels (order, class and phylum, GAMs, $P>0.05$, Chi-square
test). The positive correlation between polymorphism rate and richness held at all taxonomic
levels (GAMs, $P<0.05$, Chi-square test) (**Figure 2-figure supplement 1, Supplementary File
1c**, **Supplementary File 2 section 2 and 3**). When estimated at nonsynonymous sites,
polymorphism rate was not significantly correlated with Shannon diversity at any taxonomic
level (GAMs, $P>0.05$, Chi-square test), but was positively correlated with richness at the highest
levels (phyla, class and order, $P=3e-04$, $P=0.017$ and $P=6.11e-04$ respectively, Chi-square test
from GAMs) (**Figure 2-figure supplement 2**, **Supplementary File 1d**, **Supplementary File 2
section 5 and 6**). Even when not statistically significant, the diversity slopes were generally
positive at all taxonomic levels for both synonymous and nonsynonymous polymorphism
(**Figure 2-figure supplements 1 and 2**). Overall, these results are consistent with the predictions
of DBD at most taxonomic levels. However, slightly different relationships are observed when
considering different measures of community diversity (Shannon or richness) and different
components of within-species diversity (nonsynonymous or synonymous).
223

Different measures of community diversity have contrasting associations with intra-species
strain diversity
Within host polymorphism rates span several orders of magnitude (10^{-5}/bp to 10^{-2}/bp), largely
due to the fact that strain content is variable across hosts. As previously argued (Garud et al.,
2019), with conservatively high estimates for mutation rate ($\mu \sim 10^{-9}$) (Sung et al., 2012),
generation times (~ 10 / day) (Poulsen et al., 1995), and time since colonization (<100 years),
polymorphism rates of ~10^{-2}/bp or more are inconsistent with within-host diversification of a
single colonizing lineage. Therefore, hosts with relatively high intra-host polymorphism rates are
likely colonized by mixtures of multiple strains that diverged long before colonizing a host.
Moreover, recent work suggests that the numbers and genetic composition of strains colonizing a

host can vary from host to host (Garud et al., 2019; Olm et al., 2017; Russell and Cavanaugh,
2017; Truong et al., 2017; Verster et al., 2017). The associations between polymorphism and
community diversity (**Figure 2**) are likely driven by a combination of *de novo* mutation and co-
colonization by multiple strains.
238
To separate these two sources of diversity and to explicitly account for the strain structure within
hosts, we inferred the number of strains per focal species with StrainFinder (Smillie et al., 2018)
(Methods) and used strain number as another quantifier of intra-species diversity. We found that
the number of strains per focal species follows an approximately linear relationship with
community diversity in the nine most prevalent species in HMP (**Figure 3A, B**). Because of
these approximately linear relationships, we used generalized linear mixed models (GLMMs) to
investigate the relationship between the number of strains per focal species and community
diversity, while taking into account coverage per sample as a covariate and variation between
species, hosts and samples as random effects (Methods). The number of strains per focal species
was positively correlated with community Shannon diversity (GLMM, *P*=3.58e-07, likelihood
ratio test (LRT)) (**Fig 3C**, **Supplementary File 1e**, **Supplementary File 2 section 7.1**). This
suggests that the positive correlation between polymorphism rate and Shannon diversity **(Figure
2**) is due at least in part to strain diversity.

Figure 3. Associations between community diversity and strain number in cross-sectional Human Microbiome Project samples. (A) Scatter plots showing the relationship between Shannon diversity and the inferred number of strains within each of the nine most prevalent species in HMP. (B) Scatter plots showing the relationship between species richness and the inferred number of strains within each of the nine most prevalent species in HMP. Significant linear correlations are shown with red trendlines (Pearson correlation, *P*<0.05); non-significant trend lines are in gray. **Results of generalized linear mixed models (GLMMs)** predicting strain count in a focal species as a function of (C) Shannon diversity, (D) species richness estimated on all data, and (E) species richness estimated on rarefied sequence data. Diversity estimates (x-axis) are standardized to zero mean and unit variance in the models. The Y-axis shows the mean number of strains per focal species predicted by the GLMM. GLMMs are based on data from 184 bacterial species across 249 HMP stool donors. *P*-values (likelihood ratio test) are displayed in each panel. Shaded areas show the 95% confidence interval of each model prediction. See **Supplementary File 1e and Supplementary File 2 section 7** for detailed model outputs.

By contrast, species richness was negatively correlated with strain number (GLMM, *P*=1.67e-06, LRT) (**Fig 3D**, **Supplementary File 1e**, **Supplementary File 2 section 7.2**). The negative relationship with richness was unlikely to be confounded by sequencing depth, since the same result was obtained using rarefied data (**Fig 3E**, **Supplementary File 1e**, **Supplementary File 2**

section 7.3). The negative strain number-richness relationship also held at all other taxonomic

ranks (GLMM, *P*<0.05, LRT), while the strain number-Shannon diversity relationship was

generally positive (**Fig 3-Figure supplement 1, Supplementary File 1f, Supplementary File 2**

section 8-9). These effects also appear to be species-specific: for example, the number of

Bacteroides vulgatus strains per host is positively correlated with both Shannon diversity and

richness (consistent with DBD predictions) whereas *B. ovatus* has no relationship with Shannon

diversity but a negative correlation with richness (consistent with EC; **Fig 2A, B**). Together,

these results reveal that different components of community diversity can have contrasting

effects on the diversity slope.

Community Shannon diversity is a predictor of intra-species polymorphism and gene loss

in time series data

Our analyses thus far have considered only individual time points, which represent static

snapshots of the dynamic processes of community assembly and evolution in the microbiome. To

interrogate these phenomena over time, we analyzed 160 HMP subjects who were sampled 2-3

times ~6 months apart. Under a DBD model, we expect community diversity at an earlier time

point to result in higher within-species polymorphism at a future time point. To test this

expectation, we defined 'polymorphism change' as the difference between polymorphism rates at

the two time points (Methods). We also investigated the effects of community diversity on gene

loss and gain events within a focal species, as such changes in gene content are known to occur

frequently within host gut microbiomes (Garud et al., 2019; Groussin et al., 2021; Yaffe and

Relman, 2020; Zhao et al., 2019). Here a gene was considered absent if its copy number (*c*) was

<0.05 and present if $0.6 \leq c \leq 1.2$. As in the cross-sectional analyses above, we also controlled

for sequencing depth of the sample and excluded genes with aberrant coverage or presence in

multiple species (Methods).

In HMP samples, polymorphism change showed no significant relationships with community

diversity at the earlier time point, whether it was estimated with Shannon index or species

richness (GAM, *P*>0.05) (**Supplementary File 2 section 10.1**). These results suggest that DBD

is negligible or undetectable over ~6-month time lags in the human gut. By contrast, we found

that gene loss in a focal species between two consecutive time points was positively correlated

with community diversity at the earlier time point (**Figure 4;** GLMM, *P*=0.028, *P*=0.034 and
P=0.049, LRT for Shannon, richness and rarefied richness respectively) (**Supplementary File**
1g, **Supplementary file 2 section 10.3**). Gene gains did not show any significant relationships
with community diversity (GLMM, *P*>0.05). Selection for gene loss in more diverse
communities is a prediction of the Black Queen Hypothesis (BQH), provided that higher
community diversity results in more redundant gene functions that compensate for losses in a
focal species (Morris et al., 2012). Most species in HMP samples lost fewer than ten genes over
~6 months – consistent with *de novo* deletion events of a few genes – but occasionally hundreds
of genes were lost from a host, suggesting that strains with smaller genomes were selected in
more diverse communities (**Figure 4A**, **4B**).

Figure 4. Positive association between community diversity and gene loss in Human Microbiome Project time
series. (A) Scatter plots showing the relationship between Shannon diversity at time point 1 (tp1) and gene loss
between tp1 and tp2 within each of the nine most prevalent species in HMP. (B) Scatter plots showing the

relationship between species richness at tp1 and gene loss between tp1 and tp2 within each of the nine most
prevalent species in HMP. Significant linear correlations are shown with red trendlines (Pearson correlation,
P<0.05); non-significant trend lines are in gray. The Y-axis is plotted on a log10 scale for clarity. **Results of**
generalized linear mixed models (GLMMs) predicting gene loss in a focal species as a function of (C) Shannon
diversity, (D) species richness estimated on all data, and (E) species richness estimated on rarefied sequence data. *P*-
values (likelihood ratio test) are displayed in each panel. Shaded areas show the 95% confidence interval of each
model prediction. The Y-axis is plotted on the link scale, which corresponds to log for negative binomial GLMMs
with a count response. GLMMs are based on data from 54 bacterial species across 154 HMP stool donors sampled at
more than one time point. See Supplementary file 1g and Supplementary File 2 section 10 for detailed model
outputs.
325
326
To study these dynamics at higher temporal resolution, we analyzed shotgun metagenomic data

from four more frequently sampled healthy individuals from a previous study (Poyet et al.,

2019). Stool from donor *am* was sequenced over 18 months with a median of one day between

samples; *an* over 12 months (median 2 days between samples); *ao* over 5 months (median 1 day

between samples); and *ae* over 7 months (median 2 days between samples). In this data, we

tracked both polymorphism change and gene gains and losses between two successive time

points in 15 species with a minimal marker gene coverage of 10 in at least ten samples. These

include seven species of *Bacteroides*, two *Eubacterium,* two *Faecalibacterium*, two

Ruminococcus, as well as *Alistipes putredinis* and *Parabacteroides merdae*.

Using the Poyet dataset, we asked whether community diversity in the gut microbiome at one

time point could predict polymorphism change at a future time point by fitting GAMs with the

change in polymorphism rate as a function of the interaction between community diversity at the

first time point and the number of days between the two time points. Shannon diversity at the

earlier time point was correlated with increases in polymorphism (consistent with DBD) up to

~150 days (~4.5 months) into the future (**Fig 5-Figure supplement 1**), but this relationship

became weaker and then inverted (consistent with EC) at longer time lags (**Fig 5A**,

Supplementary File 1h, GAM, *P*=0.023, Chi-square test). The diversity slope is approximately

flat for time lags between four and six months, which could explain why no significant

relationship was found in HMP, where samples were collected every ~6 months. No relationship

was observed between community richness and changes in polymorphism (**Supplementary File**

1h, GAM, *P*>0.05).

We next asked if community diversity at one time point could predict gene gains or losses at

future time points by fitting GLMMs (analogous to the GAMs above, but more appropriate for

gain/loss count data). Our method does not explicitly distinguish between gene gain/loss arising
from recombination or deletion versus replacement of strains with different gene content. We
found that community Shannon diversity predicted future gene loss in a focal species, and this
effect became stronger with longer time lags (**Fig 5B**, **Supplementary File 1i**, GLMM, *P*=0.006,
LRT for the effect of the interaction between the initial Shannon diversity and time lag on the
number of genes lost). The model predicts that increasing Shannon diversity from its minimum
to its maximum would result in the loss of 0.075 genes from a focal species after 250 days. In
other words, about one of the 15 focal species considered would be expected to lose a gene in
this time frame.

Higher Shannon diversity was also associated with fewer gene gains, and this relationship also
became stronger over time (**Fig 5C**, **Supplementary File 1i**, GLMM, *P*=1.11e-09, LRT). We
found a similar relationship between community species richness and gene gains, although the
relationship was slightly positive at shorter time lags (**Fig 5D**, **Supplementary File 1i**, GLMM,
P=3.41e-04, LRT). No significant relationship was observed between richness and gene loss
(**Supplementary File 1i,** GLMM, *P*>0.05). Taken together with the HMP results (**Fig 4**), these
longer time series reveal how the sign of the diversity slope can vary over time and how
community diversity is generally predictive of reduced focal species gene content.

371
Figure 5. Community diversity is associated with increases in focal species polymorphism over short time
lags and net gene loss in dense gut microbiome time series. (A) Results of a GAM predicting polymorphism
change in a focal species as a function of the interaction between Shannon diversity at the first time point and the
time lag (days) between two time points in data from Poyet et al. The response (Y-axis) was log transformed in the
Gaussian GAM. Results of GLMMs predicting (B) Number of genes lost and (C) Number of genes gained between

two time points in a focal species as a function of the interaction between Shannon diversity at the first time point
and the time lag between the two time points. (D) Results of the GLMM predicting the number of genes gained in a
focal species as a function of the interaction between rarefied species richness at the first time point and the time lag
between the two time points. The illustrated time lags correspond to the first quartile (50 days), the median (130
days), and the third quartile (250 days). See Supplementary Files 1h and i and Supplementary File 2 section 11 for
detailed model outputs. These analyses are based on data from 15 bacterial species across 4 stool donors from Poyet
et al. Only statistically significant relationships are plotted. Non-significant relationships are not shown: the GAM
predicting polymorphism change as a function of rarefied richness ($P>0.05$) and the GLMM predicting the number
of genes lost as a function of rarefied richness ($P>0.05$).

388 Discussion

How eco-evolutionary feedbacks shape biological communities is an open question that
to date has received substantial experimental and theoretical attention but is challenging to
address in nature. In our previous study using 16S rRNA amplicon sequences from the Earth
Microbiome Project, we found generally positive diversity slopes that eventually flattened at
high levels of community diversity (Madi et al., 2020). This pattern is generally consistent with
the predictions of DBD during the early stages of community assembly, but at later stages
becomes more consistent with EC as niches become filled. Based on the time series
metagenomic data analyzed here, the predictions of DBD also tend to hold over short time scales
but fail over longer time scales of several months. Whether this leads to a terminal plateau of
diversity, or whether ecological disturbances lead to cycles of DBD and EC, deserves further
study.

In our previous study, the animal gut microbiome had one of the highest positive
diversity slopes, making it an ideal candidate for investigating eco-evolutionary interactions at
greater intra-species resolution using metagenomic data. In this follow-up study, we investigate
the same phenomenon at a subspecies level, with results that are broadly consistent with the
predictions of DBD giving way to EC over long time scales. We note that experiments
supporting DBD have generally been conducted over short time scales ranging from two to 20
days (Estrela et al., 2022; Jousset et al., 2016), consistent with the importance of DBD early in
community assembly. We also identify several nuances and caveats to this general conclusion,
which are discussed below in detail.

Another recent study also found evidence for eco-evolutionary feedbacks in the HMP, in
the form of a positive relationship between evolutionary modifications or strain replacements in
a focal species and community diversity (Good and Rosenfeld, 2022). Using a model, they

further showed that these eco-evolutionary dynamics could be explained by resource competition
and did not require the cross-feeding interactions previously invoked (Estrela et al., 2022; San
Roman and Wagner, 2021, 2018) to explain DBD at higher taxonomic levels. This could be
because cross-feeding operates at the family- or genus- level and is less relevant at finer
evolutionary scales.

There are several noteworthy caveats to our study. First, using metagenomic data from
human microbiomes allowed us to study genetic diversity, but limited us to considering only
relatively abundant species with genomes that were well-covered by short sequence reads.
Deeper or more targeted sequencing may permit us to determine whether the same patterns hold
for rarer members of the microbiome. However, it is notable that the majority of the dozens of
species across the two datasets analyzed support DBD, suggesting that the phenomenon may
generalize.

Second, we cannot establish causal relationships without controlled experiments. We are
therefore careful to conclude that positive diversity slopes are consistent with the predictions of
DBD, and negative slopes with EC, but unmeasured environmental drivers could be at play. For
example, increased dietary diversity could simultaneously select for higher community diversity
and also higher intra-species diversity. In our previous study, we found that positive diversity
slopes persisted even after controlling for potential abiotic drivers such as pH and temperature
(Madi et al., 2020), but a similar analysis was not possible here due to a lack of metadata.
Neutral processes can account for several ecological patterns such as species-area relationships
(Hubbell, 2001), and must be rejected in favor of niche-centric models like DBD or EC. Using
neutral models without DBD or EC, we found generally flat or negative diversity slopes due to
sampling processes alone and that positive slopes were hard to explain with a neutral model
(Madi et al., 2020). These models were intended mainly for 16S rRNA gene sequence data, but
we expect the general conclusions to extend to metagenomic data. Nevertheless, further
modeling and experimental work will be required to fully exclude a neutral explanation for the
diversity slopes we report in the human gut microbiome.

Based on controlled experiments (Estrela et al., 2022) and modeling studies (San Roman
and Wagner, 2021), DBD is a plausible causal explanation for positive diversity slopes in the gut
microbiome. Although they also note that causality is difficult to establish, Good and Rosenfeld
(2022) suggest the importance of focal species evolution as a driver of changes in community

structure, as shown in an experimental study of *Pseudomonas* in compost communities (Padfield et al., 2020). Clearly, further work is needed to establish the extent and relative rates of eco-evolutionary feedback in both directions. How these feedbacks among bacteria are influenced by abiotic factors and by interactions with fungi, archaea, and phages also deserve further study.

Third, the diversity slope changes depending on which component of within-species diversity or community diversity is considered. Notably, the number of strains within a focal species is positively correlated with Shannon diversity, but inversely correlated with species richness, suggesting that the ability of strains to colonize a host may be associated with higher community evenness rather than total species count. Higher evenness might maximize the chance of inter-species interactions, whereas higher richness might be driven by rare species that are less likely to interact. Although Shannon diversity is considered to be more robust and informative than richness in estimating bacterial diversity (He et al., 2013; Reese and Dunn, 2018), we observe the same contrasting results between Shannon diversity and richness when community diversity is calculated at higher taxonomic levels, suggesting that this pattern is not due to artifacts such as sequencing effort.

Our measures of intra-species diversity included both synonymous and nonsynonymous single nucleotide variants, inferred strain richness, and gene content. Synonymous nucleotide variation was consistently and positively associated with both community richness and Shannon diversity at all taxonomic levels (although not always with statistical significance). Nonsynonymous variation also tended to track positively with both measures of community diversity but was only statistically significantly associated with phylum and class richness. This suggests that evolutionarily older, less selectively constrained synonymous mutations and more recent nonsynonymous mutations that affect protein structure both track similarly with measures of community diversity. Nonetheless, a parsimonious explanation for possible differences between the two classes is that while they are affected similarly, we have more statistical power to identify correlations in the more numerous synonymous mutations. This merits further investigation.

Metagenomes from the same individual sampled over time allowed us to detect gene gain and loss events. In both HMP and Poyet et al. time series, community diversity was predictive of future gene loss in a focal species. This phenomenon is not explicitly predicted by either DBD or EC but it is compatible with aspects of the Black Queen Hypothesis, with some caveats. BQH

predicts that a focal species will be less likely to encode genes with functions provided by other
members of the surrounding community if such functions are "leaky" and available as diffusible
public goods (Morris et al., 2012). The BQH could also act as a driver of polymorphism within a
species (Morris et al., 2014). Gene loss may be adaptive, provided that there is a cost to encoding
and expressing the relevant genes (Albalat and Cañestro, 2016; Koskiniemi et al., 2012;
Simonsen, 2022). The tendency for reductive genome evolution in bacteria is well established
(Albalat and Cañestro, 2016; Koskiniemi et al., 2012; Puigbò et al., 2014). Genome reduction is
a particular hallmark of endosymbiotic bacteria, which depend on their hosts for many metabolic
gene products (McCutcheon and Moran, 2012; Nikoh et al., 2011). It has been shown that
uncultivated bacteria from the gut have undergone considerable genome reduction, which may
be an adaptive process that results from reliance on public goods (Nayfach et al., 2019). In the
gut microbiome, the BQH has been invoked to explain the distribution of genes involved in
vitamin B metabolism (Sharma et al., 2019) and iron acquisition (Vatanen et al., 2019).

Our findings in human gut metagenomes are compatible with the BQH under the
assumption that increasing community diversity also increases the availability of leaky gene
products – which may not be the case if genomes in the gut microbiome are functionally
redundant, as inferred in a recent study (Tian et al., 2020). This study found that species in the
gut microbiome were highly redundant at the level of annotated metabolic pathways (KEGG
orthologs) and that more functionally redundant microbiomes were more resistant to colonization
by fecal transplants. Relatively low-redundancy microbiomes could therefore be more easily
colonized but might also require migrants to encode more gene functions in order to persist.
Importantly, functional redundancy may be high at the level of well-annotated metabolic
functions, but low at the finer level of individual gene families, as demonstrated in marine
microbiomes (Galand et al., 2018) but not yet studied explicitly in the gut. Here we report that
genome reduction in the gut is higher in more diverse gut communities. This could be due to *de*
novo gene loss, preferential establishment of migrant strains encoding fewer genes, or a
combination of the two. The mechanisms underlying this correlation remain unclear and could be
due to biotic interactions – including metabolic cross-feeding as posited by some models (Estrela
et al., 2022; San Roman and Wagner, 2021, 2018) but not others (Good and Rosenfeld, 2022) –
or due to unknown abiotic drivers of both community diversity and gene loss. Finally, we
measured community diversity from the phylum to the species level, not below. We therefore did

not investigate how the BQH could extend to maintain gene content variation within a species, as
has been shown experimentally in *E. coli* (Morris et al., 2014). This could be an avenue for
future work.

In our previous analysis of lower-resolution 16S rRNA amplicon sequences, we reported
a tendency for focal genera with larger genomes to have higher diversity slopes, perhaps because
they experience stronger DBD (Madi et al., 2020). At face value, this tendency seems at odds
with the BQH, which predicts genome reduction in more diverse communities. This apparent
contradiction may be reconciled by considering eco-evolutionary dynamics on different time
scales. A recent study used phylogenetic and metabolic reconstructions to show that gene gains
often drive metabolic dependencies among bacteria (Goyal, 2022), potentially explaining why
genera with larger maximum genome size could experience stronger DBD. Our earlier study
only had the genetic resolution to consider focal taxa down to the genus level, and by using the
maximum genome size observed in a public database we did not capture the dynamic process of
gene gain and loss within a species, as was possible in the current metagenomic study. It is
therefore possible that on longer (ecological) time scales, larger genomes have more metabolic
interactions and thus experience stronger DBD, while genome reduction in more diverse
communities occurs on shorter (evolutionary) time scales.

In summary, we demonstrate how metagenomic data can be used to test the predictions of
eco-evolutionary theory, including DBD, EC, and the BQH. It remains to be seen whether the
distinct eco-evolutionary processes proposed by DBD and the BQH operate orthogonally or
whether they interact. If BQH leads to gene losses that remain polymorphic rather than being lost
entirely from the species (Morris et al., 2014) – or invasions of strains with fewer genes that
remain incomplete and do not replace the resident strain – this could be viewed as a form of
diversification and perhaps a special case of DBD. Here we considered gene loss as a directional
process; we did not attempt to distinguish between directional changes in gene copy number and
the complete extinction of a gene, which is difficult to show using metagenomic data. Future
work could attempt to resolve this point and to potentially combine DBD and BQH into a unified
theory.

Data and materials availability

The raw sequencing reads for the metagenomic samples used in this study were downloaded from the Human Microbiome Project Consortium 2012 and Lloyd-Price et al. (2017) (URL: https://aws.amazon.com/datasets/human-microbiome-project/); and Poyet et al. 2019 (NCBI accession number PRJNA544527). All computer code for this paper is available at https://github.com/Naima16/DBD_in_gut_microbiome.

Methods

Metagenomic analyses

Estimation of species, gene, and SNV content of metagenomic samples

We used MIDAS (Metagenomic Intra-Species Diversity Analysis System, version 1.2, downloaded on November 21, 2016) (Nayfach et al., 2016) to estimate within-species nucleotide and gene content of raw metagenomic whole genome shotgun sequencing data for HMP1-2 and Poyet et al. 2019 data. MIDAS relies on a reference database comprised of 31,007 bacterial genomes that are clustered into 5,952 species, covering roughly 50% of species found in human stool metagenomes from "urban" individuals. Described below are the parameters used to estimate species abundances, single nucleotide variants (SNVs), and gene copy number variants (CNVs) with MIDAS.

Estimation of species content

We estimated species abundances, SNVs and CNVs by mapping metagenomic shotgun reads to reference genomes. Since a component of this work relies on quantifying polymorphism and CNV changes over time, we constructed a "personal" reference database to avoid spurious inferences of allele frequency and CNV changes due to errors in mapping of reads to regions of the genome shared by multiple species (Garud et al., 2019). This per-host reference database was comprised of the union of all species present at one or more timepoints so as to be as inclusive as possible to prevent reads from being "donated" to reference genome, while also being selective to prevent a reference genome from "stealing" reads from a species truly present.

To estimate the species relative abundances for each host x timepoint sample, we mapped
reads to 15 universal single-copy marker genes that are a part of the MIDAS pipeline (Nayfach et
al., 2016; Wu et al., 2013) and belong to the 5,952 species in the MIDAS reference database. A
species with an average marker gene coverage ≥ 3 was considered present for the purposes of
building a per-host database for mapping reads to infer SNVs and CNVs below. The per-host
database was constructed by including all species present at one or more timepoints with
coverage ≥ 3. However, more stringent thresholds were imposed for calling SNVs and CNVs, as
described below.

Estimation of copy number variation

To estimate gene copy number variation (CNV), we mapped reads to the pangenomes of
species present in a host's personal database using Bowtie2 (Langmead and Salzberg, 2012) with
default MIDAS settings (local alignment, MAPID$\geq$94.0%, READQ$\geq$20, and ALN_COV$\geq$0.75).
Each gene's coverage was estimated by dividing the total number of reads mapped to a given
gene by the gene length. These genes included the aforementioned 15 universal single-copy
marker genes. A given gene's copy number (c) was estimated by taking the ratio of its coverage
and the median coverage of the species' single-copy marker genes.

With these copy number values, we estimated the prevalence of genes in the between-
host population, defined as the fraction of samples with copy number $c \leq 3$ and $c \geq 0.3$ (conditional
on the mean single gene marker coverage being $\geq 5x$). For each species, we computed "core
genes", defined as genes in the MIDAS reference database that are present in at least 90% of
samples within a given cohort. Within-host polymorphism rates were computed in core genes.

Orthologous genes present in multiple species can result in read "stealing" and read
"donating" to species from which the reads did not originate. Thus, we excluded a set of genes
belonging to a 'blacklist' composed of genes present in multiple species. This blacklist was
constructed in (Garud et al., 2019) using USEARCH (Edgar, 2010) to cluster all genes in human-
associated reference genomes with a 95% nucleotide identity threshold. Since some genes may
be absent from the MIDAS database but may nevertheless be shared across species, we
implemented another filter (as in Garud et al. 2019) in which genes with $c \geq 3$ in at least one
sample in our cohort were excluded from analysis of polymorphism rate or gene changes over
time.

Inferring single nucleotide variants (SNVs) within bacterial species

To call SNVs, we mapped reads to a single representative reference genome as per the

default MIDAS software. Reads were mapped with Bowtie2, with default MIDAS mapping

thresholds: global alignment, MAPID≥94.0%, READQ≥20, ALN_COV≥0.75, and MAPQ≥20.

Species were excluded from further analysis if reads mapped to ≤ 40% of their genome. We

additionally excluded samples from further analysis if they had low median read coverage ($\underline{D}$) at

protein coding sites. Specifically, samples with $\underline{D} < 5$ across all protein coding sites with nonzero

coverage were excluded. This MIDAS SNV output was then used for computing within-species

polymorphism rates and inferring the number of strains present for each species in each sample

(see below).

To compute polymorphism rates, additional bioinformatic filters were imposed to avoid

read stealing and donating across different species. First, we did not call SNVs in blacklisted

genes present in multiple species. Additionally, we excluded sites in a given sample if $D < 0.3\underline{D}$

or $D > 3\underline{D}$ as these sites harbor anomalously low or high coverage compared to the genome-wide

average $\underline{D}$. Additional filters are described below.

Shannon diversity, species richness and polymorphism rate calculations

Shannon diversity and richness were computed within each sample by including any

species with abundance greater than zero. Rarefied species richness estimates are based on

HMP1-2 samples rarefied to 20 million reads and Poyet samples rarefied to 5 million reads. SNV

and gene content variation within a focal species were ascertained only from the full dataset and

not the rarefied dataset.

The polymorphism rate of a species in a sample was computed as the proportion of

synonymous sites in core genes with intermediate allele frequencies ($0.2 \leq f \leq 0.8$), as was done

in Garud et al. 2019. Only species with a MIDAS marker gene coverage of 10 or more in 10 or

more samples were included, yielding 69 species in 249 HMP stool donors and 15 species in four

Poyet et al. 2019 donors. As explained in SI text 1 in Garud et al. 2019, this is quantitatively

similar to the more traditional population genetic measure of heterozygosity, $H=E[2f(1-f)]$, in

which intermediate frequency alleles contribute the most weight to heterozygosity. By

computing polymorphism with the criteria $0.2 \leq f \leq 0.8$, we avoid inclusion of low frequency

sequencing errors, which can otherwise greatly influence the mean heterozygosity.
Polymorphism rates were computed separately for synonymous (4-fold degenerate) and
nonsynonymous (1-fold degenerate) sites. The degeneracy of sites was determined based on
MIDAS output.
630

Temporal changes in polymorphism rates and gene content
Polymorphism change was computed as the difference in polymorphism rates between
time points within a host. Gene gains and losses between time points were computed in species
with sufficient prevalence (at least 10 samples with marker gene coverage of at least 10, as in the
polymorphism analysis above) by identifying genes with copy number c ≤0.05 (indicating gene
absence) in one sample and $0.6 \leq c \leq 1.2$ (with marker coverage ≥20x) in another (indicating
single copy gene presence). These thresholds were used in Garud et al. 2019 when inferring gene
changes in temporal data and reflect a range of copy numbers expected in either the absence of a
gene or presence of a single copy of a gene given typical coverage values in growing cells
(Korem et al., 2015). These copy number cutoffs were chosen to avoid spuriously analyzing
genes linked to multiple species. In such cases, mapping artifacts in which reads can be
arbitrarily assigned to multiple species cannot be disentangled. For example, a gene present in
multiple species would likely have copy number significantly deviating from 1 (including values
that lie in an ambiguous zone of 0.05 to 0.6, as well as >>1), reflecting the joint abundances of
the multiple species. Thus, although we may miss many biologically interesting multi-copy
genes (e.g. transporter genes in Bacteroides (Wexler and Goodman, 2017)), our thresholds avoid
confounding our analysis with read stealing or donating among different species. Filters for
coverage and blacklisted genes were applied as described above.
649

Strain number inference
We used StrainFinder (Smillie et al., 2018) to infer the number of strains present within
each species in each HMP1-2 metagenomic sample. To do so, we used allele frequencies from
MIDAS SNV output, generated as described above. For each species in each host, all multi-
allelic sites with coverage of 20x or greater were passed as input to StrainFinder. Species/host
pairs which had fewer than 100 sites with 20x coverage were removed from the analysis.
StrainFinder was then run on each sample separately for strain number 1, 2, 3, and 4, and the

optimal strain number was chosen based on the Bayesian Information Criterion (BIC). This

range of strain number was chosen for biological reasons. A number of studies have

demonstrated that at most a small handful of strains (between one and four) not sharing a

common ancestor within the host are ever observed within a single gut microbiome at any one

time (Garud et al., 2019; Truong et al., 2017; Verster et al., 2017; Yassour et al., 2018).

Additionally, for the four densely longitudinally sampled hosts in Poyet et al. 2019, multiple

analyses employing distinct sequencing strategies and strain phasing techniques have similarly

concluded that a maximum of four strains were present at any one time within a host for the ~30

most prevalent species (Poyet et al., 2019; Wolff et al., 2021; Zheng et al., 2022). Thus, four

strains were chosen as the maximum to accommodate the range of observed possibilities.

Statistical analyses

Model construction and evaluation
Using data from the Human Microbiome Project (HMP) and Poyet et al. 2019, we examined the

relationship between within-species genetic diversity and the gut microbiome community

diversity. Within-species genetic diversity was estimated with polymorphism rate and strain

richness. Community diversity was estimated with the Shannon index, species richness estimated

on the whole data, and species richness calculated on the data rarefied to an equal number of

reads per sample (as described above). When the relationship between the response variable

(within-species genetic diversity) and the predictor (community diversity) was approximately

linear by visual inspection, we fit generalized linear mixed models (GLMMs) (glmmTMB

function from the glmmTMB R package - RStudio version 1.2.5042) with community diversity

as the predictor of within-species genetic diversity. Otherwise we fit generalized additive mixed

models (GAMs) (mgcv function from the mgcv R package - RStudio version 1.2.5042) to

account for the non-linearity of the relationships.

To account for variation in sequencing depth, which can affect estimates of both community

diversity and within-species genetic diversity, we added read count per sample (coverage) as a

covariate to all generalized mixed models. Species name, subject identifier and sample identifier

were added as random effects to account for variation between different species and subjects,

and to account for non-independence between observations. The R syntax and statistics of all

generalized models are reported in **Supplementary File 2**.

In generalized mixed models, the predictors were standardized to zero mean and unit variance

before analyses. We first assessed random effects significance by comparing nested models

where each random effect was dropped one at a time using the likelihood-ratio test (LRT, anova

function from the R stats package) and only significant random effects were included in the final

models. We then assessed the fixed effects' significance with likelihood-ratio tests implemented

in the drop1 function in the R stats package. This function drops individual terms from the full

model and reports the AIC and the LRT P-value. All the P-values reported for the GLMMs

correspond to LRT and not to the Wald P-values reported by glmm.summary function from the

R package glmmTMB, as was recommended in https://bbolker.github.io/mixedmodels-

misc/glmmFAQ.html. We again used LRTs to compare the full significant models to null models

including all random effects but no fixed effects other than the intercept. The difference in

Akaike information criterion (ΔAIC) between full and null model and their associated P-values

are reported in **Supplementary File 1e,f,g**. As an additional evaluation of the goodness of fits,

we estimated the coefficient of determination (R^2) using the r2 function from the performance R

package. Two values are reported: the marginal R^2, a measure of the variance explained only by

fixed effects, and the conditional R^2, a measure of the variance explained by the entire model.

We evaluated model fits by inspecting the residuals using the DHARMa library in R

(simulateResiduals and plot functions) for the GLMMs and by inspecting residual distributions

and fitted-observed value plots using the gam.check function from the mgcv R package for the

GAMs. Adjusted R^2 values (from gam.summary function from the mgcv R package) are reported

as a goodness of fit for the GAMs. All model outputs (summary function from mgcv and

glmmTMB R packages) are reported in the **Supplementary File 2**.

To study the relationship between focal species polymorphism and community diversity

calculated at higher taxonomic ranks (from genus to phylum), we used GTDBK and the Genome

Taxonomy Database (GTDB) (Chaumeil et al., 2020) to annotate MIDAS reference genomes.

Richness at each level was estimated with the total number of distinct taxonomic units in the

sample. The Shannon index was calculated based on the relative abundances table from MIDAS:
at each taxonomic level, we used the sum of the abundances of all species belonging to that
taxonomic level to calculate the Shannon index (using the diversity function from the R vegan
library). We then fit two separate GAMs for each taxonomic rank (from genus to phylum) with
either Shannon diversity or richness as the predictors of within-species polymorphism rate (with
the coverage per sample as a covariate and species name, sample and subject identifiers as
random effects). These GAMs were fitted with a beta error distribution with logit-link function,
chosen because polymorphism rate is a continuous value strictly bounded by 1, and all the terms
were smoothed terms (See **Supplementary File 1c** and **Supplementary File 2 section 1-3** for
additional model details).

We repeated the same methods for focal species synonymous and nonsynonymous
polymorphism separately. See **Supplementary File 1b and d** and **Supplementary File 2**
section 4-6 for details of the models applied to nonsynonymous polymorphism.

Analysis of strain count

To study the relationship between community diversity and the number of strains within a focal
species in the HMP, we restricted the analysis to 184 focal species genomes with at least 100
nucleotide sites with 20x coverage in a sample. We fit separate GLMMs with strain count in a
focal species as a function of community diversity estimated with Shannon diversity, species
richness, or rarefied species richness. Since strain number is positive count data, we compared
many zero-truncated count models based on the Akaike information criterion (AIC) score
(AICtab function from bbmle R library) (Brooks et al., 2017). We fit the model with the
truncated negative binomial distribution (truncated_nbinom2 or truncated_nbinom1 in
glmmTMB; the second best fit) in order to resolve the overdispersion detected in the best fit (the
truncated Poisson model). Overdispersion was tested using the check_overdispersion function
from the performance R package as described here: https://bbolker.github.io/mixedmodels-
misc/glmmFAQ.html.

As described above for focal species polymorphism, we tested the relationship between focal
species strain count and community diversity at higher taxonomic levels from genus to phylum,

fitting a separate GLMM with strain count in a focal species as a function of each metric of

diversity (Shannon and richness) at higher taxonomic levels (from genus to phylum). All GLMM

details are reported in **Supplementary File 1f** and **Supplementary File 2 section 7-9**.

Analysis of time series data

To test the predictions of DBD over time, we used HMP samples with multiple time points from

the same person to look at the relationship between within-species polymorphism change,

defined as the difference between polymorphism rate at two time points, and community

diversity at the earlier time point. To account for non-linearity of the relationships, we fit GAMs

with log transformed polymorphism change as a function of community diversity at the earlier

time point, and added the coverage per sample at the earlier time point as a covariate as well as

species name, sample and subject identifiers as random effects (**Supplementary File 2 section**

10.1).

In addition, we investigated the effect of community diversity at one time point on gene content

variation (gains and losses considered separately) at the subsequent time point. The relationships

were generally linear, so we used separate negative binomial generalized linear mixed models

with gene gain as the response and each of the metrics of community diversity as the predictor

with the same covariates and random effects used in the previous models (**Supplementary File 2**

section 10.2). The same method was used to test how gene loss was related to community

diversity (**Supplementary File 1g**, **Supplementary File 2 section 10.3**).

HMP longitudinal data consisted of hosts sampled at a time lag of ~6 months. To assess the

relationship between within-species genetic diversity and community diversity at higher

temporal resolution, we used the same methods to analyze longitudinal metagenomic data from

four more frequently sampled healthy stool donors (hosts *am*, *an*, *ao* and *ae*) (Poyet et al., 2019).

Stool from donor *am* was sequenced over 18 months with a median of one day between samples;

an over 12 months (median 2 days between samples); *ao* over 5 months (median 1 day between

samples); and *ae* over 7 months (median 2 days between samples). We looked at polymorphism

change and gene gains and losses between two time points in the 15 species with a minimal

marker gene coverage of ten in at least ten samples. Community diversity was estimated with

Shannon diversity (unrarefied) and richness calculated on rarefied data to 5 million reads per
sample.

To study the relationship between community diversity at the initial time point and
polymorphism change between the initial time point and all the future time points, we fit
Gaussian generalized additive mixed models with log-transformed polymorphism change as the
response and the interaction between community diversity at the first time point and the number
of days between time points as the predictor. Covariates included coverage, species name,
sample, and subject identifiers as random effects (**Supplementary File 1h**, **Supplementary File**
2 section 11.1 and 11.2). To study the relationship between gene variation (gains and losses
separately) and diversity at the first time point, we fit negative binomial generalized linear mixed
models with gene variation as a function of the interaction between diversity at the first time
point and the number of days between the two time points, with the same covariate and random
effects as used above for polymorphism change over time (**Supplementary File 1i**,
Supplementary File 2 section 11.3-11.6).

Acknowledgements
We sincerely thank members of the Garud and Shapiro labs for their feedback during the
development of this paper. NRG received support from the Paul Allen Frontiers Group, a
University of California Hellman fellowship, a UCLA Faculty Career Development award, and
the Research Corporation for Science Advancement. DWC received funding support from
NIH R25 MH 109172. BJS was supported by a Natural Sciences and Engineering Research
Council of Canada (NSERC) Discovery Grant and a Canada Research Chair. We also thank
Djordje Bajić and two anonymous reviewers for their constructive suggestions that substantially
improved the manuscript.

Supplementary files

810
811 **Supplementary File 1.** Supplementary tables a-i.

812
813 **Supplementary File 2.** R syntax and statistics of all generalized models.

816 # Supplementary figure legends
817

818
819
Figure 2-Supplement 1. Results of generalized additive models predicting within-species polymorphism rate
(at synonymous sites) as a function of community diversity at higher taxonomic levels (HMP data). (A1-E1)
The predictor is Shannon diversity. (A2-E2) The predictor is richness. Adjusted r-squared (R^2) and Chi-squared *P*-
values corresponding to the predictor are displayed in each panel (gam.summary function from mgcv R package).
Shaded areas show the 95% confidence interval of each model prediction. See Supplementary File 1c and
supplementary file 2 sections 2-3 for further details about model outputs.
826
827
828

Figure 2-Supplement 2. Results of generalized additive models predicting within-species polymorphism rate
(at nonsynonymous sites) in a focal species as a function of community diversity at higher taxonomic levels
(HMP data). (A1-E1) The predictor is Shannon diversity. (A2-E2) The predictor is richness. Adjusted r-squared
(R^2) and Chi-squared *P*-values corresponding to the predictor are displayed in each panel (gam.summary function
from mgcv R package). Shaded areas show the 95% confidence interval of each model prediction. See
Supplementary File 1d and supplementary file 2 sections 5-6 for further details about model outputs.
836
837

838
839

Figure 3-Supplement 1. Results of generalized linear mixed models predicting strain count in a focal species
as a function of community diversity at higher taxonomic levels (HMP data). Strain number in a focal species is
positively correlated with Shannon (A1-E1) whereas its correlation with richness remains negative (A2-E2) through
all taxonomic levels. The Y-axis is the predicted mean number of strains within a focal species. *P*-values (drop1
function from R stats package, LRT). Shaded areas show the 95% confidence interval of each model prediction. See
Supplementary File 1f and supplementary file 2 section 9 for model details.
846

848 **Figure 5-Supplement 1.** Results of a GAM predicting polymorphism change in a focal species as a function of the
849 interaction between Shannon diversity at the first time point and the time lag (days) between two time points in the
850 Poyet time series. The response (Y-axis) was log transformed in the Gaussian GAM. Several different time lags are
851 shown to illustrate the inversion of the relationship around a lag time of 150 days. See Supplementary File 1h and
852 supplementary file 2 section 11 for further model details.

References

Albalat R, Cañestro C. 2016. Evolution by gene loss. *Nat Rev Genet*. doi:10.1038/nrg.2016.39

Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Mächler M, Bolker BM. 2017. Modeling zero-inflated count data with glmmTMB. *BioRxiv*.

Calcagno V, Jarne P, Loreau M, Mouquet N, David P. 2017. Diversity spurs diversification in ecological communities. *Nat Commun* **8**. doi:10.1038/ncomms15810

Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: A toolkit to classify genomes with the genome taxonomy database. *Bioinformatics* **36**. doi:10.1093/bioinformatics/btz848

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**. doi:10.1093/bioinformatics/btq461

Estrela S, Diaz-Colunga J, Vila JCC, Sanchez-Gorostiaga A, Sanchez A. 2022. Diversity begets diversity under microbial niche construction. *BioRxiv*.

Galand PE, Pereira O, Hochart C, Auguet JC, Debroas D. 2018. A strong link between marine microbial community composition and function challenges the idea of functional redundancy. *ISME J* **12**. doi:10.1038/s41396-018-0158-1

Garud NR, Good BH, Hallatschek O, Pollard KS. 2019. Evolutionary dynamics of bacteria in the gut microbiome within and across hosts. *PLoS Biol* **17**:e3000102. doi:10.1371/JOURNAL.PBIO.3000102

Garud NR, Pollard KS. 2020. Population Genetics in the Human Microbiome. *Trends Genet* **36**:53–67. doi:10.1016/j.tig.2019.10.010

Good BH, Rosenfeld LB. 2022. Eco-evolutionary feedbacks in the human gut microbiome. *bioRxiv*.

Goyal A. 2021. Horizontal Gene Transfer Drives the Evolution of Dependencies in Bacteria. *iScience* **25**: 104312

Goyal A, Bittleston LS, Leventhal GE, Lu L, Cordero OX. 2022. Interactions between strains govern the eco-evolutionary dynamics of microbial communities. *Elife* **11**. doi:10.7554/ELIFE.74987

Groussin M, Poyet M, Sistiaga A, Kearney SM, Moniz K, Noel M, Hooker J, Gibbons SM, Segurel L, Froment A, Mohamed RS, Fezeu A, Juimo VA, Lafosse S, Tabe FE, Girard C, Iqaluk D, Nguyen LTT, Shapiro BJ, Lehtimäki J, Ruokolainen L, Kettunen PP, Vatanen T,

884  Sigwazi S, Mabulla A, Domínguez-Rodrigo M, Nartey YA, Agyei-Nkansah A, Duah A,

885   Awuku YA, Valles KA, Asibey SO, Afihene MY, Roberts LR, Plymoth A, Onyekwere CA,

886   Summons RE, Xavier RJ, Alm EJ. 2021. Elevated rates of horizontal gene transfer in the

887   industrialized human microbiome. *Cell* **184**. doi:10.1016/j.cell.2021.02.052

888 He Y, Zhou BJ, Deng GH, Jiang XT, Zhang H, Zhou HW. 2013. Comparison of microbial

889   diversity determined with the same variable tag sequence extracted from two different PCR

890   amplicons. *BMC Microbiol* **13**. doi:10.1186/1471-2180-13-208

891 Hehemann JH, Correc G, Barbeyron T, Helbert W, Czjzek M, Michel G. 2010. Transfer of

892   carbohydrate-active enzymes from marine bacteria to Japanese gut microbiota. *Nature*

893   **464**:908–912. doi:10.1038/nature08937

894 Hibbing ME, Fuqua C, Parsek MR, Peterson SB. 2010. Bacterial competition: Surviving and

895   thriving in the microbial jungle. *Nat Rev Microbiol*. doi:10.1038/nrmicro2259

896 Hubbell SP. 2001. The Unified Neutral Theory of Biodiversity and Biogeography. *Princeton*

897   *Univ Press*.

898 Jousset A, Eisenhauer N, Merker M, Mouquet N, Scheu S. 2016. High functional diversity

899   stimulates diversification in experimental microbial communities. *Sci Adv* **2**.

900   doi:10.1126/sciadv.1600124

901 Korem T, Zeevi D, Suez J, Weinberger A, Avnit-Sagi T, Pompan-Lotan M, Matot E, Jona G,

902   Harmelin A, Cohen N, Sirota-Madi A, Thaiss CA, Pevsner-Fischer M, Sorek R, Xavier RJ,

903   Elinav E, Segal E. 2015. Growth dynamics of gut microbiota in health and disease inferred

904   from single metagenomic samples. *Science (80-)* **349**:1101–1106.

905   doi:10.1126/science.aac4812

906 Koskiniemi S, Sun S, Berg OG, Andersson DI. 2012. Selection-driven gene loss in bacteria.

907   *PLoS Genet* **8**. doi:10.1371/journal.pgen.1002787

908 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**.

909   doi:10.1038/nmeth.1923

910 Lenski RE. 2017. Experimental evolution and the dynamics of adaptation and genome evolution

911   in microbial populations. *ISME J*. doi:10.1038/ismej.2017.69

912 Lloyd-Price J, Mahurkar A, Rahnavard G, Crabtree J, Orvis J, Hall AB, Brady A, Creasy HH,

913   McCracken C, Giglio MG, McDonald D, Franzosa EA, Knight R, White O, Huttenhower C.

914   2017. Strains, functions and dynamics in the expanded Human Microbiome Project. *Nature*

550:61–66. doi:10.1038/nature23889

Madi N, Vos M, Murall CL, Legendre P, Shapiro BJ. 2020. Does diversity beget diversity in
microbiomes? *Elife* **9**. doi:10.7554/eLife.58999

Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A
phylogenetic perspective. *Science (80-)*. doi:10.1126/science.aac9323

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. *Nat Rev*
Microbiol. doi:10.1038/nrmicro2670

Meyer JR, Kassen R. 2007. The effects of competition and predation on diversification in a
model adaptive radiation. *Nature* **446**. doi:10.1038/nature05599

Mitri S, Foster KR. 2013. The genotypic view of social interactions in microbial communities.
Annu Rev Genet. doi:10.1146/annurev-genet-111212-133307

Morris JJ, Lenski RE, Zinser ER. 2012. The black queen hypothesis: Evolution of dependencies
through adaptive gene loss. *MBio* **3**. doi:10.1128/mBio.00036-12

Morris JJ, Papoulis SE, Lenski RE. 2014. Coexistence of evolving bacteria stabilized by a shared
Black Queen function. *Evolution* **68**. doi:10.1111/evo.12485

Nayfach S, Rodriguez-Mueller B, Garud N, Pollard KS. 2016. An integrated metagenomics
pipeline for strain profiling reveals novel patterns of bacterial transmission and
biogeography. *Genome Res* **26**:1612–1625. doi:10.1101/gr.201863.115

Nayfach S, Shi ZJ, Seshadri R, Pollard KS, Kyrpides NC. 2019. New insights from uncultivated
genomes of the global human gut microbiome. *Nature* **568**:505–510. doi:10.1038/s41586-
019-1058-x

Nikoh N, Hosokawa T, Oshima K, Hattori M, Fukatsu T. 2011. Reductive evolution of bacterial
genome in insect gut environment. *Genome Biol Evol* **3**. doi:10.1093/gbe/evr064

Olm MR, Brown CT, Brooks B, Firek B, Baker R, Burstein D, Soenjoyo K, Thomas BC,
Morowitz M, Banfield JF. 2017. Identical bacterial populations colonize premature infant
gut, skin, & oral microbiomes & exhibit different in situ growth rates. *Genome Res* **27**:601–
612. doi:10.1101/gr.213256.116

Padfield D, Vujakovic A, Paterson S, Griffiths R, Buckling A, Hesse E. 2020. Evolution of
diversity explains the impact of pre-adaptation of a focal species on the structure of a
natural microbial community. *ISME J* **14**. doi:10.1038/s41396-020-00755-3

Poulsen LK, Licht TR, Rang C, Krogfelt KA, Molin S. 1995. Physiological state of Escherichia

coli BJ4 growing in the large intestines of streptomycin-treated mice. *J Bacteriol* **177**.

doi:10.1128/jb.177.20.5840-5845.1995

Poyet M, Groussin M, Gibbons SM, Avila-Pacheco J, Jiang X, Kearney SM, Perrotta AR, Berdy

B, Zhao S, Lieberman TD, Swanson PK, Smith M, Roesemann S, Alexander JE, Rich SA,

Livny J, Vlamakis H, Clish C, Bullock K, Deik A, Scott J, Pierce KA, Xavier RJ, Alm EJ.

2019. A library of human gut bacterial isolates paired with longitudinal multiomics data

enables mechanistic microbiome research. *Nat Med* **25**:1442–1452. doi:10.1038/s41591-

019-0559-3

Puigbò P, Lobkovsky AE, Kristensen DM, Wolf YI, Koonin E V. 2014. Genomes in turmoil:

Quantification of genome dynamics in prokaryote supergenomes. *BMC Med* **12**.

doi:10.1186/s12915-014-0066-4

Reese AT, Dunn RR. 2018. Drivers of microbiome biodiversity: A review of general rules, feces,

and ignorance. *MBio* **9**. doi:10.1128/mBio.01294-18

Russell SL, Cavanaugh CM. 2017. Intrahost genetic diversity of bacterial symbionts exhibits

evidence of mixed infections and recombinant haplotypes. *Mol Biol Evol* **34**:2747–2761.

doi:10.1093/molbev/msx188

San Roman M, Wagner A. 2021. Diversity begets diversity during community assembly until

ecological limits impose a diversity ceiling. *Mol Ecol* **30**. doi:10.1111/mec.16161

San Roman M, Wagner A. 2018. An enormous potential for niche construction through bacterial

cross-feeding in a homogeneous environment. *PLoS Comput Biol* **14**.

doi:10.1371/journal.pcbi.1006340

Schluter D. 2000. The Ecology of Adaptive Radiation. *Oxford Ser Ecol Evol*.

Schluter D, Pennell MW. 2017. Speciation gradients and the distribution of biodiversity. *Nature*.

doi:10.1038/nature22897

Sharma V, Rodionov DA, Leyn SA, Tran D, Iablokov SN, Ding H, Peterson DA, Osterman AL,

Peterson SN. 2019. B-Vitamin sharing promotes stability of gut microbial communities.

Front Microbiol **10**. doi:10.3389/fmicb.2019.01485

Simonsen AK. 2022. Environmental stress leads to genome streamlining in a widely distributed

species of soil bacteria. *ISME J* **16**. doi:10.1038/s41396-021-01082-x

Smillie CS, Sauk J, Gevers D, Friedman J, Sung J, Youngster I, Hohmann EL, Staley C, Khoruts

A, Sadowsky MJ, Allegretti JR, Smith MB, Xavier RJ, Alm EJ. 2018. Strain Tracking

Reveals the Determinants of Bacterial Engraftment in the Human Gut Following Fecal

Microbiota Transplantation. *Cell Host Microbe*. doi:10.1016/j.chom.2018.01.003

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier hypothesis and

mutation-rate evolution. *Proc Natl Acad Sci U S A* **109**. doi:10.1073/pnas.1216223109

The Human Microbiome Project Consortium. 2012. A framework for human microbiome

research. *Nature* **486**.

Tian L, Wang XW, Wu AK, Fan Y, Friedman J, Dahlin A, Waldor MK, Weinstock GM, Weiss

ST, Liu YY. 2020. Deciphering functional redundancy in the human microbiome. *Nat*

Commun **11**. doi:10.1038/s41467-020-19940-1

Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. 2017. Microbial strain-level population

structure & genetic diversity from metagenomes. *Genome Res* **27**:626–638.

doi:10.1101/gr.216242.116

Vatanen T, Plichta DR, Somani J, Münch PC, Arthur TD, Hall AB, Rudolf S, Oakeley EJ, Ke X,

Young RA, Haiser HJ, Kolde R, Yassour M, Luopajärvi K, Siljander H, Virtanen SM,

Ilonen J, Uibo R, Tillmann V, Mokurov S, Dorshakova N, Porter JA, McHardy AC,

Lähdesmäki H, Vlamakis H, Huttenhower C, Knip M, Xavier RJ. 2019. Genomic variation

and strain-specific functional adaptation in the human gut microbiome during early life. *Nat*

Microbiol **4**. doi:10.1038/s41564-018-0321-5

Venturelli OS, Carr A V, Fisher G, Hsu RH, Lau R, Bowen BP, Hromada S, Northen T, Arkin

AP. 2018. Deciphering microbial interactions in synthetic human gut microbiome

communities. *Mol Syst Biol* **14**. doi:10.15252/msb.20178157

Verster AJ, Ross BD, Radey MC, Bao Y, Goodman AL, Mougous JD, Borenstein E. 2017. The

Landscape of Type VI Secretion across Human Gut Microbiomes Reveals Its Role in

Community Composition. *Cell Host Microbe* **22**:411-419.e4.

doi:10.1016/j.chom.2017.08.010

Walters KE, Martiny JBH. 2020. Alpha-, beta-, and gamma-diversity of bacteria varies across

habitats. *PLoS One* **15**. doi:10.1371/journal.pone.0233872

Wexler AG, Goodman AL. 2017. An insider's perspective: Bacteroides as a window into the

microbiome. *Nat Microbiol*. doi:10.1038/nmicrobiol.2017.26

Wolff R, Shoemaker WR, Garud NR. 2021. Ecological Stability Emerges at the Level of Strains

in the Human Gut Microbiome. *bioRxiv*. doi:https://doi.org/10.1101/2021.09.30.462616

Wu D, Jospin G, Eisen JA. 2013. Systematic Identification of Gene Families for Use as

"Markers" for Phylogenetic and Phylogeny-Driven Ecological Studies of Bacteria and

Archaea and Their Major Subgroups. *PLoS One* **8**. doi:10.1371/journal.pone.0077033

Yaffe E, Relman DA. 2020. Tracking microbial evolution in the human gut using Hi-C reveals

extensive horizontal gene transfer, persistence and adaptation. *Nat Microbiol* **5**.

doi:10.1038/s41564-019-0625-0

Yang Y, Nguyen M, Khetrapal V, Sonnert ND, Martin AL, Chen H, Kriegel MA, Palm NW.

2022. Within-host evolution of a gut pathobiont facilitates liver translocation. *Nature*.

Yassour M, Jason E, Hogstrom LJ, Arthur TD, Tripathi S, Siljander H, Selvenius J, Oikarinen S,

Hyöty H, Virtanen SM, Ilonen J, Ferretti P, Pasolli E, Tett A, Asnicar F, Segata N,

Vlamakis H, Lander ES, Huttenhower C, Knip M, Xavier RJ. 2018. Strain-Level Analysis

of Mother-to-Child Bacterial Transmission during the First Few Months of Life. *Cell Host*

Microbe **24**:146-154.e4. doi:10.1016/j.chom.2018.06.007

Zhao S, Lieberman TD, Poyet M, Kauffman KM, Gibbons SM, Groussin M, Xavier RJ, Alm EJ.

2019. Adaptive Evolution within Gut Microbiomes of Healthy People. *Cell Host Microbe*

25:656-667.e8. doi:10.1016/j.chom.2019.03.007

Zheng W, Zhao S, Yin Y, Zhang H, Needham DM, Evans ED, Dai CL, Lu PJ, Alm EJ, Weitz

DA. 2022. High-throughput, single-microbe genomics with strain resolution, applied to a

human gut microbiome. *Science (80-)* **376**.

1027