Heritable variation in thermoregulation is associated with

reproductive success in the world's largest bird

Erik I. Svensson [*, a], Mads F. Schou [*, a], Julian Melgar [a], John Waller [b], Anel Engelbrecht [c],

Zanell Brand [c], Schalk Cloete [c, d] & Charlie K. Cornwallis [a]

[a] Department of Biology, Lund University, Lund, Sweden
[b] Global Biodiversity Information Facility (GBIF), Copenhagen, Denmark
[c] Directorate Animal Sciences, Western Cape Department of Agriculture, South Africa
[d] Department of Animal Sciences, University of Stellenbosch, Matieland, South Africa
* These authors contributed equally

Manuscript type: *Letter*

Short running title: *Evolution of thermoregulation in the ostrich*

Keywords: birds, deserts, fitness, heat stress, heritability, ostrich, *Struthio camelus*, thermal imaging, thermoregulation, thermal adaptation

Abstract word count: 159

Total word count: 4852

Abstract

Organisms inhabiting extreme thermal environments, such as desert birds, have evolved spectacular adaptations to thermoregulate during hot and cold conditions. However, our knowledge if and how selection acts on thermoregulation and the potential for evolutionary responses is limited, particularly for large organisms experiencing extreme temperature fluctuations. Here we use thermal imaging to quantify selection and genetic variation in thermoregulation in ostriches (*Struthio camelus*), the world's largest bird species that is regularly exposed to increasingly volatile temperatures. We found that females that are better at regulating their head temperatures had higher egg-laying rates under hotter conditions. Thermoregulatory capacity was both heritable and showed signatures of local adaptation. Females originating from populations exposed to more unpredictable climates were better at regulating their head temperatures in response to temperature fluctuations. Together these results reveal that past and present evolutionary processes have shaped genetic variation in thermoregulatory capacity, which appears to protect critical organs, such as the brain, from extreme temperatures during reproduction.

Impact Summary

Large animals inhabiting extreme thermal environments, such as deserts, are predicted to be particularly vulnerable to the sustained heat waves expected in their future climate. However, previous research on the evolution of thermoregulation has focused on the effect of increasing temperatures on survival in ectotherms. Here we study the ostrich (*Struthio camelus*), the world's largest bird, that inhabits some of the hottest and driest regions on Earth. We show that the ability of females to reproduce during hot conditions is associated with their thermoregulatory capacity. Furthermore, variation in thermoregulation is heritable and related to past climatic conditions: Females from populations originating from parts of Africa with more extreme temperature fluctuations were better able to thermoregulate, indicating local adaptation to different climatic conditions. Together, these results suggest that thermoregulation in this large desert bird has evolved in response to past climatic conditions, remains genetically variable and is currently under selection due to its effect on reproduction.

Introduction

Organisms need to manage heat and cold stress to survive and reproduce in variable climates (Angiletta, 2009). The universal challenge of coping with thermal stress (Hurley *et al.*, 2018; Sales *et al.*, 2018; Walsh *et al.*, 2019; Schou *et al.*, 2021) has promoted the evolution of adaptations to regulate body temperature (Huey *et al.*, 2003; Angiletta, 2009). For example, physiological mechanisms of selective brain cooling in birds (Fuller *et al.*, 2003), sweating in humans (Périard *et al.*, 2015) and morphological traits such as the ears of African Elephants (Weissenböck *et al.*, 2010) and the bills (Tattersall *et al.*, 2009; Symonds & Tattersall, 2010) and featherless skin patches (Szafrańska *et al.*, 2020) in different species of birds can all help to regulate temperature (McCafferty *et al.*, 2011). The different mechanisms by which animals thermoregulate and how these mechanisms could promote survival have been intensely studied. However, much less is known about the heritability of thermoregulatory traits and how such traits affect reproductive success, particularly for organisms living in the regions most affected by climate change (Walsh *et al.*, 2019), such as deserts (Bourne *et al.*, 2020; Schou *et al.*, 2021).

One group of animals expected to be particularly vulnerable to changing temperatures are large endotherms (birds and mammals). Larger bodies have a higher thermal inertia and slower rate of body temperature change compared to small bodies (Bogert, 1949; Angiletta, 2009). Thermal inertia can help maintain body temperatures during cold conditions, but it can also increase physiological stress under sustained heat, jeopardising survival and reproduction (Hurley *et al.*, 2018; Sales *et al.*, 2018; Bourne *et al.*, 2020; Parratt *et al.*, 2021; Schou *et al.*, 2021, 2022). Large animals are also expected to be particularly vulnerable to changing climates because their rate of adaptative evolution is potentially limited by longer generation times and lower population sizes (Cardillo *et al.*, 2005; Atwood *et al.*, 2020). Understanding how large-bodied

animals cope with thermal stress in fluctuating thermal environments therefore requires

particular consideration.

The evolutionary potential of populations to adapt to thermal stressors such as heatwaves and

cold snaps, depends on genetic variance in traits involved in thermoregulation (Hurley *et al.*,

2018; Sales *et al.*, 2018; Bourne *et al.*, 2020). Current empirical evidence suggests that genetic

variation in thermoregulatory traits is often low, constraining species responses to shifting

temperature regimes (Hoffmann *et al.*, 2003; Kellermann *et al.*, 2009). For instance, body

temperatures and thermal adaptations are typically strongly phylogenetically conserved

(Kellermann *et al.*, 2012a, 2012b) and evolve slowly compared to other traits (Moreira *et al.*,

2021). The late evolutionary biologist George C. Williams even questioned if endothermic body

temperatures could evolve at all, suggesting that there may be evolutionary stasis in body

temperatures and thermal adaptations (Williams, 1992). However, it is unclear if genetic

variance in thermoregulatory traits is genuinely low, setting constraints on evolutionary

responses, or whether such variation is just difficult to measure, given the challenges in

quantifying individual variation in thermoregulatory capacity. This is a genuine empirical

challengefor large vertebrates with their small population sizes and associated difficulties in

obtaining large enough sample sizes.

Here we study the potential for evolutionary responses in thermoregulation in the world's

largest bird, the ostrich (*Struthio camelus*) (Fig. 1). Estimating additive genetic variance in

thermoregulation, and its relationship to fitness, requires measuring these traits across hundreds

of individuals of known pedigree. To do this we studied a captive population of ostriches in the

Klein Karoo region of South Africa, where a large number of individuals have been reared in a

semi-natural environment over 25-years to produce a large nine-generation pedigree (Schou *et*

al., 2021, 2022). To quantify thermoregulation, we used thermal imaging (infrared camera

technology) as this technique allows the surface temperature of hundreds of individuals to be

non-invasively measured (Tattersall & Cadena, 2010). The temperature profiles of 423 adult

females were repeatedly measured an average of six times (total images=2744). Such surface

temperatures have been shown to be positively correlated with internal body temperatures in

several species of birds (Andreasson *et al.*, 2016; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022)

and insects (Tsubaki *et al.*, 2010; Svensson *et al.*, 2020). Thermal imaging has therefore

emerged as a powerful technique that has been used to quantify body temperature variation and

thermoregulatory capacity across diverse taxa under a variety of natural contexts (Tattersall *et*

al., 2009; Svensson & Waller, 2013; Andreasson *et al.*, 2016; Moore *et al.*, 2019; Svensson *et*

al., 2020; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022).

We focus on the surface temperature of the head, as shielding the brain from extreme

temperatures is of crucial importance for animals (Kilgore *et al.*, 1976; Fuller *et al.*, 2003;

Beltrán *et al.*, 2021). Combining our thermal image data with daily measurements of

reproductive success and pedigree information, we test: (1) if head temperatures are regulated

by blood circulating around the surface of the neck (Fig. 1B), and if the neck could thus serve

as a so called 'thermal radiator' as shown for other morphological traits in other animal species

(Tattersall *et al.*, 2009; Weissenböck *et al.*, 2010; Darnell & Munguia, 2011); (2) if variation in

the efficiency of head thermoregulation influences female reproductive success, which is highly

sensitive to temperature (Schou *et al.*, 2021, 2022); (3) if there is genetic variation in the

efficiency of head thermoregulation; and (4) if populations originating from regions with

different climatic regimes have evolved different head thermoregulatory capacities.

Results

Thermoregulation across naturally fluctuating temperatures

The surface temperatures (T) of the head and neck were both positively correlated with air temperature (Fig. 2; Tables S1-2). The rate of temperature change with increasing and decreasing temperatures was, however, significantly higher for the neck compared to the head (Increasing air T_{Neck} vs Increasing air T_{Head} (CI) = 1.6 (1.2, 1.9), pMCMC = 0.001; Decreasing air T_{Neck} vs Decreasing air T_{Head} (CI) = -2.1 (-3.1, -1.2), pMCMC = 0.001, Fig. 2C; Table S3). Consequently, there were large differences between neck and head temperatures at extreme temperatures (low < 20°C and high > 30°C), but at benign air temperatures (20 to 30°C), where the need for thermoregulation is minimal, neck and head temperatures were similar (Fig. 2D, Table S4).

Despite head and neck temperatures differing in their rates of change, we found no direct evidence that the neck actively regulates the surface temperature of the head. There was no accelerating change in surface temperature, as would be expected if blood flow was actively directed to the neck at critical air temperature inflection points (Fig. S1) (Tattersall *et al.*, 2009; Janse van Vuuren *et al.*, 2020). Instead, neck temperatures increased linearly with air temperature (Figs. 2A-B). This suggests that the deviation in head and neck temperature ('relative head temperature') captures the ability of individuals to thermoregulate and protect vital organs, such as the brain, from thermal stress. Furthermore, given the neck does not appear to actively regulate head temperatures, it likely provides useful information as a control on the current exposure of individuals to temperature stress. Such information can be difficult to quantify in free ranging animals due to microclimatic factors and previous behavioural activity.

Reproductive consequences and evolutionary potential of head thermoregulation

To examine if head thermoregulation has evolutionary potential, we investigated whether variation in thermoregulatory capacity, as measured by relative head temperature was heritable and related to reproductive success. We found that females with lower relative head temperatures during hot periods had higher egg-laying rates (Fig. 3A. Hot$_{\text{Head-Neck}}$ (CI) = -0.16 (-0.30,0.00); pMCMC = 0.027; Table S5), which was not the case under more benign temperatures (Benign$_{\text{Head-Neck}}$ (CI) = -0.06 (-0.18,0.05); pMCMC = 0.277; Table S5). This suggests that there is selection for the ability to cool the head during high temperatures due to its effects in promoting reproduction.

We further found that the capacity of females to regulate head temperature was heritable as revealed by positive and significant heritabilities and evolvabilities(Hansen *et al.*, 2011) (h^2 ranged from 0.06 to 0.08. Fig. 3B, Table S6, see also Table S7-S8 for separate models of head and neck). Heritability estimates were similar and significantly different from zero across cold, benign and hot conditions, although measures of additive genetic variances and repeatabilities were slightly higher under cold and hot conditions. Heritabilities are capped by how repeatable measurements are across individuals, which can often be low for labile physiological traits, such as thermal profiles when measured non-invasively under semi-natural settings where few variables are controlled (Figs. 1-2). Our repeatability estimates of the relative head temperature ranged from 0.12 to 0.17 (Fig. 3B, Table S6) which is quite low. It is therefore plausible that our heritability, while significant could underestimate the true heritability of relative head temperature due to the inherent difficulties in quantifying individual variation in thermoregulatory traits.

Thermoregulatory ability differs between populations from different climatic regions

We investigated whether head thermoregulation has evolved in response to past climate conditions by comparing the thermoregulatory ability of females from populations originating from different geographic regions (Fig. 4A-B). At our study site in South Africa, there are three different ostrich populations from different regions of Africa: South African Blacks (SAB: *Struthio camelus*), Zimbabwean Blues (ZB: *S. c. australis*) and Kenyan Reds (KR: *S. c. massaicus*) that have evolved under different climatic regimes (Fig. 4A-C). The climate in the East Africa, where KR naturally occur, is less seasonal, exhibiting lower fluctuations in temperature and precipitation, than the regions where ZB and SAB populations occur (Fig. 4C). Comparing individuals from the three populations that have all been raised in South Africa allowed us to examine population variation in head thermoregulation in a 'common garden' setting.

There were pronounced differences in head thermoregulation between the populations in response to increasing temperatures that corresponded to the stability of their past climates. The relative head temperatures of KR females that come from a stable thermal environment, changed very little in response to temperature increases (Fig. 4C, Table S11). This indicates similar rates of change in head and neck temperature with air temperature, and therefore only modest thermoregulation in response to increasing temperatures. In contrast, the SAB and ZB populations, that originate from the more variable thermal environments of southern Africa, showed a decrease in relative head temperature as temperatures increased, suggesting a greater capacity to regulate head temperatures (i.e. head temperature did not increase as fast as neck temperature, Fig. 4D). In contrast, thermoregulation during decreasing temperatures was similar across KRs, SABs and ZABs. These population differences in thermoregulatory

capacity were not confounded by differences in body size between the three populations (Tables

S12-13, Figs. S2-3).

Discussion

Our study has revealed that the capacity to regulate head temperature is important for

reproductive success under fluctuating temperatures (Fig. 3).   The efficiency of

thermoregulation appears to have a genetic basis and enables reproduction under a greater range

of temperatures (Fig. 3). Ostriches from populations that experience more pronounced

temperature fluctuations were also more efficient at regulating head temperatures, reducing

head temperatures relative to the neck. These results suggest that thermoregulation of the head

has evolved in response to past climates and may evolve in response to future climatic change

(Fig. 4).

Previous research has identified a variety of mechanisms that aid temperature regulation

(Kilgore *et al.*, 1976; Fuller *et al.*, 2003; Angiletta, 2009; Tattersall *et al.*, 2009; Symonds &

Tattersall, 2010; Weissenböck *et al.*, 2010; Périard *et al.*, 2015). While such studies have

provided compelling evidence for the physiological consequences of various thermoregulatory

traits, they have not provided direct estimates of their effect on fitness and underlying genetic

variation. Such estimates require more quantitative approaches and are necessary to fully

understand how thermoregulatory traits influence future evolutionary responses to changing

climatic conditions. Our results demonstrate the power of combining measurements of

individual reproductive success with thermal profiles with pedigree data from individuals that

have been exposed to natural temperature variation in realistic environmental settings. . The

differences (Fig. 4) between phenotypically differentiated populations from different climatic

regions together with significant additive genetic variance in head thermoregulation within

populations (Fig. 3), suggests that selection to reproduce under fluctuating past and present

climates has shaped the evolution of the ostrich's remarkable capacity for head cooling under

extreme temperatures.

Decades of research on birds in temperate zones has focused on food availability in altricial

birds as a major limiting factor for reproduction (Lack, 1954; Williams, 1966). However, for

precocial birds inhabiting tropical and subtropical areas, like the ostrich, temperature stress

during reproduction might pose a more severe challenge than food limitation (Bourne *et al.*,

2020). Thermoregulation can maintain non-lethal body temperatures (Tattersall *et al.*, 2009;

Symonds & Tattersall, 2010; Tattersall & Cadena, 2010; Darnell & Munguia, 2011; McCafferty

et al., 2011; Szafrańska *et al.*, 2020), but heat waves can still compromise population viability

by decreasing individual survival or reproductive success (Walsh *et al.*, 2019; Bourne *et al.*,

2020; Parratt *et al.*, 2021). Recent research from several taxa suggest that climate-mediated

local extinctions might already be common (Sinervo *et al.*, 2010; Wiens, 2016), with signs of

collapse of some desert bird communities being documented (Riddell *et al.*, 2019). Whether

genetic variation in adaptations underlying thermoregulation is sufficient to enable evolutionary

responses to increasingly hot and fluctuating conditions remains an open question.

The low heritability estimates of thermal profiles in this study make it difficult to predict the

evolutionary potential of ostrich populations to adapt to future temperature change. Previous

research on ectotherms such as *Drosophila* have also found low levels of additive genetic

variance and low heritabilities of thermal tolerance, revealing the challenges to obtain precise

quantitative genetic estimates (Hoffmann *et al.*, 2003; Kellermann *et al.*, 2009). Variation in

body temperatures can also be nonadaptive or maladaptive, a contention supported by recent

empirical research (Ghalambor *et al.*, 2015; Radersma *et al.*, 2020; Stamp & Hadfield, 2020;

Svensson *et al.*, 2020). For example, changes in both core body temperature and external body

parts have been shown to be related to lower fitness in insects (Svensson *et al.*, 2020), reptiles

(Campbell-Staton *et al.*, 2021), and birds (Nilsson *et al.*, 2016). Such maladaptive thermal

plasticity may reflect costs of maintaining homeostasis and stable body temperatures under

thermally stressful conditions, leading to selection for thermal robustness (Svensson *et al.*,

2020; Campbell-Staton *et al.*, 2021). It is likely that selection for thermal robustness on one

side, and costs of maintaining a stable body temperature on the other side, are shaping genetic

variance in thermoregulation in both the ostrich and other organisms inhabiting thermally

stressful environments like deserts. While challenging, combining studies on the genetics of

thermal tolerance with long-term population monitoring of reproduction and survival is key to

forecasting the potential damage caused by climate change. This is especially the case for large,

sub-tropical endotherms like the ostrich.

Methods

Study site and study populations

The study was conducted at the Oudtshoorn Research Farm in the arid Klein Karoo region of

South Africa (GPS: 33° 38' 21.5"S, 22° 15' 17.4"E). Fenced enclosures (N=181) were used to

monitor the reproductive success of ostriches that included 156 (~0.25 ha) for male-female pairs

(Cloete *et al.*, 2008) and 22 for groups (~0.47 ha). All individuals had access to *ad libitum* food

and water. The ostrich individuals in this study belong to three different subspecies, hereafter

referred to as populations: 1) the Masai ostrich (*Struthio camelus massaicus*), or Kenyan Red

(KR), 2) the Southern African ostrich, (*S. c. australis*), sometimes referred to as the

Zimbabwean Blue (ZB) because of its origin in Namibia and Zimbabwe, and 3) the South

African Blacks (SAB), that is thought to be of mixed origin, but is genetically very similar to

ZB (Davies et al 2012; unpublished data). SAB are also referred to as *S. c. var. domesticus*.

Individuals that had less than 85% expected relatedness to one population, as determined by the

pedigree (see below), were considered hybrids. Breeding birds were recruited from surviving

chicks from previous years, and parentage data were used to compile a 9-generation pedigree

with 139 founding individuals. Ethical clearance was obtained from the Western Cape

Department of Agriculture (DECRA R12/48).

Thermal imaging data

From 2012 to 2018 we took thermal images of ostriches in the enclosures using an infrared

thermography camera (H2640, NEC Avio Infrared Technologies). Pictures were taken during

October to December, usually from distances between 2m and 25m. We used the software

InfRec Analyzer to draw separate polygons within the head and neck of the ostrich in each

image, and the average temperature of these polygons were extracted as individual head and

neck surface temperatures, respectively. We used the same procedures and same default settings

of the software as in our previous work (Svensson & Waller, 2013; Svensson *et al.*, 2020),

assuming an emissivity of 1. As a measure of air temperature, we used estimates of hourly

temperatures from a weather station positioned 600 m from the field site. We fitted a cubic

spline to the hourly temperature estimates of each day using the R-package *mgcv* v.1.8 (Wood,

2004), from which we extracted the predicted air temperature at the time-points when thermal

images were taken.

Datasets and analyses

Quantifying rate of change in head and neck surface temperatures

From 2012 to 2017, we took 2744 pictures of 423 females from early morning till late afternoon

(average images per female = 6.5). This dataset was designed with the aim of having a high

number of individuals with repeated sampling within and across years, such that each individual

was monitored in different thermal environments, but with little repeated sampling within days

(8% of the pictures).

With this dataset we modelled the change in surface temperatures of the head and neck in

response to air temperature using generalized linear mixed models (GLMMs). A previous

investigation showed reproductive success of ostriches is highest at a daily maximum

temperature of ~20°C (Schou *et al.*, 2021). We therefore defined 20°C as the optimum air

temperature for ostriches, and calculated absolute *temperature change* away from this optimum,

defining the factor *direction* to denote whether the change in air temperature was due to a

decrease or increase from the optimum. To make the intercept of statistical models represent

the most benign air temperature, we set 20°C to 0 and calculated deviations above (increases)

and below (decreases) this value. The variance of slopes (see below) depends on the scale of

the environmental parameter, and we therefore standardized our data by dividing it by the

maximum of the air temperature range, resulting in 1 being the maximum temperature change.

Head and neck temperatures were modeled as Gaussian traits in separate models. Models

included the fixed effects of temperature change (ranging from 0 to 1) and direction (decreases

or increases). The interaction between temperature change and direction was modelled with a

common intercept for decreases and increases, as the way temperature change was calculated

dictated that the intercepts were identical. We included the fixed effects of population (SAB,

ZB, KR or hybrids) and both the linear and quadratic terms of time of day (scaled and centered

to a mean of zero and unit variance). We included interactions between population, temperature

change and direction.

We accounted for environmental effects that varied across years, such as diet, by including year

as a random effect. Photographs were taken across 48 days, and we therefore included date as

a random effect. We also added enclosure as a random effect as the enclosures varied in

vegetation cover, potentially impacting on the local climatic conditions, and were repeatedly

used across years. Temperature change and direction were interacted with individual ID, to

allow independent rates of change in surface temperature of each female. This was modelled as

a 3x3 unstructured variance-covariance matrix.

GLMMs were run in R v.3.6.0 (R Core Team, 2020) using the Bayesian framework

implemented in the R-package MCMCglmm v.2.29 (Hadfield, 2010). For random terms we

used the weakly informative inverse-Gamma distribution (scale = 0.001, shape = 0.001, i.e. *V*

= diag(n), nu =n-1+0.002, with n being the dimension of the matrix) as priors. Each model was

run for 5,100,000 iterations of which the initial 100,000 were discarded and only every 4,000th

iteration was used for estimating posterior probabilities. The number of iterations was based on

inspection of autocorrelation among posterior samples in preliminary runs. Convergence of the

estimates was checked by running the model three times and inspecting the overlap of estimates

in trace plots and the level of autocorrelation among posterior samples. Posterior mode and 95%

credible intervals are reported for random effects.

3.2) Determining the impact of body mass on variation in surface temperatures

To test if differences in the rate of change in surface temperatures with air temperature were

caused by differences in body mass, we ran a separate set of models including individual body

mass as an additional fixed effect. We had records of body mass for all individuals and when

multiple records were available for one individual, we used the record closest to the time of the

thermal image. Models for surface temperature of the head and neck were implemented using

the model structure described above, but with body mass (scaled and centered to a mean of zero

and unit variance) interacted with all the previously described fixed effects.

Comparing the neck and the head

We compared the rate of surface temperature change between the neck and head. This was done

by running a model that included body part (neck or head) as a fixed effect. The models were

implemented using the model structure described above, but with body part interacted with all

the previously described fixed effects. We also added image as an additional random effect

because head and neck surface temperatures were derived as pairs from the same image.

The change in relative head temperature when hot or cold

Results from the analyses above indicated that the head is more robust to changes in air

temperature than the neck, potentially indicating the head is protected from temperature

fluctuations. However microclimatic factors (i.e. the operative environmental temperature) and

the behaviour up until the time of the thermal image will influence estimates of absolute head

surface temperatures. To better estimate the capacity of individuals to regulate head

temperature, we compared the temperature of the head to the neck (head-neck surface

temperature) to control for differences in operative environmental temperatures. Positive values

therefore indicated that the neck was warmer than the head, and negative values that the head

was warmer than the neck. To test if the head-neck temperature at high air temperatures differs

from benign and cold air temperatures, we grouped air temperature into three categories: Cold

(air temperatures <= 20°C, n_{images} = 849), Benign (air temperatures > 20°C & < 30°C, n_{images} =

1069) and Hot (air temperatures >= 30°C, n_{images} = 826). The grouping of air temperature into

these three categories was based on the thermal neutral zone of the emu (Maloney, 2008), and

to ensure roughly equal replication within the cold and hot categories. These categories do not

capture the effects of small deviations in air temperature, and we therefore added a continuous

measure of the deviation from the mean air temperature in each category. This variable was

constructed by centering and scaling the air temperature records within each air temperature

category. We modelled the response of the head-neck temperature (Gaussian) to air temperature

category by following the same general approach as above. The major difference was that air

temperature category was included as a fixed factor (instead of air temperature change and

direction) and interacted with female ID to generate a 3x3 unstructured variance-covariance

matrix composed of the cold, benign and hot temperature categories. We also estimated the

residual variance separately for each air temperature category.

Effect of neck and head surface temperatures on reproductive success

To test if the capacity to regulate head temperature influences reproductive success we analysed

its relationship with rates of egg-laying. We connected thermal measurements of individual

females to their egg-laying records. From previous investigations we know that when daily

maximum temperature is above or below 20°C ostrich egg laying-rate starts to slowly decrease

two to four days later, possibly because this is the time it takes for the egg to travel down the

oviduct (Schou *et al.*, 2021). The maximum temperature was below 20°C in just two of the days

during which we monitored thermoregulation, but maximum temperature was above 20°C in

41 days. We therefore monitored whether any eggs were laid two to four days after females

were thermal imaged at days reaching more than 20°C. If females with higher neck than head

surface temperatures have a higher probability of laying an egg during this time window, this

would indicate that the thermoregulatory capacity of the head is indeed important for

reproductive success. We are, however, careful with inferring causation, as it is possible that a

third confounding factor, such as metabolic rate, may generate an autocorrelation between head

regulation and laying rates. To overcome this, we grouped the data into photographs taken

during hot and benign times of the days. If an elevated neck to head temperature ratio reflects

increased tolerance to heat, then its positive relationship with egg-laying should be most

pronounced when hot.

Egg laying rates were analysed using the model structure described above, but with the

following modifications: a) the probability of laying (binary, model type: "threshold") was used

as the response variable, b) the fixed effects included were head-neck temperature difference

(scaled and centered to a mean of zero and unit variance), air temperature category (hot or

benign), and population (KR, SAB, ZB or Hybrid), as well as the interaction between head-

neck temperature and temperature category, c) as females of two years of age lay fewer eggs

than older females (Schou *et al.*, 2021), a factor of age (2 versus >2) was also included as a

fixed effect, and d) year, date, enclosure and individual ID were included as random effects.

The first 45 days of the breeding season were removed as this is the average time it takes for

pairs to acclimate to each other and their enclosure (Schou *et al.*, 2021). We also removed

females that laid fewer than ten eggs per year (5% of all females) to avoid including females

from incompatible pairs, and individuals that did not enter the breeding state. Each model was

run for 31,500,000 iterations of which the initial 1,500,000 were discarded and only every

10,000th iteration was used for estimating posterior probabilities.

Quantifying the evolutionary potential of relative head temperature

Using the model of the change in head-neck temperature in each air temperature category (see

above), we estimated the repeatability (R) of the head-neck temperature under different

temperature conditions. This was done using the estimates of permanent individual variances

(*pe*) estimated in the variance-covariance matrix of individual ID by air temperature category:

$$(1)\ R = \frac{\sigma^2_{pe}}{\sigma^2_{pe}+\sigma^2_{year}+\sigma^2_{enclosure}+\sigma^2_{res}}.$$

The individual variance and covariance in head-neck temperatures may originate from both

environmental and genetic factors. To partition the among-individual variance that is due to

additive genetic effects we added a second 3x3 unstructured variance-covariance matrix of

individual ID linked to the pedigree (*a*). With these variance components, we estimated the

narrow sense heritability (h^2) of the head-neck temperature in each air temperature category as

the proportion of phenotypic variance attributable to additive genetic variance:

$$(2)\ h^2 = \frac{\sigma^2_a}{\sigma^2_{pe} + \sigma^2_a + \sigma^2_{year} + \sigma^2_{camp}+\sigma^2_{res}}$$

We also estimated evolvability (I_A) (Houle, 1992):

$$(3)\ I_A = \frac{\sigma^2_a}{trait\ mean^2} * 100$$

One characteristic of evolvability is that it increases very fast as trait mean approaches zero.

When we initially used the posterior of the trait mean to estimate evolvability, this caused near

infinity estimates of evolvability for some of the samples in the posterior, causing biased

estimates of the posterior mode and mean of evolvability. To avoid this, we used the posterior

mode of trait mean in the denominator, such that only the uncertainty of additive genetic

variance is included in the reported estimate of evolvability. We also ran identical models with

head or neck surface temperature as the response variable. The outcome of these analyses is

available in the supplementary materials (Tables S7-S8).

Finally, we modelled head-neck temperature in a random regression model, using the model

structure described above. This approach was taken to investigate if the head-neck temperature

at the optimum temperature (the intercept) influences the rate of change in head-neck

temperatures as air temperatures increase or decrease (the slopes). To test if such a relationship

is driven by different families, we added a second 3x3 unstructured variance-covariance matrix

of individual ID linked to the pedigree and interacted with temperature change and direction.

The genetic variance and co-variance was then used to estimate the genetic correlation between

the slopes and intercepts (correlation = $\text{covariance}_{\text{trait1,trait2}}$ / sqrt($\text{var}_{\text{trait1}}$*$\text{var}_{\text{trait2}}$)) (Table S9).

Current distribution of ostrich populations

We obtained estimates of current distributions of *S. c. massaicus* (KR) and *S. c. australis* (ZB).

This was done by downloading region-based presence/absence data from Avibase

(https://avibase.bsc-eoc.org, September, 2020) and plotting these using the R-package

"rnaturalearth" v. 0.1.0. To identify climatic differences in their distributions we downloaded

19 bioclimatic variables (10min) from WorldClim v. 2 (Fick & Hijmans, 2017). We performed

a Principal Component Analyses (PCA) and inspected the loadings of the first four principal

components after varimax transformation (Table S10). Based on this inspection we described

each principal component by one or two bioclimatic variables to characterize climatic

differences.

Acknowledgements

We thank the staff and workers at Oudtshoorn Research Farm for assistance with data collection and maintenance of the birds and the Western Cape Government for use of their resources. We also thank members of "Svensson Lab" (students and assistants) who helped with analyzing the thermal images, and Andreas Nord and Fredrik Andreasson for valuable feedback on an earlier version of this manuscript. The computations were performed on resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under Project SNIC 2018/8-359.

Author Contribution

Conceptualization: E.I.S, C.K.S., M.F.S.; Data curation: E.I.S., C.K.C., M.F.S., A.E., Z.B., S.C.; Formal analysis: M.F.S.; Funding acquisition: C.K.C, E.I.S., M.F.S, S.C.; Investigation: E.I.S., C.K.C., J.M., J.W., M.F.S., S.C., Z.B; Methodology: E.I.S., C.K.C., M.F.S.; Project administration: E.I.S., C.K.C., M.F.S.; Writing- original draft E.I.S, C.K.C., M.F.S.; Writing – reviewing and editing: A.E., CKC, E.I.S., J.M., J.W., M.F.S, S.C., Z.B.

Funding

Carlsberg Foundation (MFS)

Swedish Research Council grant 2017-03880 (CKC)

Knut and Alice Wallenberg Foundation grant 2018.0138 (CKC)

Carl Tryggers grant 12: 92 & 19: 71 (CKC)

Carl Tryggers grant 19: 71 (CKC)

Swedish Research Council grant 2016-03356 (EIS)

Western Cape Agricultural Research Trust grant 0070/000VOLSTRUISE (SC)

Technology and Human Resources for Industry program of the South African National

Research Foundation grant TP14081390585) (SC)

Data Accessibility

All data extracted from thermal images are available at https://osf.io/fu2wx/ and will be

uploaded in an open digital repository such as Dryad Digital Repository

(https://datadryad.org/stash) upon acceptance on this manuscript. The remaining data used

support the findings of this study are available from the Western Cape Department of

Agriculture in South Africa (WCDA). Restrictions apply to the use of some of these data, which

are thus not publicly available. These data are however available from the WCDA

upon reasonable request. Code for analyses is available on Github:

www.github.com/abumadsen/thermal-images-ostrich/tree/main

Competing interests

Authors declare that they have no competing interests.

Supporting Information

Tables S1-S15

Figures S1-S8

References

Andreasson, F., Nord, A. & Nilsson, J.-Å. (2016) Brood size constrains the development of endothermy in blue tits. *Journal of Experimental Biology*, **219**, 2212–2219.

Angiletta, M.J. (2009) *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press, Oxford.

Atwood, T.B., Valentine, S.A., Hammill, E., McCauley, D.J., Madin, E.M.P., Beard, K.H., *et al.* (2020) Herbivores at the highest risk of extinction among mammals, birds, and reptiles. *Science Advances*, **6**, eabb8458.

Beltrán, I., Herculano-Houzel, S., Sinervo, B. & Whiting, M.J. (2021) Are ectotherm brains vulnerable to global warming? *Trends in Ecology & Evolution*, **36**, 691–699.

Bogert, C.M. (1949) Thermoregulation in reptiles, a factor in evolution. *Evolution*, **3**, 195–211.

Bourne, A.R., Cunningham, S.J., Spottiswoode, C.N. & Ridley, A.R. (2020) Hot droughts compromise interannual survival across all group sizes in a cooperatively breeding bird. *Ecology Letters*, **23**, 1776–1788.

Campbell-Staton, S.C., Velotta, J.P. & Winchell, K.M. (2021) Selection on adaptive and maladaptive gene expression plasticity during thermal adaptation to urban heat islands. *Nature Communications*, **12**, 6195.

Cardillo, M., Mace, G.M., Jones, K.E., Bielby, J., Bininda-Emonds, O.R.P., Sechrest, W., *et al.* (2005) Multiple causes of high extinction risk in large mammal species. *Science*, **309**, 1239–1241.

Cloete, S.W.P., Engelbrecht, A., Olivier, J.J. & Bunter, K.L. (2008) Deriving a preliminary breeding objective for commercial ostriches: an overview. *Australian Journal of Experimental Agriculture*, **48**, 1247–1256.

Darnell, M.Z. & Munguia, P. (2011) Thermoregulation as an alternate function of the sexually dimorphic Fiddler Crab claw. *American Naturalist*, **178**, 419–428.

Fick, S.E. & Hijmans, R.J. (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, **37**, 4302–4315.

Fuller, A., Kamerman, P.R., Maloney, S.K., Mitchell, G. & Mitchell, D. (2003) Variability in brain and arterial blood temperatures in free-ranging ostriches in their natural habitat. *Journal of Experimental Biology*, **206**, 1171–1181.

Gauchet, L., Jaeger, A. & Grémillet, D. (2022) Using facial infrared thermography to infer avian body temperatures in the wild. *Marine Biology*, **169**, 57.

Ghalambor, C.K., Hoke, K.L., Ruell, E.W., Fischer, E.K., Reznick, D.N. & Hughes, K.A. (2015) Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*, **525**, 372–375.

Hadfield, J.D. (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The **MCMCglmm** R Package. *Journal of Statistical Software*, **33**, 1–22.

Hansen, T.F., Pelabon, C. & Houle, D. (2011) Heritability is not evolvability. *Evolutionary Biology*, **38**, 258–277.

Hoffmann, A.A., Hallas, R.J., Dean, J.A. & Schiffer, M. (2003) Low potential for climatic stress adaptation in a rainforest *Drosophila* species. *Science*, **301**, 100–102.

Houle, D. (1992) Comparing evolvability and variability of quantitative traits. *Genetics*, **130**, 195–204.

Huey, R.B., Hertz, P.E. & Sinervo, B. (2003) Behavioral drive versus behavioral inertia in evolution: a null model approach. *American Naturalist*, **161**, 357–366.

Hurley, L.L., McDiarmid, C.S., Friesen, C.R., Griffith, S.C. & Rowe, M. (2018) Experimental heatwaves negatively impact sperm quality in the zebra finch. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20172547.

Janse van Vuuren, A.K., Kemp, L.V. & McKechnie, A.E. (2020) The beak and unfeathered skin as heat radiators in the southern ground-hornbill. *Journal of Avian Biology*, **51**.

Kellermann, V., Heerwaarden, B. van, Sgro, C.M. & Hoffmann, A.A. (2009) Fundamental evolutionary limits in ecological traits drive *Drosophila* species distributions. *Science*, **325**, 1244–1246.

Kellermann, V., Loeschcke, V., Hogenkamp, D., Kristensen, T.N., Flojgaard, C., David, J.R., *et al.* (2012a) Phylogenetic constraints in key functional traits behind species' climate niches: patterns of dessication and cold resistance across 95 *Drosophila* species. *Evolution*, **66**, 3377–3389.

Kellermann, V., Overgaard, J., Hoffmann, A.A., Flojgaard, C., Svenning, J.C. & Loeschcke, V. (2012b) Upper thermal limits of Drosophila are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 16228–16233.

Kilgore, D.L., Bernstein, M.H. & Hudson, D.M. (1976) Brain temperatures in birds. *Journal of comparative physiology*, **110**, 209–215.

Lack, D. (1954) *The natural regulation of animal numbers*. Clarendon, Oxford.

Maloney, S.K. (2008) Thermoregulation in ratites: a review. *Australian Journal of Experimental Agriculture*, **48**, 1293–1301.

McCafferty, D.J., Gilbert, C., Paterson, W., Pomeroy, P.P., Thompson, D., Currie, J.I., *et al.* (2011) Estimating metabolic heat loss in birds and mammals by combining infrared thermography with biophysical modelling. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, The challenge of measuring energy expenditure: current field and laboratory methods, **158**, 337–345.

Moore, M.P., Lis, C., Gherghel, I. & Martin, R.A. (2019) Temperature shapes the costs, benefits and geographic diversification of sexual coloration in a dragonfly. *Ecology Letters*, **22**, 437–446.

Moreira, M.O., Qu, Y.-F. & Wiens, J.J. (2021) Large-scale evolution of body temperatures in land vertebrates. *Evolution Letters*, **5**, 484–494.

Nilsson, J.-Å., Molokwu, M.N. & Olsson, O. (2016) Body temperature regulation in hot environments. *PLoS ONE*, **11**, e0161481.

Parratt, S.R., Walsh, B.S., Metelmann, S., White, N., Manser, A., Bretman, A.J., *et al.* (2021) Temperatures that sterilize males better match global species distributions than lethal temperatures. *Nature Climate Change*, **11**, 481–484.

Périard, J.D., Racinais, S. & Sawka, M.N. (2015) Adaptations and mechanisms of human heat acclimation: Applications for competitive athletes and sports. *Scandinavian Journal of Medicine & Science in Sports*, **25**, 20–38.

R Core Team. (2020) *R: A language and environment for statistical computing*. R foundation for statistical computing, Vienna, Austria.

Radersma, R., Noble, D.W.A. & Uller, T. (2020) Plasticity leaves a phenotypic signature during local adaptation. *Evolution Letters*, **4**, 360–370.

Riddell, E.A., Iknayan, K.J., Wolf, B.O., Sinervo, B. & Beissinger, S.R. (2019) Cooling requirements fueled the collapse of a desert bird community from climate change. *Proceedings of the National Academy of Sciences of the United States of America*, **116**, 21609–21615.

Sales, K., Vasudeva, R., Dickinson, M.E., Godwin, J.L., Lumley, A.J., Michalczyk, Ł., *et al.* (2018) Experimental heatwaves compromise sperm function and cause transgenerational damage in a model insect. *Nature Communications*, **9**, 4771.

Schou, M.F., Bonato, M., Engelbrecht, A., Brand, Z., Svensson, E.I., Melgar, J., *et al.* (2021) Extreme temperatures compromise male and female fertility in a large desert bird. *Nature Communications*, **12**, 666.

Schou, M.F., Engelbrecht, A., Brand, Z., Svensson, E.I., Cloete, S. & Cornwallis, C.K. (2022) Evolutionary trade-offs between heat and cold tolerance limit responses to fluctuating climates. *Science Advances*, **8**, eabn9580.

Sinervo, B., Mendez-de-la-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagran-Santa, C.M., *et al.* (2010) Erosion of lizard diversity by climate change and altered thermal niches. *Science*, **328**, 894–899.

Stamp, M.A. & Hadfield, J.D. (2020) The relative importance of plasticity versus genetic differentiation in explaining between population differences; a meta-analysis. *Ecology Letters*, **23**, 1432–144.

Svensson, E.I., Gomez-Llano, M.A. & Waller, J.T. (2020) Selection on phenotypic plasticity favors thermal canalization. *Proceedings of the National Academy of Sciences of the United States of America*, **117**, 29767–29774.

Svensson, E.I. & Waller, J.T. (2013) Ecology and sexual selection: evolution of wing pigmentation in calopterygid damselflies in relation to latitude, sexual dimorphism and speciation. *American Naturalist*, **182**, E174–E195.

Symonds, M.R.E. & Tattersall, G.J. (2010) Geographical variation in bill size across bird species provides evidence for Allen's Rule. *American Naturalist*, **176**, 188–197.

Szafrańska, P.A., Andreasson, F., Nord, A. & Nilsson, J.-Å. (2020) Deep body and surface temperature responses to hot and cold environments in the zebra finch. *Journal of Thermal Biology*, **94**, 102776.

Tattersall, G.J., Andrade, D.V. & Abe, A.S. (2009) Heat exchange from the Toucan bill reveals a controllable vascular thermal radiator. *Science*, **325**, 468–470.

Tattersall, G.J. & Cadena, V. (2010) Insights into animal temperature adaptations revealed through thermal imaging. *Imaging Science Journal*, **58**, 261–268.

Tsubaki, Y., Samejima, Y. & Siva-Jothy, M.T. (2010) Damselfly females prefer hot males: higher courtship success in males in sunspots. *Behavioral Ecology and Sociobiology*, **64**, 1547–1554.

Walsh, B.S., Parratt, S.R., Hoffmann, A.A., Atkinson, D., Snook, R.R., Bretman, A., *et al.* (2019) The impact of climate change on fertility. *Trends in Ecology & Evolution*, **34**, 249–259.

Weissenböck, N.M., Weiss, C.M., Schwammer, H.M. & Kratochvil, H. (2010) Thermal windows on the body surface of African elephants (Loxodonta africana) studied by infrared thermography. *Journal of Thermal Biology*, **35**, 182–188.

Wiens, J.J. (2016) Climate-related local extinctions are already widespread among plant and animal species. *PLOS Biology*, **14**, e2001104.

Williams, G.C. (1966) Natural selection, the costs of reproduction, and a refinement of Lack's principle. *American Naturalist*, **100**, 687–690.

Williams, G.C. (1992) *Natural selection: Domains, levels and challenges*. Oxford University Press, Oxford.

Wood, S.N. (2004) Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association*, **99**, 673–686.

Figures and captions

Fig. 1. The ostrich inhabits thermally challenging desert environments such as the Karoo in South Africa.

A. The ostrich (*Struthio camelus*) is the world's largest living bird with a feathered body and featherless neck and

head. *Left*: female. *Right:* male (photograph by C. K. Cornwallis in Karoo National Park, Western Cape Province

of South Africa). **B.** Thermal image of a female (left) and male (right) ostrich in our study population at Oudtshoorn

(photograph by E. I. Svensson). Note how the long neck is hot and emits excess heat (red, warm colour). **C.** The

dry and tree-less semi-desert environment around the study site is characterized by extensive temperature

fluctuations ranging from -5°C to 45°C, causing extreme thermal stress (Schou *et al.*, 2021) (photograph by E. I.

Svensson).

Fig. 2. Neck temperatures show how ostriches thermoregulate their head temperatures. A) When air temperatures increase above 20°C, the surface temperatures of the head increase slower than the neck (n_{images} = 1895, Tables S1-S2). **B)** Similarly, when air temperatures decrease below 20°C the reduction in surface temperatures are greatest in the neck (n_{images} = 849). Three extreme datapoints in A were removed for graphical purposes (see Fig. S4). The larger points are averages with standard errors binned according to temperature. Point size illustrates relative number of individuals: smallest point = 2 and largest point = 71. **(C)** The rate of surface temperature change was steeper for the neck compared to the head (Table S3). This difference in rates of temperature increase between the two body parts was consistent across both decreasing and increasing temperatures ($n_{individuals}$ = 423). **(D)** The difference in rates of temperature increase between the head and neck led to higher discrepancy between neck to head surface temperatures during hot conditions (air temperatures >= 30°C, n_{images} = 826) and cold (air temperatures <= 20°C, n_{images} = 849) compared to benign (air temperatures > 20°C & < 30°C, n_{images} = 1069) conditions (Table S4).

Fig. 3. The evolutionary potential for head temperature regulation. The capacity of individuals to regulate their head temperature was estimated as the difference between head and neck temperatures ('relative head temperature') in relation to air temperature. **A)** On hot days (daily maximum > 20°C, n_{days} = 35), when temperatures were highest (air temperatures >= 30°C, n_{images} = 471, $n_{females}$ = 228) an increased capacity to regulate head temperature (i.e. a lower head than neck temperature) was associated with a higher egg-laying rate two days later. There is no such association when examining thermoregulation at times when temperatures were benign of the hot days (air temperatures = 20-30°C, n_{images} = 615, $n_{females}$ = 220) (Table S5). Maximum temperatures below 20°C were recorded only on two days, preventing us from similar analyses on importance of thermoregulation during the cold times of cold days. **B)** Repeatability and heritability of relative head temperature were estimated at different air temperatures measured at the time when thermal images were taken (cold < 20°C, benign = 20-30°C and hot >= 30°C, Table S6). Note that the heritability of head temperatures under cold conditions could be estimated because there were cold periods (typically mornings) on days where temperatures exceeded 20°C. This contrasts to selection analyses where temperature per day had to be used because of the two day time lag between egg formation and egg laying.

Fig. 4. Ostrich populations originating from more variable climates exhibit larger decreases in relative head temperatures when hot. **(A-C)** Kenyan Reds (KR) inhabit eastern Africa that is less seasonal and has lower temperature fluctuations compared to Southern Africa where Zimbabwean Blues (ZB) and South African Blacks (SAB) occur. Distribution ranges in A were estimated from regional presence/absence data from Avibase (https://avibase.bsc-eoc.org) and climatic data was obtained from WorldClim (Fick & Hijmans, 2017). Average temperature and seasonality were the main bioclimatic variables of PC1 (43%) and PC2 (29%) in a PCA of 19 bioclimatic variables (Fig. S5 & Table S10). **(D)** For both the SAB ($n_{individuals}$ = 264) and ZB ($n_{individuals}$ = 36) populations the relative head temperatures (head-neck) was reduced when temperatures increased, but this was not the case for KR ($n_{individuals}$ = 23) (Table S11, see also Figs. S6-7 which includes hybrid individuals).