Heritable variation in thermoregulation is associated with reproductive success in the world's largest bird

Erik I. Svensson [*, a], Mads F. Schou[*,a,b], Julian Melgar [a], John Waller [c], Anel Engelbrecht [d], Zanell Brand [d], Schalk Cloete [d, e] & Charlie K. Cornwallis [a]

[a] Department of Biology, Lund University, Lund, Sweden
[b] Department of Biology, Aarhus University, Aarhus, Denmark
[c] Global Biodiversity Information Facility (GBIF), Copenhagen, Denmark
[d] Directorate Animal Sciences, Western Cape Department of Agriculture, South Africa
[e] Department of Animal Sciences, University of Stellenbosch, Matieland, South Africa
* These authors contributed equally

Manuscript type: *Letter*

Short running title: *Evolution of thermoregulation in the ostrich*

Keywords: birds, deserts, fitness, heat stress, heritability, ostrich, *Struthio camelus*, thermal imaging, thermoregulation, thermal adaptation

Abstract word count: 153

Total word count: 6458

Abstract

Organisms inhabiting extreme thermal environments, such as desert birds, have evolved spectacular adaptations to thermoregulate during hot and cold conditions. However, our knowledge of selection for thermoregulation and the potential for evolutionary responses is limited, particularly for large organisms experiencing extreme temperature fluctuations. Here we use thermal imaging to quantify selection and genetic variation in thermoregulation in ostriches (*Struthio camelus*), the world's largest bird species that is experiencing increasingly volatile temperatures. We found that females that are better at regulating their head temperatures ('thermoregulatory capacity') had higher egg-laying rates under hotter conditions. Thermoregulatory capacity was both heritable and showed signatures of local adaptation: females originating from more unpredictable climates were better at regulating their head temperatures in response to temperature fluctuations. Together these results reveal that past and present evolutionary processes have shaped genetic variation in thermoregulatory capacity, which appears to protect critical organs, such as the brain, from extreme temperatures during reproduction.

Impact Summary

Large animals inhabiting extreme thermal environments, such as deserts, are predicted to be

particularly vulnerable to the increasing temperature fluctuations expected in the future.

However, previous work on the evolutionary potential of thermoregulation has primarily

focused on the effect of hot temperatures on the survival of small ectotherms. We know little

about how large endothermic vertebrates, such as birds, will respond to changing temperatures.

Here we study the ostrich (*Struthio camelus*), the world's largest bird, that inhabits some of the

hottest and driest regions on Earth. We show that the ability of females to reproduce during hot

conditions is associated with the regulation of their head temperatures ('thermoregulatory

capacity'). Furthermore, variation in thermoregulation is heritable and related to past climatic

conditions: females originating from parts of Africa with more extreme temperature

fluctuations were better able to thermoregulate, indicating local adaptation to different climatic

conditions. Together, these results suggest that thermoregulation in this large desert bird has

evolved in response to past climatic conditions, remains genetically variable and is currently

under selection through its effect on reproduction.

Introduction

Organisms need to manage heat and cold stress to survive and reproduce in variable climates (Angiletta, 2009). The universal challenge of coping with thermal stress (Hurley *et al.*, 2018; Sales *et al.*, 2018; Walsh *et al.*, 2019; Schou *et al.*, 2021) has promoted the evolution of adaptations to regulate body temperature. For example, body temperature can be maintained by behavioural thermoregulation mechanisms, such as habitat selection (Bogert, 1949; Huey *et al.*, 2003; Angiletta, 2009; Muñoz & Losos, 2018; Muñoz, 2022), or by physiological thermoregulation, such as selective brain cooling (Fuller *et al.*, 2003) and sweating (Périard *et al.*, 2015). Other thermoregulatory mechanisms involve morphological traits, such as the ears of African Elephants (Weissenböck *et al.*, 2010) and the bills (Tattersall *et al.*, 2009, 2018; Symonds & Tattersall, 2010) and featherless skin patches of birds (McCafferty *et al.*, 2011; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022). Such morphological structures can act as so called 'thermal radiators' or 'thermal windows" that disperse excess heat in an adaptive fashion (Tattersall *et al.*, 2009; Symonds & Tattersall, 2010; Weissenböck *et al.*, 2010; Darnell & Munguia, 2011; Eastick *et al.*, 2019).

Here, we define thermoregulation as the ability of an organism to maintain body temperatures within a relatively narrow thermal zone (Deutsch *et al.*, 2008) favourable for survival or reproduction (Huey *et al.*, 2003; Muñoz, 2022). This general definition is applicable to both ecto- and endothermic animals and to both physiological and behavioural mechanisms of thermoregulation (Angiletta, 2009). The mechanisms by which animals thermoregulate and how these mechanisms promote survival have been intensely studied. However, we know much less about the quantitative genetics of thermoregulatory traits and how such traits affect reproductive success, particularly for organisms living in the regions most affected by climate change (Walsh *et al.*, 2019), such as deserts (Bourne *et al.*, 2020; Schou *et al.*, 2021).

The evolutionary potential of populations to adapt to thermal stressors, such as heatwaves and

cold snaps, depends on genetic variance in traits involved in thermoregulation (Hurley *et al.*,

2018; Sales *et al.*, 2018; Bourne *et al.*, 2020). Current empirical evidence from insects and

reptiles suggests that additive genetic variances and hence evolutionary potential in thermal

tolerance traits can be high (Ma *et al.*, 2014), but are more often low (Hoffmann *et al.*, 2003;

Kellermann *et al.*, 2009; Logan *et al.*, 2018; Castañeda *et al.*, 2019). In addition, body

temperatures and thermal adaptations are typically strongly phylogenetically conserved

(Kellermann *et al.*, 2012a, 2012b) and evolve slowly compared to other traits (Moreira *et al.*,

2021). The late evolutionary biologist George C. Williams even questioned if endothermic body

temperatures could evolve at all, suggesting that there may be evolutionary stasis in body

temperatures and thermal adaptations (Williams, 1992). However, it is unclear if genetic

variance in thermal traits is genuinely low, setting constraints on evolutionary responses, or

whether such variation is just difficult to measure, given the challenges in quantifying variation

in thermoregulatory capacity across large numbers of individuals. This is a genuine empirical

challenge for research on large vertebrates with small population sizes where it is difficult to

obtain large sample sizes. To our knowledge there is a lack of quantitative genetic studies on

large endothermic animals, which is unfortunate, as they are likely to be particularly vulnerable

to changing temperatures.

Large animal bodies have a higher thermal inertia and slower rate of body temperature change

compared to small bodies (Bogert, 1949; Angiletta, 2009). Thermal inertia can help maintain

body temperatures during cold conditions, but it can also increase physiological stress under

sustained heat, jeopardising survival and reproduction (Hurley *et al.*, 2018; Sales *et al.*, 2018;

Bourne *et al.*, 2020; Parratt *et al.*, 2021; Schou *et al.*, 2021, 2022). Large animals are also

predicted to be particularly vulnerable to changing climates because their rate of adaptive

evolution is potentially limited by longer generation times and lower population sizes (Cardillo

et al., 2005; Atwood *et al.*, 2020). Understanding how large-bodied endothermic animals evolve

to cope with thermal stress in fluctuating thermal environments therefore requires particular

attention.

Here we study the evolutionary potential of thermoregulation in the world's largest bird, the

ostrich (*Struthio camelus*) (Fig. 1). Estimating additive genetic variance in thermoregulation,

and its relationship to fitness, requires measuring these traits across hundreds of individuals of

known pedigree. To do this we studied captive breeding populations of ostriches in the Klein

Karoo region of South Africa, where a large number of individuals have been reared in a semi-

natural environment over 25-years to produce a large nine-generation pedigree (Schou *et al.*,

2021, 2022). To quantify thermoregulation, we used thermal imaging (infrared camera

technology) which makes it possible to measure the surface temperature of hundreds of

individuals in a non-invasive way (Tattersall & Cadena, 2010; McCafferty *et al.*, 2015; Gauchet

et al., 2022). The temperature profiles of 423 adult females were repeatedly measured an

average of six times (n_{images} = 2744). Skin surface temperatures have been shown to be

significantly and positively correlated with internal body temperatures in studies on both

vertebrates and invertebrates (Tsubaki *et al.*, 2010; Andreasson *et al.*, 2016; Svensson *et al.*,

2020; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022). In some species of birds, such as red-footed

boobies (*Sula sula*), the correlation between skin surface temperatures and internal body

temperatures (T_b) have been found to be very strong, with body temperatures explaining 76-82

% of the variation (Gauchet *et al.*, 2022). Thermal imaging has therefore emerged as a powerful

and non-invasive technique that has been used to quantify individual variation in thermal

profiles and thermoregulatory capacity across diverse taxa under a variety of natural contexts

(Tattersall *et al.*, 2009; Svensson & Waller, 2013; Andreasson *et al.*, 2016; Moore *et al.*, 2019;

Svensson *et al.*, 2020; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022).

We combined thermal image data with daily measurements of reproductive success and

pedigree information to: (1) examine the regulation of head and body surface temperatures

across ambient temperatures. We focus specifically on the surface temperature of the head, as

shielding the brain from extreme temperatures is of crucial importance for animals (Kilgore *et*

al., 1976; Fuller *et al.*, 2003; Beltrán *et al.*, 2021); (2) test if variation in the efficiency of head

thermoregulation influences female reproductive success, which is highly sensitive to

temperature (Schou *et al.*, 2021, 2022); (3) quantify genetic variation in the efficiency of head

thermoregulation; and (4) investigate if ostriches originating from regions with different

climatic regimes have evolved differences in their head thermoregulation.

One complication with measuring variation in thermoregulation in free-ranging animals under

natural conditions is that it is not possible to control previous exposure to temperature stress,

for example, due to variation in microclimatic and biotic factors. A high temperature

measurement may therefore be because the animal was exposed to heat, has poor

thermoregulatory ability for a given temperature, or both. However, by calculating temperature

differences between the head and other body parts it is possible to estimate the ability of

individuals to regulate head temperatures for a given thermal stress load (as indicated by the

temperature of that body part). A high head temperature *relative to the body* cannot be due to

high heat exposure per se as both body parts are exposed to similar external conditions.

Consequently, the difference between head and body is likely to capture the specific ability of

individuals to regulate their head temperature, a proxy for vital brain functioning. In ostriches,

the neck provides an appropriate body part for comparison to the head as it is featherless,

reflecting skin temperature, moves in synchrony with the head, and individuals typically

orientate themselves such that the side of these two organs can be repeatedly captured by

thermal imaging. This provides control over distance, angle in the image, and similar skin

exposure to sun/shade at the time when being photographed (see Supplementary Information

for analyses to support that this approach minimises the effect of distance and angle of imaging

on measurements).

Results

Thermoregulation across naturally fluctuating temperatures

The surface temperatures (T) of the head and neck were both positively correlated with air

temperature (Fig. 2; Tables S1-2). The change in surface temperatures with increasing and

decreasing air temperatures was, however, significantly higher for the neck compared to the

head (Increasing air T_{Neck} vs Increasing air T_{Head} (CI) = 1.6 (1.2, 1.9), pMCMC = 0.001;

Decreasing air T_{Neck} vs Decreasing air T_{Head} (CI) = -2.1 (-3.1, -1.2), pMCMC = 0.001, Fig. 2C;

Table S3). Consequently, there were large differences between neck and head temperatures at

extreme temperatures (low < 20°C and high > 30°C), but not at benign air temperatures (20 to

30°C), where the need for thermoregulation is minimal (Fig. 2D, Table S4). Furthermore, even

though the head and neck differed in their rates of temperature change, they both increased

linearly with air temperature (Figs. 2A-B). There was no accelerating change in the surface

temperature of the neck (Fig. S1), as would be expected if it acted as a so-called 'thermal

radiator', directing heat away from the head at critical air temperature inflection points

(Tattersall *et al.*, 2009; Janse van Vuuren *et al.*, 2020). Therefore, head surface temperatures

appear to be more tightly regulated than neck surface temperatures at temperature extremes,

and there was no evidence that the neck is actively involved in head thermoregulation.

Reproductive consequences and evolutionary potential of head thermoregulation

Next, we examined if the ability to regulate head temperatures under a given thermal load, measured as the difference between head and neck temperatures, has the potential to evolve. We did this by first investigating the relationship of the head-neck temperature with reproductive success and secondly testing if variation across individuals was heritable. We found that females with lower head-neck temperatures during hot periods had higher egg-laying rates (Fig. 3A. $\text{Hot}_{\text{Head-Neck}}$ (CI) = -0.16 (-0.30, 0.00); pMCMC = 0.027; Table S6). Under more benign temperatures, however, there was no significant relationship between head-neck temperature and egg laying rates ($\text{Benign}_{\text{Head-Neck}}$ (CI) = -0.06 (-0.18, 0.05); pMCMC = 0.277; Table S6). This suggests that there is selection for the ability to cool the head during high temperatures via its effects on fecundity.

We further found that head-neck temperature had small but positive and significant heritabilities and evolvabilities (Hansen *et al.*, 2011) (h^2 ranged from 0.06 to 0.08. Fig. 3B, Table S7, see also Table S8-S9 for separate models of head and neck). Heritability estimates were similar and significantly different from zero across cold, benign and hot conditions, although measures of additive genetic variances and repeatabilities were slightly higher under cold and hot conditions. Heritabilities are capped by how repeatable measurements are across individuals, which can often be low for labile physiological traits, such as thermal profiles, when measured non-invasively under semi-natural settings where it is difficult to control for other potentially confounding variables (Figs. 1-2). Our repeatability estimates of the head-neck temperature deviation ranged from 0.12 to 0.17 (Fig. 3B, Table S7). It is therefore plausible that our estimates of heritability, while significant, could underestimate the true heritability of head-neck temperature due to the inherent difficulties in quantifying individual variation in

thermoregulatory traits.

Thermoregulatory ability differs between subspecies from different climatic regions

Finally, we investigated whether head-neck temperature showed some evolutionary signatures

in response to past climate conditions by comparing females from different ostrich subspecies

that originated from different geographic regions (Fig. 4A-B). At our study site in South Africa,

there are three ostrich subspecies from different regions of Africa: South African Blacks (SAB:

Struthio camelus), Zimbabwean Blues (ZB: *S. c. australis*) and Kenyan Reds (KR: *S. c.*

massaicus) that have evolved under different climatic regimes (Fig. 4A-C). The climate in East

Africa, where KR naturally occurs, is less seasonal, exhibiting lower fluctuations in temperature

and precipitation, than the regions where ZB and SAB subspecies occur (Fig. 4C). Although

the number of subspecies is limited, comparing individuals from these three subspecies raised

at the same site in South Africa allowed us to examine variation in head-neck temperature in

this 'common garden' setting.

There were pronounced differences in head-neck temperature between the subspecies in

response to increasing temperatures that corresponded to the stability of their past climates. The

head-neck temperature of KR females, which come from a stable thermal environment, changed

little in response to temperature increases (Fig. 4C, Table S12). This indicates similar rates of

change in head and neck temperature with air temperature, and therefore only modest

thermoregulation in response to increasing temperatures. In contrast, females from ZB and SAB

populations, which have their origins in the more variable thermal environments of southern

Africa, had head temperatures that did not increase as fast as neck temperatures with rising air

temperatures, suggesting a greater capacity to regulate their head temperature (Fig. 4D). In

contrast, head-neck temperatures during decreasing temperatures were similar across KRs,

SABs and ZABs. These differences in thermoregulatory capacity were not confounded by

differences in body size between the three subspecies (Tables S13-14, Figs. S2-3).

Discussion

Our study has revealed that the capacity to regulate head temperature is important for

reproductive success under fluctuating temperatures (Fig. 3). The efficiency of

thermoregulation appears to have a genetic basis and enables reproduction under a greater range

of temperatures (Fig. 3). Ostriches originating from geographic areas with more pronounced

temperature fluctuations were also more efficient at regulating head temperatures, revealed by

lower head relative to the neck temperatures. These results suggest that thermoregulation of the

head has evolved in response to past climates and may evolve in response to future climatic

change (Fig. 4).

Previous research has identified a variety of physiological and behavioural thermoregulatory

mechanisms (Kilgore *et al.*, 1976; Fuller *et al.*, 2003; Huey *et al.*, 2003; Angiletta, 2009, 2009;

Tattersall *et al.*, 2009; Symonds & Tattersall, 2010; Weissenböck *et al.*, 2010; Périard *et al.*,

2015; Muñoz & Losos, 2018). While such studies have provided compelling evidence for the

influence of various traits on thermal physiology, they have not provided direct estimates of

genetic variation in such traits or their effect on fitness. Such data is required to fully understand

how thermoregulatory traits influence future evolutionary responses to changing climatic

conditions. Our results demonstrate the power of combining measurements of reproductive

success and pedigree data with thermal profiles from individuals exposed to natural temperature

variation in realistic environmental settings. The differences (Fig. 4) between phenotypically

different subspecies from different climatic regions in combination with our findings of

significant additive genetic variance in head thermoregulation (Fig. 3), suggest that selection

for reproducing under fluctuating climates has shaped the evolution of the ostrich's remarkable

capacity for head cooling under hot conditions.

Decades of research on birds in temperate zones has focused on food availability in altricial

birds as a major limiting factor for reproduction (Lack, 1954; Williams, 1966). However, for

precocial birds inhabiting tropical and subtropical areas, like the ostrich, temperature stress

during reproduction might pose a more severe challenge than food limitation (Bourne *et al.*,

2020). Thermoregulation can keep body temperatures within non-lethal limits (Tattersall *et al.*,

2009; Symonds & Tattersall, 2010; Tattersall & Cadena, 2010; Darnell & Munguia, 2011;

McCafferty *et al.*, 2011; Szafrańska *et al.*, 2020), but heat waves can still compromise

population viability by decreasing individual reproductive success (Walsh *et al.*, 2019; Bourne

et al., 2020; Parratt *et al.*, 2021). Recent research from different taxa suggests that climate-

mediated local extinctions might already be common (Sinervo *et al.*, 2010; Wiens, 2016), and

there are worrying signs of recent collapses of some desert bird communities (Riddell *et al.*,

2019). Whether genetic variation in thermal tolerance is sufficient to drive evolutionary

responses to increasingly hot and fluctuating conditions remains an open question.

The low heritability estimates of thermal profiles in this study suggest a low but significant

evolutionary potential of ostrich populations to adapt to future temperature change. Previous

research on ectotherms, such as insects and reptiles, has also found thermal tolerance to have

low additive genetic variances or heritabilities, indicating that obtaining precise quantitative

genetic estimates for thermal tolerance may be difficult (Hoffmann *et al.*, 2003; Logan *et al.*,

2018; Castañeda *et al.*, 2019). Variation in body temperatures can also be nonadaptive or

maladaptive, a contention supported by recent empirical research (Ghalambor *et al.*, 2015;

Radersma *et al.*, 2020; Stamp & Hadfield, 2020; Svensson *et al.*, 2020). For example, plastic

changes in either core body temperatures or surface temperatures have been shown to reduce

fitness in insects (Svensson *et al.*, 2020), reptiles (Campbell-Staton *et al.*, 2021), and birds

(Nilsson *et al.*, 2016). Such maladaptive thermal plasticity may reflect costs of maintaining

homeostasis and stable body temperatures under thermally stressful conditions (Svensson *et al.*,

2020; Campbell-Staton *et al.*, 2021). It is likely that selection for thermal robustness on one

side, and costs of maintaining a stable body temperature on the other side, are shaping genetic

variance in thermoregulation in both the ostrich and other organisms inhabiting thermally

stressful environments like deserts. While challenging, combining studies on the genetics of

thermal tolerance with long-term population monitoring of reproduction and survival is key to

forecasting the potential effects of climate change on population viability. This is especially the

case for large, sub-tropical endotherms like the ostrich.

Methods

Study site, subspecies and general settings of enclosures

The study was conducted at the Oudtshoorn Research Farm in the arid Klein Karoo region of

South Africa (GPS: 33° 38' 21.5"S, 22° 15' 17.4"E). Fenced enclosures (N=170) were used for

thermal imaging and to monitor the reproductive success of ostriches in male-female pairs

($N_{enclosures}$ = 148, ~0.25 ha per enclosure) (Cloete *et al.*, 2008) and groups ($N_{enclosures}$ = 22, ~0.47

ha per enclosure). These enclosures contained natural vegetation in the form of bushes and

small trees, making it possible for the animals to take shelter when needed. All individuals had

access to *ad libitum* food and water. The ostriches in this study belong to three different

subspecies: 1) the Masai ostrich (*Struthio camelus massaicus*), or Kenyan Red (KR), 2) the

Southern African ostrich, (*S. c. australis*), sometimes referred to as the Zimbabwean Blue (ZB)

because of its origin in Namibia and Zimbabwe, and 3) the South African Blacks (SAB), that

is thought to be of mixed origin, but is genetically similar to ZB (Davies et al 2012; unpublished

data). SAB are also referred to as *S. c. var. domesticus*. Individuals that had less than 85%

expected relatedness to a particular subspecies, as determined by the pedigree (see below), were

considered hybrids. Breeding birds were recruited from surviving chicks from previous years,

and parentage data were used to compile a 9-generation pedigree with 139 founding individuals.

Although 9 generations have passed since the first individuals were used to establish the study

populations, not all individuals are from F9. More than 85% of the photographed ostriches were

six years or younger. Ostriches are reproductively mature at age 2-3 years, whereby typically

three generations were present in any given year. Unlike the SAB and ZB populations which

were founded more than 30 years ago, the KR population was founded just 16 years ago

reducing the number of generations across its cohort substantially. Ethical clearance was

obtained from the Western Cape Department of Agriculture (DECRA R12/48).

Thermal imaging data

We took thermal images of ostriches in the enclosures using an infrared thermography camera

(H2640, NEC Avio Infrared Technologies; purchased from the company Senso-Test in

Sweden). Pictures were taken in the field, usually from distances between 2 m and 25 m. We

regularly calibrated the thermal camera against a black body during our field sessions, following

the instructions from the camera supplier, which also provided regular recalibration services of

our equipment.

Images that were out of focus were discarded prior to analysis, and we used the software InfRec

Analyzer to draw separate polygons within the head and neck regions of the ostrich in each

image. The average temperature of these polygons was extracted as individual head and neck

surface temperatures, respectively. We used the same procedures and default software settings

as in our previous work (Svensson & Waller, 2013; Svensson *et al.*, 2020), which assumes an

emissivity of 1. Biological structures, such as bird skin and feathers, typically have emissivity

values ranging between 0.94 -1.0 (Gerken *et al.*, 2006; Yahav & Giloh, 2012). However, the

exact emissivity value used does not change our results as we focused on relative temperature

measures that are independent of emissivity values. Hourly air temperatures were measured at

a weather station positioned 600 m from the study populations. We fitted a cubic spline to the

hourly temperature estimates of each day using the R-package *mgcv* v.1.8 (Wood, 2004), from

which we extracted the predicted air temperature at the time-points when thermal images were

taken.

Datasets and analyses

3.1) Thermal imaging dataset

From 2012 to 2017 during October to December, we took 2744 pictures of 423 females between

5 am and 6 pm (average images per female = 6.5). This dataset was designed to maximize the

number of individuals with repeated measures within and across years, in order to characterize

individual thermal profiles across different thermal environments. Consequently, there was

little repeated sampling within days (8% of the pictures).

3.2) General modelling details

We constructed and ran generalized linear mixed models (GLMMs) in R v.3.6.0 (R Core Team,

2020) using the Bayesian framework implemented in the R-package MCMCglmm v.2.29

(Hadfield, 2010). For random terms, we used the weakly informative inverse-Gamma

distribution (scale = 0.001, shape = 0.001, i.e. *V = diag(n), nu =n-1+0.002, with n being the*

dimension of the matrix) as priors. Unless otherwise stated, each model was run for 5,100,000

iterations of which the initial 100,000 were discarded and only every 4,000th iteration was used

for estimating posterior probabilities. The number of iterations was based on the inspection of

autocorrelation among posterior samples in preliminary runs. Convergence of the estimates was

checked by running the model three times and inspecting the overlap of estimates in trace plots

and the level of autocorrelation among posterior samples. We report the results from the first

of these three runs. Posterior mode and 95% credible intervals are reported for random effects.

In all analyses, we accounted for environmental effects that varied across years, such as diet,

by including year as a random effect. Photographs were taken across 48 days, and we therefore

included date as a random effect. We also added enclosure as a random effect as they were

repeatedly used across years and varied in vegetation cover, potentially impacting on the local

climatic conditions. Individual ID was also included as a random effect, as all analyses

contained multiple records per individual.

3.3) Are head surface temperatures more tightly regulated than neck surface temperatures?

To estimate the change in surface temperatures with changing air temperatures we first

constructed models with either head or neck temperatures as the Gaussian response variable.

Hereafter we ran a model including both head and neck temperatures to allow comparison of

the two body parts. We examined increases and decreases in air temperature from a thermal

optimum as this allows us to estimate separate parameters for the responses to decreases in

temperature (cold tolerance) and increases in temperature (heat tolerance). A previous

investigation showed the reproductive success of ostriches is highest at a daily maximum

temperature of ~20°C (Schou *et al.*, 2021). We therefore defined 20°C as the optimum air

temperature for ostriches, and calculated absolute *temperature change* away from this optimum,

with a factor *direction of change* used to denote if it was a decrease or an increase in temperature

from the optimum. To make the intercept in statistical models represent the most benign air

temperature, we set 20°C to 0 and calculated deviations above (increases) and below (decreases)

this value. The variance of slopes (e.g. *temperature change,* see below) depends on the scale of

the parameter. We therefore standardized our data by dividing it by the maximum of the air

temperature range, resulting in 1 being the maximum temperature change.

Models included the fixed effects of temperature change (ranging from 0 to 1) and direction

(decreases or increases). The interaction between temperature change and direction was

modelled with a common intercept for decreases and increases, as the way temperature change

was calculated dictated that the intercepts were identical. We included the fixed effects of

subspecies (SAB, ZB, KR or hybrids) and both the linear and quadratic terms of time of day

(scaled and centered to a mean of zero and unit variance). We included interactions between

subspecies, temperature change and direction. In addition to the random effects common for all

models (year, enclosure, date and individual ID), we also interacted individual ID with

temperature change and direction, to allow independent rates of change in surface temperature

of each female. This was modelled as a 3x3 unstructured variance-covariance matrix.

The model with both head and neck surface temperatures as response variable contained

additional fixed and random effects. To identify the source of the surface temperature we

included the fixed effect body part (factorial: neck or head) which was interacted with all the

previously described fixed effects. We also added image as an additional random effect because

head and neck surface temperatures of an individual at a given time were derived from the same

image.

3.4) Does the neck actively regulate the temperature of the head?

To investigate if the neck acts as a thermal radiator actively regulating head temperature, we

inspected the relationship between neck surface temperature and air temperature. When air

temperature increases towards the body temperature of the ostrich, the difference between air

and surface temperature will decrease. If the neck is a thermal radiator, we would expect that

as the air temperature reaches certain inflection points, blood is shunted to the surface of the

neck to emit more heat, increasing the neck temperature and thereby increasing the difference

between neck surface temperatures and air temperature (Tattersall *et al.*, 2009; Janse van

Vuuren *et al.*, 2020). If instead the neck does not function as an actively regulated thermal

radiator, we expect a linear relationship between air temperature and neck temperature.

3.5) Effect of head-neck temperature on reproductive success

To test if the capacity to regulate head temperature influences reproductive success we analysed

its relationship with rates of egg-laying. Microclimatic factors (i.e. the operative environmental

temperature) and the behaviour up to the time of the thermal image being taken will influence

the absolute head surface temperatures. To better be able to compare head temperature

regulation across individuals, that always will differ in their recent behaviour and microclimatic

exposure, we used the neck, which moves in synchrony with the head, as a proxy for the heat

stress the animal is experiencing and what it has been exposed to (abiotic and biotic activity).

This was done by estimating the head-neck temperature (head surface temperature – neck

surface temperature) as a standardised measure of head thermoregulation. Individuals that show

lower head-neck temperatures when hot compared to others therefore regulate their head

temperature to a larger extent.

After estimating the head-neck temperature of individual females, we connected it to their egg-

laying records. From previous investigations we know that when the daily maximum

temperature is above or below 20°C, ostrich egg-laying rates start to decrease two to four days

later, possibly because this is the time it takes for the egg to travel down the oviduct (Schou *et*

al., 2021). The maximum temperature was below 20°C on just two of the days when thermal

images were taken, preventing us from examining the relationship between reproduction and

temperature regulation under cold conditions. In contrast, maximum temperatures were above

20°C in 41 days when thermal images were taken allowing responses to heat and reproduction

to be analysed.

We examined the probability of eggs being laid two to four days after females were thermal

imaged on days exceeding 20°C. If females with lower head-neck temperatures have a higher

probability of laying an egg during this time window, this would indicate that the

thermoregulatory capacity of the head is associated with reproductive success. Any such

associations may not be causal, as it is possible that reproduction itself causes an increase in

surface temperatures, for example by impacting metabolic rate. Assuming that any impact of

reproduction on surface temperatures is similar during hot and benign times of the day, we

grouped the data into photographs taken during benign (air temperatures > 20°C & < 30°C,

n_{images} = 615) and hot (air temperatures >= 30°C, n_{images} = 471) times of the day. Only if the

relationship between head-neck temperature and egg-laying is most pronounced when hot,

would we then expect it to be a consequence of improved thermoregulation under heat stress.

The data was analysed using a model with the probability of laying (binary, model type:

"threshold") as the response variable. The fixed effects included were head-neck temperature

(scaled and centered to a mean of zero and unit variance), air temperature category (hot or

benign), and subspecies (KR, SAB, ZB or Hybrid), as well as the interaction between head-

neck temperature and temperature category. As females of two years of age lay fewer eggs than

older females (Schou *et al.*, 2021), a factor of age (2 versus >2) was also included as a fixed

effect. We included year, enclosure, date and individual ID as random effects. The first 45 days

of the breeding season were removed as this is the average time it takes for pairs to acclimate

to each other and their enclosure (Schou *et al.*, 2021). We also removed females that laid fewer

than ten eggs per year (5% of all females) to avoid including females from incompatible pairs,

and individuals that were not in breeding condition. The models were run for 31,500,000

iterations of which the initial 1,500,000 were discarded and only every 10,000th iteration was

used for estimating posterior probabilities.

3.6) Is there genetic variation in the efficiency of head thermoregulation?

To investigate the evolutionary potential of head thermoregulation we constructed models to

estimate the repeatability and heritability when cold (air temperatures <= 20°C, n_{images} = 849),

benign (air temperatures > 20°C & < 30°C, n_{images} = 1069) and hot (air temperatures >= 30°C,

n_{images} = 826). The grouping of air temperature into these three categories was based on the

thermal neutral zone of the emu (Maloney, 2008), and to ensure roughly equal replication within

the cold and hot categories. Models had head-neck temperature (Gaussian) as the response

variable, and we included the fixed effect temperature category (factorial: cold, benign or hot).

These categories do not capture the effects of small deviations in air temperature, and we

therefore added a continuous measure of the deviation from the mean air temperature in each

category. This fixed effect was constructed by centering and scaling the air temperature records

within each air temperature category. We also included the fixed effects of subspecies (SAB,

ZB, KR or hybrids) and both the linear and quadratic terms of time of day (scaled and centered

to a mean of zero and unit variance). We included interactions between subspecies and

temperature category. In addition to the random effects common for all models (year, enclosure,

date and individual ID), we also interacted individual ID with air temperature category, to allow

independent rates of change in surface temperature of each female. This was modelled as a 3x3

unstructured variance-covariance matrix. We also estimated the residual variance separately for

each air temperature category.

Repeatability of head-neck temperatures under different temperature categories were estimated

using the estimates of permanent individual variances (*pe*) from the variance-covariance matrix

of individual ID by air temperature category:

$$(1)\ R = \frac{\sigma^2_{pe}}{\sigma^2_{pe} + \sigma^2_{year} + \sigma^2_{enclosure} + \sigma^2_{res}}.$$

The individual variances and covariances in head-neck temperature may originate from both

environmental and genetic factors. To control for maternal effects post laying, eggs are removed

from the nest twice a day and artificially incubated. Hereafter chicks were reared in small pens

in mixed groups with access to shade, and with *ad libitum* food and water - but without adult

ostriches. These experimental procedures should experimentally remove the majority of any

maternal effects. We also tried to estimate maternal effects by fitting dam identity in our

statistical models. However, such dam effects were close to zero and were therefore excluded

from our final models, as they did not significantly improve the explanatory power of our

models. This suggests that such maternal effects – even if present – do not persist in to

adulthood and are therefore not affecting adult thermal profiles.

To partition the among-individual variance that is due to additive genetic effects we added a

second 3x3 unstructured variance-covariance matrix of individual ID linked to the pedigree (*a*).

With these variance components, we estimated the narrow sense heritability (h^2) of head-neck

temperature in each air temperature category as the proportion of phenotypic variance

attributable to additive genetic variance:

$$(2)\ h^2 = \frac{\sigma^2_a}{\sigma^2_{pe} + \sigma^2_a + \sigma^2_{year} + \sigma^2_{enclosure} + \sigma^2_{res}}$$

We also estimated evolvability (I_A) (Houle, 1992):

$$(3)\ I_A = \frac{\sigma^2{}_a}{trait\ mean^2} * 100$$

One characteristic of evolvability is that it increases very fast as trait mean approaches zero.

When we initially used the posterior of the trait mean to estimate evolvability, this caused near-

infinity estimates of evolvability for some of the samples in the posterior, causing biased

estimates of the posterior mode and mean of evolvability. To avoid this, we used the posterior

mode of trait mean in the denominator, such that only the uncertainty of additive genetic

variance is included in the reported estimate of evolvability. We also ran identical models with

head or neck surface temperature as the response variable. The outcome of these analyses is

available in the supplementary materials (Tables S8-S9). Finally, we also used a similar model

to investigate if average feather temperatures on the side of the abdomen better captures

temperature exposure of the animal than recorded air temperatures. This model was run without

the animal term, and had temperature categories based on individual feather temperature instead

of recorded air temperatures. However, temperature measurements of feathers were highly

variable (Figure S8), likely because the feathers could have been facing the sun or away from

the sun in the time up until the picture. Indeed, estimated repeatabilities of the head-neck

difference was not higher when using feather temperatures instead of air temperatures (Table

S5).

3.7) Current distribution of the ostrich subspecies

The different ostrich subspecies come from different regions of Africa that may differ in climate

and therefore impact the need for thermoregulation. We obtained estimates of the current

distributions of *S. c. massaicus* (KR) and *S. c. australis* (ZB) were obtained by downloading

region-based presence/absence data from Avibase (https://avibase.bsc-eoc.org, September,

2020) and plotting these using the R-package "rnaturalearth" v. 0.1.0. To characterise the

climate of the distributions of each subspecies, we downloaded 19 bioclimatic variables

(10min) from WorldClim v. 2 (Fick & Hijmans, 2017). We performed a Principal Component

Analyses (PCA) and inspected the loadings of the first four principal components after varimax

transformation (Table S11). Based on this inspection we described each principal component

by one or two bioclimatic variables to characterize climatic differences in among the

distributions of the ostrich subspecies.

3.8) Do ostriches originating from different climatic regions differ in thermoregulation?

To test if the subspecies differ in their responses to increasing and decreasing temperatures, we

modelled head-neck temperature in a random regression model. This approach also allows us

to test if the head-neck temperature at the optimum temperature (the intercept) influences the

rate of change in head-neck temperatures as air temperatures increase or decrease (the slopes),

and if such a relationship is genetically based. Models included the fixed effects of temperature

change (ranging from 0 to 1) and direction (decreases or increases). The interaction between

temperature change and direction was modelled with a common intercept for decreases and

increases, as the way temperature change was calculated dictated that the intercepts were

identical. We included the fixed effects of subspecies (SAB, ZB, KR or hybrids) and both the

linear and quadratic terms of time of day (scaled and centered to a mean of zero and unit

variance). We included interactions between subspecies, temperature change and direction. In

addition to the random effects common for all models (year, enclosure, date and individual ID),

we also interacted individual ID with temperature change and direction, to allow independent

rates of change in surface temperature of each female. This was modelled as a 3x3 unstructured

variance-covariance matrix. To partition genetic and non-genetic effects contributing to the

relationship between temperature at the optimum and temperature change, we added a second

3x3 unstructured variance-covariance matrix of individual ID linked to the pedigree and

interacted with temperature change and direction. The genetic variance and co-variance was

then used to estimate the genetic correlation between the slopes and intercepts (correlation =

$covariance_{trait1,trait2}$ / sqrt(var_{trait1}*var_{trait2})) (Table S10).

To test if differences in the surface temperatures of the head and neck under increasing or

decreasing air temperatures across subspecies were caused by differences in body mass, we ran

a set of identical models also including individual body mass (scaled and centered to a mean of

zero and unit variance) as a fixed effect. We had records of body mass for all individuals and

when multiple records were available for one individual, we used the record closest to the time

of the thermal image.

3.9) The effects of distance and angle on surface temperatures

Animal surface temperatures measured using thermal imaging can be influenced by

confounding factors such as distance to the animal and animal orientation to the sun and wind

speed. We quantified the impact of distance to the animal on estimated surface temperatures,

using the number of pixels in the head and neck region as a proxy for distance. When we

inspected the relationship between number of pixels and absolute head or neck surface

temperatures, we found indications of more error in images where birds are further away.

However, in pictures taken far away, both head and neck are small, resulting in any biases being

accounted for by the head-neck metric (Fig S9). The angle of the animal is also important, and

ostriches typically orientate themselves side on to people so that the side of the head and neck

consistently face the investigator. This ensures that the majority of pictures are taken from the

same angle. To quantify this, we recorded the angle of the head in a subset of pictures and found

87% included the side of the head, and found no effect of the angle of the head on the head-

neck metric (Fig S10). In conclusion, our measures of head-neck temperature appear robust to

confounding variables such as distance and angle. This is in line with several independent

studies that have shown to surface temperatures to correlate with internal body temperatures

(McCafferty *et al.*, 2015; Andreasson *et al.*, 2016; Szafrańska *et al.*, 2020; Gauchet *et al.*, 2022)

and that the effect of confounding variables on measurements of internal body temperatures are

relatively modest (Gauchet *et al.*, 2022). Finally, additional details about our statistical analyses

of thermal imaging data and results are provided in the Supplementary Materials (Tables S1-

S16 and Figs. S1-S10).

Acknowledgements

We thank the staff and workers at Oudtshoorn Research Farm for assistance with data collection

and maintenance of the birds and the Western Cape Government for use of their resources. We

also thank members of "Svensson Lab" (students and assistants) who helped with analyzing the

thermal images, and Andreas Nord and Fredrik Andreasson for valuable feedback on an earlier

version of this manuscript. The computations were performed on resources provided by SNIC

through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX)

under Project SNIC 2018/8-359.

Author Contribution

610 Conceptualization: E.I.S, C.K.S., M.F.S.; Data curation: E.I.S., C.K.C., M.F.S., A.E., Z.B.,

611 S.C.; Formal analysis: M.F.S.; Funding acquisition: C.K.C, E.I.S., M.F.S, S.C.; Investigation:

612 E.I.S., C.K.C., J.M., J.W., M.F.S., S.C., Z.B; Methodology: E.I.S., C.K.C., M.F.S.; Project

613 administration: E.I.S., C.K.C., M.F.S.; Writing- original draft E.I.S, C.K.C., M.F.S.; Writing –

614 reviewing and editing: A.E., CKC, E.I.S., J.M., J.W., M.F.S, S.C., Z.B.

Funding

Carlsberg Foundation (MFS)

Swedish Research Council grant 2017-03880 (CKC)

Knut and Alice Wallenberg Foundation grant 2018.0138 (CKC)

Carl Tryggers grant 12: 92 & 19: 71 (CKC)

Carl Tryggers grant 19: 71 (CKC)

Swedish Research Council grant 2016-03356 (EIS)

Western Cape Agricultural Research Trust grant 0070/000VOLSTRUISE (SC)

Technology and Human Resources for Industry program of the South African National

Research Foundation grant TP14081390585) (SC)

Data Accessibility

All data extracted from thermal images are available at https://osf.io/fu2wx/ and will be uploaded in an open digital repository such as Dryad Digital Repository (https://datadryad.org/stash) upon acceptance on this manuscript. The remaining data used support the findings of this study are available from the Western Cape Department of Agriculture in South Africa (WCDA). Restrictions apply to the use of some of these data, which are thus not publicly available. These data are however available from the WCDA upon reasonable request. Code for analyses is available on Github: www.github.com/abumadsen/thermal-images-ostrich/tree/main

Competing interests

Authors declare that they have no competing interests.

Supporting Information

640   Tables S1-S16

641   Figures S1-S10

References

Andreasson, F., Nord, A. & Nilsson, J.-Å. (2016) Brood size constrains the development of endothermy in blue tits. *Journal of Experimental Biology*, **219**, 2212–2219.

Angiletta, M.J. (2009) *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press, Oxford.

Atwood, T.B., Valentine, S.A., Hammill, E., McCauley, D.J., Madin, E.M.P., Beard, K.H., *et al.* (2020) Herbivores at the highest risk of extinction among mammals, birds, and reptiles. *Science Advances*, **6**, eabb8458.

Beltrán, I., Herculano-Houzel, S., Sinervo, B. & Whiting, M.J. (2021) Are ectotherm brains vulnerable to global warming? *Trends in Ecology & Evolution*, **36**, 691–699.

Bogert, C.M. (1949) Thermoregulation in reptiles, a factor in evolution. *Evolution*, **3**, 195–211.

Bourne, A.R., Cunningham, S.J., Spottiswoode, C.N. & Ridley, A.R. (2020) Hot droughts compromise interannual survival across all group sizes in a cooperatively breeding bird. *Ecology Letters*, **23**, 1776–1788.

Campbell-Staton, S.C., Velotta, J.P. & Winchell, K.M. (2021) Selection on adaptive and maladaptive gene expression plasticity during thermal adaptation to urban heat islands. *Nature Communications*, **12**, 6195.

Cardillo, M., Mace, G.M., Jones, K.E., Bielby, J., Bininda-Emonds, O.R.P., Sechrest, W., *et al.* (2005) Multiple causes of high extinction risk in large mammal species. *Science*, **309**, 1239–1241.

Castañeda, L.E., Romero-Soriano, V., Mesas, A., Roff, D.A. & Santos, M. (2019) Evolutionary potential of thermal preference and heat tolerance in Drosophila subobscura. *Journal of Evolutionary Biology*, **32**, 818–824.

Cloete, S.W.P., Engelbrecht, A., Olivier, J.J. & Bunter, K.L. (2008) Deriving a preliminary breeding objective for commercial ostriches: an overview. *Australian Journal of Experimental Agriculture*, **48**, 1247–1256.

Darnell, M.Z. & Munguia, P. (2011) Thermoregulation as an alternate function of the sexually dimorphic Fiddler Crab claw. *American Naturalist*, **178**, 419–428.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., *et al.* (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 6668–6672.

Eastick, D.L., Tattersall, G.J., Watson, S.J., Lesku, J.A. & Robert, K.A. (2019) Cassowary casques act as thermal windows. *Scientific Reports*, **9**, 1966.

Fick, S.E. & Hijmans, R.J. (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, **37**, 4302–4315.

Fuller, A., Kamerman, P.R., Maloney, S.K., Mitchell, G. & Mitchell, D. (2003) Variability in brain and arterial blood temperatures in free-ranging ostriches in their natural habitat. *Journal of Experimental Biology*, **206**, 1171–1181.

Gauchet, L., Jaeger, A. & Grémillet, D. (2022) Using facial infrared thermography to infer avian body temperatures in the wild. *Marine Biology*, **169**, 57.

Gerken, M., Afnan, R. & Dörl, J. (2006) Adaptive behaviour in chickens in relation to thermoregulation. *Poultry Science*, **70**, 199–207.

Ghalambor, C.K., Hoke, K.L., Ruell, E.W., Fischer, E.K., Reznick, D.N. & Hughes, K.A. (2015) Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*, **525**, 372–375.

Hadfield, J.D. (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The **MCMCglmm** *R* Package. *Journal of Statistical Software*, **33**, 1–22.

Hansen, T.F., Pelabon, C. & Houle, D. (2011) Heritability is not Evolvability. *Evolutionary Biology*, **38**, 258–277.

Hoffmann, A.A., Hallas, R.J., Dean, J.A. & Schiffer, M. (2003) Low potential for climatic stress adaptation in a rainforest Drosophila species. *Science*, **301**, 100–102.

Houle, D. (1992) Comparing evolvability and variability of quantitative traits. *Genetics*, **130**, 195–204.

Huey, R.B., Hertz, P.E. & Sinervo, B. (2003) Behavioral drive versus behavioral inertia in evolution: a null model approach. *American Naturalist*, **161**, 357–366.

Hurley, L.L., McDiarmid, C.S., Friesen, C.R., Griffith, S.C. & Rowe, M. (2018) Experimental heatwaves negatively impact sperm quality in the zebra finch. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20172547.

Janse van Vuuren, A.K., Kemp, L.V. & McKechnie, A.E. (2020) The beak and unfeathered skin as heat radiators in the southern ground-hornbill. *Journal of Avian Biology*, **51**.

Kellermann, V., Heerwaarden, B. van, Sgro, C.M. & Hoffmann, A.A. (2009) Fundamental evolutionary limits in ecological traits drive Drosophila species distributions. *Science*, **325**, 1244–1246.

Kellermann, V., Loeschcke, V., Hogenkamp, D., Kristensen, T.N., Flojgaard, C., David, J.R., *et al.* (2012a) Phylogenetic constraints in key functional traits behind species' climate niches: patterns of dessication and cold resistance across 95 Drosophila species. *Evolution*, **66**, 3377–3389.

Kellermann, V., Overgaard, J., Hoffmann, A.A., Flojgaard, C., Svenning, J.C. & Loeschcke, V. (2012b) Upper thermal limits of Drosophila are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 16228–16233.

Kilgore, D.L., Bernstein, M.H. & Hudson, D.M. (1976) Brain temperatures in birds. *Journal of comparative physiology*, **110**, 209–215.

Lack, D. (1954) *The natural regulation of animal numbers*. Clarendon, Oxford.

Logan, M.L., Curlis, J.D., Gilbert, A.L., Miles, D.B., Chung, A.K., McGlothlin, J.W., *et al.* (2018) Thermal physiology and thermoregulatory behaviour exhibit low heritability despite genetic divergence between lizard populations. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20180697.

Ma, F.-Z., Lü, Z.-C., Wang, R. & Wan, F.-H. (2014) Heritability and evolutionary potential in thermal tolerance traits in the invasive Mediterranean cryptic species of Bemisia tabaci (Hemiptera: Aleyrodidae). *PLOS ONE*, **9**, e103279.

Maloney, S.K. (2008) Thermoregulation in ratites: a review. *Australian Journal of Experimental Agriculture*, **48**, 1293–1301.

McCafferty, D.J., Gallon, S. & Nord, A. (2015) Challenges of measuring body temperatures of free-ranging birds and mammals. *Animal Biotelemetry*, **3**, 33.

McCafferty, D.J., Gilbert, C., Paterson, W., Pomeroy, P.P., Thompson, D., Currie, J.I., *et al.* (2011) Estimating metabolic heat loss in birds and mammals by combining infrared thermography with biophysical modelling. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, The challenge of measuring energy expenditure: current field and laboratory methods, **158**, 337–345.

Moore, M.P., Lis, C., Gherghel, I. & Martin, R.A. (2019) Temperature shapes the costs, benefits and geographic diversification of sexual coloration in a dragonfly. *Ecology Letters*, **22**, 437–446.

Moreira, M.O., Qu, Y.-F. & Wiens, J.J. (2021) Large-scale evolution of body temperatures in land vertebrates. *Evolution Letters*, **5**, 484–494.

Muñoz, M.M. (2022) The Bogert effect, a factor in evolution. *Evolution*, **76**, 49–66.

Muñoz, M.M. & Losos, J.B. (2018) Thermoregulatory behavior simultaneously promotes and forestalls evolution in a tropical lizard. *The American Naturalist*, **191**, E15–E26.

Nilsson, J.-Å., Molokwu, M.N. & Olsson, O. (2016) Body temperature regulation in hot environments. *PLoS ONE*, **11**, e0161481.

Parratt, S.R., Walsh, B.S., Metelmann, S., White, N., Manser, A., Bretman, A.J., *et al.* (2021) Temperatures that sterilize males better match global species distributions than lethal temperatures. *Nature Climate Change*, **11**, 481–484.

Périard, J.D., Racinais, S. & Sawka, M.N. (2015) Adaptations and mechanisms of human heat acclimation: Applications for competitive athletes and sports. *Scandinavian Journal of Medicine & Science in Sports*, **25**, 20–38.

R Core Team. (2020) *R: A language and environment for statistical computing*. R foundation for statistical computing, Vienna, Austria.

Radersma, R., Noble, D.W.A. & Uller, T. (2020) Plasticity leaves a phenotypic signature during local adaptation. *Evolution Letters*, **4**, 360–370.

Riddell, E.A., Iknayan, K.J., Wolf, B.O., Sinervo, B. & Beissinger, S.R. (2019) Cooling requirements fueled the collapse of a desert bird community from climate change. *Proceedings of the National Academy of Sciences of the United States of America*, **116**, 21609–21615.

Sales, K., Vasudeva, R., Dickinson, M.E., Godwin, J.L., Lumley, A.J., Michalczyk, Ł., *et al.* (2018) Experimental heatwaves compromise sperm function and cause transgenerational damage in a model insect. *Nature Communications*, **9**, 4771.

Schou, M.F., Bonato, M., Engelbrecht, A., Brand, Z., Svensson, E.I., Melgar, J., *et al.* (2021) Extreme temperatures compromise male and female fertility in a large desert bird. *Nature Communications*, **12**, 666.

Schou, M.F., Engelbrecht, A., Brand, Z., Svensson, E.I., Cloete, S. & Cornwallis, C.K. (2022) Evolutionary trade-offs between heat and cold tolerance limit responses to fluctuating climates. *Science Advances*, **8**, eabn9580.

Sinervo, B., Mendez-de-la-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagran-Santa, C.M., *et al.* (2010) Erosion of lizard diversity by climate change and altered thermal niches. *Science*, **328**, 894–899.

Stamp, M.A. & Hadfield, J.D. (2020) The relative importance of plasticity versus genetic differentiation in explaining between population differences; a meta-analysis. *Ecology Letters*, **23**, 1432–144.

Svensson, E.I., Gomez-Llano, M.A. & Waller, J.T. (2020) Selection on phenotypic plasticity favors thermal canalization. *Proceedings of the National Academy of Sciences of the United States of America*, **117**, 29767–29774.

Svensson, E.I. & Waller, J.T. (2013) Ecology and sexual selection: evolution of wing pigmentation in calopterygid damselflies in relation to latitude, sexual dimorphism and speciation. *American Naturalist*, **182**, E174–E195.

Symonds, M.R.E. & Tattersall, G.J. (2010) Geographical variation in bill size across bird species provides evidence for Allen's Rule. *American Naturalist*, **176**, 188–197.

Szafrańska, P.A., Andreasson, F., Nord, A. & Nilsson, J.-Å. (2020) Deep body and surface temperature responses to hot and cold environments in the zebra finch. *Journal of Thermal Biology*, **94**, 102776.

Tattersall, G.J., Andrade, D.V. & Abe, A.S. (2009) Heat exchange from the Toucan bill reveals a controllable vascular thermal radiator. *Science*, **325**, 468–470.

Tattersall, G.J. & Cadena, V. (2010) Insights into animal temperature adaptations revealed through thermal imaging. *Imaging Science Journal*, **58**, 261–268.

Tattersall, G.J., Chaves, J.A. & Danner, R.M. (2018) Thermoregulatory windows in Darwin's finches. *Functional Ecology*, **32**, 358–368.

Tsubaki, Y., Samejima, Y. & Siva-Jothy, M.T. (2010) Damselfly females prefer hot males:
higher courtship success in males in sunspots. *Behavioral Ecology and Sociobiology*, **64**,
1547–1554.
Walsh, B.S., Parratt, S.R., Hoffmann, A.A., Atkinson, D., Snook, R.R., Bretman, A., *et al.*
(2019) The impact of climate change on fertility. *Trends in Ecology & Evolution*, **34**, 249–
259.
Weissenböck, N.M., Weiss, C.M., Schwammer, H.M. & Kratochvil, H. (2010) Thermal
windows on the body surface of African elephants (Loxodonta africana) studied by infrared
thermography. *Journal of Thermal Biology*, **35**, 182–188.
Wiens, J.J. (2016) Climate-related local extinctions are already widespread among plant and
animal species. *PLOS Biology*, **14**, e2001104.
Williams, G.C. (1966) Natural selection, the costs of reproduction, and a refinement of Lack's
principle. *American Naturalist*, **100**, 687–690.
Williams, G.C. (1992) *Natural selection: Domains, levels and challenges*. Oxford University
Press, Oxford.
Wood, S.N. (2004) Stable and efficient multiple smoothing parameter estimation for
generalized additive models. *Journal of the American Statistical Association*, **99**, 673–686.
Yahav, S. & Giloh, M. (2012) Infrared Thermography - Applications in Poultry Biological
Research. In *Infrared Thermography* (ed. by Prakash, R.V.). InTech.
809

Figures and captions

Fig. 1. The ostrich inhabits thermally challenging desert environments such as the Karoo in South Africa.

A. The ostrich (*Struthio camelus*) is the world's largest living bird with a feathered body and featherless neck and

head. *Left*: female. *Right:* male (photograph by C. K. Cornwallis in Karoo National Park, Western Cape Province

of South Africa). **B.** Thermal image of a female (left) and male (right) ostrich at our study site in Oudtshoorn

(photograph by E. I. Svensson). Note how the long neck is hot and emits excess heat (red, warm colour). **C.** The

dry and tree-less semi-desert environment around the study site is characterized by extensive temperature

fluctuations ranging from -5°C to 45°C, causing extreme thermal stress (Schou *et al.*, 2021) (photograph by E. I.

Svensson).

Fig. 2. Neck temperatures show how ostriches thermoregulate their head temperatures. **A)** When air temperatures increase above 20°C, the surface temperatures of the head increase slower than the neck (n_{images} = 1895, Tables S1-S2). **B)** Similarly, when air temperatures decrease below 20°C the reduction in surface temperatures are greatest in the neck (n_{images} = 849). Three extreme datapoints in A were removed for graphical purposes (see Fig. S4). The larger points are averages with standard errors binned according to temperature. Point size illustrates relative number of individuals: smallest point = 2 and largest point = 71. Fitted lines and 95% credible intervals (shaded area) were extracted from the statistical models **(C)** The rate of surface temperature change was steeper for the neck compared to the head (Table S3). This difference in rates of temperature increase between the two body parts was consistent across both decreasing and increasing temperatures ($n_{individuals}$ = 423). **(D)** The difference in rates of temperature increase between the head and neck led to higher discrepancy between neck to head surface temperatures during hot conditions (air temperatures >= 30°C, n_{images} = 826) and cold (air temperatures <= 20°C, n_{images} = 849) compared to benign (air temperatures > 20°C & < 30°C, n_{images} = 1069)

833 conditions (Table S4).

Fig. 3. The evolutionary potential for head temperature regulation. The capacity of individuals to regulate

their head temperature was estimated as the difference between head and neck temperatures in relation to air

temperature. **A)** On hot days (daily maximum > 20°C, n_{days} = 35), when temperatures were highest (air

temperatures >= 30°C, n_{images} = 471, $n_{females}$ = 228) an increased capacity to regulate head temperature (i.e. a lower

head than neck temperature) was associated with a higher egg-laying rate two days later. There is no such

association when examining thermoregulation at times when temperatures were benign of the hot days (air

temperatures = 20-30°C, n_{images} = 615, $n_{females}$ = 220) (Table S6). Maximum temperatures below 20°C were

recorded only on two days, preventing us from similar analyses on importance of thermoregulation during the cold

times of cold days. Fitted lines and 95% credible intervals (shaded area) were extracted from the statistical model

B) Repeatability and heritability of the head-neck temperature deviation were estimated at different air

temperatures measured at the time when thermal images were taken (cold < 20°C, benign = 20-30°C and hot >=

30°C, Table S7). Note that the heritability of these thermal profiles under cold conditions could be estimated

because there were cold periods (typically mornings) on days where temperatures exceeded 20°C. This contrasts

to selection analyses where temperature per day had to be used because of the two day time lag between egg

formation and egg laying.

Fig. 4. Ostriches originating from more variable climates exhibit larger decreases in head-neck temperatures when hot. **(A-C)** Kenyan Reds (KR) inhabit eastern Africa that is less seasonal and has lower temperature fluctuations compared to Southern Africa where Zimbabwean Blues (ZB) and South African Blacks (SAB) occur. Distribution ranges in A were estimated from regional presence/absence data from Avibase (https://avibase.bsc-eoc.org) and climatic data was obtained from WorldClim (Fick & Hijmans, 2017). Average temperature and seasonality were the main bioclimatic variables of PC1 (43%) and PC2 (29%) in a PCA of 19 bioclimatic variables (Fig. S5 & Table S11). **(D)** For both the SAB ($n_{individuals}$ = 264) and ZB ($n_{individuals}$ = 36) ostriches, the head temperature deviation (head-neck) was reduced when temperatures increased, but this was not the case for KR ($n_{individuals}$ = 23) (Table S12, see also Figs. S6-7 which includes hybrid individuals).